The DNA Replication Checkpoint Targets the Kinetochore for Relocation of Collapsed Forks to the Nuclear Periphery

Running Title: DRC targets Cep3 for fork relocation

Keywords: Nuclear Pore, Collapsed Fork, DNA Replication Checkpoint, Damage Induced Microtubule, DNA Repair, Nuclear Periphery, Rad53, Dun1, TNR, Non-B DNA, Huntington's Disease, CAG Repeat

Authors: Tyler Maclay[*,1], Jenna Whalen[*,1], Matthew Johnson[1] & Catherine H. Freudenreich[1]
[1]Department of Biology, Tufts University, Medford, MA 02155
[*]Authors contributed equally

Abstract

Hairpin forming expanded CAG/CTG repeats pose significant challenges to DNA replication which can lead to replication fork collapse. Long CAG/CTG repeat tracts relocate to the nuclear pore complex to maintain their integrity. Forks impeded by DNA structures are known to activate the DNA damage checkpoint, thus we asked whether checkpoint proteins play a role in relocation of collapsed forks to the nuclear periphery in *S. cerevisiae*. We show that relocation of a $(CAG/CTG)_{130}$ tract is dependent on activation of the Mrc1/Rad53 replication checkpoint. Further, checkpoint-mediated phosphorylation of the kinetochore protein Cep3 is required for relocation, implicating detachment of the centromere from the spindle pole body. Activation of this pathway leads to DNA damage-induced microtubule recruitment to the repeat. These data suggest a role for the DNA replication checkpoint in facilitating movement of collapsed replication forks to the nuclear periphery by centromere release and microtubule-directed motion.

Highlights

- The DNA replication checkpoint initiates relocation of a structure-forming CAG repeat tract to the nuclear pore complex (NPC)
- The importance of Mrc1 (hClaspin) implicates fork uncoupling as the initial checkpoint signal
- Phosphorylation of the Cep3 kinetochore protein by Dun1 kinase allows for centromere release, which is critical for collapsed fork repositioning
- Damage-inducible nuclear microtubules (DIMs) colocalize with the repeat locus and are required for relocation to the NPC
- Establishes a new role for the DNA replication and DNA damage checkpoint response to trigger repositioning of collapsed forks within the nucleus.

Introduction

The nuclear periphery (NP) has emerged as an important site for the maintenance of genome stability when normal replication is impaired. Several types of replication stress have been shown to provoke relocation to the nuclear periphery in both yeast and human cells. These include drugs known to stall or collapse replication forks such as hydroxyurea combined with methyl methanesulfonate (MMS) in yeast [1], expanded CAG repeats that form secondary

structures [2], the polymerase inhibitor aphidicolin in human cells [3], protein mediated stalls by
replication fork barriers (RFBs) [4,5], and replication stalls within telomeres [6,7]. Some persistent
types of DNA damage such as double-strand breaks (DSBs) lacking a readily available template
for repair or in sequestered heterochromatin domains and eroded telomeres also relocate the
nuclear periphery (for review, see [8–10]).
We previously elucidated a detailed pathway for how collapsed forks caused by an
expanded CAG/CTG tract of 130 repeat units (abbreviated $(CAG)_{130}$) relocates to the *S.*
cerevisiae nuclear pore complex (NPC) in late S phase in a manner dependent on replication [2,11].
CAG/CTG repeats over 35 repeat units have a propensity to form hairpin structures that can
impair replication leading to fork collapse [12,13]. Defects in relocation to the NPC result in
increased chromosome end loss and repeat instability, implicating this pathway in protecting
against fork breakage and inaccurate repair [2]. The DNA structure induced relocation pathway has
many similarities with pathways that have been described for persistent or heterochromatic DSBs
and eroded telomeres, but also several differences (reviewed in [8]). We determined that relocation
of collapsed forks caused by CAG repeats was dependent on resection, the binding and
SUMOylation of key fork associated repair proteins (RPA, Rad52, Rad59, Smc5) by the Smc5-6
associated SUMO ligase Mms21, and interaction with the SUMO interacting motifs (SIMs) of
Slx5 which targets the collapsed fork to the NPC [11].
In addition to the SUMOylation requirements, we wanted to investigate other processes
that could be involved in relocation of collapsed forks caused by CAG repeats to the NPC. One
important pathway known to respond to stalled and collapsed forks is the DNA damage
response. The DNA damage response can be subdivided into two pathways: the DNA replication
checkpoint (DRC) and the DNA damage checkpoint (DDC), defined by their mediator proteins
Mrc1 or Rad9, respectively. Our previous findings showed that expanded CAG repeats activate
the DDC [14], and long range resection is required for $(CAG)_{130}$ relocation [11], which would
produce ssDNA to stimulate DDC activation. In addition, members of the DDC and the DRC
were important for maintaining repeat stability and preventing chromosome breaks at long CAG
tracts [15–17].
The checkpoint is composed of several sensor and mediator components that activate
downstream effectors, reviewed in [18,19] and summarized in (Figure 1A). Sensor proteins identify
processed DNA lesions, such as resected DNA substrates, and then recruit other proteins to the
damage site. In *S. cerevisiae*, the sensor Ddc2 (hATRIP) interacts with RPA bound to ssDNA to
recruit the kinase Mec1 (ATR). Mec1 is also activated by the PCNA-like 9-1-1 clamp (Ddc1,
Rad17, Mec3 in budding yeast) which is loaded by the Rad24-RFC complex onto ds/ssDNA
junctions. Downstream effector proteins such as Rad53 (hChk2) and Chk1 elicit the DNA
damage response through further phosphorylation signaling. At the replication fork there is an
additional Rad53 activation mechanism involving the mediator, Mrc1 (hClaspin), which travels
with the replisome and is phosphorylated upon replisome uncoupling [20,21]. Another pathway
more specific to DSBs involves Tel1 (hATM), which senses DNA ends through its interaction
with the yeast MRX (hMRN) complex. Finally, the yeast mediator protein Rad9 (h53BP1)
amplifies Rad53 phosphorylation. Rad9 is recruited to histone modifications including
methylated H3 (H3-K79) and damage-specific H2A-S129 phosphorylation, which can occur at
either collapsed forks or DSBs. Due to the different DNA damage sensors, identification of the
checkpoint pathway that is activated can give insight into the lesion or lesions that are created by
a particular type of DNA damage.

The relocation of persistent DSBs in *S. cerevisiae* and heterochromatic DSBs in
Drosophila required both Mec1/Tel1 and ATR/ATM, respectively [1,22,23]. In human U2OS and
IMR90 cells that were treated with aphidicolin to inhibit polymerases, ATR inhibition prevented
the localization of stalled replication forks to the nuclear periphery [3]. In contrast, single deletion
of either Mec1 or Tel1 did not prevent the relocation of an expanded $(CAG)_{130}$ repeat to the NPC
[2], leaving it an open question whether the checkpoint is involved in the relocation of DNA
damage caused by DNA structures to the nuclear periphery, and if so what pathway is involved.
In yeast, Mec1/ATR is not only required for DSB relocation, but has been linked to increasing
chromatin mobility in response to DSBs [24]. Yet an expanded $(CAG)_{130}$ repeat not only failed to
increase mobility of a chromosome harboring this fork-stalling element, it decreased mobility
due to the added constraint of NPC interaction [2]. Therefore, it was not clear whether the
checkpoint pathway would be involved in relocation of DNA structure-mediated collapsed forks
to the NPC, and if so, in what manner.
In this study, we show that the activation of the effector kinase Rad53 by phosphorylation
of the replisome associated Mrc1 protein is needed for the relocation of a $(CAG)_{130}$ tract to the
nuclear periphery. We identified that the Rad53 target Dun1 is also required for relocation.
Through investigating several Dun1 substrates, we determined that the phosphorylation of the
kinetochore protein Cep3 at serine 575 is the important target to allow for repositioning of the
DNA structure-induced collapsed fork in the nucleus. We investigated several possible
mechanisms by which Cep3 phosphorylation could act and identified the disruption of
centromere attachment to the spindle pole body (the yeast microtubule organizing center which
is functionally equivalent to the mammalian centrosome) as a critical factor. In contrast, Cep3
phosphorylation-dependent increases in global chromosome movement were not detectable.
Finally, we show that damage-induced microtubules (DIMs) [25] are formed in response to damage
at the CAG tract, colocalize with the CAG tract locus, and are dependent on this checkpoint
pathway. We speculate that the phosphorylation of Cep3-S575 allows for relocation of collapsed
forks to the NPC by relieving centromere constraint and supporting DIM-directed motion
towards the nuclear periphery.
120
121
122 **Results**
123
Relocation of a $(CAG)_{130}$ tract to the nuclear pore complex depends on the activation of
Rad53 and Mrc1.
We used a previously described zoning assay [2] to visualize the CAG repeat in relation to
the nuclear periphery. Briefly, 130 CAG/CTG repeats are integrated on chromosome VI 6.4 kb
away from a LacO array which can be visualized by binding of GFP-LacI; in some experiments
the repeat was placed 6.4 kb from a TetO/mCherry-TetR array (Figure 1B). GFP-Nup49 was
used to mark NPCs to allow for visualization of the nuclear periphery. The nucleus was divided
into three equal areas to determine the frequency of CAG repeat occupancy in each zone (see
Methods). Zone 1 represents the nuclear periphery (where the NPCs are located), zone 3 is the
middle of the nucleus, and zone 2 is the area in between (Figure 1B). The presence of the
$(CAG)_{130}$ tract caused an increase in the occupancy of the nearby tagged locus in zone 1, from
31% in the $(CAG)_0$ control to 48% for the chromosome containing $(CAG)_{130}$ (Fig. 1C). The
distribution between zones 1, 2 and 3 was consistent between both wild-type $(CAG)_{130}$ strains,

whether the LacO/GFP-LacI or TetO/mCherry-TetR array was used to follow the CAG repeat
locus (Figure S1).
Previously, it was shown that single deletions in Mec1 or Tel1 sensor kinases (ATR and
ATM in mammals, respectively) do not affect relocation of the $(CAG)_{130}$ tract to the nuclear
periphery during S phase [2]. Simultaneous disruption of Mec1 and Tel1 could not be tested by the
zoning assay due to severely abnormal nuclear morphology. As there are many factors involved
in the activation of the DNA damage response that can act redundantly (Figure 1A), we
investigated if there was any role for the checkpoint in CAG tract relocation by testing the
effector kinases. The two effector kinases of the yeast DNA damage checkpoint are Chk1 and
Rad53 (reviewed in [18]). Deletion of Chk1 showed no significant decrease in the frequency the
CAG tract was found at the nuclear periphery (Figure 1C). However, deletion of Rad53 (viable
with the added deletion of Sml1 [26]) did result in a significant decrease in $(CAG)_{130}$ occupancy in
zone 1. This result supports that the DNA damage response is required for relocation. This was
further confirmed by testing $rad53^{K227A}$, a kinase-defective mutant of Rad53 that is checkpoint-
deficient [27]. $rad53^{K227A}$ showed the same significant decrease in relocation as $rad53\Delta sml1\Delta$
(Figure 1C). We conclude that activation of the DNA damage response and the kinase activity of
Rad53 is needed for relocation of the collapsed fork to the NPC.
There are multiple sensor and mediator proteins that can activate Rad53 in response to
different types of DNA damage. We had previously shown that long CAG tracts were especially
prone to breakage and exhibited increased fork stalling in the absence of the Mrc1 protein [17],
which couples polymerase epsilon to the CMG helicase. Mrc1 contains multiple SQ/TQ motifs
that are targeted by kinases of the PIKK family, such as Mec1 (hATR), Tel1 (hATM) or Rad53
(hChk2) for phosphorylation upon fork uncoupling [28,29]. The Mrc1^{AQ} mutant, with its 17 SQ/TQ
sites mutated to alanine [29], had a modest effect on chromosome fragility but exhibited increased
repeat instability [17], suggesting a possible role for Mrc1 in sensing forks stalled at the $(CAG)_{130}$
tract. The deletion of Mrc1 had severely abnormal nuclear morphology precluding zoning
analysis. However, complementation with the checkpoint-deficient Mrc1^{AQ} mutant abrogated
these abnormal morphologies and showed a significant reduction in relocation to the nuclear
periphery to a similar level as a Rad53 mutant (Figure 1C). Thus, the checkpoint signal relevant
for repositioning to the NPC is channeled through Mrc1, which suggests that the signal
originates from an uncoupled replication fork.
168
Mec1 and Rad9 act redundantly to sense damage at the $(CAG)_{130}$ tract and signal
repositioning within the nucleus.
Though single mutations in either of the main checkpoint sensor kinases, Mec1 and Tel1,
did not result in a decrease in relocation to the nuclear periphery (see Figure 1D) the presence of
the Mre11 protein, which senses and processes DSB ends, is required [11]. Deletion of the Rad9
mediator protein (h53BP1 ortholog), which acts downstream of Tel1 to facilitate Rad53
autophosphorylation [30,31], showed no effect on the relocation of the repeat to the nuclear
periphery (Figure 1D). Similarly, deletion of the 9-1-1 clamp component Ddc1 (9-1-1 activates
Mec1) or the clamp loader Rad24 also had no effect on the percent of the CAG locus in zone 1
(Figure 1D). These results were somewhat surprising given the important role of Mrc1, which is
phosphorylated by Mec1, and our previous findings that both Mec1 and Rad9 were important for
preventing breaks and instability at expanded CAG tracts [15]. Since there is a well-described
redundancy between these sensing pathways (reviewed in [18]), we explored whether there was
redundancy in the sensing of lesions to signal relocation of the $(CAG)_{130}$ to the nuclear

periphery. Deletion of both Mec1 and Rad9 resulted in a decrease in relocation of the $(CAG)_{130}$
to the nuclear periphery (Figure 1D). In addition, co-disruption of Rad9 and Rad24 also led to a
decrease in relocation to the NP, and deletion of all 3 genes (*mec1Δrad24Δrad9Δ*) led to the
lowest % zone 1 occupancy (Figure 1D). These effects were not made worse by additional
disruption of Tel1 (Figure 1D), suggesting that the role of Tel1 is not independent of Rad9 in this
situation. We previously showed that both H2A-S129 and H2A-T126 phosphorylation could be
detected at a $(CAG)_{155}$ tract 40 minutes into S phase [32,33], which is shortly before relocation
occurs [11] and could serve to recruit Rad9. Importantly, DNA damage dependent phosphorylation
of H2A requires either Mec1 or Tel1 [34,35]. So, in the absence of Mec1, the recruitment of Rad9
would be expected to be downstream of Tel1 activity. Thus, our data indicate that there are
multiple ways to activate Rad53 to enable relocation of the $(CAG)_{130}$ tract: a Tel1/Rad9 mediated
pathway, or a Rad24/Mec1 pathway, acting redundantly with the Rad9 pathway. In addition,
Mrc1 phosphorylation, which facilitates recruitment of Rad53 [36], is an important mediator of
Rad53 activation at the CAG-stalled fork.
Altogether, this analysis suggests that there may be multiple types of damage at the CAG
tract that can provoke the checkpoint response and are important for nuclear repositioning. Based
on our data, repeat mediated checkpoint activation and relocation to the NPC could be due to an
uncoupled fork (sensed by Mrc1), a region of single-stranded DNA (sensed by the Ddc2-Mec1
and 9-1-1 complexes), or a DSB end (sensed by the MRX-Tel1-Rad9 pathway). Single-strand
DNA could occur at an uncoupled fork, at a reversed fork, or at a post-replicative gap, and a
DSB end could result either from fork reversal or fork breakage/cleavage. All of these signals
would be present at an uncoupled fork that transitions to either a reversed or broken fork, which
we refer to collectively as "collapsed forks" (Figure 1E). Alternatively, there could be a mixture
of possible species that occur in different cells.
207
Phosphorylation of Cep3 Serine 575 by Dun1 is needed for relocation of collapsed forks to
the nuclear pore complex.
After determining that activation of the transducing kinase Rad53 by the DNA replication
checkpoint is required for $(CAG)_{130}$ relocation, we investigated what targets of Rad53 are
needed. Dun1 is a well characterized target of Rad53 (Figure 2A) [37–40]. A *dun1Δ* showed a
significant decrease in CAG tract relocation to the N.P. similar to *rad53Δsml1Δ* and *rad53^{K227A}*
mutants (Figure 2B). Therefore, Rad53 activation of Dun1 is a crucial step in the pathway
leading to repositioning of collapsed forks to the NPC.
Next, we investigated what targets of Dun1 are required for relocation (Figure 2A). Dun1
is required for the phosphorylation of the Rad51 associated repair protein Rad55 [37].
Phosphorylation of Rad55 allows for it to dimerize with Rad57 and effectively stabilize Rad51
binding to promote more successful strand invasion [41]. Therefore, Dun1 plays a direct role in
recombination mediated repair. However, neither *rad55Δ* nor *rad55Δrad57Δ* mutants showed a
decrease in the percentage of CAG tract loci detect in zone 1 (Figure 2B). Dun1 has also been
well studied as an activator of ribonucleotide reductase (RNR) activity and functions to regulate
nucleotide metabolism in response to replicative stress. Dun1 inhibits Rfx1, Sml1, and Dif1
through phosphorylation which in turn prevents them from downregulating nucleotide pools [42–
44]. Since Dun1 inhibits Sml1 activity, in order to determine if Dun1's role in relocation is
through maintaining nucleotide pools, Sml1 was deleted in the *dun1Δ* strain. If the deletion of
Sml1 rescues the defect in CAG tract repositioning in the *dun1Δ* strain, it would indicate that this
activity of Dun1 is required. However, *dun1Δsml1Δ* showed a similar percent S phase zone 1

occupancy as *dun1Δ* strains (Figure 2B). We conclude that the role for Dun1 in relocation to the
NP is neither to promote recombination mediated repair nor to increase nucleotide pools.
Another phosphorylation target of Dun1 is the kinetochore protein Cep3 [45]. Cep3 is a
DNA-binding protein that recognizes the centromere sequence as a member of the CBF3
kinetochore complex [46,47]. Phosphorylation of Cep3 at serine 575 (S575) occurs in response to an
induced DSB or zeocin-induced DNA damage in a manner dependent on both Rad53 and Dun1
[45]. The yeast kinetochore interacts with the spindle pole body (SPB), the yeast microtubule
organizing center which is functionally equivalent to the mammalian centrosome. Cep3-S575
phosphorylation has been reported to allow for an increase in the dynamics of the SPB-CEN
attachment after DSB induction on a different chromosome [45]. Mutation of serine 575 to a non-
phosphorylatable alanine residue resulted in a significant decrease in the CAG locus occupancy
in zone 1, showing that Cep3 phosphorylation is required for relocation of the CAG tract to the
NPC (Figure 2B).
To confirm that Cep3 phosphorylation at S575 is required for relocation we constructed a
phosphomimetic mutant of Cep3 by mutating serine 575 to glutamic acid. The *cep3^{S575E}*
mutation bypassed the need for Dun1 as the *dun1Δcep3^{S575E}* mutant showed a normal level of
CAG locus S phase occupancy in zone 1 (Figure 2C). We conclude that phosphorylation of
Cep3-S575 mediated by Dun1 is needed for relocation of collapsed forks to the NPC.
To further address the role of Cep3 phosphorylation we assessed the timing of CAG locus
relocation to the NP in *cep3^{S575E}* cells over the course of an S-phase. As previously reported, in
wild-type cells the $(CAG)_{130}$ tract is rarely at the NPC in early S-phase and is maximal in late S
phase at 60 min post release from G1 (Figure 2D) [11]. However Rad52 is found at the repeat locus
40 minutes into S phase [2], suggesting that fork collapse has occurred by then and that there is a
delay between the occurrence of damage and anchoring at the pore. The *cep3^{S575E}* mutation led to
a significant increase in zone 1 occupancy of the $(CAG)_{130}$ repeat tract at the 40- and 50-minute
time points. (Figure 2D). This suggests that the *cep3^{S575E}* phosphomimetic is bypassing the
normal signaling pathway, priming the fork for repositioning as soon as enough RPA-coated
ssDNA has occurred to recruit Rad52.
257
Phosphorylation of Cep3-S575 allows for relocation of collapsed forks to the NPC by
modulating kinetochore-SPB attachment, but not by increasing global chromosome
movement or activating the spindle assembly checkpoint.
It has been shown that forcing transcription through the centromere with a galactose-
inducible promoter causes detachment of the centromere from the SPB [48,49]. Cep3-S575
phosphorylation is needed for kinetochore declustering and increased SPB-CEN dynamics in
response to DSBs and has therefore been suggested to relieve attachment of centromeres to the
SPB [45], though the *cep3^{S575A}* mutation was not observed to increase SPB-CEN distance or
position with respect to the nuclear periphery after DSB induction [45,50]. We reasoned that if
centromere attachment to the SPB was preventing relocation to the NPC in the *cep3^{S575A}* mutant,
then forcing transcription through the centromere and thus interrupting that attachment could
allow for relocation of the repeat tract in the *cep3^{S575A}* mutant. We integrated the *GAL1* promoter
upstream of the centromere on chromosome 6 (*pGAL1-CEN6*) [51], which is the chromosome
containing the $(CAG)_{130}$ tract. Upon galactose induction and transcription through the
centromere, the reduced zone 1 occupancy of the CAG locus in the *cep3^{S575A}* strain was rescued
to normal wild-type levels (Figure 3A). We conclude that phosphorylation of Cep3-S575 may

allow for relocation of collapsed forks to the NPC by relieving centromere attachment to the
SPB.
Detachment of the centromere from the SPB has a number of consequences. It has been
shown that inactivating a centromere can increase the mobility of that chromosome [45]. It is also
known that induced DSBs increase both local [24] and global [52] chromosome mobility (reviewed in
[53]). This DSB-induced mobility was shown to be dependent on Cep3 phosphorylation in one
study [45], but not another [50]. We have previously published that the $(CAG)_{130}$ tract, in contrast to a
DSB, does not increase either local or global chromosome mobility, and instead causes
decreased locus mobility relative to chromosomes lacking the repeat tract [2]. However, those data
were collected from unsynchronized cells in S phase, and the locus likely becomes constrained
only after anchoring to the NPC. We reasoned that an increase in chromosome mobility due to a
collapsed fork at the $(CAG)_{130}$ tract and subsequent Cep3-S575 phosphorylation could be
transient. The anchoring at the NPC occurs in a distinct time window in late S phase, between
about 55-75 minutes after release from G1 and peaking at ~60 minutes (Figure 2D and [11]),
therefore any chromosome VI movement could be confined to a short time period preceding
NPC anchoring. The radius of constraint, a measurement of locus mobility [54], was measured for
the $(CAG)_{130}$ locus in cells that were synchronously released from G1 phase over a 100 minute
time course segmented into 5 minute intervals. During each 5 minute interval, images were
acquired every 1.5 seconds, closely mirroring the conditions used in [45], and MSD curves for each
5 minute window were plotted to obtain a radius of constraint. An example of the MSD curve for
the first 100 seconds of the 60-minute timepoint is depicted (Figure 3B). For each timepoint, the
radius of constraint was calculated based on the plateau reached within $\Delta t = 100s$ (Figure 3C;
Table S3). No difference in mobility of chromosome VI adjacent to the $(CAG)_{130}$ tract was
observed between wild-type and $cep3^{S575A}$ cells (Figure 3C). Though a fast directed motion may
not have been detected by this method, this result argues against a local increase in chromosome
movement triggered by Cep3 phosphorylation in response to a collapsed fork.
Another driver of DSB mobility is local nucleosome depletion [50]. Loss of Nhp6, encoded
in two loci Nhp6a and Nhp6b, has been shown to reduce histone occupancy on DNA [55], and this
leads to increased chromosome mobility [50,56].To test the mobility hypothesis in a different way,
we reasoned that if increasing chromosome mobility by phosphorylating $Cep3^{S575}$ was a
requirement for $(CAG)_{130}$ relocation, then causing global nucleosome depletion by deletion of
Nhp6a and Nhp6b could rescue the $cep3^{S575A}$ relocation defect. Deletion of Nhp6a and Nhp6b
$(nhp6\Delta\Delta)$ did not cause a decrease in relocation, and when combined with $cep3^{S575A}$ did not
rescue the relocation defect of those cells (Figure 3D). Altogether, we conclude that an increase
in chromosome mobility does not explain the importance of Cep3 phosphorylation for relocation
of the $(CAG)_{130}$ tract to the NP.
Centromeres unattached to the spindle can activate the spindle assembly checkpoint
(SAC) via Mad2 [57], and phosphorylation of Cep3 has been shown to activate the SAC [45].
Therefore, we tested whether activation of the SAC by Mad2 is required for relocation, however
there was no relocation defect in $mad2\Delta$ cells (Figure 3D). We conclude that the role of Cep3
phosphorylation in collapsed fork repositioning to the NP is not due to a defect in SAC
activation.
316
Expanded CAG repeat loci associate with DIMs in a Dun1 and Cep3-S575
phosphorylation-dependent manner.

In yeast, microtubule structures termed "Damage-Inducible Microtubules" or "DIMs"
form in response to double stranded breaks in a Rad9 checkpoint-dependent manner [25].
Interestingly, inactivating a centromere is sufficient to drive formation of DNA damage-
inducible microtubules (DIMs) [25]. Since our data supported that centromere inactivation and
Cep3 phosphorylation were important to allow relocation of the CAG locus to the NP, but not
through increasing chromosome mobility, we wondered whether DIM formation might be
involved. We reasoned that if DIMs were important for relocation to the periphery, then the
formation of microtubules in general would be important. To test this, cells were treated for 30
minutes with either 15 μg/mL of the tubulin depolymerizing drug nocodazole or an equivalent
amount of the vehicle (DMSO) and the zoning assay was performed. Addition of nocodazole
significantly reduced the percentage of the $(CAG)_{130}$ tract localized to the nuclear periphery in
mid-to-late S phase cells (Figure 4A). It is known that the minus end-directed microtubule motor
Kar3 promotes, though is not strictly required, for DIM formation [25]. Therefore, we tested
whether deletion of Kar3 had an effect on the frequency of $(CAG)_{130}$ locus association with the
NP. In $kar3\Delta$ cells in S phase, relocation of the repeat to zone 1 was reduced (though not
eliminated; Figure 4B), in line with the reported $kar3\Delta$ effect on DIMs.
To determine if DIMs were being recruited to the CAG repeat locus, we integrated an
additional copy of the tubulin subunit Tub1 tagged with the YFP variant mVenus at the *TUB1*
genomic locus [58]. GFP-LacI was replaced with CFP-LacI and GFP-Nup49 with Nup49-mCherry
to allow simultaneous visualization of all 3 nuclear components (Figure 4C). Consistent with
previous data [25], treatment with 0.03% MMS caused DIM formation in 62% of S-phase cells
(Figure 4D). To test whether DIMs were also induced by fork collapse, they were measured in
cells treated with 0.03%MMS + 0.2M Hydroxyurea for 60 minutes, which causes genome-wide
fork collapse and relocation of marked chromosome loci to the nuclear periphery [1,11]. Under fork
collapsing conditions, 87% of mid-to-late S phase cells contained a DIM (Figure 4D). We then
tested whether a single expanded $(CAG)_{130}$ repeat locus was sufficient to cause DIM formation
in cells. We observed that there is an increase in prevalence of DIM formation in cells with an
expanded CAG repeat to 36% compared to 18% for the no-tract control (Figure 4E). The
majority of these DIMs (74%) colocalized with the repeat locus (Figure 4E; see Figures 4C and
S2 examples). DIM formation was dependent upon Dun1 and Cep3 phosphorylation as deletion
of Dun1 or mutation to $cep3^{S575A}$ caused a significant reduction of DIMs (Figure 4E). The repeat
tract-induced DIMs were specifically affected in these mutants and were reduced to 7% in the
$cep3^{S575A}$ strain (Figure 4E). We conclude that there is a checkpoint-dependent induction of
DIMs and recruitment of a DIM to the site of fork collapse. These DIMs appear to be required to
mediate relocation of the collapsed fork to the NPC as deletion of *KAR3* or treatment with
nocodazole, both of which perturb DIM formation, led to a decrease in the frequency of
relocation of the CAG tract to the nuclear periphery.
356
<u>Discussion</u>
In this study we aimed to determine if the DNA damage checkpoint response plays a role
in the relocation of collapsed forks to the nuclear pore complex (NPC). We previously showed
that long CAG/CTG repeat tracts activate the DNA damage response and that the fork protection
complex containing Mrc1 is crucial for preventing CAG fragility and instability and promoting
fork progression through the repeat [14,15,17]. Other systems showed varying checkpoint
requirements for relocation of chromosomes with persistent DSBs, aphidicolin stalled forks or
eroded telomeres [1,3,22,23]. It was unknown what, if any, role the checkpoint played in the

relocation of stalled/collapsed forks caused by expanded CAG repeats to the nuclear periphery.
Even for types of damage where a requirement for checkpoint proteins had been established, the
mechanism by which the checkpoint pathway mediates repositioning of damaged chromosomal
loci had not been elucidated.
369   Our results indicate that the Mrc1/Rad53 axis is required for the repositioning of CAG
repeats to the NPC in mid-to-late S phase. The $(CAG)_{130}$ repeat forms hairpin structures that
interfere with DNA replication [59,60] and CAG repeats have similar relocation requirements as
cells which have been treated with both HU and MMS which causes fork collapse [1,2,11], thus it
was speculated that the relocation event was a consequence of a collapsed fork [2,11]. Since the
phosphorylation of Mrc1 is required for the efficient sensing and stabilizing of stalled or
uncoupled forks and the checkpoint activity of Mrc1 is critical for the relocation of the expanded
repeat to the nuclear periphery, this implies that a stalled or uncoupled fork is the initial lesion
being sensed and then processed for relocation to the NPC.
378   We found that there is redundancy in the sensing of the lesion caused by the $(CAG)_{130}$
tract, suggesting that by-products of fork uncoupling such as ssDNA or an exposed DSB end
caused by fork reversal or breakage also come into play. It was revealing that both the Mec1 and
Rad9 pathways needed to be inactivated in order to abolish relocation to the NP. Long-range
resection and RPA SUMOylation are required for CAG repeat relocation to the NPC [11], thus we
would expect RPA-coated ssDNA, which recruits Mec1, to be available prior to relocation.
Rad9, which is activated by both Mec1 and Tel1, was required only when Mec1 and/or the 9-1-1
complex are compromised, indirectly suggesting a role for Tel1. Tel1 has been shown to bind to
the MRX complex [61], which binds DSB ends and is required for relocation [11]. These results
imply that either ssDNA or a DSB end can activate the pathway needed for repositioning,
suggesting that there could be a transition from the initial uncoupled fork to a reversed or broken
fork that provides a DSB end (Figure 1E). This redundancy in signaling leads to a robust
mechanism to activate the Rad53 checkpoint in response to the initial fork stall caused by an
expanded CAG repeat.
392   Rad53 is the master regulator of the checkpoint in *S. cerevisiae* and has multiple roles in
the response to stalled or collapsed replication forks [26,62–64]. One well-studied target of Rad53 is
the kinase Dun1 [37], which we determined was needed for relocation (Figure 2B). It is important
to note that while Dun1 is a required target of Rad53 for relocation, it may not be the only target
as the kinase-deficient $rad53^{K227A}$ mutant is proficient for Dun1 activation [65], yet does not
relocate. Dun1 phosphorylates several downstream targets including the recombination factor
Rad55 [37], RNR inhibition factors (Crt1, Sml1, and Dif1 reviewed in [38]), and the kinetochore
protein Cep3 [45]. Of these targets, the phosphorylation of Cep3 was required for relocation
(Figure 2C).
401   The yeast centromere is only ~120bp in length and has three sub-sections: CDEI, CDEII,
and CDEIII. Kinetochore proteins at each centromere bind to a single microtubule, which is
attached to the spindle pole body (SPB) throughout most of the cell cycle [66], with a transient
release during centromere replication [67]. Cep3 has a zinc finger DNA binding domain that
interacts with the CDEIII region as a member of the CBF3 complex (reviewed in [68]).
Phosphorylation of the kinetochore has been widely studied and has varying functions (reviewed
in [69,70]. For example, phosphorylation of the kinetochore proteins, such as Dam1, by Aurora B
kinase (Ipl1) destabilizes erroneous microtubule attachments in mitosis [71,72]. However, the
consequences of Cep3 phosphorylation have been comparatively poorly described. It was
previously suggested that Cep3 phosphorylation led to a release of centromere constraint to

allow the increased global mobility of chromosomes observed in response to an HO-induced
DSB at the *MAT* locus in the G2/M phase of the cell cycle [45]. However, another group was
unable to show a Cep3-dependent increased radius of constraint [50] using similar conditions. Our
results indicate that Cep3 phosphorylation may act through another mechanism besides
increasing local or global chromatin mobility, at least in response to replication fork collapse.
We propose that the DNA damage dependent phosphorylation of Cep3 is causing centromere
release from the SPB attachment normally present during S phase. Consistent with this model,
the requirement for Cep3 phosphorylation can be obviated by centromere inactivation (Figure
3A).
DIMs form in response to DNA damage induced by MMS, zeocin, camptothecin, or I-
SceI-induced DSBs [25]. With this work, we've now shown that while DIM formation is common
after treatment with MMS, which stalls forks, it is dramatically increased after treatment with
HU + MMS, which has been shown to be fork collapsing. DIM formation even occurs as a result
of a single fork collapse caused by an expanded CAG repeat tract. Interestingly, much like how
CEN inactivation eliminates the need for Cep3 phosphorylation for relocation, CEN inactivation
can cause spontaneous formation of DIMs [25]. This suggests that while the initial damage
(whether a DSB or uncoupled/reversed/broken fork) is responsible for centromere release from
the SPB microtubule through Cep3 phosphorylation, the DIM may form as a result of that release
rather than directly responding to the damage. Thus, we hypothesize that release of the
centromere from the end of the SPB microtubule triggers the formation of a DIM.
The microtubule motor protein Kar3 has been shown to promote the capture and
directional movement of damaged loci by DIMs [25]. When centromeres are inactivated,
kinetochores that were previously attached to the microtubule plus end reposition and bind the
microtubule side-on [73,74]. Kar3, as a minus end directed motor, is involved in repositioning the
centromere on the microtubule, by hindering tethering of the kinetochore to the plus end of the
microtubule [73–75]. During this kinetochore release and repositioning process, the damaged repeat
locus could be freed up to move to the NPC. If this is the case, then we might expect that the
presence of Kar3 could increase the duration of this "relocation permissive" state by
antagonizing reattachment of the kinetochore to the microtubule plus end, which could explain
why a *kar3Δ* reduces but does not completely abrogate relocation to the NPC. What remains
unclear is whether the microtubule interacts directly with the collapsed fork, or if the DIM allows
for repositioning of the lesion in an indirect manner, for example by allowing either free
movement of the damaged locus or for the centromere to slide along the DIM. The former is an
attractive hypothesis due to the increased colocalization of the repeat locus with the DIM (Figure
4E). It has recently been shown that DIMs serve as sites of repair center nucleation, with liquid
droplets of Rad52 colocalizing with DIMs [76], and we've previously shown that many repair
proteins, including Rad52 [2] and RPA [11] are recruited to the site of the collapsed fork prior to
relocation. These factors have been shown to become SUMOylated, and their SUMOylation has
been shown to be critically required for relocation to the NPC through an interaction with Slx5/8
[11]. Slx5/8 in turn is critical for anchoring of damaged DNA at the NPC (reviewed in [8]).
SUMOylation has also been shown to promote condensation of repair factors [77]. Possibly,
SUMOylation or factors within SUMOylation-induced repair centers could play a role in the
association between the collapsed fork at the CAG tract and the DIM.
Thus, we propose that after fork collapse at the CAG repeat, the DNA replication
checkpoint results in the phosphorylation of Cep3. This phosphorylation causes centromere
release from the SPB, which promotes DIM formation. The DIM allows movement of the

collapsed fork locus to the periphery in a manner yet to be fully elucidated, but which may
involve interaction of the damaged locus with the DIM. Once at the periphery, interactions
between SUMOylated repair factors, Slx5/8, and the nucleoporins tether the locus to the NPC for
repair or fork restart (Figure 4F).
There is prior evidence in other types of eukaryotic cells for a physical mechanism by
which movement of lesions to repair centers within the nucleus occurs. Heterochromatic DSBs
in *Drosophila* require nuclear F-actin and myosin for directed movement to the nuclear periphery
[78]. In human U2OS cells the relocation of collapsed forks caused by aphidicolin is dependent on
the polymerization of nuclear F-actin, and late-replication foci were observed to move along
actin filaments to the nuclear periphery [3]. Further it's been shown that recruitment of nuclear
actin in response to damage can control pathway choice in response to replication stress [79]. It is
unclear if yeast have equivalent nuclear actin filaments or if directed movement to the NP would
be detectable in the smaller yeast nucleus. However, in yeast, the kinesin-14 motor protein
complex is required for movement of telomeric DSBs from their usual attachment site on the
nuclear membrane to NPCs and these were observed to move along DIMs [25,80]. Our results
suggest that collapsed forks may also utilize DIMs to facilitate their movement to the yeast NPC.
In human cells, it was recently reported that cytoplasmic microtubules that form in response to
DSBs push on the nuclear envelope to form nuclear envelope tubules (NETs) that elongate into
the nucleus and bring the nuclear envelope into the proximity of DSBs within the nuclear
interior, and the DSBs may also move toward the dsbNETs [81]. The formation of these dsbNETs
serves to mobilize repair proteins to sites of DSBs and is dependent on the ATM/ATR/DNA-PK
checkpoint response [81]. Thus, the involvement of microtubules in mobilizing a repair response
seems to be conserved, though the details of where in the nucleus the damage-inducible
microtubules form and how they interact with the damaged locus may differ between organisms.
It will be interesting to elucidate if DIMs are utilized in multicellular eukaryotic cells or if their
function has been replaced by a combination of nuclear actin filaments and dsbNETs in these
larger nuclei.
In summary, our results illuminate a new role for the DNA replication and DNA damage
checkpoint response, which is to phosphorylate a kinetochore component to allow for
repositioning of a collapsed fork within the nucleus. Since this repositioning is important for
repair or fork restart, we identify it as a crucial new role for the DNA damage checkpoint
response in *S. cerevisiae*. Repositioning of damaged loci to a repair center is present across many
organisms. Where it has been studied, the checkpoint is critical to facilitate this relocation, but
the targets involved have not been elucidated. Given the importance of recovery from replication
fork collapse, a mechanism to allow centromere release to facilitate damaged locus repositioning
might be especially critical at regions with fork-stalling DNA structures and for cancerous or
aging cells experiencing high levels of replication stress.
494

495 **Acknowledgements**
496 We would like to thank Stéphane Marcand for the gift of the *pGAL1*-CEN6 plasmid and Wei-Lih
497 Lee for the *p*HIS3 *p*:Venus-Tub1+3'UTR::LEU2 plasmid.
498

499 **Funding**
500 This work was funded by the National Institute of General Medical Sciences of the National
501 Institutes of Health under Award Number R35 GM144215 to CHF.
502

Author Contributions
JW: Conceptualization, Investigation, Analysis, Supervision, Writing
TM: Conceptualization, Investigation, Analysis, Software, Writing
MJ: Investigation, Analysis
CHF: Supervision, Resources, Writing, Administration, Conception

References

1. Nagai, S., Dubrana, K., Tsai-Pflugfelder, M., Davidson, M.B., Roberts, T.M., Brown, G.W., Varela, E., Hediger, F., Gasser, S.M., and Krogan, N.J. (2008). Functional Targeting of DNA Damage to a Nuclear Pore–Associated SUMO-Dependent Ubiquitin Ligase. Science *322*, 597–602. https://doi.org/10.1126/science.1162790.

2. Su, X.A., Dion, V., Gasser, S.M., and Freudenreich, C.H. (2015). Regulation of recombination at yeast nuclear pores controls repair and triplet repeat stability. Genes Dev. *29*, 1006–1017. https://doi.org/10.1101/gad.256404.114.

3. Lamm, N., Read, M.N., Nobis, M., Van Ly, D., Page, S.G., Masamsetti, V.P., Timpson, P., Biro, M., and Cesare, A.J. (2020). Nuclear F-actin counteracts nuclear deformation and promotes fork repair during replication stress. Nat. Cell Biol. *22*, 1460–1470. https://doi.org/10.1038/s41556-020-00605-6.

4. Horigome, C., Unozawa, E., Ooki, T., and Kobayashi, T. (2019). Ribosomal RNA gene repeats associate with the nuclear pore complex for maintenance after DNA damage. PLOS Genet. *15*, e1008103. https://doi.org/10.1371/journal.pgen.1008103.

5. Kramarz, K., Schirmeisen, K., Boucherit, V., Ait Saada, A., Lovo, C., Palancade, B., Freudenreich, C., and Lambert, S.A.E. (2020). The nuclear pore primes recombination-dependent DNA synthesis at arrested forks by promoting SUMO removal. Nat. Commun. *11*, 5643. https://doi.org/10.1038/s41467-020-19516-z.

6. Aguilera, P., Whalen, J., Minguet, C., Churikov, D., Freudenreich, C., Simon, M.-N., and Géli, V. (2020). The nuclear pore complex prevents sister chromatid recombination during replicative senescence. Nat. Commun. *11*, 160. https://doi.org/10.1038/s41467-019-13979-5.

7. Pinzaru, A.M., Kareh, M., Lamm, N., Lazzerini-Denchi, E., Cesare, A.J., and Sfeir, A. (2020). Replication stress conferred by POT1 dysfunction promotes telomere relocalization to the nuclear pore. Genes Dev. *34*, 1619–1636. https://doi.org/10.1101/gad.337287.120.

8. Whalen, J.M., and Freudenreich, C.H. (2020). Location, Location, Location: The Role of Nuclear Positioning in the Repair of Collapsed Forks and Protection of Genome Stability. Genes *11*, 635. https://doi.org/10.3390/genes11060635.

9. Lamm, N., Rogers, S., and Cesare, A.J. (2021). Chromatin mobility and relocation in DNA repair. Trends Cell Biol. *31*, 843–855. https://doi.org/10.1016/j.tcb.2021.06.002.

10. Gasser, S.M., and Stutz, F. (2023). SUMO in the regulation of DNA repair and transcription at nuclear pores. FEBS Lett. *597*, 2833–2850. https://doi.org/10.1002/1873-3468.14751.

11. Whalen, J.M., Dhingra, N., Wei, L., Zhao, X., and Freudenreich, C.H. (2020). Relocation of
Collapsed Forks to the Nuclear Pore Complex Depends on Sumoylation of DNA Repair
Proteins and Permits Rad51 Association. Cell Rep. *31*, 107635–107635.
https://doi.org/10.1016/j.celrep.2020.107635.

12. Usdin, K., House, N.C.M., and Freudenreich, C.H. (2015). Repeat instability during DNA
repair: Insights from model systems. Crit. Rev. Biochem. Mol. Biol. *50*, 142–167.
https://doi.org/10.3109/10409238.2014.999192.

13. Khristich, A.N., and Mirkin, S.M. (2020). On the wrong DNA track: Molecular mechanisms
of repeat-mediated genome instability. J. Biol. Chem. *295*, 4134–4170.
https://doi.org/10.1074/jbc.REV119.007678.

14. Sundararajan, R., and Freudenreich, C.H. (2011). Expanded CAG/CTG Repeat DNA
Induces a Checkpoint Response That Impacts Cell Proliferation in Saccharomyces
cerevisiae. PLoS Genet. *7*, e1001339. https://doi.org/10.1371/journal.pgen.1001339.

15. Lahiri, M., Gustafson, T.L., Majors, E.R., and Freudenreich, C.H. (2004). Expanded CAG
Repeats Activate the DNA Damage Checkpoint Pathway. Mol. Cell *15*, 287–293.
https://doi.org/10.1016/j.molcel.2004.06.034.

16. Freudenreich, C.H., and Lahiri, M. (2004). Structure-Forming CAG/CTG Repeat Sequences
are Sensitive to Breakage in the Absence of Mrc1 Checkpoint Function and S-Phase
Checkpoint Signaling: Implications for Trinucleotide Repeat Expansion Diseases. Cell Cycle
3, 1370–1374. https://doi.org/10.4161/cc.3.11.1246.

17. Gellon, L., Kaushal, S., Cebrián, J., Lahiri, M., Mirkin, S.M., and Freudenreich, C.H. (2019).
Mrc1 and Tof1 prevent fragility and instability at long CAG repeats by their fork stabilizing
function. Nucleic Acids Res. *47*, 794–805. https://doi.org/10.1093/nar/gky1195.

18. Hustedt, N., Gasser, S.M., and Shimada, K. (2013). Replication Checkpoint: Tuning and
Coordination of Replication Forks in S Phase. Genes *4*, 388–434.
https://doi.org/10.3390/genes4030388.

19. McClure, A.W., Canal, B., and Diffley, J.F.X. (2022). A DNA replication fork-centric view
of the budding yeast DNA damage response. DNA Repair *119*, 103393.
https://doi.org/10.1016/j.dnarep.2022.103393.

20. Katou, Y., Kanoh, Y., Bando, M., Noguchi, H., Tanaka, H., Ashikari, T., Sugimoto, K., and
Shirahige, K. (2003). S-phase checkpoint proteins Tof1 and Mrc1 form a stable replication-
pausing complex. Nature *424*, 1078–1083. https://doi.org/10.1038/nature01900.

21. Lou, H., Komata, M., Katou, Y., Guan, Z., Reis, C.C., Budd, M., Shirahige, K., and
Campbell, J.L. (2008). Mrc1 and DNA Polymerase □ Function Together in Linking DNA
Replication and the S Phase Checkpoint. Mol. Cell *32*, 106–117.
https://doi.org/10.1016/j.molcel.2008.08.020.

22. Khadaroo, B., Teixeira, M.T., Luciano, P., Eckert-Boulet, N., Germann, S.M., Simon, M.N.,
Gallina, I., Abdallah, P., Gilson, E., Géli, V., et al. (2009). The DNA damage response at
eroded telomeres and tethering to the nuclear pore complex. Nat. Cell Biol. *11*, 980–987.
https://doi.org/10.1038/ncb1910.

23. Chiolo, I., Minoda, A., Colmenares, S.U., Polyzos, A., Costes, S.V., and Karpen, G.H.
(2011). Double-Strand Breaks in Heterochromatin Move Outside of a Dynamic HP1a
Domain to Complete Recombinational Repair. Cell *144*, 732–744.
https://doi.org/10.1016/j.cell.2011.02.012.

24. Dion, V., Kalck, V., Horigome, C., Towbin, B.D., and Gasser, S.M. (2012). Increased
mobility of double-strand breaks requires Mec1, Rad9 and the homologous recombination
machinery. Nat. Cell Biol. *14*, 502–509. https://doi.org/10.1038/ncb2465.

25. Oshidari, R., Strecker, J., Chung, D.K.C., Abraham, K.J., Chan, J.N.Y., Damaren, C.J., and
Mekhail, K. (2018). Nuclear microtubule filaments mediate non-linear directional motion of
chromatin and promote DNA repair. Nat. Commun. *9*, 2567. https://doi.org/10.1038/s41467-
018-05009-7.

26. Zhao, X., Muller, E.G.D., and Rothstein, R. (1998). A Suppressor of Two Essential
Checkpoint Genes Identifies a Novel Protein that Negatively Affects dNTP Pools. Mol. Cell
2, 329–340. https://doi.org/10.1016/S1097-2765(00)80277-4.

27. Fay, D.S., Sun, Z., and Stern, D.F. (1997). Mutations in SPK1/RAD53 that specifically
abolish checkpoint but not growth-related functions. Curr. Genet. *31*, 97–105.
https://doi.org/10.1007/s002940050181.

28. Naylor, M.L., Li, J., Osborn, A.J., and Elledge, S.J. (2009). Mrc1 phosphorylation in
response to DNA replication stress is required for Mec1 accumulation at the stalled fork.
Proc. Natl. Acad. Sci. U. S. A. *106*, 12765–12770. https://doi.org/10.1073/pnas.0904623106.

29. Osborn, A.J., and Elledge, S.J. (2003). Mrc1 is a replication fork component whose
phosphorylation in response to DNA replication stress activates Rad53. Genes Dev. *17*,
1755–1767. https://doi.org/10.1101/gad.1098303.

30. Emili, A. (1998). MEC1-Dependent Phosphorylation of Rad9p in Response to DNA
Damage. Mol. Cell *2*, 183–189. https://doi.org/10.1016/S1097-2765(00)80128-8.

31. Gilbert, C.S., Green, C.M., and Lowndes, N.F. (2001). Budding Yeast Rad9 Is an ATP-
Dependent Rad53 Activating Machine. Mol. Cell *8*, 129–136. https://doi.org/10.1016/S1097-
2765(01)00267-2.

32. House, N.C.M., Yang, J.H., Walsh, S.C., Moy, J.M., and Freudenreich, C.H. (2014). NuA4
Initiates Dynamic Histone H4 Acetylation to Promote High-Fidelity Sister Chromatid
Recombination at Postreplication Gaps. Mol. Cell *55*, 818–828.
https://doi.org/10.1016/j.molcel.2014.07.007.

33. House, N.C., Polleys, E.J., Quasem, I., De la Rosa Mejia, M., Joyce, C.E., Takacsi-Nagy, O., Krebs, J.E., Fuchs, S.M., and Freudenreich, C.H. (2019). Distinct roles for S. cerevisiae H2A copies in recombination and repeat stability, with a role for H2A.1 threonine 126. eLife *8*, e53362. https://doi.org/10.7554/eLife.53362.

34. Downs, J.A., Lowndes, N.F., and Jackson, S.P. (2000). A role for Saccharomyces cerevisiae histone H2A in DNA repair. Nature *408*, 1001–1004. https://doi.org/10.1038/35050000.

35. Shroff, R., Arbel-Eden, A., Pilch, D., Ira, G., Bonner, W.M., Petrini, J.H., Haber, J.E., and Lichten, M. (2004). Distribution and Dynamics of Chromatin Modification Induced by a Defined DNA Double-Strand Break. Curr. Biol. CB *14*, 1703–1711. https://doi.org/10.1016/j.cub.2004.09.047.

36. Alcasabas, A.A., Osborn, A.J., Bachant, J., Hu, F., Werler, P.J.H., Bousset, K., Furuya, K., Diffley, J.F.X., Carr, A.M., and Elledge, S.J. (2001). Mrc1 transduces signals of DNA replication stress to activate Rad53. Nat. Cell Biol. *3*, 958–965. https://doi.org/10.1038/ncb1101-958.

37. Bashkirov, V.I., Bashkirova, E.V., Haghnazari, E., and Heyer, W.-D. (2003). Direct Kinase-to-Kinase Signaling Mediated by the FHA Phosphoprotein Recognition Domain of the Dun1 DNA Damage Checkpoint Kinase. Mol. Cell. Biol. *23*, 1441–1452. https://doi.org/10.1128/MCB.23.4.1441-1452.2003.

38. Nordlund, P., and Reichard, P. (2006). Ribonucleotide Reductases. Annu. Rev. Biochem. *75*, 681–706. https://doi.org/10.1146/annurev.biochem.75.103004.142443.

39. de la Torre Ruiz, M.-A., and Lowndes, N.F. (2000). DUN1 defines one branch downstream of RAD53 for transcription and DNA damage repair in Saccharomyces cerevisiae. FEBS Lett. *485*, 205–206. https://doi.org/10.1016/S0014-5793(00)02198-0.

40. Zhou, Z., and Elledge, S.J. (1993). DUN1 encodes a protein kinase that controls the DNA damage response in yeast. Cell *75*, 1119–1127. https://doi.org/10.1016/0092-8674(93)90321-G.

41. Fortin, G.S., and Symington, L.S. (2002). Mutations in yeast Rad51 that partially bypass the requirement for Rad55 and Rad57 in DNA repair by increasing the stability of Rad51–DNA complexes. EMBO J. *21*, 3160–3170. https://doi.org/10.1093/emboj/cdf293.

42. Huang, M., Zhou, Z., and Elledge, S.J. (1998). The DNA Replication and Damage Checkpoint Pathways Induce Transcription by Inhibition of the Crt1 Repressor. Cell *94*, 595–605. https://doi.org/10.1016/S0092-8674(00)81601-3.

43. Wu, X., and Huang, M. (2008). Dif1 Controls Subcellular Localization of Ribonucleotide Reductase by Mediating Nuclear Import of the R2 Subunit. Mol. Cell. Biol. *28*, 7156–7167. https://doi.org/10.1128/MCB.01388-08.

44. Zhao, X., and Rothstein, R. (2002). The Dun1 checkpoint kinase phosphorylates and
regulates the ribonucleotide reductase inhibitor Sml1. Proc. Natl. Acad. Sci. *99*, 3746–3751.
https://doi.org/10.1073/pnas.062502299.

45. Strecker, J., Gupta, G.D., Zhang, W., Bashkurov, M., Landry, M.-C., Pelletier, L., and
Durocher, D. (2016). DNA damage signalling targets the kinetochore to promote chromatin
mobility. Nat. Cell Biol. *18*, 281–290. https://doi.org/10.1038/ncb3308.

46. Lechner, J., and Carbon, J. (1991). A 240 kd multisubunit protein complex, CBF3, is a major
component of the budding yeast centromere. Cell *64*, 717–725. https://doi.org/10.1016/0092-
8674(91)90501-o.

47. Lechner, J. (1994). A zinc finger protein, essential for chromosome segregation, constitutes a
putative DNA binding subunit of the Saccharomyces cerevisiae kinetochore complex, Cbf3.
EMBO J. *13*, 5203–5211. https://doi.org/10.1002/j.1460-2075.1994.tb06851.x.

48. Collins, K.A., Castillo, A.R., Tatsutani, S.Y., and Biggins, S. (2005). De Novo Kinetochore
Assembly Requires the Centromeric Histone H3 Variant. Mol. Biol. Cell *16*, 5649–5660.
https://doi.org/10.1091/mbc.E05-08-0771.

49. Hill, A., and Bloom, K. (1987). Genetic manipulation of centromere function. Mol. Cell.
Biol. *7*, 2397–2405.

50. Cheblal, A., Challa, K., Seeber, A., Shimada, K., Yoshida, H., Ferreira, H.C., Amitai, A., and
Gasser, S.M. (2020). DNA Damage-Induced Nucleosome Depletion Enhances Homology
Search Independently of Local Break Movement. Mol. Cell *80*, 311-326.e4.
https://doi.org/10.1016/j.molcel.2020.09.002.

51. Pobiega, S., and Marcand, S. (2010). Dicentric breakage at telomere fusions. Genes Dev. *24*,
720–733. https://doi.org/10.1101/gad.571510.

52. Miné-Hattab, J., and Rothstein, R. (2012). Increased chromosome mobility facilitates
homology search during recombination. Nat. Cell Biol. *14*, 510–517.
https://doi.org/10.1038/ncb2472.

53. García Fernández, F., and Fabre, E. (2022). The Dynamic Behavior of Chromatin in
Response to DNA Double-Strand Breaks. Genes *13*, 215.
https://doi.org/10.3390/genes13020215.

54. Dion, V., and Gasser, S.M. (2013). Chromatin Movement in the Maintenance of Genome
Stability. Cell *152*, 1355–1364. https://doi.org/10.1016/j.cell.2013.02.010.

55. Celona, B., Weiner, A., Felice, F.D., Mancuso, F.M., Cesarini, E., Rossi, R.L., Gregory, L.,
Baban, D., Rossetti, G., Grianti, P., et al. (2011). Substantial Histone Reduction Modulates
Genomewide Nucleosomal Occupancy and Global Transcriptional Output. PLOS Biol. *9*,
e1001086. https://doi.org/10.1371/journal.pbio.1001086.

56. Hauer, M.H., Seeber, A., Singh, V., Thierry, R., Sack, R., Amitai, A., Kryzhanovska, M.,
Eglinger, J., Holcman, D., Owen-Hughes, T., et al. (2017). Histone degradation in response
to DNA damage enhances chromatin dynamics and recombination rates. Nat. Struct. Mol.
Biol. *24*, 99–107. https://doi.org/10.1038/nsmb.3347.

57. Hoyt, M.A., Totis, L., and Roberts, B.T. (1991). S. cerevisiae genes required for cell cycle
arrest in response to loss of microtubule function. Cell *66*, 507–517.
https://doi.org/10.1016/0092-8674(81)90014-3.

58. Markus, S.M., Omer, S., Baranowski, K., and Lee, W.-L. (2015). Improved Plasmids for
Fluorescent Protein Tagging of Microtubules in Saccharomyces cerevisiae. Traffic Cph.
Den. *16*, 773–786. https://doi.org/10.1111/tra.12276.

59. Pelletier, R., Krasilnikova, M.M., Samadashwily, G.M., Lahue, R., and Mirkin, S.M. (2003).
Replication and Expansion of Trinucleotide Repeats in Yeast. Mol. Cell. Biol. *23*, 1349–
1357. https://doi.org/10.1128/MCB.23.4.1349-1357.2003.

60. Viterbo, D., Michoud, G., Mosbach, V., Dujon, B., and Richard, G.-F. (2016). Replication
stalling and heteroduplex formation within CAG/CTG trinucleotide repeats by mismatch
repair. DNA Repair *42*, 94–106. https://doi.org/10.1016/j.dnarep.2016.03.002.

61. Cassani, C., Gobbini, E., Wang, W., Niu, H., Clerici, M., Sung, P., and Longhese, M.P.
(2016). Tel1 and Rif2 Regulate MRX Functions in End-Tethering and Repair of DNA
Double-Strand Breaks. PLOS Biol. *14*, e1002387.
https://doi.org/10.1371/journal.pbio.1002387.

62. Santocanale, C., and Diffley, J.F.X. (1998). A Mec1- and Rad53-dependent checkpoint
controls late-firing origins of DNA replication. Nature *395*, 615–618.
https://doi.org/10.1038/27001.

63. Herzberg, K., Bashkirov, V.I., Rolfsmeier, M., Haghnazari, E., McDonald, W.H., Anderson,
S., Bashkirova, E.V., Yates, J.R., and Heyer, W.-D. (2006). Phosphorylation of Rad55 on
Serines 2, 8, and 14 Is Required for Efficient Homologous Recombination in the Recovery of
Stalled Replication Forks. Mol. Cell. Biol. *26*, 8396–8409.
https://doi.org/10.1128/MCB.01317-06.

64. McClure, A.W., and Diffley, J.F. (2021). Rad53 checkpoint kinase regulation of DNA
replication fork rate via Mrc1 phosphorylation. eLife *10*, e69726.
https://doi.org/10.7554/eLife.69726.

65. Hoch, N.C., Chen, E.S.-W., Buckland, R., Wang, S.-C., Fazio, A., Hammet, A., Pellicioli,
A., Chabes, A., Tsai, M.-D., and Heierhorst, J. (2013). Molecular basis of the essential s
phase function of the rad53 checkpoint kinase. Mol. Cell. Biol. *33*, 3202–3213.
https://doi.org/10.1128/MCB.00474-13.

66. Heun, P., Laroche, T., Raghuraman, M.K., and Gasser, S.M. (2001). The Positioning and
Dynamics of Origins of Replication in the Budding Yeast Nucleus. J. Cell Biol. *152*, 385–
400.

67. Kitamura, E., Tanaka, K., Kitamura, Y., and Tanaka, T.U. (2007). Kinetochore–microtubule
interaction during S phase in Saccharomyces cerevisiae. Genes Dev. *21*, 3319–3330.
https://doi.org/10.1101/gad.449407.

68. Biggins, S. (2013). The Composition, Functions, and Regulation of the Budding Yeast
Kinetochore. Genetics *194*, 817–846. https://doi.org/10.1534/genetics.112.145276.

69. Funabiki, H., and Wynne, D.J. (2013). Making an effective switch at the kinetochore by
phosphorylation and dephosphorylation. Chromosoma *122*, 135–158.
https://doi.org/10.1007/s00412-013-0401-5.

70. Klemm, C., Thorpe, P.H., and Ólafsson, G. (2021). Cell-cycle phospho-regulation of the
kinetochore. Curr. Genet. *67*, 177–193. https://doi.org/10.1007/s00294-020-01127-2.

71. Tanaka, T.U., Rachidi, N., Janke, C., Pereira, G., Galova, M., Schiebel, E., Stark, M.J.R.,
and Nasmyth, K. (2002). Evidence that the Ipl1-Sli15 (Aurora kinase-INCENP) complex
promotes chromosome bi-orientation by altering kinetochore-spindle pole connections. Cell
108, 317–329. https://doi.org/10.1016/s0092-8674(02)00633-5.

72. Pinsky, B.A., Kung, C., Shokat, K.M., and Biggins, S. (2006). The Ipl1-Aurora protein
kinase activates the spindle checkpoint by creating unattached kinetochores. Nat. Cell Biol.
8, 78–83. https://doi.org/10.1038/ncb1341.

73. Tanaka, K., Kitamura, E., Kitamura, Y., and Tanaka, T.U. (2007). Molecular mechanisms of
microtubule-dependent kinetochore transport toward spindle poles. J. Cell Biol. *178*, 269–
281. https://doi.org/10.1083/jcb.200702141.

74. Tanaka, T.U., Stark, M.J.R., and Tanaka, K. (2005). Kinetochore capture and bi-orientation
on the mitotic spindle. Nat. Rev. Mol. Cell Biol. *6*, 929–942.
https://doi.org/10.1038/nrm1764.

75. Tytell, J.D., and Sorger, P.K. (2006). Analysis of kinesin motor function at budding yeast
kinetochores. J. Cell Biol. *172*, 861–874. https://doi.org/10.1083/jcb.200509101.

76. Oshidari, R., Huang, R., Medghalchi, M., Tse, E.Y.W., Ashgriz, N., Lee, H.O., Wyatt, H.,
and Mekhail, K. (2020). DNA repair by Rad52 liquid droplets. Nat. Commun. *11*, 695.
https://doi.org/10.1038/s41467-020-14546-z.

77. Keiten-Schmitz, J., Schunck, K., and Müller, S. (2020). SUMO Chains Rule on Chromatin
Occupancy. Front. Cell Dev. Biol. *7*. https://doi.org/10.3389/fcell.2019.00343.

78. Caridi, C.P., D'Agostino, C., Ryu, T., Zapotoczny, G., Delabaere, L., Li, X., Khodaverdian,
V.Y., Amaral, N., Lin, E., Rau, A.R., et al. (2018). Nuclear F-actin and myosins drive
relocalization of heterochromatic breaks. Nature *559*, 54–60. https://doi.org/10.1038/s41586-
018-0242-8.

79. Palumbieri, M.D., Merigliano, C., González-Acosta, D., Kuster, D., Krietsch, J., Stoy, H.,
von Känel, T., Ulferts, S., Welter, B., Frey, J., et al. (2023). Nuclear actin polymerization

rapidly mediates replication fork remodeling upon stress by limiting PrimPol activity. Nat.
Commun. *14*, 1–15. https://doi.org/10.1038/s41467-023-43183-5.

80. Chung, D.K.C., Chan, J.N.Y., Strecker, J., Zhang, W., Ebrahimi-Ardebili, S., Lu, T.,
Abraham, K.J., Durocher, D., and Mekhail, K. (2015). Perinuclear tethers license telomeric
DSBs for a broad kinesin- and NPC-dependent DNA repair process. Nat. Commun. *6*, 7742.
https://doi.org/10.1038/ncomms8742.

81. Shokrollahi, M., Stanic, M., Hundal, A., Chan, J.N.Y., Urman, D., Jordan, C.A., Hakem, A.,
Espin, R., Hao, J., Krishnan, R., et al. (2024). DNA double-strand break–capturing nuclear
envelope tubules drive DNA repair. Nat. Struct. Mol. Biol., 1–12.
https://doi.org/10.1038/s41594-024-01286-7.

82. Pike, B.L., Hammet, A., and Heierhorst, J. (2001). Role of the N-terminal Forkhead-
associated Domain in the Cell Cycle Checkpoint Function of the Rad53 Kinase*210. J. Biol.
Chem. *276*, 14019–14026. https://doi.org/10.1074/jbc.M009558200.

83. Generoso, W.C., Gottardi, M., Oreb, M., and Boles, E. (2016). Simplified CRISPR-Cas
genome editing for Saccharomyces cerevisiae. J. Microbiol. Methods *127*, 203–205.
https://doi.org/10.1016/j.mimet.2016.06.020.

84. Meister, P., Gehlen, L.R., Varela, E., Kalck, V., and Gasser, S.M. (2010). Chapter 21 -
Visualizing Yeast Chromosomes and Nuclear Architecture. In Methods in Enzymology
Guide to Yeast Genetics: Functional Genomics, Proteomics, and Other Systems Analysis.
(Academic Press), pp. 535–567. https://doi.org/10.1016/S0076-6879(10)70021-5.

85. Hediger, F., Neumann, F.R., Van Houwe, G., Dubrana, K., and Gasser, S.M. (2002). Live
Imaging of Telomeres: yKu and Sir Proteins Define Redundant Telomere-Anchoring
Pathways in Yeast. Curr. Biol. *12*, 2076–2089. https://doi.org/10.1016/S0960-
9822(02)01338-6.

<u>Figure Legends</u>

Figure 1: Relocation of (CAG)$_{130}$ locus to the nuclear periphery depends on the activation
of the DNA replication or DNA damage checkpoint. A) Flow chart of the DNA damage
checkpoint. B) Yeast chromosome VI containing integrated 130 CAG/CTG repeat tract
((CAG)$_{130}$) and a *lacO* or *tetO* array that binds the LacI-GFP or TetR-RFP protein, respectively.
Mid/late S phase cells were imaged, and the location of the foci were scored into one of three
zones of equal area, using the GFP-Nup49 signal to mark the nuclear periphery. Representative
images are shown. C) Percent of cells with LacO loci in zone 1 for mid/late S-phase cells. Cells
have either (CAG)$_0$ or (CAG)$_{130}$ as marked. D) Percent of zone 1 foci for mid/late S-phase cells
for indicated strains. All strains have (CAG)$_{130}$. E) Model illustrating activation of the DRC in
response to an uncoupled fork (left) or activation of the DDC following fork reversal and
recruitment of the MRX complex (right). (*) P≤0.05, (**) P≤0.01 (***) P≤0.001 compared with

wild-type $(CAG)_{130}$ strain by Fisher's exact test. More than 100 cells were analyzed per strain.
See Table S1 for the exact number of cells analyzed, percentages, and P-values.

798
Figure 2: Phosphorylation of Cep3^{S575} by Dun1 is needed for relocation of $(CAG)_{130}$ to the
nuclear periphery. A) Diagram of Dun1 targets. The colors used for mutants in this figure
correspond with colors used in this diagram. B) Percent of zone 1 foci for $(CAG)_{130}$ mid/late S-
phase cells for strains deleted for Dun1 or its targets. C) Percent of zone 1 foci for $(CAG)_{130}$
mid/late S-phase cells for a phosphomimetic mutant of Cep3 (*cep3^{S575E}*) individually and in the
dun1Δ mutant. D) Percentage of zone 1 foci for $(CAG)_{130}$ cells at indicated time points after
release from alpha factor arrest in wild-type and *cep3^{S575E}* strains. See Table S2 for the exact
number of cells analyzed, percentages, and P-values. (*) P≤0.05, (**) P≤0.01 (***) P≤0.001
compared with wild-type by Fisher's exact test. (^^^) P ≤ 0.001 compared as indicated by
Fisher's Exact Test. More than 100 cells were analyzed per strain. See Table S1 for the exact
number of cells analyzed, percentages, and P-values.

810
Figure 3: Cep3 phosphorylation allows for $(CAG)_{130}$ relocation through release from
centromeric constraint. A) Percent zone 1 foci for $(CAG)_{130}$ mid/late S-phase cells for
cep3^{S575A} and P_{GAL1}-*CEN6 cep3^{S575A}* strains. Cultures were grown in YP-lactate media then split
into media containing either glucose or galactose to induce transcription through CEN6. B) MSD
track for Cep3^{S575A} and WT strains containing $(CAG)_{130}$. Cells were imaged every 1.5 seconds
for 5 minutes, first Δt =[0,100] seconds shown. Data was captured 60 minutes post-release from
alpha factor arrest. The lines represent the 95% Confidence intervals of the mean square
displacement at a particular timepoint. C) Summary of radii of constraint calculated for WT and
cep3^{S575A} over 100 minutes post release from alpha factor arrest. Points represent calculated
Radius of constraint for a 5-minute interval imaged every 1.5s as in B, error bars are standard
error of the calculated radius of constraint., specific details in Table S3. D) Percent of zone 1 foci
for $(CAG)_{130}$ mid/late S-phase cells >100 cells per strain analyzed, see Table S1 for exact
values(***) P≤0.001 compared with wild-type by Fisher's exact test. (^^^) P= 0.0004 by
Fisher's exact test.

825
Figure 4: Damage-Inducible Microtubules associate with expanded CAG repeats in a Dun1
and Cep3 phosphorylation-dependent manner. A) Percentage of zone 1 $(CAG)_{130}$ cells in
mid/late S-phase. Cells were treated with either 15µg/mL nocodazole or an equivalent volume of
DMSO (0.1%). B) Percent zone 1 $(CAG)_{130}$ cells for cells lacking Kar3. C) Example images of
CFP-LacI colocalizing with DIMs. Bottom left image is a cell without a DIM for comparison.
The yellow channel was contrast-adjusted for clarity, raw images can be found in Figure S2A. D)
Fraction of mid-late S phase cells (no CAG tract strain) treated as indicated with at least 1 DIM.
E) Fraction of DIM containing cells of the indicated genotype. DIMs, if present, were scored as
colocalized or not colocalized with the CFP-LacI focus. (*) P≤0.05, (***) P≤0.001, (****) P ≤
0.0001 compared with wild-type by Fisher's exact test (zoning assay) or chi-square test (DIM
assay). See Table S1 (for zoning) or Table S4 (for DIM) for the exact number of cells analyzed,
percentages, and P-values. F) Proposed model for the relocation of a collapsed fork at the CAG
repeat to the nuclear pore complex. Collapsed or uncoupled forks at a CAG repeat result in a
Mrc1 phosphorylation followed by Rad53 activation. Rad53 activated Dun1 leads to
phosphorylation of Cep3. This phosphorylation alters the SPB-CEN attachment, leading to the

formation of DIMs. This may allow repositioning of the centromere on the DIM, for example in
a side-on attachment to permit relocation of the repeat to the NPC. Created with BioRender.com
843
844
845 **Figure S1: Comparison between wild-type strains.** Percent of zone 1 foci for $(CAG)_{130}$
846 mid/late S-phase cells for wild-type cells with the LacO/GFP-LacI array or TetO/mCherry-TetR
847 array ~6.4kb away from $(CAG)_{130}$ near ARS0607 on Chr VI. The number of cells analyzed per
848 strain were 157 and 162 respectively.
849
850 **Figure S2: DIM Examples:** A) Example images of Damage-inducible microtubules. Defined as
851 monopolar microtubule extrusions during S-phase which exhibit rapidly decaying fluorescence
852 intensity. Cell lacking a DIM shown for context. B) Example movie of a DIM associating with
853 an expanded CAG repeat locus. Video acquired at 0.2 frame/s.
854
855

Method Details
Yeast strains and genetic manipulation
Standard procedures were used in cell growth and medium preparation. All *S. cerevisiae* strains
were made in the W303 (RAD5+) background and are listed in Table S5. Point mutant alleles
have been previously characterized: *rad53*[K227A82], *cep3*[S575A45]. The point mutants were generated
using CRISPR-Cas9 methods as described previously[83]. See Table S6 for primers used to create
each mutation. Correct mutation was screened by both restriction enzyme digest and sequencing.
The P_{GAL1}-*CEN6* construct has been previously characterized [51]. Centromere inactivation was
confirmed by plating cells onto galactose containing media and observing a decrease in viability
compared to glucose. Venus-Tub1 was introduced as an additional copy using plasmid
pHIS3p:Venus-Tub1+3'UTR::LEU2, which was a gift from Wei-Lih Lee [58].
867

Microscopy
Colonies were checked for 130-CAG repeats by PCR with primers flanking the repeat
(CTGRev2/T720 or 2StepCAG_pAG32-F/R). PCR products separated on a 2% agarose gel
(MetaPhor, Lonza). Cells from colonies with the correct repeat length were inoculated into YC
media, grown overnight at 30C with shaking, diluted back to OD=0.2 and grown to
approximately 5×10^6 cells per mL in YC media. Cells were fixed with 4% paraformaldehyde
and washed 3 times with 1X PBS. 1.4% YC agar was used to make agar pads on depression
slides and 5 µL of cells were added on top with a coverslip. Z-stack images were taken using a
Zeiss AX10 fluorescent microscope and/or a Delta Vision Ultra High-Resolution using an
Olympus 100X with 1.45NA. Step interval size was 0.15-0.2um and approximately 25 Z-planes
were taken per field of cells. Exposure time was approximately DIC: 100ms; GFP: 300ms and
mCherry: 500ms (for strains with mCherry-TetR). Images were deconvolved, and three-zoning
criteria was used to evaluate the location of the GFP foci for mid/late S-phase cells with the
ImageJ point picker program as described in [84]. Only the middle two-thirds of the stacks were
used for analysis. Mid/Late S-phase cells were determined by yeast morphology using bud size
criteria of 2/3[rd] or less, and at least 15% the size of the mother cell [85]. For drug treatments, cells
were grown to approximately 1×10^7, split into 2 cultures and grown for 30 minutes in either
0.1% dimethyl sulfoxide (DMSO) or 15ug/mL nocodazole. The cells were then fixed and imaged
as described above. For DIM experiments, cells were prepared as described above and imaged

with the filters (and exposure): Cyan (200ms), Yellow (200ms), Red (200ms). Cells were scored
for colocalization determined by any overlap (not inclusive of "touching") of the repeat and the
nuclear periphery, and of the repeat with DIMs. DIMs were defined as monopolar microtubule
structures emanating from the SPB during mid-late S phase (See Figure 4C for examples).
891
P_{GAL1}-CEN6 Inactivation
Colonies were checked for 130-CAG repeats by PCR with primers flanking the repeat
(CTGRev2/T720). Cells from colonies with the correct repeat length were grown overnight in
synthetic complete media +2% glucose. Cultures were spun down and resuspended in YEP-
lactate media to an OD_{600} of 0.3 and grown overnight. Cultures were spun down, split in half,
and resuspended in synthetic complete media +2% galactose or +2% glucose to an OD_{600} of 0.3.
Cultures were grown for 3 hours followed by the zoning assay protocol as outlined above.
899
MSD Analysis
Colonies were checked for 130-CAG repeats by PCR with primers flanking the repeat
(2StepCAG_pAG32-F/R). Cells with correct repeat length were grown to approximately OD_{600}
of 0.5 before being transferred to a concanavalin A-coated chamber slide for attachment. From
there, cells were synchronized by addition of alpha factor (1µM final concentration) then
released by washing 3X with YPD media. A single field of cells was imaged with DeltaVision
Ultra High Resolution microscope using a 100X objective using optical axis integration every
1.5s for 5 minutes. This was repeated using cells in different areas of the slide until a total of 100
minutes of timepoints were taken. Exposure was: Red 400ms, Green 400ms, Pol 50ms (every
50^{th} frame). Images were analyzed with CellProfiler, tracking the LacO/LacI-mCherry focus and
correcting for nuclear translation by tracking Nup49-GFP. Tracked foci were quantified using
$MSD = < \llbracket (x^- - (x\_dt)^-) \rrbracket ^2 >$ and radius of constraint = sqrt(5/4 x MSD plateau) (Dion &
Gasser, 2013). The plateau was calculated by regression in Graphpad PRISM software using the
first 100 Δt timepoints.
914
915   **Quantification and statistical analysis**
916   Prism software was used to calculate statistical significance. For microscopy experiments
917   Fisher's exact test (Relocation experiments) or chi square test (DIM experiments) were used.
918
919   **Data and software availability-** All raw data is available in the supplemental tables

A

	CFP-LacI	mVenus-Tub1	Nup49-mCherry	Merge	Contrast Adjust mVenus-Tub1	Merge

DIM, Repeat @ NPC

No DIM

DIM, Repeat @ NPC

DIM, Repeat not @ NPC

DIM, Repeat not @ NPC

DIM, Repeat not @ NPC

B

	t=0	t=5s	t=10s	t=15s	t=20s	t=25s

Raw

Contrast adjusted