Temporal dynamics of gene expression during metamorphosis in two distant *Drosophila* species

Ozerova AM[1,*], Kulikova DA[2], Evgen'ev MB[3], Gelfand MS[1]

[1] Center for Molecular and Cellular Biology

[2] N.K.Koltsov Institute of Developmental Biology RAS

[3] V.A.Engelhardt Institute of Molecular Biology RAS

* correspondence: alexapogorelskaya@gmail.com

Abstract:

Complete metamorphosis of holometabolous insects is a complex biological process characterized by profound morphological, physiological, and transcriptional changes. To reveal the temporal dynamics of gene expression during this critical developmental transition, a detailed analysis of the developmental transcriptomes of two Drosophila species, *Drosophila melanogaster* and *Drosophila virilis*, was conducted. We confirm partial recapitulation of the embryonic transcriptional program in pupae, but instead of the traditional hourglass model, which posits maximal conservation at mid-embryonic stages, at different stages of pupae we observe a more complicated pattern of alternating low and high diversity, resembling an inverted hourglass, or "spindle", that underscores the complexity of developmental processes during complete metamorphosis. Notably, recently formed genes (specific to insects) exhibit pronounced expression peaks during mid-pupal development, indicating their potential role in developmental transitions.

Keywords:

metamorphosis, development, transcriptomics

Significance Statement:

This study describes the transcriptomic dynamics of complete metamorphosis in holometabolous insects. By analyzing developmental transcriptomes of *Drosophila melanogaster* and *Drosophila virilis*, we characterize the spindle pattern of gene expression during pupal stages, contrasting with the classical hourglass model of embryogenesis. This alternating pattern of low and high transcriptional diversity highlights the unique complexity of metamorphose processes.

Introduction

Insects undergoing the complete metamorphosis (holometabolous insects) represent around 60% of all animals (Mora et al. 2011), suggesting evolutionary success. Their intricate life cycle has been proposed to be a key contributing factor to this accomplishment. One of the adaptive benefits is the occupation of multiple niches which allows switching between alternative resources (Ten Brink, de Roos, and Dieckmann 2019)

Upon hatching, the larvae reach a considerable size within a short timeframe (Texada, Koyama, and Rewitz 2020). Through successive molts, they experience rapid growth, eventually culminating in the stage of transformation, the pupa. Rather than resembling miniature adults, as seen in nymphs of hemimetabolous species, holometabolous insects undergo a radical change that endows them with the appearance of an entirely distinct organism. This process involves rebuilding of virtually all larval organs with those required for adult life.

Several mechanisms underlie terminal differentiation of tissues and organ formation. One such mechanism is dedifferentiation, followed by redifferentiation into adult tissues, a distinguished example being restructuring of the insect longitudinal gut muscles. In the course of this process syncytial larval muscle cells dissociate into mononuclear cells before redifferentiating to form the adult muscles (Schultheis and Frasch 2023).

Unlike the musculature, the intestinal epithelium undergoes a complete renewal process during metamorphosis, representing another mechanism of rebuilding the organs. Adult midgut precursors are maintained on the basal side of the larval midgut throughout the larval development and undergo rapid divisions only after formation of the puparium (Micchelli et al. 2011).

Another example of a complex process during metamorphosis is the development of the adult brain. While larval neurons are retained, the peripheral and central nervous systems are significantly expanded through the activation of numerous neurons that remained in the state of developmental arrest throughout larval growth (Truman 1990).

The intricate processes inherent in pupal development share a level of complexity comparable to embryogenesis, demanding precise orchestration of genes providing the development of emerging adult organs and systems. Transcriptomic analyses across various insect orders revealed reactivation of the embryonic gene expression program at pupal stages (Ozerova and Gelfand 2022). While the developmental hourglass model has been traditionally applied in the context of embryogenesis, its potential extension to the metamorphic phase was not considered so far. It is somewhat parallel to phenomena observed in plants, where the hourglass pattern emerges in transcriptomes at several crucial stages such as germination and floral transition (Drost et al. 2017).

The developmental hourglass model postulates that mid-embryogenesis stages exhibit the highest level of conservation, while early and late stages are more diverged (Duboule 1994). These mid-embryogenesis stages, often referred to as the phylotypic period, play a crucial role in establishing the body plan of the future organism. The term "phylotypic" originates from the period's association with the maximum similarity observed among species within each animal phylum (Richardson 1995). This

conservation is believed to stem from the network of interactions between genes and developmental processes, where any evolutionary modifications could prove deleterious due to potential side effects (Raff 2012). On average, genes that are expressed during the phylotypic stage tend to have higher predicted age compared to genes expressed earlier and later in the course of the development (as measured by the transcriptome age index in (Domazet-Lošo and Tautz 2010)). Another explanation is that the phylotypic period is essential for the precise coordination between growth and patterning (Duboule 1994).

The hourglass pattern was observed on morphological and molecular levels across various groups of organisms, including mammals (based on the expression of human and mouse orthologous genes) (Hazkani-Covo, Wool, and Graur 2005), *Drosophila* spp. *(Kalinka et al. 2010)*, *Caenorhabditis* spp. *(Levin et al. 2012)*. Despite this evidence, the hourglass hypothesis remains a subject of controversy, with some studies challenging both the hourglass model and the existence of the phylotypic stage (Irie 2017). A recent study, employing single-cell resolution to analyze gastrulation in rabbits and mice, demonstrated clear differences in mid-phases between the two species (Mayshar et al. 2023). In this case, earlier and later developmental phases were more similar in terms of cell type composition as revealed by single-cell transcriptome analysis. This observation supports a *spindle model* of the pupal development, proposed here, that is a pattern characterized by low diversity at early stages, increasing in mid-development, and decreasing again at the later stages i.e. an apparent inversion of the known hourglass pattern.

Here, we explore the developmental transcriptomes of two distant cosmopolitan species *Drosophila melanogaster* and *Drosophila virilis*, aiming to characterize the dynamics and conservation of gene expression throughout metamorphosis. A divergence of 50 million years between the species enables quantitative comparisons across orthologous genes.

Results

Reproducibility and preprocessing of RNA sequencing data

To investigate the dynamics of gene expression during development, we performed RNA sequencing (RNA-seq) experiments on the developmental time course of *Drosophila melanogaster* and *Drosophila virilis*. These well-studied organisms are divergent enough to have distinct gene expression profiles, while being close enough to observe an identifiable embryonic hourglass pattern (based on data from (Kalinka et al. 2010), not shown). A comprehensive analysis was conducted across ten distinct developmental stages in each species, including the embryonic phylotypic stage, larvae, seven stages spanning pre-pupae and pupae development, and the adult (imago) stage. Each developmental stage was sampled in three biological replicates to capture the inherent variability in gene expression.

Whole body samples rather than specific tissues were analyzed to allow consistent sampling conditions across all developmental stages, as specific tissues do not persist throughout each stage. This approach enabled the capture of the overall gene expression dynamics reflective of the developmental program as a whole. While tissue-specific expression differences could provide additional insights, a whole-organism approach allowed to maintain comparability across stages, facilitating the identification of broader developmental patterns.

To evaluate the reproducibility of the RNA-seq data, the concordance among the biological replicates for each sample was assessed. The replicates demonstrated a high degree of similarity, indicating robust and reproducible measurements across biological replicates (the Spearman correlation coefficient equal to 0.97 on average, Supplementary Figure 1). The Spearman correlation coefficient was higher than 0.84 for all replicate pairs in both species, the lowest example being larval samples of *D. melanogaster*. Consequently, gene expression values were averaged across the replicates, generating a unified gene expression matrix for the downstream analyses.

In the *D. virilis* dataset, one of the replicates from the P3 and P4 pupal stages showed increased variation compared to the other two replicates, with the greatest divergence observed along the first principal component (Supplementary Fig. 3). To prevent potential bias in the results, downstream analyses were conducted excluding these outlier replicates. All observed dynamics of expression conservation between species remained consistent when the full dataset was analyzed (see below and Supplementary Fig. 4).

An average of 25,000 transcripts per sample for *D. melanogaster* and 20,000 transcripts per sample for *D. virilis* were identified. To simplify the data and facilitate comparative analysis, the transcripts were aggregated at the gene level, resulting in an average of 12,000 genes for each developmental stage in both species. Grouping of genes into orthologous groups yielded 10,000 one-to-one orthologous groups shared by *D. melanogaster* and *D. virilis*. To ensure robust and informative downstream analyses and focus on genes that exhibited significant expression changes during development, the dataset was refined by filtering out genes with low expression values and low variance across the developmental stages (see *Materials and Methods*). The final dataset

comprised 6,200 orthologous groups of genes that demonstrated dynamic expression profiles across the developmental time course in both *D. melanogaster* and *D. virilis*.

Intra-species analysis of transcription profiles during pupal development in *D. melanogaster* and *D. virilis*

To characterize processes occurring in each species, initial analyses of the transcription profiles were performed for *D. melanogaster* and *D. virilis* separately. During the pupal development phase, gradual modifications in the expression profiles were observed, with the transcriptome at each stage exhibiting higher correlation with neighboring stages rather than more distant ones, as determined by the Spearman correlation coefficient (Fig. 1a).

However, several pupal stages were closer to the embryonic state than the larval state. Vice versa, the embryonic transcriptome was more similar to the pupal transcriptomes than to the larval ones, suggesting a partial reversion of the embryonic developmental programs after pupation. These findings are consistent with our previous study conducted on several holometabolous species belonging to various orders (Ozerova and Gelfand 2022), providing further evidence for the reversion of the transcriptional program during pupation. Additionally, a decreased similarity between the P4 stage and the embryo was observed in both *D. melanogaster* and *D. virilis*, indicating the activation of some pupal-specific transcriptional programs at the P4 stage, while earlier and subsequent stages of pupal development are more similar to the embryonic state.

Figure 1. (a) Heatmap of the Spearman correlation coefficients for *D. melanogaster* (above the diagonal) and *D. virilis* (below the diagonal) developmental stages for each species separately. Each square represents the correlation coefficient between the respective stages of development. The color ranges from grey for low correlation coefficients (close to zero) to red for coefficients of 0.5, with more intense colors indicating higher correlation coefficients. **(b, c)** Principal Component Analysis (PCA) of developmental stages in *D. melanogaster* **(c)** and *D. virilis* **(c)**. Each dot represents a distinct developmental stage. The first three components are shown. **(d, e)** Expression dynamics in the course of development for genes homologous to the Yamanaka factors for *D. melanogaster* (d) and *D. virilis* (e). For each gene, mean expression values are shown in various colors with shaded areas representing standard deviation of the expression across replicates.

Principal component analysis (PCA) further corroborates these findings (Fig. 1b,c). In both *D. melanogaster* and *D. virilis*, the first principal component, explaining 34% and 45% of the variance, respectively, effectively separates the four main developmental stages in the sequence of ontogeny. The second principal component, accounting for 28% of the variance in *D. melanogaster* and 18% in *D. virilis*, distinguishes the active larvae and imago stages from the developmental embryo and pupae stages, when the embryonic transcriptome program is activated. The third principal component (13% and 16% of the explained variance, respectively) primarily reflects temporal progression, with pupal stages aligning along this axis. These observations are consistent in both species, with *D. melanogaster* PCA results shown in Figure 1b and *D. virilis* results, in Figure 1c. The percentages represent the proportion of the total variance in the data.

Identification of stage-specific genes and their associated biological processes

Differential expression analysis was performed aiming to identify genes that exhibited significant upregulation at a specific stage compared to other stages.

The larval and imago stages exhibited the largest number of stage-specific genes, with over 1,000 genes identified for each stage in both *D. melanogaster* and *D. virilis*. We also identified approximately 900 genes that were specifically expressed during embryogenesis and all pupal stages.

To gain insights into the functions and roles of genes specifically expressed during the pupal stage, we performed gene ontology (GO) enrichment analysis (see Supplementary Tables 1-6).

Among genes upregulated during the P3 stage (the *bubble prepupa*), we found significant enrichment of terms related to *isoprenoid metabolic process* and *isoprenoid biosynthetic process* (GO:0006720 and GO:0008299, *p*-value<10^{-3} in both species). This observation suggests that lipid metabolism and protein modification are important during the P3 stage of the *Drosophila* development. However, the precise functions of isoprenoids during this stage are not yet fully understood, although the mevalonate pathway and heart formation may be linked by isoprenylation (Yi et al. 2006). Genes upregulated during the P4 stage (the *buoyant*) are enriched in the molecular function termed "*structural constituent of cuticle*" (GO:0042302, *p*-value<10^{-4}). Statistically significant intersections between GO terms enriched in two species were not observed for other pupal stages, likely due to a high similarity in gene expression between temporarily close pupal stages (Fig. 1a), yielding minimal statistical power to detect differences in the GO-term enrichment.

We also specifically analyzed several groups of genes known to be important for development. The Yamanaka factors were used as a general example of genes associated with cellular reprogramming and differentiation. Homologs of the Yamanaka factors in flies were analyzed (Fig. 1de) (Kaur et al. 2018). As expected, these factors exhibited increased expression in pupae in both species. The *Sox* genes had a strong expression peak during early pupal stages that was particularly prominent during the P1-P3 stages in *D. melanogaster*. On the contrary, the *Luna* genes had stable expression values after the onset of pupal development.

Comparative analysis of developmental conservation and divergence in the *Drosophila* species

At the next step, a comparative analysis was performed with the goal to investigate similarities and differences in the developmental gene expression patterns and to identify a potential phylotypic stage.

The comparison between the two species, as shown in Figure 2 and Supplementary Figures 2 and 4, revealed high Spearman correlation coefficients along the diagonal, indicating that each developmental stage demonstrated close similarity to the corresponding stage in the other species, and hence the stages had been correctly assigned during sample collection.

Figure 2. (a) Heatmap of the Spearman correlation coefficients for *D. melanogaster* and *D. virilis* developmental stages. Each square represents the correlation coefficient between the expression profiles of the corresponding stages between the two species. The horizontal axis lists the stages of *D. melanogaster*, while the vertical axis lists the stages of *D. virilis*. The color scheme is as in Fig. 1. **(b)** Spearman correlation coefficients are shown as a separate line plot, representing the correlation between corresponding stages of *D. melanogaster* and *D. virilis*, that is, the main diagonal of the heatmap in (a). The x-axis represents the developmental stages, and the y-axis represents the correlation coefficients. The shaded area shows the standard deviation across replicates. **(c)** Standard deviation (STD) values for gene expression are shown for each developmental stage. The x-axis represents the developmental stages, and the y-axis represents the STD values, for details see the *Results* section. **(d)** The median STD values for each pupal stage shown as a line plot. The x-axis represents the developmental stages, and the y-axis represents the median STD value.

Consistent with pupal samples being closer to embryos than to larvae in the same species, *D. melanogaster* embryos were closer to *D. virilis* pupae than to larvae, and vice versa, but again, with a distinct expression profile prominent at the P4 stage.

The main diagonal of the heatmap in Figure 2a represents the correlation between the same stages during development and is shown separately as Figure 2b. The strongest correlation was observed at the P3 stage, followed by a decrease at the P4 stage and a subsequent peak at the P5 stage. These observations suggest convergence in expression profiles during early and late pupal development.

The standard deviation (STD) from the mean expression across replicates was used as an additional metric of gene expression stability. Genes with more conserved expression have lower STD indicating more consistent expression across replicates, while less constrained genes would have higher STD. The distributions of STD values for each developmental stage (Figure 2c,d) exhibited dynamics similar to the Spearman correlation. Variation in gene expression increased at early pupal stages, specifically at the P3 and P4 stages, indicative of decreased constraints on expression during these stages.

The P4 (buoyant) stage shows the lowest level of interspecies conservation, with both the correlation coefficient and standard deviation indicating minimal similarity compared to neighboring pupal stages, which display higher conservation. While mid and late pupal stages suggest a gradual decline in conservation, further sampling around these timepoints is needed to accurately capture the dynamics of this trend.

Clusters of co-expressed genes

To study gene groups with distinct developmental expression patterns, a hierarchical clustering analysis was performed using the Spearman correlation coefficient as the distance metric (Fig. 3). To focus on the dynamics of pupae, only pupal stages were considered at the clustering step, yielding eight clusters.

Clusters were sorted according to the number of common orthologous genes, so that similar pairs of clusters had the same names in both species (Fig. 3b). In most cases the median gene trajectories of paired clusters turned out to be highly similar between the two species, despite independent clustering in the species (Supplementary Fig. 5).

Cluster pair 1 were the largest and had the largest set of common genes and were composed of genes with expression gradually increasing throughout the lifespan in both *D. melanogaster* and *D. virilis*.

The observed similarities demonstrate that orthologous genes tend to have similar dynamics (statistics provided in Supplementary Fig. 5), supporting the biological relevance of the clusters. Hence, we selected pairs of clusters exhibiting high correlation of expression profiles and containing common orthologous genes, and performed GO enrichment analysis to reveal the functional role of these clusters.

Expression of genes comprising cluster pair 2 increased during the early stages of pupal development followed by a sequential gradual decrease. The enriched GO terms for this cluster were related to the regulation of growth, development and cell migration (Supplementary Fig. 6). Genes in this cluster pair have a pupae-specific expression, since in embryogenesis their expression is relatively low.

Genes in cluster pair 3 had higher correlation of gene expression between the species for all developmental stages compared to the average correlation (Supplementary Fig. 7). Expression of these genes increased during the P5 and mid-pupae stages and also

significantly increased during the embryonic stage. This suggests that these genes are involved in both embryogenesis and metamorphosis transcriptomic programs. The GO enrichment analysis revealed enrichment of terms related to chromatin organization, cell cycle, and developmental processes for these gene clusters.

Cluster 7 in both species has an expression peak at the later pupal stages while also maintaining notable expression in larvae. However, the overlap of genes within these clusters between the two species is relatively small (Figure 3c), indicating that these genes are characterized by species-specific expression patterns rather than a conserved expression pattern across species. Thus, we have not pursued a detailed analysis of these clusters.

GO enrichment results for other cluster pairs with a significant number of intersecting genes and statistically reliable outcomes are given in Supplementary Fig. 8-11.

Figure 3. Clustering of expression profiles. Developmental expression profiles for each of the constructed clusters for D. melanogaster **(a)** and D. virilis **(b)**. Each gene is represented by a gray line, and the median expression profile for each cluster is shown in red. **(c)** Heatmap illustrating the intersection of gene sets in each cluster pair. Each square represents a cluster pair, with the color scheme shown at the right; lighter color indicates a higher ratio of intersecting genes.

Gene age influence on the developmental transcriptome

To gain insights into the evolutionary origin of the observed transcriptome dynamics, the phylostrata analysis was performed. Each gene was assigned to a phylostratum corresponding to its predicted time of emergence (for details, see *Materials and Methods* section).

Genes belonging to different phylostrata exhibit specific temporal dynamics of expression, as shown in Figure 4, while in general having the trend to increase the expression rate during ontogeny. The most ancient genes (gray) tend to have higher expression during the larval stage, suggesting their involvement in housekeeping

functions associated with larval feeding and growth. Intermediate-aged genes (blue), common to *Arthropoda*, *Pancrustacea*, and *Insecta*, have an additional prominent expression peak during the P4 stage and late pupae. Following the trend, the group of youngest, *Holometabola*-specific genes has only pupae-specific peaks, indicating that these newly acquired genes could have been selected in *Holometabola* to perform metamorphosis-related functions.

Many young genes up-regulated during the P4 stage encode proteins with candidate signal peptides that control protein secretion (see *Materials and Methods* for the details). However, signal peptides tend to appear in young genes in general (Supplementary Fig. 12). The distributions of all genes and signal peptide-encoding genes across the phylostrata are statistically different (*p*-value = 8.3×10^{-75}, statistic = 384.2, the Chi-squared test), P4-specific genes follow the same trend with no observed statistically significant difference (the Chi-squared statistic = 0.2).

Figure 4. Phylostrata-specific gene expression dynamics. Horizontal axis — the developmental stage, vertical axis — the median log-transformed expression. The shaded area represents the standard deviation (STD) of gene expression. The oldest phylostrata (see the legend) are shown in gray, middle phylostrata, in blue, and the youngest ones, in green.

Genes from different phylostrata exhibit varying impact on the overall similarity of transcriptomes during metamorphosis, as seen in the plots of correlation between gene expression at each stage, calculated separately for each phylostratum (Fig. 5). For intermediate-aged genes (Fig. 5a, blue), the correlation between expression in the two species was larger at the later pupal stages. Hence, the expression of these genes is conserved closer to the end of the pupal development. Genes common to *Pancrustacea* have an additional conservation peak during earlier stages. Conversely, younger genes (Fig. 5a, green) exhibited higher correlation between species primarily at the onset of the pupal development, when imaginal disks are everted and active differentiation occurs. Notably, however, *Holometabola*-specific genes within this group deviate from this general trend, showing a pattern more similar to that of genes of intermediate ages.

Looking at another angle, that is, comparing expression of individual genes throughout development (Fig. 5b, c), we observed that recently emerged genes from *Diptera* and *Drosophila* phylostrata tended to have higher correlation between species (*p*-values for the one-sided Mann–Whitney U test are 9.7×10^{-3} and 1.2×10^{-5}, respectively). This suggests that either these genes had less evolutionary time to diverge, or their function is crucial for pupae, leading to negative selection on changes in expression. In contrast, genes belonging to the *Pancrustacea* phylostratum exhibited statistically lower correlation than the average (*p*-value equal to 4×10^{-6}).

The median correlation coefficient for genes comprising different phylostrata tends to decrease with the decrease of the predicted age, see old (grey) and middle (blue) phylostrata in Fig. 5c. However a breakpoint is observed after the transition from *Pancrustacea* to *Insecta* and this tendency is reversed, so that the younger the phylostratum (green in Fig. 5c), the higher median correlation between the species it has. The most similar expression throughout the development was observed for the oldest and the youngest genes.

Clusters 4 and 6 (with increased expression at mid-pupae) were enriched with both young, *Drosophila*-specific genes), genes from the *Insecta* phylostratum, and old genes (Fig. 6). Genes in cluster 4 had an expression burst at the P4 stage, while cluster 6 featured higher expression at the early pupal and adult stages. Cluster 5 consisting of genes expressed at early pupal stages was dominated by middle phylostrata. Cluster 1, characterized by a constantly increasing expression profile throughout the development, was strongly enriched with oldest and Bilateria genes.

Figure 5. Phylostrata-based analysis of conservation of gene expression. **(a)** Correlation of gene expression profiles in two species, calculated separately for each phylostratum. Each line represents an average correlation coefficient between species with the shaded areas showing the standard deviation across replicates. **(b)** Density plot for the Spearman correlation coefficients for gene expression in two species across development, calculated separately for each phylostratum. The density plot for all genes is shown as the shaded area. **(c)** Median values for the distribution of correlation coefficients from (b). The color code as in Figure 4.

Figure 6. Intersection between phylostrata and clusters. The color of the square corresponds to the significance of the intersection (see *Materials and Methods* for details).

Discussion

The hourglass model of gene expression, characterized by a phase of higher difference in gene expression at the beginning and end of development with a period of higher conservation in the middle, phylotypic, stage has been well-established in the context of embryogenesis (Irie and Kuratani 2014; Drost et al. 2017). This pattern has been documented across a wide range of species ranging from insects and nematodes to mammals (Kalinka et al. 2010; Levin et al. 2012; Hazkani-Covo, Wool, and Graur 2005). Furthermore, it may extend beyond embryogenesis and, in plants, is also observed in processes like floral transition (Drost et al. 2016), suggesting that this rule has broader implications than initially thought. One hypothesis has been that the pupal development of holometabolous insects, when organogenesis re-starts, adheres to a similar hourglass-like pattern. This was based on our observation that pupae tend to recapitulate embryogenesis transcription patterns (Ozerova and Gelfand 2022). The expectation was that a phase of increased similarity in gene expression might occur during the mid-pupal stages.

Contrary to the classic hourglass model, our findings indicate that while there are stages of conserved gene expression, they predominantly occur during early metamorphosis stages, rather than the middle (Fig. 2). Moreover, we observe that the P4 stage is the most divergent stage in terms of gene expression between species, as measured by both correlation coefficients and standard deviation (Fig. 2a,b). A previous microarray study also identified the P4 stage as being the least similar to the embryonic stages compared to other pupal stages in *D. melanogaster* (Arbeitman et al. 2002). This suggests a spindle-like pattern, where the transcriptome divergence is accentuated during this phase. The major process during this stage is the eversion of the imaginal head sac. Reorganization of the respiratory system could be visually observed since lateral trunk tracheae becomes obscured. Also the oral armature of the larva is expelled, signaling the complete removal of larval feeding structures as the organism prepares for its final differentiation into the adult form. Overall, this underscores the dynamic nature of gene expression during pupal transformation, likely driven by the relatively recent emergence of the latter.

An intriguing observation is that genes expressed during the P4 stage, a period characterized by high divergence, tend to be evolutionarily new. For example, clusters 4 and 6, that have increased expression on this stage, are enriched with young genes (Fig. 6). Moreover, while being relatively highly expressed, these recently emerged genes (specific for *Diptera* and *Drosophila*) demonstrate rather low correlation in expression at the P4 stage compared to the adjacent stages (Fig. 5a).

To place these findings in the evolutionary context, the *dN/dS* ratio for *D. melanogaster* genes (Mensch et al. 2013) was used as a measure of natural selection and evolutionary rate. While genes with higher sequence conservation (low *dN/dS* values) have higher expression levels for all stages except for imago (Supplementary Figure 13a), genes from cluster 4 with expression burst during the P4 stage demonstrate significantly higher substitution rates compared to the complete gene set (*p*-value = 0.006, Supplementary Figure 13c). However, genes that are upregulated exclusively during P4 do not feature a statistically significant difference in the *dN/dS* ratio, likely due to the small number of genes specifically upregulated during this short stage.

Conversely, genes upregulated at the earliest prepupal stage (P1) when the expression levels are highly coordinated (Fig. 2) (*p*-value = 0.012, Supplementary Fig. 13b) and genes from cluster 2, which show peak expression at P1 (*p*-value = 0.002, Supplementary Fig. 13c), have significantly lower *dN/dS* ratios and hence are more conserved.

Remarkably, we find that genes with a middle predicted age (emerging in *Arthropoda* and *Pancrustacea*) have conserved expression patterns in the studied *Drosophila* species during later pupal stages (Fig. 5a). The expression range of these genes remains conserved, suggesting that they did not diverge after the emergence. Indeed, genes from these phylostrata are outliers in the *dN/dS* ratio distribution (Supplementary Fig. 12d), whereas genes from younger phylostrata tend to display higher *dN/dS* ratios.

Our study provides insights into the transcriptional dynamics of Drosophila metamorphosis, shedding light on the temporal patterns of gene expression during this critical developmental process. However, delving deeper into the temporal dynamics by analyzing additional stages could offer a more comprehensive understanding of metamorphosis. By including more initial and intermediate stages, we could capture finer transitions and potentially identify novel regulatory mechanisms underlying metamorphic events. On the other dimension, analyzing a wider range of *Drosophila* species would enable us to refine the timing of key pupal developmental stages and enhance the statistical power of our comparative analyses.

Additionally, expanding the scope of comparisons beyond *Drosophila* species — studying insects with mobile pupae, such as mosquitoes, presents a unique opportunity to investigate the preservation of larval structures during pupal development. Unlike *Drosophila*, whose immobile pupae undergo extensive tissue remodeling, insects with mobile pupae should retain larval organs due to the need for locomotion and feeding during the pupal stage. Investigating the molecular mechanisms and gene expression dynamics underlying these differences between species with fixed and mobile pupae could uncover fundamental insights into the evolutionary diversity of metamorphic strategies across insects.

Materials and Methods

Sample collection

Wild-type *D. melanogaster* (Oregon-R) and wild-type *D. virilis* (Strain 9) were used for RNA sequencing. These two distant species were selected for the analysis because they belong to two different subgenera, *Sophophora* and *Drosophila*, respectively (O'Grady and DeSalle 2018). Therefore, the observed developmental patterns should be common for the whole *Drosophila* genus.

The flies, spanning all developmental stages, were reared on the standard *Drosophila* wheat meal-yeast-sugar-agar medium and maintained at a constant temperature of 25°C. The *D. virilis* developmental rate being approximately 1.5 times slower than that of *D. melanogaster*, to account for the intrinsic differences in developmental timing between the two species, the temporal synchrony of developmental stages was monitored by visual observation.

In order to synchronize the ages of collected embryos, adult flies were allowed to reside in a Petri dish containing a 3% agarose substrate covered with apple juice and yeast for one hour. Subsequently, fresh dishes were employed for embryo collection. The collection process transpired over a one-hour interval, followed by incubation of the dishes at an appropriate temperature until the embryos reached the phylotypic stage, which required 10 hours for *D. melanogaster* and 15 hours for *D. virilis*. Larvae were individually collected with developmental stages monitored visually. All biological replicates for *D. virilis* larvae were second instar larvae, while *D. melanogaster* larvae were sequenced at all three ages. Pupae were also collected individually with selective individual monitoring, without distinguishing males and females, including stages P1, P2, P3, P4, and P5, as per the developmental stage descriptions outlined by (Bainbridge and Bownes 1981). Furthermore, intermediate (24 and 36 hours of development for *D. melanogaster* and *D. virilis*, respectively) and late (48 and 72 hours of development for *D. melanogaster* and *D. virilis*, respectively) pupal stages were collected. A particular focus was placed on the early pupal stages (P1-P5), as they encompass key developmental transitions, including the reorganization of larval tissues and the differentiation of adult structures. These early stages are particularly dynamic and critical for metamorphosis, marking major shifts in gene expression and tissue remodeling. In contrast, later pupal stages primarily involve growth and refinement processes, with the developing organism already resembling the adult form. Adult samples consisted of five-day-old males. Adult females were excluded from the experiment to avoid the influence of the ovaries with eggs, as in this case the full transcriptome would be skewed towards the egg state. The influence of sex on other stages was considered to be negligible, and each biological replicate was expected to consist of almost equal proportion of males and females, although the sex ratio in non-adult stages was not directly verified.

All samples were collected in three independent biological replicates. Subsequently, the samples underwent a thorough rinsing process to remove any residual food and yeast, using the PBS buffer. Following this step, total RNA was extracted from the samples using RNAzol RT reagent (MRC, Cincinnati, OH, USA), following the manufacturer's recommended protocols. RNA quality assessments were performed through gel

electrophoresis, and the RNA samples were carefully stored in ethanol at −70°C, in preparation for subsequent experimental procedures.

Sequencing

Ribosomal RNA depletion was conducted using the NEBNext® Poly(A) mRNA Magnetic Isolation Module. Subsequently, RNA library preparation was carried out with the NEBNext® Ultra™ II Directional RNA Library Prep Kit, and the prepared libraries were sequenced on the Illumina HiSeq4000 platform in the Skoltech Genomics Core Facility. The sequencing yielded paired-end reads with the length of 150 nucleotides. Due to suboptimal quality observed in the replicates of *D. melanogaster* 2nd instar larvae, we made the decision to use the 1st and 3rd instar samples for the downstream analyses.

Processing of transcriptome data

Low-quality nucleotides and reads were systematically eliminated using the `fastp` tool with the default parameters. Subsequently, the filtered reads were aligned to the respective reference genomes with the `kallisto` pseudomapping procedure and the BDGP6.46 and DvirRS2 assemblies as references for *D. melanogaster* and *D. virilis*, respectively. Transcript-level data were aggregated to the gene level using a custom Python script. Based on the observed high reproducibility, gene expression values were averaged across the replicates.

To mitigate noise, genes expressed at low levels and exhibiting limited variance across the developmental stages were excluded from the subsequent analyses. The threshold for the expression values was set as the inflection point of the expression values density plot, separating the bell-shaped part of the distribution from low values (95% and 93% genes are preserved for *D. melanogaster* and *D. virilis* respectively). The variance threshold was set on the first inflection point of the cumulative distribution. Overall, 69% of genes passed specified filters for both species.

Orthologous groups and functional analysis

Orthologous gene groups were downloaded from the `Ensembl` database using the `Biomart` tool. Subsequently, only genes belonging to orthologous groups shared by both species were considered for the subsequent analyses. Gene ontology (GO) term annotations were obtained from the `AmiGO2` database for *D. melanogaster* and from the `Ensembl` database for *D. virilis*.

GO term enrichment analysis was performed using the `goatools` Python package (Klopfenstein et al. 2018) with the default settings and an adjusted *p*-value threshold of 0.05. The analysis used a reference set of all genes showing positive expression values in the analyzed dataset. The semantic analysis of the enriched GO terms was visualized using the `REVIGO` tool (Supek et al. 2011) and plotted with the R `ggplot` package.

To predict the presence of signal peptides in the protein sequences, the `SignalP-6.0` tool was used with the default eukaryotic settings (Teufel et al. 2022).

Data analysis

Ad hoc `Python` scripts were used for data analysis and visualization, and figures were prepared using the matplotlib and seaborn packages. Statistical tests were performed using the `Scipy` package.

Analysis of phylostrata

For the phylostrata annotation of *D. melanogaster* genes, a phylostratigraphy approach was used (Domazet-Loso, Brajković, and Tautz 2007). For example, a gene was assigned to the *Holometabola* phylostratum if its homologs were found exclusively within the *Holometabola* group, indicating an estimated age of approximately 340 million years.

To assess the significance of enrichment for each phylostratum in each cluster, the following ratio was used:

$$\rho_{i,j} = \frac{(observed_{i,j} - expected_{i,j})^2}{expected_{i,j}}$$

where $observed_{i,j}$ is the number of common genes in the intersection of cluster *i* and phylostratum *j*, and $expected_{i,j} = \frac{(gene\ count\ in\ cluster\ i) \times (gene\ count\ in\ phylostratum\ j)}{total\ gene\ count}$. This metric highlights substantial intersections between clusters and phylostrata, while adjusting for the overall gene count within each group.

Identification of stage-specific genes

A gene was classified as stage-specific if it met two criteria: (1) it was significantly upregulated at a particular stage, with an adjusted *p*-value < 0.05 and a positive log-fold change, as determined by the DESeq2 algorithm, and (2) it was not upregulated at any other stage, except for the adjacent ones.

Acknowledgements

This study was supported by the Russian Science Foundation via grant 23-14-00136.

We acknowledge the Skoltech Genomics Core Facility for library preparation and sequencing supported by the Skoltech Center of Molecular and Cellular Biology grant for the students' projects.

Experiments were conducted using the equipment of the Core Centrum of the Institute of Developmental Biology RAS, supported by the IDB RAS research program 0088-2024-0010.

We also acknowledge Ilya Borisovitch Mertsalov for technical assistance in the collection of specimens.

Data availability

The RNA sequencing data that support the findings of this study are available in the NCBI GEO data repository under the accession GSE269951.

Competing interests

The authors declare no competing interests.

Author contributions statement

Ozerova AM* - Data curation, Formal analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – original draft
Kulikova DA - Investigation, Methodology, Writing – review & editing
Evgen'ev MB - Resources, Supervision, Writing – review & editing
Gelfand MS - Conceptualization, Funding acquisition, Methodology, Project administration, Resources, Supervision, Validation, Writing – review & editing

Arbeitman, Michelle N., Eileen E. M. Furlong, Farhad Imam, Eric Johnson, Brian H. Null, Bruce S. Baker, Mark A. Krasnow, Matthew P. Scott, Ronald W. Davis, and Kevin P. White. 2002. "Gene Expression during the Life Cycle of Drosophila Melanogaster." *Science* 297 (5590): 2270–75.

Bainbridge, S. P., and M. Bownes. 1981. "Staging the Metamorphosis of Drosophila Melanogaster." *Journal of Embryology and Experimental Morphology* 66 (December):57–80.

Domazet-Loso, Tomislav, Josip Brajković, and Diethard Tautz. 2007. "A Phylostratigraphy Approach to Uncover the Genomic History of Major Adaptations

in Metazoan Lineages." *Trends in Genetics: TIG* 23 (11): 533–39.

Domazet-Lošo, Tomislav, and Diethard Tautz. 2010. "A Phylogenetically Based Transcriptome Age Index Mirrors Ontogenetic Divergence Patterns." *Nature* 468 (7325): 815–18.

Drost, Hajk-Georg, Julia Bellstädt, Diarmuid S. Ó'Maoiléidigh, Anderson T. Silva, Alexander Gabel, Claus Weinholdt, Patrick T. Ryan, et al. 2016. "Post-Embryonic Hourglass Patterns Mark Ontogenetic Transitions in Plant Development." *Molecular Biology and Evolution* 33 (5): 1158–63.

Drost, Hajk-Georg, Philipp Janitza, Ivo Grosse, and Marcel Quint. 2017. "Cross-Kingdom Comparison of the Developmental Hourglass." *Current Opinion in Genetics & Development* 45 (August):69–75.

Duboule, D. 1994. "Temporal Colinearity and the Phylotypic Progression: A Basis for the Stability of a Vertebrate Bauplan and the Evolution of Morphologies through Heterochrony." *Development . Supplement*, 135–42.

Hazkani-Covo, Einat, David Wool, and Dan Graur. 2005. "In Search of the Vertebrate Phylotypic Stage: A Molecular Examination of the Developmental Hourglass Model and von Baer's Third Law." *Journal of Experimental Zoology. Part B, Molecular and Developmental Evolution* 304 (2): 150–58.

Irie, Naoki. 2017. "Remaining Questions Related to the Hourglass Model in Vertebrate Evolution." *Current Opinion in Genetics & Development* 45 (August):103–7.

Irie, Naoki, and Shigeru Kuratani. 2014. "The Developmental Hourglass Model: A Predictor of the Basic Body Plan?" *Development* 141 (24): 4649–55.

Kalinka, Alex T., Karolina M. Varga, Dave T. Gerrard, Stephan Preibisch, David L. Corcoran, Julia Jarrells, Uwe Ohler, Casey M. Bergman, and Pavel Tomancak. 2010. "Gene Expression Divergence Recapitulates the Developmental Hourglass Model." *Nature* 468 (7325): 811–14.

Kaur, Prameet, Helen Jingshu Jin, Jay B. Lusk, and Nicholas S. Tolwinski. 2018. "Modeling the Role of Wnt Signaling in Human and Drosophila Stem Cells." *Genes* 9 (2). https://doi.org/10.3390/genes9020101.

Klopfenstein, D. V., Liangsheng Zhang, Brent S. Pedersen, Fidel Ramírez, Alex Warwick Vesztrocy, Aurélien Naldi, Christopher J. Mungall, et al. 2018. "GOATOOLS: A Python Library for Gene Ontology Analyses." *Scientific Reports* 8 (1): 10872.

Levin, Michal, Tamar Hashimshony, Florian Wagner, and Itai Yanai. 2012. "Developmental Milestones Punctuate Gene Expression in the Caenorhabditis Embryo." *Developmental Cell* 22 (5): 1101–8.

Mayshar, Yoav, Ofir Raz, Saifeng Cheng, Raz Ben-Yair, Ron Hadas, Netta Reines, Markus Mittnenzweig, et al. 2023. "Time-Aligned Hourglass Gastrulation Models in Rabbit and Mouse." *Cell* 186 (12): 2610–27.e18.

Mensch, Julián, François Serra, Nicolás José Lavagnino, Hernán Dopazo, and Esteban Hasson. 2013. "Positive Selection in Nucleoporins Challenges Constraints on Early Expressed Genes in Drosophila Development." *Genome Biology and Evolution* 5 (11): 2231–41.

Micchelli, Craig A., Lisa Sudmeier, Norbert Perrimon, Shan Tang, and Ryan Beehler-Evans. 2011. "Identification of Adult Midgut Precursors in Drosophila." *Gene Expression Patterns: GEP* 11 (1-2): 12–21.

Mora, Camilo, Derek P. Tittensor, Sina Adl, Alastair G. B. Simpson, and Boris Worm. 2011. "How Many Species Are There on Earth and in the Ocean?" *PLoS Biology* 9 (8): e1001127.

O'Grady, Patrick M., and Rob DeSalle. 2018. "Phylogeny of the Genus Drosophila." *Genetics* 209 (1): 1–25.

Ozerova, Alexandra M., and Mikhail S. Gelfand. 2022. "Recapitulation of the Embryonic Transcriptional Program in Holometabolous Insect Pupae." *Scientific Reports* 12 (1): 17570.

Raff, Rudolf A. 2012. *The Shape of Life: Genes, Development, and the Evolution of Animal Form*. University of Chicago Press.

Richardson, M. K. 1995. "Heterochrony and the Phylotypic Period." *Developmental Biology* 172 (2): 412–21.

Schultheis, Dorothea, and Manfred Frasch. 2023. "Longitudinal Visceral Muscles in Drosophila Fully Dedifferentiate and Fragment prior to Their Reestablishment during Metamorphosis." *microPublication Biology* 2023 (March). https://doi.org/10.17912/micropub.biology.000756.

Supek, Fran, Matko Bošnjak, Nives Škunca, and Tomislav Šmuc. 2011. "REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms." *PLoS One* 6 (7): e21800.

Ten Brink, Hanna, André M. de Roos, and Ulf Dieckmann. 2019. "The Evolutionary Ecology of Metamorphosis." *The American Naturalist* 193 (5): E116–31.

Teufel, Felix, José Juan Almagro Armenteros, Alexander Rosenberg Johansen, Magnús Halldór Gíslason, Silas Irby Pihl, Konstantinos D. Tsirigos, Ole Winther, Søren Brunak, Gunnar von Heijne, and Henrik Nielsen. 2022. "SignalP 6.0 Predicts All Five Types of Signal Peptides Using Protein Language Models." *Nature Biotechnology* 40 (7): 1023–25.

Texada, Michael J., Takashi Koyama, and Kim Rewitz. 2020. "Regulation of Body Size and Growth Control." *Genetics* 216 (2): 269–313.

Truman, J. W. 1990. "Metamorphosis of the Central Nervous System of Drosophila." *Journal of Neurobiology* 21 (7): 1072–84.

Yi, Peng, Zhe Han, Xiumin Li, and Eric N. Olson. 2006. "The Mevalonate Pathway Controls Heart Formation in Drosophila by Isoprenylation of Ggamma1." *Science* 313 (5791): 1301–3.