Temporal dynamics of gene expression during metamorphosis in two distant *Drosophila* species

Ozerova AM[1,*], Kulikova DA[2], Evgen'ev MB[3], Gelfand MS[1]

[1] Skolkovo Institute of Science and Technology

[2] N.K.Koltsov Institute of Developmental Biology RAS

[3] V.A.Engelhardt Institute of Molecular Biology RAS

* correspondence: alexapogorelskaya@gmail.com

Abstract:

Complete metamorphosis of holometabolous insects is a complex biological process characterized by profound morphological, physiological, and transcriptional changes. To reveal the temporal dynamics of gene expression during this critical developmental transition, a detailed analysis of the developmental transcriptomes of two Drosophila species, *Drosophila melanogaster* and *Drosophila virilis*, was conducted. We confirm partial recapitulation of the embryonic transcriptional program in pupae, but instead of the traditional hourglass model, which posits maximal conservation at mid-embryonic stages, at different stages of pupae we observe a more complicated pattern of alternating low and high diversity, resembling an inverted hourglass, or "spindle". This observation challenges the notion of a singular conserved phylotypic period during holometabola ontogeny and underscores the complexity of developmental processes during complete metamorphosis. Notably, recently formed genes (specific to insects) exhibit pronounced expression peaks during mid-pupal development, underscoring their potential role in developmental transitions.

Introduction

Insects undergoing the complete metamorphosis (holometabolous insects) represent around 60% of all animals (Mora et al. 2011), suggesting evolutionary success and fast adaptation. Their intricate life cycle has been proposed to be a key contributing factor to this accomplishment. One of the adaptive benefits is the occupation of multiple niches which allows switching between alternative resources (Ten Brink, de Roos, and Dieckmann 2019)

Upon hatching, the larvae reach a considerable size within a short timeframe (Texada, Koyama, and Rewitz 2020). Through successive molts, they experience rapid growth, eventually culminating in the stage of transformation, the pupa. Rather than resembling miniature adults, as seen in larvae of hemimetabolous species, holometabolous insects undergo a radical change that endows them with the appearance of an entirely distinct organism. This process involves replacement of virtually all larval organs with those required for adult life.

Several mechanisms underlie terminal differentiation of tissues and organ formation. One such mechanism is dedifferentiation, followed by redifferentiation into adult tissues, a distinguished example being restructuring of the insect longitudinal gut muscles. In the course of this process syncytial larval muscle cells dissociate into mononuclear cells before redifferentiating to form the adult muscles (Schultheis and Frasch 2023).

Unlike the musculature, the intestinal epithelium undergoes a complete renewal process during metamorphosis, representing another mechanism of cell differentiation. Adult midgut precursors are maintained on the basal side of the larval midgut throughout the larval development and undergo rapid divisions only after formation of the puparium (Micchelli et al. 2011).

The intricate processes inherent in pupal development share a level of complexity comparable to embryogenesis, demanding precise orchestration of genes providing the development of emerging adult organs and systems. Transcriptomic analyses across various insect orders revealed reactivation of the embryonic gene expression program at pupal stages (Ozerova and Gelfand 2022). While the developmental hourglass model has been traditionally applied in the context of embryogenesis, its potential extension to the metamorphic phase was not considered so far. It is somewhat parallel to phenomena observed in plants, where the hourglass pattern emerges in transcriptomes at several crucial stages such as germination and floral transition (Drost et al. 2017).

The developmental hourglass model postulates that mid-embryogenesis stages exhibit the highest level of conservation, while early and late stages are more diverged (Duboule 1994). These mid-embryogenesis stages, often referred to as the phylotypic period, play a crucial role in establishing the body plan of the future organism. The term "phylotypic" originates from the period's association with the maximum similarity observed among species within each animal phylum (Richardson 1995). This conservation is believed to stem from the network of interactions between genes and developmental processes, where any evolutionary modifications could prove deleterious due to potential side effects (Raff 2012). Another explanation is that the phylotypic period is essential for the precise coordination between growth and patterning (Duboule 1994).

The hourglass pattern has found support on both morphological and molecular levels across various groups of organisms, including mammals (based on mouse and human orthologous genes) (Hazkani-Covo, Wool, and Graur 2005), *Drosophila (Kalinka et al. 2010)*, *Caenorhabditis (Levin et al. 2012)*, zebrafish (Irmler, Schmidt, and Starck 2004). Despite this evidence, the hourglass hypothesis remains a subject of controversy, with some studies challenging both the hourglass model and the existence of the phylotypic stage (Irie 2017). A recent study, employing single-cell resolution to analyze gastrulation in rabbits and mice, demonstrated clear differences in mid-phases between the two species (Mayshar et al. 2023). In this case, earlier and later developmental phases were more similar in terms of cell type composition as revealed by single-cell transcriptome analysis. This observation supports a *spindle model* proposed by us, a pattern characterized by low diversity at early stages, increasing in mid-development, and decreasing again at the later stages i.e. an apparent inversion of the known hourglass pattern.

Here, we explore the developmental transcriptomes of two distant cosmopolitan species *Drosophila melanogaster* and *Drosophila virilis*, aiming to characterize the dynamics and conservation of gene expression throughout metamorphosis. A divergence of 50 million years between the species enables quantitative comparisons across orthologous genes.

Materials and Methods

Sample collection

Wild-type *D. melanogaster* (Oregon-R) and wild-type *D. virilis* (Strain 9) were used for RNA sequencing. These two distant species were selected for the analysis because they belong to two different subgenera, *Sophophora* and *Drosophila*, respectively (O'Grady and DeSalle 2018). Therefore, the observed developmental patterns should be common for the whole *Drosophila* genus.

The flies, spanning all developmental stages, were reared on the standard *Drosophila* wheat meal-yeast-sugar-agar medium and maintained at a constant temperature of 25°C. The *D. virilis* developmental rate being approximately 1.5 times slower than that of *D. melanogaster*, to account for the intrinsic differences in developmental timing between the two species, the temporal synchrony of developmental stages was monitored by visual observation.

In order to synchronize the ages of collected embryos, adult flies were allowed to reside in a Petri dish containing a 3% agarose substrate covered with apple juice and yeast for one hour. Subsequently, fresh dishes were employed for embryo collection. The collection process transpired over a one-hour interval, followed by incubation of the dishes at an appropriate temperature until the embryos reached the phylotypic stage, which required 10 hours for *D. melanogaster* and 15 hours for *D. virilis*. Larvae were individually collected with developmental stages monitored visually. Pupae were also collected individually with selective individual monitoring, without distinguishing males and females, including stages P1, P2, P3, P4, and P5, as per the developmental stage descriptions outlined by (Bainbridge and Bownes 1981). Furthermore, intermediate (24 and 36 hours of development for *D. melanogaster* and *D. virilis*, respectively) and late (48 and 72 hours of development for *D. melanogaster* and *D. virilis*, respectively) pupal stages were collected. Adult samples consisted of five-day-old males.

All samples were collected in three independent biological replicates. Subsequently, the samples underwent a thorough rinsing process to remove any residual food and yeast, using the PBS buffer. Following this step, total RNA was extracted from the samples using RNAzol RT reagent (MRC, Cincinnati, OH, USA), following the manufacturer's recommended protocols. RNA quality assessments were performed through gel electrophoresis, and the RNA samples were carefully stored in ethanol at -70°C, in preparation for subsequent experimental procedures.

Sequencing

Ribosomal RNA depletion was conducted using the NEBNext® Poly(A) mRNA Magnetic Isolation Module. Subsequently, RNA library preparation was carried out with the NEBNext® Ultra™ II Directional RNA Library Prep Kit, and the prepared libraries were sequenced on the Illumina HiSeq4000 platform in the Skoltech Genomics Core Facility. The sequencing yielded paired-end reads with the length of 150 nucleotides. Due to suboptimal quality observed in the replicates of *D. melanogaster* 2[nd] instar larvae, we made the decision to use the 1[st] and 3[rd] instar samples for the downstream analyses.

Processing of transcriptome data

Low-quality nucleotides and reads were systematically eliminated using the `fastp` tool
with the default parameters. Subsequently, the filtered reads were aligned to the
respective reference genomes with the `kallisto` pseudomapping procedure and the
BDGP6.46 and DvirRS2 assemblies as references for *D. melanogaster* and *D. virilis*,
respectively. Transcript-level data were aggregated to the gene level using a custom
Python script. Based on the observed high reproducibility, gene expression values were
averaged across the replicates.

To mitigate noise, genes expressed at low levels and exhibiting limited variance across
the developmental stages were excluded from the subsequent analyses. The threshold
for the expression values was set as the inflection point of the expression values density
plot, separating the bell-shaped part of the distribution from low values (95% and 93%
genes are preserved for *D. melanogaster* and *D. virilis* respectively). The variance
threshold was set on the first inflection point of the cumulative distribution. Overall, 69%
of genes passed specified filters for both species.

Orthologous groups and functional analysis

Orthologous gene groups were downloaded from the `Ensembl` database using the
`Biomart` tool. Subsequently, only genes belonging to orthologous groups shared by
both species were considered for the subsequent analyses. Gene ontology (GO) term
annotations were obtained from the `AmiGO2` database for *D. melanogaster* and from the
`Ensembl` database for *D. virilis*.

GO term enrichment analysis was performed using the `goatools` Python package
(Klopfenstein et al. 2018) with the default settings and an adjusted p-value threshold of
0.05. The analysis used a reference set of all genes showing positive expression values
in the analyzed dataset. The semantic analysis of the enriched GO terms was visualized
using the `REVIGO` tool (Supek et al. 2011) and plotted with the R `ggplot` package.

To predict the presence of signal peptides in the protein sequences, the `SignalP-6.0`
tool was used with the default eukaryotic settings (Teufel et al. 2022).

Data analysis

Ad hoc `Python` scripts were used for data analysis and visualization, and figures were
prepared using the matplotlib and seaborn packages. Statistical tests were performed
using the `Scipy` package.

Analysis of phylostrata

For the phylostrata annotation of *D. melanogaster* genes, a phylostratigraphy approach
was used (Domazet-Loso, Brajković, and Tautz 2007). For example, a gene was
assigned to the *Holometabola* phylostratum if its homologs were found exclusively
within the *Holometabola* group, indicating an estimated age of approximately 340 million
years.

To assess the significance of enrichment for each phylostratum in each cluster, the
following ratio was used:

$$\rho_{i,j} = \frac{(observed_{i,j} - expected_{i,j})^2}{expected_{i,j}}$$

where $observed_{i,j}$ is the number of common genes in the intersection of cluster *i* and

phylostratum *j*, and $expected_{i,j} = \frac{(gene\ count\ in\ cluster\ i)\ *\ (gene\ count\ in\ phylostratum\ j)}{total\ gene\ count}$. This metric

allows differentiation of statistically significant intersection sizes.

Results

Reproducibility and preprocessing of RNA sequencing data

To investigate the dynamics of gene expression during development, we performed RNA sequencing (RNA-seq) experiments on the developmental time course of *Drosophila melanogaster* and *Drosophila virilis*. A comprehensive analysis was conducted across ten distinct developmental stages in each species, including the embryonic phylotypic stage, larvae, seven stages spanning pre-pupae and pupae development, and the adult (imago) stage. Each developmental stage was sampled in three biological replicates to capture the inherent variability in gene expression.

To evaluate the reproducibility of the RNA-seq data, the concordance among the biological replicates for each sample was assessed. The replicates demonstrated a high degree of similarity, indicating robust and reproducible measurements across biological replicates (the Spearman correlation coefficient equal to 0.97 on average, Supplementary Figure 1). The Spearman correlation coefficient was higher than 0.84 for all replicate pairs in both species, the lowest example being larval samples of *D. melanogaster*. Consequently, gene expression values were averaged across the replicates, generating a unified gene expression matrix for the downstream analyses.

An average of 25,000 transcripts per sample for *D. melanogaster* and 20,000 transcripts per sample for *D. virilis* were identified. To simplify the data and facilitate comparative analysis, the transcripts were aggregated at the gene level, resulting in an average of 12,000 genes for each developmental stage in both species. Grouping of genes into orthologous groups yielded 10,000 one-to-one orthologous groups shared by *D. melanogaster* and *D. virilis*. To ensure robust and informative downstream analyses and focus on genes that exhibited significant expression changes during development, the dataset was refined by filtering out genes with low expression values and low variance across the developmental stages (see *Materials and Methods*). The final dataset comprised 6,200 orthologous groups of genes that demonstrated dynamic expression profiles across the developmental time course in both *D. melanogaster* and *D. virilis*.

Intra-species analysis of transcription profiles during pupal development in *D. melanogaster* and *D. virilis*

To characterize processes occurring in each species, initial analyses of the transcription profiles were performed for *D. melanogaster* and *D. virilis* separately. During the pupal development phase, gradual modifications in the expression profiles were observed, with the transcriptome at each stage exhibiting higher correlation with neighboring stages rather than more distant ones, as determined by the Spearman correlation coefficient (Fig. 1a).

However, several pupal stages were closer to the embryonic state than the larval state. Vice versa, the embryonic transcriptome was more similar to the pupal transcriptomes than to the larval ones, suggesting a partial reversion of the transcriptional program after pupation. These findings are consistent with our previous study conducted on several holometabolous species belonging to various orders (Ozerova and Gelfand 2022), providing further evidence for the reversion of the transcriptional program during pupation. Additionally, a decreased similarity between the P4 stage and the embryo was

observed in both *D. melanogaster* and *D. virilis*, indicating the activation of a pupal-specific transcriptional program at the P4 stage, while earlier and subsequent stages of pupal development tended to invoke embryonic-like programs.

Figure 1. (a) Heatmap of the Spearman correlation coefficients for *D. melanogaster* (above the diagonal) and *D. virilis* (below the diagonal) developmental stages. Each square represents the correlation coefficient between the respective stages of development. The color ranges from black for low correlation coefficients (close to zero) to red for coefficients of 0.5, with lighter colors indicating higher correlation coefficients. **(b, c)** Principal Component analysis (PCA) of developmental stages in *D. melanogaster* **(c)** and *D. virilis* **(c)**. Each dot represents a distinct developmental stage. The first three components are shown. **(d, e)** Expression dynamics in the course of development for genes homologous to the Yamanaka factors for *D. melanogaster* (d) and *D. virilis* (e). For each gene, mean expression values are shown in various colors with shaded areas representing standard deviation of the expression across replicates.

Principal component analysis (PCA) further corroborates these findings (Fig. 1b,c). In both *D. melanogaster* and *D. virilis*, the first principal component, explaining 34% and 45% of the variance, respectively, effectively separates the four main developmental stages in the sequence of ontogeny. The second principal component, accounting for 28% of the variance in *D. melanogaster* and 18% in *D. virilis*, distinguishes the active larvae and imago stages from the developmental embryo and pupae stages, when the embryonic transcriptome program is activated. The third principal component (13% and 16% of the explained variance) primarily reflects temporal progression, with pupal stages aligning along this axis. These observations are consistent in both species.

Identification of stage-specific genes and their associated biological processes

Differential expression analysis was performed aiming to identify genes that exhibited significant upregulation at a specific stage compared to other stages.

The larval and imago stages exhibited the largest number of stage-specific genes, with over 1,000 genes identified for each stage in both *D. melanogaster* and *D. virilis*. We also identified approximately 900 genes that were specifically expressed during embryogenesis and all pupal stages.

To gain insights into the functions and roles of genes specifically expressed during the pupal stage, we performed gene ontology (GO) enrichment analysis (see Supplementary Tables 1-6).

Among genes upregulated during the P3 stage (the *bubble prepupa*), we found significant enrichment of terms related to *isoprenoid metabolic process* and *isoprenoid biosynthetic process* (GO:0006720 and GO:0008299, p-value<10^{-3} in both species). This observation suggests that lipid metabolism and protein modification are important during the P3 stage of the *Drosophila* development. However, the precise functions of isoprenoids during this stage are not yet fully understood, although the mevalonate pathway and heart formation may be linked by isoprenylation (Yi et al. 2006). Genes upregulated during the P4 stage (the *buoyant*) are enriched in the molecular function termed "*structural constituent of cuticle*" (GO:0042302, p-value<10^{-4}). Statistically significant intersections between GO terms enriched in the two species were not observed for other pupal stages.

We also specifically analyzed several groups of genes known to be important for development. Thus, homologs of the Yamanaka factors that are associated with cell pluripotency were analyzed (Fig. 1de) (Kaur et al. 2018). As expected, these factors exhibited increased expression in pupae in both species. The *Sox* genes had a strong expression peak during early pupal stages that was particularly prominent during the P1-P3 stages in *D. melanogaster*. On the contrary, the *Luna* genes had stable expression values after the onset of pupal development.

Comparative analysis of developmental conservation and divergence in the *Drosophila* species

At the next step, a comparative analysis was performed with the goal to investigate similarities and differences in the developmental gene expression patterns and to identify a potential phylotypic stage.

The comparison between the two species, as shown in Figure 2, revealed high Spearman correlation coefficients along the diagonal, indicating that each developmental stage demonstrated close similarity to the corresponding stage in the other species, and hence the stages had been correctly assigned during sample collection.

Figure 2. (a) Heatmap of the Spearman correlation coefficients for *D. melanogaster* and *D. virilis*
developmental stages. Each square represents the correlation coefficient between the expression profiles
of the corresponding stages of the two species. The color scheme is as in Fig. 1. **(b)** Spearman
correlation coefficients are shown as a separate line plot, representing the correlation between
corresponding stages of *D. melanogaster* and *D. virilis*, that is, the main diagonal of the heatmap in (a).
The x-axis represents the developmental stages, and the y-axis represents the correlation coefficients.
The shaded area shows the standard deviation across replicates. **(c)** Standard deviation (STD) values for
gene expression are shown for each developmental stage. The x-axis represents the developmental
stages, and the y-axis represents the STD values, for details see the *Results* section. **(d)** The median
STD values for each pupal stage shown as a line plot. The x-axis represents the developmental stages,
and the y-axis represents the median STD value.

Consistent with pupal samples being closer to embryos than to larvae in the same
species, *D. melanogaster* embryos were closer to *D. virilis* pupae than to larvae, and
vice versa, but again, with a distinct expression profile prominent at the P4 stage.

The main diagonal of the heatmap in Figure 2a represents the correlation between the
same stages during development and is shown separately as Figure 2b. The strongest
correlation was observed at the P3 stage, followed by a decrease at the P4 stage and a
subsequent peak at the P5 stage. These observations suggest convergence in
expression profiles during early and late pupal development.

The standard deviation (STD) from the mean expression across replicates was used as
an additional metric of gene expression stability. Genes with more conserved
expression have lower STD indicating more consistent expression across replicates,
while less constrained genes would have STD. The distributions of STD values for each
developmental stage (Figure 2c,d) exhibited dynamics similar to the Spearman
correlation. Variation in gene expression increased at mid-pupae, specifically at the P3
and P4 stages, indicative of decreased constraints on expression during these stages.

Hence the P4 (buoyant) stage featured the least interspecies similarity (among the
pupal stages) with both metrics, correlation coefficient and standard deviation.

Clusters of co-expressed genes

To study gene groups with distinct developmental expression patterns, a hierarchical
clustering analysis was performed using the Spearman correlation coefficient as the
distance metric (Fig. 3). To focus on the dynamics of pupae, only pupal stages were
considered at the clustering step, yielding eight clusters.

Clusters were sorted according to the number of common orthologous genes, so that
similar pairs of clusters had the same names in both species (Fig. 3b). In most cases
the median gene trajectories of paired clusters turned out to be highly similar between
the two species, despite independent clustering in the species (Supplementary Fig. 2).

Cluster pair 1 were the largest and had the largest set of common genes and were
composed of genes with expression gradually increasing throughout the lifespan in both
D. melanogaster and *D. virilis*.

The observed similarities demonstrate that orthologous genes tend to have similar
dynamics (statistics provided in Supplementary Fig. 2), supporting the biological
relevance of the clusters. Hence, we selected pairs of clusters exhibiting high
correlation of expression profiles and containing common orthologous genes, and
performed GO enrichment analysis to reveal the functional role of these clusters.

Expression of genes comprising cluster pair 2 increased during the early stages of
pupal development followed by a sequential gradual decrease. The enriched GO terms
for this cluster were related to the regulation of growth, development and cell migration
(Supplementary Fig. 3). Genes in this cluster pair have a pupae-specific expression,
since in embryogenesis their expression is relatively low.

Genes in cluster pair 3 had higher correlation of gene expression between the species
for all developmental stages compared to the average correlation (Supplementary Fig.
4). Expression of these genes increased during the P5 and mid-pupae stages and also
significantly increased during the embryonic stage. This suggests that these genes are
involved in both embryogenesis and metamorphosis transcriptomic programs. The GO
enrichment analysis revealed enrichment of terms related to chromatin organization, cell
cycle, and developmental processes for these gene clusters.

GO enrichment results for other cluster pairs with a significant number of intersecting
genes and statistically reliable outcomes are given in Supplementary Fig. 5-8.

Figure 3. Clustering of expression profiles. **(a)** Developmental expression profiles for each of the constructed clusters. Each gene is represented by a gray line, and the median expression profile for each cluster is shown in red. **(b)** Heatmap illustrating the intersection of gene sets in each cluster pair. Each square represents a cluster pair, with the color scheme shown at the right; lighter color indicates a higher ratio of intersecting genes.

Gene age influence on the developmental transcriptome

To gain insights into the evolutionary origin of the observed transcriptome dynamics, the phylostrata analysis was performed. Each gene was assigned to a phylostratum corresponding to its predicted time of emergence (for details, see *Materials and Methods* section).

Genes belonging to different phylostrata exhibit specific temporal dynamics of expression, as shown in Figure 4. The most ancient genes (gray) tend to have higher expression during the larval stage, suggesting their involvement in housekeeping functions associated with larval feeding and growth. Intermediate-aged genes (blue), common to *Arthropoda*, *Pancrustacea*, and *Insecta*, have an additional prominent expression peak during the P4 stage. Following the trend, the group of youngest, *Holometabola*-specific genes has a single peak during the P4 stage, indicating that these newly acquired genes could have been selected in *Holometabola* to perform metamorphosis-related functions.

Many young genes up-regulated during the P4 stage (mid-pupal stage) encode proteins with candidate signal peptides that control protein secretion (see *Materials and Methods* for the details). However, signal peptides tend to appear in young genes in general (Supplementary Fig. 9). The distributions of all genes and signal peptide-encoding genes across the phylostrata are statistically different (p-value = 8.3×10^{-75}, statistic = 384.2, the Chi-squared test), P4-specific genes follow the same trend with no observed statistically significant difference (the Chi-squared statistic = 0.2).

Figure 4. Phylostrata-specific gene dynamics. Horizontal axis — the developmental stage, vertical axis — the median log-transformed expression. The shaded area represents the standard deviation (STD) of gene expression. The oldest phylostrata (see the legend) are shown in gray, middle phylostrata, in blue, and the youngest ones, in green.

Genes from different phylostrata exhibit varying impact on the overall similarity of transcriptomes during metamorphosis, as seen in the plots of correlation between gene expression at each stage, calculated separately for each phylostratum (Fig. 5). For intermediate-aged genes (Fig. 5a, blue), the correlation between expression in the two species was larger at the later pupal stages. Hence, the expression of these genes is conserved closer to the end of the pupal development. Conversely, younger genes (Fig. 5a, green) exhibited higher correlation between species primarily at the onset of the pupal development, when imaginal disks are everted and active differentiation occurs.

Looking at another angle, that is, comparing expression of individual genes throughout development (Fig. 5b, c), we observed that recently emerged genes from *Diptera* and *Drosophila* phylostrata tended to have higher correlation between species (p-values for the one-sided Mann–Whitney U test are $9.7×10^{-3}$ and $1.2×10^{-5}$, respectively). This suggests that either these genes had less evolutionary time to diverge, or their function is crucial for pupae, leading to negative selection on changes in expression. In contrast, genes belonging to the *Pancrustacea* phylostratum exhibited statistically lower correlation than the average (p-value equal to $4×10^{-6}$).

The median correlation coefficient for genes comprising different phylostrata tends to decrease with the decrease of the predicted age, see old (grey) and middle (blue) phylostrata in Fig. 5c. However a breakpoint is observed after the transition from *Pancrustacea* to *Insecta* and this tendency is reversed, so that the younger the phylostratum (green in Fig. 5c), the higher median correlation between the species it has. The most similar expression throughout the development was observed for the oldest and the youngest genes.

Clusters 4 and 6 (with increased expression at mid-pupae) were enriched with both young, *Drosophila*-specific genes), genes from the *Insecta* phylostratum, and old genes (Fig. 6). Genes in cluster 4 had an expression burst at the P4 stage, while cluster 6 featured higher expression at the early pupal and adult stages. Cluster 5 consisting of genes expressed at early pupal stages was dominated by middle phylostrata. Cluster 1, characterized by a constantly increasing expression profile throughout the development, was strongly enriched with oldest and Bilateria genes.

Figure 5. Phylostrata-based analysis of conservation of gene expression. **(a)** Correlation of gene expression profiles in two species, calculated separately for each phylostratum. Each line represents an average correlation coefficient between species with the shaded areas showing the standard deviation across replicates. **(b)** Density plot for the Spearman correlation coefficients for gene expression in two species across development, calculated separately for each phylostratum. The density plot for all genes is shown as the shaded area. **(c)** Median values for the distribution of correlation coefficients from (b). The color code as in Figure 4.

Figure 6. Intersection between phylostrata and clusters. The color of the square corresponds to the significance of the intersection (see *Materials and Methods* for details).

Discussion

The hourglass model of gene expression, characterized by a phase of higher difference in gene expression at the beginning and end of development with a period of higher conservation in the middle, phylotypic, stage has been well-established in the context of embryogenesis (Irie and Kuratani 2014; Drost et al. 2017). This pattern has been documented across a wide range of species ranging from insects and nematodes to mammals (Kalinka et al. 2010; Levin et al. 2012; Hazkani-Covo, Wool, and Graur 2005). Furthermore, it may extend beyond embryogenesis and, in plants, is also observed in processes like floral transition (Drost et al. 2016), suggesting that this rule has broader implications than initially thought. One hypothesis has been that the pupal development of holometabolous insects, when organogenesis happens, adheres to a similar hourglass-like pattern. This was based on our observation that pupae tend to recapitulate embryogenesis transcription patterns (Ozerova and Gelfand 2022). The expectation was that a phase of increased similarity in gene expression might occur during the mid-pupal stages.

Contrary to the classic hourglass model, our findings indicate that while there are stages of conserved gene expression, they predominantly occur at the beginning of pupal development, rather than the middle (Fig. 2). Moreover, we observe that mid-pupae P4 stage is the most divergent stage in terms of gene expression between species, as measured by both correlation coefficients and standard deviation (Fig. 2a,b). A previous microarray study also identified the P4 stage as being the least similar to the embryonic stages compared to other pupal stages in *D. melanogaster* (Arbeitman et al. 2002). This suggests a spindle-like pattern, where the transcriptome divergence is accentuated during this phase. This finding underscores the dynamic nature of gene expression during pupal transformation, likely driven by the relatively recent emergence of the latter.

An intriguing observation is that genes expressed during the P4 stage, a period characterized by high divergence, tend to be evolutionarily new. For example, clusters 4 and 6, that have increased expression on this stage, are enriched with young genes (Fig. 6). Moreover, while being relatively highly expressed, these recently emerged genes (specific for *Diptera* and *Drosophila*) demonstrate rather low correlation in expression at the P4 stage compared to the adjacent stages (Fig. 5a).

Remarkably, we find that genes with a middle predicted age (emerging in *Arthropoda* and *Pancrustacea*) have conserved expression patterns in the studied *Drosophila* species during later pupal stages (Fig. 5a). The expression range of these genes remains conserved, suggesting that they did not diverge after the emergence.

Our study provides insights into the transcriptional dynamics of Drosophila metamorphosis, shedding light on the temporal patterns of gene expression during this critical developmental process. However, delving deeper into the temporal dynamics by analyzing additional stages could offer a more comprehensive understanding of metamorphosis. By including more initial and intermediate stages, we could capture finer transitions and potentially identify novel regulatory mechanisms underlying metamorphic events. On the other dimension, analyzing a wider range of *Drosophila*

species would enable us to refine the timing of key pupal developmental stages and enhance the statistical power of our comparative analyses.

Additionally, expanding the scope of comparisons beyond *Drosophila* species — studying insects with mobile pupae, such as mosquitoes, presents a unique opportunity to investigate the preservation of larval structures during pupal development. Unlike *Drosophila*, whose immobile pupae undergo extensive tissue remodeling, insects with mobile pupae should retain larval organs due to the need for locomotion and feeding during the pupal stage. Investigating the molecular mechanisms and gene expression dynamics underlying these differences between species with fixed and mobile pupae could uncover fundamental insights into the evolutionary diversity of metamorphic strategies across insects.

Acknowledgements

This study was supported by the Russian Science Foundation via grant 23-14-00136.

We acknowledge the Skoltech Genomics Core Facility for library preparation and sequencing supported by the Skoltech Center of Molecular and Cellular Biology grant for the students' projects.

Experiments were conducted using the equipment of the Core Centrum of the Institute of Developmental Biology RAS, supported by the IDB RAS research program 0088-2024-0010.

We also acknowledge Mertsalov IB for technical assistance in the collection of specimens.

Data availability

The RNA sequencing data that support the findings of this study are available in the NCBI GEO data repository under the accession GSE269951.

Competing interests

The authors declare no competing interests.

Author contributions statement

Ozerova AM* - Data curation, Formal analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – original draft

Kulikova DA - Investigation, Methodology, Writing – review & editing

Evgen'ev MB - Resources, Supervision, Writing – review & editing

Gelfand MS - Conceptualization, Funding acquisition, Methodology, Project administration, Resources, Supervision, Validation, Writing – review & editing

Arbeitman, Michelle N., Eileen E. M. Furlong, Farhad Imam, Eric Johnson, Brian H. Null, Bruce S. Baker, Mark A. Krasnow, Matthew P. Scott, Ronald W. Davis, and Kevin P. White. 2002. "Gene Expression during the Life Cycle of Drosophila Melanogaster." *Science* 297 (5590): 2270–75.

Bainbridge, S. P., and M. Bownes. 1981. "Staging the Metamorphosis of Drosophila Melanogaster." *Journal of Embryology and Experimental Morphology* 66 (December): 57–80.

Domazet-Loso, Tomislav, Josip Brajković, and Diethard Tautz. 2007. "A Phylostratigraphy Approach to Uncover the Genomic History of Major Adaptations

in Metazoan Lineages." *Trends in Genetics: TIG* 23 (11): 533–39.

Drost, Hajk-Georg, Julia Bellstädt, Diarmuid S. Ó'Maoiléidigh, Anderson T. Silva, Alexander Gabel, Claus Weinholdt, Patrick T. Ryan, et al. 2016. "Post-Embryonic Hourglass Patterns Mark Ontogenetic Transitions in Plant Development." *Molecular Biology and Evolution* 33 (5): 1158–63.

Drost, Hajk-Georg, Philipp Janitza, Ivo Grosse, and Marcel Quint. 2017. "Cross-Kingdom Comparison of the Developmental Hourglass." *Current Opinion in Genetics & Development* 45 (August): 69–75.

Duboule, D. 1994. "Temporal Colinearity and the Phylotypic Progression: A Basis for the Stability of a Vertebrate Bauplan and the Evolution of Morphologies through Heterochrony." *Development . Supplement*, 135–42.

Hazkani-Covo, Einat, David Wool, and Dan Graur. 2005. "In Search of the Vertebrate Phylotypic Stage: A Molecular Examination of the Developmental Hourglass Model and von Baer's Third Law." *Journal of Experimental Zoology. Part B, Molecular and Developmental Evolution* 304 (2): 150–58.

Irie, Naoki. 2017. "Remaining Questions Related to the Hourglass Model in Vertebrate Evolution." *Current Opinion in Genetics & Development* 45 (August): 103–7.

Irie, Naoki, and Shigeru Kuratani. 2014. "The Developmental Hourglass Model: A Predictor of the Basic Body Plan?" *Development* 141 (24): 4649–55.

Irmler, Ingo, Kai Schmidt, and J. Matthias Starck. 2004. "Developmental Variability during Early Embryonic Development of Zebra Fish, Danio Rerio." *Journal of Experimental Zoology. Part B, Molecular and Developmental Evolution* 302 (5): 446–57.

Kalinka, Alex T., Karolina M. Varga, Dave T. Gerrard, Stephan Preibisch, David L. Corcoran, Julia Jarrells, Uwe Ohler, Casey M. Bergman, and Pavel Tomancak. 2010. "Gene Expression Divergence Recapitulates the Developmental Hourglass Model." *Nature* 468 (7325): 811–14.

Kaur, Prameet, Helen Jingshu Jin, Jay B. Lusk, and Nicholas S. Tolwinski. 2018. "Modeling the Role of Wnt Signaling in Human and Drosophila Stem Cells." *Genes* 9 (2). https://doi.org/10.3390/genes9020101.

Klopfenstein, D. V., Liangsheng Zhang, Brent S. Pedersen, Fidel Ramírez, Alex Warwick Vesztrocy, Aurélien Naldi, Christopher J. Mungall, et al. 2018. "GOATOOLS: A Python Library for Gene Ontology Analyses." *Scientific Reports* 8 (1): 10872.

Levin, Michal, Tamar Hashimshony, Florian Wagner, and Itai Yanai. 2012. "Developmental Milestones Punctuate Gene Expression in the Caenorhabditis Embryo." *Developmental Cell* 22 (5): 1101–8.

Mayshar, Yoav, Ofir Raz, Saifeng Cheng, Raz Ben-Yair, Ron Hadas, Netta Reines, Markus Mittnenzweig, et al. 2023. "Time-Aligned Hourglass Gastrulation Models in Rabbit and Mouse." *Cell* 186 (12): 2610–27.e18.

Micchelli, Craig A., Lisa Sudmeier, Norbert Perrimon, Shan Tang, and Ryan Beehler-Evans. 2011. "Identification of Adult Midgut Precursors in Drosophila." *Gene Expression Patterns: GEP* 11 (1-2): 12–21.

Mora, Camilo, Derek P. Tittensor, Sina Adl, Alastair G. B. Simpson, and Boris Worm. 2011. "How Many Species Are There on Earth and in the Ocean?" *PLoS Biology* 9 (8): e1001127.

O'Grady, Patrick M., and Rob DeSalle. 2018. "Phylogeny of the Genus Drosophila."
Genetics 209 (1): 1–25.
Ozerova, Alexandra M., and Mikhail S. Gelfand. 2022. "Recapitulation of the Embryonic
Transcriptional Program in Holometabolous Insect Pupae." *Scientific Reports* 12
(1): 17570.
Raff, Rudolf A. 2012. *The Shape of Life: Genes, Development, and the Evolution of*
Animal Form. University of Chicago Press.
Richardson, M. K. 1995. "Heterochrony and the Phylotypic Period." *Developmental*
Biology 172 (2): 412–21.
Schultheis, Dorothea, and Manfred Frasch. 2023. "Longitudinal Visceral Muscles in
Drosophila Fully Dedifferentiate and Fragment prior to Their Reestablishment
during Metamorphosis." *microPublication Biology* 2023 (March).
https://doi.org/10.17912/micropub.biology.000756.
Supek, Fran, Matko Bošnjak, Nives Škunca, and Tomislav Šmuc. 2011. "REVIGO
Summarizes and Visualizes Long Lists of Gene Ontology Terms." *PloS One* 6 (7):
e21800.
Ten Brink, Hanna, André M. de Roos, and Ulf Dieckmann. 2019. "The Evolutionary
Ecology of Metamorphosis." *The American Naturalist* 193 (5): E116–31.
Teufel, Felix, José Juan Almagro Armenteros, Alexander Rosenberg Johansen, Magnús
Halldór Gíslason, Silas Irby Pihl, Konstantinos D. Tsirigos, Ole Winther, Søren
Brunak, Gunnar von Heijne, and Henrik Nielsen. 2022. "SignalP 6.0 Predicts All
Five Types of Signal Peptides Using Protein Language Models." *Nature*
Biotechnology 40 (7): 1023–25.
Texada, Michael J., Takashi Koyama, and Kim Rewitz. 2020. "Regulation of Body Size
and Growth Control." *Genetics* 216 (2): 269–313.
Yi, Peng, Zhe Han, Xiumin Li, and Eric N. Olson. 2006. "The Mevalonate Pathway
Controls Heart Formation in Drosophila by Isoprenylation of Ggamma1." *Science*
313 (5791): 1301–3.