Lack of an atypical PDR transporter generates an immunogenic *Cryptococcus neoformans*

strain that drives a dysregulated and lethal immune response in murine lungs

Christopher J. Winski[1,#], Peter V. Stuckey[1], Armando M. Marrufo[1], Georgina Agyei[1,$], Robbi L.

Ross[1,2,$], Tamanna Urmi[1,$], Sarah Chapman[5], and Felipe H. Santiago-Tirado[1,2,3,4,*]

[1] Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana, USA

[2] Integrated Biomedical Sciences, University of Notre Dame, Notre Dame, Indiana, USA

[3] Eck Institute for Global Health, University of Notre Dame, Notre Dame, Indiana, USA

[4] Warren Center for Drug Discovery, University of Notre Dame, Notre Dame, Indiana, USA

[5] Integrated Imaging Facility, University of Notre Dame, Notre Dame, Indiana, USA

Running Head: PDR transporter's role during cryptococcal infection

[#]Current address: Laboratory of Virology, National Institute of Allergy and Infectious Diseases,

National Institutes of Health, Hamilton, MT, USA

[$]Equal contributions, name order determined alphabetically

*Address correspondence to Felipe H. Santiago-Tirado, fsantiago@nd.edu

Word Counts

Abstract: 236

Importance: 149

Text (excluding Methods, References, and captions): 6,785

ABSTRACT

Cryptococcus neoformans is an opportunistic fungal pathogen responsible for >150,000 deaths every year with a mortality rate as high as 81%. This high medical burden is due, in part, to an incomplete understanding of its pathogenesis. In a previous study, we identified a cryptococcal atypical ATP-binding cassette (ABC) pleiotropic drug resistance (PDR) transporter, *PDR6*, that affected antifungal resistance and host interactions. Here, we follow-up on the role of *PDR6* in cryptococcal virulence. *In vivo*, mice infected with the *pdr6Δ* strain display altered symptomatology and disease progression. Specifically, we observed a significant increase in the innate immune cell populations in the *pdr6Δ*-infected mice when compared to their WT-infected littermates. Furthermore, quantification of pulmonary cytokines/chemokines revealed a robust increase of pro-inflammatory cytokines in mice infected with the *pdr6Δ* mutant strain. Despite the documented sensitivity of the *pdr6Δ* strain to azole antifungal drugs, the treatment of *pdr6Δ*-infected animals with antifungals did not affect survival, yet treatment with a corticosteroid significantly extended survival, highlighting the importance of a balanced/controlled host immune response. Results with mice that mount opposing immune responses supports out hypothesis that the *pdr6Δ* strain induces a hyper-inflammatory immune response, and that the mice succumb to immune-dependent tissue damage rather than the fungal burden. This altered immune response is driven, in part, by changes in the mutant's surface. Taken together, this study provides insights regarding cryptococcal pathogenesis and highlights additional functions of PDR-type ABC transporters in pathogenic fungi.

IMPORTANCE

Yeasts of the *Cryptococcus* genus, especially *C. neoformans*, can cause disease with unacceptably high mortality. This is due to delays in diagnostics, ineffective treatments, and an incomplete understanding of the interactions between this fungus and our immune system. In this study, we expand our knowledge of the biological function of the *PDR6* gene, particularly its effect on modulating the host's immune response. Normally, *C. neoformans*'s infections are characterized by an anti-inflammatory response that is unable to control the yeast. In the absence of *PDR6*, the response to the infection is a dysregulated pro-inflammatory response that initially controls the fungi but eventually results in death of the host due to too much tissue damage. This is due, in part, to an altered fungal surface. Given the dual role of *PDR6* in modulating antifungal sensitivity and immune responses, this work provides important insights that may lead to new or improved therapeutics.

INTRODUCTION

Commonly associated with soil and bird guano, *Cryptococcus neoformans* is an environmental, opportunistic yeast responsible for >194,000 infections every year (1). Upon inhalation of spores or desiccated yeast, *C. neoformans* reaches the alveoli where it encounters alveolar macrophages and other host immune cells (2); however, this fungus is capable of establishing a unique intracellular niche that allows it to survive and replicate within these host cells (3). Under certain immune conditions, this neurotropic fungus can then escape the lungs, traverse the blood-brain-barrier, and enter the central nervous system (CNS), resulting in cryptococcal meningoencephalitis (CME) with a mortality rate as high as 81% (1, 4, 5). Several lines of evidence indicate that in addition to its use of host immune cells as replication havens, *C. neoformans* can use infected cells as vehicles for dissemination and brain invasion (5, 6). Hence, understanding this complex cryptococcal-phagocyte interaction from both fungal and host perspectives represents a challenging but clinically relevant avenue of research given its role in determining the course and outcome(s) of infection.

In addition to its ability to survive intracellularly, *C. neoformans* has several other virulence factors including an antiphagocytic polysaccharide capsule that provides protection from host stressors and masks several cryptococcal pathogen-associated molecular patterns (PAMPs) including chitin, chitosan, mannoproteins, and β-glucans from host phagocytes (7). The capsule, structurally composed of glucuronoxylomannan (GXM) and glucuronoxylomannogalactan (GXMGal), is constitutively secreted/shed and is known to modulate the host immune response in ways that promote infection (8). Additionally, *C. neoformans* synthesizes cell wall-localized melanin that protects the yeast from ultraviolet

radiation and oxidative/nitrosative stress (9). Finally, the fungus displays thermotolerance, able

to grow at physiological-relevant conditions (37°C + 5% CO_2) (10-12). These virulence factors are

integral to the pathogenesis of *C. neoformans* and as such, have been well-studied. However, *C.*

neoformans utilizes additional virulence factors which remain significantly understudied or

completely unknown. For example, we previously identified *PDR6* (also known as *AFR3*), a novel

ATP-binding cassette (ABC) transporter of the pleiotropic drug resistance (PDR) class, and

showed that it had a role in various cryptococcal virulence facets (13, 14). Present only in plants

and fungi, PDR transporters hydrolyze ATP to transport substrates across biological membranes

and have been primarily implicated in antifungal resistance (15, 16). Indeed, deletion of *PDR6*

(*pdr6Δ*) results in hyper-susceptibility to the azole-class of antifungals, but also showed *in vitro*

and *in vivo* defects in virulence. Thus, the role of *PDR6* in fungal pathogenesis and modulating

the host immune response warranted further investigation.

During microbial infections, the host's immune system is tasked with mounting an

appropriate and balanced immune response that will clear the infection while ideally limiting

tissue damage (17). An exaggerated Th1, or pro-inflammatory, response will likely clear a

microbial infection but also result in excessive cellular and tissue damage. Opposingly, an

inappropriate Th2, or anti-inflammatory, response will limit host damage but allow for

uncontrolled microbial proliferation and dissemination. Many pathogens take advantage of this

dichotomy to promote their infection. For example, early in HIV infection, this virus promotes

the secretion of pro-inflammatory cytokines IFN-γ and TNF-α, generating a strong Th1 response

that is permissive to viral replication (18, 19). In the context of *C. neoformans* infections, the

fungus promotes an anti-inflammatory Th2 response that allows for intracellular survival and

increased replication and dissemination, ultimately resulting in patient mortality (20, 21). It is

evident, however, that between these two immune response extremes, there is a whole

spectrum of responses to microbial infections, making this topic extremely complex but also an

important target for therapeutics. Hence, increased understanding of the immune response

against *Cryptococcus* can inform future therapeutic strategies where the immune response

could be modulated to benefit the host.

Our previous work characterizing *PDR6* showed that it had important roles in

cryptococcal biology and suggested that it was necessary for full virulence (13). Here, we study

the role of *PDR6* in the establishment and progression of disease using a murine inhalation

model of cryptococcosis. We show that a lower inoculum or treatment with antifungal drugs,

significantly decreased *pdr6Δ* fungal burdens, but do not change the outcome of the infection

by this mutant. Instead, the survival studies and immunological analyses suggest that the *pdr6Δ*

mutant elicits an altered host immune response characterized by significant increases in pro-

inflammatory cytokines, chemokines and innate immune cells. We treated the *pdr6Δ*-infected

mice with a corticosteroid and significantly extended their survival despite little to no change in

fungal burden. Moreover, using mouse breeds that are known to establish opposing immune

responses, we show that indeed the host's immune response is the main determinant of the

pdr6Δ infection outcome. This dysregulated immune response might be a consequence of

fungal surface alterations that expose antigenic and immunogenic components that normally

would be hidden from the immune system. Collectively, this study furthers our understanding

of cryptococcal pathogenesis, highlights the clinical importance of the host immune response in

controlling fungal infections, while also providing novel functions and insights of fungal PDR-

type transporters in the context of disease progression.

RESULTS

***pdr6Δ*-infected mice succumb to infection despite decreased inoculum and fungal burden.** In

our previous study we determined that *PDR6* was required for complete cryptococcal virulence

as mice infected with the *pdr6Δ* mutant lived significantly longer and had decreased fungal

burdens when compared to their H99-infected littermates (13). Consistently, this finding was

recapitulated by *Oliveira et al.* using the *Galleria mellonella* invertebrate infection model (14).

Notably, the *pdr6Δ*-infected animals displayed altered symptomatology compared to their H99-

infected littermates. Specifically, at endpoint, the WT-infected mice displayed altered behavior

associated with neurological deficiencies, traditional of cryptococcal meningitis; in contrast, the

pdr6Δ-infected mice exhibited respiratory symptoms characterized by rapid and labored

breathing, suggestive of pneumonia. We hypothesized that the altered symptomatology could

be explained, in part, by differences in the establishment and progression of disease.

Previously, fungal burdens were only quantified at time of death (TOD), potentially overlooking

differences earlier on in the infection. To obtain a comprehensive view of the infection over

time, we inoculated mice with 5 x 10^4 cells of H99 or *pdr6Δ* strains, and quantified fungal

burden every seven days over the course of the infection (Fig. 1A). Interestingly, we found that

at 7 days post infection (dpi) there were no differences in the fungal burdens between the H99

and *pdr6Δ*-infected animals in all organs tested, suggesting that the strains established infection

similarly. Following this initial establishment, however, at 14 dpi there was a significant

decrease in fungal burden for the $pdr6\Delta$ strain compared to the WT. At 17 dpi, the H99-infected

mice succumbed to the infection with average lung, brain, and spleen burdens of 3.7 x 10^8, 1.9 x

10^6, and 1.3 x 10^4 CFU while the $pdr6\Delta$-infected mice CFUs were 3.6 x 10^7, 5.1 x 10^3, and 677,

respectively. Between 14 and 28 dpi, the fungal burden of the $pdr6\Delta$-infected animals remains

relatively constant in the brain and spleen, with a gradual, although small, increase in the lungs.

Then, after 28 dpi, there are gradual increases in all organs with final average CFU counts of 5.5

x 10^7, 1.1 x 10^5, and 1.5 x 10^3 in the lungs, brain, and spleen, respectively. The differences in

CFUs highlight key differences between H99 and $pdr6\Delta$ disease progression and suggests a

potential role of the adaptive immune response given the changes at 14 dpi.

Experimentally, the mice in our previous studies were inoculated with 5 x 10^4 cells, a

common inoculum for the murine inhalation model of cryptococcosis. Given the *in vitro* defects

of the $pdr6\Delta$ strain (13), and the fact that initially, the mice were able to control the infection

(Fig. 1A), we hypothesized that by lowering the fungal inoculum, the survival of the $pdr6\Delta$-

infected mice would be extended with a potential for complete clearance of the infection. To

assess this, we decreased the initial inoculum 10-fold and infected mice with 5 x 10^3 cells and

monitored survival (Fig. 1B). The H99-infected mice had a mean survival of 21 days and the

$pdr6\Delta$-infected mice displayed a mean survival of 36 days, similar to our studies with 10-fold

higher inoculum. Quantification of the fungal burden showed similar CFUs in all organs

between the animals infected with low and high WT inoculum (Fig. 1C compared to 1A);

however, despite having similar lungs CFUs, the animals infected with low $pdr6\Delta$ inoculum had

lower brain and spleen CFUs at TOD relative to the higher inoculum (Fig. 1C compared to 1A).

Evidently, using a lower inoculum did not change the trend we have seen before: the $pdr6\Delta$-

infected mice still survive longer with significant decreases in fungal burden across all organs.

However, notably, in the *pdr6Δ*-infected mice the lower inoculum results in lower overall organ

burdens, with less dissemination, yet the *pdr6Δ*-infected mice still succumbed to the infection

with the same dynamics as when using a higher inoculum (35-38 days; Fig. 1B).

Infection with *pdr6Δ* results in increased tissue damage. Despite initially colonizing the

pulmonary environment, the *Cryptococcus*-associated fatal pathology occurs in the brain due to

the meningoencephalitis (22, 23). The altered symptomatology coupled with the differential

disease progression suggests that the *pdr6Δ* strain likely affects the host immune response.

Furthermore, when performing the previous fungal burden experiments, we observed

significant phenotypic differences in lung sizes in H99 v. *pdr6Δ*-infected mice (Fig. 2). To

qualitatively assess any potential differences at the cellular and tissue level, we harvested the

lungs of mock, H99, and *pdr6Δ*-infected mice, stained with hematoxylin and eosin (H&E), and

assessed tissue damage (Fig. 2 and Table 1). At 14 dpi, the gross anatomical images clearly show

increased size and inflammation of the H99-infected lungs when compared to the *pdr6Δ*-

infected. Additionally, the 10X and 40X images display deterioration of the lung parenchyma

and infiltration of *C. neoformans* which is not observed in the *pdr6Δ*-infected lungs (Fig. 2A). At

35 dpi, the *pdr6Δ*-infected lungs are severely inflamed, infiltrated with yeast, and characterized

by the presence of many inflammatory immune cells in the peri-bronchial and perivascular

space (Fig. 2B). Table 1 provides a quantitative summary comparing the H99 and *pdr6Δ*-infected

pulmonary tissue over time. Consistent with our observations in the fungal burden

experiments, there were little to no differences at 7 dpi. However, at 14 dpi the *pdr6Δ*-infected

lung contains the yeast in granuloma-like regions and have 30% less *C. neoformans* than the

WT-infected tissues. Furthermore, an increase in lymphocytes is observed in the *pdr6Δ*-infected

lungs relative to WT (105 to 26, respectively; Table 1). Finally, as the *pdr6Δ*-infected mice

approach TOD, the fungal infiltration, inflammation, and number of inflammatory immune cells

in the lung tissue all increase to levels well above those observed in the WT-infected animals at

TOD. Altogether, these findings suggest that the *pdr6Δ* strain elicits a differential immune

response in the pulmonary environment thus affecting disease progression and

symptomatology.

Fluconazole treatment improves fungal burden but not outcome in *pdr6Δ*-infected mice.

Currently, there a very few antifungal drugs available to treat cryptococcal infections and the

compounds that are available are either not effective, extremely expensive, or associated with

deleterious side effects (24). Fluconazole (FLC), a fungistatic azole class antifungal, is the

mainstay treatment option in low-resource regions where cryptococcal infections are endemic.

Mechanistically, FLC inhibits lanosterol 14α-demethylase, an enzyme in the ergosterol

biosynthesis pathway encoded by *ERG11* (25). In our previous study, we showed that the

deletion of *PDR6* increased the susceptibility of *C. neoformans* to azole class antifungals

including FLC (13). Given this phenotype, we hypothesized that treatment of *pdr6Δ*-infected

mice with FLC would result in total clearance of the infection, preventing the lethal immune

pathology. To test this, mice infected with either the WT or *pdr6Δ* strains, or mock infected,

were treated daily (starting 1 dpi) with either a low (8 mg/kg) or high dose (16 mg/kg) of FLC

and monitored for survival (Fig. 3A). The FLC treatment significantly extended survival in both

the H99 and *pdr6Δ*-infected animals compared to the untreated controls, albeit to the same

extent. Numerically, the mean survival for the untreated H99-infected mice was 17 days and

the low and high doses extended survival 4.5 and 6 days, respectively. Similarly, the mean

survival for the untreated *pdr6Δ*-infected mice was 34 days and the low and high doses

extended survival 5 and 7 days, respectively. Surprisingly, although the survival was similarly

extended between strains, quantification of the fungal burden revealed significant differences

(Fig. 3B-G). At 14 dpi, both doses of the FLC treatment significantly decreased the fungal burden

in the lungs, brain, and spleen only in the *pdr6Δ*-infected mice (Fig. 3B-D). Similarly, at TOD, the

daily FLC treatment resulted in an additional substantial decrease in the fungal burden of the

pdr6Δ-infected mice across all three organs, with several mice where the brain and spleen CFU

counts were below the limit of detection (LOD) (Fig. 3E-G). Not surprisingly, neither of the FLC

doses had a significant effect on WT-infected mice. This observation is interesting because

despite significantly decreasing the fungal burden with the FLC treatment, the *pdr6Δ*-infected

animals still succumbed to the infection.

To understand how the FLC treatment was affecting the pulmonary environment, we

prepared slides and performed qualitative histological analysis at TOD for both WT and *pdr6Δ*-

infected mice (Fig. 4 and Table 2). The H99-infected lungs at 14 dpi treated with FLC displayed

similar features to the previously pictured H99 untreated lungs, namely a high percentage of

fungal infiltration and tissue breakdown. In contrast, there was a clear decrease in fungi in the

FLC-treated *pdr6Δ*-infected lungs at this timepoint. Consistently, the FLC-treated *pdr6Δ*-infected

lungs at 35 dpi appear to have the infection controlled in granuloma-like regions (Fig. 4, inset 2)

with widespread immune cell infiltration (Fig. 4, inset 1). Interestingly, despite reduced

pulmonary CFUs (Fig. 3 and Table 2) and decreased inflammation (Fig. 4 and Table 2), the

pdr6Δ-infected mice were still succumbing to the infection. Together, these data suggest that

the mice infected with the *pdr6Δ*-strain are not necessarily dying directly because of the *C.*

neoformans infection, but rather indirectly because of an inappropriate host immune response.

C. neoformans pdr6Δ strain elicits a strong pro-inflammatory immune response. During

microbial infections, the host's immune response ultimately determines the clinical outcome

(26). Ideally, a balanced Th1 pro-inflammatory and Th2 anti-inflammatory immune response

will allow for microbial clearance and subsequent tissue repair. Previous studies have

determined that *C. neoformans* skews the immune response towards a permissive anti-

inflammatory Th2 response, thus benefiting the yeast and allowing for fungal replication and

subsequent dissemination (27). Given our previous observations, we hypothesize that the

pdr6Δ strain is instead generating a strong pro-inflammatory immune response that is able to

control the infection initially, but also damaging large areas of pulmonary tissue. To assess this,

we quantified the secreted cytokines and chemokines profile in the lungs and brains of H99 and

pdr6Δ-infected mice at their respective TODs (Fig. 5 and Fig. S1). Interestingly, we found that

several of the pro-inflammatory cytokines are significantly upregulated in the *pdr6Δ*-infected

lungs relative to WT (Fig. 5A). Specifically, the production of IL-1α, IL-1β, and MIP-1β (CCL4) is

increased 13-fold, 25-fold, and 37-fold respectively, when compared to the lungs of the H99-

infected mice. Interestingly, the expression of IL-13 and IL-33, anti-inflammatory cytokines

associated with allergic diseases, where also statistically upregulated in the *pdr6Δ*-infected mice

relative to WT. A similar trend was observed in the brain, although relative to the pulmonary

tissues, it was more muted (Fig. 5B), with statistically significant upregulation of G-CSF and IL-3

relative to the H99-infected brain. This weakened expression of inflammatory cytokines in the

brain might be due to the decreased capacity of the *pdr6Δ* mutant to disseminate coupled with

the immune-privileged status of the CNS. During a typical cryptococcal infection, the expression

of anti-inflammatory cytokines and chemokines are increased thus generating a Th2 response,

however, in the context of an infection with the *pdr6Δ* strain, the secretion of pro-inflammatory

cytokines and chemokines is significantly increased resulting in a damaging Th1 immune

response.

Corticosteroid treatment extends survival in *pdr6Δ*-infected mice. Clinically, corticosteroids

like dexamethasone are prescribed to downregulate the immune system. Mechanistically,

these anti-inflammatory drugs function by suppressing the migration of neutrophils and

decreasing lymphocyte colony proliferation (28). The potential of dexamethasone as a

treatment strategy during cryptococcal infection was investigated in a clinical trial but was

prematurely concluded because of adverse effects and increased mortality (29, 30); however,

corticosteroids have been used successfully to treat CME-associated inflammatory disorders

such as postinfectious inflammatory response syndrome and immune reconstitution

inflammatory syndrome (IRIS)(22). Given the robust pro-inflammatory immune response

observed in the *pdr6Δ*-infected mice, akin to fatal IRIS in humans, we hypothesized

dexamethasone treatment would dampen the aggressive Th1 response resulting in fungal

clearance and improved survival. Experimentally, we infected mice with either the H99 or

pdr6Δ strain, or mock infected, and treated them daily (starting at 1 dpi) with either 0.3 mg/kg

dexamethasone or a dexamethasone and 16 mg/kg FLC combination and assessed survival (Fig.

6A). The H99-infected mice that received the dexamethasone treatment displayed a significant

reduction in survival (mean survival of 13 days vs 17 days) when compared to the untreated

H99 control, while the combo therapy restored survival to that of the untreated H99-infected

group (mean survival of 16 days vs 17 days). This result corroborates the corticosteroid finding

reported in the human clinical trial (29, 30). In terms of the $pdr6\Delta$-infected mice, both the

dexamethasone alone and in combination with FLC significantly extended survival. Numerically,

the mean survival of the untreated $pdr6\Delta$-infected control group was 34 days and was extended

to 55 and 58 days in the dexamethasone and combination treated groups, respectively. The

survival studies were stopped after 60 dpi because the remaining mice, including the mock-

infected receiving the daily dexamethasone treatment (not shown), appeared sick despite not

meeting endpoint criteria. We concluded that after two months of daily corticosteroid

injections the immune systems of the mice had been ablated resulting in their poor

appearances. Despite the dexamethasone treatment extending survival in the $pdr6\Delta$-infected

animals, the fungal burdens were differentially affected over the course of the experiment (Fig.

6B-G). Specifically, at 14 dpi the addition of dexamethasone significantly decreased the fungal

burden in the lungs in both the H99 and $pdr6\Delta$-infected mice and little to no changes were

observed in either strain in the brain or spleen CFUs (Fig. 6B-D). At TOD the dexamethasone

treatment in the $pdr6\Delta$-infected animals significantly increased the lung CFUs and the

combination therapy returned it to levels comparable to the untreated $pdr6\Delta$-infected control

(Fig. 6E). Although not statistically significant, a similar trend was observed in the brain and

spleen of the $pdr6\Delta$-infected mice (Fig. 6F & G). These effects were not seen in the WT-infected

mice. The dynamic changes in fungal burdens in the lungs of the dexamethasone and

combination treated *pdr6Δ*-infected mice can be explained by the absence or lack of

inflammatory immune cells to control the infection and the addition of FLC lowers the CFUs as

the fungistatic nature of the antifungal is unable to clear the initial infection. Collectively, these

data further confirm and highlight the altered disease progression elicited by the *pdr6Δ* mutant

and, more importantly, demonstrate the positive clinical outcome associated with controlling

the inappropriate pro-inflammatory immune response.

Combination dexamethasone and fluconazole treatment reduces pulmonary tissue damage in

***pdr6Δ*-infected animals.** Based on our previous results, it is evident that the addition of the

corticosteroid is benefiting the *pdr6Δ*-infected mice by significantly extending survival.

Histological analysis of the lung tissue at TOD for both WT and *pdr6Δ*-infected mice was

performed to visualize changes in fungal infiltration, tissue inflammation, and immune cell

recruitment (Fig. 7 and Table 3). Visually, the treatment with dexamethasone had a profound

effect given the qualitative reduction in inflammatory cells across all conditions. The lungs of

the H99-infected mice that received dexamethasone are not as inflamed as those from

untreated mice but the destruction of the lung tissue is devastating (Fig. 7A). The lack of

immune cells allowed for the uncontrolled proliferation of *C. neoformans* resulting in increased

damage of the pulmonary tissue. Similarly, the dexamethasone treated lungs of the *pdr6Δ*-

infected display intense fungal proliferation but there is qualitatively less damage when

compared to the H99-infected animals (Fig. 7B and Table 3). The combination dexamethasone

and FLC treatment reduced the fungal infiltration in both the H99 and *pdr6Δ*-infected mice

when compared to the dexamethasone alone (Fig. 8 and Table 4). In fact, despite the presence

of some yeast, the *pdr6Δ*-infected lung tissue appears relatively healthy with little to no

damage and inflammation suggesting that another mechanism may be responsible for the mice

reaching endpoint (Fig. 8B and Table 4).

pdr6Δ-infected mice display an increase in innate immune cells populations. During microbial

infections, the cells of the innate immune system represent the first line of defense which is

followed by the activation and recruitment of the adaptive immune response (31). In a *C.*

neoformans infection, lung macrophages represent the innate population primarily responsible

for initial clearance, although some recent literature suggests neutrophils also play a supportive

role (32-34). However, CME is an AIDS-defining disease, and low $CD4^+$ T-cell counts is a major

susceptibility for CME, indicating that the adaptive immune response also plays an important

role controlling cryptococcal infections (35). Provided the altered immune response to the

pdr6Δ strain, we expected alterations to both the quantity and the composition of innate and

adaptive inflammatory immune cells. To test this, we harvested leukocytes from the lungs of

H99 and *pdr6Δ*-infected mice at 14 dpi (close to TOD of WT-infected mice) and at 35 dpi (close

to TOD of *pdr6Δ*-infected mice) and quantified the different populations using flow cytometry

(Fig. 9 and Fig. S2). Quantification of the total innate immune cell populations at 14 dpi showed

that the FLC, dexamethasone, and combination treatments led to a significant decrease in the

pdr6Δ-infected animals whereas there were no significant changes in the H99-infected mice

(Fig. 9A). However, no reduction was observed when compared as a percentage of $CD45^+$ cells

(Fig. 9B), although, interestingly, a significant increase in the ratio of neutrophils and

macrophages was seen in the combination-treated *pdr6Δ*-infected mice (Fig. 9B). At this time

(14 dpi), quantification of the total adaptive immune cells showed a marked increase in the

pdr6Δ-infected animals when compared to their H99-infected counterparts, although only the

CD8$^+$ T-cells were statistically significant (Fig. 9C, untreated bars). Furthermore, all treatments

decrease the number of lymphocytes in the *pdr6Δ*-infected animals relative to the WT, specially

the CD8$^+$ T-cells (Fig. 9C, +FLC, +Dexa, and +Dexa +FLC bars). The same trend is observed when

compared as percentage of CD45+ cells (Fig. 9D). At 35 dpi there are some minor differences in

the adaptive immune cell populations when comparing the treated groups (Fig. 9G & H).

Interestingly, when comparing the untreated *pdr6Δ*-infected mice at 35 dpi with 14 dpi there is

a significant reduction in the total number of innate immune cells (Fig. 9D & H, untreated bars).

Moreover, a significant increase in macrophages is observed in the *pdr6Δ*-infected mice when

compared to the H99-infected (Fig. 9E & F, untreated bars). Using CD68 as a marker of

inflammatory macrophages in histological slices confirmed this influx of immune cells specific

for the *pdr6Δ*-infected (Fig. S3). Notably, the dexamethasone treatment (alone or with FLC) in

the *pdr6Δ*-infected animals significantly decreases the number of macrophages which is

replaced by a significant increase in the number of neutrophils (Fig. 9E & F, +Dexa and +Dexa

+FLC bars). It appears that in the absence of Pdr6, a robust and sustained macrophage response

is mediating the tissue damage, whereas in the WT-infected mice, neutrophils are the main

innate cells accumulating in the lungs. Collectively, these results showed that infection with the

pdr6Δ strain results in the increased recruitment of adaptive immune cells early on during the

infection while an increase in innate immune cells, specifically macrophages, was observed in

the later stages of the disease. This further explains the strong pro-inflammatory response,

altered disease progression, and differential symptomatology specific to the *pdr6Δ* strain.

The magnitude of the host response to *pdr6Δ* strain infection influences the disease outcome.

Up to this point, our results indicate that infection with the *pdr6Δ* strain results in the

recruitment of more inflammatory immune cells to the lungs, that initially can control the

infection but, when sustained over several weeks, leads to increased pulmonary damage and

mortality. This is supported by the dexamethasone treatment experiments, where dampening

the immune response significantly extended survival in the *pdr6Δ*-infected animals. However,

these experiments were done in the A/J mouse background, which is known to be "muted" in

response to cryptococcal infection. We wondered if the same would happen in mice that are

known to mount either strong pro-inflammatory (C57BL/6) or anti-inflammatory (BALB/c)

responses against WT *C. neoformans*. With C57BL/6 mice, we envisioned two potential

outcomes: the already skewed pro-inflammatory response of this mouse in combination with

the pro-inflammatory response induced by *pdr6Δ* could result in complete clearance, or

augment the tissue damage and result in hypervirulence. With the BALB/c mice, we had one

hypothesis: the strong anti-inflammatory response of this mouse could counter the pro-

inflammatory response induced by *pdr6Δ* enough to reach a balance, allowing for clearance or

latency. To test this, we infected either C57BL/6 or BALB/c mice with 5 x 10^4 cells of H99 or

pdr6Δ strains and followed their survival as before. Whenever a mouse reached humane

endpoints, organs were harvested for CFU and cytokine quantification. Interestingly, the

C57BL/6 survival curve was similar to the A/J curve (Fig. 10A), with a median survival time of 18

days for H99-infected mice and 42 days for *pdr6Δ*-infected mice, and decreased *pdr6Δ* burden

in all organs (Fig. 10B − D). Indeed, the cytokine profile of the *pdr6Δ*-infected C57BL/6 was a

more pronounced pro-inflammatory A/J profile, with the same upregulation of IL-1α, IL-1β, and

MIP-1β, but additionally, there was strong upregulation of IL-6, MIP-1α, and G-CSF (Fig. 10E and

Fig. S4). In contrast, 70% of the *pdr6Δ*-infected BALB/c mice survived to the end of the

experiment (60 dpi), whereas all of the H99-infected mice succumbed by day 18 (Fig. 10F).

Surprisingly, the surviving mice had fungal burdens higher than the H99-infected mice at TOD

(Fig. 10G − I), yet they showed no symptoms. Comparison of the immune profiles between

C57BL/6 and BALB/c showed expected similarities (IL-1α, IL-1β, and MIP-1α) but also clear

differences (lack of IL-6 and G-CSF signaling and stronger induction of IL-13 in BALB/c)(Fig. 10E

& J, and Fig. S4). Similarly to the dexamethasone treatment in A/J mice, the *pdr6Δ*-infected

BALB/c mice survived longer despite having higher fungal burdens, strongly suggesting that the

fatal pathology is driven by the immune response and not by the fungus.

C. neoformans *pdr6Δ* strain exhibits altered PAMP exposure. Experimentally, we have studied

the host response to this cryptococcal mutant but the question remains as to how is this

mutant modulating the immune system and generating a robust Th1 response. In our previous

work, we showed that *PDR6* encodes an atypical transporter localized to the endoplasmic

reticulum (ER) and plasma membrane (PM) involved in ergosterol trafficking (13). Others have

suggested a direct relationship between altered ergosterol levels and host immune responses

(36, 37) as well as altered surface composition due to defects in PM fluidity and lipid rafts (38,

39). We hypothesize that in the absence of *PDR6* there would be less ergosterol in the PM,

potentially resulting in downstream alterations to the cryptococcal surface that would expose

immunogenic components. To assess this, we grew H99 and *pdr6Δ* cells in YPD, stained with

fluorescent dyes that recognize and bind to fungal cell wall components, and visualized the cells

using microscopy. Under this condition, we did not observe any difference between the H99

and *pdr6Δ* stained cells (data not shown). However, previous studies have reported a

correlation between growth conditions and the architecture of the fungal cell wall (40). This led

us to repeat the experiment with cells grown under host-like conditions (DMEM at 37°C + 5%

CO_2) (Fig. 11A). Under these conditions, the *pdr6Δ* strain displayed increase staining with

calcofluor white (CFW) and concanavalin A (ConA) suggestive of increased exposure of chitin

and mannoproteins, respectively. Similarly, there was some staining with the β-(1,3)-glucan

antibody suggestive of more exposed β-glucans in the *pdr6Δ* mutant (Fig. 11A). Despite the

differences observed microscopically, quantification of the staining intensity of these dyes, and

of eosin Y (EoY) which stains chitosan, using flow cytometry did not reveal statistically

significant increases in intensity in the *pdr6Δ* strain when compared to the H99 (Fig. 11B – E).

The positive control, *cap59Δ*, deficient in polysaccharide capsule synthesis and thus with

increased exposure of its cell wall, showed the expected increases in intensity. This was

confirmed using three independently made *pdr6Δ* mutants (Fig. 11B – E and Fig. S5). Having

previously shown that the *pdr6Δ* mutant has no defect in capsule induction, we hypothesized

that alterations in the capsular architecture could explain the visually increased exposed

cryptococcal PAMPs. To assess this, we used several anti-polysaccharide capsule antibodies to

stain the capsules of H99, *pdr6Δ*, and *cap59Δ* cells (Fig. 11F). A significant increase in staining

and penetration of the antibody was observed in the *pdr6Δ* mutant when using the mAb 302

(Fig. 11F). This antibody only recognizes and binds acetylated GXM (41) and suggests that the

integrity and/or porousness of the polysaccharide capsule in the *pdr6Δ* strain is altered when

compared to the structurally-sound H99 capsule. This could justify the high-uptake phenotype

of the *pdr6Δ* mutant (13, 42). Lastly, we tested if the *pdr6Δ* mutant could form titan cells using

a recent *in vitro* induction protocol (43), as titan cells are restricted to the lungs, have an

altered cell surface, and are related to latency. Although there was a trend to form less titan

cells, it was not statistically significant relative to H99 (Fig. 11G). Altogether, these findings

suggest that the deletion of *PDR6* disrupts the cryptococcal cell wall composition and the

capsule density (since it is induced to normal size in the *pdr6Δ* mutant), increasing the potential

exposure of fungal PAMPs, which can drive the strong pro-inflammatory response by the host

immune system.

DISCUSSION

Fungal PDR transporters have been primarily studied for their role in antifungal

resistance. For example, the overexpression of *AFR1* in *C. neoformans* and *CDR1* in *C. albicans*

clinical strains increase their resistance to many of the azole class antifungals and complicates

their treatment strategy (44, 45). Structurally, these two proteins (and all other well-

characterized fungal PDR transporters) are 'full-length' consisting of two nucleotide-binding

domains (NBDs) and two transmembrane domains (TMDs), which is required for a functional

fungal PDR protein (15, 16, 46). Here, we continue the characterization of *C. neoformans PDR6*,

an atypical and novel 'half-size' PDR transporter since it only possesses one NBD and one TMD.

Despite this, various lines of evidence suggest it forms a functional transporter (13, 14).

However, we found that in addition to azole resistance, *PDR6* modulates the host immune

response, a phenotype that may be unique to 'half-size' PDR transporters (13).

The *pdr6Δ*-dependent alterations in survival, fungal burden, and symptoms all suggested

a differential host response. Moreover, the fact that despite the significant decrease in fungal

burden across organs, and the use of a lower inoculum, the mice still died, suggesting that the

pdr6Δ-infected mice were not succumbing to uncontrolled fungal growth, but rather to an

inappropriate and damaging immune response. This was further supported by the FLC and

dexamethasone studies. Treatment of the *pdr6Δ*-infected mice with FLC decreased the fungal

burden in all organs, and in some to the limit of detection, and yet the mice still died. In

contrast, but consistent with the immune response as the culprit, dexamethasone treatment in

the *pdr6Δ*-infected mice significantly extended their survival despite having fungal burdens that

were higher than the WT-infected mice at TOD. Lastly, the divergence in symptoms, histological

analysis, immune cell quantification, and cytokine/chemokine expression profiles, all were

consistent with an aggressive lung pro-inflammatory immune response against the *pdr6Δ*

mutant, resulting in their death by pneumonia rather than by CME. We recognize, however,

that the *pdr6Δ*-infected mice still had CFUs in the brain, so CME cannot be ruled out, but we

believe that it was below the threshold to trigger the neurological symptoms characteristic of

CME. This is supported by the "muted" immune response in the brain tissue, indicative of little

inflammation, and by the fact that when treated with FLC, there was minimal extrapulmonary

dissemination but no change in the outcome of the infection. If we consider that the *pdr6Δ*

mutant grows slower under host-like conditions (13), is recognized more avidly by THP-1 cells

(alveolar macrophages)(13, 42), and has reduced intracellular survival (13, 14), then the

massive lung tissue damage seen in the *pdr6Δ*-infected mice at TOD has to be due to an

inappropriate, dysregulated, Th1 immune response.

Although treatment with a corticosteroid supported our hypothesis, we acknowledge

there are some limitations. Dexamethasone is a strong immunosuppressant, dampening the

immune response in the whole body, especially given the length of the study. To address that,

we purposefully used doses that corresponded to less than those used in human clinical trials

(29). This extended treatment might explain why, even in the combo treatment with FLC, which

reduced both inflammation and fungal burden in the lungs and restored tissue architecture

almost to normal, the mice appeared so sickly. Optimizing the timing of the dexamethasone

treatments, or employing ways to target the drug specifically to the lungs, as with the use of an

inhaler, might results in improved outcomes and represents a future direction of investigation.

However, we reasoned that we could counter the pro-inflammatory response induced by the

pdr6Δ mutant using mouse breeds that naturally mount anti-inflammatory, or Th2-skewed,

responses to infection, such as the BALB/c mouse. As a control, we also infected C57BL/6 mice,

which are known to be Th1-skewed in their immune responses. Again, consistent with our

hypothesis, 70% of the BALB/c mice survived until the experimental endpoint at 60 dpi despite

having significantly higher lung fungal burdens at this timepoint compared to the H99-infected

mice at TOD.

The *in vivo* experimentation allowed us to comprehend what was occurring within the

host, but why was this specifically occurring with the *pdr6Δ* strain? Our data showing that the

pdr6Δ mutant displays altered PAMP exposure is a reasonable explanation for the pro-

inflammatory response. However, it may not be the only reason as the acapsular *cap59Δ*

mutant also exhibits increased PAMP exposure and it is avirulent in the same mouse model.

Furthermore, we previously showed that conditioned media (CM) from *pdr6Δ* cultures

modulates the phagocytosis of *C. neoformans* by macrophages, whereas the CM from *cap59Δ*

or other PDR transporter mutants had little to no effect (13), suggestive of an

immunomodulatory Pdr6-dependent secreted factor. Furthermore, other cryptococcal mutants

with altered PAMP exposure have been reported but none of them induce an immune response

as that seen here with the *pdr6Δ* mutant (47-50). The *mar1Δ* mutant has a disorganized cell

wall with increased chitin exposure, less glucans and mannans, and little capsule (49). Like our

case, the *mar1Δ* mutant causes a protracted death in C57BL/6 but shows no mortality in BALB/c

(47). However, in the *mar1Δ*-infected BALB/c mice, the immune response was controlled in the

form of GM-CSF-dependent granulomas and there was no overt pro-inflammatory response,

which is different from *pdr6Δ*-infected mice. The *pdr6Δ*-induced mortality is more similar to

that seen with the *usv101Δ* strain, which causes fatal pneumonia in A/J mice with little to no

extrapulmonary dissemination (50). However, pathologically, the *usv101Δ* mutant did not

generate granulomas like the *pdr6Δ* mutant, and in our case, we see much more

extrapulmonary dissemination than that with the *usv101Δ* mutant. The *usv101Δ* strain has a

larger capsule than WT, thus it might not be exposing the same antigens as the *pdr6Δ* (which

has a normal-sized capsule), generating a muted immune response relative to the response

induced by *pdr6Δ*. Lastly, the *rim101Δ* mutant also induces a strong pro-inflammatory response

in A/J mice as a result of an altered cell surface, but unlike the *pdr6Δ* mutant, it results in

hypervirulence rather than a protracted death. Furthermore, the *rim101Δ*-induced immune

response is mediated by neutrophils and eosinophils, hence, does not involve granulomas like

in the *mar1Δ* or *pdr6Δ* cases. Therefore, an altered exposure of PAMPs alone cannot explain the

pdr6Δ modulation of the immune response. Consistently, there are examples of cryptococcal

mutants that induce altered immune responses without any detectable surface changes, like

the F-box protein mutant *fbp1Δ* (48). Overall, this work and the examples discussed indicate

that a granulomatous inflammatory response is protective against *Cryptococcus* lung infection.

Interestingly, early clinical studies in cryptococcosis patients showed that T cells are needed for

an effective containment of fungal cells in a granuloma (51). This is consistent with our finding

that at 14 dpi, when the mice are controlling the *pdr6Δ* infection, there is a considerably

increase in lymphocytes relative to the WT-infection, while there are no significant changes in

innate cells. However, at 35 dpi, percentage-wise, there are almost no lymphocytes in the

pdr6Δ-infected lungs while there is a significant increase in macrophages (Fig. 11H). Although

further studies depleting these cell populations are needed to confirm this, it is conceivable

that a failed sustained lymphocyte response is responsible for the eventual death in the *pdr6Δ*-

infected mice. Curiously, a recent study from the Nielsen lab found an association between

hypervirulence and increased Th1/Th17 immune responses specific to the A/J mouse breed

(52). However, in that study, the hypervirulent strains induced a neutrophil-based response

while the strong inflammatory response induced by *pdr6Δ* seems to be mediated by

macrophages and not by neutrophils (Fig. 11H).

Taken together, these data provide important insights for improving the management of

CME. Anticryptococcal treatments are limited by toxicity, costs, and rising microbial resistance,

but FLC, despite being sub-optimal, still is the antifungal of choice in many regions of the world

because it is cheap, relatively less toxic, and broad spectrum. Since the deletion of *PDR6* results

in hypersensitivity to FLC, the design and synthesis of a novel compound that targets Pdr6,

coupled with FLC and/or a corticosteroid, may have clinical potential. Alternatively, several

groups in the cryptococcal and immunology fields have tried to identify potential vaccine

candidates against this fungus (53). Collectively, these preliminary vaccines, elicit a pro-

inflammatory response allowing for protection against subsequent challenges (54-56).

However, this is hampered because the immune mechanisms that elicit a protective immune

response are not known yet. Thus, the knowledge generated in this study has potential to help

inform cryptococcal vaccine development by providing novel insights into what constitutes a

deleterious pro-inflammatory response against *C. neoformans* infection.

Finally, the full complexities of PDR half-size transporters remain unknown. It is

understood that these proteins are promiscuous and transport a diverse array of substrates,

and with the expanded repertoire of PDR genes in *Cryptococcus*, it is not surprising to find roles

other than traditional efflux pumps and subsequent resistance to xenobiotics. *PDR6* has also

been implicated in aging and replicative life span (14), and most recently, *PDR9*, a full-length

PDR gene in the same clade as *AFR1*, was found to play a role in CO_2 tolerance via an unknown

mechanism that may involve phospholipid flipping (57). Hence, *PDR6* may act through multiple

direct and indirect mechanisms to modulate the inflammatory host response, such as by

secreting immunomodulatory molecules, aiding in maintaining the correct PM fluidity by

regulating ergosterol transport, and preventing exposure of surface PAMPs by supporting

correct cell wall and/or capsule architecture. All of these functions could be mediated by Pdr6-

dependent metabolites, lipids, or proteins, supporting future studies in these areas such as

comparative metabolomics of WT and *pdr6Δ* CM.

In conclusion, we have further characterized a novel PDR transporter in *C. neoformans*

by studying the role of *PDR6* in a mouse model of cryptococcosis and examining the altered

immune response. Mice infected with the *pdr6Δ* strain develop an overexuberant inflammatory

response that is effective at controlling the infection in the lungs short-term, but eventually

causes a fulminant pneumonia. Elucidating the role of *PDR6* in the context of infection

improves our understanding of cryptococcal pathogenesis while also informing potential

clinically-relevant therapeutic studies such as the search for novel drug targets or the

development of a vaccine. More broadly, this study expands the field of fungal PDR

transporters and given the conserved nature of these proteins, these findings may be

applicable to other fungal pathogens.

MATERIALS & METHODS

Strains, media, and growth conditions.

All *C. neoformans* strains used in this study were in the serotype A H99α background. All fungal

strains were maintained at -80ºC and grown at 30ºC on yeast peptone dextrose (YPD) with

antibiotics (NAT or G418) as appropriate. The *pdr6Δ* strain used for most of these studies was

obtained from the Madhani deletion collection (58) and the recreated strains were made in this

background by biolistics (13). The genome of all four independently created *pdr6Δ* mutants was

sequenced to confirm proper deletion of the gene without additional gross changes.

The human monocyte cell line THP-1 (ATCC TIB-202) was grown in THP-1 complete medium and

differentiated with phorbol 12-myristate 12-acetate (PMA, Sigma) as described (13). The hAELVi

cell line is an immortalized normal human alveolar epithelial cell obtained commercially

(InSCREENeX GmbH, Germany) and was grown and used as described (59).

Murine survival studies

In vivo fungal virulence was tested using the murine inhalation model of cryptococcosis (13, 42,

50, 60). Briefly, 5- to 6-week-old female inbred mice (A/J, C57BL/6, or BALB/c, all from The

Jackson Laboratories) were anesthetized with a ketamine rodent cocktail delivered i.p. and

intranasally inoculated, unless otherwise stated, with 5.0×10^4 cryptococcal cells in 40 μL of

sterile PBS. Animals were closely monitored and humanely euthanized if they lost >20% weight

relative to peak weight, or at predetermined timepoints, or at the end of the experiment

(maximum of 60 days). For drug treatment studies, cryptococcal-infected mice were randomly

assigned into treated or non-treated control groups. Stock solutions of fluconazole (1.6 mg/mL),

dexamethasone (0.03 mg/mL), and a fluconazole/dexamethasone combo were prepared in

sterile PBS. The drug(s) were administered intraperitoneally at doses of 16 mg/kg/day or 8

mg/kg/day for fluconazole, and of 0.3 mg/kg/day for dexamethasone. Treatment with these

compounds began 24 h after infection and continued throughout the experiment until endpoint

was reached. Statistically significant differences between survival curves were determined by

log rank Mantel-Cox test (GraphPad Prism). All of the mouse infections and procedures were

approved by the University of Notre Dame Institutional Animal Care and Use Committee

(protocol #23-05-7918).

Fungal Burden Quantification

Infected mice were humanely euthanized at predetermined time points or TOD by isoflurane

overdose followed by cervical dislocation, and lungs, brain, and spleens were harvested. Tissues

were homogenized in 5 mL of sterile PBS, diluted, and plated on YPD agar medium

supplemented with ampicillin. CFUs were counted following incubation at 30ºC for 48 h.

Statistical significance was determined using ANOVA Kruskal-Wallis test (GraphPad Prism).

Histological analyses

A/J female mice (The Jackson Laboratories) were infected via the intranasal route as described

above with 5 x 10^4 CFU of either WT (H99) or the *pdr6*Δ mutant strain. Mice were sacrificed as

predetermined time points (14, 35, 50 days post inoculation) by isoflurane overdose followed

by cervical dislocation. Lungs were perfused with sterile PBS and 10% neutralized formalin.

Lungs were subsequently paraffin embedded, sectioned, mounted, and stained with H&E by the

University of Notre Dame Histology Core facility. H&E-stained slides were imaged at 10X and

40X using an inverted Zeiss microscope. Slides were scored in a blind manner by the resident

pathologist at the Harper Cancer Research Institute.

Cytokine Analyses

Female mice (A/J, C57BL/6, or BALB/c, all from The Jackson Labs) were infected via the

intranasal route with WT (H99) or the *pdr6*Δ mutant strain as described above. Mice were

euthanized at predetermined time points by isoflurane overdose followed by cervical

dislocation. Cytokine levels within the A/J mice lung and brain homogenates were analyzed

using the Proteome Profiler Mouse XL Cytokine Array (R&D Systems Inc., Minneapolis, MN). The

relative expression levels of 111 soluble mouse proteins were determined following the

manufacturer's protocol. Array membranes were imaged using LI-COR Odyssey Infrared

Imaging System and quantified using ImageStudio software. Cytokine/chemokine expression

levels were further confirmed in A/J mice, as well as in C57BL/6 and BALB/c mice, using the Bio-

Plex Multi-Plex Assay Kit (Bio-Rad Laboratories, Hercules, CA). Supernatants from organ

homogenates were used to determine presence of 22 proteins including eotaxin, G-CSF, GM-

CSF, IFN-γ, IL-1α, IL-1β, IL-2, IL-3, IL-4, IL-6, IL-9, IL-10, IL-12 (p40), IL-12 (p70), IL-13, IL-17A, KC,

MCP-1, MCP-1α, MIP-1β, RANTES, and TNF-α.

Pulmonary leukocyte isolation and flow cytometric analysis

H99 and *pdr6*Δ-infected A/J mice were euthanized at predetermined time points and TOD by

isoflurane overdose followed by cervical dislocation. Lungs of infected mice were excised and

digested enzymatically at 37°C for 60 min in 10 mL of digest buffer (0.34 U/mL Liberase^{TM}

(Roche) and 100 ug/mL DNaseI (ThermoFisher)) in 1X DPBS. After 60 min, digestion was

arrested with 1X DPBS supplemented with 2% FBS and 0.2 um EDTA. The cell suspension was

washed and passed through 40-μm cell strainers. The single cell suspensions were stained with

a live/dead far red viability dye (Invitrogen), washed, and incubated on ice for 20 min with

purified anti-mouse CD16/CD32 antibodies (BioLegend) to block Fc receptors. Following

blocking, cells were incubated with fluorochrome-conjugated antibodies to allow for

multistaining for 45 min on ice. The innate panel consisted of PE-Cy5 conjugated anti-CD45 (30-

111), Brilliant Violet (BV) 650-conjugated anti-CD11b (M1/70), Super Bright (SB) 780-conjugated

anti-F4/80 (BM8), BV711-conjugated anti-Ly6G (1A8), PE-Cy7-conjugated anti-Ly6C (HK1.4), PE-

conjugated anti-SiglecF (1RNM44N), and APC-eFluor 780-conjugated anti-CD11c (N418). The

adaptive panel included PE-Cy5 conjugated anti-CD45 (30-111), AF700-conjugated anti-CD3

(145-2C11), BV421-conjugated anti-CD4 (GK1.5), BV510-conjugated anti-CD25 (PC61-5), PE-

conjugated anti-CD8 (53-6.7), and AF488-conjugated anti-FoxP3 (FJK-16s). Cells were washed

and fixed with 2% formalin in FACS buffer. UltraComp eBeads (Invitrogen) were stained with

each respective fluorochrome-conjugated antibody and used to determine and

spillover/compensation calculations. Data was analyzed using FlowJo software.

Cell wall and capsule stains

Starter cultures were grown overnight in YPD, washed with DMEM, and 1 x 10^6 cells/mL were

added to 24-well tissue culture plates. Plates were incubated at 37°C + 5% CO2 for 48 h. The

cells were collected by centrifugation at 3,000 x g for 5 min. For staining, Calcofluor White

(CFW), Alexa Fluor 488-conjugated concanavalin A (ConA), β-(1,3)-glucan antibody 2GB

(ab233743; Abcam, USA), eosin Y (EoY), and the capsular antibody 302 were used. Cells were

washed and stained with CFW, ConA, β-(1,3)-glucan antibody, EoY, and Mab 302 at 100 μg/mL,

50 μg/mL, 20 μg/mL, 500 μg/mL, and 10 μg/mL, respectively. Cells were incubated with the

stains for 30 min, washed, fixed, and imaged at 100X using an inverted Zeiss microscope.

Fluorescence of mAb 302 was calculated using Cell Profiler software. Flow cytometry data was

acquired using a BD LSRFortessa cytometer. The CFW signal was detected using a 405 nm laser

and the ConA and EoY signals were detected using a 488 nm laser. Flow cytometry data was

analyzed with FlowJo software.

ACKNOWLEDGMENTS

We acknowledge assistance and support on the murine studies from the University of Notre

Dame Freimann Life Science Center (FLSC). We give a special thanks to William J. Kaliney, M.D.,

pathologist from Harper Cancer Research Center, for assistance with histological analysis and

scoring. We acknowledge financial support from the Warren Center for Drug Discovery (pilot

grant) and NIH grant R21AI71742. Other related work in the Santiago-Tirado lab is supported by

NIH grant R01AI177875. We also thank members of the Santiago-Tirado lab for their thoughtful

comments and feedback on this work.

REFERENCES

1.   Denning DW. 2024. Global incidence and mortality of severe fungal disease. Lancet Infect Dis
doi:10.1016/S1473-3099(23)00692-8.
2.   Gaylord EA, Choy HL, Doering TL. 2020. Dangerous Liaisons: Interactions of Cryptococcus
neoformans with Host Phagocytes. Pathogens 9.
3.   Stuckey PV, Santiago-Tirado FH. 2023. Fungal mechanisms of intracellular survival: what can we
learn from bacterial pathogens? Infect Immun 91:e0043422.
4.   Rajasingham R, Govender NP, Jordan A, Loyse A, Shroufi A, Denning DW, Meya DB, Chiller TM,
Boulware DR. 2022. The global burden of HIV-associated cryptococcal infection in adults in
2020: a modelling analysis. Lancet Infect Dis doi:10.1016/S1473-3099(22)00499-6.
5.   Santiago-Tirado FH, Doering TL. 2017. False friends: Phagocytes as Trojan horses in microbial
brain infections. PLoS Pathog 13:e1006680.
6.   Denham ST, Brown JCS. 2018. Mechanisms of Pulmonary Escape and Dissemination by
Cryptococcus neoformans. J Fungi (Basel) 4.
7.   Zaragoza O, Rodrigues ML, De Jesus M, Frases S, Dadachova E, Casadevall A. 2009. The capsule
of the fungal pathogen Cryptococcus neoformans. Adv Appl Microbiol 68:133-216.
8.   Denham ST, Verma S, Reynolds RC, Worne CL, Daugherty JM, Lane TE, Brown JCS. 2018.
Regulated Release of Cryptococcal Polysaccharide Drives Virulence and Suppresses Immune Cell
Infiltration into the Central Nervous System. Infect Immun 86.
9.   Eisenman HC, Casadevall A. 2012. Synthesis and assembly of fungal melanin. Appl Microbiol
Biotechnol 93:931-40.
10.   Alspaugh JA. 2015. Virulence mechanisms and Cryptococcus neoformans pathogenesis. Fungal
Genet Biol 78:55-8.
11.   Pacelli C, Bryan RA, Onofri S, Selbmann L, Shuryak I, Dadachova E. 2017. Melanin is effective in
protecting fast and slow growing fungi from various types of ionizing radiation. Environ
Microbiol 19:1612-1624.
12.   Ueno K, Nagamori A, Honkyu NO, Kataoka M, Shimizu K, Chang YC, Kwon-Chung KJ, Miyazaki Y.
2023. Cryptococcus neoformans requires the TVF1 gene for thermotolerance and virulence.
Med Mycol 61.
13.   Winski CJ, Qian Y, Mobashery S, Santiago-Tirado FH. 2022. An Atypical ABC Transporter Is
Involved in Antifungal Resistance and Host Interactions in the Pathogenic Fungus *Cryptococcus*
neoformans. mBio 13:e0153922.
14.   Oliveira NK, Bhattacharya S, Gambhir R, Joshi M, Fries BC. 2022. Novel ABC Transporter
Associated with Fluconazole Resistance in Aging of Cryptococcus neoformans. J Fungi (Basel) 8.
15.   Moreno A, Banerjee A, Prasad R, Falson P. 2019. PDR-like ABC systems in pathogenic fungi. Res
Microbiol 170:417-425.
16.   Khunweeraphong N, Kuchler K. 2021. Multidrug Resistance in Mammals and Fungi-From MDR to
PDR: A Rocky Road from Atomic Structures to Transport Mechanisms. Int J Mol Sci 22.
17.   Jankovic D, Liu Z, Gause WC. 2001. Th1- and Th2-cell commitment during infectious disease:
asymmetry in divergent pathways. Trends Immunol 22:450-7.

18. Stacey AR, Norris PJ, Qin L, Haygreen EA, Taylor E, Heitman J, Lebedeva M, DeCamp A, Li D,
Grove D, Self SG, Borrow P. 2009. Induction of a striking systemic cytokine cascade prior to peak
viremia in acute human immunodeficiency virus type 1 infection, in contrast to more modest
and delayed responses in acute hepatitis B and C virus infections. J Virol 83:3719-33.
19. McMichael AJ, Borrow P, Tomaras GD, Goonetilleke N, Haynes BF. 2010. The immune response
during acute HIV-1 infection: clues for vaccine development. Nat Rev Immunol 10:11-23.
20. Wozniak KL, Ravi S, Macias S, Young ML, Olszewski MA, Steele C, Wormley FL. 2009. Insights into
the mechanisms of protective immunity against Cryptococcus neoformans infection using a
mouse model of pulmonary cryptococcosis. PLoS One 4:e6854.
21. Wozniak KL, Hardison S, Olszewski M, Wormley FL, Jr. 2012. Induction of protective immunity
against cryptococcosis. Mycopathologia 173:387-94.
22. Chang CC, Harrison TS, Bicanic TA, Chayakulkeeree M, Sorrell TC, Warris A, Hagen F, Spec A,
Oladele R, Govender NP, Chen SC, Mody CH, Groll AH, Chen YC, Lionakis MS, Alanio A, Castaneda
E, Lizarazo J, Vidal JE, Takazono T, Hoenigl M, Alffenaar JW, Gangneux JP, Soman R, Zhu LP,
Bonifaz A, Jarvis JN, Day JN, Klimko N, Salmanton-Garcia J, Jouvion G, Meya DB, Lawrence D,
Rahn S, Bongomin F, McMullan BJ, Sprute R, Nyazika TK, Beardsley J, Carlesse F, Heath CH,
Ayanlowo OO, Mashedi OM, Queiroz-Telles Filho F, Hosseinipour MC, Patel AK, Temfack E, Singh
N, Cornely OA, Boulware DR, et al. 2024. Global guideline for the diagnosis and management of
cryptococcosis: an initiative of the ECMM and ISHAM in cooperation with the ASM. Lancet Infect
Dis doi:10.1016/S1473-3099(23)00731-4.
23. Rothe C, Sloan DJ, Goodson P, Chikafa J, Mukaka M, Denis B, Harrison T, van Oosterhout JJ,
Heyderman RS, Lalloo DG, Allain T, Feasey NA. 2013. A prospective longitudinal study of the
clinical outcomes from cryptococcal meningitis following treatment induction with 800 mg oral
fluconazole in Blantyre, Malawi. PLoS One 8:e67311.
24. Coelho C, Casadevall A. 2016. Cryptococcal therapies and drug targets: the old, the new and the
promising. Cell Microbiol 18:792-9.
25. Zhang J, Li L, Lv Q, Yan L, Wang Y, Jiang Y. 2019. The Fungal CYP51s: Their Functions, Structures,
Related Drug Resistance, and Inhibitors. Front Microbiol 10:691.
26. Casadevall A. 2022. Immunity to Invasive Fungal Diseases. Annu Rev Immunol 40:121-141.
27. Qiu Y, Dayrit JK, Davis MJ, Carolan JF, Osterholzer JJ, Curtis JL, Olszewski MA. 2013. Scavenger
receptor A modulates the immune response to pulmonary Cryptococcus neoformans infection. J
Immunol 191:238-48.
28. Brinks J, van Dijk EHC, Habeeb M, Nikolaou A, Tsonaka R, Peters HAB, Sips HCM, van de Merbel
AF, de Jong EK, Notenboom RGE, Kielbasa SM, van der Maarel SM, Quax PHA, Meijer OC, Boon
CJF. 2018. The Effect of Corticosteroids on Human Choroidal Endothelial Cells: A Model to Study
Central Serous Chorioretinopathy. Invest Ophthalmol Vis Sci 59:5682-5692.
29. Day J, Imran D, Ganiem AR, Tjahjani N, Wahyuningsih R, Adawiyah R, Dance D, Mayxay M,
Newton P, Phetsouvanh R, Rattanavong S, Chan AK, Heyderman R, van Oosterhout JJ, Chierakul
W, Day N, Kamali A, Kibengo F, Ruzagira E, Gray A, Lalloo DG, Beardsley J, Binh TQ, Chau TT,
Chau NV, Cuc NT, Farrar J, Hien TT, Van Kinh N, Merson L, Phuong L, Tho LT, Thuy PT, Thwaites
G, Wertheim H, Wolbers M. 2014. CryptoDex: a randomised, double-blind, placebo-controlled
phase III trial of adjunctive dexamethasone in HIV-infected adults with cryptococcal meningitis:
study protocol for a randomised control trial. Trials 15:441.
30. Kitonsa J, Nsubuga R, Mayanja Y, Kiwanuka J, Nikweri Y, Onyango M, Anywaine Z, Ggayi AB,
Kibengo FM, Kaleebu P, Day J. 2020. Determinants of two-year mortality among HIV positive
patients with Cryptococcal meningitis initiating standard antifungal treatment with or without
adjunctive dexamethasone in Uganda. PLoS Negl Trop Dis 14:e0008823.

31.   Netea MG, Schlitzer A, Placek K, Joosten LAB, Schultze JL. 2019. Innate and Adaptive Immune
Memory: an Evolutionary Continuum in the Host's Response to Pathogens. Cell Host Microbe
25:13-26.
32.   Rudman J, Evans RJ, Johnston SA. 2019. Are macrophages the heroes or villains during
cryptococcosis? Fungal Genet Biol 132:103261.
33.   Hole C, Wormley FL, Jr. 2016. Innate host defenses against Cryptococcus neoformans. J
Microbiol 54:202-11.
34.   Hole CR, Lam WC, Upadhya R, Lodge JK. 2020. Cryptococcus neoformans Chitin Synthase 3 Plays
a Critical Role in Dampening Host Inflammatory Responses. mBio 11.
35.   O'Halloran JA, Powderly WG, Spec A. 2017. Cryptococcosis today: It is not all about HIV
infection. Curr Clin Microbiol Rep 4:88-95.
36.   Koselny K, Mutlu N, Minard AY, Kumar A, Krysan DJ, Wellington M. 2018. A Genome-Wide
Screen of Deletion Mutants in the Filamentous Saccharomyces cerevisiae Background Identifies
Ergosterol as a Direct Trigger of Macrophage Pyroptosis. mBio 9.
37.   Rodrigues ML. 2018. The Multifunctional Fungal Ergosterol. mBio 9.
38.   Pasrija R, Panwar SL, Prasad R. 2008. Multidrug transporters CaCdr1p and CaMdr1p of Candida
albicans display different lipid specificities: both ergosterol and sphingolipids are essential for
targeting of CaCdr1p to membrane rafts. Antimicrob Agents Chemother 52:694-704.
39.   Xu X, Bittman R, Duportail G, Heissler D, Vilcheze C, London E. 2001. Effect of the structure of
natural sterols and sphingolipids on the formation of ordered sphingolipid/sterol domains
(rafts). Comparison of cholesterol to plant, fungal, and disease-associated sterols and
comparison of sphingomyelin, cerebrosides, and ceramide. J Biol Chem 276:33540-6.
40.   Upadhya R, Lam WC, Hole CR, Vasselli JG, Lodge JK. 2023. Cell wall composition in Cryptococcus
neoformans is media dependent and alters host response, inducing protective immunity. Front
Fungal Biol 4.
41.   Kozel TR, Levitz SM, Dromer F, Gates MA, Thorkildson P, Janbon G. 2003. Antigenic and
biological characteristics of mutant strains of Cryptococcus neoformans lacking capsular O
acetylation or xylosyl side chains. Infect Immun 71:2868-75.
42.   Santiago-Tirado FH, Peng T, Yang M, Hang HC, Doering TL. 2015. A Single Protein S-acyl
Transferase Acts through Diverse Substrates to Determine Cryptococcal Morphology, Stress
Tolerance, and Pathogenic Outcome. PLoS Pathog 11:e1004908.
43.   Dambuza IM, Drake T, Chapuis A, Zhou X, Correia J, Taylor-Smith L, LeGrave N, Rasmussen T,
Fisher MC, Bicanic T, Harrison TS, Jaspars M, May RC, Brown GD, Yuecel R, MacCallum DM,
Ballou ER. 2018. The Cryptococcus neoformans Titan cell is an inducible and regulated
morphotype underlying pathogenesis. PLoS Pathog 14:e1006978.
44.   Posteraro B, Sanguinetti M, Sanglard D, La Sorda M, Boccia S, Romano L, Morace G, Fadda G.
2003. Identification and characterization of a Cryptococcus neoformans ATP binding cassette
(ABC) transporter-encoding gene, CnAFR1, involved in the resistance to fluconazole. Mol
Microbiol 47:357-71.
45.   Prasad R, De Wergifosse P, Goffeau A, Balzi E. 1995. Molecular cloning and characterization of a
novel gene of Candida albicans, CDR1, conferring multiple resistance to drugs and antifungals.
Curr Genet 27:320-9.
46.   Lamping E, Baret PV, Holmes AR, Monk BC, Goffeau A, Cannon RD. 2010. Fungal PDR
transporters: Phylogeny, topology, motifs and function. Fungal Genet Biol 47:127-42.
47.   Telzrow CL, Esher Righi S, Castro-Lopez N, Campuzano A, Brooks JT, Carney JM, Wormley FL, Jr.,
Alspaugh JA. 2022. An Immunogenic and Slow-Growing Cryptococcal Strain Induces a Chronic
Granulomatous Infection in Murine Lungs. Infect Immun 90:e0058021.

48. Masso-Silva J, Espinosa V, Liu TB, Wang Y, Xue C, Rivera A. 2018. The F-Box Protein Fbp1 Shapes
the Immunogenic Potential of Cryptococcus neoformans. mBio 9.
49. Esher SK, Ost KS, Kohlbrenner MA, Pianalto KM, Telzrow CL, Campuzano A, Nichols CB, Munro C,
Wormley FL, Jr., Alspaugh JA. 2018. Defects in intracellular trafficking of fungal cell wall
synthases lead to aberrant host immune recognition. PLoS Pathog 14:e1007126.
50. Gish SR, Maier EJ, Haynes BC, Santiago-Tirado FH, Srikanta DL, Ma CZ, Li LX, Williams M, Crouch
EC, Khader SA, Brent MR, Doering TL. 2016. Computational Analysis Reveals a Key Regulator of
Cryptococcal Virulence and Determinant of Host Response. MBio 7:e00313-16.
51. Hill JO. 1992. CD4+ T cells cause multinucleated giant cells to form around Cryptococcus
neoformans and confine the yeast within the primary site of infection in the respiratory tract. J
Exp Med 175:1685-95.
52. Ding M, Jackson KM, Harris-Gordon M, Dileepan T, Meya DB, Nielsen K. 2025. The hypervirulent
Type-1/Type-17 phenotype of Cryptococcus neoformans clinical isolates is specific to A/J mice.
Infect Immun 93:e0058524.
53. Del Poeta M, Wormley FL, Jr., Lin X. 2023. Host populations, challenges, and commercialization
of cryptococcal vaccines. PLoS Pathog 19:e1011115.
54. Wang Y, Wang K, Rivera A, Xue C. 2023. Development of a Heat-Killed fbp1 Mutant Strain as a
Therapeutic Agent To Treat Invasive Cryptococcus Infection. Microbiol Spectr 11:e0495522.
55. Pham T, Li Y, Watford W, Lin X. 2023. Vaccination with a ZNF2(oe) Strain of Cryptococcus
Provides Long-Lasting Protection against Cryptococcosis and Is Effective in
Immunocompromised Hosts. Infect Immun 91:e0019823.
56. Wormley FL, Jr., Perfect JR, Steele C, Cox GM. 2007. Protection against cryptococcosis by using a
murine gamma interferon-producing *Cryptococcus neoformans* strain. Infect Immun 75:1453-62.
57. Ristow LC, Jezewski AJ, Chadwick BJ, Stamnes MA, Lin X, Krysan DJ. 2023. Cryptococcus
neoformans adapts to the host environment through TOR-mediated remodeling of phospholipid
asymmetry. Nat Commun 14:6587.
58. Liu OW, Chun CD, Chow ED, Chen C, Madhani HD, Noble SM. 2008. Systematic genetic analysis
of virulence in the human fungal pathogen *Cryptococcus neoformans*. Cell 135:174-188.
59. Kuehn A, Kletting S, de Souza Carvalho-Wodarz C, Repnik U, Griffiths G, Fischer U, Meese E,
Huwer H, Wirth D, May T, Schneider-Daum N, Lehr CM. 2016. Human alveolar epithelial cells
expressing tight junctions to model the air-blood barrier. ALTEX 33:251-60.
60. Santiago-Tirado FH, Hurtaux T, Geddes-McAlister J, Nguyen D, Helms V, Doering TL, Romisch K.
2023. The ER Protein Translocation Channel Subunit Sbh1 Controls Virulence of Cryptococcus
neoformans. mBio 14:e0338422.

Figure legends and Table captions

Fig. 1. *C. neoformans pdr6Δ*-infected mice succumb to infection despite decreased fungal
burden. (A) Quantification of fungal burden over time to monitor disease progression. Mice

were inoculated with 5 x 10^4 cells and sacrificed every 7 days and organs were harvested to
determine CFUs. Significance was determined using Mann-Whitney test, *P<0.05. **(B)** Survival
studies of animals infected with H99 and *pdr6Δ* strains, or mock-infected (PBS). 4 A/J mice per
group were intranasally infected with 5 x 10^3 cryptococcal cells or mock infected with sterile
PBS and monitored for 38 days (experiment was terminated when the last infected mouse
reached endpoint). Significance was determined using Mantel-Cox test, **P<0.01. **(C)**
Quantified organ burdens of H99 and *pdr6Δ*-infected mice at TOD. Organs were harvested,
homogenized, and dilutions were plated on YPD. Plates were incubated for 48 h and CFUs were
counted. Each data point represents a mouse and black bars represent the mean. Significance
was determined using Mann-Whitney test, *P<0.05; **P<0.01.

Fig. 2. Infection with *C. neoformans pdr6Δ* results in altered histopathology. (A & B) Dissected
lungs of female A/J mice infected with 5 x 10^4 cells of the WT strain (H99) or *pdr6Δ* mutant
strain, or mock-infected, were harvested at predetermined endpoints of 14 **(A)** and 35 **(B)** dpi.
Gross organ examination revealed significant changes in inflammation and tissue damage. H&E
staining was used to visualize microscopic lung pathology (inset depicted by the green boxes is
shown in the 40X column). The 10X scale bar is 200 μm and the 40X scale bar is 50 μm.

Table 1. Quantitative analysis of H99 and *pdr6Δ*-infected lung sections over time.
*All H99-infected mice had succumbed to infection by this timepoint.
Histiocytes refer to leukocytes of the monocyte/macrophage/dendritic cell lineage.
hpf, high power field.

Fig. 3. FLC treatment significantly reduces CFUs in *pdr6Δ*-infected mice without changing the
outcome. (A) Survival curve of H99 and *pdr6Δ*-infected mice treated daily with either 8 or 16
mg/kg FLC. At least 4 A/J mice per group were intranasally infected with 5 x 10^4 cells, or mock-
infected with DPBS and monitored for up to 50 dpi (mock-infected and FLC-treated mice not
shown for clarity). Significance was determined using Mantel-Cox test, *P<0.05; **P<0.01;
***P<0.001; ****P<0.0001. **(B-G)** Calculated organ burdens at 14 dpi **(B-D)** or endpoint **(E-G)**.
Organs were harvested, homogenized, and dilutions were plated on YPD. Plates were incubated
for 48 hr and CFUs were enumerated. Each data point represents a mouse and black bars
represent the mean. Significance was determined using one-way ANOVA with multiple
comparisons (Welch's corrected), *P<0.05; **P<0.01; ****P<0.0001.

Fig. 4. FLC treatment improves lung pathology in *pdr6Δ*-infected animals. Dissected lungs of
female A/J mice infected with 5 x 10^4 cells of the WT strain (H99) or *pdr6Δ* mutant strain and
treated daily with 16 mg/kg FLC were harvested at 14 and 35 dpi. Gross organ examination

revealed alterations in fungal containment and disease progression. H&E staining was used to
visualize microscopic lung pathology. Insets 1 & 2 highlight differential histopathology in the
FLC-treated *pdr6Δ*-infected mice at 35 dpi (inset depicted by the green boxes is shown in the
40X column). Inset 1 shows an intact lung parenchyma with several fungi and immune cells
while inset 2 is characterized as a granuloma. The 10X scale bar is 200 μm and the 40X scale bar
is 50 μm.

Table 2. Quantitative analysis of Fluconazole treated H99 and *pdr6Δ*-infected lung sections
over time
Histiocytes refer to leukocytes of the monocyte/macrophage/dendritic cell lineage. hpf, high
power field.

Fig. 5 Infection with the *pdr6Δ* mutant elicits a robust pro-inflammatory immune response. (A
& B) Heatmap of the pro-inflammatory and anti-inflammatory cytokines/chemokines in the
lungs **(A)** and brains **(B)** of *C. neoformans*-infected mice. Female A/J mice were intranasally
infected with 5 x 10^4 cells of the H99 or *pdr6Δ* strain and euthanized at endpoint. Homogenates
from the lungs and brain of at least 4 mice per group were used to perform Proteome Profiler
Mouse XL Cytokine Arrays and Bio-Plex Multi-Plex Arrays. **(C & D)** Expression of selected
cytokines/chemokines in the lungs **(C)** and brains **(D)** of *pdr6Δ*-infected mice was normalized to
H99 (dotted line). Significance was determined by Ordinary 2-way ANOVA with multiple
comparisons, *$P<0.05$; **$P<0.01$; ***$P<0.001$; ****$P<0.0001$.

Fig. 6 Dexamethasone treatment significantly extends survival of *pdr6Δ*-infected mice. (A)
Survival curve of H99 and *pdr6Δ*-infected mice treated daily with either 0.03 mg/kg Dexa or
0.03 mg/kg Dexa and 16 mg/kg FLC. At least 3 A/J mice per group were intranasally infected
with 5 x 10^4 cells, or mock-infected with DPBS and monitored for up to 60 dpi (mock-infected
and treated mice not shown for clarity). Significance was determined using Mantel-Cox test,
$P<0.01$; ****$P<0.0001$. **(B-G) Calculated organ burdens at 14 dpi **(B-D)** or endpoint **(E-G)**.
Organs were harvested, homogenized, and dilutions were plated on YPD. Plates were incubated
for 48 hr and CFUs were enumerated. Each data point represents a mouse and black bars
represent the mean. Significance was determined using one-way ANOVA with multiple
comparisons (Welch's corrected) *$P<0.05$; **$P<0.01$; ***$P<0.001$.

Fig. 7. Effect of dexamethasone treatment on lung histopathology. Dissected lungs of female
A/J mice infected with 5 x 10^4 cells of the WT strain (H99) or *pdr6Δ* mutant strain and treated
daily with 0.3 mg/kg Dexa and harvested at 14 and 35 dpi. Gross organ examination revealed

alterations in levels of inflammation. H&E staining was used to visualize microscopic lung
pathology (inset [green boxes]). The 10X scale bar is 200 μm and the 40X scale bar is 50 μm.

Table 3. Quantitative analysis of dexamethasone-treated H99 and *pdr6Δ*-infected lung
sections over time
Histiocytes refer to leukocytes of the monocyte/macrophage/dendritic cell lineage.
hpf, high power field.

Fig. 8. Effect of dexamethasone/FLC combination treatment on lung histopathology.
Dissected lungs of female A/J mice infected with 5 x 10^4 cells of the WT strain (H99) or *pdr6Δ*
mutant strain and treated daily with a combination of 0.3 mg/kg Dexa and 16 mg/kg FLC and
harvested at 14 and 35 dpi. Gross organ examination revealed alterations in levels of
inflammation. H&E staining was used to visualize microscopic lung pathology (inset [green
boxes]). The 10X scale bar is 200 μm and the 40X scale bar is 50 μm.

Table 4. Quantitative analysis of combination Dexamethasone + Fluconazole treated H99 and
***pdr6Δ*-infected lung sections over time**
Histiocytes refer to leukocytes of the monocyte/macrophage/dendritic cell lineage.
hpf, high power field.

Fig. 9 Infection with the *C. neoformans pdr6Δ* mutant generates an increase in innate immune
cells, specifically macrophages. (A-H) Quantification of innate and adaptive immune cell
populations at 14 dpi **(A-D)** and endpoint (35 dpi for *pdr6Δ*-infected mice)**(E-H)**. Female A/J
mice were intranasally infected with 5 x 10^4 cells of the WT type (H99) or *pdr6Δ* strain and
euthanized at clinical endpoints corresponding to imminent mortality. Lungs were harvested
and single-cell suspensions were generated and stained with immune panels to quantify
monocytes (CD45+F4/80-CD11b+Ly6C+), neutrophils (CD45+F4/80-CD11b+Ly6G+),
macrophages (CD45+F4/80+CD11b+), CD4$^+$ Th (CD45+CD3+CD4+), CD4$^+$ T-regs
(CD45+CD3+CD4+CD25+FoxP3+), and CD8$^+$ (CD45+CD3+CD8+) cells. Samples were processed
using flow cytometry and are presented as total # cells **(A, C, E, G)** or as % CD45$^+$ cells **(B, D, F,**
H). Significance was determined using one-way ANOVA with multiple comparisons (Welch's
corrected) *P<0.05; **P<0.01. Note: for **G** and **H**, the CD8$^+$ T cell panel could not be run for the
day 35 samples, hence they were removed from the H99 bars.

Fig. 10 Mice breeds with skewed immune responses influence the outcome of the *pdr6Δ*
infection. Survival curves and organ burdens for C57BL/6 **(A − D)** or BALB/c **(F − I)** mice infected

with 5 x 10^4 cells of WT (H99) or *pdr6Δ* strains. At least 5 mice were infected for each group.
Significance in the survival curves was determined using Mantel-Cox test, **P<0.01;
***P<0.001. For fungal burdens, organs were harvested, homogenized, and dilutions were
plated on YPD. Plates were incubated for 48 hr and CFUs were enumerated. Each data point
represents a mouse and black bars represent the mean. In panels G-I, crossed data points
represent the values from the mice that succumbed to infection. Significance was determined
using one-way ANOVA with multiple comparisons (Welch's corrected), *P<0.05. **(E & J)**
Heatmap of the pro-inflammatory and anti-inflammatory cytokines/chemokines in the lungs of
C57BL/6 **(E)** and BALB/c **(J)** infected mice. Values are shown normalized to age-match mock-
infected mice.

Fig. 11 *C. neoformans pdr6Δ* cells display altered cell wall features and capsule density. (A)
Visualization of calcofluor (CFW), Alexa488-conjugated Concanavalin A (ConA), and β-(1,3)-
glucan staining in H99, *pdr6Δ*-L (from commercial deletion collection), and *cap59Δ* cells.
Representative images of stained cells. Scale bar, 10 μm. **(B - E)** Quantification of CFW **(B)**, ConA
(C), Eosin Y **(D)**, and β-glucan **(E)** staining intensity in H99, *pdr6Δ*, and *cap59Δ* strains. Staining
intensity was measured by flow cytometry and results normalized to H99 (dotted line). Four
independent *pdr6Δ* strains were tested (*pdr6Δ*-L and *pdr6Δ*-1 to -3). Significance was
determined using one-way ANOVA (Welch's corrected), *P<0.05; ***P<0.001; ****P<0.0001.
(F) Visualization and quantification of capsular antibody staining of H99, *pdr6Δ*, and *cap59Δ*
strains. Cells were stained with anti-capsule mAb 302 and imaged using an inverted Zeiss
microscope. Representative images of stained cells. Staining intensity was quantified using a
Cell Profiler pipeline. Significance was determined using one-way ANOVA (Welch's corrected)
***P<0.001. **(G)** The % of titanization of the indicated strains was determined as described in
the methods. Each data point represents the average % of titanization in independent biological
experiments each with at least 100 cells quantified. One-way ANOVA with multiple
comparisons, ***P<0.001; ****P<0.0001. **(H)** Model explaining how *pdr6Δ* strains alter disease
progression in A/J and C57BL/6 mice, causing fatal pneumonia rather than the typical
meningitis caused by H99. The *pdr6Δ* strain has an altered cell surface, including a less dense
capsule that is more permeable to antibodies, that results in exposure of pro-inflammatory
antigens. This results in an initial immune response that can control the infection, consisting on
an increased number of CD4 lymphocytes. However, that initial response is not enough to clear
the infection, and eventually results in dysregulation, with a drop in the number of CD4
lymphocytes and a dramatic increase in inflammatory macrophages. Despite having less fungal
burden in the lungs and little dissemination, the mice succumb to the infection due to too much
lung tissue damage. This illustration was originally made in BioRender
https://BioRender.com/vpa8dnh) and subsequently edited in Photoshop.

Figure 1

Figure 2

Table 1

DPI	Strain	% inflammation	Fungal infiltration	Quantitation of inflammatory immune cells/hpf 400X	Observations
7	H99	5 – 10%	20% alveoli contain encapsulated yeast	8 neutrophils 33 lymphocytes 40 histiocytes	Immune cells in peri-bronchial and perivascular space.
	pdr6Δ	5 – 10%	15 – 20% alveoli contain encapsulated yeast	9 neutrophils 26 lymphocytes 19 histiocytes	Immune cells in peri-bronchial and perivascular space. Some yeast has been phagocytized by histiocytes.
14	H99	10%	80 – 90% alveoli contain encapsulated yeast	52 neutrophils 26 lymphocytes 57 histiocytes	Immune cells in peri-bronchial and perivascular space.
	pdr6Δ	10%	50 – 60% alveoli contain encapsulated yeast	43 neutrophils 105 lymphocytes 71 histiocytes	Immune cells in peri-bronchial and perivascular space. Many histiocytes have phagocytized yeast. Areas of accumulated yeast/immune cells but no true parenchymal granulomas.
21*	pdr6Δ	10%	90% alveoli contain encapsulated yeast	50 neutrophils 49 lymphocytes 55 histiocytes	Immune cells in peri-bronchial and perivascular space.
28*	pdr6Δ	5 – 10%	80% alveoli contain encapsulated yeast	8 neutrophils 33 lymphocytes 55 histiocytes	Immune cells in peri-bronchial and perivascular space and single parenchymal granuloma (1.2mm)
35*	pdr6Δ	60 – 70%	100% present throughout	Too many to quantify	Consolidated parenchyma with many coalescent granulomas (histiocytes w/phagocytized yeast) and nests of neutrophils and lymphocytes with necrosis.

Figure 3

Figure 4

Table 2

DPI	Strain	% inflammation	Fungal infiltration	Quantitation of inflammatory immune cells/hpf 400X	Observations
14	Mock +FLC	0%	None	-	-
	H99 +FLC	5 – 10%	~ 60% alveoli contain encapsulated yeast	50 – 60 neutrophils 40 – 50 lymphocytes 15 histiocytes	Moderate number of inflammatory cells predominately present in perivascular & peribronchial areas.
	pdr6Δ +FLC	< 5%	30 – 40% alveoli contain encapsulated yeast	1 – 5 neutrophils 0 – 15 lymphocytes 1 – 5 histiocytes	Mild infiltrate of inflammatory cells in focal peribronchial & perivascular areas.
35	Mock +FLC	0%	None	-	-
	pdr6Δ +FLC	5 – 10%	45 – 50% alveoli contain encapsulated yeast	5 – 10 neutrophils 10 – 20 lymphocytes < 5 histiocytes	Mild inflammatory cell infiltrate in many peribronchial and perivascular areas. Consolidation of adjacent alveoli in solid areas.

Figure 5

And Fig. S1

Figure 6

Figure 7

Table 3

DPI	Strain	% inflammation	Fungal infiltration	Quantitation of inflammatory immune cells/hpf 400X	Observations
14	Mock +Dexa	0%	None	-	-
	H99 +Dexa	10%	85 – 90% alveoli contain encapsulated yeast	10 – 20 neutrophils 10 – 30 lymphocytes < 5 histiocytes	Perivascular and peribronchial aggregates of admixed neutrophils and lymphocytes.
	pdr6Δ +Dexa	< 5%	20 – 30% alveoli contain encapsulated yeast	5 – 10 neutrophils 5 lymphocytes < 5 histiocytes	Focal inflammatory cell nests in perivascular and peribronchial space.
50	Mock +Dexa	0%	None	-	Inflammatory cells uncommon. Rare small nests of lymphocytes on or near pleural surface.
	pdr6Δ +Dexa	20%	> 80% alveoli contain encapsulated yeast	5 – 10 neutrophils 5 – 10 lymphocytes 20 – 30 histiocytes	Focal consolidated areas of dense inflammatory tissue up to 4 – 5 mm in dimension.

Figure 8

Table 4

DPI	Strain	% inflammation	Fungal infiltration	Quantitation of inflammatory immune cells/hpf 400X	Observations
14	Mock +Dexa +FLC	0%	None	-	-
	H99 +Dexa +FLC	15 – 20%	45 – 50% alveoli contain encapsulated yeast	15 – 30 neutrophils 15 – 30 lymphocytes < 5 histiocytes	Small to medium sized nests of inflammatory cells.
	pdr6Δ +Dexa +FLC	0%	10 – 15% alveoli contain encapsulated yeast	-	Inflammatory cells are uncommon with rare perivascular and peribronchial nests of small lymphocytes and few histiocytes.
50	Mock +Dexa +FLC	0%	None	Rare clusters of 2 – 4 histiocytes	Mildly dilated capillaries.
	pdr6Δ +Dexa +FLC	0%	75 – 80% alveoli contain encapsulated yeast	-	Few small foci of perivascular lymphocytes and rare histiocytes.

14 dpi

TOD

Figure 10   And Fig. S3 and S4.

Figure 11

Figure S2

Innate Immune Cell Gating Strategy

Adaptive Immune Cell Gating Strategy

Figure S3

Mock - 14 dpi   H99 - 14 dpi   *pdr6Δ* - 35 dpi

DAPI

α-CD68

Figure S4

Figure S4

Figure S5