Lack of an atypical PDR transporter generates an immunogenic *Cryptococcus neoformans* strain

that drives a dysregulated and lethal immune response in murine lungs

Christopher J. Winski[1], Peter V. Stuckey[1], Armando M. Marrufo[1], Robbi L. Ross[1,2], Georgina Agyei[1],

Sarah Chapman[5], and Felipe H. Santiago-Tirado[1,2,3,4,#]

[1] Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana, USA

[2] Integrated Biomedical Sciences, University of Notre Dame, Notre Dame, Indiana, USA

[3] Eck Institute for Global Health, University of Notre Dame, Notre Dame, Indiana, USA

[4] Warren Center for Drug Discovery, University of Notre Dame, Notre Dame, Indiana, USA

[5] Integrated Imaging Facility, University of Notre Dame, Notre Dame, Indiana, USA

Running Head: PDR transporter role during cryptococcal infection

#Address correspondence to Felipe H. Santiago-Tirado, fsantiago@nd.edu

Word Counts

Abstract: 215

Importance: 149

Text (excluding Methods, References, and captions): 5,836

ABSTRACT

Cryptococcus neoformans is an opportunistic fungal pathogen responsible for >150,000 deaths

every year with a mortality rate as high as 81%. This high medical burden is due, in part, to an

incomplete understanding of its pathogenesis. In a previous study, we identified a cryptococcal

atypical pleiotropic drug resistance (PDR) transporter, *PDR6*, that regulated antifungal resistance

and host interactions. Here, we follow-up on the role of *PDR6* in cryptococcal virulence. *In vivo*,

mice infected with the *pdr6Δ* strain display altered symptomatology and disease progression.

Specifically, we observed a significant increase in the innate immune cell populations in the

pdr6Δ-infected mice when compared to their WT-infected littermates. Furthermore,

quantification of pulmonary cytokines/chemokines revealed a robust increase of pro-

inflammatory cytokines in mice infected with the *pdr6Δ* mutant strain. Whereas antifungal

treatment of *pdr6Δ*-infected animals did not affect survival, treatment with a corticosteroid

significantly extended survival, highlighting the importance of a balanced/controlled host

immune response. We determined that the hyper-inflammatory immune response occurs, in

part, because the loss of the Pdr6 transporter indirectly alters the cryptococcal cell wall

architecture and results in the increased exposure of chitin, β-glucan, and other cryptococcal-

specific pathogen associated molecular patterns. Taken together, this study provides clinical

insights regarding cryptococcal pathogenesis while also providing additional functions of PDR-

type ATP-binding cassette (ABC) transporters in pathogenic fungi.

IMPORTANCE

Yeasts of the *Cryptococcus* genus, especially *C. neoformans*, can cause disease with unacceptably high mortality. This is due to delays in diagnostics, ineffective treatments, and an incomplete understanding of the interactions between this fungus and our immune system. In this study, we expand our knowledge of the biological function of the *PDR6* gene, particularly its effect on modulating the host's immune response. Normally, *C. neoformans*'s infections are characterized by an anti-inflammatory response that is unable to control the yeast. In the absence of *PDR6*, the response to the infection is a dysregulated pro-inflammatory response that initially controls the fungi but eventually results in death of the host due to too much tissue damage. This is due, in part, to an altered fungal surface. Given the dual role of *PDR6* in modulating antifungal sensitivity and immune responses, this work provides important insights that may lead to new or improved therapeutics.

INTRODUCTION

Commonly associated with soil and bird guano, *Cryptococcus neoformans* is an environmental, opportunistic yeast responsible for >194,000 infections every year (1). Upon inhalation of spores or desiccated yeast, *C. neoformans* reaches the alveoli where it encounters alveolar macrophages and other host immune cells (2); however, this fungus is capable of establishing a unique intracellular niche that allows it to survive and replicate within these host cells (3). The neurotropic fungus can then disseminate to extrapulmonary organs, traverse the blood-brain-barrier and enter the central nervous system (CNS), resulting in cryptococcal meningoencephalitis (CME) with a mortality rate as high as 81% (1, 4, 5). Several lines of evidence indicate that in addition to its use of host immune cells as replication havens, *C. neoformans* can use infected cells as vehicles for dissemination and brain invasion (5, 6). Hence, understanding this complex cryptococcal-phagocyte interaction from both fungal and host perspectives represents a challenging but clinically relevant avenue of research given its role in determining the course and outcome(s) of infection.

In addition to its ability to survive intracellularly, *C. neoformans* has several other virulence factors including an antiphagocytic polysaccharide capsule that provides protection from host stressors and masks several cryptococcal pathogen-associated molecular patterns (PAMPs) including chitin, chitosan, mannoproteins, and β-glucans from host phagocytes (7). Structurally composed of glucuronoxylomannan (GXM) and glucuronoxylomannogalactan (GXMGal), the capsule is also constitutively secreted/shed and it modulates the host immune response in ways that promote infection (8). Additionally, *C. neoformans* synthesizes cell wall-localized melanin that protects the yeast from ultraviolet radiation and oxidative/nitrosative

stress. Finally, the fungus displays thermotolerance, able to grow at physiological-relevant

conditions (37°C + 5% CO_2) (9-11). These virulence factors are integral to the pathogenesis of *C.*

neoformans and as such, have been well-studied. However, *C. neoformans* utilizes additional

virulence factors which remain significantly understudied or completely unknown. For example,

we previously identified *PDR6* (also known as *AFR3*), a novel ATP-binding cassette (ABC)

transporter of the pleiotropic drug resistance (PDR) class, and showed that it had a role in various

cryptococcal virulence facets (12, 13). Present only in plants and fungi, PDR transporters

hydrolyze ATP to transport substrates across biological membranes and have been primarily

implicated in antifungal resistance (14, 15). Indeed, deletion of *PDR6* (*pdr6Δ*) results in hyper-

susceptibility to the azole-class of antifungals, but also showed *in vitro* and *in vivo* defects in

virulence. Thus, the role of *PDR6* in fungal pathogenesis and modulating the host immune

response warranted further investigation.

During microbial infections, the host's immune system is tasked with mounting an

appropriate and balanced immune response that will clear the infection while hopefully limiting

tissue damage (16). An exaggerated Th1, or pro-inflammatory, response will likely clear a

microbial infection but also result in excessive cellular and tissue damage. Opposingly, an

inappropriate Th2, or anti-inflammatory, response will certainly limit host damage but allow for

uncontrolled microbial proliferation and dissemination. Many pathogens take advantage of this

dichotomy to promote their infection. For example, early in HIV infection, this virus promotes the

secretion of pro-inflammatory cytokines IFN-γ and TNF-α, generating a strong Th1 response that

is permissive to viral replication (17, 18). In the context of *C. neoformans* infections, the fungus

promotes an anti-inflammatory Th2 response that allows for intracellular survival and increased

replication and dissemination, ultimately resulting in patient mortality (19, 20). It is evident,

however, that between these two immune response extremes, there is a whole spectrum of

responses to microbial infections, making this topic extremely complex but also an important

target for therapeutics. Hence, increased understanding of the immune response against

Cryptococcus can inform future therapeutic strategies where the immune response could be

modulated to benefit the host.

Our previous work characterized a novel PDR transporter with important roles in

cryptococcal biology and presented evidence consistent with *PDR6* being a new fungal virulence

factor (12). Here, we study the role of *PDR6* in the establishment and progression of disease using

a murine inhalation model of cryptococcosis. We show that a lower inoculum and significantly

decreased *pdr6Δ* fungal burdens are still not sufficient to clear infection by this mutant. Instead,

the survival studies and histological analysis suggests that the *pdr6Δ* mutant elicits an altered

host immune response characterized by significant increases in pro-inflammatory

cytokines/chemokines and innate immune cells. We treated the *pdr6Δ*-infected mice with a

corticosteroid and significantly extended their survival despite little to no change in fungal

burden. This dysregulated immune response might be a consequence of fungal surface

alterations that expose antigenic and immunogenic components that normally would be hidden

from the immune system. Collectively, this study furthers our understanding of cryptococcal

pathogenesis, highlights the clinical importance of the host immune response in controlling

fungal infections, while also providing novel functions and insights of fungal PDR-type

transporters in the context of disease progression.

RESULTS

***pdr6*Δ-infected mice succumb to infection despite decreased inoculum and fungal burden.** In our previous study we determined that *PDR6* was required for complete cryptococcal virulence as mice infected with the *pdr6*Δ mutant lived significantly longer and had decreased fungal burdens when compared to their H99-infected littermates (12). Consistently, this finding was recapitulated by *Oliveira et al.* using the *Galleria mellonella* invertebrate infection model (13). Notably, the *pdr6*Δ-infected animals displayed altered symptomatology compared to their H99-infected littermates. Specifically, at endpoint, the WT-infected mice displayed altered behavior associated with neurological deficiencies, traditional of cryptococcal meningitis; in contrast, the *pdr6*Δ-infected mice exhibited respiratory symptoms characterized by rapid and labored breathing, suggestive of pneumonia. We hypothesized that the altered symptomatology could be explained, in part, by differences in the establishment and progression of disease. Previously, fungal burdens were only quantified at time of death (TOD), potentially overlooking differences earlier on in the infection. To obtain a comprehensive view of the infection over time, we inoculated mice with 5×10^4 cells of H99 or *pdr6*Δ strains, and quantified fungal burden every seven days over the course of the infection (Fig. 1A). Interestingly, we found that at 7 days post infection (dpi) there were no differences in the fungal burdens between the H99 and *pdr6*Δ-infected animals in all organs tested, suggesting that the strains established infection similarly. Following this initial establishment, however, at 14 dpi there was a significant decrease in fungal burden for the *pdr6*Δ strain compared to the WT. At 17 dpi, the H99-infected mice succumbed to the infection with lung, brain, and spleen burdens of 3.7×10^8, 1.9×10^6, and 1.3×10^4 CFU while the *pdr6*Δ-infected mice CFUs were 3.6×10^7, 2.3×10^4, and 677, respectively. Between 14 and

28 dpi, the fungal burden of the *pdr6Δ*-infected animals remains relatively constant in the brain

and spleen, with a gradual, although small, increase in the lungs. Then, after 28 dpi, there are

gradual increases in all organs with final CFU counts of 5.5 x 10^7, 1.1 x 10^5, and 1.5 x 10^3 in the

lungs, brain, and spleen, respectively. The differences in CFUs highlight key differences between

H99 and *pdr6Δ* disease progression and suggests a potential role of the adaptive immune

response given the changes at 14 dpi.

Experimentally, the mice in our previous studies were inoculated with 5 x 10^4 cells, a

common inoculum for the murine inhalation model of cryptococcosis. Given the *in vitro* defects

of the *pdr6Δ* strain (12), and the fact that initially, the mice were able to control the infection

(Fig. 1A), we hypothesized that by lowering the fungal inoculum, the survival of the *pdr6Δ*-

infected mice would be extended with a potential for complete clearance of the infection. To

assess this, we decreased the initial inoculum 10-fold and infected mice with 5 x 10^3 cells and

monitored survival (Fig. 1B). The H99-infected mice had a mean survival of 21 days and the *pdr6Δ*-

infected mice displayed a mean survival of 36 days, similar to our studies with 10-fold higher

inoculum. Quantification of the fungal burden showed similar CFUs in all organs between the

animals infected with low and high WT inoculum (Fig. 1C compared to 1A); however, despite

having similar lungs CFUs, the animals infected with low *pdr6Δ* inoculum had lower brain and

spleen CFUs at TOD relative to the higher inoculum (Fig. 1C compared to 1A). Evidently, using a

lower inoculum did not change the trend we have seen before: the *pdr6Δ*-infected mice still

survive longer with significant decreases in fungal burden across all organs. However, notably, in

the *pdr6Δ*-infected mice the lower inoculum results in lower overall organ burdens, with less

dissemination, yet the *pdr6Δ*-infected mice still succumbed to the infection with the same

dynamics as when using a higher inoculum (35-38 days; Fig. 1B).

Infection with *pdr6Δ* results in increased tissue damage. Despite initially colonizing the

pulmonary environment, the *Cryptococcus*-associated fatal pathology occurs in the brain due to

the meningoencephalitis (21, 22). The altered symptomatology coupled with the differential

disease progression suggests that the *pdr6Δ* strain likely affects the host immune response.

Furthermore, when performing the previous fungal burden experiments, we observed significant

phenotypic differences in lung sizes in H99 v. *pdr6Δ*-infected mice (Fig. 2). To qualitatively assess

any potential differences at the cellular and tissue level, we harvested the lungs of mock, H99,

and *pdr6Δ*-infected mice, stained with hematoxylin and eosin (H&E), and assessed tissue damage

(Fig. 2 and Table 1). At 14 dpi, the gross anatomical images clearly show increased size and

inflammation of the H99-infected lungs when compared to the *pdr6Δ*-infected. Additionally, the

10X and 40X images display deterioration of the lung parenchyma and infiltration of *C.*

neoformans which is not observed in the *pdr6Δ*-infected lungs (Fig. 2A). At 35 dpi, the *pdr6Δ*-

infected lungs are severely inflamed, infiltrated with yeast, and characterized by the presence of

many inflammatory immune cells in the peri-bronchial and perivascular space (Fig. 2B). Table 1

provides a quantitative summary comparing the H99 and *pdr6Δ*-infected pulmonary tissue over

time. Consistent with our observations in the fungal burden experiments, there were little to no

differences at 7 dpi. However, at 14 dpi the *pdr6Δ*-infected lung contains the yeast in granuloma-

like regions and have 30% less *C. neoformans* than the WT-infected tissues. Furthermore, an

increase in lymphocytes is observed in the *pdr6Δ*-infected lungs relative to WT (105 to 26,

respectively; Table 1). Finally, as the *pdr6Δ*-infected mice approach TOD, the fungal infiltration,

inflammation, and number of inflammatory immune cells in the lung tissue all increase to levels

well above those observed in the WT-infected animals at TOD. Altogether, these findings suggest

that the *pdr6Δ* strain elicits a differential immune response in the pulmonary environment thus

affecting disease progression and symptomatology.

Fluconazole treatment improves fungal burden but not outcome in *pdr6Δ*-infected mice.

Currently, there a very few antifungal drugs available to treat cryptococcal infections and the

compounds that are available are either not effective, extremely expensive, or associated with

deleterious side effects (23). Fluconazole (FLC), a fungistatic azole class antifungal, is the mainstay

treatment option in low-resource regions where cryptococcal infections are endemic.

Mechanistically, FLC inhibits lanosterol 14α-demethylase, an enzyme in the ergosterol

biosynthesis pathway encoded by *ERG11* (24). In our previous study, we showed that the deletion

of *PDR6* increased the susceptibility of *C. neoformans* to azole class antifungals including FLC (12).

Given this phenotype, we hypothesized that treatment of *C. neoformans pdr6Δ*-infected mice

with FLC would result in total clearance of the infection, preventing the lethal immune pathology.

To test this, mice infected with either the WT or *pdr6Δ* strains, or mock infected, were treated

daily (starting 1 dpi) with either a low (8 mg/kg) or high dose (16 mg/kg) of FLC and monitored

for survival (Fig. 3A). The FLC treatment significantly extended survival in both the H99 and *pdr6Δ*-

infected animals compared to the untreated controls. Unfortunately, survival was extended to

the same extent for both groups. Numerically, the mean survival for the untreated H99-infected

mice was 17 days and the low and high doses extended survival 4.5 and 6 days, respectively.

Similarly, the mean survival for the untreated *pdr6Δ*-infected mice was 34 days and the low and

high doses extended survival 5 and 7 days, respectively. Surprisingly, although the survival was

similarly extended between strains, quantification of the fungal burden revealed significant

differences (Fig. 3B-G). At 14 dpi, both doses of the FLC treatment significantly decreased the

fungal burden in the lungs, brain, and spleen only in the *pdr6Δ*-infected mice (Fig. 3B-D). Similarly,

at TOD, the daily FLC treatment continued to lower the fungal burden of the *pdr6Δ*-infected mice

even more across all three organs, with several mice where the brain and spleen CFU counts were

at the limit of detection (LOD) (Fig. 3E-G). Not surprisingly, neither of the FLC doses had a

significant effect on WT-infected mice. This observation is interesting because despite

significantly decreasing the fungal burden with the FLC treatment, the *pdr6Δ*-infected animals

were still succumbing to the infection.

To understand how the FLC treatment was affecting the pulmonary environment, we

prepared slides and performed qualitative histological analysis at TOD for both WT and *pdr6Δ*-

infected mice (Fig. 4 and Table 2). The H99-infected lungs at 14 dpi treated with FLC displayed

similar features to the previously pictured H99 untreated lungs, namely a high percentage of

fungal infiltration and tissue breakdown. In contrast, the *pdr6Δ*-infected lungs at 35 dpi treated

with FLC appear to have the infection controlled in granuloma-like regions (Fig. 4, inset 2) with

widespread immune cell infiltration (Fig. 4, inset 1). Interestingly, despite reduced pulmonary

CFUs (Fig. 3 and Table 2) and decreased inflammation (Fig. 4 and Table 2), the *pdr6Δ*-infected

mice were still succumbing to the infection. Together, these data suggest that the mice infected

with the *pdr6Δ*-strain are not necessarily dying directly because of the *C. neoformans* infection,

but rather indirectly because of an inappropriate host immune response.

***C. neoformans pdr6Δ* strain elicits a strong pro-inflammatory immune response.** During

microbial infections, the host's immune response ultimately determines the clinical outcome

(25). Ideally, a balanced Th1 pro-inflammatory and Th2 anti-inflammatory immune response will

allow for microbial clearance and subsequent tissue repair. Previous studies have determined

that *C. neoformans* skews the immune response towards a permissive anti-inflammatory Th2

response, thus benefiting the yeast and allowing for fungal replication and subsequent

dissemination (26). Given our previous observations, we hypothesize that the *pdr6Δ* strain is

instead generating a strong pro-inflammatory immune response that is able to control the

infection initially, but also damaging large areas of pulmonary tissue. To assess this, we quantified

the secreted cytokines and chemokines profile in the lungs and brains of H99 and *pdr6Δ*-infected

mice at their respective TODs (Fig. 5).   Interestingly, we found that several of the pro-

inflammatory cytokines are significantly upregulated in the *pdr6Δ*-infected lungs relative to WT

(Fig. 5A). Specifically, the production of IL-1α, IL-1β, and MIP-1β (macrophage inflammatory

protein 1β) is increased 13-fold, 23-fold, and 29-fold respectively, when compared to the lungs

of the H99-infected mice. Additionally, the expression of several anti-inflammatory cytokines like

IL-9, IL-13, and IL-33 are lower in the *pdr6Δ*-infected mice relative to WT. A similar trend was

observed in the brain although not to the same extent as that observed in the pulmonary tissues

(Fig. 5B). Specifically, the pro-inflammatory cytokines/chemokines IL-1β, KC (keratinocyte

chemoattractant), and TNF-α (tumor necrosis factor α) are all highly expressed when normalized

to the H99-infected brain, and the anti-inflammatory cytokines IL-4 and IL-9 display decreased

expression in the brain when compared to WT. The decreased capacity of the *pdr6Δ* mutant to

survive in host phagocytes and disseminate coupled with the immune-privileged status of the

CNS may explain the weakened expression of inflammatory cytokines in the brain. During a

typical cryptococcal infection, the expression of anti-inflammatory cytokines/chemokines are

increased thus generating a Th2 response, however, in the context of an infection with the *pdr6Δ*

strain, the secretion of pro-inflammatory cytokines/chemokines is significantly increased

resulting in a damaging Th1 immune response.

Corticosteroid treatment extends survival in *pdr6Δ*-infected mice. Clinically, corticosteroids like

dexamethasone are prescribed to downregulate the immune system. Mechanistically, these anti-

inflammatory drugs function by suppressing the migration of neutrophils and decreasing

lymphocyte colony proliferation (27). The potential of dexamethasone as a treatment strategy

during cryptococcal infection was investigated in a clinical trial but was prematurely concluded

because of adverse effects and increased mortality (28, 29); however, corticosteroids have been

used successfully to treat CME-associated inflammatory disorders such as postinfectious

inflammatory response syndrome and immune reconstitution inflammatory syndrome (21).

Given the robust pro-inflammatory immune response observed in the *pdr6Δ*-infected mice, we

hypothesized dexamethasone treatment would dampen or knock down the aggressive Th1

response resulting in fungal clearance and improved survival. Experimentally, we infected mice

with either the H99 or *pdr6Δ* strain, or mock infected, and treated them daily (starting at 1 dpi)

with either 0.3 mg/kg dexamethasone or a dexamethasone and 16 mg/kg FLC combination and

assessed survival (Fig. 6A). The H99-infected mice that received the dexamethasone treatment

displayed a significant reduction in survival (mean survival of 13 days vs 17 days) when compared

to the untreated H99 control, while the combo therapy restored survival to that of the untreated

H99-infected group (mean survival of 16 days vs 17 days). This result corroborates the

corticosteroid finding reported in the human clinical trial. In terms of the *pdr6Δ*-infected mice,

both the dexamethasone alone and in combination with FLC significantly extended survival.

Numerically, the mean survival of the untreated *pdr6Δ*-infected control group was 34 days and

was extended to 55 and 58 days in the dexamethasone and combination treated groups,

respectively. The survival studies were stopped after 60 dpi because the remaining mice,

including the mock-infected receiving the daily dexamethasone treatment (not shown),

appeared sick despite not meeting endpoint criteria. We concluded that after two months of

daily corticosteroid injections the immune systems of the mice had been ablated resulting in their

poor appearances. Despite the dexamethasone treatment extending survival in the *pdr6Δ*-

infected animals, the fungal burdens were differentially affected over the course of the

experiment (Fig. 6B-G). Specifically, at 14 dpi the addition of dexamethasone significantly

decreased the fungal burden in the lungs in both the H99 and *pdr6Δ*-infected mice and little to

no changes were observed in either strain in the brain or spleen CFUs (Fig. 6B-D). At TOD the

dexamethasone treatment in the *pdr6Δ*-infected animals significantly increased the lung CFUs

and the combination therapy returned it to levels comparable to the untreated *pdr6Δ*-infected

control (Fig. 6E). Although not statistically significant, a similar trend was observed in the brain

and spleen of the *pdr6Δ*-infected mice (Fig. 6F & G). These effects were not seen in the WT-

infected mice. The dynamic changes in fungal burdens in the lungs of the dexamethasone and

combination treated *pdr6Δ*-infected mice can be explained by the absence or lack of

inflammatory immune cells to control the infection and the addition of FLC lowers the CFU as the

fungistatic nature of the antifungal is unable to clear the initial infection. Collectively, these data

further confirm and highlight the altered disease progression elicited by the *pdr6Δ* mutant and

more importantly demonstrate the positive clinical outcome associated with controlling the

inappropriate pro-inflammatory immune response.

Combination dexamethasone and fluconazole treatment reduces pulmonary tissue damage in

***pdr6Δ*-infected animals.** Based on our previous results, it is evident that the addition of the

corticosteroid is benefiting the *pdr6Δ*-infected mice by significantly extending survival.

Histological analysis of the lung tissue at TOD for both WT and *pdr6Δ*-infected mice was

performed to visualize changes in fungal infiltration, tissue inflammation, and immune cell

recruitment (Fig. 7 and Table 3). Visually, the treatment with dexamethasone had a profound

effect given the qualitative reduction in inflammatory cells across all conditions. The lungs of the

H99-infected mice that received dexamethasone are not as inflamed as what was observed on

our previous slides but the destruction of the lung tissue is devastating (Fig. 7A). Clearly the lack

of immune cells allowed for the uncontrolled proliferation of *C. neoformans* resulting in increased

damage of the pulmonary tissue. Similarly, the dexamethasone treated lungs of the *pdr6Δ*-

infected display intense fungal proliferation but there is qualitatively less damage when

compared to the H99-infected animals (Fig. 7B and Table 3). The combination dexamethasone

and FLC treatment reduced the fungal infiltration in both the H99 and *pdr6Δ*-infected mice when

compared to the dexamethasone alone (Fig. 8 and Table 4). In fact, despite the presence of some

yeast, the *pdr6Δ*-infected lung tissue appears relatively healthy with little to no damage and

inflammation suggesting that another mechanism may be responsible for the mice reaching

endpoint (Fig. 8B and Table 4).

***pdr6Δ*-infected mice display an increase in innate immune cells populations.** During microbial

infections, the cells of the innate immune system represent the first line of defense which is

followed by the activation and recruitment of the adaptive immune response (30). In a *C.*

neoformans infection, lung macrophages represent the innate population primarily responsible

for initial clearance, although some recent literature suggests neutrophils also play an important

role (31-33). However, CME is an AIDS-defining disease, and low $CD4^+$ T-cell counts is a major

susceptibility for CME, indicating that the adaptive immune response also plays an important role

controlling cryptococcal infections (34). Provided the altered immune response to the *pdr6Δ*

strain, we expected alterations to both the quantity and the composition of innate and adaptive

inflammatory immune cells. To test this, we harvested leukocytes from the lungs of H99 and

pdr6Δ-infected mice at 14 dpi (close to TOD of WT-infected mice) and at 35 dpi (close to TOD of

pdr6Δ-infected mice) and quantified the different populations using flow cytometry (Fig. 9).

Quantification of the total innate immune cell populations at 14 dpi showed that the FLC,

dexamethasone, and combination treatments led to a significant decrease in the *pdr6Δ*-infected

animals whereas there were no significant changes in the H99-infected mice (Fig. 9A). A similar

trend was observed when compared as a percentage of $CD45^+$ cells. Interestingly, a significant

increase in neutrophils and macrophages was seen in the combination-treated *pdr6Δ*-infected

mice (Fig. 9B). At this time (14 dpi), quantification of the total adaptive immune cells showed a

marked increase in the *pdr6Δ*-infected animals when compared their H99-infected counterparts,

although only the CD8[+] T-cells were statistically significant (Fig. 9C, untreated bars). Furthermore,

all treatments decrease the number of adaptive immune cells in the *pdr6Δ*-infected animals

relative to the WT, specially the CD8[+] T-cells (Fig. 9C, FLC, Dexa, and +Dexa +FLC bars). The same

trend is observed when compared as percentage of CD45+ cells (Fig. 9D). At 35 dpi there are

some minor differences in the adaptive immune cell populations when comparing the treated

groups (Fig. 9G & H) but the interesting finding is observed in the innate immune cell populations

(Fig. 9E & F). Specifically, a significant increase in macrophages is observed in the *pdr6Δ*-infected

mice when compared to the H99-infected (Fig. 9E & F, untreated bars). Furthermore, the

dexamethasone treatment in the *pdr6Δ*-infected animals significantly decreases the number of

macrophages which is replaced by a significant increase in the number of neutrophils (Fig. 9E &

F, Dexa bars). It appears that in the absence of Pdr6, the neutrophil response is more robust and

sustained than the macrophage response, whereas is the opposite in the WT-infected mice.

Collectively, these results showed that infection with the *pdr6Δ* strain results in the increased

recruitment of adaptive immune cells early on during the infection while an increase in innate

immune cells, specifically macrophages, was observed in the later stages of the disease. This

further explains the strong pro-inflammatory response, altered disease progression, and

differential symptomatology specific to the *pdr6Δ* strain.

***C. neoformans pdr6Δ* strain exhibits increased PAMP exposure.** Up to this point we have shown

that the infection of our mouse model with the *C. neoformans pdr6Δ* strain results in the

recruitment of more inflammatory immune cells leading to increased pulmonary damage and

mortality. Further support was provided in the dexamethasone treatment experiments, where

dampening the immune response significantly extended survival in the *pdr6Δ*-infected animals.

Experimentally, we have studied the host response to this cryptococcal mutant but the question

that remains is how is this mutant modulating the immune system and generating a robust Th1

response? In our previous work, we showed that *PDR6* encodes an atypical transporter localized

to the endoplasmic reticulum (ER) and plasma membrane (PM) involved in ergosterol trafficking

(12). Others have suggested a direct relationship between altered ergosterol levels and host

immune responses (35, 36) as well as altered surface composition due to defects in PM fluidity

and lipid rafts (37, 38) We hypothesized that in deleting this transport protein there would be

less ergosterol in the PM, potentially resulting in downstream alterations to the cryptococcal

surface and an increase in exposed immunogenic components. To assess this, we grew H99 and

pdr6Δ cells in YPD, stained with fluorescent dyes that recognize and bind to fungal cell wall

components, and visualized the cells using microscopy. Under this condition, we did not observe

any difference between the H99 and *pdr6Δ* stained cells (data not shown). However, previous

studies have reported a correlation between growth conditions and the architecture of the fungal

cell wall (39). This led us to repeat the experiment with cells grown under host-like conditions

(DMEM at 37^{o}C + 5% CO_2) (Fig. 10A). Under these conditions, the *pdr6Δ* strain displayed increase

staining with calcofluor white (CFW) and concanavalin A (ConA) suggestive of increased exposure

of chitin and mannoproteins, respectively. Similarly, there was an increase in binding of the β-

(1,3)-glucan antibody suggestive of more exposed β-glucans in the *pdr6Δ* mutant (Fig. 10A).

Quantification of CFW, ConA, and eosin Y (EoY), which binds/stains chitosan, using flow

cytometry revealed a significant increase in the binding of all the stains in the *pdr6Δ* strain when

compared to the WT (Fig. 10B). The *cap59Δ* mutant, deficient in polysaccharide capsule synthesis

and thus with increased exposure of its cell wall, was used as a positive control. Having previously

shown that the pdr6 mutant has no defect in capsule induction, we hypothesized that alterations

in the capsular architecture could justify the increase in exposed cryptococcal PAMPs. To assess

this, we used several anti-polysaccharide capsule antibodies to stain the capsules of H99, *pdr6Δ*,

and *cap59Δ* cells (Fig. 10C). A significant increase in staining and penetration of the antibody was

observed in the *pdr6Δ* mutant when using the mAb 302. This antibody only recognizes and binds

acetylated GXM (40) and suggests that the integrity and/or porousness of the polysaccharide

capsule in the *pdr6Δ* strain is altered when compared to the structurally-sound WT capsule.

Altogether, these findings suggest that the deletion of *PDR6* disrupts the cryptococcal cell wall

composition and the capsule density (since it is induced to normal size in the *pdr6Δ* mutant),

allowing for increased exposure of fungal PAMPs, namely chitin, mannoproteins, and β-glucans.

This observed increase in PAMP exposure in the *pdr6Δ* strain can mechanistically explain the

increased pro-inflammatory response by the host immune system.

DISCUSSION

Fungal PDR transporters are primarily considered virulence factors because of their role in

antifungal resistance. For example, the overexpression of *AFR1* in *C. neoformans* and *CDR1* in *C.*

albicans clinical strains increase their resistance to many of the azole class antifungals and

complicates the treatment strategy (41, 42). Structurally, these two proteins (and all other well-

characterized fungal PDR transporters) are 'full-length' consisting of two nucleotide-binding

domains (NBDs) and two transmembrane domains (TMDs), which is required for a functional

fungal PDR protein (15, 43). Here, we continue the characterization of *C. neoformans PDR6*, an

atypical and novel 'half-size' PDR transporter since it only possesses one NBD and one TMD.

Given the asymmetrical nature of PDR transporters and the conserved 'degenerated' sequence

in one of the NBDs, it is unlikely that Pdr6 homo-dimerizes and instead hetero-dimerizes with

another half-size PDR protein present in the cryptococcal genome (of which there are two more)

to form a functional transporter (14). An interesting future direction includes the identification

and subsequent deletion of Pdr6's binding partner which should generate phenotypes similar to

those observed in the *pdr6Δ* strain. Nevertheless, we found that in addition to azole resistance,

PDR6 modulates the host immune response, a phenotype that may be unique to 'half-size' PDR

transporters (12).

The *pdr6Δ*-dependent alterations in survival, fungal burden, and symptoms all suggested

a differential host response. Specifically, the fact that the significant decrease in fungal burden

across all organs coupled with the lower inoculum still resulted in mortality showed that the mice

infected with the *pdr6Δ* mutant were not succumbing because of the fungus, but because of an

inappropriate and damaging immune response. Consistently, the treatment of the *pdr6Δ*-

infected mice with FLC decreased the fungal burden in all organs, and in some to the limit of

detection, and yet the mice still died, implicating the immune system. Furthermore, the

divergence in symptoms between the WT and *pdr6Δ*-infected animals at TOD was clinically

relevant. Although the *pdr6Δ*-infected mice displayed a pneumonia-like symptomatology rather

than the expected neurological symptoms, they still had CFUs in the brain so CME cannot be ruled

out; however, we think that it was below the threshold to trigger the symptoms and the

neurological deficiencies characteristic of CME. The histological analysis, immune cell

quantification, and cytokine/chemokine expression profiles all showed an aggressive lung pro-

inflammatory immune response against the *pdr6Δ* mutant, while the response in the brain was

more muted, substantiating our claim. After understanding that the *pdr6Δ*-infected mice were

dying because of a sustained Th1 immune response, we expected the corticosteroid treatment

experiment would result in total clearance of the infection. The dosages of the dexamethasone

and FLC were based off clinical reports and we purposefully used the lower dexamethasone dose

given the extended length of the proposed experiment (28). We show that the addition of

dexamethasone significantly extended the survival of the *pdr6Δ*-infected mice although the

infection was not cleared even with the addition of FLC. Notably, the combo treatment reduced

both inflammation and fungal burden in the lungs and restored tissue architecture almost to

normal, but the outcome was not significantly different. Optimizing the timing of the

dexamethasone treatments, such that it is not administered for the entirety of the experiment,

or the doses are tapered after an initial pulse, warrants further investigation and represents a

future direction. Since corticosteroids are immunosuppressive, and the i.p. injection of

dexamethasone will dampen the immune response in the whole body, it is possible that while

eliminating the tissue damage in the lungs, the treatment also inhibits the clearance of the yeast

in other organs (most likely kidneys, liver, or heart), explaining the eventual death after extending

survival for more than three weeks. In that sense, if the dexamethasone could be targeted to

specifically affect the lung tissue, as with the use of an inhaler, we would expect a better

outcome.

The *in vivo* experimentation allowed us to comprehend what was occurring within the

host, but why was this specifically occurring with the *pdr6Δ* strain? Our data showing that the

pdr6Δ mutant displays increased PAMP exposure is a reasonable explanation for the pro-

inflammatory response. However, it may not be the only reason as the acapsular *cap59Δ* mutant

also exhibits increased PAMP exposure and it is avirulent in the same mouse model. Furthermore,

we previously showed that conditioned media (CM) from *pdr6Δ* cultures modulates the

phagocytosis of *C. neoformans* by macrophages, whereas the CM from *cap59Δ* or other PDR

transporter mutants had little to no effect (12), suggestive of an immunomodulatory Pdr6-

dependent secreted factor. Nevertheless, all these data identify and support Pdr6 as a potential

therapeutic target. Anticryptococcal treatments are limited by toxicity, costs, and rising microbial

resistance, but FLC, despite being sub-optimal, still is the antifungal of choice in many regions of

the world because it is cheap, relatively less toxic, and broad spectrum. Since the deletion of

PDR6 results in hypersensitivity to FLC, the design and synthesis of a novel compound that targets

Pdr6, coupled with FLC and/or a corticosteroid, may have clinical potential. Alternatively, several

groups in the cryptococcal and immunology fields have tried to identify potential vaccine

candidates against this fungus (44). Collectively, these preliminary vaccines, elicit a pro-

inflammatory response allowing for protection against subsequent challenges (45-47). However,

this is hampered because the immune mechanisms that elicit a protective immune response are

not known yet. Thus, the knowledge generated in this study has potential to help inform

cryptococcal vaccine development by providing novel insights into what constitutes a deleterious

pro-inflammatory response against *C. neoformans* infection.

Finally, the complexities of PDR half-size transporters still remain unknown. It is

understood that these proteins are promiscuous and transport a diverse array of substrates, and

coupled with the expanded repertoire of PDR genes in *Cryptococcus*, it is not surprising to find

roles other than traditional efflux pumps and subsequent resistance to xenobiotics. *PDR6* has

also been implicated in aging and replicative life span (13), and most recently, *PDR9*, a full-length

PDR gene in the same clade as *AFR1*, was found to play a role in CO_2 tolerance via an unknown

mechanism that may involve phospholipid flipping (48). Hence, *PDR6* may act through multiple

direct and indirect mechanisms to modulate the inflammatory host response, such as by

secreting immunomodulatory molecules, aiding in maintaining the correct PM fluidity by

regulating ergosterol transport, and preventing exposure of surface PAMPs by supporting correct

cell wall and/or capsule architecture. All of these functions could be mediated by Pdr6-

dependent metabolites, lipids, or proteins, supporting future studies in these areas such as

comparative metabolomics of WT and *pdr6Δ* CM.

In conclusion, we have further characterized a novel virulence factor in *C. neoformans* by

studying the role of *PDR6* in a mouse model of cryptococcosis and examining the altered immune

response. Mice infected with the *pdr6Δ* strain still perished despite a lower initial inoculum, lower

organ burdens over the course of infection, significantly decreased fungal CFUs when treated

with FLC, and muting of an overexuberant inflammatory response when treated with

dexamethasone. *PDR6* is not necessarily involved in the establishment of disease, as there was

no difference in CFUs 7 dpi relative to WT, but it is critical in the progression of cryptococcosis as

mice succumbed to a fulminant pneumonia rather than CME. This is because in its absence, there

is increased exposure of immunomodulatory chitin, chitosan, mannoproteins, and β-glucans,

contributing to the secretion of pro-inflammatory cytokines and chemokines which increase the

number of macrophages and other inflammatory immune cells in the lungs, containing the

infection but causing devasting tissue damage and inflammation. Elucidating the role of *PDR6* in

the context of infection improves our understanding of cryptococcal pathogenesis while also

informing potential clinically relevant therapeutic studies such as the search for novel drug

targets or the development of a vaccine. More broadly, this study expands the field of fungal PDR

transporters and given the conserved nature of these proteins, these findings may be applicable

to other fungal pathogens.

MATERIALS & METHODS

Strains, media, and growth conditions.

All *C. neoformans* strains used in this study were in the serotype A H99α background. All fungal

strains were maintained at -80°C and grown at 30°C on yeast peptone dextrose (YPD) with

antibiotics (NAT or G418) as appropriate.

Murine survival studies

In vivo fungal virulence was tested in our murine inhalation model of cryptococcosis (12, 49-51).

Briefly, 5- to 6-week-old female A/J inbred mice (The Jackson Laboratories) were anesthetized

with a ketamine rodent cocktail delivered i.p. and intranasally inoculated, unless otherwise

stated, with 5.0 x 10^4 cryptococcal cells in 40 μL of sterile PBS. Animals were closely monitored

and humanely euthanized if they lost >20% weight relative to peak weight, or at predetermined

timepoints, or at the end of the experiment (maximum of 60 days).

For drug treatment studies, cryptococcal-infected mice were randomly assigned into treated or

non-treated control groups. Stock solutions of fluconazole (1.6 mg/mL), dexamethasone (0.03

mg/mL), and a fluconazole/dexamethasone combo were prepared in sterile PBS. The drug(s)

were administered intraperitoneally at doses of 16 mg/kg/day and 8 mg/kg/day for fluconazole

and 0.3 mg/kg/day for dexamethasone. In survival studies, treatment began 24 h after infection

and continued throughout the experiment until endpoint was reached. Statistically significant

differences between survival curves were determined by log rank Mantel-Cox test (GraphPad

Prism).

Fungal Burden Quantification

Infected mice were humanely euthanized at predetermined time points or TOD by isoflurane

overdose followed by cervical dislocation, and lungs, brain, and spleens were harvested. Tissues

were homogenized in 5 mL of sterile PBS, diluted, and plated on YPD agar medium supplemented

with ampicillin. CFUs were counted following incubation at 30℃ for 48 h. Statistical significance

was determined using ANOVA Kruskal-Wallis test (GraphPad Prism).

Histological analyses

A/J female mice (The Jackson Laboratories) were infected via the intranasal route as described

above with 5 x 10^4 CFU of either WT (H99) or the *pdr6Δ* mutant strain. Mice were sacrificed as

predetermined time points (14, 35, 50 days post inoculation) by isoflurane overdose followed by

cervical dislocation. Lungs were perfused with sterile PBS and 10% neutralized formalin. Lungs

were subsequently paraffin embedded, sectioned, mounted, and stained with H&E by the

University of Notre Dame Histology Core facility. H&E-stained slides were imaged at 10X and 40X

using an inverted Zeiss microscope. Slides were scored in a blind manner by the resident

pathologist at the Harper Cancer Research Institute [W.J.K.].

Cytokine Analyses

A/J female mice from Jackson Laboratories were infected via the intranasal route with WT (H99)

or the *pdr6Δ* mutant strain as described above. Mice were euthanized at predetermined time

points by isoflurane overdose followed by cervical dislocation. Cytokine levels within the lung and

brain homogenates of infected mice were analyzed using the Proteome Profiler Mouse XL

Cytokine Array (R&D Systems Inc., Minneapolis, MN). The relative expression levels of 111 soluble

mouse proteins were determined following the manufacturer's protocol. Array membranes were

imaged using LI-COR Odyssey Infrared Imaging System and quantified using ImageStudio

software.

Cytokine/chemokine expression levels were further confirmed using the Bio-Plex Multi-Plex

Assay Kit (Bio-Rad Laboratories, Hercules, CA). Supernatants from organ homogenates were

used to determine presence of 22 proteins including eotaxin, G-CSF, GM-CSF, IFN-γ, IL-1α, IL-1β,

IL-2, IL-3, IL-4, IL-6, IL-9, IL-10, IL-12 (p40), IL-12 (p70), IL-13, IL-17A, KC, MCP-1, MCP-1α, MIP-1β,

RANTES, and TNF-α.

Pulmonary leukocyte isolation and flow cytometric analysis

H99 and *pdr6Δ*-infected mice were euthanized at predetermined time points and TOD by

isoflurane overdose followed by cervical dislocation. Lungs of infected mice were excised and

digested enzymatically at 37^{o}C for 60 min in 10 mL of digest buffer (0.34 U/mL Liberase^{TM} (Roche)

and 100 ug/mL DNaseI (ThermoFisher)) in 1X DPBS. After 60 min, digestion was arrested with 1X

DPBS supplemented with 2% FBS and 0.2 um EDTA. The cell suspension was washed and passed

through 40-μm cell strainers.

The single cell suspensions were stained with a live/dead far red viability dye (Invitrogen),

washed, and incubated on ice for 20 min with purified anti-mouse CD16/CD32 antibodies

(BioLegend) to block Fc receptors. Following blocking, cells were incubated with fluorochrome-

conjugated antibodies to allow for multistaining for 45 min on ice. The innate panel consisted of

PE-Cy5 conjugated anti-CD45 (30-111), Brilliant Violet (BV) 650-conjugated anti-CD11b (M1/70),

Super Bright (SB) 780-conjugated anti-F4/80 (BM8), BV711-conjugated anti-Ly6G (1A8), PE-Cy7-

conjugated anti-Ly6C (HK1.4), PE-conjugated anti-SiglecF (1RNM44N), and APC-eFluor 780-

conjugated anti-CD11c (N418). The adaptive panel included PE-Cy5 conjugated anti-CD45 (30-

111), AF700-conjugated anti-CD3 (145-2C11), BV421-conjugated anti-CD4 (GK1.5), BV510-

conjugated anti-CD25 (PC61-5), PE-conjugated anti-CD8 (53-6.7), and AF488-conjugated anti-

FoxP3 (FJK-16s). Cells were washed and fixed with 2% formalin in FACS buffer. UltraComp eBeads

(Invitrogen) were stained with each respective fluorochrome-conjugated antibody and used to

determine and spillover/compensation calculations. Data was analyzed using FlowJo software.

Cell wall and capsule stains

Starter cultures were grown overnight in YPD, washed with DMEM, and 1 x 10^6 cells/mL were

added to 24-well tissue culture plates. Plates were incubated at 37°C + 5% CO2 for 48 h. The cells

were collected by centrifugation at 3,000 x g for 5 min. For staining, Calcofluor White (CFW),

Alexa Fluor 488-conjugated concanavalin A (ConA), β-(1,3)-glucan antibody 2GB (ab233743;

Abcam, USA), eosin Y (EoY), and the capsular antibody 302 were used. Cells were washed and

stained with CFW, ConA, β-(1,3)-glucan antibody, EoY, and Mab 302 at 100 μg/mL, 50 μg/mL, 20

μg/mL, 500 μg/mL, and 10 μg/mL, respectively. Cells were incubated with the stains for 30 min,

washed, fixed, and imaged at 100X using an inverted Zeiss microscope. Fluorescence of mAb 302

was calculated using Cell Profiler software. Flow cytometry data was acquired using a BD

LSRFortessa cytometer. The CFW signal was detected using a 405 nm laser and the ConA and EoY

signals were detected using a 488 nm laser. Flow cytometry data was analyzed with FlowJo

software.

ACKNOWLEDGMENTS

We acknowledge assistance and support on the murine studies from the University of Notre

Dame Freimann Life Science Center (FLSC). We give a special thanks to William J. Kaliney, M.D.,

pathologist from Harper Cancer Research Center, for assistance with histological analysis and

scoring. We acknowledge financial support from the Warren Center for Drug Discovery (pilot

grant) and NIH grant R21AI71742. We also thank members of the Santiago-Tirado lab for their

thoughtful comments and feedback on this work.

REFERENCES

1.   Denning DW. 2024. Global incidence and mortality of severe fungal disease. Lancet Infect Dis
doi:10.1016/S1473-3099(23)00692-8.
2.   Gaylord EA, Choy HL, Doering TL. 2020. Dangerous Liaisons: Interactions of Cryptococcus
neoformans with Host Phagocytes. Pathogens 9.
3.   Stuckey PV, Santiago-Tirado FH. 2023. Fungal mechanisms of intracellular survival: what can we
learn from bacterial pathogens? Infect Immun 91:e0043422.
4.   Rajasingham R, Govender NP, Jordan A, Loyse A, Shroufi A, Denning DW, Meya DB, Chiller TM,
Boulware DR. 2022. The global burden of HIV-associated cryptococcal infection in adults in
2020: a modelling analysis. Lancet Infect Dis doi:10.1016/S1473-3099(22)00499-6.
5.   Santiago-Tirado FH, Doering TL. 2017. False friends: Phagocytes as Trojan horses in microbial
brain infections. PLoS Pathog 13:e1006680.
6.   Denham ST, Brown JCS. 2018. Mechanisms of Pulmonary Escape and Dissemination by
Cryptococcus neoformans. J Fungi (Basel) 4.
7.   Zaragoza O, Rodrigues ML, De Jesus M, Frases S, Dadachova E, Casadevall A. 2009. The capsule
of the fungal pathogen Cryptococcus neoformans. Adv Appl Microbiol 68:133-216.
8.   Denham ST, Verma S, Reynolds RC, Worne CL, Daugherty JM, Lane TE, Brown JCS. 2018.
Regulated Release of Cryptococcal Polysaccharide Drives Virulence and Suppresses Immune Cell
Infiltration into the Central Nervous System. Infect Immun 86.
9.   Alspaugh JA. 2015. Virulence mechanisms and Cryptococcus neoformans pathogenesis. Fungal
Genet Biol 78:55-8.
10.   Pacelli C, Bryan RA, Onofri S, Selbmann L, Shuryak I, Dadachova E. 2017. Melanin is effective in
protecting fast and slow growing fungi from various types of ionizing radiation. Environ
Microbiol 19:1612-1624.
11.   Ueno K, Nagamori A, Honkyu NO, Kataoka M, Shimizu K, Chang YC, Kwon-Chung KJ, Miyazaki Y.
2023. Cryptococcus neoformans requires the TVF1 gene for thermotolerance and virulence.
Med Mycol 61.

12.   Winski CJ, Qian Y, Mobashery S, Santiago-Tirado FH. 2022. An Atypical ABC Transporter Is
Involved in Antifungal Resistance and Host Interactions in the Pathogenic Fungus Cryptococcus
neoformans. mBio 13:e0153922.
13.   Oliveira NK, Bhattacharya S, Gambhir R, Joshi M, Fries BC. 2022. Novel ABC Transporter
Associated with Fluconazole Resistance in Aging of Cryptococcus neoformans. J Fungi (Basel) 8.
14.   Moreno A, Banerjee A, Prasad R, Falson P. 2019. PDR-like ABC systems in pathogenic fungi. Res
Microbiol 170:417-425.
15.   Khunweeraphong N, Kuchler K. 2021. Multidrug Resistance in Mammals and Fungi-From MDR to
PDR: A Rocky Road from Atomic Structures to Transport Mechanisms. Int J Mol Sci 22.
16.   Jankovic D, Liu Z, Gause WC. 2001. Th1- and Th2-cell commitment during infectious disease:
asymmetry in divergent pathways. Trends Immunol 22:450-7.
17.   Stacey AR, Norris PJ, Qin L, Haygreen EA, Taylor E, Heitman J, Lebedeva M, DeCamp A, Li D,
Grove D, Self SG, Borrow P. 2009. Induction of a striking systemic cytokine cascade prior to peak
viremia in acute human immunodeficiency virus type 1 infection, in contrast to more modest
and delayed responses in acute hepatitis B and C virus infections. J Virol 83:3719-33.
18.   McMichael AJ, Borrow P, Tomaras GD, Goonetilleke N, Haynes BF. 2010. The immune response
during acute HIV-1 infection: clues for vaccine development. Nat Rev Immunol 10:11-23.
19.   Wozniak KL, Ravi S, Macias S, Young ML, Olszewski MA, Steele C, Wormley FL. 2009. Insights into
the mechanisms of protective immunity against Cryptococcus neoformans infection using a
mouse model of pulmonary cryptococcosis. PLoS One 4:e6854.
20.   Wozniak KL, Hardison S, Olszewski M, Wormley FL, Jr. 2012. Induction of protective immunity
against cryptococcosis. Mycopathologia 173:387-94.
21.   Chang CC, Harrison TS, Bicanic TA, Chayakulkeeree M, Sorrell TC, Warris A, Hagen F, Spec A,
Oladele R, Govender NP, Chen SC, Mody CH, Groll AH, Chen YC, Lionakis MS, Alanio A, Castaneda
E, Lizarazo J, Vidal JE, Takazono T, Hoenigl M, Alffenaar JW, Gangneux JP, Soman R, Zhu LP,
Bonifaz A, Jarvis JN, Day JN, Klimko N, Salmanton-Garcia J, Jouvion G, Meya DB, Lawrence D,
Rahn S, Bongomin F, McMullan BJ, Sprute R, Nyazika TK, Beardsley J, Carlesse F, Heath CH,
Ayanlowo OO, Mashedi OM, Queiroz-Telles Filho F, Hosseinipour MC, Patel AK, Temfack E, Singh
N, Cornely OA, Boulware DR, et al. 2024. Global guideline for the diagnosis and management of
cryptococcosis: an initiative of the ECMM and ISHAM in cooperation with the ASM. Lancet Infect
Dis doi:10.1016/S1473-3099(23)00731-4.
22.   Rothe C, Sloan DJ, Goodson P, Chikafa J, Mukaka M, Denis B, Harrison T, van Oosterhout JJ,
Heyderman RS, Lalloo DG, Allain T, Feasey NA. 2013. A prospective longitudinal study of the
clinical outcomes from cryptococcal meningitis following treatment induction with 800 mg oral
fluconazole in Blantyre, Malawi. PLoS One 8:e67311.
23.   Coelho C, Casadevall A. 2016. Cryptococcal therapies and drug targets: the old, the new and the
promising. Cell Microbiol 18:792-9.
24.   Zhang J, Li L, Lv Q, Yan L, Wang Y, Jiang Y. 2019. The Fungal CYP51s: Their Functions, Structures,
Related Drug Resistance, and Inhibitors. Front Microbiol 10:691.
25.   Pulendran B. 2004. Modulating TH1/TH2 responses with microbes, dendritic cells, and pathogen
recognition receptors. Immunol Res 29:187-96.
26.   Qiu Y, Dayrit JK, Davis MJ, Carolan JF, Osterholzer JJ, Curtis JL, Olszewski MA. 2013. Scavenger
receptor A modulates the immune response to pulmonary Cryptococcus neoformans infection. J
Immunol 191:238-48.
27.   Brinks J, van Dijk EHC, Habeeb M, Nikolaou A, Tsonaka R, Peters HAB, Sips HCM, van de Merbel
AF, de Jong EK, Notenboom RGE, Kielbasa SM, van der Maarel SM, Quax PHA, Meijer OC, Boon
CJF. 2018. The Effect of Corticosteroids on Human Choroidal Endothelial Cells: A Model to Study
Central Serous Chorioretinopathy. Invest Ophthalmol Vis Sci 59:5682-5692.

28. Day J, Imran D, Ganiem AR, Tjahjani N, Wahyuningsih R, Adawiyah R, Dance D, Mayxay M,
Newton P, Phetsouvanh R, Rattanavong S, Chan AK, Heyderman R, van Oosterhout JJ, Chierakul
W, Day N, Kamali A, Kibengo F, Ruzagira E, Gray A, Lalloo DG, Beardsley J, Binh TQ, Chau TT,
Chau NV, Cuc NT, Farrar J, Hien TT, Van Kinh N, Merson L, Phuong L, Tho LT, Thuy PT, Thwaites
G, Wertheim H, Wolbers M. 2014. CryptoDex: a randomised, double-blind, placebo-controlled
phase III trial of adjunctive dexamethasone in HIV-infected adults with cryptococcal meningitis:
study protocol for a randomised control trial. Trials 15:441.
29. Kitonsa J, Nsubuga R, Mayanja Y, Kiwanuka J, Nikweri Y, Onyango M, Anywaine Z, Ggayi AB,
Kibengo FM, Kaleebu P, Day J. 2020. Determinants of two-year mortality among HIV positive
patients with Cryptococcal meningitis initiating standard antifungal treatment with or without
adjunctive dexamethasone in Uganda. PLoS Negl Trop Dis 14:e0008823.
30. Netea MG, Schlitzer A, Placek K, Joosten LAB, Schultze JL. 2019. Innate and Adaptive Immune
Memory: an Evolutionary Continuum in the Host's Response to Pathogens. Cell Host Microbe
25:13-26.
31. Rudman J, Evans RJ, Johnston SA. 2019. Are macrophages the heroes or villains during
cryptococcosis? Fungal Genet Biol 132:103261.
32. Hole C, Wormley FL, Jr. 2016. Innate host defenses against Cryptococcus neoformans. J
Microbiol 54:202-11.
33. Hole CR, Lam WC, Upadhya R, Lodge JK. 2020. Cryptococcus neoformans Chitin Synthase 3 Plays
a Critical Role in Dampening Host Inflammatory Responses. mBio 11.
34. O'Halloran JA, Powderly WG, Spec A. 2017. Cryptococcosis today: It is not all about HIV
infection. Curr Clin Microbiol Rep 4:88-95.
35. Koselny K, Mutlu N, Minard AY, Kumar A, Krysan DJ, Wellington M. 2018. A Genome-Wide
Screen of Deletion Mutants in the Filamentous Saccharomyces cerevisiae Background Identifies
Ergosterol as a Direct Trigger of Macrophage Pyroptosis. mBio 9.
36. Rodrigues ML. 2018. The Multifunctional Fungal Ergosterol. mBio 9.
37. Pasrija R, Panwar SL, Prasad R. 2008. Multidrug transporters CaCdr1p and CaMdr1p of Candida
albicans display different lipid specificities: both ergosterol and sphingolipids are essential for
targeting of CaCdr1p to membrane rafts. Antimicrob Agents Chemother 52:694-704.
38. Xu X, Bittman R, Duportail G, Heissler D, Vilcheze C, London E. 2001. Effect of the structure of
natural sterols and sphingolipids on the formation of ordered sphingolipid/sterol domains
(rafts). Comparison of cholesterol to plant, fungal, and disease-associated sterols and
comparison of sphingomyelin, cerebrosides, and ceramide. J Biol Chem 276:33540-6.
39. Upadhya R, Lam WC, Hole CR, Vasselli JG, Lodge JK. 2023. Cell wall composition in Cryptococcus
neoformans is media dependent and alters host response, inducing protective immunity. Front
Fungal Biol 4.
40. Kozel TR, Levitz SM, Dromer F, Gates MA, Thorkildson P, Janbon G. 2003. Antigenic and
biological characteristics of mutant strains of Cryptococcus neoformans lacking capsular O
acetylation or xylosyl side chains. Infect Immun 71:2868-75.
41. Posteraro B, Sanguinetti M, Sanglard D, La Sorda M, Boccia S, Romano L, Morace G, Fadda G.
2003. Identification and characterization of a Cryptococcus neoformans ATP binding cassette
(ABC) transporter-encoding gene, CnAFR1, involved in the resistance to fluconazole. Mol
Microbiol 47:357-71.
42. Prasad R, De Wergifosse P, Goffeau A, Balzi E. 1995. Molecular cloning and characterization of a
novel gene of Candida albicans, CDR1, conferring multiple resistance to drugs and antifungals.
Curr Genet 27:320-9.
43. Lamping E, Baret PV, Holmes AR, Monk BC, Goffeau A, Cannon RD. 2010. Fungal PDR
transporters: Phylogeny, topology, motifs and function. Fungal Genet Biol 47:127-42.

44. Del Poeta M, Wormley FL, Jr., Lin X. 2023. Host populations, challenges, and commercialization
of cryptococcal vaccines. PLoS Pathog 19:e1011115.
45. Wang Y, Wang K, Rivera A, Xue C. 2023. Development of a Heat-Killed fbp1 Mutant Strain as a
Therapeutic Agent To Treat Invasive Cryptococcus Infection. Microbiol Spectr 11:e0495522.
46. Pham T, Li Y, Watford W, Lin X. 2023. Vaccination with a ZNF2(oe) Strain of Cryptococcus
Provides Long-Lasting Protection against Cryptococcosis and Is Effective in
Immunocompromised Hosts. Infect Immun 91:e0019823.
47. Wormley FL, Jr., Perfect JR, Steele C, Cox GM. 2007. Protection against cryptococcosis by using a
murine gamma interferon-producing Cryptococcus neoformans strain. Infect Immun 75:1453-
62.
48. Ristow LC, Jezewski AJ, Chadwick BJ, Stamnes MA, Lin X, Krysan DJ. 2023. Cryptococcus
neoformans adapts to the host environment through TOR-mediated remodeling of phospholipid
asymmetry. Nat Commun 14:6587.
49. Santiago-Tirado FH, Peng T, Yang M, Hang HC, Doering TL. 2015. A Single Protein S-acyl
Transferase Acts through Diverse Substrates to Determine Cryptococcal Morphology, Stress
Tolerance, and Pathogenic Outcome. PLoS Pathog 11:e1004908.
50. Santiago-Tirado FH, Hurtaux T, Geddes-McAlister J, Nguyen D, Helms V, Doering TL, Romisch K.
2023. The ER Protein Translocation Channel Subunit Sbh1 Controls Virulence of Cryptococcus
neoformans. mBio 14:e0338422.
51. Gish SR, Maier EJ, Haynes BC, Santiago-Tirado FH, Srikanta DL, Ma CZ, Li LX, Williams M, Crouch
EC, Khader SA, Brent MR, Doering TL. 2016. Computational Analysis Reveals a Key Regulator of
Cryptococcal Virulence and Determinant of Host Response. mBio 7:e00313-16.

Figures and figure legends

Fig. 1. *C. neoformans pdr6Δ*-**infected mice succumb to infection despite decreased fungal burden. (A)** Quantified fungal burden over time to monitor disease progression. Mice were inoculated with 5 x 10^4 cells and sacrificed every 7 days and organs were harvested to determine CFUs. Significance was determined using Mann-Whitney test, *P<0.05. **(B)** In vivo survival of H99 and *pdr6Δ* strains. 4 A/J mice per group were intranasally infected with 5 x 10^3 cryptococcal cells or mock infected with sterile PBS and monitored for 38 days (experiment was terminated when the last infected mouse reached endpoint). Significance was determined using Mantel-Cox test, **P<0.01. **(C)** Quantified organ burdens of H99 and *pdr6Δ*-infected mice at TOD. Organs were harvested, homogenized, and dilutions were plated on YPD. Plates were incubated for 48 h and CFUs were counted. Each data point represents a mouse and black bars represent the mean. Significance was determined using Mann-Whitney test, *P<0.05; **P<0.01.

Fig. 2. Infection with *C. neoformans pdr6Δ* results in altered histopathology. (A & B) Dissected

lungs of female A/J mice infected with 5 x 10⁴ cells of the WT strain or *pdr6*Δ mutant strain

were harvested at predetermined endpoints of 14 **(A)** and 35 **(B)** dpi. Gross organ examination

revealed significant changes in inflammation and tissue damage. H&E staining was used to

772 visualize microscopic lung pathology (inset [green boxes]). The 10X scale bar is 200 μm and the
773 40X scale bar is 50 μm.

DPI	Strain	% inflammation	Fungal infiltration	Quantitation of inflammatory immune cells/hpf 400X	Observations
7	H99	5 – 10%	20% alveoli contain encapsulated yeast	8 neutrophils 33 lymphocytes 40 histiocytes	Immune cells in peri-bronchial and perivascular space.
	pdr6Δ	5 – 10%	15 – 20% alveoli contain encapsulated yeast	9 neutrophils 26 lymphocytes 19 histiocytes	Immune cells in peri-bronchial and perivascular space. Some yeast has been phagocytized by histiocytes.
14	H99	10%	80 – 90% alveoli contain encapsulated yeast	52 neutrophils 26 lymphocytes 57 histiocytes	Immune cells in peri-bronchial and perivascular space.
	pdr6Δ	10%	50 – 60% alveoli contain encapsulated yeast	43 neutrophils 105 lymphocytes 71 histiocytes	Immune cells in peri-bronchial and perivascular space. Many histiocytes have phagocytized yeast. Areas of accumulated yeast/immune cells but no true parenchymal granulomas.
21*	pdr6Δ	10%	90% alveoli contain encapsulated yeast	50 neutrophils 49 lymphocytes 55 histiocytes	Immune cells in peri-bronchial and perivascular space.
28*	pdr6Δ	5 – 10%	80% alveoli contain encapsulated yeast	8 neutrophils 33 lymphocytes 55 histiocytes	Immune cells in peri-bronchial and perivascular space and single parenchymal granuloma (1.2mm)
35*	pdr6Δ	60 – 70%	100% present throughout	Too many to quantify	Consolidated parenchyma with many coalescent granulomas (histiocytes w/phagocytized yeast) and nests of neutrophils and lymphocytes with necrosis.

776 **Table 1. Quantitative analysis of H99 and *pdr6*Δ-infected lung sections over time.**
777 *All H99-infected mice had succumbed to infection by timepoint.
778 Histiocytes refer to leukocytes of the monocyte/macrophage/dendritic cell lineage.
779 hpf, high power field.

Fig. 3. FLC treatment significantly reduces CFUs in *pdr6*Δ-infected mice. (A) *In vivo* survival
curve of H99 and *pdr6*Δ-infected mice treated daily with either 8 or 16 mg/kg FLC. At least 4 A/J
mice per group were intranasally infected with 5 x 10^4 cells, or mock-infected with DPBS and
monitored for up to 50 dpi (mock-infected and FLC-treated mice not shown for clarity).
Significance was determined using Mantel-Cox test, *P<0.05; **P<0.01; ***P<0.001;
****P<0.0001. **(B-G)** Calculated organ burdens at 14 dpi **(B-D)** or endpoint **(E-G)**. Organs were
harvested, homogenized, and dilutions were plated on YPD. Plates were incubated for 48 hr
and CFUs were enumerated. Each data point represents a mouse and black bars represent the
mean. Significance was determined using one-way ANOVA with multiple comparisons (Welch's
corrected) *P<0.05; **P<0.01; ****P<0.0001.

Fig. 4. FLC treatment improves lung pathology in *pdr6Δ*-infected animals. Dissected lungs of
female A/J mice infected with 5 x 10^4 cells of the WT strain or *pdr6Δ* mutant strain and treated

daily with 16 mg/kg FLC were harvested at 14 and 35 dpi. Gross organ examination revealed
alterations in fungal containment and disease progression. H&E staining was used to visualize
microscopic lung pathology. Insets 1 & 2 highlight differential histopathology in the FLC-treated
pdr6Δ-infected mice at 35 dpi (inset [green boxes]). Inset 1 shows an intact lung parenchyma
with several fungi and immune cells while inset 2 is characterized as a granuloma. The 10X scale
bar is 200 μm and the 40X scale bar is 50 μm.

DPI	Strain	% inflammation	Fungal infiltration	Quantitation of inflammatory immune cells/hpf 400X	Observations
14	Mock +FLC	0%	None	-	-
	H99 +FLC	5 – 10%	~ 60% alveoli contain encapsulated yeast	50 – 60 neutrophils 40 – 50 lymphocytes 15 histiocytes	Moderate number of inflammatory cells predominately present in perivascular & peribronchial areas.
	pdr6Δ +FLC	< 5%	30 – 40% alveoli contain encapsulated yeast	1 – 5 neutrophils 0 – 15 lymphocytes 1 – 5 histiocytes	Mild infiltrate of inflammatory cells in focal peribronchial & perivascular areas.
35	Mock +FLC	0%	None	-	-
	pdr6Δ +FLC	5 – 10%	45 – 50% alveoli contain encapsulated yeast	5 – 10 neutrophils 10 – 20 lymphocytes < 5 histiocytes	Mild inflammatory cell infiltrate in many peribronchial and perivascular areas. Consolidation of adjacent alveoli in solid areas.

804 **Table 2. Quantitative analysis of Fluconazole treated H99 and *pdr6Δ*-infected lung sections**
805 **over time**
806 Histiocytes refer to leukocytes of the monocyte/macrophage/dendritic cell lineage.
807 hpf, high power field.

Fig. 5 Infection with the *pdr6Δ* mutant elicits a robust pro-inflammatory immune response. (A
& B) Quantification of pro-inflammatory and anti-inflammatory cytokines in the lungs **(A)** and
brains **(B)** of *C. neoformans*-infected mice. Female A/J mice were intranasally infected with 5 x
10^5 cells of the H99 or *pdr6Δ* strain and euthanized at endpoint. Homogenates from the lungs
and brain were used to perform Proteome Profiler Mouse XL Cytokine Arrays and Bio-Plex
Multi-Plex Arrays. Expression of cytokines/chemokines in *pdr6Δ*-infected mice was normalized
to H99 (dotted line).

Fig. 6 Dexamethasone treatment significantly extends survival in *pdr6*Δ-infected mice. (A) *In vivo* survival curve of H99 and *pdr6*Δ-infected mice treated daily with either 0.03 mg/kg Dexa or 0.03 mg/kg Dexa and 16 mg/kg FLC. At least 3 A/J mice per group were intranasally infected with 5 x 10^4 cells, or mock-infected with DPBS and monitored for up to 60 dpi (mock-infected and treated mice not shown for clarity). Significance was determined using Mantel-Cox test, **P<0.01; ****P<0.0001. **(B-G)** Calculated organ burdens at 14 dpi **(B-D)** or endpoint **(E-G)**. Organs were harvested, homogenized, and dilutions were plated on YPD. Plates were incubated for 48 hr and CFUs were enumerated. Each data point represents a mouse and black bars represent the mean. Significance was determined using one-way ANOVA with multiple comparisons (Welch's corrected) *P<0.05; **P<0.01; ***P<0.001.

Fig. 7. Effect of dexamethasone treatment on lung histopathology. Dissected lungs of female A/J mice infected with 5 x 10^4 cells of the WT strain or *pdr6Δ* mutant strain and treated daily with 0.3 mg/kg Dexa and harvested at 14 and 35 dpi. Gross organ examination revealed alterations inflammation. H&E staining was used to visualize microscopic lung pathology (inset [green boxes]). The 10X scale bar is 200 μm and the 40X scale bar is 50 μm.

DPI	Strain	% inflammation	Fungal infiltration	Quantitation of inflammatory immune cells/hpf 400X	Observations
14	Mock +Dexa	0%	None	-	-
	H99 +Dexa	10%	85 – 90% alveoli contain encapsulated yeast	10 – 20 neutrophils 10 – 30 lymphocytes < 5 histiocytes	Perivascular and peribronchial aggregates of admixed neutrophils and lymphocytes.
	pdr6Δ +Dexa	< 5%	20 – 30% alveoli contain encapsulated yeast	5 – 10 neutrophils 5 lymphocytes < 5 histiocytes	Focal inflammatory cell nests in perivascular and peribronchial space.
50	Mock +Dexa	0%	None	-	Inflammatory cells uncommon. Rare small nests of lymphocytes on or near pleural surface.
	pdr6Δ +Dexa	20%	> 80% alveoli contain encapsulated yeast	5 – 10 neutrophils 5 – 10 lymphocytes 20 – 30 histiocytes	Focal consolidated areas of dense inflammatory tissue up to 4 – 5 mm in dimension.

Table 3. Quantitative analysis of Dexamethasone treated H99 and *pdr6*Δ-infected lung sections over time

Histiocytes refer to leukocytes of the monocyte/macrophage/dendritic cell lineage.

hpf, high power field.

Fig. 8. Effect of dexamethasone/FLC combination treatment on lung histopathology.
Dissected lungs of female A/J mice infected with 5 x 10^4 cells of the WT strain or *pdr6Δ* mutant strain and treated daily with a combination of 0.3 mg/kg Dexa and 16 mg/kg FLC and harvested at 14 and 35 dpi. Gross organ examination revealed alterations inflammation. H&E staining was used to visualize microscopic lung pathology (inset [green boxes]). The 10X scale bar is 200 μm and the 40X scale bar is 50 μm.

DPI	Strain	% inflammation	Fungal infiltration	Quantitation of inflammatory immune cells/hpf 400X	Observations
14	Mock +Dexa +FLC	0%	None	-	-
	H99 +Dexa +FLC	15 – 20%	45 – 50% alveoli contain encapsulated yeast	15 – 30 neutrophils 15 – 30 lymphocytes < 5 histiocytes	Small to medium sized nests of inflammatory cells.
	pdr6Δ +Dexa +FLC	0%	10 – 15% alveoli contain encapsulated yeast	-	Inflammatory cells are uncommon with rare perivascular and peribronchial nests of small lymphocytes and few histiocytes.
50	Mock +Dexa +FLC	0%	None	Rare clusters of 2 – 4 histiocytes	Mildly dilated capillaries.
	pdr6Δ +Dexa +FLC	0%	75 – 80% alveoli contain encapsulated yeast	-	Few small foci of perivascular lymphocytes and rare histiocytes.

Table 4. Quantitative analysis of combination Dexamethasone + Fluconazole treated H99 and pdr6Δ-infected lung sections over time

Histiocytes refer to leukocytes of the monocyte/macrophage/dendritic cell lineage.

hpf, high power field.

Fig. 9 Infection with the *C. neoformans pdr6Δ* mutant generates an increase in innate immune cells, specifically macrophages. (A-H) Quantification of innate and adaptive immune cell populations at 14 dpi **(A-D)** and endpoint **(E-H)**. Female A/J mice were intranasally infected with 5 x 10⁴ cells of the H99 or *pdr6Δ* strain and euthanized at clinical endpoints corresponding to imminent mortality. Lungs were harvested and single-cell suspensions were generated and stained with immune panels to quantify monocytes, neutrophils, macrophages, CD4⁺ Th, CD4⁺ T-regs, and CD8⁺ cells. Samples were processed using flow cytometry and are presented as total # cells **(A, C, E, G)** or as % CD45⁺ cells **(B, D, F, H)**. Significance was determined using one-way ANOVA with multiple comparisons (Welch's corrected) *P<0.05; **P<0.01. Note: for **G** and **H**, the CD8⁺ T cell panel could not be run for the day 35 samples, hence they were removed from the H99 bars.

Fig. 10 *C. neoformans pdr6Δ* cells display increased PAMP exposure and altered capsule density. (A) Visualization of CFW, ConA, and β-(1,3)-glucan staining in H99, *pdr6Δ*, and *cap59Δ* cells. Representative images of stained cells. Scale bar, 10 μm. **(B)** Quantification of CFW, ConA, and EoY in H99, *pdr6Δ*, and *cap59Δ* strains. Cells were stained and measured using flow cytometry. Significance was determined using one-way ANOVA (Welch's corrected) *P<0.05; **P<0.01; ****P<0.0001. **(C)** Visualization and quantification of capsular antibody staining of H99, *pdr6Δ*, and *cap59Δ* strains. Cells were stained with anti-capsule mAb 302 and imaged using an inverted Zeiss microscope. Representative images of stained cells. Staining was quantified using a Cell Profiler pipeline. Significance was determined using one-way ANOVA (Welch's corrected) ***P<0.001.