# Rapid rebalancing of co-tuned ensemble activity in the auditory cortex

HiJee Kang[1], Travis A. Babola[1], & Patrick O. Kanold[1,2*]

**Affiliations:**

[1] Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD 20215

[2] Kavli Neuroscience Discovery Institute, Johns Hopkins University, Baltimore, MD 20215

*Corresponding author. Email: pkanold@jhu.edu

9
10 **Keywords:** cortex, optogenetics, holography, balance, plasticity
11
12

**Abstract:** Sensory information is represented by small varying neuronal ensembles in sensory cortices. In the auditory cortex (AC) repeated presentations of the same sound activate differing ensembles indicating high trial-by trial variability in activity even though the sounds activate the same percept. Efficient processing of complex acoustic signals requires that these sparsely distributed neuronal ensembles actively interact in order to provide a constant percept. Thus, the differing ensembles might interact to process the incoming sound inputs. Here, we probe interactions within and across ensembles by combining *in vivo* 2-photon $Ca^{2+}$ imaging and holographic optogenetic stimulation to study how increased activity of single cells level affects the cortical network. We stimulated a small number of neurons sharing the same frequency preference alongside the presentation of a target pure tone, further increasing their tone-evoked activity. We found that other non-stimulated co-tuned neurons decreased their tone-evoked activity when the frequency of the presented pure tone matched to their tuning property, while non co-tuned neurons were unaffected. Activity decrease was greater for non-stimulated co-tuned neurons with higher frequency selectivity. Co-tuned and non co-tuned neurons were spatially intermingled. Our results shows that co-tuned ensembles communicated and balanced their total activity across the larger network. The rebalanced network activity due to external stimulation remained constant. These effects suggest that co-tuned ensembles in AC interact and rapidly rebalance their activity to maintain encoding network dynamics, and that the rebalanced network is persistent.

## Introduction

Sensory perception requires fast encoding of relevant stimuli from a mixture of complex signals. Sensory cortices play a vital role in such sensory processing. In the auditory domain, for example, small neuronal ensembles in the auditory cortex (AC) are actively engaged to efficiently perceive relevant acoustic information [1-4]. The AC contains multiple ensembles of neurons that can be functionally identified, e.g., those formed by subsets of neurons preferring the same frequency also referred to as co-tuned neurons [5-7]. Repeated presentation of the same acoustic stimulus, e.g., a tone of the same frequency leads to a stable percept, but in the AC different ensembles of neurons are activated together at each repeat indicating a high trial-by-trial variability [5,8-10]. Activation of these different subsets of co-tuned neurons at each presentation of a stimulus reflects a sparse encoding of sound stimuli. Such sparse representation of co-activated neurons enables efficient coding with reduced metabolic energy to process complex information [11-16]. The sparse neuronal representation raises key questions of how activation of different ensembles leads to the same percept and how the overall activity within the cortical network is balanced across ensembles of co-tuned neurons. In particular, when a specific sound is present, a subset of co-tuned neurons will be activated, but not all co-tuned neurons [17]. Given that the percept of a repeating stimulus is constant, we speculated that neural activity is balanced across co-tuned as well as non co-tuned ensembles.

While neuronal ensembles constantly update their activities based on incoming information, how the activation of a particular sparse neuronal ensemble affects other neurons within the network to maintain the overall network balance for processing specific sensory information *in vivo* is largely unknown. *In vivo* optogenetic stimulation studies in the visual cortex (VC) suggested that inhibitory processes play a role in balancing network activity. In particular, *in vivo* single-cell holographic stimulation on a group of target cells, which induced increased response amplitude, resulted in changes in the response amplitudes of neighboring non-target neurons in the primary visual cortex (V1) [18], with similarly tuned neurons' activity being suppressed. Moreover, *in vivo* holographic optogenetic stimulation showed that visually-suppressed neurons had attenuated response amplitudes when holographic stimulation was given along with the visual stimulus presentation, which was not observed in visually activated neurons [19]. This suggested that neurons exhibit supralinear-to-linear input-output (IO) functions *in vivo*, rather than threshold-linear IO functions observed *in vitro*. These studies suggest that inhibitory influence from additional neuronal activation in the VC seems to play a major role during *in vivo* sensory processing, likely to maintain the activity balance of the network by modulating activities of neighboring neurons that share a similar tuning property.

One major difference between VC and AC is that the frequency tuning of neurons in the AC is less spatially localized, especially in a superficial layer (Layer 2/3) [20]. The local frequency preferences in the AC are diverse, thus neighboring neurons can show widely differing tuning properties [5]. To test how activity in specific AC cells among an intermingled and spatially distributed co-tuned and non co-tuned cell population is balanced during auditory processing, we stimulated a small group of AC cells using *in vivo* holographic optogenetic stimulation [21,22] while imaging AC population activity using 2-photon $Ca^{2+}$ imaging in awake mice. We further tested whether any activity changes induced by holographic stimulation persist, as recurrent cortical networks engage homeostatic plasticity to stabilize overall network activity levels [23,24]. Stimulating small ensembles of co-tuned neurons together with the presentation of a pure tone in their preferred frequency increased their tone-evoked activity. Furthermore, we observed that non-stimulated co-tuned neurons decreased their tone-evoked activity. Non co-tuned ensembles did not exhibit such changes in tone-evoked responses, regardless of the pure tone frequency. Thus, the increased activity in the stimulation-targeted ensemble had caused a decrease in activity in the non-stimulated co-tuned ensembles, specifically when the stimulation-paired pure tone was their preferred

frequency. Non-target co-tuned neurons exhibiting such effects were not necessarily
neighboring the targeted cells, suggesting specific interactions between co-tuned but not co-
located neurons. Lastly, the decreased activity in the non-stimulated co-tuned ensembles
persisted in the subsequent imaging session, even in the absence of holographic stimulation.
These results suggest that co-tuned ensembles form interacting overall networks that balance
their activity.
93
**Results**
**Optogenetic holographic stimulation increases activity in small ensembles *in vivo***
To sparsely manipulate neuronal ensembles, we used *in vivo* holographic stimulation.
To achieve reliable and selective *in vivo* holographic optogenetic stimulation of small
ensembles of neurons with single cell precision, we generated an AAV co-expressing the red-
shifted opsin rsChRmine and GCaMP8s (AAV9-hSyn-GCaMP8s-T2A-rsChRmine), as
rsChRmine minimizes the optical cross talk reducing a possible activation from the imaging
laser (940 nm excitation wavelength) [25]. Injecting AAV9-hSyn-GCaMP8s-T2A-rsChRmine
into AC yielded cells expressing both GCaMP and opsin (Fig. 1A). We first tested the
efficiency and reliability holographic stimulation, by targeting either a single cell or a small
ensemble of five cells. For single cell stimulation, we varied the stimulation point from the
target cell position by 10, 20, and 30 μm along either the x-axis or y-axis of the fields of view
(FOV; n = 15 cells, 3 animals). This results in a rapid decay of response amplitudes to
stimulation by the distance shift from the original cell position, confirming reliable
holographic stimulation at the single-cell level (~15 μm diameter) (mixed-effect model, $p <$
0.05; Fig. 1B). Furthermore, the stimulation effect of target cells was specific to the targeted
z-plane, showing no stimulation effect when the stimulation z-plane was off by 20 μm (Fig.
S2). For 5-cell stimulation, a majority of cells reliably responded to photo-stimulated *in vivo*
(5 mW/cell, 15 μm spiral, 30 revolutions, 6 s inter-stimulus interval (ISI), 5 trials) and
exhibited robust $Ca^{2+}$ responses (Fig. 1CD; permutation test, all $p < 0.05$), comparable with
responses to other opsins [26-29]. Thus, *in vivo* holographic stimulation enables precise targeting
and activation of groups of single neurons in AC.
116
**Optogenetic holographic stimulation increases sound evoked activity in A1 ensembles**
Since repeated sound stimulation activates different ensembles while resulting in the
same percept, we reasoned that ensembles interacted and speculated that increased activity in
one ensemble would prevent or reduce activity in co-tuned ensembles. We thus next sought
to investigate how increased neural activity in small co-tuned ensembles during sound
presentation affected sound-evoked responses in targeted and non-target co-tuned and non
co-tuned ensembles. To achieve this, we first needed to identify the tuning properties of
single neurons and then target a subset of co-tuned neurons for stimulation. To study how the
increased activity of a small number of neurons influences the activity of other neurons
according to their frequency tuning properties, we designed an experimental paradigm
comprising four sequential imaging sessions (Fig. 1E):
First, in the cell selection session (Fig. 1E), we identified tuned ensembles in primary
auditory cortex (A1) layer 2/3 (L2/3) by assessing frequency tuning properties of neurons
within the FOVs covering 550 μm$^2$ (total cells = 7344, sound responsive cells = 1331, FOVs
= 23; Fig. 1D). We presented pure tones of three different frequencies spanning the hearing
range of the mouse (4 kHz, 16 kHz, and 54 kHz, 100 ms duration, 2 sec. ISIs, 10 repeats for
each frequency). We chose 16 kHz and 54 kHz as the representative target ensemble tone
frequencies, as 16 kHz is within the most sensitive frequency range of mice [30] and 54 kHz is
within the range of mouse ultrasonic vocalization [31]. By selecting target ensembles in two
different frequencies, we ensured that effects of stimulation were not specific to one
particular population. For each condition (16 kHz or 54 kHz target ensemble for stimulation),

we selected 5 target cells to stimulate. To ensure that all cells in the ensemble were selective
for the target tone, we chose the most responsive cells in each condition. Thus, for the 16 kHz
target ensemble condition, we selected five cells (*target cells*) among the top 30% most
responsive cells to the 16 kHz tone. Similarly, for the 54 kHz target ensemble condition, we
selected five of the top 30% most 54 kHz tone responsive cells. By selecting target cells
sharing the same frequency preference, we aimed at investigating how activity changes from
co-tuned neuronal ensembles alter the processing of the target frequency in other co-tuned
and non co-tuned cells.
Second, in the baseline session (Fig. 1E), we determined the sound-evoked responses
of all imaged cells by presenting a series of 16 kHz and 54 kHz pure tones in a random order
(100 ms duration, 5.8-6.5 sec. ISI; baseline session, 30 repeats for each frequency). Exemplar
responses of cells from a 16 kHz and a 54 kHz ensemble are shown as black traces in Figure
2A and 2C.
Third, in the stimulation session (Fig. 1E), we examined how all sound-responsive
cells change their responses when a small group of cells in the network increases their
activity. We presented the same tones (16 kHz and 54 kHz in a random order), in tandem
with the optogenetic stimulation of five target cells (stimulation session, 100 ms stimulation
duration). We performed different sessions for the 16 kHz and 54 kHz target ensembles,
varying FOVs for each session (18 FOVs for 16 kHz target ensemble condition and 12 FOVs
for 54 kHz target ensemble condition). Figure 2A and C show two example FOV with
targeted neurons for a 16 kHz and 54 kHz ensemble.
Since both imaging and optogenetic stimulation involve optomechanical components,
we wanted to ensure that effects were not due to artifacts caused by our stimulation or
imaging setup. Moreover, cells can adapt their responses to repeated sound presentation.
Thus, to confirm any response changes observed from the stimulation session is due to the
optogenetic stimulation rather than simple response change due to acoustic sound
presentation, we added an additional control condition. For this control condition, we
performed the "stimulation" session with five target cells but with 0 mW laser power (i.e., no
stimulation) to verify that any response changes occurring in the stimulation session
compared to the baseline session were not simply due to the eventual response adaptation of
neurons to the tuned frequency (control condition; 13 FOVs). By selecting cells and
presenting 0 mW laser power, instead of no target cell selection or selecting any other no-cell
area within the FOV, we ensured that the laser power given to selected cells was the only
difference between the actual stimulation and control conditions, keeping any noise caused
by the imaging and stimulation setup the same.
Fourth, after the stimulation session (Fig. 1E), we performed an additional imaging
session (post-stimulation session), presenting another series of 16 kHz and 54 kHz pure tones
in a random order to examine whether changes in the sound-evoked responses persisted or
reverted back after the stimulation session.
177
**Optogenetic holographic stimulation increases activity in targeted ensembles**
We first identified the effect of the optogenetic stimulation on the targeted ensembles.
Figures 2A and 2D show fluorescence traces of exemplar cells from 16 kHz and 54 kHz
target ensembles. Optogenetic stimulation increased the sound-evoked fluorescence
amplitude in these individual cells. To quantify the effect of the optogenetic stimulation, we
compared the tone-evoked fluorescence responses of the targeted cells with and without
stimulation (Stimulation effect $= \Delta F/F_{(stimulation\ session)} - \Delta F/F_{(baseline\ session)}$). Around
72% of target cells (66 out of 90 cells over 18 FOVs for 16 kHz target stimulation and 42 out
of 60 cells 12 FOVs for 54 kHz target stimulation) showed increased response amplitude
during the stimulation session compared to the baseline session, regardless of the tone
presented (Fig. 2BD; permutation tests, all $p < 0.001$). These results indicate that holographic

stimulation was able to reliably increase activity in small populations of neurons. Moreover,
given that the target cells we selected were most responsive to their preferred tone frequency,
this increase in fluorescence indicates that the cells' responses to their preferred tone were not
saturated.
193
**Optogenetic holographic stimulation decreases activity in non-target co-tuned**
**ensembles**
We next investigated whether the optogenetically enhanced sound-evoked activity of
a small group of cells would cause activity changes in other non-stimulated cells. During
holographic optogenetic stimulation of the targeted cells, the non-target, but sound-
responsive cells (n = 995 cells for 16 kHz target ensemble condition and n = 675 cells for 54
kHz target ensemble condition), also changed their activity, showing either increased or
decreased response amplitudes (Fig. 2AC).
If cortical networks rebalance their activity, we speculated that the increased tone-
evoked activity in the targeted ensemble would lead to a decrease in tone-evoked activity in
coupled ensembles. Such rebalancing would keep the activity within the cortical network
stable. Moreover, given that we increased the activity to the preferred sound frequency, if this
rebalancing happens, it should occur only for the distinct sound frequency related to the cell's
tuning property. For example, stimulation of a 16 kHz ensemble should cause a greater
reduction in the 16 kHz tone response of non-targeted 16 kHz cells compared to their
response to the 54 kHz tone.
To address these questions, we investigated whether increased activity in the targeted
cells influenced the activity of non-target cells and how these changes were related to the
tuning properties of the cells. We first defined each sound-responsive cell's frequency
selectivity by computing a difference between response amplitude to 16 kHz and 54 kHz
from the baseline session (frequency preference = $(\Delta F/F_{(16kHz)}) - (\Delta F/F_{(54kHz)})$). We then
divided these cells into either 16 kHz preferring or 54 kHz preferring groups, taking 0 (i.e.,
no preference) as a criterion (Fig. S2). Both subgroups exhibited stronger tone-evoked
responses to their preferred frequency, independent of the condition ($t(5700) = 4.79$, $p <$
0.0001; Fig. S2). This confirms that the criterion for cell group threshold is valid.
We then focused on our main question by comparing the stimulation effect of the two
target ensemble groups to the control condition to identify whether stimulation decreased the
response of non-target co-tuned neurons. Neural activity in AC rapidly shows stimulus-
specific adaptation to the repeated presentation of the stimulus[32-34], which can obscure
stimulation related changes. We thus used the response amplitude change between the
baseline and the "stimulation" control session as a representative threshold to test the effect
of the stimulation. We once again used the difference in response amplitude between the
baseline and stimulation sessions as the measure of the stimulation effect
$(\Delta F/F_{(stimulation\ session)} - \Delta F/F_{(baseline\ session)})$. Neighboring cells within 20 μm from the
target stimulation point were removed from the analysis since they could have been directly
affected by the stimulation.
We compared the stimulation effect between non-target co-tuned and non co-tuned
cells across conditions (16 kHz and 54 kHz target ensembles as well as control conditions)
for different pure tone presentations. Since our primary interest was how non-target cells
respond to increased activity in target ensembles, we focused on conditions where the pure
tone frequency matched or did not match the tuning properties of the non-target cells. Since
we stimulated during tone presentation the effects of the holographic stimulation and
stimulus-specific adaptation co-occurred. To isolate these components, we used a linear
mixed-effect model with cell group, condition, and pure tone frequency as fixed factors, and
FOVs as a random factor. We then performed ANOVA on the model to assess the main
effects and interactions.

A marginal significant main effect of the condition ($F(2,37.1) = 2.983$, $p = 0.0628$) on
the response change in the stimulation session relative to the baseline session (i.e.,
stimulation effect) was observed, indicating that these changes may depend on the
stimulation condition. We further analyzed the data to better understand how the different
factors interacted in the response amplitude changes. A significant interaction between the
pure tone frequency and cell group ($F(1,4397.6) = 186.967$, $p < 0.0001$) suggests that each
cell group responded differently to the two pure tone frequencies. Specifically, the response
amplitude decreases in the stimulation session relative to the baseline session were more
pronounced for each cell group when the played pure tone matched to their tuning property.
This interaction between pure tone frequency and cell group highlights the importance of
frequency tuning in modulating response amplitudes. Such response amplitude decreases of
non-target cells to their preferred pure tone presentation further aligns with the stimulus
specific adaptation due to repeatedly presented pure tones [32]. Additionally, a significant three-
way interaction across condition, cell groups, and pure tone frequency ($F(2,4397.6) = 3.517$,
$p = 0.0298$) suggests the combined effects of the stimulation condition and the cell group on
response amplitude depend on the pure tone frequency. The stimulation effect is not uniform
across cell groups and depends heavily on the frequency, highlighting a complex interplay
between the tuning property of cells, stimulation condition, and presented pure tone
frequency.
Consequently, we analyzed post-hoc comparisons estimated marginal means with
contrasts, as our focus was how co-tuned cells change their responses due to the increased
activity in the target cells along with the frequency of the presented pure tone.
For 16 kHz preferring cell group (n = 537), we observed a greater stimulation effect
(i.e., decrease in response amplitude) for 16 kHz tone presentation when 16 kHz target
ensemble was stimulated compared to the control condition ($t(124) = 3.114$, $p = 0.0064$). For
all other pairs, no significant stimulation effect was observed. This suggests that non-target
16 kHz co-tuned cells reduce their response amplitudes when target ensembles share the
same tuning property. Furthermore, such response change occurs only when they process
their preferred frequency (Fig. 3B, left).
We repeated the experiments and the analysis with 54kHz cells as the target group. In
general, we observed similar results. The stimulation effect was significantly more
pronounced for 54 kHz tone presentation when 54 kHz target ensemble (n = 359) was
stimulated compared to the control condition ($t(168) = 3.074$, $p = 0.0069$; all $p$-values were
adjusted for multiple comparisons using the Tukey method). All other pairs did not show any
stimulation effect (Fig. 3B, right).
To further explore whether the stimulation effect could be explained by activity
rebalancing within the co-tuned network, we implemented a simple model in which a
suppression term was applied either to all neurons or specifically to non-target co-tuned cells.
By comparing two different model outcomes and the real data, we observed a significant
effect of the model type ($F(2, 2535) = 34.943$, $p < 0.0001$). Moreover, an interaction between
the model type and cell groups was observed ($F(2, 2535) = 36.348$, $p < 0.0001$). Applying
suppression to only non-target co-tuned cells during the stimulation session yielded a
significant response amplitude decrease for co-tuned cells compared to non co-tuned cells
($F(1, 2535) = 45.62$, $p < 0.0001$), which resembles the real data. In contrast, applying
suppression to all non-target cells led to similar amplitude changes in both co-tuned and non
co-tuned neurons ($F(1, 2535) = 0.87$, $p = 0.35$), which was not observed in either the real data
or the simulated data restricted to co-tuned cell suppression. These results suggest that the
target cell stimulation induces a selective activity suppression within the co-tuned network
for processing their preferred frequency.
Together, these results indicate that the effect of holographic optogenetic stimulation
depends not on the specific tuning of cells, but on the co-tuning between stimulated and non-

stimulated neurons. Also, this effect is not driven solely by a few non-target cells with large
response changes. Rather, the overall population of cells shows relative response changes due
to the stimulation when synchronized with their preferred frequency.
Overall, these results further suggest that when neural activity is increased in a subset
of target cells due to photostimulation in addition to the target sound presentation, other co-
tuned cells selectively reduce their tone-evoked responses to their preferred tone presentation,
indicating that the network rebalances to maintain network activity within a certain range.
298
**Rebalanced network responses are stable**
We then questioned whether such response amplitude changes due to stimulation
within the local network are persistent. To test whether the rebalanced status of the neuronal
ensemble is persistent, we examined the tone-evoked response amplitude changes between
the    post-stimulation    and    the    stimulation    sessions    (post-stimulation    effect:
$\Delta F/F_{(post-stimulation\,session)} - \Delta F/F_{(stimulation\,session)}$). Response amplitudes were similar
across conditions and tone presentation frequencies for both groups of cells ($F(2, 4056) =$
$1.83, p = 0.16$; Fig. 3B). These results indicate that pairing exogenous stimulation on a subset
of neurons along with sounds can instantaneously change the network responses to sounds,
and this change can persist at least for many minutes during the experimental session.
309
**Neurons with higher frequency selectivity show greater response changes**
Our results demonstrate that response changes on non-target cells are significantly
influenced by the frequency tuning of stimulation-target cells as well as the frequency of the
presented pure tone along with the stimulation. However, we also observed a marginal
stimulation effect in the 54 kHz non-target cell group during 54 kHz pure tone presentation,
even when 16 kHz target cells were stimulated. We reasoned that this effect might be due to
some weak sound activation of 54 kHz cells by 16 kHz tones potentially due to the
asymmetric shapes of many auditory tuning curves in AC [35,36]. Indeed, many cells exhibited
broad tuning properties, responding to both 16 kHz and 54 kHz (Fig. 3C). Thus, this marginal
stimulation effect could be attributed to cells grouped as 54 kHz preferring cells, yet still
showing sound evoked responses to 16 kHz, particularly given that 16 kHz is within the
sensitive frequency range in mice [30].
Building on our findings of a rebalanced cortical network, we next aimed to identify
whether frequency tuning selectivity influences response amplitude changes in the non-
targeted co-tuned neurons. For each cell, we calculated the frequency preference index
$(\Delta F/F_{(16kHz)}) - (\Delta F/F_{(54kHz)})$  and divided the cells into three categories of frequency
selectivity: low, mid, and high. We removed cells with extreme frequency preference values,
where the index values exceed $\pm$ 4 standard deviations from the median, prior to dividing
them into three categories. This removed about 1% of cells from the dataset for further
analyses. This grouping was based on the 33% quartile ranges, with each category
representing one-third of the data distribution (Fig. 3C). Values closer to 0 indicate more
broadly tuned cells across frequencies while extreme positive and negative values to indicate
sharply tuned cells to either frequency.
We then tested whether cells with higher frequency selectivity to one frequency
exhibited greater response amplitude changes. We performed a three-way ANOVA to
examine the effect of frequency selectivity (low, mid, high selectivity), stimulation condition
(control, 16 kHz target stim, 54 kHz target stim), and cell groups (16 kHz vs. 54 kHz
preferring cells) on the response amplitude change. There were significant main effects of
frequency selectivity ($F(2, 2183) = 23.52, p < 0.0001$) and stimulation condition ($F(2, 2183)$
$= 11.03, p < 0.0001$). No significant main effect of cell group was observed ($F(1, 2183) =$
$0.77, p = 0.379$). Thus, neither the interaction between frequency selectivity and cell group
($F(2, 2183) = 0.69, p = 0.503$), nor the interaction between condition and cell group ($F(2,$

2183) = 2.64, $p$ = 0.072) was significant. The three-way interaction between frequency
selectivity, stimulation condition, and cell group was also not significant ($F(4, 2183) = 0.86$,
$p = 0.487$).

However, the interaction between frequency selectivity and stimulation condition was
significant ($F(4, 2183) = 2.82$, $p = 0.0238$), indicating that the effect of frequency selectivity
depended on the condition. These results suggests that the response amplitude changes across
conditions were more prominent for cells with higher frequency selectivity.

To identify where the significant response difference occurred across conditions, we
further performed post-hoc pairwise comparisons between conditions within each frequency
selectivity category for each cell groups. For 16 kHz preferring cells, we observed a
significant difference in the response change between control and 16 kHz stim conditions
only from the high frequency selectivity category ($p < 0.027$, Holm-Bonferroni correction for
multiple comparisons). In parallel, for 54 kHz preferring cells, the significant effect was
observed between control and 54 kHz stim conditions ($p = 0.033$, Holm-Bonferroni
correction for multiple comparisons; Fig. 3D). These results indicate that non-targeted cells
with higher frequency selectivity exhibit the greatest response amplitude changes, only when
the target stimulated cells were co-tuned with them. These results also suggest that frequency
selective neurons form co-tuned networks.

**Sparsely distributed non-target co-tuned ensembles immediately rebalance their**
**activities to maintain the network balance**
Network balance can be achieved by multiple mechanisms operating on different
timescales. To get insight into the potential mechanisms underlying the observed rebalancing,
we next investigated how rapidly cells start adjusting their responses during the stimulation
condition. We thus examined the stimulation effect (changes in response amplitude due to
stimulation of target cells) for non-target co-tuned ensembles at the single-trial level. We
observed decreased response amplitudes from the first trial, with no significant decay across
trials (Fig. 4A), regardless of cell groups, frequency presentation, and conditions (sum-of-
squares $F$-test, all $p > 0.05$). The absence of trial-related response amplitude changes in non-
target co-tuned ensembles indicates that non-target co-tuned cells immediately change their
activity whenever targeted cells increased their activity due to stimulation, to maintain the
network balance.

**Non-target co-tuned ensembles that show rebalancing are spatially distributed**
Activity rebalancing could be driven by local, e.g. changes in excitatory-inhibitory
(E/I) balance [37], or distributed changes. To identify whether co-tuned ensembles that changed
their activities are locally or widely distributed, we computed the center of mass distance
between each non-target cell to any of the target cells. For the stimulation condition, we
excluded non-target cells that were within 20 μm distance of the target cells to ensure that
any effects from those neighboring cells with their increased response amplitudes could have
been not, even partially, due to photostimulation (Fig. 1B-D). We observed that non-target
co-tuned ensembles were widely distributed within the FOV, similar to non-target non co-
tuned ensembles as well as those from the control condition (sum-of-squares $F$-test, $p > 0.05$;
Fig. 4B). This indicates that activity changes of non-target co-tuned ensembles are not merely
the result of direct input from external photostimulation within a tight localized network.
Rather, widespread, sparsely represented co-tuned ensembles continuously update incoming
information based on their tuning properties.

**Discussion**
Trial-by-trial variability in neuronal activity is ubiquitous in the brain, with sensory
stimuli evoking activity in different sparse co-tuned ensembles at different times. How

sensory-evoked activity is distributed and coordinated across sparsely distributed co-tuned
networks has been unknown. Here, we leveraged the capability for selective *in vivo*
stimulation via holographic optogenetics to investigate how functionally related neuronal
ensembles in AC coordinate activity. Our results show that manipulating a small subset of
target co-tuned neurons alters the auditory-evoked responses of other non-target co-tuned
neurons. Increased activity in one subset of neurons is balanced by decreased activity in the
rest of the co-tuned population of neurons.
Importantly, such network rebalancing occurs only for processing acoustic features
specific to their tuning. Our analysis shows that the most selective non-targeted neurons show
the strongest effect after stimulation, suggesting that selective neurons form functionally
interacting sub-networks consistent with in vivo correlation analyses [38] and in vitro studies in
visual cortex [39]. Functionally related cells might form these subnetworks during development
likely due to lineage relationships and Hebbian processes [40-42]. Together, our findings suggest
that neuronal ensembles with strengthened connectivity across neuronal ensembles sharing
similar functional tuning properties actively interact and update their status to maintain the
overall network, enabling energy-efficient sensory processing.
The present work applied holographic optogenetic stimulation to manipulate neuronal
activity at a single-cell resolution in the AC for the first time. Similar to previous findings in
VC [18,19], our study further supports the idea that extra activation within the network exerts an
inhibitory influence on a subset of neurons. In the VC, single cell stimulation resulted in
suppression of neighboring co-tuned neurons [18]. Our results here show that feature-specific
suppression occurs in spatially dispersed ensembles of non-target co-tuned neurons. The
widely distributed response amplitude decreases in those neurons suggest that this
phenomenon is not limited to local neighboring cells but involved widespread networks [8].
Thus, the effect is not solely due to inhibition caused by neighboring interneurons from the
optogenetic stimulation. Instead, neurons with similar functional characteristics, sparsely
distributed throughout the AC, actively interact, with more sharply tuned neurons to modulate
their activity the most.
Cortical networks are shaped by dynamic changes in neural activity driven by various
factors. Neurons rapidly modulate their responses based on their functional roles in sensory
processing. Recurrent cortical networks are thought to update their activity based on
incoming information to maintain homeostatic balance [23,24]. Concurrently, co-activated
neurons processing similar acoustic properties strengthen their connectivity by Hebbian
learning [43-45]. Thus, cortical networks are shaped by an active interplay of synaptic plasticity,
homeostasis and Hebbian learning, rather than by a single dominant mechanism [46-49].
Rebalancing of network activity is often attributed to homeostatic rebalancing of individual
cell's activity [24,50] or an E/I balance: increased activity of inhibitory neurons resulting in
reduced activity of excitatory neurons [37]. Our results suggest that rebalancing is tuning-
specific.
Given that inhibitory neurons in A1 are generally less frequency selective than
excitatory neurons [51], changes in inhibition are unlikely to be the only contributor to the
observed effect. While our AAV was not cell-type specific, it is also unlikely that many
selected target neurons would be inhibitory interneurons as a greater proportion of neurons in
sensory cortices are excitatory (about 80% compared to 20% inhibitory [52-55]; AAV-hSyn is
expressed in both excitatory and inhibitory neurons in similar proportions [56]). Furthermore,
our primary cell selection criterion for stimulation yielded a subgroup of a strong specific
frequency-responsive cells (top 30% of cells that show the biggest evoked responses to the
target tone after excluding cells that show responses to multiple tones). This criterion likely
selected more excitatory neurons, as they generally show greater stimulus-selective responses
than inhibitory neurons in sensory cortices [51,57].

It is noteworthy that not all target neurons showed clear activation in response to
holographic stimulation. Although we attempted to pre-select cells responsive to stimulation,
some of them seemed to exhibit reduced activation during the experiment, potentially due to
motion artifacts, response adaptation, network suppression, or trial-by-trial variability.
Nevertheless, the frequency-specific suppression on co-tuned neurons observed in this study
suggests that this effect can be driven by activation of a very small number of neurons. One
caveat is that, while we presume most target cells were likely excitatory, inhibitory neurons
may have been included in the target cell group. Inhibitory neurons could show a reliable,
strong responses to the optogenetics compared to more variable responses that excitatory
neurons could show [58], thus may have been included in the target cell group. However, if
more inhibitory cells were included, this would have reduced trial-by-trial variability from
the stimulation yielding higher probability of target cell activation, which is different from
what we observed. Additionally, re-occurring sounds evoke activity changes in both
excitatory and inhibitory neurons in the same direction, leaving the E/I balance unchanged
[59,60]. Moreover, the fraction of inhibitory cells is 10 ~ 20% of all cortical neurons [61-63], thus
the chance of stimulating them is small. Together, these considerations suggest that our
results are unlikely the effect of the activation of inhibitory neurons. Rather, other balancing
mechanisms such as short-term depression at thalamocortical synapses may play a role [64].
In contrast to slow changes of homeostatic plasticity related to the E/I balance,
plasticity in cortical cellular responses can occur quickly and are cell-specific, thereby tuned
to their functional response properties [65-67]. Based on this, we speculate that the decrease in
response amplitudes of non-target neurons likely reflects rapid activity-dependent synaptic
changes in excitatory cells. Future work using Cre-dependent virus expression or a cell type
specific labeling approach will be required to confirm cell-type-specific roles in this
phenomenon and underlying mechanisms.
Lastly, no additional response changes were observed in the post-stimulation session
indicating that the rebalanced network status remained constant. Indeed, constant amplitudes
were observed regardless of conditions, subgroups, and frequencies, suggesting that the
network persistence was achieved through repeated acoustic stimuli presentation and
photostimulation. The persistence after the stimulation condition further suggests that once
the newly learned rebalanced network status is achieved, the network response stabilizes and
remains persistent. The mechanisms behind this stabilization are unclear but may involve an
active interplay of homeostasis and Hebbian learning to form co-active networks [68].
Taken together, the present study reveals how neuronal ensembles in the AC
rebalance to maintain a homeostatic processing equilibrium for a given sensory input, and
that rebalanced networks remain persistent. Moreover, our results show that network activity
can be controlled by even a small subset of neurons, and the network changes are closely tied
to the functional tuning properties of neurons.
481
**Materials and Methods**
Methods
*Animals*
A total of 14 mice over 8 weeks old (8 – 30 weeks, 6 males and 8 females) were used in the
experiments. To retain high-frequency hearing at experimental ages, offspring from
C57BL6/J and B6.CAST-*Cdh23*$^{Ahl+}$/Kjn (Jax 002756) were used for all experiments.
Animals were housed on 12-hr reversed light/dark cycle. All experimental procedures were
approved by Johns Hopkins Institutional Animal Care and Use Committee.
490
*Preparation of AAV9-hSyn-GCaMP8s-T2A-rsChRmine*
To generate the AAV9-hSyn-GCaMP8s-T2A-rsChRmine virus, a gene fragment containing
T2A, rsChRmine, Kv2.1 soma-targeted localization motif, and 3xHA tag (synthesized by

Twist Biosciences) was subcloned into the pGP-AAV-syn-jGCaMP8s plasmid vector
(Addgene: 162374). Viral vectors were commercially prepared (Virovek) to a concentration
of 1 x 10$^{13}$ vg/mL.
497
*Surgery and virus injection*
Surgery was performed as described in previous studies [9,10]. We injected dexamethasone
(1mg/kg, VetOne) subcutaneously (s.c.) to minimize brain swelling prior to surgery. 4%
isoflurane (Fluriso, VetOne) with a calibrated vaporizer (Matrix VIP 3000) was used to
induce anesthesia, which was then reduced down to 1.5 – 2% for maintenance. Body
temperature was monitored and maintained at around 36□ throughout the surgery (Harvard
Apparatus Homeothermic monitor). We first removed the hair on the head using a hair
removal product (Nair) to expose the skin. Betadine and 70% ethanol were applied three
times to the exposed skin. Skin and tissues were then removed, and muscles were scraped to
expose the left temporal side where the craniotomy was conducted. Unilateral craniotomy
was performed to expose about 3.5 mm diameter region over the left AC. Virus (AAV9-
hSyn-GCaMP8s-TSA-rsChRmine titer of 1:2) injection was performed at 2-3 sites near
tentative A1 area at about 300 μm depth from the surface, using a glass pipette controlled by
a micromanipulator (Sutter Instrument MPC-200 and ROE-200 controller). We injected 300
nL on each site at the rate of 180 nL/min (Nanoject3). Once virus injection was completed,
two circular glass coverslips (one of 3 mm and one of 4 mm in diameter) were affixed with a
clear silicone elastomer (Kwik-Sil, World Precision Instruments). An extra layer of dental
acrylic (C&B Metabond) was applied around the edge of the cranial window to further secure
it, cover the exposed skull, and adhere a custom 3D-printed stainless steel headpost.
Carprofen (5 mg/kg) and cefazolin (300mg/kg) were injected (s.c.) post-operatively. Animals
were given at least 3-4 weeks of recovery and viral expression time before any imaging was
performed.
520
*Experimental procedures*
Animals were head-fixed on a custom-made stage, where a free field speaker (TDT
ED1) was faced towards the right ear at 45 degrees. All sound stimuli were driven by TDT
RX6 multiprocessor and we imaged GCaMP8s responses with a resonant scanning two-
photon microscope (Bruker Ultima 2Pplus; 940 nm excitation wavelength) at A1 L2/3 (160 –
200 μm). A1 was identified by its tonotopy gradient using widefield imaging (Fig. 1E) [69].
During 2-photon imaging sessions, we first conducted a short imaging session (about 1 min.)
presenting 100-ms pure tones at 3 different frequencies (4, 16, 54 kHz) at 70 dB SPL,
covering the mouse hearing range, for 10 times each in a randomized order (ISI: 2 sec). This
was to identify initial tuning properties of sound-responsive cells (cell selection session; Fig.
1E). The acquired imaging data was immediately processed using 'suite2p' and a custom-
written Matlab script to identify tone-responsive cells for each frequency. We continued only
when at least 50% of cells within the FOV showed sound-evoked responses. We took 16 kHz
or 54 kHz as our target functional properties. Target frequency was randomly assigned. We
manually tested the response changes to stimulation of the top 30% of target frequency
responsive cells (~ 20 cells) selected from the cell selection session by using a stimulation
laser (Light Conversion Carbide; 1040 nm excitation wavelength). Stimulation laser power
was set around 5 mW per cell. We selected 5 representative cells (target cells) that showed
visible fluorescence changes to the stimulation.
Prior to the main experimental session, we ran a short (~ 1 min.) stimulation session
without any sound presentation that comprised 5 trials of 100-ms stimulation with 6-second
ISIs for a rapid check of the stimulation effect. This session was restricted to 5 trials of
stimulation given the limited time of the imaging session, leading to larger variability of the

observed stimulation effect. We further verified the stimulation effect from the experimental
stimulation session where 30 trials were given.
For the main experiment, three consecutive imaging sessions were followed by the
presentation of either 16 kHz or 54 kHz 100 ms pure tones with random ISIs between 5.8 –
6.5 sec. for 30 trials each, as baseline session, stimulation session, and post-stimulation
session (Fig. 1E). Only the second imaging session (i.e., stimulation session) received
holographic stimulation on the pre-selected 5 target cells for 30 revolutions of 15 um spiral
for about 100 ms at 5 mW laser power per cell (16 kHz target cell or 54 kHz target cell
stimulation conditions) or 0 mW laser power per cell (control condition). The default
mechanical setup (Bruker PrairieView version 5.7) of the microscope opens and closes the
uncaging shutter to enable the stimulation laser path for every single stimulation time point,
which causes an external mechanic sound that can trigger neural activation of the AC. To
minimize any effect of external mechanic sounds to cells in the AC, we kept the uncaging
shutter open during the imaging session. Number of imaging sessions per animal varied
depending on the virus expression. Regardless, all animals were used for control and at least
one stimulation condition with minimum 2 days apart by varying FOVs to avoid imaging the
same cells multiple times. The order of conditions and the imaging depth presented to the
same animal were randomized.
An additional single-cell stimulation only session was conducted on a subset of
animals, to further test a reliable holographic stimulation at a single-cell level (n = 3 animals).
We varied a stimulation position from the original target stimulation point to 10, 20, or 30 μm
shifted along x-axis or y-axis, generating 7 different stimulation point (original cell position,
10, 20, or 30 μm shifted along the x-axis, 10, 20, or 30 μm shifted along the y-axis). We
stimulated each stimulation point for 10 times in a randomized order across stimulation
points with 8-sec. ISIs.
569
*Analysis*
Imaging data were processed with 'suite2p' for motion correction, cell detection, and cell
fluorescence trace extraction [70]. We applied neuropil correction to the fluorescence traces
using the following equation: $F_{(cell\_corrected)} = F_{(cell)} - (0.8 * F_{(neuropil)})$. We then computed ΔF/F
normalized to the baseline, by following the equation: $\Delta F/F = (F_{(trace)} - F_{(baseline)}) / F_{(baseline)}$,
where baseline is about 300 ms before the sound onset. For the single-cell level stimulation
session, we computed peak ΔF/F per each stimulation point and applied a mixed-effect model
by taking peak ΔF/F as dependent variables, stimulation point as independent variables, and
cells as random factor. For the 5-cell stimulation validation session, we computed a
proportion of activated cells (any cell with peak ΔF/F > 0) among stimulated cells per each
FOV as well as peak ΔF/F and ran a random permutation with 100 iterations.
For experimental sessions, to select sound responsive cells, we compared sound-
evoked activity (160 ms – 660 ms after sound onset capturing the sound-evoked response due
to slow calcium transient) and the baseline activity (300 ms – 0 ms before the sound onset).
We considered cells sound responsive when the amplitude of the average sound-evoked
activity exceeded two standard deviations of the amplitude of the average baseline activity.
Sound responsive cells were selected based on fluorescence traces only from the baseline
session to minimize any potential effect of stimulation on cell selection. We then computed
the ratio of the evoked activity between two frequencies as an index of the frequency
preference ($\Delta F/F_{(16kHz)} - \Delta F/F_{(54kHz)}$). We subgrouped sound-responsive cells into either
16 kHz preferring cells or 54 kHz preferring cells based on the frequency preference, taking 0
as a subgroup criterion. As our main interest was changes to non-target cells, we excluded
target cells for further analyses. To compare response changes due to stimulation for each
group of cells, average sound-evoked activity of sound-responsive cells from the baseline
session was subtracted from the stimulation session (stimulation effect:

$\Delta F/F_{(stimulation\ session)} - \Delta F/F_{(baseline\ session)})$ for each condition. We further compared
response changes between post-stimulation session and stimulation session, again by
subtracting the response amplitudes between two sessions for each group and condition (post-
stimulation effect: $\Delta F/F_{(post-stimulation\ session)} - \Delta F/F_{(stimulation\ session)}$). To quantify the
stimulation effect based on functional properties for conditions and groups, we applied a
mixed-effect model by taking amplitude changes as dependent variables, frequency,
conditions, and cell groups as independent variables, and FOVs as random factor. We then
computed Type III analysis of variance (ANOVA) to examine whether the effect of
functional property (frequency) to the amplitude changes was specific to the target frequency
presentation synchronized with the stimulation (i.e., 16 kHz for 16 kHz target cell stimulation
or 54 kHz for 54 kHz target cell stimulation). To quantify whether the response amplitude
changes due to stimulation differ across trials, we fitted a dataset of average stimulation
effect across each trial per each condition (stimluation or control), each cell group (16 kHz
preferring or 54 kHz preferring cells), and each tone presentation (16 kHz or 54 kHz) to the
three-parameter model and computed the extra sum-of-squares $F$-test to compare whether
response amplitude changes across trials were different from a constant line [71]. To quantify a
relationship between response amplitude changes of non-target cells and their distances to
target cells, we computed a center of mass distance of each cell position relative to target
cells, and fitted the dataset of the response amplitude changes across distance to the three-
parameter model to compute the extra sum-of-squares $F$-test, per each condition (16 kHz
target stimulation, 54 kHz target stimulation, or control), each cell group (16 kHz preferring
or 54 kHz preferring cells), and each tone presentation (16 kHz or 54 kHz). For the control
condition, as there were no stimulated target cells, we chose top five most tone-responsive
cells from the baseline session as "target" cells.
We then generated a simple model in which a suppression term was applied either to
all neurons or specifically to non-target co-tuned cells to test our results from the data. We
took a similar range of number of neurons and FOVs to closely simulate the model to the real
dataset structure. On 50 simulated calcium traces of neurons (n),
623
624 $$\text{Trace}_{n(t)} = R_{n(t)} - \text{theta}_n + \text{epsilon}_{n(t)}$$
625
Where $R_{n(t)}$ is a response amplitude from either baseline or stimulation session, $\text{theta}_n$
is a suppression term applied either to all neurons or only to non-target co-tuned neurons,
only during the stimulation session, and $\text{epsilon}_{n(t)}$ is additive noise. Theta was defined as
proportional to the average stimulation strength from target neurons, derived from the real
dataset, and scaled by a factor $\alpha = 0.3$ in the current simulation. To introduce neuron-level
variability, an additional jitter ($\text{epsilon}_n$) was applied as follow:
632
633 $$\text{Theta}_n = \alpha * \text{mean(target stimulation amplitude)} * (1 + \text{epsilon}_n)$$
634
635 　　　　Similar to the real data analyses, we compared the response change between the
636 stimulation and baseline sessions' trace amplitudes.
637
638
*Histology*
Animals were deeply anesthetized with 4% isoflurane to perform transcardial perfusion with
4% paraformaldehyde (PFA) in 0.1 M phosphate buffer saline (PBS). The extracted brains
were post-fixed in 4% PFA for additional 12-24 hours. Coronal brain sections at 50 μm
containing the AC were stained with primary antibodies of HA-Tag (1:500) and chicken
Green Fluorescent Protein (GFP, 1:500) for GCaMP8s, and secondary antibodies of 594-

conjugated anti-rabbit IgG (1:1000) and 488-conjugated anti-chicken IgG (1:1000) for red-
shifted opsins.

**References**

1. Read, H., Winer, J., and Schreiner, C. (2001). Modular organization of intrinsic
connections associated with spectral tuning in cat auditory cortex. Proceedings of the
National Academy of Sciences of the United States of America *98*, 8042-8047.
10.1073/pnas.131591898.

2. Schnupp, J., Mrsic-Flogel, T., and King, A. (2001). Linear processing of spatial cues in
primary auditory cortex. Nature *414*, 200-204. 10.1038/35102568.

3. Lu, K., Liu, W., Zan, P., David, S.V., Fritz, J.B., and Shamma, S.A. (2018). Implicit
Memory for Complex Sounds in Higher Auditory Cortex of the Ferret. J Neurosci *38*,
9955-9966. 10.1523/JNEUROSCI.2118-18.2018.

4. Fritz, J., Shamma, S., Elhilali, M., and Klein, D. (2003). Rapid task-related plasticity of
spectrotemporal receptive fields in primary auditory cortex. Nature Neuroscience *6*,
1216-1223. 10.1038/nn1141.

5. Bandyopadhyay, S., Shamma, S.A., and Kanold, P.O. (2010). Dichotomy of functional
organization in the mouse auditory cortex. Nat Neurosci *13*, 361-368.
10.1038/nn.2490.

6. Rothschild, G., Nelken, I., and Mizrahi, A. (2010). Functional organization and
population dynamics in the mouse primary auditory cortex. Nature Neuroscience *13*,
353-U321. 10.1038/nn.2484.

7. Panzeri, S., Macke, J., Gross, J., and Kayser, C. (2015). Neural population coding:
combining insights from microscopic and mass signals. Trends in Cognitive Sciences
*19*, 162-172. 10.1016/j.tics.2015.01.002.

8. Bowen, Z., Winkowski, D., and Kanold, P. (2020). Functional organization of mouse
primary auditory cortex in adult C57BL/6 and F1 (CBAxC57) mice. Scientific Reports
*10*, ARTN 10905. 10.1038/s41598-020-67819-4.

9. Francis, N., Winkowski, D., Sheikhattar, A., Armengol, K., Babadi, B., and Kanold, P.
(2018). Small Networks Encode Decision-Making in Primary Auditory Cortex. Neuron
*97*, 885-+. 10.1016/j.neuron.2018.01.019.

10. Francis, N., Mukherjee, S., Kocillari, L., Panzeri, S., Babadi, B., and Kanold, P. (2022).
Sequential transmission of task-relevant information in cortical neuronal networks.
Cell Reports *39*, ARTN 110878. 10.1016/j.celrep.2022.110878.

11. Hromádka, T., Deweese, M.R., and Zador, A.M. (2008). Sparse representation of
sounds in the unanesthetized auditory cortex. PLoS Biol *6*, e16.
10.1371/journal.pbio.0060016.

12. Hromadka, T., and Zador, A. (2009). Representations in auditory cortex. Current
Opinion in Neurobiology *19*, 430-433. 10.1016/j.conb.2009.07.009.

13. Liang, F., Li, H., Chou, X., Zhou, M., Zhang, N., Xiao, Z., Zhang, K., Tao, H., and Zhang, L.
(2019). Sparse Representation in Awake Auditory Cortex: Cell-type Dependence,
Synaptic Mechanisms, Developmental Emergence, and Modulation. Cerebral Cortex
*29*, 3796-3812. 10.1093/cercor/bhy260.

14. Yu, L., and Yu, Y. (2017). Energy-Efficient Neural Information Processing in Individual
Neurons and Neuronal Networks. Journal of Neuroscience Research *95*, 2253-2266.
10.1002/jnr.24131.

15. Jadi, M., and Sejnowski, T. (2014). Cortical oscillations arise from contextual
interactions that regulate sparse coding. Proceedings of the National Academy of
Sciences of the United States of America *111*, 6780-6785. 10.1073/pnas.1405300111.

16.    Grabner, R., Neubauer, A., and Stern, E. (2006). Superior performance and neural
efficiency: The impact of intelligence and expertise. Brain Research Bulletin *69*, 422-
439. 10.1016/j.brainresbull.2006.02.009.
17.    Bowen, Z., Winkowski, D.E., Seshadri, S., Plenz, D., and Kanold, P.O. (2019). Neuronal
Avalanches in Input and Associative Layers of Auditory Cortex. Front Syst Neurosci *13*,
45. 10.3389/fnsys.2019.00045.
18.    Chettih, S.N., and Harvey, C.D. (2019). Single-neuron perturbations reveal feature-
specific competition in V1. Nature *567*, 334-340. 10.1038/s41586-019-0997-6.
19.    LaFosse, P.K., Zhou, Z., O'Rawe, J.F., Friedman, N.G., Scott, V.M., Deng, Y., and Histed,
M.H. (2024). Single-cell optogenetics reveals attenuation-by-suppression in visual
cortical neurons. bioRxiv. 10.1101/2023.09.13.557650.
20.    Winkowski, D.E., and Kanold, P.O. (2013). Laminar transformation of frequency
organization in auditory cortex. J Neurosci *33*, 1498-1508. 10.1523/JNEUROSCI.3101-
12.2013.
21.    Adesnik, H., and Abdeladim, L. (2021). Probing neural codes with two-photon
holographic optogenetics. Nat Neurosci *24*, 1356-1366. 10.1038/s41593-021-00902-
9.
22.    Yang, W., and Yuste, R. (2021). Holographic Imaging and Stimulation of Neural
Circuits. Optogenetics: Light-Sensing Proteins and Their Applications in Neuroscience
and Beyond, 613-639.
23.    Karmarkar, U., and Buonomano, D. (2007). Timing in the absence of clocks: Encoding
time in neural network states. Neuron *53*, 427-438. 10.1016/j.neuron.2007.01.006.
24.    Liu, B., Seay, M., and Buonomano, D. (2023). Creation of Neuronal Ensembles and
Cell-Specific Homeostatic Plasticity through Chronic Sparse Optogenetic Stimulation.
Journal of Neuroscience *43*, 82-92. 10.1523/JNEUROSCI.1104-22.2022.
25.    Kishi, K.E., Kim, Y.S., Fukuda, M., Inoue, M., Kusakizako, T., Wang, P.Y., Ramakrishnan,
C., Byrne, E.F.X., Thadhani, E., Paggi, J.M., et al. (2022). Structural basis for channel
conduction in the pump-like channelrhodopsin ChRmine. Cell *185*, 672-689.e623.
10.1016/j.cell.2022.01.007.
26.    LaFosse, P., Zhou, Z., Friedman, N., Deng, Y., Li, A., Akitake, B., and Histed, M. (2023).
Bicistronic Expression of a High-Performance Calcium Indicator and Opsin for All-
Optical Stimulation and Imaging at Cellular Resolution. Eneuro *10*.
10.1523/ENEURO.0378-22.2023.
27.    Sridharan, S., Gajowa, M., Ogando, M., Jagadisan, U., Abdeladim, L., Sadahiro, M.,
Bounds, H., Hendricks, W., Turney, T., Tayler, I., et al. (2022). High-performance
microbial opsins for spatially and temporally precise perturbations of large neuronal
networks. Neuron *110*, 1139-+. 10.1016/j.neuron.2022.01.008.
28.    Mardinly, A., Oldenburg, I., Pegard, N., Sridharan, S., Lyall, E., Chesnov, K., Brohawn,
S., Waller, L., and Adesnik, H. (2018). Precise multimodal optical control of neural
ensemble activity. Nature Neuroscience *21*, 881-+. 10.1038/s41593-018-0139-8.
29.    Daie, K., Svoboda, K., and Druckmann, S. (2021). Targeted photostimulation uncovers
circuit motifs supporting short-term memory. Nature Neuroscience *24*, 259-265.
10.1038/s41593-020-00776-3.
30.    Kane, K.L., Longo-Guess, C.M., Gagnon, L.H., Ding, D., Salvi, R.J., and Johnson, K.R.
(2012). Genetic background effects on age-related hearing loss associated with
Cdh23 variants in mice. Hear Res *283*, 80-88. 10.1016/j.heares.2011.11.007.

31.    Branchi, I., Santucci, D., and Alleva, E. (2001). Ultrasonic vocalisation emitted by
infant rodents: a tool for assessment of neurobehavioural development. Behav Brain
Res *125*, 49-56. 10.1016/s0166-4328(01)00277-7.

32.    Ulanovsky, N., Las, L., Farkas, D., and Nelken, I. (2004). Multiple time scales of
adaptation in auditory cortex neurons. J Neurosci *24*, 10440-10453.
10.1523/JNEUROSCI.1905-04.2004.

33.    Yarden, T.S., and Nelken, I. (2017). Stimulus-specific adaptation in a recurrent
network model of primary auditory cortex. PLoS Comput Biol *13*, e1005437.
10.1371/journal.pcbi.1005437.

34.    Malmierca, M.S., Sanchez-Vives, M.V., Escera, C., and Bendixen, A. (2014). Neuronal
adaptation, novelty detection and regularity encoding in audition. Front Syst
Neurosci *8*, 111. 10.3389/fnsys.2014.00111.

35.    Schreiner, C.E., Read, H.L., and Sutter, M.L. (2000). Modular organization of frequency
integration in primary auditory cortex. Annu Rev Neurosci *23*, 501-529.
10.1146/annurev.neuro.23.1.501.

36.    Liu, J., and Kanold, P.O. (2021). Diversity of Receptive Fields and Sideband Inhibition
with Complex Thalamocortical and Intracortical Origin in L2/3 of Mouse Primary
Auditory Cortex. J Neurosci *41*, 3142-3162. 10.1523/JNEUROSCI.1732-20.2021.

37.    Froemke, R., and Hyman, S. (2015). Plasticity of Cortical Excitatory-Inhibitory Balance.
Annual Review of Neuroscience, Vol 38 *38*, 195-219. 10.1146/annurev-neuro-
071714-034002.

38.    Rupasinghe, A., Francis, N., Liu, J., Bowen, Z., Kanold, P., and Babadi, B. (2021). Direct
extraction of signal and noise correlations from two-photon calcium imaging of
ensemble neuronal activity. Elife *10*, ARTN e68046. 10.7554/eLife.68046.

39.    Ko, H., Mrsic-Flogel, T., and Hofer, S. (2014). Emergence of Feature-Specific
Connectivity in Cortical Microcircuits in the Absence of Visual Experience. Journal of
Neuroscience *34*, 9812-9816. 10.1523/JNEUROSCI.0875-14.2014.

40.    Katz, L., and Shatz, C. (1996). Synaptic activity and the construction of cortical circuits.
Science *274*, 1133-1138. 10.1126/science.274.5290.1133.

41.    Ohtsuki, G., Nishiyama, M., Yoshida, T., Murakami, T., Histed, M., Lois, C., and Ohki, K.
(2012). Similarity of visual selectivity among clonally related neurons in visual cortex.
Neuron *75*, 65-72. 10.1016/j.neuron.2012.05.023.

42.    Li, Y., Lu, H., Cheng, P.L., Ge, S., Xu, H., Shi, S.H., and Dan, Y. (2012). Clonally related
visual cortical neurons show similar stimulus feature selectivity. Nature *486*, 118-121.
10.1038/nature11110.

43.    Hebb, D.O. (2005). The organization of behavior: A neuropsychological theory
(Psychology press).

44.    Malenka, R.C., and Nicoll, R.A. (1999). Long-term potentiation--a decade of progress?
Science *285*, 1870-1874. 10.1126/science.285.5435.1870.

45.    Brown, T.H., Kairiss, E.W., and Keenan, C.L. (1990). Hebbian synapses: biophysical
mechanisms and algorithms. Annual review of neuroscience *13*, 475-511.

46.    Vitureira, N., and Goda, Y. (2013). The interplay between Hebbian and homeostatic
synaptic plasticity. Journal of Cell Biology *203*, 175-186. 10.1083/jcb.201306030.

47.    Fox, K., and Stryker, M. (2017). Integrating Hebbian and homeostatic plasticity:
introduction. Philosophical Transactions of the Royal Society B-Biological Sciences
*372*, ARTN 20160413. 10.1098/rstb.2016.0413.

48. Toyoizumi, T., Kaneko, M., Stryker, M., and Miller, K. (2014). Modeling the Dynamic
Interaction of Hebbian and Homeostatic Plasticity. Neuron *84*, 497-510.
10.1016/j.neuron.2014.09.036.
49. Turrigiano, G., and Nelson, S. (2000). Hebb and homeostasis in neuronal plasticity.
Current Opinion in Neurobiology *10*, 358-364. 10.1016/S0959-4388(00)00091-X.
50. Yin, J., and Yuan, Q. (2015). Structural homeostasis in the nervous system: a
balancing act for wiring plasticity and stability. Frontiers in Cellular Neuroscience *8*,
ARTN 439. 10.3389/fncel.2014.00439.
51. Maor, I., Shalev, A., and Mizrahi, A. (2016). Distinct Spatiotemporal Response
Properties of Excitatory Versus Inhibitory Neurons in the Mouse Auditory Cortex.
Cerebral Cortex *26*, 4242-4252. 10.1093/cercor/bhw266.
52. Wang, Y., Ye, M., Kuang, X., Li, Y., and Hu, S. (2018). A simplified morphological
classification scheme for pyramidal cells in six layers of primary somatosensory
cortex of juvenile rats. Ibro Reports *5*, 74-90. 10.1016/j.ibror.2018.10.001.
53. Marin, O. (2012). Interneuron dysfunction in psychiatric disorders. Nature Reviews
Neuroscience *13*, 107-120. 10.1038/nrn3155.
54. Xu, J., Fan, J., Wang, X., Eacker, S., Kam, T., Chen, L., Yin, X., Zhu, J., Chi, Z., Jiang, H., et
al. (2016). Cultured networks of excitatory projection neurons and inhibitory
interneurons for studying human cortical neurotoxicity. Science Translational
Medicine *8*, ARTN 333ra48. 10.1126/scitranslmed.aad0623.
55. Wonders, C., and Anderson, S. (2006). The origin and specification of cortical
interneurons. Nature Reviews Neuroscience *7*, 687-696. 10.1038/nrn1954.
56. Nathanson, J., Yanagawa, Y., Obata, K., and Callaway, E. (2009). Preferential Labeling
of Inhibitory and Excitatory Cortical Neurons by Endogenous Tropism of Adeno-
associated Virus and Lentivirus Vectors. Neuroscience *161*, 441-450.
10.1016/j.neuroscience.2009.03.032.
57. Kerlin, A.M., Andermann, M.L., Berezovskii, V.K., and Reid, R.C. (2010). Broadly tuned
response properties of diverse inhibitory neuron subtypes in mouse visual cortex.
Neuron *67*, 858-871. 10.1016/j.neuron.2010.08.002.
58. Mateo, C., Avermann, M., Gentet, L.J., Zhang, F., Deisseroth, K., and Petersen, C.C.
(2011). In vivo optogenetic stimulation of neocortical excitatory neurons drives brain-
state-dependent inhibition. Curr Biol *21*, 1593-1602. 10.1016/j.cub.2011.08.028.
59. Wehr, M., and Zador, A.M. (2003). Balanced inhibition underlies tuning and sharpens
spike timing in auditory cortex. Nature *426*, 442-446.
60. Chen, I.W., Helmchen, F., and Lütcke, H. (2015). Specific early and late oddball-evoked
responses in excitatory and inhibitory neurons of mouse auditory cortex. Journal of
Neuroscience *35*, 12560-12573.
61. Meyer, H., Schwarz, D., Wimmer, V., Schmitt, A., Kerr, J., Sakmann, B., and
Helmstaedter, M. (2011). Inhibitory interneurons in a cortical column form hot zones
of inhibition in layers 2 and 5A. Proceedings of the National Academy of Sciences of
the United States of America *108*, 16807-16812. 10.1073/pnas.1113648108.
62. Tasaka, G.I., Maggi, C., Taha, E., and Mizrahi, A. (2023). The local and long-range
input landscape of inhibitory neurons in mouse auditory cortex. J Comp Neurol *531*,
502-514. 10.1002/cne.25437.
63. Markram, H., Toledo-Rodriguez, M., Wang, Y., Gupta, A., Silberberg, G., and Wu, C.
(2004). Interneurons of the neocortical inhibitory system. Nat Rev Neurosci *5*, 793-
807. 10.1038/nrn1519.

64.    Li, J.Y., and Glickfeld, L.L. (2023). Input-specific synaptic depression shapes temporal
integration in mouse visual cortex. Neuron. 10.1016/j.neuron.2023.07.003.

65.    Davis, G. (2013). Homeostatic Signaling and the Stabilization of Neural Function.
Neuron *80*, 718-728. 10.1016/j.neuron.2013.09.044.

66.    Tien, N., Soto, F., and Kerschensteiner, D. (2017). Homeostatic Plasticity Shapes Cell-
Type-Specific Wiring in the Retina. Neuron *94*, 656-+. 10.1016/j.neuron.2017.04.016.

67.    Maffei, A., and Fontanini, A. (2009). Network homeostasis: a matter of coordination.
Current Opinion in Neurobiology *19*, 168-173. 10.1016/j.conb.2009.05.012.

68.    Barak, O., and Tsodyks, M. (2007). Persistent activity in neural networks with dynamic
synapses. Plos Computational Biology *3*, 323-332, ARTN e35.
10.1371/journal.pcbi.0030035.

69.    Liu, J., Whiteway, M.R., Sheikhattar, A., Butts, D.A., Babadi, B., and Kanold, P.O. (2019).
Parallel Processing of Sound Dynamics across Mouse Auditory Cortex via Spatially
Patterned Thalamic Inputs and Distinct Areal Intracortical Circuits. Cell Rep *27*, 872-
885.e877. 10.1016/j.celrep.2019.03.069.

70.    Pachitariu, M., Stringer, C., Schröder, S., Dipoppa, M., Rossi, L.F., Carandini, M., and
Harris, K.D. (2016). Suite2p: beyond 10,000 neurons with standard two-photon
microscopy. BioRxiv, 061507.

71.    Motulsky, H., and Christopoulos, A. (2004). Fitting models to biological data using
linear and nonlinear regression: a practical guide to curve fitting (Oxford University
Press).

853
854

855 **Acknowledgments:** Supported by U19 NS107464 (POK), NIH RO1DC017785 (POK), NIH
856 F32DC019842 (TAB).
857
858 **Author Contributions:** Conceptualization: HK, TAB, POK; Methodology: HK, TAB, POK;
859 Investigation: HK, TAB; Writing: HK, POK
860
861 **Competing interest:** Authors declare that they have no competing interests.
862
863 **Data and materials availability:** All preprocessed imaging data and relevant analyses
864 scripts will be deposited at Johns Hopkins University research data repository, available for
865 open access.
866
867

**Figures**

**Figure 1. Holographic optogenetic stimulation in AC and experimental procedure. A**: An example brain slice showing cells in AC expressing both GCaMP and opsin (AAV9-hSyn-GC8s-T2A-rsChRmine). **B**: An example field of view (FOV) where single cell targeting precision was tested and response traces to the holographic stimulation from an example cell. Stimulation was offset from the original position (red circle) to distance-shifted positions in 10 μm increments (gray dashed circles in the x-axis or y-axis of the FOV). Responses were the greatest when the stimulation was performed on the original cell position (red solid line trace). Rapid amplitude decay along the position shift was observed (red dashed line traces). Grey error shades indicate SEM across trials. A right inset errorbar plot shows a grand average amplitude change per stimulation point across all tested cells (n = 15 cells, 3 animals). Error bars indicate SEM across cells. **C**: An example FOV showing a population of cells (left) and amplitude changes to 5-cell stimulation as a stimulation effect ($\Delta$ F/F$_{stim}$ - $\Delta$ F/F$_{spont., right}$). Filled squares indicate each cell. White circles indicate stimulation targeted cells. **D**: (left) Proportion of stimulated cells that showed an increase in fluorescence following photostimulation. Error bars indicate SEM across FOVs. A horizontal

dashed line indicates average permutation results (random permutation test on 100 iterations,
$p < 0.0001$). (right) Grand average of the stimulation effect across imaging sessions. Error
bars indicate SEM across FOVs. A horizontal dashed line indicates average permutation
results (random permutation test on 100 iterations, $p < 0.0001$). **E**: Experimental procedure.
A total of four consecutive imaging sessions were acquired: 1) A cell selection session to
identify neurons selective for 16 kHz pure tones, 2) a baseline imaging session to acquire
tone-evoked activity response to either 16 or 54 kHz pure tone, 3) a stimulation session
representing five cells of either 16 kHz or 54 kHz responsive cells as target stimulation to
examine the effect of stimulation synchronized to tone presentations, and 4) a post-
stimulation session to examine network persistence after stimulation-related changes.
896

**Figure 2. Targeted cells and non-target cells show response changes due to stimulation.**
**A**: (top) An example FOV showing the stimulation effect ($\Delta F/F_{stim} - \Delta F/F_{baseline}$) of sound responsive cells for 16 kHz target cell stimulation (filled squares). Black circles indicate stimulated target cells. (bottom) Mean response traces of an example target and non-target cell in baseline session (black) and stimulation session (red). Error bars indicate SEM across trials. An example target cell shows an increased response due to the stimulation. An example non-target cell shows a decreased response due to stimulation on the target cells. **B**: (left) A violin plot of the proportion of stimulated cells that showed increased activity due to stimulation across FOVs. Horizontal solid line indicates mean proportion, empty circle indicates median proportion, and gray filled circles indicate individual FOVs. (right) Mean amplitude changes of target cells for stimulation session and post-stimulation session normalized to the baseline session. Error bars indicate SEM across cells. Dashed horizontal lines on both panels indicate average permutation results (random permutation test on 100 iterations, $p < 0.0001$). **CD**: Same as ABC for 54 kHz target cell stimulation.

914
**Figure 3. Non-target co-tuned cells show more decreased response amplitudes due to**
**stimulation when synchronized with their preferred tones. A**: Stimulation effect ($\Delta$ F/F$_{stim}$
- $\Delta$ F/F$_{baseline}$) in all sound responsive cells, including both target and non-target cells,
responding to either 16 kHz (blue) or 54 kHz (orange) pure tones, representing global activity
changes due to the stimulation. No significant differences between stimulation conditions and
responses to different frequencies were observed (all $p > 0.05$). **B**: Stimulation effect ($\Delta$
F/F$_{stim}$ - $\Delta$ F/F$_{baseline}$) in 16 kHz (blue) and 54 kHz (orange) preferring cells. Both cell groups
show decreased amplitude to their preferred frequency regardless of conditions due to
acoustic stimulus-specific adaptation. Only co-tuned cells (16 kHz preferring cells for 16 kHz
stimulation or 54 kHz preferring cells for 54 kHz stimulation) show a further decrease in
response amplitudes due to the stimulation, when the preferred pure tone (PT) frequency was
synchronized. Error bars indicate SEM across FOVs (*: $p < 0.0001$). **C**: Stimulation effect
from the model prediction. Amplitude changes computed from simulated data by applying
cell suppression to all cells (All supp.) or only co-tuned cells (Co-tuned supp.) were
compared with real data. Only the Co-tuned supp. model showed a significant amplitude
decrease for co-tuned neurons compared to non co-tuned neurons, similar to the result from
the real data ($p < 0.05$; see texts for more detail). **D**: Post-stimulation effect ($\Delta$ F/F$_{post}$ - $\Delta$
F/F$_{stim}$) 16 kHz (blue) and 54 kHz (orange) preferring cells. No significant response
amplitude changes were observed. Error bars indicate SEM across FOVs. **E**: Sub-
categorization of cells based on the frequency selectivity for each target stimulation condition
(left: 16 kHz stim, right 54 kHz stim). Cells were first grouped into either 16 kHz preferring
cells (blue) or 54 kHz preferring cells (orange). Within each cell group, cells were further
subdivided into low, mid, and high frequency selectivity categories based on their 33%

quartile ranges. Note that the frequency preference was log-transformed for visualization, but
the x-axis labels kept the original frequency selectivity values before transformation. Vertical
dashed lines indicate 33% quartile ranges. **F**: Response amplitude change based on the
frequency selectivity category for each cell groups (blue: 16 kHz preferring cells, orange: 54
kHz preferring cells). Significant response amplitude changes relative to the control condition
were observed only for high frequency selectivity category when target stimulated cells were
co-tuned (*: $p < 0.05$).
945

946
**Figure 4. Rebalanced response changes on non-target 16 kHz cells are immediate and**
**widely distributed. A**: Stimulation effect ($\Delta$ F/F$_{stim}$ - $\Delta$ F/F$_{baseline}$) in 16 kHz (blue) and 54
kHz (orange) preferring cells per each trial for the 5-cell 16 kHz stimulation condition (left),
54 kHz stimulation condition (middle), and the no-cell control condition (right). Each circle
represents average stimulation effect per each trial. Decreased amplitudes to preferred
frequencies were observed from as early as trial 1 with no significant further changes across
trials, regardless of frequencies and conditions (sum-of-squares *F*-test, all *p* > 0.05). Solid
lines indicate fitted curves and dashed lines indicate 95% confidence intervals. **B**: Stimulation
effect ($\Delta$ F/F$_{stim}$ - $\Delta$ F/F$_{baseline}$) of each non-target 16 kHz (blue) or 54 kHz (orange) preferring
cells for either 16 kHz (top row) or 54 kHz (bottom row) presentation in relation to the mass
of center distance to any target cells for the stimulation condition (left), 54 kHz stimulation
condition (middle), and the control condition (right). Each circle represents each cell. For the
control condition, we considered top 5 most tone-responsive cells from the baseline session
as "*target*" cells, as there was no stimulation. Non-target cells are widely distributed within
the FOV (550 $\mu$m$^2$), regardless of cell groups, frequencies, and conditions. Gray lines
indicate fitted curves, excluding cells that are closer than 15 $\mu$m (vertical green lines; cells <
15 $\mu$m marked in lighter shades), and dashed lines indicate 95% confidence intervals.
964