Rapid rebalancing of co-tuned ensemble activity in the auditory cortex

HiJee Kang[1], Travis A. Babola[1], & Patrick O. Kanold[1,2*]

Affiliations:

[1] Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD 21205

[2] Kavli Neuroscience Discovery Institute, Johns Hopkins University, Baltimore, MD 21205

*Corresponding author. Email: pkanold@jhu.edu

Keywords: cortex, optogenetics, holography, balance, plasticity

Abstract: Sensory information is represented by small varying neuronal ensembles in sensory cortices. In the auditory cortex (AC) repeated presentations of the same sound activate differing ensembles indicating high trial-by trial variability in activity even though the sounds activate the same percept. Efficient processing of complex acoustic signals requires that these sparsely distributed neuronal ensembles actively interact in order to provide a constant percept. Thus, the differing ensembles might interact to process the incoming sound inputs. Here, we probe interactions within and across ensembles by combining *in vivo* 2-photon Ca^{2+} imaging and holographic optogenetic stimulation to study how increased activity of single cells level affects the cortical network. We stimulated a small number of neurons sharing the same frequency preference alongside the presentation of a target pure tone, further increasing their tone-evoked activity. We found that other non-stimulated co-tuned neurons decreased their tone-evoked activity when the frequency of the presented pure tone matched to their tuning property, while non co-tuned neurons were unaffected. Activity decrease was greater for non-stimulated co-tuned neurons with higher frequency selectivity. Co-tuned and non co-tuned neurons were spatially intermingled. Our results shows that co-tuned ensembles communicated and balanced their total activity across the larger network. The rebalanced network activity due to external stimulation remained constant. These effects suggest that co-tuned ensembles in AC interact and rapidly rebalance their activity to maintain encoding network dynamics, and that the rebalanced network is persistent.

Introduction

Sensory perception requires fast encoding of relevant stimuli from a mixture of complex signals. Sensory cortices play a vital role in such sensory processing. In the auditory domain, for example, small neuronal ensembles in the auditory cortex (AC) are actively engaged to efficiently perceive relevant acoustic information [1-4]. The AC contains multiple ensembles of neurons that can be functionally identified, e.g., those formed by subsets of neurons preferring the same frequency also referred to as co-tuned neurons [5-7]. Repeated presentation of the same acoustic stimulus, e.g., a tone of the same frequency leads to a stable percept, but in the AC different ensembles of neurons are activated together at each repeat indicating a high trial-by-trial variability [5,8-10]. Activation of these different subsets of co-tuned neurons at each presentation of a stimulus reflects a sparse encoding of sound stimuli. Such sparse representation of co-activated neurons enables efficient coding with reduced metabolic energy to process complex information [11-16]. The sparse neuronal representation raises key questions of how activation of different ensembles leads to the same percept and how the overall activity within the cortical network is balanced across ensembles of co-tuned neurons. In particular, when a specific sound is present, a subset of co-tuned neurons will be activated, but not all co-tuned neurons [17]. Given that the percept of a repeating stimulus is constant, we speculated that neural activity is balanced across co-tuned as well as non co-tuned ensembles.

While neuronal ensembles constantly update their activities based on incoming information, how the activation of a particular sparse neuronal ensemble affects other neurons within the network to maintain the overall network balance for processing specific sensory information *in vivo* is largely unknown. *In vivo* optogenetic stimulation studies in the visual cortex (VC) suggested that inhibitory processes play a role in balancing network activity. In particular, *in vivo* single-cell holographic stimulation on a group of target cells, which induced increased response amplitude, resulted in changes in the response amplitudes of neighboring non-target neurons in the primary visual cortex (V1) [18], with similarly tuned neurons' activity being suppressed. Moreover, *in vivo* holographic optogenetic stimulation showed that visually-suppressed neurons had attenuated response amplitudes when holographic stimulation was given along with the visual stimulus presentation, which was not observed in visually activated neurons [19]. This suggested that neurons exhibit supralinear-to-linear input-output (IO) functions *in vivo*, rather than threshold-linear IO functions observed *in vitro*. These studies suggest that inhibitory influence from additional neuronal activation in the VC seems to play a major role during *in vivo* sensory processing, likely to maintain the activity balance of the network by modulating activities of neighboring neurons that share a similar tuning property.

One major difference between VC and AC is that the frequency tuning of neurons in the AC is less spatially localized, especially in a superficial layer (Layer 2/3) [20]. The local frequency preferences in the AC are diverse, thus neighboring neurons can show widely differing tuning properties [5]. To test how activity in specific AC cells among an intermingled and spatially distributed co-tuned and non co-tuned cell population is balanced during auditory processing, we stimulated a small group of AC cells using *in vivo* holographic optogenetic stimulation [21,22] while imaging AC population activity using 2-photon Ca^{2+} imaging in awake mice. We further tested whether any activity changes induced by holographic stimulation persist, as recurrent cortical networks engage homeostatic plasticity to stabilize overall network activity levels [23,24]. Stimulating small ensembles of co-tuned neurons together with the presentation of a pure tone in their preferred frequency increased their tone-evoked activity. Furthermore, we observed that non-stimulated co-tuned neurons decreased their tone-evoked activity. Non co-tuned ensembles did not exhibit such changes in tone-evoked responses, regardless of the pure tone frequency. Thus, the increased activity in the stimulation-targeted ensemble had caused a decrease in activity in the non-stimulated co-tuned ensembles, specifically when the stimulation-paired pure tone was their preferred frequency. Non-target co-tuned neurons exhibiting such effects were not necessarily neighboring the targeted cells,

86 suggesting specific interactions between co-tuned but not co-located neurons. Lastly, the
87 decreased activity in the non-stimulated co-tuned ensembles persisted in the subsequent
88 imaging session, even in the absence of holographic stimulation. These results suggest that co-
89 tuned ensembles form interacting overall networks that balance their activity.
90
Results
Optogenetic holographic stimulation increases activity in small ensembles *in vivo*
To sparsely manipulate neuronal ensembles, we used *in vivo* holographic stimulation.
To achieve reliable and selective *in vivo* holographic optogenetic stimulation of small
ensembles of neurons with single cell precision, we generated an AAV co-expressing the red-
shifted opsin rsChRmine and GCaMP8s (AAV9-hSyn-GCaMP8s-T2A-rsChRmine), as
rsChRmine minimizes the optical cross talk reducing a possible activation from the imaging
laser (940 nm excitation wavelength) [25]. Injecting AAV9-hSyn-GCaMP8s-T2A-rsChRmine
into AC yielded cells expressing both GCaMP and opsin (Fig. 1A). We first tested the
efficiency and reliability holographic stimulation, by targeting either a single cell or a small
ensemble of five cells. For single cell stimulation, we varied the stimulation point from the
target cell position by 10, 20, and 30 μm along either the x-axis or y-axis of the fields of view
(FOV; n = 15 cells, 3 animals). This results in a rapid decay of response amplitudes to
stimulation by the distance shift from the original cell position, confirming reliable holographic
stimulation at the single-cell level (~15 μm diameter) (mixed-effect model, $p < 0.05$; Fig. 1B).
Furthermore, the stimulation effect of target cells was specific to the targeted z-plane, showing
no stimulation effect when the stimulation z-plane was off by 20 μm (Fig. S1). For 5-cell
stimulation, a majority of cells reliably responded to photo-stimulated *in vivo* (5 mW/cell, 15
μm spiral, 30 revolutions, 6 s inter-stimulus interval (ISI), 5 trials) and exhibited robust Ca^{2+}
responses (Fig. 1CD; permutation test, all $p < 0.05$), comparable with responses to other opsins
[26-29]. Thus, *in vivo* holographic stimulation enables precise targeting and activation of groups
of single neurons in AC.
113
Optogenetic holographic stimulation increases sound evoked activity in A1 ensembles
Since repeated sound stimulation activates different ensembles while resulting in the
same percept, we reasoned that ensembles interacted and speculated that increased activity in
one ensemble would prevent or reduce activity in co-tuned ensembles. We thus next sought to
investigate how increased neural activity in small co-tuned ensembles during sound
presentation affected sound-evoked responses in targeted and non-target co-tuned and non co-
tuned ensembles. To achieve this, we first needed to identify the tuning properties of single
neurons and then target a subset of co-tuned neurons for stimulation. To study how the
increased activity of a small number of neurons influences the activity of other neurons
according to their frequency tuning properties, we designed an experimental paradigm
comprising four sequential imaging sessions (Fig. 1E):
First, in the cell selection session (Fig. 1E), we identified tuned ensembles in primary
auditory cortex (A1) layer 2/3 (L2/3) by assessing frequency tuning properties of neurons
within the FOVs covering 550 μm² (total cells = 7344, sound responsive cells = 1331, FOVs =
23; Fig. 1E). We presented pure tones of three different frequencies spanning the hearing range
of the mouse (4 kHz, 16 kHz, and 54 kHz, 100 ms duration, 2 sec. ISIs, 10 repeats for each
frequency). We chose 16 kHz and 54 kHz as the representative target ensemble tone
frequencies, as 16 kHz is within the most sensitive frequency range of mice [30] and 54 kHz is
within the range of mouse ultrasonic vocalization [31]. By selecting target ensembles in two
different frequencies, we ensured that effects of stimulation were not specific to one particular
population. For each condition (16 kHz or 54 kHz target ensemble for stimulation), we selected
5 target cells to stimulate. To ensure that all cells in the ensemble were selective for the target
tone, we chose the most responsive cells in each condition. Thus, for the 16 kHz target

ensemble condition, we selected five cells (*target cells*) among the top 30% most responsive
cells to the 16 kHz tone. Similarly, for the 54 kHz target ensemble condition, we selected five
of the top 30% most 54 kHz tone responsive cells. By selecting target cells sharing the same
frequency preference, we aimed at investigating how activity changes from co-tuned neuronal
ensembles alter the processing of the target frequency in other co-tuned and non co-tuned cells.
Second, in the baseline session (Fig. 1E), we determined the sound-evoked responses
of all imaged cells by presenting a series of 16 kHz and 54 kHz pure tones in a random order
(100 ms duration, 5.8-6.5 sec. ISI; baseline session, 30 repeats for each frequency). Exemplar
responses of cells from a 16 kHz and a 54 kHz ensemble are shown as black traces in Figure
2A and 2C.
Third, in the stimulation session (Fig. 1E), we examined how all sound-responsive cells
change their responses when a small group of cells in the network increases their activity. We
presented the same tones (16 kHz and 54 kHz in a random order), in tandem with the
optogenetic stimulation of five target cells (stimulation session, 100 ms stimulation duration).
We performed different sessions for the 16 kHz and 54 kHz target ensembles, varying FOVs
for each session (18 FOVs for 16 kHz target ensemble condition and 15 FOVs for 54 kHz target
ensemble condition). Figure 2A and C show two example FOVs with targeted neurons for a 16
kHz and 54 kHz ensemble, respectively.
Since both imaging and optogenetic stimulation involve optomechanical components,
we wanted to ensure that effects were not due to artifacts caused by our stimulation or imaging
setup. Moreover, cells can adapt their responses to repeated sound presentation. Thus, to
confirm any response changes observed from the stimulation session is due to the optogenetic
stimulation rather than simple response change due to acoustic sound presentation, we added
an additional control condition. For this control condition, we performed the "stimulation"
session with five target cells but with 0 mW laser power (i.e., no stimulation) to verify that any
response changes occurring in the stimulation session compared to the baseline session were
not simply due to the eventual response adaptation of neurons to the tuned frequency (control
condition; 13 FOVs). By selecting cells and presenting 0 mW laser power, instead of no target
cell selection or selecting any other no-cell area within the FOV, we ensured that the laser
power given to selected cells was the only difference between the actual stimulation and control
conditions, keeping any noise caused by the imaging and stimulation setup the same.
Fourth, after the stimulation session (Fig. 1E), we performed an additional imaging
session (post-stimulation session), presenting another series of 16 kHz and 54 kHz pure tones
in a random order to examine whether changes in the sound-evoked responses persisted or
reverted back after the stimulation session.
172

Optogenetic holographic stimulation increases activity in targeted ensembles
We first identified the effect of the optogenetic stimulation on the targeted ensembles.
Figures 2A and 2C show fluorescence traces of exemplar cells from 16 kHz and 54 kHz target
ensembles. Optogenetic stimulation increased the sound-evoked fluorescence amplitude in
these individual cells. To quantify the effect of the optogenetic stimulation, we compared the
tone-evoked fluorescence responses of the targeted cells with and without stimulation
(Stimulation effect $= \Delta F/F_{(stimulation\ session)} - \Delta F/F_{(baseline\ session)}$). Around 72% of target
cells (66 out of 90 cells over 18 FOVs for 16 kHz target stimulation and 42 out of 60 cells 15
FOVs for 54 kHz target stimulation) showed increased response amplitude during the
stimulation session compared to the baseline session, regardless of the tone presented (Fig.
2BD; permutation tests, all $p < 0.001$). These results indicate that holographic stimulation was
able to reliably increase activity in small populations of neurons. Moreover, given that the
target cells we selected were most responsive to their preferred tone frequency, this increase in
fluorescence indicates that the cells' responses to their preferred tone were not saturated.
187

Optogenetic holographic stimulation decreases activity in non-target co-tuned ensembles

We next investigated whether the optogenetically enhanced sound-evoked activity of a
small group of cells would cause activity changes in other non-stimulated cells. During
holographic optogenetic stimulation of the targeted cells, the non-target, but sound-responsive
cells (n = 995 cells for 16 kHz target ensemble condition and n = 675 cells for 54 kHz target
ensemble condition), also changed their activity, showing either increased or decreased
response amplitudes (Fig. 2A bottom and 2C bottom).
If cortical networks rebalance their activity, we speculated that the increased tone-
evoked activity in the targeted ensemble would lead to a decrease in tone-evoked activity in
coupled ensembles. Such rebalancing would keep the activity within the cortical network
stable. Moreover, given that we increased the activity to the preferred sound frequency, if this
rebalancing happens, it should occur only for the distinct sound frequency related to the cell's
tuning property. For example, stimulation of a 16 kHz ensemble should cause a greater
reduction in the 16 kHz tone response of non-targeted 16 kHz cells compared to their response
to the 54 kHz tone.
To address these questions, we investigated whether increased activity in the targeted
cells influenced the activity of non-target cells and how these changes were related to the tuning
properties of the cells. We first confirmed that the overall population activity from sound
responsive cells, including both target and non-target cells, did not differ across conditions
(control, 16 kHz, or 54 kHz target ensemble conditions; all $p > 0.05$, Fig. 3A). This suggests
that non-target cells may adjust their activities during the target cell stimulation to maintain the
global network activity level. To identify the activity changes based on functional
characteristics of cells, we defined each sound-responsive cell's frequency selectivity by
computing a difference between response amplitude to 16 kHz and 54 kHz from the baseline
session (frequency preference $= (\Delta F/F_{(16kHz)}) - (\Delta F/F_{(54kHz)})$). We then divided these cells
into either 16 kHz preferring or 54 kHz preferring groups, taking 0 (i.e., no preference) as a
criterion (Fig. 3B). Both subgroups exhibited stronger tone-evoked responses to their preferred
frequency, independent of the condition ($t(5700) = 4.79$, $p < 0.0001$; Fig. S2). This confirms
that the criterion for cell group threshold is valid.
We then focused on our main question by comparing the stimulation effect of the two
target ensemble groups to the control condition to identify whether stimulation decreased the
response of non-target co-tuned neurons. Neural activity in AC rapidly shows stimulus-specific
adaptation to the repeated presentation of the stimulus[32-34], which can obscure stimulation
related changes. We thus used the response amplitude change between the baseline and the
"stimulation" control session as a representative threshold to test the effect of the stimulation.
We once again used the difference in response amplitude between the baseline and stimulation
sessions as the measure of the stimulation effect ($\Delta F/F_{(stimulation\ session)} - \Delta F/$
$F_{(baseline\ session)}$). Neighboring cells within 20 μm from the target stimulation point were
removed from the analysis since they could have been directly affected by the stimulation.
We compared the stimulation effect between non-target co-tuned and non co-tuned cells
across conditions (16 kHz and 54 kHz target ensembles as well as control conditions) for
different pure tone presentations. Since our primary interest was how non-target cells respond
to increased activity in target ensembles, we focused on conditions where the pure tone
frequency matched or did not match the tuning properties of the non-target cells. Since we
stimulated during tone presentation the effects of the holographic stimulation and stimulus-
specific adaptation co-occurred. To isolate these components, we used a linear mixed-effect
model with cell group, condition, and pure tone frequency as fixed factors, and FOVs as a
random factor. We then performed ANOVA on the model to assess the main effects and
interactions.
A marginal significant main effect of the condition ($F(2,37.1) = 2.983$, $p = 0.0628$) on
the response change in the stimulation session relative to the baseline session (i.e., stimulation

effect) was observed, indicating that these changes may depend on the stimulation condition.
We further analyzed the data to better understand how the different factors interacted in the
response amplitude changes. A significant interaction between the pure tone frequency and cell
group ($F(1,4397.6) = 186.967$, $p < 0.0001$) suggests that each cell group responded differently
to the two pure tone frequencies. Specifically, the response amplitude decreases in the
stimulation session relative to the baseline session were more pronounced for each cell group
when the played pure tone matched to their tuning property. This interaction between pure tone
frequency and cell group highlights the importance of frequency tuning in modulating response
amplitudes. Such response amplitude decreases of non-target cells to their preferred pure tone
presentation further aligns with the stimulus specific adaptation due to repeatedly presented
pure tones [32]. Additionally, a significant three-way interaction across condition, cell groups,
and pure tone frequency ($F(2,4397.6) = 3.517$, $p = 0.0298$) suggests the combined effects of
the stimulation condition and the cell group on response amplitude depend on the pure tone
frequency. The stimulation effect is not uniform across cell groups and depends heavily on the
frequency, highlighting a complex interplay between the tuning property of cells, stimulation
condition, and presented pure tone frequency.
Consequently, we analyzed post-hoc comparisons estimated marginal means with
contrasts, as our focus was how co-tuned cells change their responses due to the increased
activity in the target cells along with the frequency of the presented pure tone.
For 16 kHz preferring cell group (n = 537), we observed a greater stimulation effect
(i.e., decrease in response amplitude) for 16 kHz tone presentation when 16 kHz target
ensemble was stimulated compared to the control condition ($t(124) = 3.114$, $p = 0.0064$). For
all other pairs, no significant stimulation effect was observed. This suggests that non-target 16
kHz co-tuned cells reduce their response amplitudes when target ensembles share the same
tuning property. Furthermore, such response change occurs only when they process their
preferred frequency (Fig. 3C, left).
We repeated the experiments and the analysis with 54kHz cells as the target group. In
general, we observed similar results. The stimulation effect was significantly more pronounced
for 54 kHz tone presentation when 54 kHz target ensemble (n = 359) was stimulated compared
to the control condition ($t(168) = 3.074$, $p = 0.0069$; all p-values were adjusted for multiple
comparisons using the Tukey method). All other pairs did not show any stimulation effect (Fig.
3C, right).
To further explore whether the stimulation effect could be explained by activity
rebalancing within the co-tuned network, we implemented a simple model in which a
suppression term was applied either to all non-target cells, randomly selected non-target cells,
or specifically to non-target co-tuned cells. By comparing three different model outcomes and
the real data, we observed a significant effect of the model type ($F(3, 3343) = 56.243$, $p <$
0.0001). Moreover, an interaction between the model type and cell groups was observed ($F(3,$
$3343) = 49.635$, $p < 0.0001$). Applying suppression to only non-target co-tuned cells during the
stimulation session yielded a significant response amplitude decrease for co-tuned cells
compared to non co-tuned cells ($F(1, 3343) = 48.68$, $p < 0.0001$), which resembles the real
data. In contrast, applying suppression to all non-target cells and random non-target cells led
to similar amplitude changes in both co-tuned and non co-tuned neurons ($F(1, 3343) = 0.01$, p
$= 0.925$ for all suppression; $F(1, 3343) = 0.05$, $p = 1$), which was not observed in either the real
data or the simulated data restricted to co-tuned cell suppression (Fig. 3D). These results
suggest that the target cell stimulation induces a selective activity suppression within the co-
tuned network for processing their preferred frequency.
Together, these results indicate that the effect of holographic optogenetic stimulation
depends not on the specific tuning of cells, but on the co-tuning between stimulated and non-
stimulated neurons. Also, this effect is not driven solely by a few non-target cells with large

response changes. Rather, the overall population of cells shows relative response changes due
to the stimulation when synchronized with their preferred frequency.
Overall, these results further suggest that when neural activity is increased in a subset
of target cells due to photostimulation in addition to the target sound presentation, other co-
tuned cells selectively reduce their tone-evoked responses to their preferred tone presentation,
indicating that the network rebalances to maintain network activity within a certain range.
295
Rebalanced network responses are stable
We then questioned whether such response amplitude changes due to stimulation within
the local network are persistent. To test whether the rebalanced status of the neuronal ensemble
is persistent, we examined the tone-evoked response amplitude changes between the post-
stimulation   and   the   stimulation   sessions   (post-stimulation   effect: $\Delta F/$
$F_{(post-stimulation\ session)} - \Delta F/F_{(stimulation\ session)}$). Response amplitudes were similar
across conditions and tone presentation frequencies for both groups of cells ($F(2, 4056) = 1.83$,
$p = 0.16$; Fig. 3E). These results indicate that pairing exogenous stimulation on a subset of
neurons along with sounds can instantaneously change the network responses to sounds, and
this change can persist at least for many minutes during the experimental session.
306
Neurons with higher frequency selectivity show greater response changes
Our results demonstrate that response changes on non-target cells are significantly
influenced by the frequency tuning of stimulation-target cells as well as the frequency of the
presented pure tone along with the stimulation. However, we also observed a marginal
stimulation effect in the 54 kHz non-target cell group during 54 kHz pure tone presentation,
even when 16 kHz target cells were stimulated. We reasoned that this effect might be due to
some weak sound activation of 54 kHz cells by 16 kHz tones potentially due to the asymmetric
shapes of many auditory tuning curves in AC [35,36]. Indeed, many cells exhibited broad tuning
properties, responding to both 16 kHz and 54 kHz (Fig. 3B). Thus, this marginal stimulation
effect could be attributed to cells grouped as 54 kHz preferring cells, yet still showing sound
evoked responses to 16 kHz, particularly given that 16 kHz is within the sensitive frequency
range in mice [30].
Building on our findings of a rebalanced cortical network, we next aimed to identify
whether frequency tuning selectivity influences response amplitude changes in the non-
targeted co-tuned neurons. For each cell, we calculated the frequency preference index
$(\Delta F/F_{(16kHz)}) - (\Delta F/F_{(54kHz)})$ and divided the cells into three categories of frequency
selectivity: low, mid, and high. We removed cells with extreme frequency preference values,
where the index values exceed ± 4 standard deviations from the median, prior to dividing them
into three categories. This removed about 1% of cells from the dataset for further analyses.
This grouping was based on the 33% quartile ranges, with each category representing one-third
of the data distribution (Fig. 3B). Values closer to 0 indicate more broadly tuned cells across
frequencies while extreme positive and negative values to indicate sharply tuned cells to either
frequency.
We then tested whether cells with higher frequency selectivity to one frequency
exhibited greater response amplitude changes. We performed a three-way ANOVA to examine
the effect of frequency selectivity (low, mid, high selectivity), stimulation condition (control,
16 kHz target stim, 54 kHz target stim), and cell groups (16 kHz vs. 54 kHz preferring cells)
on the response amplitude change. There were significant main effects of frequency selectivity
($F(2, 2183) = 23.52$, $p < 0.0001$) and stimulation condition ($F(2, 2183) = 11.03$, $p < 0.0001$).
No significant main effect of cell group was observed ($F(1, 2183) = 0.77$, $p = 0.379$). Thus,
neither the interaction between frequency selectivity and cell group ($F(2, 2183) = 0.69$, $p =$
0.503), nor the interaction between condition and cell group ($F(2, 2183) = 2.64$, $p = 0.072$) was

significant. The three-way interaction between frequency selectivity, stimulation condition,
and cell group was also not significant ($F(4, 2183) = 0.86$, $p = 0.487$).
However, the interaction between frequency selectivity and stimulation condition was
significant ($F(4, 2183) = 2.82$, $p = 0.0238$), indicating that the effect of frequency selectivity
depended on the condition. These results suggests that the response amplitude changes across
conditions were more prominent for cells with higher frequency selectivity.
To identify where the significant response difference occurred across conditions, we
further performed post-hoc pairwise comparisons between conditions within each frequency
selectivity category for each cell groups. For 16 kHz preferring cells, we observed a significant
difference in the response change between control and 16 kHz stim conditions only from the
high frequency selectivity category ($p < 0.027$, Holm-Bonferroni correction for multiple
comparisons). In parallel, for 54 kHz preferring cells, the significant effect was observed
between control and 54 kHz stim conditions ($p = 0.033$, Holm-Bonferroni correction for
multiple comparisons; Fig. 3F). These results indicate that non-targeted cells with higher
frequency selectivity exhibit the greatest response amplitude changes, only when the target
stimulated cells were co-tuned with them (Fig. S3). These results also suggest that frequency
selective neurons form co-tuned networks.
356
Sparsely distributed non-target co-tuned ensembles immediately rebalance their
activities to maintain the network balance
Network balance can be achieved by multiple mechanisms operating on different
timescales. To get insight into the potential mechanisms underlying the observed rebalancing,
we next investigated how rapidly cells start adjusting their responses during the stimulation
condition. We thus examined the stimulation effect (changes in response amplitude due to
stimulation of target cells) for non-target co-tuned ensembles at the single-trial level. We
observed decreased response amplitudes from the first trial, with no significant decay across
trials (Fig. 4A), regardless of cell groups, frequency presentation, and conditions (sum-of-
squares F-test, all $p > 0.05$). The absence of trial-related response amplitude changes in non-
target co-tuned ensembles indicates that non-target co-tuned cells immediately change their
activity whenever targeted cells increased their activity due to stimulation, to maintain the
network balance.
370
Non-target co-tuned ensembles that show rebalancing are spatially distributed
Activity rebalancing could be driven by local, e.g. changes in excitatory-inhibitory (E/I)
balance [37], or distributed changes. To identify whether co-tuned ensembles that changed their
activities are locally or widely distributed, we computed the center of mass distance between
each non-target cell to any of the target cells. For the stimulation condition, we excluded non-
target cells that were within 20 μm distance of the target cells to ensure that any effects from
those neighboring cells with their increased response amplitudes could have been not, even
partially, due to photostimulation (Fig. 1B-D). We observed that non-target co-tuned
ensembles were widely distributed within the FOV, similar to non-target non co-tuned
ensembles as well as those from the control condition (sum-of-squares F-test, $p > 0.05$; Fig.
4B). This indicates that activity changes of non-target co-tuned ensembles are not merely the
result of direct input from external photostimulation within a tight localized network. Rather,
widespread, sparsely represented co-tuned ensembles continuously update incoming
information based on their tuning properties.
385
Discussion
Trial-by-trial variability in neuronal activity is ubiquitous in the brain, with sensory
stimuli evoking activity in different sparse co-tuned ensembles at different times. How sensory-
evoked activity is distributed and coordinated across sparsely distributed co-tuned networks

has been unknown. Here, we leveraged the capability for selective *in vivo* stimulation via
holographic optogenetics to investigate how functionally related neuronal ensembles in AC
coordinate activity. Our results show that manipulating a small subset of target co-tuned
neurons alters the auditory-evoked responses of other non-target co-tuned neurons. Increased
activity in one subset of neurons is balanced by decreased activity in the rest of the co-tuned
population of neurons.
Importantly, such network rebalancing occurs only for processing acoustic features
specific to their tuning. Our analysis shows that the most selective non-targeted neurons show
the strongest effect after stimulation, suggesting that selective neurons form functionally
interacting sub-networks consistent with in vivo correlation analyses [38] and in vitro studies in
visual cortex [39]. Functionally related cells might form these subnetworks during development
likely due to lineage relationships and Hebbian processes [40-42]. Together, our findings suggest
that neuronal ensembles with strengthened connectivity across neuronal ensembles sharing
similar functional tuning properties actively interact and update their status to maintain the
overall network, enabling energy-efficient sensory processing.
The present work applied holographic optogenetic stimulation to manipulate neuronal
activity at a single-cell resolution in the AC for the first time. Similar to previous findings in
VC [18,19], our study further supports the idea that extra activation within the network exerts an
inhibitory influence on a subset of neurons. In the VC, single cell stimulation resulted in
suppression of neighboring co-tuned neurons [18]. Our results here show that feature-specific
suppression occurs in spatially dispersed ensembles of non-target co-tuned neurons. The
widely distributed response amplitude decreases in those neurons suggest that this phenomenon
is not limited to local neighboring cells but involved widespread networks [8]. Thus, the effect is
not solely due to inhibition caused by neighboring interneurons from the optogenetic
stimulation. Instead, neurons with similar functional characteristics, sparsely distributed
throughout the AC, actively interact, with more sharply tuned neurons to modulate their activity
the most.
Cortical networks are shaped by dynamic changes in neural activity driven by various
factors. Neurons rapidly modulate their responses based on their functional roles in sensory
processing. Recurrent cortical networks are thought to update their activity based on incoming
information to maintain homeostatic balance [23,24]. Concurrently, co-activated neurons
processing similar acoustic properties strengthen their connectivity by Hebbian learning [43-45].
Thus, cortical networks are shaped by an active interplay of synaptic plasticity, homeostasis
and Hebbian learning, rather than by a single dominant mechanism [46-49]. Rebalancing of
network activity is often attributed to homeostatic rebalancing of individual cell's activity [24,50]
or an E/I balance: increased activity of inhibitory neurons resulting in reduced activity of
excitatory neurons [37]. Our results suggest that rebalancing is tuning-specific.
Given that inhibitory neurons in A1 are generally less frequency selective than
excitatory neurons [51], changes in inhibition are unlikely to be the only contributor to the
observed effect. While our AAV was not cell-type specific, it is also unlikely that many
selected target neurons would be inhibitory interneurons as a greater proportion of neurons in
sensory cortices are excitatory (about 80% compared to 20% inhibitory [52-55]; AAV-hSyn is
expressed in both excitatory and inhibitory neurons in similar proportions [56]). Furthermore, our
primary cell selection criterion for stimulation yielded a subgroup of a strong specific
frequency-responsive cells (top 30% of cells that show the biggest evoked responses to the
target tone after excluding cells that show responses to multiple tones). This criterion likely
selected more excitatory neurons, as they generally show greater stimulus-selective responses
than inhibitory neurons in sensory cortices [51,57].
It is noteworthy that not all target neurons showed clear activation in response to
holographic stimulation. Although we attempted to pre-select cells responsive to stimulation,
some of them seemed to exhibit reduced activation during the experiment, potentially due to

motion artifacts, response adaptation, network suppression, or trial-by-trial variability.
Nevertheless, the frequency-specific suppression on co-tuned neurons observed in this study
suggests that this effect can be driven by activation of a very small number of neurons. One
caveat is that, while we presume most target cells were likely excitatory, inhibitory neurons
may have been included in the target cell group. Inhibitory neurons could show a reliable,
strong responses to the optogenetics compared to more variable responses that excitatory
neurons could show [58], thus may have been included in the target cell group. However, if more
inhibitory cells were included, this would have reduced trial-by-trial variability from the
stimulation yielding higher probability of target cell activation, which is different from what
we observed. Additionally, re-occurring sounds evoke activity changes in both excitatory and
inhibitory neurons in the same direction, leaving the E/I balance unchanged [59,60]. Moreover,
the fraction of inhibitory cells is 10 ~ 20% of all cortical neurons [61-63], thus the chance of
stimulating them is small. Together, these considerations suggest that our results are unlikely
the effect of the activation of inhibitory neurons. Rather, other balancing mechanisms such as
short-term depression at thalamocortical synapses may play a role [64].
In contrast to slow changes of homeostatic plasticity related to the E/I balance, plasticity
in cortical cellular responses can occur quickly and are cell-specific, thereby tuned to their
functional response properties [65-67]. Based on this, we speculate that the decrease in response
amplitudes of non-target neurons likely reflects rapid activity-dependent synaptic changes in
excitatory cells. Future work using Cre-dependent virus expression or a cell type specific
labeling approach will be required to confirm cell-type-specific roles in this phenomenon and
underlying mechanisms.
Lastly, no additional response changes were observed in the post-stimulation session
indicating that the rebalanced network status remained constant. Indeed, constant amplitudes
were observed regardless of conditions, subgroups, and frequencies, suggesting that the
network persistence was achieved through repeated acoustic stimuli presentation and
photostimulation. The persistence after the stimulation condition further suggests that once the
newly learned rebalanced network status is achieved, the network response stabilizes and
remains persistent. The mechanisms behind this stabilization are unclear but may involve an
active interplay of homeostasis and Hebbian learning to form co-active networks [68].
Taken together, the present study reveals how neuronal ensembles in the AC rebalance
to maintain a homeostatic processing equilibrium for a given sensory input, and that rebalanced
networks remain persistent. Moreover, our results show that network activity can be controlled
by even a small subset of neurons, and the network changes are closely tied to the functional
tuning properties of neurons.
476
Materials and Methods
Methods
Animals
A total of 14 mice over 8 weeks old (8 – 30 weeks, 6 males and 8 females) were used in the
experiments. To retain high-frequency hearing at experimental ages, offspring from C57BL6/J
and B6.CAST-*Cdh23*$^{Ahl+}$/Kjn (Jax 002756) were used for all experiments. Animals were
housed on 12-hr reversed light/dark cycle. All experimental procedures were approved by
Johns Hopkins Institutional Animal Care and Use Committee.
485
Preparation of AAV9-hSyn-GCaMP8s-T2A-rsChRmine
To generate the AAV9-hSyn-GCaMP8s-T2A-rsChRmine virus, a gene fragment containing
T2A, rsChRmine, Kv2.1 soma-targeted localization motif, and 3xHA tag (synthesized by Twist
Biosciences) was subcloned into the pGP-AAV-syn-jGCaMP8s plasmid vector (Addgene:
162374). Viral vectors were commercially prepared (Virovek) to a concentration of 1 x 10^{13}
vg/mL.

Surgery and virus injection

Surgery and virus injection
Surgery was performed as described in previous studies [9,10]. We injected dexamethasone
(1mg/kg, VetOne) subcutaneously (s.c.) to minimize brain swelling prior to surgery. 4%
isoflurane (Fluriso, VetOne) with a calibrated vaporizer (Matrix VIP 3000) was used to induce
anesthesia, which was then reduced down to 1.5 – 2% for maintenance. Body temperature was
monitored and maintained at around 36°C throughout the surgery (Harvard Apparatus
Homeothermic monitor). We first removed the hair on the head using a hair removal product
(Nair) to expose the skin. Betadine and 70% ethanol were applied three times to the exposed
skin. Skin and tissues were then removed, and muscles were scraped to expose the left temporal
side where the craniotomy was conducted. Unilateral craniotomy was performed to expose
about 3.5 mm diameter region over the left AC. Virus (AAV9-hSyn-GCaMP8s-TSA-
rsChRmine titer of 1:2) injection was performed at 2-3 sites near tentative A1 area at about 300
μm depth from the surface, using a glass pipette controlled by a micromanipulator (Sutter
Instrument MPC-200 and ROE-200 controller). We injected 300 nL on each site at the rate of
180 nL/min (Nanoject3). Once virus injection was completed, two circular glass coverslips
(one of 3 mm and one of 4 mm in diameter) were affixed with a clear silicone elastomer (Kwik-
Sil, World Precision Instruments). An extra layer of dental acrylic (C&B Metabond) was
applied around the edge of the cranial window to further secure it, cover the exposed skull, and
adhere a custom 3D-printed stainless steel headpost. Carprofen (5 mg/kg) and cefazolin
(300mg/kg) were injected (s.c.) post-operatively. Animals were given at least 3-4 weeks of
recovery and viral expression time before any imaging was performed.

Experimental procedures
Fully awake animals were head-fixed on a custom-made stage, where a free field
speaker (TDT ED1) was faced towards the right ear at 45 degrees. All sound stimuli were
driven by TDT RX6 multiprocessor and we imaged GCaMP8s responses with a resonant
scanning two-photon microscope (Bruker Ultima 2Pplus; 940 nm excitation wavelength) at A1
L2/3 (160 – 200 μm). A1 was identified by its tonotopy gradient using widefield imaging (Fig.
1E) [69]. During 2-photon imaging sessions, we first conducted a short imaging session (about 1
min.) presenting 100-ms pure tones at 3 different frequencies (4, 16, 54 kHz) at 70 dB SPL,
covering the mouse hearing range, for 10 times each in a randomized order (ISI: 2 sec). This
was to identify initial tuning properties of sound-responsive cells (cell selection session; Fig.
1E). The acquired imaging data was immediately processed using 'suite2p' and a custom-
written Matlab script to identify tone-responsive cells for each frequency. We continued only
when at least 50% of cells within the FOV showed sound-evoked responses. We took 16 kHz
or 54 kHz as our target functional properties. Target frequency was randomly assigned. We
manually tested the response changes to stimulation of the top 30% of target frequency
responsive cells (~ 20 cells) selected from the cell selection session by using a stimulation laser
(Light Conversion Carbide; 1040 nm excitation wavelength). Stimulation laser power was set
around 5 mW per cell. We selected 5 representative cells (target cells) that showed visible
fluorescence changes to the stimulation.
Prior to the main experimental session, we ran a short (~ 1 min.) stimulation session
without any sound presentation that comprised 5 trials of 100-ms stimulation with 6-second
ISIs for a rapid check of the stimulation effect. This session was restricted to 5 trials of
stimulation given the limited time of the imaging session, leading to larger variability of the
observed stimulation effect. We further verified the stimulation effect from the experimental
stimulation session where 30 trials were given.
For the main experiment, three consecutive imaging sessions were followed by the
presentation of either 16 kHz or 54 kHz 100 ms pure tones with random ISIs between 5.8 – 6.5
sec. for 30 trials each, as baseline session, stimulation session, and post-stimulation session

(Fig. 1E). Only the second imaging session (i.e., stimulation session) received holographic
stimulation on the pre-selected 5 target cells for 30 revolutions of 15 μm spiral for about 100
ms at 5 mW laser power per cell (16 kHz target cell or 54 kHz target cell stimulation conditions)
or 0 mW laser power per cell (control condition). The default mechanical setup (Bruker
PrairieView version 5.7) of the microscope opens and closes the uncaging shutter to enable the
stimulation laser path for every single stimulation time point, which causes an external
mechanic sound that can trigger neural activation of the AC. To minimize any effect of external
mechanic sounds to cells in the AC, we kept the uncaging shutter open during the imaging
session. Number of imaging sessions per animal varied depending on the virus expression.
Regardless, all animals were used for control and at least one stimulation condition with
minimum 2 days apart by varying FOVs to avoid imaging the same cells multiple times. The
order of conditions and the imaging depth presented to the same animal were randomized.
555   An additional single-cell stimulation only session was conducted on a subset of animals,
to further test a reliable holographic stimulation at a single-cell level (n = 3 animals). We varied
a stimulation position from the original target stimulation point to 10, 20, or 30 μm shifted
along x-axis or y-axis, generating 7 different stimulation point (original cell position, 10, 20,
or 30 μm shifted along the x-axis, 10, 20, or 30 μm shifted along the y-axis). We stimulated
each stimulation point for 10 times in a randomized order across stimulation points with 8-sec.
ISIs.
562
Analysis
Imaging data were processed with 'suite2p' for motion correction, cell detection, and cell
fluorescence trace extraction [70]. We applied neuropil correction to the fluorescence traces using
the following equation: $F_{(cell\_corrected)} = F_{(cell)} - (0.8 * F_{(neuropil)})$. We then computed ΔF/F
normalized to the baseline, by following the equation: $\Delta F/F = (F_{(trace)} - F_{(baseline)}) / F_{(baseline)}$,
where baseline is about 300 ms before the sound onset. For the single-cell level stimulation
session, we computed peak ΔF/F per each stimulation point and applied a mixed-effect model
by taking peak ΔF/F as dependent variables, stimulation point as independent variables, and
cells as random factor. For the 5-cell stimulation validation session, we computed a proportion
of activated cells (any cell with peak ΔF/F > 0) among stimulated cells per each FOV as well
as peak ΔF/F and ran a random permutation with 100 iterations.
574   For experimental sessions, to select sound responsive cells, we compared sound-evoked
activity (160 ms – 660 ms after sound onset capturing the sound-evoked response due to slow
calcium transient) and the baseline activity (300 ms – 0 ms before the sound onset). We
considered cells sound responsive when the amplitude of the average sound-evoked activity
exceeded two standard deviations of the amplitude of the average baseline activity. Sound
responsive cells were selected based on fluorescence traces only from the baseline session to
minimize any potential effect of stimulation on cell selection. We then computed the ratio of
the evoked activity between two frequencies as an index of the frequency preference
$(\Delta F/F_{(16kHz)} - \Delta F/F_{(54kHz)})$. We subgrouped sound-responsive cells into either 16 kHz
preferring cells or 54 kHz preferring cells based on the frequency preference, taking 0 as a
subgroup criterion. As our main interest was changes to non-target cells, we excluded target
cells for further analyses. To compare response changes due to stimulation for each group of
cells, average sound-evoked activity of sound-responsive cells from the baseline session was
subtracted from the stimulation session (stimulation effect: $\Delta F/F_{(stimulation\ session)} - \Delta F/$
$F_{(baseline\ session)}$) for each condition. We further compared response changes between post-
stimulation session and stimulation session, again by subtracting the response amplitudes
between two sessions for each group and condition (post-stimulation effect: $\Delta F/$
$F_{(post-stimulation\ session)} - \Delta F/F_{(stimulation\ session)}$). To quantify the stimulation effect
based on functional properties for conditions and groups, we applied a mixed-effect model by
taking amplitude changes as dependent variables, frequency, conditions, and cell groups as

independent variables, and FOVs as random factor. We then computed Type III analysis of
variance (ANOVA) to examine whether the effect of functional property (frequency) to the
amplitude changes was specific to the target frequency presentation synchronized with the
stimulation (i.e., 16 kHz for 16 kHz target cell stimulation or 54 kHz for 54 kHz target cell
stimulation). To quantify whether the response amplitude changes due to stimulation differ
across trials, we fitted a dataset of average stimulation effect across each trial per each condition
(stimulation or control), each cell group (16 kHz preferring or 54 kHz preferring cells), and
each tone presentation (16 kHz or 54 kHz) to the three-parameter model and computed the
extra sum-of-squares F-test to compare whether response amplitude changes across trials were
different from a constant line [71]. To quantify a relationship between response amplitude
changes of non-target cells and their distances to target cells, we computed a center of mass
distance of each cell position relative to target cells, and fitted the dataset of the response
amplitude changes across distance to the three-parameter model to compute the extra sum-of-
squares F-test, per each condition (16 kHz target stimulation, 54 kHz target stimulation, or
control), each cell group (16 kHz preferring or 54 kHz preferring cells), and each tone
presentation (16 kHz or 54 kHz). For the control condition, as there were no stimulated target
cells, we chose top five most tone-responsive cells from the baseline session as "target" cells.
We then generated a simple model in which a suppression term was applied either to
all neurons or specifically to non-target co-tuned cells to test our results from the data. We took
a similar range of number of neurons and FOVs to closely simulate the model to the real dataset
structure. On 50 neurons (n) per FOV across 18 FOVs, the simulated calcium trace of each
neuron was defined as

$$\text{Trace}_{n(t)} = R_{n(t)} - \text{theta}_n + \text{epsilon}_{n(t)}$$

where $R_{n(t)}$ is a time-varying response amplitude from the sound onset to the offset,
modeled separately for the baseline and stimulation sessions. The suppression term theta_n was
applied only during the stimulation session either to all neurons, randomly selected neurons, or
only non-target co-tuned neurons depending on the simulation condition, and $\text{epsilon}_{n(t)}$ is
additive Gaussian noise. To simulate sound-evoked calcium transients, we assigned a faster
decay time constant (200 ms) for non-target co-tuned neurons $R_{n(t)}$ and a slower decay (1000
ms) for non-target non co-tuned neurons $R_{n(t)}$ for both the baseline and stimulation sessions.
Theta was defined as proportional to the average stimulation strength from target neurons,
derived from the real dataset, and scaled by a factor $\alpha = 0.3$ in the current simulation. To
introduce neuron-level variability, an additional jitter (epsilon_n) was applied as follow:

$$\text{Theta}_n = \alpha * \text{mean(target stimulation amplitude)} * (1 + \text{epsilon}_n)$$

Similar to the real data analyses, we compared the response change between the
stimulation and baseline sessions' trace amplitudes.

Histology
Animals were deeply anesthetized with 4% isoflurane to perform transcardial perfusion with
4% paraformaldehyde (PFA) in 0.1 M phosphate buffer saline (PBS). The extracted brains
were post-fixed in 4% PFA for additional 12-24 hours. Coronal brain sections at 50 μm
containing the AC were stained with primary antibodies of HA-Tag (1:500) and chicken Green
Fluorescent Protein (GFP, 1:500) for GCaMP8s, and secondary antibodies of 594-conjugated
anti-rabbit IgG (1:1000) and 488-conjugated anti-chicken IgG (1:1000) for red-shifted opsins.

References

1. Read, H., Winer, J., and Schreiner, C. (2001). Modular organization of intrinsic connections associated with spectral tuning in cat auditory cortex. Proceedings of the National Academy of Sciences of the United States of America *98*, 8042-8047. 10.1073/pnas.131591898.

2. Schnupp, J., Mrsic-Flogel, T., and King, A. (2001). Linear processing of spatial cues in primary auditory cortex. Nature *414*, 200-204. 10.1038/35102568.

3. Lu, K., Liu, W., Zan, P., David, S.V., Fritz, J.B., and Shamma, S.A. (2018). Implicit Memory for Complex Sounds in Higher Auditory Cortex of the Ferret. J Neurosci *38*, 9955-9966. 10.1523/JNEUROSCI.2118-18.2018.

4. Fritz, J., Shamma, S., Elhilali, M., and Klein, D. (2003). Rapid task-related plasticity of spectrotemporal receptive fields in primary auditory cortex. Nature Neuroscience *6*, 1216-1223. 10.1038/nn1141.

5. Bandyopadhyay, S., Shamma, S.A., and Kanold, P.O. (2010). Dichotomy of functional organization in the mouse auditory cortex. Nat Neurosci *13*, 361-368. 10.1038/nn.2490.

6. Rothschild, G., Nelken, I., and Mizrahi, A. (2010). Functional organization and population dynamics in the mouse primary auditory cortex. Nature Neuroscience *13*, 353-U321. 10.1038/nn.2484.

7. Panzeri, S., Macke, J., Gross, J., and Kayser, C. (2015). Neural population coding: combining insights from microscopic and mass signals. Trends in Cognitive Sciences *19*, 162-172. 10.1016/j.tics.2015.01.002.

8. Bowen, Z., Winkowski, D., and Kanold, P. (2020). Functional organization of mouse primary auditory cortex in adult C57BL/6 and F1 (CBAxC57) mice. Scientific Reports *10*, ARTN 10905. 10.1038/s41598-020-67819-4.

9. Francis, N., Winkowski, D., Sheikhattar, A., Armengol, K., Babadi, B., and Kanold, P. (2018). Small Networks Encode Decision-Making in Primary Auditory Cortex. Neuron *97*, 885-+. 10.1016/j.neuron.2018.01.019.

10. Francis, N., Mukherjee, S., Kocillari, L., Panzeri, S., Babadi, B., and Kanold, P. (2022). Sequential transmission of task-relevant information in cortical neuronal networks. Cell Reports *39*, ARTN 110878. 10.1016/j.celrep.2022.110878.

11. Hromádka, T., Deweese, M.R., and Zador, A.M. (2008). Sparse representation of sounds in the unanesthetized auditory cortex. PLoS Biol *6*, e16. 10.1371/journal.pbio.0060016.

12. Hromadka, T., and Zador, A. (2009). Representations in auditory cortex. Current Opinion in Neurobiology *19*, 430-433. 10.1016/j.conb.2009.07.009.

13. Liang, F., Li, H., Chou, X., Zhou, M., Zhang, N., Xiao, Z., Zhang, K., Tao, H., and Zhang, L. (2019). Sparse Representation in Awake Auditory Cortex: Cell-type Dependence, Synaptic Mechanisms, Developmental Emergence, and Modulation. Cerebral Cortex *29*, 3796-3812. 10.1093/cercor/bhy260.

14. Yu, L., and Yu, Y. (2017). Energy-Efficient Neural Information Processing in Individual Neurons and Neuronal Networks. Journal of Neuroscience Research *95*, 2253-2266. 10.1002/jnr.24131.

15. Jadi, M., and Sejnowski, T. (2014). Cortical oscillations arise from contextual interactions that regulate sparse coding. Proceedings of the National Academy of Sciences of the United States of America *111*, 6780-6785. 10.1073/pnas.1405300111.

16. Grabner, R., Neubauer, A., and Stern, E. (2006). Superior performance and neural efficiency: The impact of intelligence and expertise. Brain Research Bulletin *69*, 422-439. 10.1016/j.brainresbull.2006.02.009.

17. Bowen, Z., Winkowski, D.E., Seshadri, S., Plenz, D., and Kanold, P.O. (2019). Neuronal Avalanches in Input and Associative Layers of Auditory Cortex. Front Syst Neurosci *13*, 45. 10.3389/fnsys.2019.00045.

18. Chettih, S.N., and Harvey, C.D. (2019). Single-neuron perturbations reveal feature-specific competition in V1. Nature *567*, 334-340. 10.1038/s41586-019-0997-6.

19. LaFosse, P.K., Zhou, Z., O'Rawe, J.F., Friedman, N.G., Scott, V.M., Deng, Y., and Histed, M.H. (2024). Single-cell optogenetics reveals attenuation-by-suppression in visual cortical neurons. bioRxiv. 10.1101/2023.09.13.557650.

20. Winkowski, D.E., and Kanold, P.O. (2013). Laminar transformation of frequency organization in auditory cortex. J Neurosci *33*, 1498-1508. 10.1523/JNEUROSCI.3101-12.2013.

21. Adesnik, H., and Abdeladim, L. (2021). Probing neural codes with two-photon holographic optogenetics. Nat Neurosci *24*, 1356-1366. 10.1038/s41593-021-00902-9.

22. Yang, W., and Yuste, R. (2021). Holographic Imaging and Stimulation of Neural Circuits. Optogenetics: Light-Sensing Proteins and Their Applications in Neuroscience and Beyond, 613-639.

23. Karmarkar, U., and Buonomano, D. (2007). Timing in the absence of clocks: Encoding time in neural network states. Neuron *53*, 427-438. 10.1016/j.neuron.2007.01.006.

24. Liu, B., Seay, M., and Buonomano, D. (2023). Creation of Neuronal Ensembles and Cell-Specific Homeostatic Plasticity through Chronic Sparse Optogenetic Stimulation. Journal of Neuroscience *43*, 82-92. 10.1523/JNEUROSCI.1104-22.2022.

25. Kishi, K.E., Kim, Y.S., Fukuda, M., Inoue, M., Kusakizako, T., Wang, P.Y., Ramakrishnan, C., Byrne, E.F.X., Thadhani, E., Paggi, J.M., et al. (2022). Structural basis for channel conduction in the pump-like channelrhodopsin ChRmine. Cell *185*, 672-689.e623. 10.1016/j.cell.2022.01.007.

26. LaFosse, P., Zhou, Z., Friedman, N., Deng, Y., Li, A., Akitake, B., and Histed, M. (2023). Bicistronic Expression of a High-Performance Calcium Indicator and Opsin for All-Optical Stimulation and Imaging at Cellular Resolution. Eneuro *10*. 10.1523/ENEURO.0378-22.2023.

27. Sridharan, S., Gajowa, M., Ogando, M., Jagadisan, U., Abdeladim, L., Sadahiro, M., Bounds, H., Hendricks, W., Turney, T., Tayler, I., et al. (2022). High-performance microbial opsins for spatially and temporally precise perturbations of large neuronal networks. Neuron *110*, 1139-+. 10.1016/j.neuron.2022.01.008.

28. Mardinly, A., Oldenburg, I., Pegard, N., Sridharan, S., Lyall, E., Chesnov, K., Brohawn, S., Waller, L., and Adesnik, H. (2018). Precise multimodal optical control of neural ensemble activity. Nature Neuroscience *21*, 881-+. 10.1038/s41593-018-0139-8.

29. Daie, K., Svoboda, K., and Druckmann, S. (2021). Targeted photostimulation uncovers circuit motifs supporting short-term memory. Nature Neuroscience *24*, 259-265. 10.1038/s41593-020-00776-3.

30. Kane, K.L., Longo-Guess, C.M., Gagnon, L.H., Ding, D., Salvi, R.J., and Johnson, K.R. (2012). Genetic background effects on age-related hearing loss associated with Cdh23 variants in mice. Hear Res *283*, 80-88. 10.1016/j.heares.2011.11.007.

31. Branchi, I., Santucci, D., and Alleva, E. (2001). Ultrasonic vocalisation emitted by infant rodents: a tool for assessment of neurobehavioural development. Behav Brain Res *125*, 49-56. 10.1016/s0166-4328(01)00277-7.

32. Ulanovsky, N., Las, L., Farkas, D., and Nelken, I. (2004). Multiple time scales of adaptation in auditory cortex neurons. J Neurosci *24*, 10440-10453. 10.1523/JNEUROSCI.1905-04.2004.

739   33.   Yarden, T.S., and Nelken, I. (2017). Stimulus-specific adaptation in a recurrent network
740   model of primary auditory cortex. PLoS Comput Biol *13*, e1005437.
741   10.1371/journal.pcbi.1005437.
742   34.   Malmierca, M.S., Sanchez-Vives, M.V., Escera, C., and Bendixen, A. (2014). Neuronal
743   adaptation, novelty detection and regularity encoding in audition. Front Syst Neurosci
744   *8*, 111. 10.3389/fnsys.2014.00111.
745   35.   Schreiner, C.E., Read, H.L., and Sutter, M.L. (2000). Modular organization of frequency
746   integration in primary auditory cortex. Annu Rev Neurosci *23*, 501-529.
747   10.1146/annurev.neuro.23.1.501.
748   36.   Liu, J., and Kanold, P.O. (2021). Diversity of Receptive Fields and Sideband Inhibition
749   with Complex Thalamocortical and Intracortical Origin in L2/3 of Mouse Primary
750   Auditory Cortex. J Neurosci *41*, 3142-3162. 10.1523/JNEUROSCI.1732-20.2021.
751   37.   Froemke, R., and Hyman, S. (2015). Plasticity of Cortical Excitatory-Inhibitory Balance.
752   Annual Review of Neuroscience, Vol 38 *38*, 195-219. 10.1146/annurev-neuro-071714-
753   034002.
754   38.   Rupasinghe, A., Francis, N., Liu, J., Bowen, Z., Kanold, P., and Babadi, B. (2021). Direct
755   extraction of signal and noise correlations from two-photon calcium imaging of
756   ensemble neuronal activity. Elife *10*, ARTN e68046. 10.7554/eLife.68046.
757   39.   Ko, H., Mrsic-Flogel, T., and Hofer, S. (2014). Emergence of Feature-Specific
758   Connectivity in Cortical Microcircuits in the Absence of Visual Experience. Journal of
759   Neuroscience *34*, 9812-9816. 10.1523/JNEUROSCI.0875-14.2014.
760   40.   Katz, L., and Shatz, C. (1996). Synaptic activity and the construction of cortical circuits.
761   Science *274*, 1133-1138. 10.1126/science.274.5290.1133.
762   41.   Ohtsuki, G., Nishiyama, M., Yoshida, T., Murakami, T., Histed, M., Lois, C., and Ohki, K.
763   (2012). Similarity of visual selectivity among clonally related neurons in visual cortex.
764   Neuron *75*, 65-72. 10.1016/j.neuron.2012.05.023.
765   42.   Li, Y., Lu, H., Cheng, P.L., Ge, S., Xu, H., Shi, S.H., and Dan, Y. (2012). Clonally related
766   visual cortical neurons show similar stimulus feature selectivity. Nature *486*, 118-121.
767   10.1038/nature11110.
768   43.   Hebb, D.O. (2005). The organization of behavior: A neuropsychological theory
769   (Psychology press).
770   44.   Malenka, R.C., and Nicoll, R.A. (1999). Long-term potentiation--a decade of progress?
771   Science *285*, 1870-1874. 10.1126/science.285.5435.1870.
772   45.   Brown, T.H., Kairiss, E.W., and Keenan, C.L. (1990). Hebbian synapses: biophysical
773   mechanisms and algorithms. Annual review of neuroscience *13*, 475-511.
774   46.   Vitureira, N., and Goda, Y. (2013). The interplay between Hebbian and homeostatic
775   synaptic plasticity. Journal of Cell Biology *203*, 175-186. 10.1083/jcb.201306030.
776   47.   Fox, K., and Stryker, M. (2017). Integrating Hebbian and homeostatic plasticity:
777   introduction. Philosophical Transactions of the Royal Society B-Biological Sciences *372*,
778   ARTN 20160413. 10.1098/rstb.2016.0413.
779   48.   Toyoizumi, T., Kaneko, M., Stryker, M., and Miller, K. (2014). Modeling the Dynamic
780   Interaction of Hebbian and Homeostatic Plasticity. Neuron *84*, 497-510.
781   10.1016/j.neuron.2014.09.036.
782   49.   Turrigiano, G., and Nelson, S. (2000). Hebb and homeostasis in neuronal plasticity.
783   Current Opinion in Neurobiology *10*, 358-364. 10.1016/S0959-4388(00)00091-X.
784   50.   Yin, J., and Yuan, Q. (2015). Structural homeostasis in the nervous system: a balancing
785   act for wiring plasticity and stability. Frontiers in Cellular Neuroscience *8*, ARTN 439.
786   10.3389/fncel.2014.00439.

51. Maor, I., Shalev, A., and Mizrahi, A. (2016). Distinct Spatiotemporal Response
Properties of Excitatory Versus Inhibitory Neurons in the Mouse Auditory Cortex.
Cerebral Cortex *26*, 4242-4252. 10.1093/cercor/bhw266.
52. Wang, Y., Ye, M., Kuang, X., Li, Y., and Hu, S. (2018). A simplified morphological
classification scheme for pyramidal cells in six layers of primary somatosensory cortex
of juvenile rats. Ibro Reports *5*, 74-90. 10.1016/j.ibror.2018.10.001.
53. Marin, O. (2012). Interneuron dysfunction in psychiatric disorders. Nature Reviews
Neuroscience *13*, 107-120. 10.1038/nrn3155.
54. Xu, J., Fan, J., Wang, X., Eacker, S., Kam, T., Chen, L., Yin, X., Zhu, J., Chi, Z., Jiang, H., et
al. (2016). Cultured networks of excitatory projection neurons and inhibitory
interneurons for studying human cortical neurotoxicity. Science Translational Medicine
8, ARTN 333ra48. 10.1126/scitranslmed.aad0623.
55. Wonders, C., and Anderson, S. (2006). The origin and specification of cortical
interneurons. Nature Reviews Neuroscience *7*, 687-696. 10.1038/nrn1954.
56. Nathanson, J., Yanagawa, Y., Obata, K., and Callaway, E. (2009). Preferential Labeling of
Inhibitory and Excitatory Cortical Neurons by Endogenous Tropism of Adeno-
associated Virus and Lentivirus Vectors. Neuroscience *161*, 441-450.
10.1016/j.neuroscience.2009.03.032.
57. Kerlin, A.M., Andermann, M.L., Berezovskii, V.K., and Reid, R.C. (2010). Broadly tuned
response properties of diverse inhibitory neuron subtypes in mouse visual cortex.
Neuron *67*, 858-871. 10.1016/j.neuron.2010.08.002.
58. Mateo, C., Avermann, M., Gentet, L.J., Zhang, F., Deisseroth, K., and Petersen, C.C.
(2011). In vivo optogenetic stimulation of neocortical excitatory neurons drives brain-
state-dependent inhibition. Curr Biol *21*, 1593-1602. 10.1016/j.cub.2011.08.028.
59. Wehr, M., and Zador, A.M. (2003). Balanced inhibition underlies tuning and sharpens
spike timing in auditory cortex. Nature *426*, 442-446.
60. Chen, I.W., Helmchen, F., and Lütcke, H. (2015). Specific early and late oddball-evoked
responses in excitatory and inhibitory neurons of mouse auditory cortex. Journal of
Neuroscience *35*, 12560-12573.
61. Meyer, H., Schwarz, D., Wimmer, V., Schmitt, A., Kerr, J., Sakmann, B., and Helmstaedter,
M. (2011). Inhibitory interneurons in a cortical column form hot zones of inhibition in
layers 2 and 5A. Proceedings of the National Academy of Sciences of the United States
of America *108*, 16807-16812. 10.1073/pnas.1113648108.
62. Tasaka, G.I., Maggi, C., Taha, E., and Mizrahi, A. (2023). The local and long-range input
landscape of inhibitory neurons in mouse auditory cortex. J Comp Neurol *531*, 502-
514. 10.1002/cne.25437.
63. Markram, H., Toledo-Rodriguez, M., Wang, Y., Gupta, A., Silberberg, G., and Wu, C.
(2004). Interneurons of the neocortical inhibitory system. Nat Rev Neurosci *5*, 793-807.
10.1038/nrn1519.
64. Li, J.Y., and Glickfeld, L.L. (2023). Input-specific synaptic depression shapes temporal
integration in mouse visual cortex. Neuron. 10.1016/j.neuron.2023.07.003.
65. Davis, G. (2013). Homeostatic Signaling and the Stabilization of Neural Function.
Neuron *80*, 718-728. 10.1016/j.neuron.2013.09.044.
66. Tien, N., Soto, F., and Kerschensteiner, D. (2017). Homeostatic Plasticity Shapes Cell-
Type-Specific Wiring in the Retina. Neuron *94*, 656-+. 10.1016/j.neuron.2017.04.016.
67. Maffei, A., and Fontanini, A. (2009). Network homeostasis: a matter of coordination.
Current Opinion in Neurobiology *19*, 168-173. 10.1016/j.conb.2009.05.012.

68. Barak, O., and Tsodyks, M. (2007). Persistent activity in neural networks with dynamic
synapses.   Plos   Computational   Biology   *3*,   323-332,   ARTN   e35.
10.1371/journal.pcbi.0030035.
69. Liu, J., Whiteway, M.R., Sheikhattar, A., Butts, D.A., Babadi, B., and Kanold, P.O. (2019).
Parallel Processing of Sound Dynamics across Mouse Auditory Cortex via Spatially
Patterned Thalamic Inputs and Distinct Areal Intracortical Circuits. Cell Rep *27*, 872-
885.e877. 10.1016/j.celrep.2019.03.069.
70. Pachitariu, M., Stringer, C., Schröder, S., Dipoppa, M., Rossi, L.F., Carandini, M., and
Harris, K.D. (2016). Suite2p: beyond 10,000 neurons with standard two-photon
microscopy. BioRxiv, 061507.
71. Motulsky, H., and Christopoulos, A. (2004). Fitting models to biological data using
linear and nonlinear regression: a practical guide to curve fitting (Oxford University
Press).
847
848

849 **Acknowledgments:** Supported by U19 NS107464 (POK), NIH RO1DC017785 (POK), NIH
850 F32DC019842 (TAB).
851
852 **Author Contributions:** Conceptualization: HK, TAB, POK; Methodology: HK, TAB, POK;
853 Investigation: HK, TAB; Writing: HK, POK
854
855 **Competing interest:** Authors declare that they have no competing interests.
856
857 **Data and materials availability:** All preprocessed imaging data and relevant analyses
858 scripts will be deposited at Johns Hopkins University research data repository, available for
859 open access.
860
861

Figures

Figure 1. Holographic optogenetic stimulation in AC and experimental procedure. A: An
example brain slice showing cells in AC expressing both GCaMP and opsin (AAV9-hSyn-
GC8s-T2A-rsChRmine). **B**: An example field of view (FOV) where single cell targeting
precision was tested and response traces to the holographic stimulation from an example cell.
Stimulation was offset from the original position (red circle) to distance-shifted positions in 10
μm increments (gray dashed circles in the x-axis or y-axis of the FOV). Responses were the
greatest when the stimulation was performed on the original cell position (red solid line trace).
Rapid amplitude decay along the position shift was observed (red dashed line traces). Grey
error shades indicate SEM across trials. A right inset errorbar plot shows a grand average
amplitude change per stimulation point across all tested cells (n = 15 cells, 3 animals). Error
bars indicate SEM across cells. **C**: An example FOV showing a population of cells (left) and
amplitude changes to 5-cell stimulation as a stimulation effect ($\Delta F/F_{stim}$ - $\Delta F/F_{spont., right}$). Filled
squares indicate each cell. White circles indicate stimulation targeted cells. **D**: (left) Proportion
of stimulated cells that showed an increase in fluorescence following photostimulation. Error
bars indicate SEM across FOVs. A horizontal dashed line indicates average permutation results

(random permutation test on 100 iterations, $p < 0.0001$). (right) Grand average of the
stimulation effect across imaging sessions. Error bars indicate SEM across FOVs. A horizontal
dashed line indicates average permutation results (random permutation test on 100 iterations,
$p < 0.0001$). **E**: Experimental procedure. A total of four consecutive imaging sessions were
acquired: 1) A cell selection session to identify neurons selective for 16 kHz pure tones, 2) a
baseline imaging session to acquire tone-evoked activity response to either 16 or 54 kHz pure
tone, 3) a stimulation session representing five cells of either 16 kHz or 54 kHz responsive
cells as target stimulation to examine the effect of stimulation synchronized to tone
presentations, and 4) a post-stimulation session to examine network persistence after
stimulation-related changes.
890

891
Figure 2. Targeted cells and non-target cells show response changes due to stimulation.
A: (top) An example FOV showing the stimulation effect (Δ F/F$_{stim}$ - Δ F/F$_{baseline}$) of sound
responsive cells for 16 kHz target cell stimulation (filled squares). Black circles indicate
stimulated target cells. (bottom) Mean response traces of an example target and non-target cell
in baseline session (black) and stimulation session (red). Error bars indicate SEM across trials.
An example target cell shows an increased response due to the stimulation. An example non-
target cell shows a decreased response due to stimulation on the target cells. **B**: (left) A violin
plot of the proportion of stimulated cells that showed increased activity due to stimulation
across FOVs. Horizontal solid line indicates mean proportion, empty circle indicates median
proportion, and gray filled circles indicate individual FOVs. (right) Mean amplitude changes
of target cells for stimulation session and post-stimulation session normalized to the baseline
session. Error bars indicate SEM across cells. Dashed horizontal lines on both panels indicate
average permutation results (random permutation test on 100 iterations, *p* < 0.0001). **CD**: Same
as ABC for 54 kHz target cell stimulation.
906
907

Figure 3. Non-target co-tuned cells show more decreased response amplitudes due to stimulation when synchronized with their preferred tones. A: Stimulation effect ($\Delta F/F_{stim}$ - $\Delta F/F_{baseline}$) in all sound responsive cells, including both target and non-target cells, responding to either 16 kHz (blue) or 54 kHz (orange) pure tones, representing global activity changes due to the stimulation. No significant differences between stimulation conditions and responses to different frequencies were observed (all $p > 0.05$). **B**: Sub-categorization of cells based on the frequency selectivity for each target stimulation condition (left: 16 kHz stim, right 54 kHz stim). Cells were first grouped into either 16 kHz preferring cells (blue) or 54 kHz preferring cells (orange). Within each cell group, cells were further subdivided into low, mid, and high frequency selectivity categories based on their 33% quartile ranges. For visualization, frequency preference was log-transformed; original frequency selectivity distributions are shown in the upper insets. Vertical dashed lines indicate 33% quartile ranges. **C**: Stimulation effect ($\Delta F/F_{stim}$ - $\Delta F/F_{baseline}$) in 16 kHz (blue) and 54 kHz (orange) preferring cells. Both cell groups show decreased amplitude to their preferred frequency regardless of conditions due to acoustic stimulus-specific adaptation. Only co-tuned cells (16 kHz preferring cells for 16 kHz stimulation or 54 kHz preferring cells for 54 kHz stimulation) show a further decrease in response amplitudes due to the stimulation, when the preferred pure tone (PT) frequency was synchronized. Error bars indicate SEM across FOVs (*: $p < 0.0001$). **D**: Stimulation effect from the model prediction. Amplitude changes computed from simulated data by applying cell suppression to all cells (All supp.), random cells (Random supp.), or only co-tuned cells (Co-tuned supp.) were compared with real data. Only the Co-tuned supp. model showed a

significant amplitude decrease for co-tuned neurons compared to non co-tuned neurons, similar
to the result from the real data ($p < 0.05$; see texts for more detail). **E**: Post-stimulation effect
(Δ F/F$_{post}$ - Δ F/F$_{stim}$) 16 kHz (blue) and 54 kHz (orange) preferring cells. No significant
response amplitude changes were observed. Error bars indicate SEM across FOVs. **F**:
Response amplitude change based on the frequency selectivity category for each cell groups
(blue: 16 kHz preferring cells, orange: 54 kHz preferring cells). Significant response amplitude
changes relative to the control condition were observed only for high frequency selectivity
category when target stimulated cells were co-tuned (*: $p < 0.05$).

Figure 4. Rebalanced response changes on non-target 16 kHz cells are immediate and widely distributed. A: Stimulation effect ($\Delta F/F_{stim} - \Delta F/F_{baseline}$) in 16 kHz (blue) and 54 kHz (orange) preferring cells per each trial for the 5-cell 16 kHz stimulation condition (left), 54 kHz stimulation condition (middle), and the no-cell control condition (right). Each circle represents average stimulation effect per each trial. Decreased amplitudes to preferred frequencies were observed from as early as trial 1 with no significant further changes across trials, regardless of frequencies and conditions (sum-of-squares F-test, all $p > 0.05$). Solid lines indicate fitted curves and dashed lines indicate 95% confidence intervals. Vertical gray dashed lines indicate 0 amplitude change. **B**: Stimulation effect ($\Delta F/F_{stim} - \Delta F/F_{baseline}$) of each non-target 16 kHz (blue) or 54 kHz (orange) preferring cells for either 16 kHz (top row) or 54 kHz (bottom row) presentation in relation to the mass of center distance to any target cells for the stimulation condition (left), 54 kHz stimulation condition (middle), and the control condition (right). Each circle represents each cell. For the control condition, we considered top 5 most tone-responsive cells from the baseline session as "*target*" cells, as there was no stimulation. Non-target cells are widely distributed within the FOV (550 μm^2), regardless of cell groups, frequencies, and conditions. Gray lines indicate fitted curves, excluding cells that are closer than 15 μm (vertical green lines; cells < 15 μm marked in lighter shades), and dashed lines indicate 95% confidence intervals.