Mitochondrial dysfunction: unraveling the elusive biology behind anti-TNF response during ulcerative colitis

Dimitrios Kioroglou[1], Ainize Peña-Cearra[2,3], Ana M. Corraliza[4], Iratxe Seoane[2,3], Janire Castelo[2], Julian Panés[4,5], Laura Gómez-Irwin[6], Iago Rodríguez-Lago[7], Jone Ortiz de Zarate[8], Miguel Fuertes[10], Itziar Martín-Ruiz[2], Monika Gonzalez[9], Ana Mª Aransay[4,9], Azucena Salas[4], Héctor Rodríguez[2], Juan Anguita[2,11], Leticia Abecia[2,3,*], Urko M. Marigorta[1,11,*]

* Corresponding authors:
Leticia Abecia: leticia.abecia@ehu.eus
Urko M. Marigorta: umartinez@cicbiogune.es

Affiliation details
1. Integrative Genomics Lab, Center for Cooperative Research in Biosciences (CIC bioGUNE), Basque Research and Technology Alliance (BRTA), Bizkaia Technology Park, Derio, Basque Country, Spain.
2. Inflammation and Macrophage Plasticity Laboratory, CIC bioGUNE-BRTA (Basque Research and Technology Alliance), Derio 48160, Spain.
3. Immunology, Microbiology and Parasitology Department, Faculty of Medicine and Nursery, University of the Basque Country, UPV/EHU, P.O. Box 699, 48080 Bilbao, Spain.
4. Centro de Investigación Biomédica en Red de Enfermedades Hepática y Digestivas (CIBERehd), ISCIII, Barcelona, Spain.
5. Department of Gastroenterology, IDIBAPS, Hospital Clínic, Barcelona, Spain.
6. Hospital Universitario de Cruces and Biocruces Bizkaia Health Research Institute, 48903 Barakaldo, Spain.
7. Hospital Universitario de Galdakao and Biocruces Bizkaia Health Research Institute, 48960 Galdakao, Spain.
8. Hospital Universitario de Basurto. 48013 Bilbao, Bizkaia, Spain.
9. GAP, CIC bioGUNE-BRTA (Basque Research and Technology Alliance), Derio 48160, Spain.
10. NEIKER-Basque Institute for Agricultural Research and Development, Basque Research and Technology Alliance (BRTA), Bizkaia Science and Technology Park, Building 812L, 48160, Derio, Spain.
11. Ikerbasque, Basque Foundation for Science, 48013 Bilbao, Bizkaia, Spain.

Abbreviations
MM = colon samples from murine disease model
WT = all wild-type mice included in the MM dataset
WT-IFX$^-$ = wild-type mice without anti-TNF treatment
WT-IFX$^+$ = wild-type mice with anti-TNF treatment
MD = all mice of MM with MCJ-deficiency
MD-IFX$^-$ = mice with MCJ-deficiency and without anti-TNF treatment
MD-IFX$^+$ = mice with MCJ-deficiency and with anti-TNF treatment
MMset = 554 genes from MM that were differentially expressed between MD-IFX$^+$ and WT-
IFX$^+$.
COL = colon samples at week-14 from a human cohort of responders and non-responders to
anti-TNF UC patients.
COLset = 127 genes that intersected with MMset and were differentially expressed in the
COL cohort.
HABset = 89 genes from Haberman *et al.* corticosteroid based gene signature that were
differentially expressed in the COL cohort.
MON = monocyte samples of peripheral blood from a human cohort of healthy individuals and
IBD patients.
MONset = 48 genes that intersected with COLset and were differentially expressed in the
MON cohort.
BLD = whole blood samples at baseline (week-0) from a human cohort of responders and
non-responders to anti-TNF UC patients.
BLDset = set of 14 genes that were used to classify response to anti-TNF from a whole blood
sample at baseline.
PCA = principal component analysis
PCs = principal components
DEG = differentially expressed gene
DSS = dextran sodium sulfate

Abstract

Background: Recent studies hint at mitochondrial genes influencing UC patient response to anti-TNF treatment. We evaluated this hypothesis by following a targeted strategy to identify gene expression that captures the relationship between mitochondrial dysregulation and response to treatment. Our objective was to initially examine this relationship in colon samples and subsequently assess whether the resulting signal persists in the bloodstream.

Methods: We analyzed the transcriptome of colon samples from an anti-TNF treated murine model characterized by impaired mitochondrial activity and treatment resistance. We then transferred the findings that linked mitochondrial dysfunction and compromised treatment response to an anti-TNF treated UC human cohort. We next matched differential expression in the blood using monocytes from peripheral blood of controls and IBD patients, and we evaluated a classification process at baseline with whole blood samples from UC patients.

Results: In human colon samples, the derived gene-set from the murine model showed differential expression, primarily enriched metabolic pathways, and exhibited similar classification capacity as genes enriching inflammatory pathways. Moreover, the evaluation of the classification signal using blood samples from UC patients at baseline highlighted the involvement of mitochondrial homeostasis in treatment response.

Conclusion: Our results highlight the involvement of metabolic pathways and mitochondrial homeostasis in determining treatment response and their ability to provide promising classification signals with detection levels in both colon and bloodstream.

Introduction

Ulcerative colitis (UC) is an inflammatory bowel disease (IBD) of increasing worldwide

incidence, characterized by relapsing and remitting periods. Lacking a definitive cure, the

focus of treatments is to alleviate and control disease symptoms. Although current anti-TNF

treatments can extend the remission periods, a considerable 10-30% of patients do not

respond to the initial treatment, or lose their ability to respond (23-46%) over time [1].

Through comparisons of gene expression profiles of responder and non-responder patients,

several genes with biomarker potential have been proposed. This includes intestinal

Oncostatin M (OSM), IL13RA2 and TREM1, which were first identified using mucosal

samples [2–4]. Similarly, Haberman *et al.* [5] provided a gene signature derived from rectal

biopsies at baseline. Although originally described for response to corticosteroids, the gene

signature had a similar efficiency on a cohort that received anti-TNF treatment, whereas

reduced mitochondrial biogenesis was reported as an aggravating factor.

To avoid the invasive nature of the patient's monitoring via biopsies, identifying whole blood

biomarkers is a priority. For instance, TREM1 expression is arising as a promising predictive

biomarker, although the evidence is not yet conclusive [4,6]. In addition, a qPCR classifier

aimed at classifying patients according to disease course is available [7].

In spite of the valuable endeavors towards classification solutions, current methodologies are

based on differential gene expression analysis that lacks a perturbation source. To derive

insights on the underlying mechanism and provide a gene signature that includes a

reasonable number of differentially expressed genes (DEGs), studies resort to pathway

enrichment analysis. However, this approach is based on using databases that may present

limitations, including outdated or missing pathway information, and, most importantly, they are

not disease specific [8–10]. Thus, the accuracy of the emerging patterns is uncertain and may

result in lack of clinical utility for guiding therapy, as has recently been shown by Noor et al. in

a study of Crohn's disease where the authors evaluated the classification efficiency of a

blood-based biomarker [11]. Thus, more refined and targeted methodologies are necessary

for advancing biomarker discovery.

Drug response in the context of UC involves widespread changes in inflammation, and as a

result pathways with significant contribution to the treatment response can be overshadowed

by inflammatory pathways. In order to unravel the significant contribution of obscured biology,

our approach utilizes a murine model coupled with a perturbation source to identify explicit

gene expression patterns, and capture underlying mechanisms that represent promising

candidates to decipher the lack of response to treatment.

Recent evidence has highlighted the potential involvement of mitochondrial dysfunction in

disease severity and treatment response [5], emphasizing the need to identify underlying

factors that contribute to mitochondrial dysregulation. In this regard, we have previously

described that the absence of Methylation-controlled J protein (MCJ), or DnaJC15, an

endogenous negative regulator of mitochondrial complex I, leads to metabolic alterations,

increases respiration and adenosine triphosphate (ATP) synthesis and stimulates the

formation of respiratory supercomplexes, limiting oxidative stress [12–14]. This qualifies MCJ

as a good perturbation source to simulate complex I mitochondrial dysfunction. Our goal was

to transfer the findings to human cohorts and assess the persistence of the emerged patterns.

In the context of experimental colitis in a murine model, we have previously shown that

deficiency of MCJ contributes to disease severity by modulating the gut microbial composition

of the host [15-17]. Moreover, recently we have reported the critical role played by

macrophage mitochondrial function in the gut ecological niche, which can substantially affect

the severity of inflammation and the ability to successfully respond to anti-TNF therapies [18].

The lack of response to the treatment was expressed as an aggravated phenotypic profile,

where treated MCJ-deficient mice displayed significantly increased weight loss and

histological damage compared to the treated wild-type mice.

In the current study, we profiled the transcriptome of colonic tissue from the aforementioned

MCJ-deficient murine model of DSS-induced colitis [18], to generate a gene set that captures

the relationship between mitochondria dysfunction and lack of response to anti-TNF

treatment. Activated monocytes are recruited to the site of inflammation induced by DSS

where they differentiate into macrophages. The latter can be polarized towards different

subpopulations that can initiate either pro-inflammatory or anti-inflammatory responses [19]. It

has been postulated that the FcγR signaling pathway is implicated in the modulation of these

diametrically opposed responses [20]. Moreover, it has been suggested that the formation of

immune complexes between soluble TNF and therapeutic anti-TNF compounds is necessary

for the FcγR-mediated suppression of pro-inflammatory cytokines. We have shown that MCJ

expression is restricted to macrophages in colonic tissue, and MCJ-deficiency leads to

upregulation of genes belonging to the FcγR signaling pathway and aggravation of the DSS-

induced pathology [18]. Thus, we hypothesize that mitochondrial dysregulation can indirectly

compromise macrophage function and especially their ability to initiate inflammation-resolving

responses.

As summarized in Figure 1, we followed a progressive strategy where the captured gene set

from the murine model was initially transferred and evaluated in a human cohort that provided

transcriptomic data of colonic tissue. Subsequently, we explored whether the captured signal

can be detected in blood samples by applying a successive refinement with two human

cohorts profiling the transcriptome of peripheral blood monocytes, and whole blood samples.

Finally, we utilized an external dataset of whole blood samples from UC patients to evaluate

the replication of the findings in blood.

This is the first study to use a perturbation source that simulates mitochondrial dysregulation

to scrutinize the involvement of the latter on the lack of response to anti-TNF treatment during

ulcerative colitis in human patients. Our findings shed new light on the role of metabolic

pathways and mitochondrial homeostasis in determining treatment response, and pave the

way for further research into the mechanisms underlying anti-TNF response in ulcerative

colitis and other inflammatory diseases.

Methods

Ethics

All procedures involving animals, including their housing and care, were carried according to

the guidelines of the European Union Council (Directive 2010/63/EU) and Spanish

Government regulations (RD 53/2013), and with the approval of the ethics committee of CIC

bioGUNE (Spain; permit number CBBA-0615) and the Competent Authority (Diputación de

Bizkaia). All human samples were processed within the context of the IBD Sample Collection

registered at the HCB-IDIBAPS Biobank and the protocol was approved by the Institutional

Ethics Committee of the Hospital Clinic de Barcelona (HCB/2023/0523). Additional approval

for the human samples was provided by the Clinical Research Ethics Committee from

Euskadi (CEIC-E) following the Helsinki convention (PI2016034). In all human cohorts,

patients provided written informed consent following extensive counseling.

Animal model and experimental design

The murine model [18] included MCJ-deficient (MD) and wild-type (WT) male mice and is

referred to as MM dataset herein. To evaluate the response to anti-TNF treatment, colitis was

induced in WT and MD mice by administering dextran sodium sulfate (DSS) [15]. The whole

experimental period lasted nine days and the mice were divided into two groups per

genotype. One group received Infliximab (IFX$^+$) treatment on the third and sixth day (WT-IFX$^+$,

MD-IFX$^+$), whereas the second group did not receive the treatment (WT-IFX$^-$, MD-IFX$^-$). The

distribution of mice per condition was 5 WT (2 WT-IFX$^-$ and 3 WT-IFX$^+$) and 5 MD mice (2

MD-IFX$^-$ and 3 MD-IFX$^+$). Colon samples were taken from all mice at the end of the

experimental period, namely nine days after the treated mice had been administered

Infliximab (day-3 and 6), and used to study RNA expression.

All procedures and specifications involving animal treatment and the design of the mouse

model are described in the supplementary methods (S1).

Human cohorts

Human cohorts are referred to with the abbreviations COL, MON and BLD. The COL cohort

(N=28) included colonic biopsies at week-14 after anti-TNF treatment of UC patients. The

MON cohort (N=31) included healthy blood donors and IBD patients that provided monocytes

from peripheral blood. The IBD patients were individuals that presented an aggravated state

of the disease and either did not respond to previous treatments or manifested severe

symptoms. As a result, the IBD patients proceeded to emergency colectomy where blood

samples were taken prior to operation. The BLD cohort (N=34) included whole blood samples

from UC patients prior to anti-TNF treatment (week-0) and their response was evaluated at

week-14. The COL and BLD cohorts provided in total 39 patients, from which 23 patients

provided whole blood samples at week-0 and colon samples at week-14, whereas 5 patients

provided only colon samples at week-14 and 11 patients only whole blood samples at week-0.

Moreover, response to treatment was based on endoscopic activity. Age and gender of

patients at inclusion are shown in Table 1 for each cohort. Cohort details and specifications

regarding sample collection and treatment are provided in the supplementary methods (S2).

RNA extraction, preparation and quantification

In the mouse model and human cohorts, gene quantification was performed with RNASeq

using the GRCm38 mouse reference and the GRCh38.p13 human reference, respectively. A

gene was characterized as differentially expressed (DEG) for a given condition contrast, if it

met the following criteria: FDR-corrected p-value (p-adj) ≤ 0.05 and absolute fold-change $|FC|$

≥ 1.5. The derived DEGs were later utilized for enrichment analysis from which only

statistically significant enriched pathways were considered (p-adj ≤ 0.05). From each human

cohort, the set of significant genes that was selected for downstream analysis was named as

COLset, MONset and BLDset denoting the cohort of origin.

Detailed specifications regarding RNA extraction and preparation protocols, as well as

RNASeq preprocessing and enrichment analysis, are given in the supplementary methods

(S3).

Incorporation of the Haberman et al. corticosteroid gene signature into the analytical pipeline

We incorporated into the analysis the gene signature provided by Haberman et al. [5]. The

signature contains 115 genes that resulted from baseline rectal biopsies of 206 pediatric UC

patients whose remission status was evaluated on the week-4 of corticosteroid treatment.

Notably, this signature exhibited similar classification efficiency in a cohort treated with anti-

TNF treatment. From the provided 115 genes, we selected 89 genes that were differentially

expressed in our COL cohort. We named the selected genes as HABset, and were used to

conduct a comparative assessment with the COLset on their ability to differentiate between

responders and non-responders in the COL cohort.

Evaluation of the persistence of the captured signal from colon tissue in

blood samples

Since MCJ expression is restricted in macrophages in colonic tissue [18], we sought to

explore whether the captured signal from the COL cohort can be detected in the blood. To do

so, we initially identified 48 genes (MONset) in the MON cohort that can distinguish IBD

patients from non-IBD individuals. Later, we utilized whole blood samples before treatment

from the BLD cohort, where our process for resolving crossover trends between the BLD and

MON cohorts provided 14 genes (BLDset) that can classify responders and non-responders

IBD patients to treatment. The process followed for the construction of the BLDset is

described in the supplementary methods along with the concept of trends (S4).

For the classification process in the BLD cohort, we initially regressed-out age and sex from

the gene expression data. To avoid overfitting and multicollinearity, we performed PCA on the

gene expression and based a non-regularized logistic regression model on the first two PCs

with Scikit-learn [21]. One additional advantage of this strategy is that it enables the visual

evaluation of the model's performance. To mitigate the impact of the imbalanced conditions,

the weights 0.3 and 0.7 were used for the non-responder and responder classes respectively.

To evaluate the training process, we performed a 12-fold cross-validation with 3 repeats

calculating at each fold the F1-score metric for the BLD training and BLD test-set. The F1-

score was chosen since it is a well-suited metric for imbalanced categorical variables. It is

interpreted as the harmonic mean of the precision and recall metrics and ranges between 0

and 1. Finally, estimation of the decision boundary standard error was based on a

bootstrapped 95% confidence interval with 1000 iterations, sampling with replacement and

sample size of 34.

Replication of the findings in blood with an external dataset

To evaluate the replication of the findings in the whole blood, we downloaded the publicly

available RNASeq data provided by Mishra *et al.* [22] (GSE191328). It is a blood-based

dataset that underwent a similar preprocessing procedure as the BLD cohort, and included

three independent cohorts of IBD patients: two that received Infliximab treatment (referred to

as discovery and replication cohorts by the authors) and one Vedolizumab. The patients were

followed longitudinally from baseline until week-14, where they were classified according to

remission status. From the available cohorts, we considered only UC patients and focused on

the following time-points: baseline, week-2, week-6 and week-14. The discovery cohort is

referred to as IFX (N=9) and the replication cohort as REP (N=9), whereas the Vedolizumab

cohort is referred to as VED (N=10) in the current study. To avoid possible integration batch

effects and to evaluate patients heterogeneity, we kept the IFX and REP cohorts separate.

Prior to model validation, the genes *SLC18A1* and *PPARGC1A* were omitted from the BLDset

since they were not quantified across all cohorts and timepoints. We also processed each

cohort with DESeq2 stabilizing the gene expression variance with the dispersion function

whose parameters were previously estimated with the BLD cohort without specifying patient

response. Thus, no patient's response information was introduced during the dispersion

estimate that could bias the classification process. Then, each cohort was projected on the

PCs that were extracted during the model's training process and the sample scores from the

resulting transformation were used as a test-set. Finally, the performance of the classification

process in each cohort was assessed using the F1-score metric.

Results

554 DEGs in the mouse model (MMset) capture the interaction between the

absence of MCJ and the response to anti-TNF

First, we performed transcriptomic analysis of the colon tissue from a murine model of

experimental colitis that included wild-type (WT) and MCJ-deficient (MD) mice. The mouse

groups were further subdivided into treated with anti-TNF (WT-IFX$^+$,MD-IFX$^+$) and untreated

mice (WT-IFX$^-$,WT-IFX$^-$). As mentioned above, response to anti-TNF treatment was

previously evaluated [18]. Similarly to the WT-IFX$^-$ and MD-IFX$^-$ mice, the MD-IFX$^+$ mice

displayed an aggravated profile characterized by significantly higher weight loss and

histological damage compared to WT-IFX$^+$. Additionally, in the current study we observed

high efficiency of the model, with an 8.4-fold and 11.3-fold changes in the mRNA levels of

Dnajc15 between the MD-IFX$^-$ and WT-IFX$^-$ and the MD-IFX$^+$ and WT-IFX$^+$ groups,

respectively (Figure S1).

To evaluate group differences, we performed principal component analysis (PCA) using all

16,186 quantified genes. The MD-IFX$^+$ mice had unique distinguished profiles compared with

the other three groups, forming a separate cluster across PC1 (45% of the observed variance,

Figure 2A, left plot). The differences between the other three groups appeared weaker,

clustering mainly across PC2 (18.8% of the observed variance).

Differential gene expression analysis corroborated these observations, with greater intra-

group differences within the MD groups than among WT groups. The contrast between MD-

IFX$^+$ and MD-IFX$^-$ resulted in 1,544 differentially expressed genes (DEGs), for only 12 DEGs

in the WT-IFX$^+$ and WT-IFX$^-$ comparison. Of note, we observed an interaction between

Dnajc15 gene expression and the response to the anti-TNF treatment. Upon administration of

Infliximab, 617 genes were differentially expressed between the treated groups (MD-IFX$^+$ and

WT-IFX$^+$), for only 12 genes among the corresponding non-treated groups. These

observations verify the utility of the mouse model, suggesting a significant interaction between

Dnajc15 expression and the response to Infliximab treatment.

We reasoned that the 617 DEGs from the MD-IFX$^+$ versus WT-IFX$^+$ comparison represent a

set of genes with the capacity to decipher the mechanisms underlying the relationship

between mitochondrial dysregulation and unresponsiveness to the treatment. After removing

genes without corresponding human ENSEMBL identification, 554 DEGs were retained,

referred to heretofore as the MMset (supplementary materials 1). As expected, PCA showed

that the MMset recapitulates the differences between the MD-IFX$^+$ and the other groups,

forming a separate cluster across PC1 (72.78% of the observed variance, Figure 2A, right

plot).

MMset is associated with mitochondrial dysfunction

We used MitoMiner's API [23] to identify genes located in the mitochondrion. We identified 50

genes from the MMset (supplementary materials 1). These genes had a median 1.21-fold

average expression change between the non-treated (WT-IFX⁻ and MD-IFX⁻) groups,

increasing to 1.86-fold change between the treated groups (Figure S2). These results

reinforce the previous idea of an interaction between anti-TNF administration and

mitochondrial dysfunction.

We identified other notable changes in genes with relevant roles in mitochondrial

homeostasis. *Phgdh,* known to compromise the electron transport chain activity and lead to

increased production of ROS [22], was significantly down-regulated in the MD-IFX⁺ group. We

observed similar changes for *Hif1a* and *Ripk3*, that play a role in modulating mitochondrial

function and metabolism [25,26]. In turn, we observed significant up-regulation of *Ppargc1a*

and *Pink1*, which are involved in regulating mitophagy and mitochondrial biogenesis [27].

The major enrichment in MMset concerned metabolic pathways

To further characterize the biological effects underlying the MMset, we performed enrichment

analysis of the full set. Subsets of 107 and 56 genes were associated with metabolic and cell

cycle pathways, respectively (supplementary materials 2). From the subset associated with

metabolic pathways, 65.4% of the genes were up-regulated in the MD-IFX⁺ (Figure 2B). In

contrast, genes associated with cell cycle pathways were down-regulated in MD-IFX⁺. The

latter represented an interesting observation and we reasoned that the excessive

downregulated gene expression concerned cells that had undergone apoptosis. If this

rationale holds true, then it is aligned with our hypothesis that mitochondrial dysfunction

disrupts the proper function of macrophages to initiate inflammation-resolving processes such

as clearance of apoptotic cells.

The enrichment of metabolic pathways involved the activation of the xenobiotic response

system in the MD-IFX$^+$ group. This includes nuclear receptors that regulate genes involved in

the metabolism of xenobiotics, such as the farnesoid X receptor (FXR/*Nr1h4*) that exhibited a

1.67-fold change between the MD-IFX$^+$ and WT-IFX$^+$ groups causing the over-expression of

the gene *Fabp6*. Moreover, the peroxisome proliferator-activated receptor α (*Ppara*), which

regulates the expression of various UGT metabolic enzymes, ABC and SLC transporters, was

also found differentially expressed. We also observed up-regulation of the peroxisome

proliferator-activated receptor-gamma coactivator (PGC-1/*Ppargc1a*) gene, which interacts

with PPARA and together regulate the expression of enzymes involved in the oxidation of

mitochondrial fatty acids.

MMset signal linking anti-TNF response and mitochondrial dysfunction

successfully transferred to a human cohort (COLset)

We next analyzed available colon samples from a human UC cohort. Out of 21,780 quantified

genes, 3,645 were differentially expressed between responders and non-responders patients.

PCA showed a separation between the two groups across PC1, accompanied by an

underlying intra-group variability predominantly influenced by PC2 (Figure 3A, "All DEGs").

We then evaluated the expression of genes that could be linked to mitochondrial dysfunction.

We used the GeneCards portal [28] to query for genes that are associated with complex I.

The search returned 165 genes, of which 25 were differentially expressed (supplementary

materials 3). A detailed inspection revealed mitochondrial alterations, with downregulation of

complex I activity components such as *MT-ND3*, *MT-ND5*, *MT-ND4*, *MT-ND4L*, *MT-TL1*, and

MT-ND2, and up-regulation of genes such as *NLRP3*, that gets activated by excessive

mitophagy [29], or *DYSF* that regulates cytosolic influx of Ca^{2+} and affects normal

mitochondrial function under certain conditions [30].

To evaluate the mitochondrial aberrations that were linked to administration of anti-TNF in the

murine model, we intersected the 3,645 DEGs with the MMset. This generated a subset of

127 DEGs, named COLset (supplementary materials 1). PCA showed that COLset was able

to maintain the separation between responders and non-responders and increased the

captured observed variance to 70% with the first two PCs (Figure 3A, "COLset"). Interestingly,

although *DNAJC15* displayed statistically significant differences between responders and

non-responders (p-adj=0.03), its fold-change of 1.3 between the patient groups did not allow it

to be considered as differentially expressed and was not included in the COLset. This

particular case exemplifies perfectly the potential pitfalls of the typical strategy followed for the

discovery of DEGs, as genes that contribute to the underlying mechanism can be overlooked

by strict predefined thresholds.

COLset performs similarly to inflammation-driven gene signatures, and

shows metabolism as the main enriched pathway

We then juxtaposed the discriminatory capability on the COL cohort of the COLset with a set

of 89 genes from the corticosteroid-based gene signature provided recently by Haberman *et*

al. [5] that we refer to as the HABset (see methods, supplementary materials 1).

Both gene sets captured above 70% of the observed variance with the first two PCs (Figure

3A), with PC1 scores showing a strong negative Pearson correlation (COLset: r=-0.69, p-

value=2.6E-05, HABset: r=-0.70, p-value=2.1E-05) with *DNAJC15* mRNA levels (Figure S3).

These results suggest a severity gradient of non-response that correlates with the extent of

the underlying MCJ-dependent mitochondrial dysfunction. Nevertheless, both sets shared

only 9 genes (*GUCA2B*, *TNIP3*, *SLC26A3*, *VSIG1*, *TRPM6*, *HMGCS2*, *MMP3*, *MMP10* and

CHP2) suggesting that the two signatures capture different underlying gene expression

regulatory components, albeit sharing functional properties.

Indeed, enrichment analysis revealed different captured pathways. The COLset enriched

pathways associated with metabolism, transportation of molecules and regulation of nuclear

receptors, whereas the HABset appeared linked to pathways related to the immune system,

IL-17 signaling and peptide receptors (supplementary materials 2).

Finally, we evaluated the classification efficiency of the two gene sets by constructing two

predictive models on the COL cohort (supplementary methods S5). The F1-score was 0.93

with the COLset and 0.97 with the HABset. As a result, the decision boundary separated well

the patients groups in both cases (Figure 4). Overall, these two gene sets, although clearly

different in regards to gene composition and pathway enrichment, yield similar classification

results on samples from colonic tissue from UC patients undergoing anti-TNF treatment.

COLset signal was detected in circulating blood monocytes (MONset)

We analyzed RNASeq data from peripheral blood monocyte samples with the aim to assess

the persistence of the captured signal from the colon samples in the blood. We quantified

18,331 genes and identified a set of 9,262 DEGs between healthy non-IBD individuals and

IBD patients with a severe disease state that were sampled right before colectomy. The

intersection of the latter with the 165 genes related to complex I from GeneCards returned 83

genes (supplementary materials 3). Fifty-three genes were up-regulated (FC M=2.89,

SD=1.89) and 30 were down-regulated (FC M=2.97, SD=2.15) in the IBD patients. Moreover,

the IBD patients exhibited up-regulation of the majority of the NDUF genes that are

associated with complex I, and down-regulation of the majority of the TMEM genes that are

directly related to ATP synthesis.

To associate differential expression related to anti-TNF treatment, we intersected the 9,262

DEGs with the COLset, generating a subset of 48 genes which we named as MONset

(supplementary materials 1). We attributed the narrow overlap in the homogeneous and

purified nature of the samples included in the MON cohort compared to those of the COL

cohort. PCA revealed a clear separation between IBD patients and non-IBD individuals,

capturing 68% of the observed variance with the first two PCs (Figure 5A). Moreover,

enrichment analysis showed metabolism as the sole enriched pathway encompassing 17

genes (Figure 5B-C, supplementary materials 2). These results indicate that dysregulated

complex I activity can be detected in circulating monocytes, even before they migrate to

colonic tissue and differentiate into macrophages.

Pre-treatment whole blood samples reveal latent classification signals for UC patient response, linked to mitochondrial dysfunction (BLDset)

Following the results from the MON cohort, our next objective was to investigate whether the

classification signal of COLset, between responders and non-responders IBD patients, can

also be detected in whole blood samples especially before treatment. To do so, we analyzed

whole blood samples at baseline of UC patients whose status was identified at a later stage of

the treatment (week-14). We quantified 24,039 genes that did not exhibit differential

expression between responders and non-responders. The similarity in their transcriptomic

profiles at baseline implies that unbiased classification at week-0 through classical gene

signature discovery is not possible. However, non-responders displayed a mean of 1.18-fold

difference (SD=0.36) compared to responders for the 165 genes associated with complex I

from GeneCards (supplementary materials 3). This suggests that perturbations of

mitochondrial genes are present before the initiation of the treatment.

Due to lack of differential expression in the BLD cohort, it was unfeasible to intersect the

COLset with a set of DEGs in the BLD cohort. Therefore, to proceed we utilized the MONset

that includes genes from the COLset that exhibit differential expression in the blood.

Nevertheless, certain genes from the MONset displayed crossover trends between the BLD

and MON cohorts that we attributed to the lack of responders in the MON cohort. Thus, we

considered a MONset subset of 14 genes to be included in the classification process which

hereafter will be referred to as BLDset (see methods section, supplementary materials 1).

During the 12-fold cross-validation of the training process, the classifier achieved a mean F1-

score of 0.79 (95% CI: 0.78-0.80) with the BLD training-set, and a mean F1-score of 0.69

(95% CI: 0.59-0.79) with the BLD test-set. The panels A and B in Figure 6 depict the decision

boundary applied by the classifier and the probability of the assigned class (responder or non-

responder) to each included patient.

External replication showed attenuated baseline BLDset classification

signal in blood, with marked post-treatment improvement

To evaluate the replication of the classification process with BLDset, we utilized publicly

available blood-based RNASeq data provided recently by Mishra *et al.* [22] (see methods).

The study profiled two cohorts that received anti-TNF treatment (IFX and REP) and one

cohort that was treated with the $\alpha 4\beta 7$ integrin inhibitor Vedolizumab (VED). Patients were

followed longitudinally from baseline until week-14, and the distribution of the patient groups

is given in the supplementary methods (S6).

At baseline we observed great variability of the classification performance that marked the

lowest mean F1-score with the IFX cohort (Figure 7 A-B). However, the classification

drastically improved after the initiation of the treatment marking the best performance on

week-14. Moreover, we observed that the classification performance for the VED cohort did

not follow the same pattern as with the IFX and REP cohorts after baseline, implying

specificity of the BLDset towards the anti-TNF treatment.

To investigate further the variability in classification performance observed at baseline, we

calculated the number of genes with matched ranks (Figure 7A) and trends (Figure 7B)

between the BLDset and the external dataset, and compared them to see how they affected

the performance. For the IFX cohort, we observed that it showed the highest number of

matched ranks with the BLDset, but with only three genes with matched trends. This

observation underscored the poor classification performance as compared to the REP and

VED cohorts.

Although the above observations allowed us to explain the poor performance on the IFX

cohort at baseline, the improvement in classification on this cohort at week-6 remained

puzzling since the number of matched ranks and trends were relatively the same as in

baseline. Initially, we observed that in both timepoints the genes *SLC18A1* and *PIF1*

displayed similar trends, and the trend differences concerned the gene *EDN1* (baseline) and

STON1 (week-6) (supplementary Figure S4). We hypothesized that this observation could be

associated with latent patterns of gene networks that are associated with the BLD cohort and

have been indirectly incorporated in the BLDset and potentially affect the classification in the

external dataset. To do so, we initially calculated a mean expression for the BLD cohort

concerning the 165 complex I related genes from GeneCards. We then performed the same

exercise for each cohort in the external dataset and compared them with the BLD cohort

(Figure 7C). Interestingly, we observed that the performance was influenced by the captured

state of the patient groups in the BLD cohort regarding these 165 complex I genes. More

specifically, the performance improved in cases where the average fold-changes related to

these genes between responders and non-responders were closer to those observed in the

BLD cohort. Since this could be observed across all timepoints, it corroborated our hypothesis

about latent gene networks that can shape the performance of a classifier.

Discussion

Mitochondrial genes have recently been highlighted as potentially critical to explain the

underlying mechanism that compromises the response to anti-TNF treatments in UC patients.

Since such signals could be overshadowed by inflammation-driven pathways, we

hypothesized that exploring a mouse model with a clear perturbation source is ideal to

unravel obscured biology. In the current study, we first analyzed a murine model of

experimental colitis characterized by mitochondrial dysfunction and lack of response to anti-

TNF. To simulate a mitochondrial dysfunction, we utilized MCJ deficiency as a perturbation

source which is a mitochondrial transmembrane protein that interacts with complex I [12-14].

The initial focus was to select a set of genes (MMset) that showed strong relation with

mitochondrial dysregulation and response to anti-TNF treatment. Our main goal was to

discover underlying expression signals associated with mitochondrial dysfunction that can be

hidden by inflammatory pathways.

Notably, we also observed deficiency in MCJ expression in the COL cohort, with significant

differences between the patient groups. This underscores the relevance of the mouse model

with regards to the mitochondrial-driven ability to respond to anti-TNF, and we successfully

transferred the classification signal captured by MMset to the COL cohort. The transferred

signal (COLset) displayed a significant enrichment of metabolic pathways, and showed

classification capacity similar to the recent gene signature provided by Haberman *et al.* [5]

(HABset), even though the latter is enriched for inflammation pathways and both sets shared

a limited number of genes. This suggests that mitochondrial genes have equal contribution to

treatment outcomes as those associated with inflammation. Indeed, compromised

mitochondria release damage-associated molecular patterns (DAMPs) and induce innate

immune responses [29], establishing an interrelationship between both that could explain the

similar classification capacity of COLset and HABset.

Given the restricted expression of MCJ to colonic tissue macrophages, it was pertinent to

investigate whether the COLset signal extends from the colon to the bloodstream. The MON

cohort allowed us to explore this aspect and we identified a significant classification signal

(MONset) that was primarily enriched in metabolic pathways. Monocytes are circulating short-

lived macrophage precursors that are recruited on demand from the blood to sites of

inflammation such as the colon, and our results indicate that circulating monocytes in IBD

patients are characterized by compromised complex I activity.

Even though classification of pre-treatment response using whole blood samples was not the

main focus of this study, the results obtained by the MON cohort prompted us to explore the

presence of a classification signal at baseline. Since significant insights from such an exercise

can lead to valuable non-invasive classification solutions regarding response, we initially

implemented a classification process using the BLD cohort and attempted to replicate the

process using an external dataset. Although we did not observe robust performance at

baseline, BLDset displayed specificity towards anti-TNF treatment and the whole exercise

revealed the impact of gene ranks and trends on the performance across different datasets,

as well as the influence of non-DEGs that are involved in the regulatory network. This kind of

classification nuances could underlie some of the incongruent results seen in drug response

signatures [4,6].

The BLDset was the classification signal that persisted from the colon samples to the

bloodstream. Its composition included *PPARGC1A*, which is related to mitochondrial

biogenesis [5], was found among the highest ranked in expression in most of the cohorts, and

is associated with more severe colitis and decreased complex IV activity in knock-out mice

[31]. It also included *PHGDH* that has been found to regulate mitochondrial homeostasis [24],

and *RASGRF2* that coordinates processes related to the activity of the N-methyl-D-aspartate

receptor, which has been reported to induce mitochondrial dysregulation [32]. Other members

of the gene set were *SLC18A1*, a transporter with xenobiotic transmembrane activity, the

detoxification gene *SULT1A1* [33], *PBLD* that is involved in the inhibition of NF-kB signaling

[34] and *LRP8* and *PADI2*, which display calcium-ion activity and could be associated with

neoplastic progression in UC [35].

We identified certain limitations associated with our study. The small sample size and the

imbalanced patient groups limited our ability to optimize the classification process further.

This limitation represents an integral part of many studies due to the difficulties of constructing

a UC based cohort attributed to the low annual incidence. As a result, the incorporation of

patients with Crohn's disease (CD) represents a common workaround in order to increase

statistical power. However, CD and UC patients exhibit different transcriptomic profiles [36]

and following such strategies increases the risk of introducing additional latent factors into the

predictive model. Furthermore, the interaction between gut microbiota and gene expression

has not been incorporated into the model, since it would require integration of additional omic

data such as metabolomics. There is an interplay between mitochondria and intestinal

microbiota, with the latter releasing metabolites that trigger mitochondrial ROS which in turn

induce an immune response [15,37]. This interaction may affect the transcriptomic profile of

the patients and hence indirectly impact the performance of the classifier.

Besides clarifying the molecular mechanisms that underlie the relationship between

mithocondrial dysfunction and treatment response, the strategy we followed entails three

insights of interest for future gene signature discovery attempts. First, we show that a well-

defined murine model of response to therapy - here in the context of mitochondrial

dysfunction - can be representative of response traits observed in patients. Second, the

inclusion of a perturbation source allows the identification of key regulatory networks that in

other works using standard approaches can be obscured due to major pathways that are

captured by enrichment analysis. Third, considering that we leveraged all publicly available

longitudinal datasets for the replication of the classification process, our findings raise caution

regarding the implementation of classification solutions at baseline, and suggest that more

longitudinal studies are needed to characterize well the involved regulatory pathways.

565 Conclusion

566 We highlighted the importance of mitochondrial homeostasis in relation to treatment response

567 in the context of UC. Our results indicate that regulatory networks associated with

568 mitochondrial dysfunction can provide promising classification signals that can be detected

569 even in the bloodstream, leading to effective non-invasive classification solutions.

Contributors

LA, UMM conceived and supervised the study. APC, AMC, IS, JC, JP, LGI, IRL, JOZ, MF, IMR, MG, AMA, AS, HR performed experiments and/or provided human samples. LA, JA supervised experiments. DK analyzed data. DK, UMM wrote the manuscript. All authors have reviewed the manuscript.

Data sharing statement

The raw transcriptomic data for the mouse model and the COL, MON and BLD human cohorts are publicly available in the SRA repository under the accession number PRJNA1075662.

Declaration of interests

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

This work was supported by grants from the Spanish Ministry of Science MCIN/AEI/10.13039/501100011033 (PID2019-108244RA-I00 to UMM, PID2021 124328OBI00 to J.A., RYC2020-030632-I to U.M.M, CEX2021-001136-S to U.M.M and D.K), "la Caixa" Foundation (ID 100010434, fellowship code LCF/BQ/LI18/11630001 to U.M.M), V Grant from GETECCU-MSD (Grupo Español de Trabajo en Enfermedad de Crohn y Colitis ulcerosa to L.A.). A.P.C. and I.S. are supported by fellowships from the University of the Basque Country (UPV/EHU) and the Basque Government, respectively.

Acknowledgements

We acknowledge all the personnel of the Hospital Clinic IBD Unit (Clinic and laboratory) who participated in patient selection and sample collection, handling and processing and we also thank all the patients who participated in this study and the Research Committee from OSI Barrualde-Galdakao.

References

Roda G, Jharap B, Neeraj N, *et al.* Loss of response to anti-TNFs: Definition, epidemiology, and management. *Clinical and translational gastroenterology* 2016;**7**:e135.

West NR, Hegazy AN, Owens BM, *et al.* Oncostatin m drives intestinal inflammation and predicts response to tumor necrosis factor–neutralizing therapy in patients with inflammatory bowel disease. *Nature medicine* 2017;**23**:579–89.

Arijs I, Li K, Toedter G, *et al.* Mucosal gene signatures to predict response to infliximab in patients with ulcerative colitis. *Gut* 2009;**58**:1612–9.

Verstockt B, Verstockt S, Dehairs J, *et al.* Low TREM1 expression in whole blood predicts anti-TNF response in inflammatory bowel disease. *EBioMedicine* 2019;**40**:733–42.

Haberman Y, Karns R, Dexheimer PJ, *et al.* Ulcerative colitis mucosal transcriptomes reveal mitochondriopathy and personalized mechanisms underlying disease severity and treatment response. *Nature communications* 2019;**10**:1–3.

Gaujoux R, Starosvetsky E, Maimon N, *et al.* Cell-centred meta-analysis reveals baseline predictors of anti-TNFα non-response in biopsy and blood of patients with IBD. *Gut* 2019;**68**:604–14.

Biasci D, Lee JC, Noor NM, *et al.* A blood-based prognostic biomarker in IBD. *Gut* 2019;**68**:1386–95.

Wadi L, Meyer M, Weiser J, *et al.* Impact of outdated gene annotations on pathway enrichment analysis. *Nature methods* 2016;**13**:705–6.

Mubeen S, Tom Kodamullil A, Hofmann-Apitius M, *et al.* On the influence of several factors on pathway enrichment analysis. *Briefings in Bioinformatics* 2022;**23**:bbac143.

Ma J, Shojaie A, Michailidis G. Network-based pathway enrichment analysis with incomplete network information. *Bioinformatics* 2016;**32**:3165–74.

Noor NM, Lee JC, Bond S, *et al.* A biomarker-stratified comparison of top-down versus accelerated step-up treatment strategies for patients with newly diagnosed Crohn's disease (PROFILE): a multicentre, open-label randomised controlled trial. *The Lancet Gastroenterology & Hepatology*. 2024.

Hatle KM, Gummadidala P, Navasa N, *et al.* MCJ/DnaJC15, an endogenous mitochondrial repressor of the respiratory chain that controls metabolic alterations. *Molecular and cellular biology* 2013;**33**:2302–14.

Navasa N, Martín I, Iglesias-Pedraz JM, *et al.* Regulation of oxidative stress by methylation-controlled j protein controls macrophage responses to inflammatory insults. *The Journal of infectious diseases* 2015;**211**:135–45.

14   Barbier-Torres L, Iruzubieta P, Fernández-Ramos D, *et al.* The mitochondrial negative
regulator MCJ is a therapeutic target for acetaminophen-induced liver injury. *Nature*
communications 2017;**8**:1–1.

15   Pascual-Itoiz MA, Peña-Cearra A, Martín-Ruiz I, *et al.* The mitochondrial negative
regulator MCJ modulates the interplay between microbiota and the host during ulcerative
colitis. *Scientific reports* 2020;**10**:1–3.

16   Peña-Cearra A, Pascual-Itoiz MA, Lavín JL, *et al.* Mitochondrial complex i dysfunction
alters the balance of soluble and membrane-bound TNF during chronic experimental colitis.
Scientific reports 2022;**12**:1–0.

17   Peña-Cearra A, Song D, Castelo J, *et al.* Mitochondrial dysfunction promotes microbial
composition that negatively impacts on ulcerative colitis development and progression. *npj*
Biofilms and Microbiomes 2023;**9(1)**:74.

18   Peña-Cearra A, Castelo J, Lavín JL, *et al.* Mitochondrial dysfunction-associated
microbiota establishes a transmissible refractory response to anti-TNF therapy during
ulcerative colitis. *Gut Microbes* 2023;**15(2)**:2266626.

19 Anderson CF, Mosser DM. Cutting edge: biasing immune responses by directing antigen
to macrophage Fcγ receptors. *The Journal of Immunology*. 2002;**168(8)**:3697-701.

20 Bloemendaal FM, Koelink PJ, van Schie KA, *et al*. TNF-anti-TNF immune complexes
inhibit IL-12/IL-23 secretion by inflammatory macrophages via an Fc-dependent mechanism.
Journal of Crohn's and Colitis. 2018;**12(9)**:1122-30.

21   Pedregosa F, Varoquaux G, Gramfort A, *et al.* Scikit-learn: Machine learning in Python.
Journal of Machine Learning Research 2011;**12**:2825–30.

22   Mishra N, Aden K, Blase JI, *et al.* Longitudinal multi-omics analysis identifies early
blood-based predictors of anti-TNF therapy response in inflammatory bowel disease. *Genome*
medicine 2022;**14**:1–20.

23   Smith AC, Robinson AJ. MitoMiner v3.1, an update on the mitochondrial proteomics
database. *Nucleic acids research* 2016;**44**:D1258–61.

24   Vandekeere S, Dubois C, Kalucka J, *et al.* Serine synthesis via PHGDH is essential for
heme production in endothelial cells. *Cell metabolism* 2018;**28**:573–87.

25   Choudhry H, Harris AL. Advances in hypoxia-inducible factor biology. *Cell metabolism*
2018;**27**:281–98.

26   Islam T, Afonso MB, Rodrigues CM. The role of RIPK3 in liver mitochondria
bioenergetics and function. *European Journal of Clinical Investigation* 2022;**52**:e13648.

27   Novak EA, Mollen KP. Mitochondrial dysfunction in inflammatory bowel disease.
Frontiers in cell and developmental biology 2015;**3**:62.

28   Safran M, Dalah I, Alexander J, *et al.* GeneCards version 3: The human gene
integrator. *Database* 2010;**2010**.

29   Chen G, Kroemer G, Kepp O. Mitophagy: An emerging role in aging and age-
associated diseases. *Frontiers in cell and developmental biology* 2020;**8**:200.

30   Vincent AE, Rosa HS, Alston CL, *et al.* Dysferlin mutations and mitochondrial
dysfunction. *Neuromuscular Disorders* 2016;**26**:782–8.

31   Cunningham KE, Vincent G, Sodhi CP, *et al.* Peroxisome proliferator-activated
receptor-γ coactivator 1-α (PGC1α) protects against experimental murine colitis. *Journal of
Biological Chemistry* 2016;**291**:10184–200.

32   Gao X, Xu X, Pang J, *et al.* NMDA receptor activation induces mitochondrial
dysfunction, oxidative stress and apoptosis in cultured neonatal rat cardiomyocytes.
Physiological research 2007;**56**:559.

33   Langmann T, Moehle C, Mauerer R, *et al.* Loss of detoxification in inflammatory bowel
disease: Dysregulation of pregnane x receptor target genes. *Gastroenterology* 2004;**127**:26–
40.

34   Chen S, Liu H, Li Z, *et al.* Epithelial PBLD attenuates intestinal inflammatory response
and improves intestinal barrier function by inhibiting NF-κb signaling. *Cell death & disease*
2021;**12**:1–5.

35   Brentnall TA, Pan S, Bronner MP, *et al.* Proteins that underlie neoplastic progression of
ulcerative colitis. *PROTEOMICS–Clinical Applications* 2009;**3**:1326–37.

36   Yang L, Tang S, Baker SS, *et al.* Difference in pathomechanism between crohn's
disease and ulcerative colitis revealed by colon transcriptome. *Inflammatory bowel diseases*
2019;**25**:722–31.

37   Özsoy M, Stummer N, Zimmermann FA, *et al.* Role of energy metabolism and
mitochondrial function in inflammatory bowel disease. *Inflammatory Bowel Diseases*
2022;**28**:1443–50.

Figures

Figure 1: Experimental design. **Phase 1**: The analysis was based initially on a murine model of experimental DSS induced colitis characterized by MCJ-deficiency (MD) and mitochondrial complex I dysfunction. A set of 554 genes (MMset) was selected that showed strong relation with mitochondrial genes and response to anti-TNF treatment. **Phase 2**: We transferred the captured signal to a human cohort of UC patients that was treated with anti-TNF and provided colon samples. We identified a set of 127 genes (COLset) that match differential expression in MMset. **Phase 3**: We identified a classification signal of 48 genes (MONset) from the COLset that can be detected in circulating monocytes by comparing healthy individuals and IBD patients with an aggravated state of the disease that proceeded to emergency colectomy. **Phase 4**: Using whole blood samples of UC patients before anti-TNF treatment, we identified

737   a signal of 14 genes from MONset (BLDset). The 14-gene based BLDset became the input of
738   a classification process regarding response before anti-TNF treatment.

746
Figure 2: A) PCA using all 16,186 quantified genes (left) from the mouse model or only the
MMset (right). **B)** Distribution of log2-fold changes (x-axis) between MD-IFX+ and WT-IFX+
for each enriched pathway based on MMset. In each pathway, up and down-regulated genes
are indicated for the MD-IFX+ in relation to WT-IFX+ group. **C)** Up and down-regulated
biological process related to each enriched pathway. Regulation is indicated for MD-IFX+ in
relation to WT-IFX+ group.

763
764

766
Figure 3: A) PCA using either all 3,645 genes that were differentially expressed between
responders and non-responders to anti-TNF UC patients, HABset or COLset. **B)** Distribution
of log2-fold changes (x-axis) between responders and non-responders to anti-TNF for each
enriched pathway based on COLset. In each pathway, up and down-regulated genes are
indicated for non-responders in relation to responders to anti-TNF UC patients. **C)** Up and

down-regulated biological process related to each enriched pathway. Regulation is indicated
for non-responders in relation to responders to anti-TNF UC patients.
774
775
776
777

779
Figure 4: Performance of the anti-TNF logistic regression classifier based on the COL cohort
and the first two PCs of the corresponding PCA that was performed using the COLset (left) or
the HABset (right). Dashed line represents classifier's decision boundary. Patients to the right
side of the decision boundary (shaded orange) have been classified as non-responders by
the model, whereas to the left side (shaded blue) as responders to anti-TNF. Patients have
been annotated based on their observed response (color) and assigned endoscopic Mayo
score (size).

Figure 5: A) PCA using all 9262 genes that were differentially expressed between healthy and IBD patients or genes included in the MONset. **B)** Distribution of log2-fold changes (x-axis) between healthy individuals and IBD patients based on MONset. In each pathway, up and down-regulated genes are indicated for the IBD patients in relation to healthy individuals. **C)** Up and down-regulated biological process related to each enriched pathway. Regulation is indicated for IBD patients in relation to healthy individuals.

Figure 6: A) Performance of the anti-TNF logistic regression classifier based on the BLD cohort and the first two PCs of the corresponding PCA that was performed using the 14 genes of BLDset. Dashed line represents classifier's decision boundary. White shaded area around decision boundary represents bootstrapped 95% CI, associated with the range within which the decision boundary could be positioned. Patients to the right side of the decision boundary (shaded orange) have been classified as non-responders by the classifer, whereas to the left side (shaded blue) as responders to anti-TNF. **B)** the probabilities assigned by the

classifer to each patient are given, regarding the possibility of no-response to anti-TNF
treatment using the BLDset. The classifier implements a probability threshold (decision
boundary) of ≥50% to classify a patient as non-responder (shaded orange area) and <50% as
responder to anti-TNF (shaded blue area). In all graphical representations, patients have
been annotated based on observed response at week-14 after treatment (color) and assigned
endoscopic Mayo score (outline).
821
822
823

825
Figure 7: A) Lines represent F1-score achieved by the classifier for the cohorts that received
anti-TNF (IFX, REP) or Vedolizumab (VED). Bars represent the numbers of BLDset genes
from the corresponding cohorts that received the same rank as those in the BLD cohort. **B)**
Lines represent F1-score achieved by the classifier for the cohorts that received anti-TNF
(IFX, REP) or Vedolizumab (VED). Bars represent the number of BLDset genes from the

corresponding cohorts that exhibit similar trend between patients groups when compared with
those in the BLD cohort. **C)** For each cohort and time-point, the mean fold-change between
the responders and non-responders UC patients is shown (error bars denote standard
deviation), concerning 165 genes related to complex I. Black horizontal line represents the
corresponding mean fold-changes that were observed in the BLD cohort. In all graphical
representations, samples at week-14 were not available for the REP cohort.

Tables

	N	Age (MIN-MAX)	Gender (F/M)
		COL cohort	
responders UC	7	41 (20-60)	3/4
non-responders UC	21	46 (27-70)	9/12
		MON cohort	
healthy	16	N/A (18-65)	N/A
IBD patients	15	49 (23-77)	4/11
		BLD cohort	
responders UC	9	49 (24-87)	3/6
non-responders UC	25	53 (32-78)	11/14

Table 1: Human cohorts specifications. Age is reported based on average and observed
minimum and maximum values. Information regarding average age and gender was not
available for the healthy individuals in the MON cohort.

Supplementary Materials

Materials 1

List of genes that were included in MMset, COLset, HABset, MONset and BLDset.

Materials 2

Pathway enrichments for genes included in the MMset, COLset, HABset and MONset.

Materials 3

List of complex I related genes that were differentially expressed in COL and MON cohorts.