Single-cell dynamics of genome-nucleolus interactions captured by nucleolar laser

microdissection (NoLMseq)

Kaivalya Walavalkar[1], Shivani Gupta[1], Jelena Kresoja-Rakic[1], Mathieu Raingeval[1,2], Chiara Mungo[1,2],

Raffaella Santoro[1]

1. Department of Molecular Mechanisms of Disease, DMMD, University of Zurich, Zurich, CH-8057,

Switzerland

2. Molecular Life Science Program, Life Science Zurich Graduate School, University of Zurich, Zurich,

CH-8057, Switzerland

*Corresponding author:

Raffaella Santoro (raffaella.santoro@dmmd.uzh.ch)

Abstract

Gene position in the nuclear space plays an important role in gene regulation. This is exemplified by repressive chromatin domains frequently contacting nuclear lamina or nucleoli. The nucleolus undergoes structural changes in response to various cellular states, potentially impacting genome organization. However, how the 3D-genome organization responds to nucleolar states has remained underinvestigated due to the lack of methods able to identify nucleolar associated domains (NADs) in single cells and under nucleolar stress. To address this, we developed NoLMseq, a method combining laser-capture microdissection and DNA sequencing to map NADs in single cells. NoLMseq identified many unexplored features of chromosome organization around single nucleoli such as NAD heterogeneity among ESCs, culminating in two major populations with distinct chromatin states. The data also revealed that NADs prevalently contact nucleoli in a monoallelic manner and allelic nucleolar contact frequency sets gene expression and chromatin states. NoLMseq also revealed how chromosomes reorganise around nucleoli under nucleolar stress conditions, highlighting the importance of nucleolus integrity in genome organization and 3D-genome response to nucleolar stress. The results demonstrated that NoLMseq accurately measures chromosome contacts around single healthy and stressed nucleoli and it will be a critical tool to study NADs within biological populations and determine how the 3D-genome responds to nucleolar stress in healthy and disease states.

Introduction

The three-dimensional (3D) organization of chromosomes in the cell's nucleus plays an important role in the regulation of gene expression programs and cell fate. One important aspect of this regulation is the position of genes in the nuclear space. This is exemplified by the frequent location of repressive chromatin domains at the nuclear periphery (i.e., lamina associated domains, LADs) or around nucleoli (nucleolar associated domains, NADs) [1-3]. However, while LADs have extensively been studied in many cell types and in single cells, NADs have remained underinvestigated mainly due to technical limitations in identifying DNA sequences associating with a compartment lacking a membrane such as the nucleolus.

The nucleolus undergoes structural changes in response to various cellular states, with a potential impact on genome organization. The reason of these changes is that the nucleolus is the compartment where ribosome biogenesis takes place, a process responsible for the assembly of the translational machinery, the ribosome, and tightly regulated according to cell state [4]. Ribosome biogenesis is initiated in the nucleolus by the RNA polymerase I (Pol I)-driven transcription of hundreds of ribosomal RNA (rRNA) genes (rDNA) that generate 45S/47S pre-rRNA. These transcripts are then modified, processed, and assembled with ribosomal proteins (RPs) in the nucleolus, forming pre-ribosomal particles [5]. rRNA genes are distributed among different chromosomes; acrocentric chromosomes in human cells and, in mouse cells, and depending on strain, they can generally be found at chromosomes 12, 16, 18, and 19 (rDNA-chromosomes) and close to centromeric regions [3]. The nucleolus structure is very dynamic and promptly responds to external stimuli that affect ribosome biogenesis (i.e., size, number, or physico-chemical properties), such that highly proliferative cells have larger nucleoli compared to quiescent cells [6-9]. Accordingly, alterations in nucleolar activities and structure have been well documented under many cellular stress conditions and in several diseases, such as cancer, neurodegenerative disorders, and premature ageing [10]. For example, nucleolar stress is defined as a condition in which abnormalities in nucleolar structure and function driven by diverse cellular insults, such as nutrient starvation or DNA damage, ultimately leads to activation of stress signalling pathways that downregulate ribosome biogenesis, including rRNA gene transcription, and alter nucleolar structure [11,12]. However, whether these structural alterations might have an impact on the genome organization around nucleoli remains unexplored, mainly due to technical limitations of current methods to map NADs.

Currently, there are two major methods to identify NADs; the biochemical purification of nucleoli

followed by sequencing and, more recently, the Nucleolar-DamID that relies on the expression of an

engineered nucleolar histone H2B fused to Dam that serves to mark NADs with adenine methylation

(H2B-Dam-NoLS) (reviewed in [13]). Although based on different methodologies, both methods revealed

that NADs are composed of repressive chromatin domains [14-18]. However, both methods have also

some limitations. First, they cannot be applied to study how genome organization around nucleoli is

affected under nucleolar stress [19,20]. Indeed, under these conditions, nucleoli become frail and cannot

be efficiently biochemically purified. On the other side, the consequent downregulation of ribosome

biogenesis decreases protein translation, thereby affecting the expression of H2B-Dam-NoLS.

Moreover, the physico-chemical alterations of nucleoli upon rDNA downregulation causes the expulsion

of the H2B-Dam-NoLS from nucleoli similarly to other nucleolar components [18]. Finally, both methods

have only been applied in cell populations and consequentially they have provided an average nucleolar

contact frequency, which might not accurately reflect the structure of the genome around single nucleoli.

Thus, so far there is no methodology able to identify NADs under nucleolar stress conditions and in

single cells.

To overcome these challenges, we developed NoLMseq, a novel approach that combines laser-capture

microdissection (LCM) and DNA sequencing to identify NADs in single cells. LCM is a technique

traditionally used for isolating single cells or specific tissue regions[21,22] that we have expanded it for the

organelle-level isolation of single nucleoli. Traditional techniques to study genome organization

predominantly rely on either microscopy or DNA sequencing. NoLMseq synergistically integrates

strengths of both these approaches, enabling precise examination of the genome architecture

surrounding the nucleolus. The application of NoLMseq in mouse embryonic stem cells (ESCs) not only

confirmed the general repressive state of NADs but also provided novel insights of genome organization

around nucleoli that could not be detected in previous NAD studies that were based on methods that

could only be applied in cell populations. These findings included the detection of different levels of

NAD heterogeneity among ESCs that culminates in two distinct cell populations with distinct chromatin

states. NoLMseq also determined the remodelling of chromosomes around nucleoli during ESC

differentiation into neural progenitors with the identification of genes linked to neuronal pathways that

detach from nucleoli and become reactivated, indicating that the nucleolus is directly involved in the

establishment of repressive chromatin states. Intriguingly, although the majority of genes contacting

nucleoli are low expressing, we detected high nucleolar contact frequency of the highly expressing

ribosomal protein genes, suggesting a potential role of nucleolus proximity to regulate ribosome

biogenesis. Moreover, NoLMseq revealed that NADs prevalently contact nucleoli in a monoallelic

manner and that allelic nucleolar contact frequency correlates with gene expression and chromatin

states. Finally, we were able to determine the genome reorganization around nucleoli under nucleolar

stress conditions, a result that highlights the importance of nucleolus integrity in genome organization

and how the genome responds to nucleolar stress conditions. Together, these results not only provided

novel insights of how chromosomes are organized around nucleoli and their remodelling during cell fate

commitment and stress-induced nucleolar structural changes but also demonstrated that NoLMseq

accurately measures chromosome contacts around single healthy and "stressed" nucleoli. Thus,

NoLMseq will be a critical tool to study NADs within biological populations and determine how genome

organization responds to nucleolar stress in healthy and disease states.

Results

NoLMseq identifies NADs in single cells

To map NADs at a single nucleoli resolution, we combined laser-capture microdissection (LCM) of

nucleoli and DNA sequencing. We named this method nucleolus laser microdissection sequencing

(NoLMseq). We applied NoLMseq to mouse embryonic stem E14 cells (ESCs), thereby enabling

comparisons with NADs from ESC populations, which have recently been identified using Nucleolar-

DamID [18]. ESCs were grown on a thin polyester membrane that can be cut with a fine laser (**Fig. 1a,b**).

We visualized the nucleolus with brightfield microscopy and used LCM to cut it out and collected it in

an isolation cap of a microcentrifuge tube using the MMI CapSure technology. The DNA content of each

nucleolus was purified and amplified via quasilinear whole-genome amplification (WGA) [23]. To prevent

any loss of DNA, WGA was carried out in the same tube in which the microdissected nucleolus was

collected. Before proceeding with library preparation and DNA sequencing, the purity of nucleolar DNA

was assessed by PCR to determine the presence of rRNA genes, which are located within nucleoli, and

the absence of *Tuba1*, an active gene that previous studies have shown to not contact nucleoli[18]

(**Extended Data Fig. 1a**).

We generated 53 high-quality single nucleolus contacts maps from ESCs, with a median of $5 \cdot 10^5$

uniquely mapped reads per nucleolus (**Fig. 1c, Extended Data Figs. 1b and 2**). We binned the genome

into 100 kb segments and called NADs using SICER2 [24] at a 100kb resolution using genomic DNA as

a control. The identified reads over the reference genome showed a bimodal distribution, indicating that

loci are either in a "contact" or "no contact" state with nucleoli (**Fig. 1c, Extended Data Fig. 1b**). Since

the microdissection of nucleoli will inevitably include genomic sequences above and below nucleoli, we

analysed the data by calculating the nucleolar contact frequency (CF) that we defined as the proportion

of microdissected nucleoli containing a defined NAD sequence (**Fig. 1d**). We found that NADs from

rDNA-chromosomes have higher CF than chromosomes not containing rRNA genes, underscoring the

specificity of NoLMseq in identifying NADs. Moreover, chromosome X, which is active in the analysed

male ESCs, showed the lowest nucleolar CF. For all downstream analyses, we selected NADs with

CF> 20%, a value that corresponds to FDR 0.0009. The analysis of NAD coverage/chromosome of all

53 nucleoli further supported the high propensity of rDNA-chromosomes to contact nucleoli relative to

the other chromosomes (**Fig. 1e**). The average NAD coverage in single nucleoli was about 15%, much

lower than the 30% NAD coverage obtained from previous NAD studies in ESC population [16,18] (**Figs.**

1e,f). The data also showed that the average length of NADs of rDNA-chromosomes was 1.85 Mb and

significantly higher than the average NAD size from the rest of the chromosomes (1.30 Mb, **Fig. 1g**).

The accuracy of NoLMseq was also supported by finding >90% NADs overlapping with previous NAD

profiles from ESC populations using Nucleolar-DamID [18] and SPRITE method [25] (**Fig. 1h**). To further

validate the data, we used an orthogonal approach by computing the CF of NADs from NoLMseq

genomic data and their distance to Fibrillarin marked nucleoli obtained by seqFISH+ microscopy [26] (**Fig.**

1i). We observed that nucleolar CFs measured by NoLMseq were inversely proportional to the distance

of NADs to the nucleolar marker Fibrillarin, indicating that NoLMseq accurately measures genomic

distance to nucleoli.

To further support the efficacy of NoLMseq, we analysed known NAD features of the identified NADs.

It is well known that centromeres tend to be positioned close to the nucleolus in various model systems.

Since in mouse cells all chromosomes are acrocentric, we measured the average NAD coverage at

centromere-proximal regions at the 5'-quintile portion of all chromosomes and found that it increases

with nucleolar CF, further supporting the accuracy of NoLMseq to identify sequences in contact with

nucleoli (**Fig. 2a**). Consistent with previous results in ESC populations obtained with Nucleolar-DamID

[18], NADs identified by NoLMseq were significantly depleted of active histone marks relative to

sequences within the active A compartment, showed a significant enrichment for the repressive

H3K9me2 but not for the other repressive marks H3K27me3 and H3K9me3, and displayed low gene

expression levels (**Fig. 2b,c**). Although NADs generally appear to have a repressive chromatin and

gene expression state, we could also find cases where highly expressed genes have high nucleolar

CFs. In particular, we found high nucleolar CFs for ribosomal protein (RP) genes, which encode proteins

for ribosome assembly in nucleoli and are transcribed by RNA Pol II, and tRNA genes, which are

transcribed by Pol III (**Fig. 2d**). In contrast, snoRNA genes, which express RNAs playing an important

role in rRNA modifications in nucleoli, did not show any preferential location to nucleoli. The nucleolar

contacts of RP and tRNA genes was not due to their location at rDNA-chromosomes since most of them

were located at chromosomes non containing rRNA genes (92% and 90%, respectively; **Extended**

Data Fig. 3). These results suggest a potential role of nucleolus proximity for the regulation of RP

genes, and consequently ribosome biogenesis, as well as tRNA, thereby influencing protein translation

in general (**Fig. 5b**). Finally, these data also suggested that not all NADs are necessarily

transcriptionally repressed.

NADs have been previously characterized in two categories; NAD-only (or Type-1) which correspond

to sequences contacting nucleoli but not the nuclear lamina (NL) and NAD/LAD (or Type-2) which are

regions that can be located both at nucleoli and NL [13,16-18]. Since this classification of NADs was

obtained from ESC population, it has remained elusive whether NAD/LAD regions contact both the

nucleoli as well as the NL in the same or different cells. The analysis of single nucleoli data revealed

that NAD-only sequences have higher CF values than NAD/LAD, suggesting that NAD/LAD domains

can contact the nucleolus in one cell and the NL in another cell, but rarely both nucleolus and NL in the

same cell (**Fig. 2e**). We also observed an inverse correlation of NAD-only CF values and H3K27ac

content, indicating that strong NADs generally exhibit low levels of active chromatin (**Fig. 2f**). Similarly,

NAD-only domains with high nucleolar CF have high H3K9me2 levels (**Fig. 2g**). However, while

NAD/LAD regions showed significantly lower gene expression levels relative to genes within the active

A compartment, the expression of genes within NAD-only regions was not significantly reduced, albeit

there was a tendency to be low expressed when located at regions with the highest nucleolar CF (**Fig.**

2h). These results are consistent with the data described above showing that some active genes can

also contact nucleoli.

ESCs are composed of two populations with distinct nucleolar contacts

NADs obtained from single nucleoli allowed to study the degree of cell heterogeneity in genome organization. We calculated the NAD coverage for each chromosome in single cells and performed hierarchical clustering (**Fig. 3a**). We found two main ESC populations that differ according to the level of nucleolar contacts among chromosomes. The major differences between these two groups were on the level of rDNA-chromosome contacts with the nucleolus, which showed a mutually exclusive relationship. rDNA-chromosomes 12 and 19 had high levels of nucleolar contacts in population 1 whereas they were almost absent in nucleoli of population 2 (**Fig. 3b**). In contrast, rDNA-chromosomes 16 and 18 displayed higher NAD coverage in population 2 than in population 1. In the case of chromosomes not bearing rRNA genes, chromosomes 7, 11, 15, and 2 showed more nucleolar contacts in population 2 than in population 1 whereas chromosomes 3 and 5 preferentially contact nucleoli of population 1. Remarkably, the analysis of histone modifications revealed that NADs enriched in population 1 were significantly depleted of active histone marks and enriched in H3K9me2 relative to NADs of population 2, suggesting different chromatin states between these two ESC populations (**Fig. 3d**). Thus, the data revealed heterogeneity in the ESC genome structure and chromatin configuration that depend on the genome organization around nucleoli.

Detachment of NADs upon differentiation into NPCs results in gene activation

Next, we asked whether the composition of genomic contacts in single nucleoli could change upon differentiation of ESCs into neural progenitors (NPCs). As in the case of ESCs, in NPCs NADs from rDNA- chromosomes consistently exhibited higher nucleolar CF than regions from other chromosomes (**Fig. 4a**). However, NPCs displayed fewer NADs than ESCs, with a large proportion (75%) being the same NADs as found in ESCs (**Fig. 4b-d**). Moreover, we observed more variability in nucleolus association in ESCs than in NPCs as indicated by the Yule's Q coefficient, which provides a measure of similarity between single-cell NADs (**Fig. 4e**). The NAD detection in single NPC nucleoli obtained with NoLMseq is consistent with previous studies performed in NPC populations showing that the architecture of chromosomes surrounding the nucleolus is in a more compact and rigid form whereas in ESCs it appears more flexible and variable [18]. Moreover, these data are consistent with previous reports indicating that ESCs harbour a more open and dynamic chromatin than differentiated cells [27,28].

Next, we investigated whether genes contacting nucleoli in ESCs but detaching from nucleoli in NPCs

(ESC specific NADs, ESCsp-NADs) could change their expression state in NPCs. We defined ESCsp-

NADs those NADs with nucleolar CF > 25% in ESCs and < 10% in NPCs. RNAseq data analysis

revealed that ESCsp-NADs are less expressed in ESCs than in NPCs, suggesting that the detachment

from nucleoli facilitates gene activation (**Fig. 4f,g**). Moreover, the absolute fold change in gene

expression in NPCs compared to ESCs was higher for the upregulated genes, with a large fraction

(60%) not expressed in ESCs (**Fig. 4h,i**). Accordingly, GO terms for these upregulated genes were

significantly linked to developmental processes, in particularly nervous system development, whereas

downregulated genes were mainly implicated in metabolic processes (**Extended Data Fig. 4**). These

results indicate that the detachment of genes from nucleoli during ESC differentiation unlocks them for

activation, further supporting a regulatory role of the nucleolus for the establishment or maintenance of

repressive chromatin states.

Nucleolar integrity is required for genomic contacts with nucleoli

Nucleolus structure is highly dynamic and promptly responds to external stimuli that affect ribosome

biogenesis and nucleolar structure. For example, downregulation of rDNA transcription by treatment

with low doses of Actinomycin D (ActD) induces a spatial reorganization of the nucleolar structure,

including the migration of rRNA genes to the nucleolar periphery, forming the so-called nucleolar caps,

and the release of the granular component (GC) protein NPM1 from nucleoli (**Fig. 5a**) [29]. However, until

now, it could not be investigated whether these structural changes can impact the organization of the

genome around nucleoli mainly due to the technical limitations of the current technologies to map NADs

under conditions in which nucleolar integrity is affected. Since nucleoli in ESCs treated with ActD

(ESCs+ActD) are still detectable with brightfield microscopy, we reasoned that NoLMseq could be

applied to determine how nucleolus integrity affects chromosome contacts with nucleoli. Compared to

untreated cells, the average NAD coverage in ESC+ActD significantly decreased from 15% to 10%

(**Fig. 5b,c, Extended Data Fig. 5**). Although NoLMseq revealed a substantial loss of nucleolar contacts

across all chromosomes, the majority of NADs retained upon ActD treatment were from the rDNA-

chromosomes (**Fig. 5d**). We validated this loss of contacts with nucleoli in ESCs+ActD by quantitative

DNA-FISH, using probes targeting four NADs at chromosomes 1, 2, 5, and 19 that were previously

identified and validated by Nucleolar-DamID [18] (**Fig. 5e-g**). These are strong NADs since they contact

nucleoli in 70-95% of ESCs (**Fig. 5f**). The DNA-FISH probe for the rDNA-chromosome 19 targeted a

NAD sequence proximal to the rDNA locus. Accordingly, this NAD remained anchored to nucleoli upon

ActD treatment, an expected result since rRNA genes remain close to nucleoli by forming nucleolar

caps (**Fig. 5f**). In contrast, DNA-FISH probes targeting NADs at chromosomes 1, 2, and 5 showed a

significant detachment of NADs from the nucleoli upon ActD treatment, indicating that nucleolar integrity

is crucial for NAD association with nucleoli and supporting the results obtained with NoLMseq. Because

of the retainment of rDNA-chromosomes to nucleoli in ESC+ActD, we reasoned that regions from rDNA-

chromosomes still able to contact nucleoli in ESC+ActD should correspond to centromere-proximal

sequences, which are linearly located close to rRNA genes. Accordingly, NAD coverage of centromere-

proximal sequences of rDNA-chromosomes was higher than the corresponding NAD coverage in

untreated ESCs, suggesting a reorganization of rDNA-chromosomes structure around nucleoli (**Fig.

5e**). In contrast, NAD coverage of centromere-proximal sequences of chromosomes not bearing rDNA

repeats showed a significant reduction compared to NADs of untreated cells, further indicating that the

anchoring of these sequences to nucleoli depends on nucleoli integrity. All these results indicate that

nucleolar integrity is an important determinant for chromosome organization around nucleoli. Moreover,

they demonstrate that NoLMseq is a unique and only genomic method for studying the role of nucleolus

in genome organization under nucleolar stress conditions that affect nucleolar integrity.

Genomic contacts with nucleoli are predominantly mono-allelic

Recent data have started to detect allelic asymmetry in gene expression and 3D-genome organization

[30,31]. So far, however, all studies on NADs were performed using cells whose maternal and paternal

chromosomes could not be distinguished from each other, as in the case of the E14 ESC line that

derives from a fully inbred homozygous mouse strain. Consequently, the information on NAD coverage

has been based on the assumption that both alleles contact nucleoli. To determine the allelic distribution

of NADs in single nucleoli, we applied NoLMseq to the male mouse F123-ESC line that was derived

from the F1 generation of two fully inbred homozygous mouse strains: *Mus musculus castaneus* (CAST,

paternal chromosomes) and *Mus musculus domesticus* S129S4/SvJae (S129, maternal chromosomes)

[32]. The F123 DNA sequence has been well characterized and contains high density of single nucleotide

variants (SNVs), with an average of 1 SNV every 124 nucleotides across autosomes [33]. We analysed

NAD profiles from 50 microdissected nucleoli of F123-ESCs by aligning them against the F123 genome

to identify allele-specific NADs (i.e., NAD_{S129} and NAD_{CAST}) (**Fig. 6a**). The vast majority (93%) of total

NADs identified in F123-ESCs corresponded to NADs identified in E14-ESCs, indicating that genomic

contacts with nucleoli are generally conserved between cell lines with the same developmental stage

but from different strains (**Extended Data Fig. 6a**). NADs of F123-ESCs showed a slightly lower

genome coverage (11.3%, calculated to haploid genome) relative to 15% NAD coverage measured in

E14-ESCs (**Fig. 6B**). Surprisingly, we found that a large fraction of NADs (86%) are mono-allelic (**Fig.**

6a-c). Monoallelic NADs were equally distributed among S129 and CAST chromosomes (**Extended**

Data Fig. 6b). We validated these results by DNA-FISH for two strong NADs, one located at

chromosome 5 and another located at the rDNA-chromosome 19 and close to the rDNA locus (**Fig.**

6d). The NAD of chromosome 5 showed monoallelic contacts in almost all the cells. Remarkably, even

the NAD of chromosome 19 located close to the rDNA locus showed mono-allelic contacts with nucleoli

in 50% of the cells, supporting the results of NoLMseq. This prevalent mono-allelic NAD distribution

resulted in a much lower NAD coverage per cell (5.6%, 297 Mbp) compared to the previously assumed

bi-allelic distribution (**Fig. 6e**). To determine whether mono- and bi-allelic NADs reflect differences in

chromatin modifications, we performed ChIPseq analysis for H3K9me2, H3K27ac, and H3K4me3 in

F123-ESCs (**Fig. 6f**). We found that bi-allelic NADs are in a more repressed chromatin states than

mono-allelic NADs since they display reduced levels of H3K27ac and H3K4me3 and higher levels of

H3K9me2, the latter the repressive histone mark characterizing NADs in ESCs [18]. Similarly, RNAseq

analyses showed a significantly lower expression of genes located at bi-allelic NADs (**Fig. 6g**). Among

genes located at bi-allelic NADs, we found cell type specific transcription factors, such as *Gata6* and

TAF4b, as well as genes linked to metabolic pathways and neuroactive ligand-receptor interactions

(**Extended Data Fig. 6c**). These results further indicate that high-frequency contact with nucleoli, even

at the allelic levels, results in a more repressive state.

Next, we asked whether the mono-allelic distribution of NADs might be parent specific. We applied

stringent criteria to assign parent specific NADs, considering, for example, that a specific NAD_{S129}

should have >25% nucleolar CF with the S129 genome and <10% nucleolar CF with the CAST genome

(**Fig. 6h**). However, very few NADs (1% on both alleles) showed a parent specific distribution. This

indicates that, generally, in one cell it is the paternal allele contacting nucleoli while in another cell it is

the maternal allele (**Fig. 6h**).

Discussion

We described NoLMseq, a method able to measure genome organization around nucleoli in single cells. NoLMseq synergistically integrates the strengths of both microscopy and DNA sequence technologies, enabling precise examination of the genome architecture surrounding the nucleolus. Relative to previous methods, NoLMseq not only detects NADs in single nucleoli but it can also be applied to study genome organization around nucleoli under conditions of nucleolar stress, which profoundly alter nucleolar structure [6-9] and that could not be examined with current methods, making elusive how genome structure around nucleoli responds to nucleolar stress. The data revealed a profound reorganization of the genome around "stressed" nucleoli, such an increase in nucleolar contacts of centromere-proximal sequences of rDNA-chromosomes while the rest of the chromosomes lose their contact. The genomic data obtained by NoLMseq are also consistent with recent image analyses showing that perturbation of nucleolus structure upon downregulation of the RNA polymerase I subunit RPA194 reduced centromere-nucleolus interactions [34]. These results indicate that nucleolar structure and integrity plays an important role in genome organization and future studies will address alterations in chromosomal interactions with nucleoli under pathological stress conditions impacting ribosome biogenesis and how these perturbations are caused and their functional significance. The results also showed that NADs are very dynamic and NoLMseq represents an unique technique able to capture NADs at the present state rather than averages of the past and present states as in the case of DamID as recently discussed [35].

The detection of NADs in single nucleoli using NoLMseq has proven to be highly precise compared to previous methods and capable of providing unexplored insights in genome organization. For example, NoLMseq determined that active genes, such as ribosomal protein genes and tRNA genes, also have high frequency contacts with nucleoli. Interestingly, previous imaging analyses in living yeast cells reported the proximity of some RP genes to nucleoli [36], suggesting that the nucleolus could act as a compartment for the regulation of the expression of ribosomal components. The presence of some active genes at nucleoli is also consistent with a recent study showing that the localization of genes at the repressive B compartment or proximal to nucleoli does not necessarily preclude transcription [37].

The accuracy of NoLMseq in detecting NADs relative to cell population studies is also shown by the identification of ESCsp-NADs that contain genes activated in NPCs, which could not be detected in previous cell population studies [18], further indicating that the nucleolus serves in general as a repressive

compartment. Because of the single nucleolus detection, NoLMseq revealed that ESCs display a certain

heterogeneity in genomic contacts, with nucleoli showing two major cell populations with distinct

chromatin features. Whether the presence of such populations, one more repressed than the other one,

might reflect differences in gene expression, pluripotency states, or other functions remains to be

investigated. These results are also consistent with recent data showing heterogeneity of nuclear

organization between individual single ESCs [38]. Finally, the application of NoLMseq in the hybrid F123-

ESC line showed that genomic contacts with nucleoli are mainly monoallelic. This result clearly provided

a more accurate value of NAD coverage, which was previously calculated based on either a haploid

genome or by assuming that all NADs are bi-allelic in the diploid genome. The average NAD coverage

5.6% (i.e., ca. 297 Mbp) per cell obtained in the F123-ESC line is much lower than the 30% NAD

coverage measured in ESC population studies using both purified nucleoli and Nucleolar-DamID

methods [16,18]. Although this calculated NAD coverage value does not include repetitive sequences such

as major and minor satellites, which are known to contact nucleoli but are not present in the reference

genome, it clearly represents a more realistic scenario than to have one third of chromosomes

contacting nucleoli. Moreover, the data showed that bi-allelic NADs are in a more repressive state than

monoallelic NADs, further supporting the efficacy of NoLMseq in identifying NADs and the nucleolus

acting as a repressive compartment.

Together, the results demonstrate that NoLMseq not only accurately measures chromosome contacts

around single nucleoli but also provides novel insights into genome organization around nucleoli that

could not be detected by previous studies. We predict that NoLMseq will not only be a critical tool to

study genome organization around the nucleolus in biological populations but also to study how the 3D-

genome organization responds to nucleolar stress in disease states and understand the functional

consequence of these alterations.

Materials and Methods:

Cell lines

129 mouse embryonic stem cells (E14 line) were cultured in serum medium containing Dulbecco's modified Eagle's medium + Glutamax (Life Technologies), 15% FCS (Life Technologies; Cat no. 10270106 FBS South American), 1× MEM NEAA (Life Technologies), 100 μM β-mercaptoethanol, recombinant leukemia inhibitory factor, LIF (Polygene, 1000 U/ml), 1× penicillin/streptomycin (Life Technologies). ESCs were seeded at a density of 50,000 cells/cm2 in culture dishes (Corning CellBIND surface) coated with 0.1% gelatin without a feeder layer. Propagation of cells was carried out every 2 days using enzymatic cell dissociation. ESCs were treated with Actinomycin D (10ng/ml) for 24h before downstream analyses.

Neural progenitor cells were generated from ESCs, according to a previously established protocol [39]. In brief, differentiation used a suspension-based embryoid bodies formation (Bacteriological Petri Dishes, Bio-one with vents, Greiner). The neural differentiation media (DMEM, 10% fetal calf serum, 1× MEM NEAA, 2 mM Pen/Strep, β-mercaptoethanol, and sodium pyruvate) was filtered through 0.22 μm filters and stored at 4 °C. During the 8-day differentiation procedure, media was exchanged every 2 days. In the last 4 days of differentiation, the media was supplemented with 2 μM retinoic acid to generate neural precursors that are Pax-6- positive radial glial cells.

F123-ESCs (mouse male, hybrid cell line, F1 S129/Jae and Cast mouse cross) [32] were cultured on a layer of mitotically inactivated feeder murine embryonic fibroblasts under standard conditions (DMEM, supplemented with 15 % KSR, 1x Glutamax, 10 mM non-essential amino acids, 50 μM beta-mercaptoethanol, 1,000 U/ml LIF). Media was changed every 24h. Before harvesting, mESCs were passaged onto feeder-free ESGRO-gelatin (EMD Millipore SF008). coated plates for at least 2 passages to remove feeder cells. Cells were harvested after approximately 48 h at 70 – 80 % confluency. F123-ESCs were a gift from Bing Ren, University of California, San Diego.

NoLMseq

For laser-microdissection, cells were grown on ESGRO-gelatin coated and UV-irradiated PET membrane slides (1.4um thick, RNase-free, MMI, 50102) for 24h. Cells were washed twice with 1X PBS and fixed with EM-grade freshly-depolymerised 3% PFA in 1X PBS for 30mins. The slides were then placed in 70% ethanol for 1min and air dried. The slides were then washed with filtered ultra-pure H_2O

(3x 1 min each) and air dried. Single nucleoli were laser microdissected with the MMI Laser Capture

Microdissection Nikon microscope (Molecular Machines & Industries) into single 0.2 ml tubes

(transparent MMI caps, 50208). Nucleoli were identified by bright-field imaging. A 0.2ml cap filled with

transparent adhesive material was lowered onto the cells. A laser was used to cut the PET membrane

surrounding each nucleolus and the cut PET membrane section along with the single nucleolus was

lifted off by the sticky cap. Each single nucleolus was collected in a separate cap and stored for further

processing.

Whole-genome amplification (WGA) was carried out using the MALBAC single cell WGA kit (YK001A)

following manufacturer's recommendations with minor modifications. The entire WGA reaction was

carried out in the same 0.2ml tube with a single nucleolus. Another tube, with an empty section was

used as a negative control whereas 30 pg of whole cell DNA were used as a positive control for the

WGA reaction. The cell lysis reaction was first carried out by using a heated lid at 50 °C for 50 mins

and then samples were collected by briefly spinning and proceeding with the cell lysis reaction again at

50 °C for 50 mins, followed by inactivation at 80 °C for 10mins. The reaction is then cooled at 4 °C. The

lysis step is followed by a pre-amplification step which involves the quasilinear amplification using

random primers at 20 °C progressively with 10 °C steps till 70 °C. After cooling, the samples were

subjected to a final amplification step by 20 cycles of PCR.

WGA-amplified DNA was purified using a Qiagen MinElute PCR Purification Kit and eluted in 140 μl

ultrapure library-grade H_2O. The amplified nucleolar DNA was checked on a 2% agarose gel to select

positive samples showing smears corresponding to amplified DNA similar to the positive control. The

positive samples were further checked for the presence of rDNA and the absence of Tubulin gene by

PCR on an agarose gel. The samples which passed this quality control check were then used for library

preparation and sequencing. 120 μl of the reaction was first sonicated in split-cap tubes by Covaris

E220 to obtain fragments around 300-500nt in length (Peak power- 75, Duty factor- 20, cycle/bursts-

1000, duration- 35s). The samples were then purified and concentrated with AMPure XP beads

(Beckman Coulter, A63881) and used for Illumina library using the NEBNext ultra II kit (NEB E7600).

Briefly, NoLMseq samples (10-100 ng) were end-repaired and polyadenylated before the ligation of

Illumina compatible adapters. The adapters contain the index for multiplexing. Library concentrations

were estimated using a Qubit® 4 fluorometer (Thermo Fisher Scientific) and the 4200 TapeStation

system and libraries were pooled together in batches averaging 40 samples. Each library pool was

sequenced in single end 100bp in one lane of Novaseq6000 to get an average 10 million reads per

library.

For data analysis, NoLM-seq reads were aligned to the mouse mm10 reference genome using Bowtie2

(version 2.5.0; [40]) with default parameters. PCR duplicates were removed using samtools [41]. Positive

NAD bins were called using SICER2 [42], using the sequenced gDNA file as the control file and the

parameters w=100000, g=100000, fdr=0.0001. Bigwig files were created from the bam files using

bamCoverage from deeptools [43]. For F123-ESCs, N-masked genome was created using the mm10

reference genome and the Variant Call *Format* (VCF) file from the mouse genome project containing

the SNP information for all strains. The SNPsplit package [44] was used to read the VCF file and extract

the SNP information for the S129/SvJae and CAST strains and N-mask for the mm10 genome. The

reads were aligned using Bowtie2 to this N-masked reference genome using default parameters.

SNPsplit was then used to tag the SNPs for S129/SvJae and CAST alleles in the bam file and sort the

haplotype-specific bam files for both the alleles. bamCoverage and Sicer2 positive NAD calling was

performed independently on the allele-specific bam files. Integrative Genome Viewer (IGV, version

2.15.4) [45] was used to visualize and extract representative single nucleolus NoLMseq bigwig and bed

tracks.

Nucleolar contact frequency was calculated at a 100kb resolution. We defined as NADs sequences with

a > 20% nucleolar contact frequency that correspond to a false discovery rate (FDR) 0.0009 or 0.09%,

meaning that more than 99% of called NADs are truly NADs. This was calculated using the cumulative

distribution function (CDF) $P(X \le k) = \sum i = 0k(in)pi(1-p)n-1$ where "n" is the number of single

nucleoli samples, "p" is the probability that a false-positive region is present in the sample, "k" is the

contact frequency threshold. For the hierarchical clustering of NADs, NAD chromosome coverages in

single nucleoli were used for clustering single nucleoli based on divisive clustering (Diana) using the

"cluster" package (version 2.1.6) in R.

RNAseq

Total RNA was purified with TRIzol reagent (Life Technologies). The quality of the isolated RNA was

determined with a Qubit® 4 fluorometer (Life Technologies, California, USA) and 4200 TapeStation

system (Agilent, Santa Clara, California, USA). Only those samples with a 260 nm/280 nm ratio between

1.8–2.1 and a 28S/18S ratio within 1.5-2 were further processed. The TruSeq Stranded mRNA (Illumina,

Inc, California, USA) was used in the succeeding steps. Briefly, total RNA samples (100–1000 ng) were polyA enriched and then reverse- transcribed into double-stranded cDNA. The cDNA samples were fragmented, end-repaired and adenylated before ligation of TruSeq adapters containing unique dual indices (UDI) for multiplexing. Fragments containing TruSeq adapters on both ends were selectively enriched with PCR. The quality and quantity of the enriched libraries were validated using Qubit® 4 fluorometer. The product is a smear with an average fragment size of approximately 260 bp. Libraries were normalized to 10 nM in Tris-Cl 10 mM, pH8.5 with 0.1% Tween 20. The NovaSeq X (Illumina, Inc, California, USA) was used for cluster generation and sequencing according to standard protocol. Sequencing was paired end at 2 × 150 bp.

The quality of the reads was checked by FastQC, a quality control tool for high throughput sequence data [46]. The quality of the reads was increased by applying: a) SortMeRNA [47] (version 2.1) tool to filter ribosomal RNA; b) Trimmomatic [48] (version 0.40) software package to trim the sorted (a) reads. N-masked genome was created using the mm10 reference genome and the VCF file from the mouse genome project containing the SNP information for all strains. The SNPsplit package [44] was used to read in the VCF file and extract the SNP information for the S129/SvJae and CAST strains and N-mask the mm10 genome. The sorted (a), trimmed (b) reads were mapped against the Nmasked mouse genome (mm10) using the default parameters of the STAR (Spliced Transcripts Alignment to a Reference, version 2.6, [49]). SNPsplit was then used to tag the SNPs for S129/SvJae and CAST alleles in the bam file and sort the haplotype-specific bam files for both the alleles. For each gene, exon coverage was calculated using a custom pipeline and then normalized in reads per kilobase per million (RPKM) [50].

ChIPseq

ChIP analysis was performed as previously described [51]. Briefly, 1% formaldehyde was added to cultured cells to cross-link proteins to DNA. Isolated nuclei were then lysed with lysis buffer (50 mM Tris-HCl, pH 8.1, 10 mM EDTA, pH 8, 1% SDS, 1X protease inhibitor complete EDTA-free cocktail, Roche). Nuclei were sonicated using a Bioruptor ultrasonic cell disruptor (Diagenode) to shear genomic DNA to an average fragment size of 200 bp. 20 µg of chromatin was diluted to a total volume of 500 µl with ChIP buffer (16.7 mM Tris-HCl, pH 8.1, 167 mM NaCl, 1.2 mM EDTA, 0.01% SDS, 1.1% Triton X-100) and incubated overnight with the ChIP-grade antibodies against H3K9me2, H3K9me3,

H3K27me3, H3K4me1, H3K4me3 and H3K27ac. After washing, bound chromatin was eluted with the

elution buffer (1% SDS, 100 mM $NaHCO_3$). Upon proteinase K digestion (50 °C for 3 h) and reversion

of cross- linking (65 °C, overnight), DNA was purified with phenol/chloroform, and ethanol precipitated.

The quality and quantity of the isolated DNA were determined with a Qubit® 4 Fluorometer (Life

Technologies, California, USA) and the 4200 TapeStation system. Briefly, ChIP samples (1 ng) were

end-repaired and polyadenylated before the ligation of Illumina compatible adapters. The adapters

contain the index for multiplexing. The quality and quantity of the enriched libraries were validated using

Qubit® 4 Fluorometer and the 4200 TapeStation system. The libraries were pooled to 10 nM in Tris-Cl

10 mM, pH8.5 with 0.1% Tween 20. Sequencing was performed on the NovaSeq 6000 for 100bp single

end reads (Illumina, Inc, California, USA).

For F123-ESCs, N-masked genome was created using the mm10 reference genome and the Variant

Call Format (VCF) file from the mouse genome project containing the SNP information for all strains.

The SNPsplit package [44] was used to read the VCF file and extract the SNP information for the

S129/SvJae and CAST strains and N-mask for the mm10 genome. ChIPseq reads were aligned to the

N-masked mm10 reference genome with Bowtie2 (version 2.5.0; (version 2.5.0; [40]) using default

parameters. PCR duplicates were removed using samtools [41]. SNPsplit was then used to tag the SNPs

for S129/SvJae and CAST alleles in the bam file and sort the haplotype-specific bam files for both the

alleles. Bigwig files were created from the bam files using bamCoverage from deeptools [43] using default

parameters. Integrative Genome Viewer (IGV, version 2.15.4) [45] was used to visualize and extract

representative ChIPseq tracks.

Gene Ontology (GO) enrichment:

GO enrichment analysis of genes was performed with DAVID 6.8 [52]. All genes in *Mus musculus* were

used as the background.

DNA-FISH

DNA-FISH probes for chromosomes 1, 2, 5, and 19 were generated as previously described [18]. The

probes were obtained with oligopaint libraries that were constructed the PaintSHOP interface created

by the Beliveau lab and were ordered from CustomArray/Genscript in the 12 K Oligopool format. Each

library contains a universal primer pair used to amplify all the probes in the library, followed by a specific

primer pair hooked to the 40-46-mer genomic sequences, for a total probe of around 124-130-mers.

Oligopaint libraries were produced by emulsion PCR from the pool, followed by a "two-step PCR" and

lambda exonuclease as described before [18,53]. Specifically, the emulsion PCR with the universal primers

allowed amplifying all pool of probes in the library. The "two-step PCR" led to the addition of a tail to the

specific probes bound to the Alexa Fluor 488 fluorochrome. All oligonucleotides were purchased from

Integrated DNA Technologies (IDT, Leuven, Belgium).

DNA-FISH was adapted from the protocol of Cavalli's lab [54]. Cells fixed with 4% PFA for 10 min at room

temperature 24h after seeding. Cells were washed three times with 1X PBS and then permeabilized

with 0.5%Triton-X100/PBS for 10 min on ice, followed by 10 min at room temperature. After three

additional washes with 1X PBS, cells were incubated with 0.1 M HCl for 10 min at room temperature,

washed twice with 2X SCCT, and twice with 50% Formamide-2X SCCT for 20min, firstly at room

temperature and then at 60°C. Probe mixture contained around 45pmol of Oligopaint probe, 0,8µL of

ribonuclease A (ThermoScientific, 10 mg/mL), and FISH hybridization buffer for a total mixture volume

of 20uL, which was directly added on the slide. Cell DNA was denatured at 83 °C for 8 min, and

hybridization was performed in a humid dark chamber overnight at 42 °C. Cells were washed with 2X

SCCT for 15 min once at 60°C and once at room temepature, once with 0.2X SCC for 10 min, twice

with 1X PBS for 2 min, and three times with 1X PBT for 2 min. Slides were then incubated for 1 h at

room temperature with blocking solution in a dark humid chamber, and overnight at 4 °C with primary

antibody against Nucleolin. Cells were washed four times with 1X PBT at increasing incubation times

(1 × 2 min, 1 × 3 min, 2 × 5 min) and then incubated for 2 h at room temperature with goat anti-mouse

IgG (H + L) Alexa Fluor 546 in a dark humid chamber. Slides were washed four times with 1X PBT at

increasing incubation times (1 × 2 min, 1 × 3 min, 2 × 5 min), three times with 1X PBS for 2 min. Then,

slides were mounted with Vectashield DAPI mounting media (Vector, H-1200) and, after drying, stored

at 4 °C.

DNA FISH/IF samples were imaged using a Zeiss LSM 980 Airyscan, with a z-stack collected for each

channel (step size, 0.3 um), using the oil objective Plan-Apochromat/1.40. Images were processed by

ImageJ (version 2.14.0/1.54F). The individual cells were identified by Hoechst/DAPI staining and cells

containing signal for DNA-FISH channel were identified manually on the corresponding fluorescent

channel. Distance between the DNA-FISH signal and the nucleolar marker immunofluorescence signal

was calculated using ImageJ (version 2.14.0/1.54F) and used to count the number of foci contacting

nucleolus and the number of cells with at least one contact with the nucleolus.

Published datasets

550   Published datasets used in the study were from: H3K27ac-, H3K4me1-, H3K27me3-ChIP-seq [39]

551   (GSE72164); H3K4me3- and H3K9me3-ChIPseq [55] (GSE23943); H3K9me2-ChIPseq [56] (GSE77420);

552   A/B compartments [57]; Nucleolar DamID and RNAseq in ESCs and NPCs [18] (GSE150822); Lamin-

553   DamID [58] (GSE17051); SPRITE [25] (GSE114242); seqFISH+ [26].

Acknowledgements

556   This work was supported by the ERC grant (ERC-AdG-787074-NucleolusChromatin), Swiss National

557   Science Foundation (31003B-201268), and EMBO Postdoctoral Fellowship (ALTF 209-2021 to K.W).

558   We thank Catherine Aquino and the Functional Genomic Center Zurich for the assistance in

559   sequencing. We thank Ana Pombo for her technical advises on laser-microdissection and library

560   preparation and Bing Ren for the F123-ESC line.

Author contributions

563   K.W. and J.K-R. established the NoLMseq. K.W. performed NoLMseq in all the described conditions,

564   ChIPseq and RNAseq in F123-ESCs and all data analyses of NADs. S.G. and M.R. performed DNA-

565   FISH analyses. C.M. performed IF analyses. K.W. and R.S. wrote the manuscript. R.S conceived and

566   supervised the project.

Declaration of interests

569   The authors declare no competing interests.

Figure legends

Figure 1

NoLMseq identifies NADs in single cells

a. Scheme representing the nucleolar laser capture microdissection (NoLMseq).

b. Representative images of ESCs before and after nucleolar laser capture microdissection.

c. Nucleolar contacts maps from 53 single nucleoli for chromosomes 6, 8, and 19. Black bars indicates NADs.

d. Cumulative histogram of genome-wide nucleolar contact frequency (CF) values with respect to their location at chromosomes containing rRNA genes (rDNA-chromosomes), without (w/o rDNA), and X and Y chromosomes.

e. Coverage of NADs with CF > 20% on each chromosome in single nucleoli. Dotted grey line indicates the mean NAD coverage in single cells. Black lines indicate the mean on each chromosome.

f. Violin plot showing NAD coverage in single cells. Red dashed line indicates the median.

g. Size of NADs located at rDNA-chromosomes and chromosomes not bearing rRNA genes (w/o rDNA). Red and blue lines indicate medians, and the dotted lines indicate quartiles. Statistical significance (*P*-values) was calculated using unpaired t-test (**** < 0.0001).

h. Coverage of NADs identified by NoLMseq relative to NADs identified in ESC populations with the Nucleolar-DamID [18] and SPRITE [25] method.

i. Genome-wide comparison of DNA seqFISH+ distance to exterior of nucleoli marked with Fibrillarin (µm) [26] and NoLMseq NAD contact frequency for 3713 paired genomic regions. Tukey boxplot where box limits represent the 25th and 75th percentiles and the black line represents the median. Statistical significance (*P*-values) was calculated using unpaired t-test (** < 0.01; **** < 0.0001).

Figure 2

NADs identified by NoLMseq correspond to repressive chromatin domains

a. NAD coverage on the centromere-proximal regions (5' -quintile) as a function of nucleolar contact frequency.

b. Levels of active histone marks (H3K27ac, H3K4me1, H3K4me3) and repressive histone marks (H3K9me2, H3K9me3, H3K27me3) at NADs relative to genomic regions located at the active A compartment (A comp.). Values are shown as average RPKM. Tukey boxplot where box limits represent

the 25th and 75th percentiles. The horizontal line within the boxes represents the median. Statistical

significance (*P*-values) was calculated using the unpaired two-tailed t test (**<0.01, ****<0.0001, ns:

nonsignificant).

c. Expression values (RPKM) of genes within the active A compartment (A Comp.) and NADs. Tukey

boxplot where box limits represent the 25th and 75th percentiles. The horizontal line within the boxes

represents the median. Statistical significance (*P*-values) was calculated using the unpaired two-tailed

t test(****<0.0001).

d. Gene densities of ribosomal protein genes (red), tRNA genes (black), snoRNA genes (pink) and all

genes (grey) as a function of nuclear CF.

e. Left panel shows nucleolar contact frequency for NAD-only and NAD/LAD sequences. Right panel

depicts how NAD-only and NAD/LAD contact nucleoli in single cells.

f,g,h. Levels of H3K27ac (**f**), H3K9me2 (**g**), and gene expression (**h**) at NAD-only and NAD/LAD as a

function of nucleolar CF relative to the active A compartment (A comp.). at the active A compartment

(A Comp.) and at NAD-only and NAD/LAD relative to nucleolar CF. Values are shown as average

RPKM. Tukey boxplot where box limits represent the 25th and 75th percentiles. The horizontal line

within the boxes represents the median. Statistical significance (*P*-values) was calculated using the

unpaired two-tailed t test (**<0.01, ***< 0.001, ****<0.0001, ns: nonsignificant).

Figure 3

ESCs showed two populations with distinct nucleolar contacts

a. Hierarchical clustering of nucleoli according to their NAD coverage/chromosome

b. NAD coverage at the indicated chromosomes in the top quartile NADs of the two ESC populations.

c. Scheme representing the contact with nucleoli of the rDNA-chromosomes with the nucleoli in the two

ESC populations.

d. Levels of active histone marks (H3K4me1, H3K4me3, H3K27ac) and repressive histone marks

(H3K9me2, H3K9me3, H3K27me3) at NADs enriched in ESC population (Pop) 1 or 2. Values are shown

as average RPKM. Tukey boxplot where box limits represent the 25th and 75th percentiles. The

horizontal line within the boxes represents the median. Statistical significance (*P*-values) was calculated

using the unpaired two-tailed t test (*<0.05, ****<0.0001, ns: nonsignificant).

Figure 4

NADs from single nucleoli of neural progenitors

a. Cumulative histogram of genome-wide of nucleolar contact frequency measured from 23 microdissected nucleoli with respect to their location at chromosomes containing rRNA genes (rDNA-chromosomes), without (w/o rDNA), and X and Y chromosomes in NPCs.

b. Coverage of NADs with CF > 20% on each chromosome in single nucleoli of NPCs. Dotted grey line indicates the average NAD coverage in single cells of NPCs. Black lines indicate the mean on each chromosome.

c. Violin plots showing NAD coverage in single ESCs and NPCs. Data from ESCs were also shown in **Figure 1f**. Red dashed lines indicate the median and black dotted lines indicate the quartiles.

d. Venn diagram showing the distribution of NADs in ESCs and NPCs.

e. Violin plots showing the distribution of Yule's Q values as a measure of NAD cell-cell similarity in ESCs and NPCs. Red dashed lines indicate the median and black dotted lines indicate the quartiles.

f. Expression levels of genes located at ESC_{sp}-NADs in ESCs and NPCs. Tukey boxplot where box limits represent the 25th and 75th percentiles. The horizontal line within the boxes represents the median. Statistical significance (*P*-values) was calculated using the paired two-tailed t test (****<0.0001).

g. Scatter plot showing the expression levels of genes at ESC_{sp}-NADs in ESCs and NPCs. Expression of genes significantly (P < 0.05) upregulated (blue) and downregulated (red) are shown.

h. Plot showing the absolute $\log_2$ fold changes of upregulated and downregulated genes at ESC_{sp}-NADs in NPCs. Black lines indicate the median. Statistical significance (*P*-values) was calculated using the unpaired two-tailed t test (****<0.0001).

i. Proportion of low expressing genes (< 0.1 RPKM) in ESCs that are located at ESC_{sp}-NADs and are up- or downregulated in NPCs.

Figure 5

Nucleolar integrity is required for genomic contacts with nucleoli

a. Left panel represents alterations in nucleolus structure upon treatment with Actinomycin D (ActD), showing the re-organization of rRNA genes, which form nucleolar caps. Right panel shows immunofluorescence images of the nucleolar markers UBF and NPM1. Scale bar is 9 μm.

b. Histogram showing the average NAD coverage/chromosome from single nucleoli of untreated ESCs

(blue) and 34 microdissected nucleoli from ESCs treated with ActD (grey).

c. Violin plots showing NAD coverage in single untreated ESCs (blue) and ESCs treated with ActD

(grey). Data from ESCs were also shown in **Figure 1F**. Red dashed lines indicate the median and black

dotted lines indicate the quartiles. Statistical significance (P-value) was calculated using the unpaired

two-tailed t test (****<0.0001).

d. Distribution of NADs in ESCs treated with ActD as a function of nucleolar CF across rDNA-

chromosomes, chromosomes without (w/o) rDNA, and NADs lost relative to untreated ESCs.

e. Representation of the DNA-FISH probes targeting NADs at chromosomes 1, 2, 5, and 19, the latter

containing rRNA genes at the 5' end, close to the centromeric region. The corresponding NADs profiles

from Nucleolar-DamID [18] are shown. iNAD: genomic domains not interacting with nucleoli.

f. Representative immuno-FISH images for NAD at chromosome 2. Nucleolin serves as nucleolar

marker. Scale bar is 5 μm.

g. Box plots represent the quantifications of immuno-DNA-FISH analyses showing the percentage of

cells displaying NADs contacting nucleoli. Error bars represent s.d. Statistical significance (*P*-values)

from three independent experiments was calculated using Mann-Whitney test (** < 0.01; *** < 0.001;

ns: non-significant).

h. NAD coverage at the centromere-proximal regions (the first 5'-quintile) between rDNA-chromosomes

and chromosomes without (w/o) rDNA. rDNA on untreated ESCs and ESCs treated with ActD. Red

dashed lines indicate the median and black dotted lines indicate the quartiles. Statistical significance

(P-value) was calculated using the unpaired two-tailed t test (**<0.01).

Figure 6

Genomic contacts with nucleoli are mono-allelic

a. NoLMseq tracks displaying NAD sequences from four microdissected single nucleoli from hybrid

F123-ESCs (S129/CAST) on the total mm10 genome (black), maternal S129 genome (pink) and

paternal CAST genome (blue). Bars below the tracks indicate NADs called on respective genomes.

b. NAD coverage from single nucleoli calculated over the haploid genome. Median and quartiles are

shown. Statistical significance (P-value) was calculated using the unpaired two-tailed t test

(****<0.0001, ns: non-significant).

c. Proportion of bi-allelic and mono-allelic NADs in single nucleoli. Black dashed lines indicate median

and black dotted lines indicate quartiles. Statistical significance (P-value) was calculated using the

unpaired two-tailed t test (****<0.0001).

d. Left panel shows DNA-FISH images of a NAD located at chromosome 19. Right panel is the

quantification of bi-allelic and mono-allelic contacts with nucleoli. Scale bar is 5 μm. Error bars indicate

s.d.

e. NAD coverage from single nucleoli over the diploid genome. Black dashed line indicates the median

and black dotted lines indicate the quartiles.

f,g. Levels of H3K9me2, H3K27ac, and H3K4me3 (**F**) and gene expression (**G**) at bi-allelic and mono-

allelic NADs. Values are shown as average RPKM. Violin plots where black dashed lines indicate the

median and black dotted lines indicate the quartiles. Statistical significance (*P*-values) was calculated

using the unpaired two-tailed t test (**<0.01; ***<0.001; ****<0.0001).

h. Pie chart showing the amounts of paternal (NAD_{CAST}) and maternal (NAD_{S129}) NADs.

i. Model showing that mono-allelic distribution of NADs at nucleoli of single cells is generally not parent

specific.

Extended Data figure legends

Extended Data Figure S1

NoLMseq identifies NADs in single cells

a. Gel electrophoresis showing whole-genome amplification (WGA) of isolated nucleoli (left panel) and the PCR amplification for rRNA genes and *Tuba1* sequences.

b. NoLMseq tracks from 6 microdissected nucleoli at chromosomes 6, 8, at 19. Black bars below the tracks indicate NADs called in respective single nucleoli.

Extended Data Figure S2

NAD contact frequency for each chromosome

Nucleolus contact frequency for the 53 ESC nucleoli along the length of each chromosome. rDNA-chromosomes are depicted with an orange line.

Extended Data Figure S3

Ribosomal protein and tRNA genes show high nucleolar contact frequency

Distribution of ribosomal protein and tRNA genes at rDNA-chromosomes and chromosomes not containing rRNA genes (other chromosomes).

Extended Data Figure S4

NADs from single nucleoli of neural progenitors

Gene ontology (GO) terms for upregulated and downregulated genes at ESC_{sp}-NADs in NPCs.

Extended Data Figure S5

Nucleolar integrity is required for genomic contacts with nucleoli

Coverage of NADs at each chromosome of single nucleoli from ESCs treated without and with Actinomycin D (ActD). Dotted grey line indicates the average NAD coverage in single cells. Black lines indicate the mean on each chromosome.

Extended Data Figure S6

Genomic contacts with nucleoli are mono-allelic, but not parent specific.

a. Pie chart showing the proportion of NADs identified in F123-ESCs that have also been found in E14-

ESCs.

b. Mono-allelic distribution of 129 and CAST genome around nucleoli of single cells. Black dashed lines

indicate median and black dotted lines indicate quartiles. Statistical significance (P-value) was

calculated using the unpaired two-tailed t test (ns: non-significant).

c. Gene ontology (GO) terms of genes located on bi-allelic NADs.

References

Willemin, A., Szabo, D. & Pombo, A. Epigenetic regulatory layers in the 3D nucleus. *Mol Cell* **84**, 415-428 (2024). https://doi.org:10.1016/j.molcel.2023.12.032

Misteli, T. The Self-Organizing Genome: Principles of Genome Architecture and Function. *Cell* **183**, 28-45 (2020). https://doi.org:10.1016/j.cell.2020.09.014

Bersaglieri, C. & Santoro, R. Genome Organization in and around the Nucleolus. *Cells* **8(6)**, 579 (2019). https://doi.org:10.3390/cells8060579

Bassler, J. & Hurt, E. Eukaryotic Ribosome Assembly. *Annu Rev Biochem* **88**, 281-306 (2019). https://doi.org:10.1146/annurev-biochem-013118-110817

Pelletier, J., Thomas, G. & Volarevic, S. Ribosome biogenesis in cancer: new players and therapeutic avenues. *Nat Rev Cancer* **18**, 51-63 (2018). https://doi.org:10.1038/nrc.2017.104

Derenzini, M. *et al.* Nucleolar size indicates the rapidity of cell proliferation in cancer tissues. *The Journal of pathology* **191**, 181-186 (2000). https://doi.org:10.1002/(SICI)1096-9896(200006)191:2<181::AID-PATH607>3.0.CO;2-V

Boisvert, F. M., van Koningsbruggen, S., Navascues, J. & Lamond, A. I. The multifunctional nucleolus. *Nat Rev Mol Cell Biol* **8**, 574-585 (2007). https://doi.org:nrm2184 [pii] 10.1038/nrm2184

Goodfellow, S. J. & Zomerdijk, J. C. Basic mechanisms in RNA polymerase I transcription of the ribosomal RNA genes. *Subcell Biochem* **61**, 211-236 (2013). https://doi.org:10.1007/978-94-007-4525-4_10

Ruggero, D. Revisiting the nucleolus: from marker to dynamic integrator of cancer signaling. *Sci Signal* **5**, pe38 (2012). https://doi.org:10.1126/scisignal.2003477

Corman, A., Sirozh, O., Lafarga, V. & Fernandez-Capetillo, O. Targeting the nucleolus as a therapeutic strategy in human disease. *Trends Biochem Sci* **48**, 274-287 (2023). https://doi.org:10.1016/j.tibs.2022.09.006

Yang, K., Yang, J. & Yi, J. Nucleolar Stress: hallmarks, sensing mechanism and diseases. *Cell Stress* **2**, 125-140 (2018). https://doi.org:10.15698/cst2018.06.139

Boukoura, S. & Larsen, D. H. Nucleolar organization and ribosomal DNA stability in response to DNA damage. *Curr Opin Cell Biol* **89**, 102380 (2024). https://doi.org:10.1016/j.ceb.2024.102380

Bersaglieri, C. & Santoro, R. Methods for mapping 3D-chromosome architecture around nucleoli. *Curr Opin Cell Biol* **81**, 102171 (2023). https://doi.org:10.1016/j.ceb.2023.102171

Nemeth, A. *et al.* Initial genomics of the human nucleolus. *PLoS Genet* **6**, e1000889 (2010). https://doi.org:10.1371/journal.pgen.1000889

Koningsbruggen, S. v. *et al.* High-Resolution Whole-Genome Sequencing Reveals That Specific Chromatin Domains from Most Human Chromosomes Associate with Nucleoli. *Molecular Biology of the Cell* **21**, 3735-3748 (2010). https://doi.org:10.1091/mbc.e10-06-0508

Bizhanova, A., Yan, A., Yu, J., Zhu, L. J. & Kaufman, P. D. Distinct features of nucleolus-associated domains in mouse embryonic stem cells. *Chromosoma* (2020). https://doi.org:10.1007/s00412-020-00734-9

Vertii, A. *et al.* Two contrasting classes of nucleolus-associated domains in mouse fibroblast heterochromatin. *Genome Res* **29**, 1235-1249 (2019). https://doi.org:10.1101/gr.247072.118

Bersaglieri, C. *et al.* Genome-wide maps of nucleolus interactions reveal distinct layers of repressive chromatin domains. *Nat Commun* **13**, 1483 (2022). https://doi.org:10.1038/s41467-022-29146-2

Shore, D. & Albert, B. Ribosome biogenesis and the cellular energy economy. *Curr Biol* **32**, R611-R617 (2022). https://doi.org:10.1016/j.cub.2022.04.083

Lempiainen, H. & Shore, D. Growth control and ribosome biogenesis. *Curr Opin Cell Biol* **21**, 855-863 (2009). https://doi.org:10.1016/j.ceb.2009.09.002

21   Datta, S. *et al.* Laser capture microdissection: Big data from small samples. *Histol Histopathol* **30**, 1255-1269
(2015). https://doi.org:10.14670/HH-11-622

22   Beagrie, R. A. *et al.* Complex multi-enhancer contacts captured by genome architecture mapping. *Nature* **543**,
519-524 (2017). https://doi.org/10.1038/nature21411

23   Baslan, T. *et al.* Genome-wide copy number analysis of single cells. *Nat Protoc* **7**, 1024-1041 (2012).
https://doi.org/10.1038/nprot.2012.039

24   Zang, C. *et al.* A clustering approach for identification of enriched domains from histone modification ChIP-Seq
data. *Bioinformatics* **25**, 1952-1958 (2009). https://doi.org/10.1093/bioinformatics/btp340

25   Quinodoz, S. A. *et al.* Higher-Order Inter-chromosomal Hubs Shape 3D Genome Organization in the Nucleus.
Cell **174**, 744-757 e724 (2018). https://doi.org/10.1016/j.cell.2018.05.024

26   Takei, Y. *et al.* Integrated spatial genomics reveals global architecture of single nuclei. *Nature* **590**, 344-350
(2021). https://doi.org/10.1038/s41586-020-03126-2

27   Meshorer, E. *et al.* Hyperdynamic plasticity of chromatin proteins in pluripotent embryonic stem cells. *Dev Cell*
10, 105-116 (2006). https://doi.org:S1534-5807(05)00421-1 [pii] 10.1016/j.devcel.2005.10.017

28   Gaspar-Maia, A., Alajem, A., Meshorer, E. & Ramalho-Santos, M. Open chromatin in pluripotency and
reprogramming. *Nature reviews. Molecular cell biology* **12**, 36-47 (2011). https://doi.org/10.1038/nrm3036

29   Reynolds, R. C., Montgomery, P. O. & Hughes, B. Nucleolar "Caps" Produced by Actinomycin D. *Cancer Res*
24, 1269-1277 (1964).

30   Sun, Y. *et al.* MSL2 ensures biallelic gene expression in mammals. *Nature* **624**, 173-181 (2023).
https://doi.org/10.1038/s41586-023-06781-3

31   Irastorza-Azcarate, I. *et al.* Extensive folding variability between homologous chromosomes in mammalian cells.
bioRxiv (2024). https://doi.org/10.1101/2024.05.08.591087

32   Gribnau, J., Hochedlinger, K., Hata, K., Li, E. & Jaenisch, R. Asynchronous replication timing of imprinted loci
is independent of DNA methylation, but consistent with differential subnuclear localization. *Genes Dev* **17**, 759-
773 (2003). https://doi.org/10.1101/gad.1059603

33   Markowski, J. *et al.* GAMIBHEAR: whole-genome haplotype reconstruction from Genome Architecture Mapping
data. *Bioinformatics* **37**, 3128-3135 (2021). https://doi.org/10.1093/bioinformatics/btab238

34   Wang, C., Ma, H., Baserga, S. J., Pederson, T. & Huang, S. Nucleolar structure connects with global nuclear
organization. *Mol Biol Cell* **34**, ar114 (2023). https://doi.org/10.1091/mbc.E23-02-0062

35   Kefalopoulou, S. *et al.* Time-resolved and multifactorial profiling in single cells resolves the order of
heterochromatin formation events during X-chromosome inactivation. *bioRxiv*, 2023.2012.2015.571749 (2023).
https://doi.org/10.1101/2023.12.15.571749

36   Berger, A. B. *et al.* High-resolution statistical mapping reveals gene territories in live yeast. *Nat Methods* **5**,
1031-1037 (2008). https://doi.org/10.1038/nmeth.1266

37   Goronzy, I. N. *et al.* Simultaneous mapping of 3D structure and nascent RNAs argues against nuclear
compartments that preclude transcription. *Cell Rep* **41**, 111730 (2022).
https://doi.org/10.1016/j.celrep.2022.111730

38   Arrastia, M. V. *et al.* Single-cell measurement of higher-order 3D genome organization with scSPRITE. *Nat
Biotechnol* **40**, 64-73 (2022). https://doi.org/10.1038/s41587-021-00998-1

39   Joshi, O. *et al.* Dynamic Reorganization of Extremely Long-Range Promoter-Promoter Interactions between
Two States of Pluripotency. *Cell Stem Cell* **17**, 748-757 (2015). https://doi.org/10.1016/j.stem.2015.11.010

40   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-359 (2012).
https://doi.org/10.1038/nmeth.1923

41 Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10** (2021).
https://doi.org:10.1093/gigascience/giab008

42 Zang, C., Wang, Y. & Peng, W. RECOGNICER: A coarse-graining approach for identifying broad domains from
ChIP-seq data. *Quant Biol* **8**, 359-368 (2020). https://doi.org:10.1007/s40484-020-0225-2

43 Ramirez, F., Dundar, F., Diehl, S., Gruning, B. A. & Manke, T. deepTools: a flexible platform for exploring deep-
sequencing data. *Nucleic Acids Res* **42**, W187-191 (2014). https://doi.org:10.1093/nar/gku365

44 Krueger, F. & Andrews, S. R. SNPsplit: Allele-specific splitting of alignments between genomes with known
SNP genotypes. *F1000Res* **5**, 1479 (2016). https://doi.org:10.12688/f1000research.9037.2

45 Robinson, J. T. *et al.* Integrative genomics viewer. *Nature Biotechnology* (2011).
https://doi.org:10.1038/nbt0111-24

46 Andrews, S. FastQC: a quality control tool for high throughput sequence data. (2010).

47 Kopylova, E. N., L.; Touzet, H. SortMeRNA: Fast and accurate filtering of ribosomal RNAs in metatranscriptomic
data. *Bioinformatics* (2012).

48 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*
30, 2114-2120 (2014). https://doi.org:10.1093/bioinformatics/btu170

49 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
https://doi.org:10.1093/bioinformatics/bts635

50 Mortazavi, A., Williams, B. A., McCue, K., Schaeffer, L. & Wold, B. Mapping and quantifying mammalian
transcriptomes by RNA-Seq. *Nat Methods* **5**, 621-628 (2008). https://doi.org:10.1038/nmeth.1226

51 Lerra, L. *et al.* An RNA-dependent and phase-separated active subnuclear compartment safeguards repressive
chromatin domains. *Mol Cell* **84**, 1667-1683 e1610 (2024). https://doi.org:10.1016/j.molcel.2024.03.015

52 Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional annotation of gene
lists (2021 update). *Nucleic Acids Res* **50**, W216-W221 (2022). https://doi.org:10.1093/nar/gkac194

53 Beliveau, B. J. *et al.* Single-molecule super-resolution imaging of chromosomes and in situ haplotype
visualization using Oligopaint FISH probes. *Nat Commun* **6**, 7147 (2015). https://doi.org:10.1038/ncomms8147

54 Szabo, Q., Cavalli, G. & Bantignies, F. in *Capturing Chromosome Conformation: Methods and Protocols* (eds
Beatrice Bodega & Chiara Lanzuolo) 221-237 (Springer US, 2021).

55 Marks, H. *et al.* The transcriptional and epigenomic foundations of ground state pluripotency. *Cell* **149**, 590-604
(2012). https://doi.org:10.1016/j.cell.2012.03.026

56 von Meyenn, F. *et al.* Impairment of DNA Methylation Maintenance Is the Main Cause of Global Demethylation
in Naive Embryonic Stem Cells. *Mol Cell* **62**, 848-861 (2016). https://doi.org:10.1016/j.molcel.2016.04.025

57 Bonev, B. *et al.* Multiscale 3D Genome Rewiring during Mouse Neural Development. *Cell* **171**, 557-572.e524
(2017). https://doi.org:10.1016/j.cell.2017.09.043

58 Peric-Hupkes, D. *et al.* Molecular maps of the reorganization of genome-nuclear lamina interactions during
differentiation. *Mol Cell* **38**, 603-613 (2010). https://doi.org:S1097-2765(10)00321-7
[pii]10.1016/j.molcel.2010.03.016
883

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Extended data Figure 1

a

WGA

b

Extended data Figure 2

Extended data Figure 3

Extended data Figure 4

GO upregulated ESCsp-NAD in NPCs

	Single-multicellular organism process
	System development
	Single-organism cellular process
	Multicellular organism development
	Neurogenesis
	Nervous system development
	Anatomical structure morphogenesis
	Anatomical structure development
	Neuron differentiation
	Generation of neurons

P value

GO downregulated ESCsp-NADs in NPCs

	RNA processing
	Nitrogen compound metabolic process
	Metabolic process
	Cellular nitrogen compound metabolic process
	Ribosome biogenesis
	Organic substance metabolic process
	Primary metabolic process
	Cellular metabolic process
	Ncrna processing
	Ribonucleoprotein complex biogenesis

P value

Extended data Figure 5

Extended data Figure 6

a

NADs only in F123 (3.2%)

NADs in both F123 and E14 (96.8%)

b

Monoallelic NAD distribution (%)

ns

S129   CAST

c

GO genes at bi-allelic NADs

Acetylcholine receptor signaling pathway

Signal transduction involved in cellular response to ammonium ion

Cellular response to acetylcholine

Response to acetylcholine

Single-organism cellular process

Localization

Response to ammonium ion

Regulation of neurotransmitter receptor activity

Cellular response to ammonium ion

Establishment of localization

10^{-8} 10^{-7} 10^{-6} 10^{-5} 10^{-4} 10^{-3} 10^{-2} 10^{-1} 10^{0}

P value