Emergence of alternative stable states in microbial communities undergoing horizontal

gene transfer

Juken Hong[1], Wenzhi Xue[1], Teng Wang[1]✉

[1]Key Laboratory of Quantitative Synthetic Biology, Shenzhen Institute of Synthetic Biology,

Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen 518055,

China.

✉Corresponding author. Email: t.wang1@siat.ac.cn.

Abstract

Microbial communities living in the same environment often display alternative stable states,

each characterized by a unique composition of species. Understanding the origin and determinants

of microbiome multistability has broad implications in environments, human health and

microbiome engineering. However, despite its conceptual importance, how multistability emerges

in complex communities remains largely unknown. Here, we focused on the role of horizontal

gene transfer (HGT), one important aspect mostly overlooked in previous studies, on the stability

landscape of microbial populations. Combining mathematical modeling and numerical simulations,

we demonstrate that, when mobile genes only affect bacterial growth rates, increasing HGT rate

in general promote the emergence of alternative stable states in complex microbiota. We further

extend our analysis to scenarios where HGT changes interspecies interactions, microbial

communities are subjected to strong environmental selections and microbes live in

metacommunities consisting of multiple local habitats. We also discuss the role of different

mechanisms, including interspecies interaction strength, the growth rate effects of MGEs, MGE

epistasis and microbial death rates in shaping the multistability of microbial communities

undergoing HGT. These results reveal how different dynamic processes collectively shape

community multistability and diversity. Our results provide key insights for the predictive control

and engineering of complex microbiota.

Introduction

Microbial communities living in similar environments often display different community types, each characterized by a unique composition of species[1-4]. Changes in environmental factors, fluctuations of species abundances, or transient invasions can trigger abrupt and irreversible transitions between two types[5-7]. Such transitions, also known as regime shifts, lead to the dramatic alterations of microbiome functions and are tightly associated with host diseases, soil fertility, marine biogeochemical flux, or bioproduction yield[8-12]. Understanding the origin of different community types has broad implications in environments, engineering and health[1,3].

Multistability, a phenomenon where multiple stable states coexist for the same set of system parameters, is one of the most important mechanisms behind alternative community types and has been widely observed in a variety of natural or experimental microbiomes[1,4,7,13-17]. Under perturbations, a multistable system can be driven away from a former stable state, cross the tipping point and reach a new steady state. The population can persist in the new state even after the perturbations are withdrawn, exhibiting a behavior of 'hysteresis'[3,13]. Despite its conceptual importance, the determinants underlying microbiota multistability have been poorly understood due to the intrinsic complexity of microbial interactions[14].

Understanding the origin and the underlying determinants of multistability in complex microbiota is an important challenge in microbial ecology and evolution. In host-associated microbiota, regime shifts between alternative stable states can trigger the critical transitions between healthy microbiome and dysbiosis[8,18]. Unravelling how different biological processes drive community multistability is important for the predictive control of complex microbiota, and can inspire novel treatments to modulate the communities towards the desired stable states or away from the unhealthy states.

Mathematical models are indispensable to decipher the emergent properties of complex populations[19]. Different ecological models have been developed to analyze the origin of multistability in microbial communities. For instance, based on the generalized Lotka-Volterra (LV)

model, Gonze D *et al.* showed that multistability could emerge in communities where different species mutually inhibited each other[1]. Dubinkina V *et al.* applied a consumer-resource model in which microbes competed for different metabolites, and showed that multistability required microbial species to have different stoichiometries of essential nutrients[14]. These works provide critical insights on the determinants of community multistability, yet the roles of many other mechanisms remain to be understood. Chief among these is horizontal gene transfer (HGT), a process where microbes share mobile genetic elements (MGEs) with their neighbors[20].

Indeed, HGT is prevalent in microbial world and mobile genes occupy a substantial proportion in microbial gene pool[21-23]. For instance, among the closely related microbial isolates in the same human body site, HGT occurs in more than 40% of the species pairs[24]. In the draft genomes of human gut microbes, tens of thousands of genes were mobilizable[22]. Depending on the encoded traits, MGEs shape microbial growth rates, host adaptability and even social interactions between microbes[25]. The dynamic gain and loss of MGEs are also a major source of evolutionary innovations of prokaryotic genomes[26]. However, despite its functional and ecological importance, the interplay between genetic exchange and community multistability has not been rigorously examined in previous studies and a general conclusion remains lacking.

In this work, we focus on communities of competing microbes, a model system for microbiome multistability[1,14,27]. By combining mathematical modelling and numerical simulations, we demonstrated that the horizontal transfer of plasmids, a major type of MGEs, could promote the emergence of alternative stable states which otherwise would not exist. We further extended our analysis to scenarios where HGT changed interspecies interactions, where microbial communities were subjected to strong environmental selections and where microbes lived in metacommunities consisting of multiple local habitats. We also analyzed the role of different mechanisms, including interspecies interaction strength, the growth rate effects of MGEs, MGE epistasis and microbial death rates in shaping the multistability of microbial communities. These results created a comprehensive framework to understand how different dynamic processes,

including but not limited to HGT rates, collectively shaped community multistability and diversity.

Our results provide valuable insights for microbial ecology, health and microbiome engineering.

Results

The role of HGT on bistability of two-species populations

To illustrate the basic concepts, let's first consider a population of two competing species.

Without HGT, the community dynamics can be captured by the classic LV model that accounts for

species growth rates (μ_1 and μ_2), interspecies interaction strengths (γ_1 and γ_2) and the dilution

or death rate (D). When all the parameters are given, one can simulate the system dynamics from

the initial abundances of the two species. The system will reach an equilibrium, characterized by

the steady-state abundances of the two species. Bistability means the coexistence of two distinct

stable states (Fig. 1a)[1,14]. Depending on the initial abundances, either one of the two species can

dominate the community. With bistability, even when the model parameters are not altered, a

change in initial species abundances is sufficient to drive the system to a different stable state (Fig.

1a). In contrast, if monostable, the population will always rest at the same steady state regardless

of the initial composition (Fig. 1a). Given the simplicity of this model, the condition of the system

being bistable can be analytically solved as $\gamma_1 > \phi_2/\phi_1$ and $\gamma_2 > \phi_1/\phi_2$ where $\phi_1 = \frac{\mu_1 - D}{\mu_1}$

and $\phi_2 = \frac{\mu_2 - D}{\mu_2}$ (see Methods for more details). This condition predicted the thresholds on the

strengths of interspecies interactions in order for the system to be bistable.

To analyze the effect of HGT on bistability, we applied a mathematical model that we

previously developed to describe the conjugative transfer of plasmids between competing species[28].

While in classic LV model μ_1 and μ_2 are constants, our model dissected species growth rates

into two components: the basal growth rates (μ_1^0 and μ_2^0) determined by chromosomal genes, and

the growth rate effects (λ_1 and λ_2) of plasmids. The two components were assumed to combine

multiplicatively: $\mu_1 = \mu_1^0(1 + \lambda_1)$, $\mu_2 = \mu_2^0(1 + \lambda_2)$. HGT creates subpopulations (denoted as

p_1 and p_2) within each species that acquire the mobilizable genes from its competitor. The

dynamics of p_1 and p_2 were governed by cell growth, plasmid transfer (the rate denoted as η)

and plasmid loss (the rate denoted as κ). The gain and loss of p_1 and p_2 in turn lead to the

temporal change of the effective growth rates (μ_1^e and μ_2^e) of each species: $\mu_1^e = \mu_1(1 + \lambda_2 \frac{p_1}{s_1})$,

$\mu_2^e = \mu_2(1 + \lambda_1 \frac{p_2}{s_2})$ (see Methods for more details). Therefore, plasmid transfer allows the two

competitors to partially exchange their growth benefits or disadvantages. Here, we assumed that

the species internal inhibition was independent of the acquired MGEs. We sought to understand

whether the process of HGT would reshape the stability landscape of the system.

After HGT being introduced, analytical solution becomes infeasible due to the increase of

modeling complexity. Instead, we determined whether the system was bistable by numerical

simulations with randomized initial compositions[27]. In general, the resolution of this approach

increases with the number of randomizations. Without loss of generality, here we randomly

initialized the species abundances for 200 times between 0 and 1 following uniform distributions

(Fig. S1a). Then we simulated the population dynamics for each initialization until steady state

and obtained their species abundances. A system is called monostable if all populations starting

from different compositions converge into the same state. In contrast, if two attractors exist, the

system is called bistable (Fig. 1a). We calculated the number of stable states for a wide range of

μ_1 and μ_2 values and drew the phase diagram (Fig. 1b). Whether it is easy for the system to

achieve bistability is reflected by the total area of bistability region on the phase diagram. Our

numerical results suggested that increasing HGT enlarged the region, creating bistability in many

systems which would otherwise be monostable (Fig. 1b-d). Here, the number of initializations was

not critical for the conclusion (Fig. S1a). The results were not fundamentally changed when the

interspecies interaction strengths or gene transfer rates were altered (Fig. S1b-d), suggesting the

robustness of our conclusion.

Our results suggested that increasing HGT made it more feasible for the system to become

bistable. However, this should not be taken as an assertion that the system will always be bistable

when HGT exists (Fig. 1c and d). For two species with large growth rate difference, the system

might remain monostable when gene transfer rate increases (Fig. 1c). In this case, the strong

competitor will always win despite the gene flow. The effect of HGT in promoting bistability is

more applicable between species with smaller growth rate difference (Fig. 1b and d).

Previous studies showed that the cell death rate determines the bistability of two interacting

species[29]. To evaluate the impact of death rate D on the interplay between HGT and system

bistability, we performed additional analysis by calculating the bistability probability under

different D values (Fig. S2). Our results suggested that varying death rate indeed changed the

bistability of the system. When the death rate equaled zero, mobile genetic elements that only

modified growth rates would have no effects on population bistability. These results highlighted

the importance of added death rate in driving multistability.

The strength of interspecies interaction also shaped the community bistability. By calculating

the bistability probability under different values of γ (Fig. S3), we showed that while changing

γ didn't fundamentally alter our conclusion, the influence of HGT on population bistability

became weak when interspecies interaction strength got smaller than 1. These results suggested

that the role of HGT was more significant in population with strong competitions.

HGT shapes the stability landscape of multi-species communities

The modeling framework can be readily generalized to more complex communities consisting

of multiple competing species by taking into account the gene transfer between every pair of

species. Consider a community of m species. Let s_i be the abundance of the i-th species. γ_{ij}

represents the interspecies interaction strength that the i-th species imposes on the j-th species

($i,j = 1,2,\ldots,m$). η_{jki} is the transfer rate of the s_j-originated plasmid from species k to

species i. κ_{ij} is the loss rate of the j-th mobile genetic element in the i-th species. The growth

rate effect of the j-th plasmid on the i-th species is denoted as λ_{ij}. p_{ij} describes the abundance

of the subpopulation in the i-th species that carries the plasmid from the j-th species. The

effective growth rate of the i-th species can be calculated based on all the plasmids that it carries:

$\mu_i^e = \mu_i \prod_{j \neq i}(1 + \lambda_{ij} \frac{p_{ij}}{s_i})$. The population dynamics can then be simulated by $m + m^2$ equations

that describe the temporal change of s_i and p_{ij}, respectively (see Methods for more details).

To understand how changing gene transfer rate affects the stability landscape of complex

microbiota, we estimated the number of alternative stable states by randomly initializing the

abundance of each species 500 times between 0 and 1 following uniform distributions. For each

initialization, we simulated the population dynamics for 2000 hours till the systems reached

equilibrium and calculated the steady-state abundances of different species (Fig. S4). Limit cycles

were not observed in our simulations (Fig. S5).

The calculate the absolute number of stable states, we grouped the steady states into different

attractors using a given threshold. Steady states with their Euclidean distances smaller than the

threshold belonged to the same group (Fig. S6). However, the estimation of this number can be

biased by the intrinsic limitations of the numerical approach. For instance, the limited number of

initializations can underestimate this number. In contrast, the inability of a system to reach

equilibrium within the limited time window of simulations can cause its overestimation. To

overcome such limitations, we developed an alternative metric χ to measure the multistability of

a system by borrowing the concept of 'entropy'. Let φ be the number of stable states and x_i be

the relative size of each domain of attraction ($\sum_i x_i = 1$, $i = 1,2,...,\varphi$). χ is calculated as $\chi =$

$exp\,(-\sum_i x_i log x_i)$, which accounts for the absolute number of stable states as well as the size of

each basin. χ equals 1 in monostable systems. χ gets larger when more stable states emerge or

the distribution of the basin sizes become more even. Numerical tests suggested that χ was robust

to the variations of distance threshold, simulation timespan or number of initializations (Fig. S7).

As shown in Fig. 2, when HGT is absent, the number of stable states was usually small in a

multi-species population. Increasing gene transfer rate reshaped the stability landscape of a

microbiome by creating additional stable states (Fig. 2a and b). Each stable state is characterized

by a different dominant species (Fig. 2a). The number of stable states is positively associated with

HGT rate (Fig. 2c, Fig. S8 and S9). This prediction is generally applicable to communities

consisting of different number of species (Fig. S8 and S9). For communities of two species,

bistability is not possible when γ_{ij}'s are smaller than 1. However, for communities of multiple

species, multistability becomes possible even if all γ_{ij} values are smaller than 1 (Fig. S10). The

effects of HGT on multistability held for different strengths of interspecies interactions (Fig. S10).

Initializing the p_{ij} subpopulations with non-zero abundances didn't change the conclusion, either

(Fig. S11).

Here we assumed the interactions among different species were homogeneous. To evaluate

whether this assumption was critical, we extended our analysis by considering heterogenous

interaction strengths in multispecies communities (Fig. S12). In particular, we randomly sampled

γ_{ij} values from uniform distributions. Our results suggested the mean value and variance of γ_{ij}

played a role in shaping multistability. The effects of HGT on community multistability becomes

stronger when the mean value of γ_{ij} gets larger than 1 and the variance of γ_{ij} is small (Fig. S12).

Nevertheless, the heterogeneous distribution of γ_{ij} didn't fundamentally change our conclusion.

MGEs can drive large growth rate differences when they encode adaptative traits like

antibiotic resistance. In many cases, however, the growth rate effects of MGEs can be small. To

evaluate the influence of λ_{ij} magnitude, we also analyzed different ranges of growth rates effects

of mobile genetic elements, by sampling λ_{ij} values from uniform distributions with given widths

(Fig. S13). Greater width led to larger magnitude of growth rate effects. We used five-species

populations as an example and tested different ranges. Our results suggested that multistability

was more feasible when the growth rate effects of MGEs were small (Fig. S13b). The qualitative

relationship between HGT and community was not dependent on the range of growth rate effects

(Fig. S13a).

So far, our analysis has been focused on communities where the interaction network is fully

connected, and each species competes with every other member. In each of such populations, it is

highly infeasible to achieve the stable coexistence of different species. While this analysis provides

the basic proof of principle, natural environments often consist of many niches[30]. Species living in

the same niche competes while species from different niches can coexist (Fig. S14a). To

understand whether our conclusion is still applicable when niche effects are considered, we

extended our analysis to communities carrying a random number of niches. Each species was

randomly allocated into one of the niches, and the stability landscape was analyzed in the same

way by randomly initializing the systems (see Methods for more details). Our results suggested

that increasing HGT rate still promoted multistability, regardless of the niche structure of the

competing communities (Fig. S14b-d).

The role of HGT on community multistability when mobile genes modify interspecies interactions

Our previous analysis has focused on the role of HGT in communities where MGEs only affect

species growth rates and have no influences on interspecies interactions. To evaluate how this

modeling structure and the underlying assumptions affected the prediction, we extended our

framework by accounting for the modifications of interspecies interactions by mobile genes.

In nature, the horizontal transfer of some genes can change the interspecies interaction

strengths between different species. For instance, the sharing of many mobile genes can promote

niche overlapping, leading to an increase of competition strength[31,32]. To understand how the

transfer of these genes would influence community multistability, we adapted the previous model

by considering the dynamic change of competition strength during HGT (see Supplementary

Materials for more details). For two-species populations, we divided the interspecies interactions

into two components: the basal competition strengths (γ_1 and γ_2) and the added parts by HGT

(δ_1 and δ_2). The overall competition strength was calculated as $\gamma_1 + \delta_1 \frac{p_1}{s_1}$ and $\gamma_2 + \delta_2 \frac{p_2}{s_2}$,

respectively. Positive δ's mean HGT promotes the inter-species competition, while negative δ's

mean HGT reduces the strength of interspecies competition (Fig. 3a-b). Numerical simulations

with randomized parameters suggested that when δ was positive, HGT still promoted the

bistability, whereas if δ was negative, gene transfer would reduce the chance of bistability (Fig.

3c-d). We further extended our analysis to multispecies communities and calculated the number of

stable states. As shown in Fig. 3e and f, the number of stable states increases with HGT rate when

MGEs promotes competition, while decreases with HGT rate when MGEs reduces competition.

These results suggested that the effects of HGT on bistability depended on how mobile genes

shaped species growth rates and competitions. To place these two factors in a comprehensive

framework, we further generalized the model structure, by accounting for the scenario where

mobile genes modified growth rates and competitions at the same time. The effect of mobile genes

on growth rates was represented by the magnitude of λ's, and the influence on competition is

described by the parameter δ (Fig. 3g to i). By varying these two parameters, we can evaluate

how the model structure and the underlying assumptions affected the baseline expectation. We

performed additional simulations with broad ranges of λ and δ values. In particular, we

analyzed whether HGT would promote the likelihood of bistability in two-species communities

compared with the scenario without gene transfer. Our results suggested that: (1) With or without

HGT, reducing λ (increasing ecological neutrality) promoted bistability (Fig. 3g and h); (2) With

HGT, increasing δ promoted bistability (Fig. 3h); (2) Compared with the population without HGT,

gene transfer promoted bistability when δ was zero or positive, while reduces bistability when δ

was largely negative (Fig. 3i). These results suggested that the interplay among HGT, species

growth and interactions added new insights into the fundamental question of how microbes

competing for a limited number of resources stably coexisted.

The role of HGT on community multistability when MGE epitasis effects are present

In previous analysis, we also assumed that the effects of MGEs on host growth (λ_1 and λ_2)

were independent of the host species. In nature, however, the same MGE can have different growth

rate effects in different genomic backgrounds, a phenomenon called epistasis[33-36]. There are two

types of epistasis: magnitude epistasis, where the host genomic background only affects the

magnitude but not the sign of λ value, and sign epistasis, where the same MGE is burdensome in

one species while beneficial in the other host. The model can be readily extended to account for

these two types of epistasis (see Supplementary Materials for more details). In particular, we

dissected the growth rates μ_1 and μ_2 into two components: the basal growth rate (μ_1^0 and μ_2^0),

and the growth rate effects (λ_{11} and λ_{22}) of the mobile genetic elements. The growth rates of p_1

and p_2 were calculated as $\mu_1(1 + \lambda_{12})$ and $\mu_2(1 + \lambda_{21})$, respectively. Here λ_{ij} ($i, j = 1,2$)

stands for the growth rate effects of the j-th mobile genetic element in the i-th species. Without

epistasis, the growth rate effects are independent of the host species ($\lambda_{11} = \lambda_{21}$ and $\lambda_{22} = \lambda_{12}$).

The epistasis can be quantified by two ratios: $\xi_1 = \lambda_{21}/\lambda_{11}$ and $\xi_2 = \lambda_{12}/\lambda_{22}$. $\xi_1 > 0$ and

$\xi_2 > 0$ represent magnitude epistasis, whereas $\xi_1 < 0$ or $\xi_2 < 0$ represent the sign epistasis.

We carried out additional numerical simulations to evaluate how different epistasis types

influenced the role of HGT in community stability. Our results suggested that with magnitude

epistasis, HGT still enlarged the area of bistability region (Fig. 4a and b). In contrast, sign epistasis

overturned HGT's role on community stability: with sign epistasis, increasing gene transfer rate

reduced the area of bistability region (Fig. 4c and d). These results suggest that MGE epistasis

might add another layer of complexities into the interplay between HGT and community stability.

Interplay between HGT and the multistability of communities under strong selections

The growth rate effect of an MGE can be extreme in some cases. For instance, under strong

antibiotic selection, only cells carrying the antibiotic resistance genes (ARGs) can survive. To

examine whether our conclusion is still applicable in this scenario, we generalized our model by

considering the transfer of an MGE in a population under strong environmental selection (see

Supplementary Materials for more details). Specifically, we assumed that the MGE was initially

carried by one species in the populations. Without HGT, only the donor species carrying the MGE

can survive due to selection (Fig. 5a). In this case, the system will always be monostable regardless

of the initial species compositions. With HGT, the MGE can be transferred to other species,

creating chances for other species to survive (Fig. 5b). To understand how HGT would affect

community stability under this condition, we first analyzed the phase diagram of the two-species

population. Our simulation results suggested that bistability became possible when there existed

gene transfer and increasing HGT rate in general enlarged the bistability area (Fig. 5c and d). Here,

without loss of generality we assumed that the MGE was initially carried only by species 1.

Bistability was more feasible when $\mu_2 > \mu_1$ (Fig. 5c), suggesting that the effect of HGT on

promoting bistability would be stronger when the weak competitor acted as the MGE donor. We

further analyzed the number of stable states in multispecies communities. As shown in Fig. 5e and

f, for these populations under antibiotic selection, increasing transfer rate of the MGE also

promoted the emergence of multistability. This conclusion equally held under different interaction

strengths or growth rate ranges (Fig. S15), suggesting the general applicability of HGT's role in

promoting multistability for communities under strong selection.

HGT-enabled multistability promotes the regional coexistence of competing species

Our results highlight the role of HGT on shaping the stability landscape of closed local

communities. Natural microbiomes, however, exist at scales broader than a localized population.

A set of local communities in similar environments, also known as 'patches', form a

metacommunity[37,38]. When multiple stable states coexist, the species compositions of different

patches may differ from each other due to the variability of their initial configurations. Collectively,

the multistability in the local patches might give rise to the regional species diversity in the

metacommunity. This intuition leads to our hypothesis that HGT allows the stable coexistence of

multiple competing species at regional scale, even when species exclusively outcompete each other

in every local population.

To examine this hypothesis, we constructed metacommunities of 100 patches connected by

dispersal of microbes. Without loss of generality, we considered a 2-D array of local patches and

assumed that the dispersal only occurred between adjacent patches (Fig. S16a, see Methods for

more details). Each local patch carried 10 competing species that exchanged mobile gene elements

with each other. The dynamics of each patch was governed by the same set of parameters but

initialized with different species compositions. We then numerically simulated the dynamics of the

metacommunity till it reached the steady state.

Without HGT, the metacommunity exhibited strong spatial homogeneity at regional scale (Fig.

S16b). Different patches rested at similar species compositions. In contrast, HGT created a spatial

pattern of species distributions across different patches, due to the emergent multistability (Fig.

S16b). To analyze the influence of HGT on regional diversity, we pooled all local populations

together and calculated the collective species diversity in the pool using Shannon index. Our results

suggested that while local diversity remained low regardless of HGT rate, the regional diversity of

microbial species increased with HGT rate (Fig. S16b and c). The conclusion is applicable for

different dispersal rates (Figs. S17 and S18). These results highlight the fundamental role of HGT

in community stability and diversity at local and regional scale.

Discussion

Our work predicts that in many scenarios horizontal gene flow promotes multistability in

communities of competing microbes. At the scale of local populations, the emergence of

alternative stable states undermines the global stability of the community, making the system less

resilient to perturbations and creating greater opportunities for regime shifts. In metacommunities,

gene flow creates spatial heterogeneity of species compositions in different patches. Despite the

low diversity in every local patch, HGT-driven multistability creates substantial diversity at

regional scale. In microbial ecology, how microbes competing for a limited number of resources

stably coexist has been a long-standing puzzle[39,40]. Our results highlight the importance of

considering gene exchange when addressing this fundamental challenge.

While this work predicts the potential role of HGT in microbiota diversity and stability, several

caveats need to be considered when applying this prediction. For MGEs with sign epistasis, or

MGEs that relax interspecies competitions, HGT might reduce the number of stable states. The

relative importance of gene flow in a real community might also be context dependent. The effect

of HGT in promoting multistability might be stronger between species with intense competition

(Fig. S3) and similar growth rates (Fig. S13). The relative importance of HGT in real environment

is also dependent on the maximum carrying capacity of the population. Simulations by scanning

the η value from 10^{-7} to 1 suggested that increasing HGT rates started to promote multistability

when η value exceeded 10^{-2} per hour (Fig. S9). This corresponds to a conjugation efficiency of

10^{-11} $cell^{-1} \cdot hr^{-1} \cdot mL$ when the maximum carrying capacity equals 10^9 $cells \cdot mL^{-1}$, or a

conjugation efficiency of 10^{-14} $cell^{-1} \cdot hr^{-1} \cdot mL$ when the maximum carrying capacity equals

10^{12} $cells \cdot mL^{-1}$. Therefore, in environments with high cell density and abundant MGEs, such

as mammal gut, the role of HGT will be more prominent[41]. In contrast, the influence of HGT on

community multistability might become less important in populations with poor cell density and

slow gene transfer.

Our work provided a comprehensive framework to explore how different parameters,

including but not limited to HGT rates, collectively shaped community multistability. For instance,

our simulations suggested that nonzero cell death or dilution rate was essential for the bistability

of two competing species. Promoting competition strength in general enlarged multistability

region. Besides, multistability was also dependent on community neutrality, *i.e.* the growth rate

similarities among different competitors. Multistability is more feasible for communities with high

neutrality. Which mechanisms contribute most to the multistability of a given community might

be context-specific, depending on the values of different parameters in the population.

Our model allowed multiple plasmids to infect a single cell, but ignored the ecological

interactions between different plasmids. These complex interactions are governed by many

different processes: from systems that exclude the entry of other plasmids, incompatibility system

that eliminates within-cell coexistence, to interference in the regulation of plasmid copy

number[42,43]. These interactions can further change the distribution of plasmids among different

species, alter the structure of gene transfer network, and reshape the stability landscape of a

microbial community. However, the general conclusion regarding how these interactions shape

population multistability remains to be established in future studies.

Previous studies documented the presence of multiple alternative community types in soil[15],

human vaginal, gut and oral microbiomes[2,44,45]. In particular, multiple groups of bacteria in human

gut exhibit robust bistable abundance distributions, resulting in over 32 different stable state

combinations[46]. Our results provide a plausible explanation for the large number of community

types in these populations, even though our interpretation doesn't exclude other mechanisms including gradients of environmental factors[1]. Disentangling the contributions of HGT from the other factors in diverse natural microbiomes is a promising topic for future research.

Controlling the multistability of complex microbiota has important applications in human health. For instance, in gut microbiome, the development of unhealthy stable states is closely related to many diseases such as obesity, type 2 diabetes mellitus and gastrointestinal disorders[47-49]. Strategies have been developed to overcome the resilience of unhealthy states, including fecal transplantation or supplementation with probiotics[18]. Our work predicts that the horizontal transfer of MGEs such as plasmids might be an additional mechanism for the emergence of the unhealthy states. Indeed, human gut is a 'hotspot' of gene flow[41]. Previous studies suggested that some small molecules such as unsaturated fatty acids can inhibit the conjugative transfer of plasmids[50-52]. By binding the type IV secretion traffic ATPase TrwD, these compounds limit the pilus biogenesis and DNA translocation[53]. Therefore, modulating MGE spread using these molecules might offer new opportunities to reshape the stability landscape and narrow down the attraction domains of the disease states[51,54].

Figures

Fig. 1| Horizontal gene transfer (HGT) promotes bistability of communities consisting of

two competing microbes.

(a) Schematics of monostable and bistable systems. A monostable system will always rest at the

same steady state no matter where the system starts from. In contrast, a bistable system can reach

two distinct steady states, depending on the initial ratio between the two competitors. The right

panel is a diagram depicting how a microbial community responds to changes on initial species

abundances. The red marble stands for the system state, which will always roll "downhill" on the

landscape towards stable states at the low points. The x axis represents the community composition

and the valleys in the landscape are the stable states (or equilibrium points) when the community

composition will stop changing. Depending on parameter values, the landscape can have one

valley (called "monostable") or two valleys (called "bistable"). When the system is bistable, the

marble can reach either valley, depending on where it starts.

(b) The phase diagram of the population with (bottom panel, $\eta = 0.2\ hr^{-1}$) or without HGT (top

panel). Different combinations of μ_1 and μ_2 between 0 and 1 hr^{-1} were tested. For each

combination, the initial abundance of each species was randomized 200 times between 0 and 1

following uniform distributions. The system was monostable if all initializations led to the same

steady state. Otherwise, the system was bistable. All combinations of μ_1 and μ_2 resulting in

bistability were marked in blue in the diagram. Other parameters are $\gamma = 1.1$, $\kappa = 0.005\ hr^{-1}$,

$D = 0.2\ hr^{-1}$, $\mu_1^0 = \mu_2^0 = 0.5\ hr^{-1}$. Two states, P ($\mu_1 = 0.3\ hr^{-1}$, $\mu_2 = 0.8\ hr^{-1}$) and Q

($\mu_1 = 0.4\ hr^{-1}$, $\mu_2 = 0.6\ hr^{-1}$), were labeled as examples.

(c-d) Population dynamics of P and Q under 100 different initializations. Here, the dynamic

changes of species 1's abundance were shown. Three different HGT rates (0, 0.1, 0.2 hr^{-1} from

top to bottom) were tested. When gene transfer rate increases, the system at Q changed from

monostable to bistable, while the system at P remained monostable.

Fig. 2| The number of alternative stable states in multi-species communities increases with HGT rate.

(a), A schematic of the stability landscape of a population composed of multiple competing species with or without HGT. Each alternative stable state is characterized by a different dominant species (shown in filled circles).

(b), The emergence of alternative stable states in a five-species community with gene transfer. We

simulated the dynamics of 500 parallel communities, each with the same kinetic parameters but

different initial compositions. The species compositions of these communities at three different

timepoints were shown by principal component analysis from left to right. Without HGT ($\eta = 0$),

all populations converged into a single stable state. In contrast, with HGT ($\eta = 0.2\ hr^{-1}$), four

different attractors appear. Here, populations reaching different stable states were marked with

different colors. Other parameters are $\mu_1^0 = \mu_2^0 = \mu_3^0 = \mu_4^0 = \mu_5^0 = 0.5\ hr^{-1}$, $\mu_1 = 0.52\ hr^{-1}$,

$\mu_2 = 0.48\ hr^{-1}$, $\mu_3 = 0.44\ hr^{-1}$, $\mu_4 = 0.60\ hr^{-1}$, $\mu_5 = 0.31\ hr^{-1}$, $\gamma = 1.05$, $\kappa = $

$0.005\ hr^{-1}$, $D = 0.2\ hr^{-1}$.

(c), The number of stable states (top panel) and the multistability coefficient χ increase with HGT

rate in communities consisting of 8 competing species. We calculated the number of stable states

by randomly initializing the species abundances 500 times between 0 and 1 following uniform

distributions. Then we simulated the steady states and clustered them into different attractors by

applying a threshold of 0.05 on their Euclidean distances. The data were presented as mean $\pm$

standard deviation of 10 replicates. Each replicate corresponded to a different combination of

randomized species growth rates.

Fig.3| The effects of HGT on population multistability when MGEs promote or reduce the strength of interspecies competition.

(a-b), The schematics of HGT promoting or reducing competition.

(c), For populations of two species, when MGEs promote competition, increasing HGT rate enlarges the area of bistability region in the phase diagram. Here, δ describes the effect of mobile genes on the competition strength. Positive δ represents HGT promoting competition. In numerical simulations, we tested three different δ values (marked in different colors). When calculating the area of bistability region, we randomized μ_1 and μ_2 500 times between 0 and 1 hr^{-1} following uniform distributions while keeping μ_1^0 and μ_2^0 constants. For each pair of growth rates, we randomized the initial abundance of each species 200 times between 0 and 1 following uniform distributions. The system was monostable if all initializations led to the same steady state. Otherwise, the system was bistable. Then we calculated the fraction of growth rate combinations

that generated bistability out of the 500 random combinations. Other parameters are $\gamma_1 = \gamma_2 =$

1.1, $\kappa = 0.005\ h^{-1}$, $D = 0.2\ h^{-1}$.

(d), When MGEs reduce competition, the area of bistability region decreases with HGT rate. Three

negative values δ were tested and shown as examples here.

(e), For populations of 5 species, when MGEs promote competition, the number of stable states

increases with HGT rate. We calculated the number of stable states by randomly initializing the

species abundances 500 times. Then we simulated the steady states and clustered them into

different attractors by applying a threshold of 0.05 on their Euclidean distances. The data were

presented as mean $\pm$ standard deviation of 10 replicates. Each replicate corresponds to a different

combination of randomized species growth rates. Other parameters are $\gamma_{i,j} = 1.1$, $\kappa =$

$0.005\ h^{-1}$, $D = 0.2\ h^{-1}, \delta = 0.5$.

(f), When MGEs reduce competition, the number of stable states decrease with HGT rate. The data

were presented as mean $\pm$ standard deviation of 10 replicates. $\delta = -0.5$ was used in the

simulation.

(g), For populations of two species without HGT, reducing the magnitude of λ (promoting

ecological neutrality) enlarges the area of bistability. When gene transfer rate equals zero, changing

δ does not influence bistability. When calculating bistability probability, we randomized λ_1 and

λ_2 500 times between $-\alpha$ and α following uniform distributions. α represents the magnitude

of λ variations. μ_1 and μ_2 were calculated as $\mu_1^0(1 + \lambda_1)$ and $\mu_2^0(1 + \lambda_2)$, respectively. The

overall competition strength was calculated as $\gamma_1 + \delta_1 \frac{p_1}{s_1}$ and $\gamma_2 + \delta_2 \frac{p_2}{s_2}$. We tested different

combinations of α (the magnitude of λ variations) and δ values. We then calculated the

fraction (f_0) of growth rate combinations that generated bistability out of the 500 random

combinations. Other parameters are $\gamma_1 = \gamma_2 = 1.1$, $\kappa = 0.005\ h^{-1}$, $D = 0.2\ h^{-1}$.

(h), In populations of two species with HGT, reducing the magnitude of λ (promoting ecological

neutrality) or promoting δ value enlarges the area of bistability (f_1). η being 0.4 was tested as

an example here.

(i), The effect of HGT on community bistability depends on how mobile genes modify growth rates and competition. Compared with the population without HGT, gene transfer promotes bistability when δ is zero or positive, while reduces bistability when δ is largely negative.

Fig. 4| The influence of epistasis on the role of HGT in mediating population bistability.

(a-b) With magnitude epistasis, increasing HGT rate still promotes bistability in populations of two competing species.

(c-d) With sign epistasis, increasing HGT rate reduces the area of bistability region. For each type of epistasis, we considered two scenarios. In a and c, host genetic background influences the growth rate effect of only one MGE, while in b and d, host genetic background influences both MGEs. ξ_1 and ξ_2 are defined as $\lambda_{21}/\lambda_{11}$ and $\lambda_{12}/\lambda_{22}$, respectively. $\xi_1 > 0$ and $\xi_2 > 0$ represent magnitude epistasis, while $\xi_1 < 0$ or $\xi_2 < 0$ represents the sign epistasis. Other parameters are $\gamma_1 = \gamma_2 = 1.1$, $\kappa = 0.005\ h^{-1}$, $D = 0.2\ h^{-1}$.

Fig. 5| HGT creates chances of multistability for communities under strong environmental selection.

(a-b), The schematics of communities under strong selection. Without HGT, only the donor species carrying the MGE can survive. HGT allows the other species to acquire MGE from the donor, creating opportunities for the other species to survive under strong selections.

(c), The phase diagram of two-species populations transferring an MGE. Different combinations of μ_1 and μ_2 between 0 and 1 hr^{-1} were tested. All combinations of μ_1 and μ_2 that led to bistability were marked in blue in the diagram. Other parameters are $\gamma = 1.1$, $\kappa = 0.005 \ hr^{-1}$, $D = 0.2 \ hr^{-1}$, $\mu_1^0 = \mu_2^0 = 0.5 \ hr^{-1}$. Here, we assumed that the ARG was initially carried only by species 1. Bistability was more feasible when $\mu_2 > \mu_1$.

(d), Increasing HGT rate enlarges the area of bistability region in the phase diagram of two competing species transferring an MGE. Three different competition strengths were tested (marked in different colors).

(e-f), For communities of 5 or 8 species under strong selection, increasing HGT rate promotes the emergence of alternative stable states. We calculated the number of stable states by randomly

initializing the species abundances 500 times. Then we simulated the steady states and clustered

them into different attractors. The carriage of the mobile gene changed the species growth rate

from 0 to a positive value μ_i. When calculating the number of stable states in multi-species

communities, we randomly drew the μ_i values from a uniform distribution between 0.3 and 0.7

hr^{-1}. The data were presented as mean $\pm$ standard deviation of 10 replicates. Each replicate

corresponds to a different combination of randomized species growth rates. Other parameters are

$\gamma_{i,j} = 1.5, \ \kappa = 0.005 \ h^{-1}, \ D = 0.2 \ h^{-1}$.

Methods

Modeling framework of two competing species sharing mobile genes

We used a mathematical model developed in a previous work to analyze the population

dynamics of two competing species transferring mobilizable plasmids with growth rate effects[28].

Briefly, the model includes four ordinary differential equations (ODEs):

519
$$\frac{dS_1}{dt} = \mu_1^e S_1 \left(1 - \frac{S_1 + R_2 S_2}{N_{m,1}} \right) - DS_1, \quad (1)$$

520
$$\frac{dS_2}{dt} = \mu_2^e S_2 \left(1 - \frac{S_2 + R_1 S_1}{N_{m,2}} \right) - DS_2, \quad (2)$$

521
$$\frac{dP_1}{dt} = \mu_1 (1 + \lambda_2) P_1 \left(1 - \frac{S_1 + R_2 S_2}{N_{m,1}} \right) + \eta_1^c (S_2 + P_1)(S_1 - P_1) - (D + \kappa_1) P_1, \quad (3)$$

522
$$\frac{dP_2}{dt} = \mu_2 (1 + \lambda_1) P_2 \left(1 - \frac{S_2 + R_1 S_1}{N_{m,2}} \right) + \eta_2^c (S_1 + P_2)(S_2 - P_2) - (D + \kappa_2) P_2. \quad (4)$$

S_1 and S_2 are the abundances of the two species. R_1 and R_2 describe the strengths of

interspecies competition. $N_{m,1}$ and $N_{m,2}$ are the maximum carrying capacities for species 1 and

2, respectively. D is the dilution rate. P_i $(i = 1, 2)$ represents the abundance of the subpopulation

in the i-th species that acquires the mobilizable genes from its competitor. Here, we assumed that

the populations would not lose their own native plasmids completely. The internal inhibition within

each species was also assumed to be independent of the acquired MGEs. μ_i represents the

maximum 'per capita' growth rate of the i-th species when interspecies HGT is absent. μ_i is

calculated as $\mu_i = \mu_i^0 (1 + \lambda_i)$ where μ_i^0 is the basal growth rate determined by the non-

mobilizable genes and λ_i is the growth rate effect of the plasmids. Interspecies gene flow allows

each species to acquire plasmids from its competitor, resulting in the dynamic change of the

effective species growth rates μ_i^e, which is calculated as: $\mu_1^e = \mu_1 (1 + \lambda_2 \frac{P_1}{S_1})$ and $\mu_2^e = \mu_2 (1 +$

$\lambda_1 \frac{P_2}{S_2})$. For instance, all cells of species 1 carry the plasmid 1. Therefore, their growth rate can be

obtained as $\mu_1^0(1 + \lambda_1)$. After HGT, the overall growth rate of species 1 changes because some of

the individuals get the plasmid from the second species. Such a change is reflected by the λ_2 term

in $\mu_1^e = \mu_1(1 + \lambda_2 \frac{P_1}{S_1})$. η_1^c and η_2^c stand for HGT rates. Here, we described the HGT process by

mass action kinetics, an assumption commonly made for plasmid transfer dynamics. We further

assumed that intraspecies and interspecies transfer occurred at the same rates. κ_1 and κ_2

represent the loss rates of the mobile genetic elements. For plasmids, κ_1 and κ_2 describe the loss

rate by segregation errors.

Let $s_1 = S_1/N_{m,1}$, $s_2 = S_2/N_{m,2}$, $p_1 = P_1/N_{m,1}$, $p_2 = P_2/N_{m,2}$, $\gamma_2 = R_2 \frac{N_{m,2}}{N_{m,1}}$, $\gamma_1 =$

$R_1 \frac{N_{m,1}}{N_{m,2}}$. The model can be further simplified as

544
$$\frac{ds_1}{dt} = \mu_1^e s_1(1 - s_1 - \gamma_2 s_2) - Ds_1, \quad (5)$$

545
$$\frac{ds_2}{dt} = \mu_2^e s_2(1 - s_2 - \gamma_1 s_1) - Ds_2, \quad (6)$$

546
$$\frac{dp_1}{dt} = \mu_1(1 + \lambda_2)p_1(1 - s_1 - \gamma_2 s_2) + \eta_1(\varrho s_2 + p_1)(s_1 - p_1) - (D + \kappa_1)p_1, \quad (7)$$

547
$$\frac{dp_2}{dt} = \mu_2(1 + \lambda_1)p_2(1 - s_2 - \gamma_1 s_1) + \eta_2(\varrho^{-1} s_1 + p_2)(s_2 - p_2) - (D + \kappa_2)p_2. \quad (8)$$

548   Here, s_1, s_2, p_1 and p_2 are dimensionless. η_i is related with η_i^c by $\eta_i = N_{m,i}\eta_i^c$. ϱ

549   describes the ratio between $N_{m,2}$ and $N_{m,1}$: $\varrho = N_{m,2}/N_{m,1}$. For simplicity, we assumed that the

550   two species had the same carrying capacity, *i.e.* $\varrho = 1$. The unit of η_1 and η_2 is $hour^{-1}$.

551   Without HGT, p_1 and p_2 become consistently 0, and the model is equivalent to the classic

552   Lotka-Volterra (LV) framework:

553
$$\frac{ds_1}{dt} = \mu_1 s_1(1 - s_1 - \gamma_2 s_2) - Ds_1, \quad (9)$$

554
$$\frac{ds_2}{dt} = \mu_2 s_2(1 - s_2 - \gamma_1 s_1) - Ds_2. \quad (10)$$

Stability analysis of the classic LV model

The classic LV model has four fixed points: $g_1 = [0,0]$, $g_2 = [0, 1 - \frac{D}{\mu_2}]$, $g_3 = [1 - \frac{D}{\mu_1}, 0]$,

$g_4 = [\frac{\gamma_2\mu_1\mu_2 - D\gamma_2\mu_1 - \mu_1\mu_2 + D\mu_2}{\mu_1\mu_2(\gamma_1\gamma_2 - 1)}, \frac{\gamma_1\mu_1\mu_2 - D\gamma_1\mu_2 - \mu_1\mu_2 + D\mu_1}{\mu_1\mu_2(\gamma_1\gamma_2 - 1)}]$. The Jacobian matrix of each fixed point

can be calculated as $J = \begin{bmatrix} \mu_1 - 2\mu_1 s_1^* - \mu_1\gamma_2 s_2^* - D & -\mu_1\gamma_2 s_1^* \\ -\mu_2\gamma_1 s_2^* & \mu_2 - 2\mu_2 s_2^* - \mu_2\gamma_1 s_1^* - D \end{bmatrix}$, where s_1^* and

s_2^* are the species abundances at each fixed point.

The system is bistable only when g_2 and g_3 are both stable. The Jacobian matrix of g_2 is

$J_2 = \begin{bmatrix} \mu_1 - D - \mu_1\gamma_2 + \frac{\mu_1\gamma_2 D}{\mu_2} & 0 \\ -\mu_2\gamma_1 + D\gamma_1 & D - \mu_2 \end{bmatrix}$. The eigenvalues of J_2 can be obtained as $\mu_1 - D -$

$\mu_1\gamma_2 + \frac{\mu_1\gamma_2 D}{\mu_2}$ and $D - \mu_2$. g_2 is stable when both eigenvalues are negative, which leads to $\gamma_2 >$

ϕ_1/ϕ_2 where $\phi_1 = \frac{\mu_1 - D}{\mu_1}$ and $\phi_2 = \frac{\mu_2 - D}{\mu_2}$. Similarly, g_3 is stable when $\gamma_1 > \phi_2/\phi_1$.

Therefore, the condition for the system being bistable is $\gamma_2 > \phi_1/\phi_2$ and $\gamma_1 > \phi_2/\phi_1$.

Bistability analysis of two-species populations by numerical simulations

We performed numerical simulations to determine the bistability of a population consisting of

two competing species transferring mobile genes. Specifically, for a system with given parameters,

we randomly initialized the species abundances between 0 and 1 following uniform distributions

for 200 times. Then we simulated the dynamics of each population till the abundance of every

species reached equilibrium, and we treated it as a steady state. We grouped the 200 steady states

into different attractors by applying the threshold of 0.01. Two steady states with their Euclidean

distance smaller than the threshold belonged to the same group. The Euclidean distance was

calculated as $d_{ij} = \sqrt{\sum_{k=1}^{m}(s_{i,k} - s_{j,k})^2}$ where k is the species index and $s_{i,k}$ represents the

abundance of the k-th species in the i-th steady state.

Theoretical framework of multiple competing species transferring mobile genes

For a community of m competing species that share mobile genes with each other, the

modeling framework consists of two groups of ODEs:

$$\frac{ds_i}{dt} = \mu_i^e s_i \left(1 - s_i - \sum_{j \neq i} \gamma_{ji} s_j\right) - Ds_i, \quad (11)$$

$$\frac{dp_{ij}}{dt} = \mu_i(1 + \lambda_{ij}) \left[\prod_{k \neq i,j} \left(1 + \lambda_{ik} \frac{p_{ik}}{s_i}\right)\right] p_{ij} \left(1 - s_i - \sum_{j \neq i} \gamma_{ji} s_j\right) + (s_i - p_{ij}) \sum_{k=1}^{n} \eta_{jki} p_{kj}$$

$$- (D + \kappa_{ij}) p_{ij}. \quad (i \neq j). \quad (12)$$

s_i is the abundance of the i-th species. p_{ij} is the size of the subpopulation in the i-th species that acquires s_j-originated MGEs. γ_{ji} is the negative interaction that s_j imposes on the i-th species. D is the dilution rate. μ_i^e, the effective growth rate of s_i, is calculated by $\mu_i^e = \mu_i \prod_{j \neq i} \left(1 + \lambda_{ij} \frac{p_{ij}}{s_i}\right)$. μ_i is defined as $\mu_i = \mu_i^0(1 + \lambda_{ii})$ where μ_i^0 is the basal growth rate, λ_{ij} is the growth rate effect of the s_j-originated MGEs in the i-th species. η_{jki} is the transfer rate of the s_j-originated plasmids from species k to species i. κ_{ij} is the loss rate of the j-th mobile genetic element in the i-th species.

Analysis of multistability of complex communities

For a population of m species with given niches and kinetic parameters, we initialized the system 500 times in the ranges of $0 < s_i < 1$ following uniform distributions. The initial conditions for p_{ij}'s are $p_{ii} = s_i$ and $p_{ij} = 0$ $(i \neq j)$. Then the dynamics of each population was simulated for 2000 hours till the system reached the steady state. For each steady state we obtained the corresponding species composition and grouped the 500 solutions into different stable states based on their Euclidean distances.

Horizontal gene transfer within communities under strong environmental selection

Under strong antibiotic selections, only cells carrying antibiotic resistant MGEs can survive. To examine whether our conclusion is still applicable in this scenario, we generalized our model by considering the transfer of an antibiotic-resistant gene (ARG) in a population of m species under strong antibiotic selection. Within each species, only cells carrying the MGE can grow, with

600   μ_i being their maximum growth rate. The cells without the MGE will be eventually depleted by

601   dilution D. The community dynamics can be described by the following ODEs ($i = 1,2, \dots, m$):

602
$$\frac{ds_i}{dt} = \mu_i p_i \left(1 - s_i - \sum_{j \neq i} \gamma_{ji} s_j \right) - D s_i, \quad (13)$$

603
$$\frac{dp_i}{dt} = \mu_i p_i \left(1 - s_i - \sum_{j \neq i} \gamma_{ji} s_j \right) + (s_i - p_i) \sum_{j=1}^{m} \eta_{ji} p_j - \kappa_i p_i - D p_i. \quad (14)$$

s_i is the total abundance of the i-th species, and p_i is the abundance of cells carrying the MGE

in s_i. In the first equation, the first term describes the overall growth of the i-th species, which is

only contributed by p_i since the MGE-free cells are not able to grow. η_{ji} is the MGE transfer

rate from the j-th to the i-th species. κ_i is MGE loss rate in the i-th species. γ_{ji} describes the

inter-species competition strength.

Without loss of generality, we assumed the species 1 was the MGE donor and p_1 was equal

with s_1. Without HGT, the MGE only resides in the donor species, while HGT allows the MGE

to be shared with other species. To understand how HGT rate influences community multistability,

we calculated the phase diagram of populations of two competing species. Increasing HGT rate

enlarged the bistability region. We further analyzed the number of alternative stable states in

communities of 5 or 8 species. In particular, we randomly initialized the species abundances 500

times, simulated the steady states and clustered them into different attractors. When calculating

the number of stable states in multi-species communities, we randomly drew the μ_i values from

a uniform distribution between 0.3 and 0.7 hr^{-1}. Our results suggested that the number of stable

states in general increased with HGT rate.

Niche-based model of multi-species communities

For a community consisting of m competing species, let the number of niches be l. Each

niche has a maximum carrying capacity $N_m^{(i)}$ ($i = 1,2, \dots, l$). Each species lives in one of the

niches. Let M be the map from species index to the index of its niche. $M_i = k$ means that the i-

th species lives in the k-th niche. For species i and j, if $M_i = M_j$, these two species belong to

the same niche and compete with each other. Species from different niche do not directly interact.

Therefore, the model becomes:

$$\frac{ds_i}{dt} = \mu_i^e s_i \left(1 - \frac{s_i + \sum_{\{j:\, M_j = M_i\}} \gamma_{ji}\, s_j}{N_m^{(M_i)}}\right) - D s_i, \quad (15)$$

$$\frac{dp_{ij}}{dt} = \mu_i(1 + \lambda_{ij}) \left[\prod_{k \neq i,j} \left(1 + \lambda_{ik} \frac{p_{ik}}{s_i}\right)\right] P_{ij} \left(1 - \frac{s_i + \sum_{\{j:\, M_j = M_i\}} \gamma_{ji}\, s_j}{N_m^{(M_i)}}\right) + (s_i - p_{ij}) \sum_{k=1}^{n} \eta_{jki} p_{kj}$$

$$\qquad - (D + \kappa_{ij}) p_{ij}. \quad (i \neq j). \quad (16)$$

Here, s_i is the abundance of the i-th species. p_{ij} is the size of the subpopulation in the i-th

species that acquires s_j-originated mobile genes. γ_{ji} is the negative interaction that s_j imposes

on the i-th species. η_{jki} is the transfer rate of the s_j-originated genes from species k to species

i. In simulations, the number of niches was randomized between 2 and $m/2$. The maximum

carrying capacity of each niche was randomized between 0 and 1 following uniform distributions.

Then each species was randomly allocated into one of the niches.

Theoretical framework of metacommunity dynamics

Consider a metacommunity composed of $u \times v$ local patches. We assumed a grid-like

arrangement of patched connected by dispersal terms $\mathcal{H}$. The location of each patch can be

described by two coordinates $[a, b]$ $(a = 1,2, \dots, u$ and $b = 1,2, \dots, v)$. Let ω be the dispersal

rate between the adjacent patches. The dynamics of the metacommunity can then be described by

generalizing equations (11) and (12) to each local patch:

$$\frac{ds_{a,b,i}}{dt} = \mu_{a,b,i}^e s_{a,b,i} \left(1 - s_{a,b,i} - \sum_{j \neq i} \gamma_{ji}\, s_{a,b,j}\right) - D s_{a,b,i} + \mathcal{H}_s^{(a,b)}, \quad (17)$$

$$\frac{dp_{a,b,i,j}}{dt} = \mu_i(1 + \lambda_{ij}) \left[\prod_{k \neq i,j} \left(1 + \lambda_{ik} \frac{p_{a,b,i,k}}{s_{a,b,i}}\right)\right] p_{a,b,i,j} \left(1 - s_{a,b,i} - \sum_{j \neq i} \gamma_{ji}\, s_{a,b,j}\right)$$

$$\qquad + (s_{a,b,i} - p_{a,b,i,j}) \sum_{k=1}^{n} \eta_{jki} p_{a,b,k,j} - (D + \kappa_{ij}) p_{a,b,i,j} + \mathcal{H}_p^{(a,b)}. \quad (i \neq j). \quad (18)$$

For the patch located at $[a, b]$, $s_{a,b,i}$ is the abundance of the i-th species, and $p_{a,b,i,j}$ is the size

of the subpopulation in the i-th species that acquires s_j-originated mobile genes. The effective

growth rate $\mu^e_{a,b,i}$ is calculated as $\mu^e_{a,b,i} = \mu_i \prod_{j \neq i} \left(1 + \lambda_{ij} \frac{p_{a,b,i,j}}{s_{a,b,i}}\right)$.

$\mathcal{H}_s^{(a,b)}$ and $\mathcal{H}_p^{(a,b)}$ are the gains of $s_{a,b,i}$ and $p_{a,b,i,j}$ by dispersal, respectively. We

assumed that microbial dispersal only occurred between adjacent patches. Therefore, $\mathcal{H}_s^{(a,b)}$ was

calculated by summing the dispersal terms at four directions: left, right, up, and down. The left

component can be calculated as

651
$$\mathcal{H}_{s,left}^{(a,b)} = \begin{cases} \omega(s_{a-1,b,i} - s_{a,b,i}), & if \ a-1 > 0 \\ 0, & if \ a-1 \leq 0 \end{cases}.$$

652   Similarly, $\mathcal{H}_{s,right}^{(a,b)}$, $\mathcal{H}_{s,up}^{(a,b)}$ and $\mathcal{H}_{s,down}^{(a,b)}$ is calculated as:

653
$$\mathcal{H}_{s,right}^{(a,b)} = \begin{cases} \omega(s_{a+1,b,i} - s_{a,b,i}), & if \ a+1 < u+1 \\ 0, & if \ a+1 \geq u+1 \end{cases},$$

654
$$\mathcal{H}_{s,up}^{(a,b)} = \begin{cases} \omega(s_{a,b+1,i} - s_{a,b,i}), & if \ b+1 < v+1 \\ 0, & if \ b+1 \geq v+1 \end{cases}$$

655
$$\mathcal{H}_{s,down}^{(a,b)} = \begin{cases} \omega(s_{a,b-1,i} - s_{a,b,i}), & if \ b-1 > 0 \\ 0, & if \ b-1 \leq 0 \end{cases}.$$

$\mathcal{H}_s^{(a,b)}$ can then be obtained as $\mathcal{H}_{s,left}^{(a,b)} + \mathcal{H}_{s,right}^{(a,b)} + \mathcal{H}_{s,up}^{(a,b)} + \mathcal{H}_{s,down}^{(a,b)}$. $\mathcal{H}_p^{(a,b)}$ can be

formulated in the similar way.

The collective species diversity in a metacommunity

To analyze the total species diversity in a metacommunity, we first calculated the total

abundance of each species across all patches: $s_i = \sum_{a=1,2,\ldots,u} \sum_{b=1,2,\ldots,v} s_{a,b,i}$. Then we measured

the diversity using Shannon index, a common metric used in ecology. Shannon index can be

calculated as $exp\left[-\sum_{i=1}^{m}\frac{s_i}{s_T}log\frac{s_i}{s_T}\right]$. s_T is the total species abundance: $s_T = \sum_{i=1}^{m}s_i$.

Biological relevance of our analysis

The empirically measured gene transfer rates (denoted as η^c, with the unit of $cells^{-1} \cdot mL \cdot$

$hour^{-1}$) should be multiplied by N_m, the maximum carrying capacity of the population, before

being used in our model. Therefore, the transfer rate η in our analysis are several orders of

magnitude higher than the experimentally measured rates[55,56]. In this work, we focused on HGT

rate in the range of $0 < \eta_i < 0.4\ hour^{-1}$. Whether this range is biologically relevant depends on

N_m of the population. Considering the typical estimates of plasmid transfer rates cross species

($10^{-13}\ cells^{-1} \cdot mL \cdot hour^{-1}$, from *Klebsiella pneumoniae* to *Escherichia coli*) and within

species ($10^{-7}\ cells^{-1} \cdot mL \cdot hour^{-1}$, between *E. coli* strains) provided by a previous study[55],

HGT rates in our analysis are relevant in various environments from human colon ($\sim 10^{12}$ cells per

gram) [57] to deep oceanic subsurface ($10^6 \sim 10^8$ cells per gram)[58].

References

Gonze, D., Lahti, L., Raes, J. & Faust, K. Multi-stability and the origin of microbial community types. *The ISME Journal* **11**, 2159-2166 (2017).

Arumugam, M. *et al.* Enterotypes of the human gut microbiome. *Nature* **473**, 174-180 (2011).

Gibson, T. E., Bashan, A., Cao, H.-T., Weiss, S. T. & Liu, Y.-Y. On the origins and control of community types in the human microbiome. *PLoS Computational Biology* **12**, e1004688 (2016).

Estrela, S. *et al.* Functional attractors in microbial community assembly. *Cell Systems* **13**, 29-42. e27 (2022).

Chaparro-Pedraza, P. C. & de Roos, A. M. Ecological changes with minor effect initiate evolution to delayed regime shifts. *Nature Ecology and Evolution* **4**, 412-418 (2020).

Beardmore, R. E. *et al.* Drug-mediated metabolic tipping between antibiotic resistant states in a mixed-species community. *Nature Ecology and Evolution* **2**, 1312-1320 (2018).

Amor, D. R., Ratzke, C. & Gore, J. Transient invaders can induce shifts between alternative stable states of microbial communities. *Science Advances* **6**, eaay8676 (2020).

Relman, D. A. The human microbiome: ecosystem resilience and health. *Nutrition Reviews* **70**, S2-S9 (2012).

Hartmann, M. & Six, J. Soil structure and microbiome functions in agroecosystems. *Nature Reviews Earth and Environment* **4**, 4-18 (2023).

Sun, L. *et al.* The shift of soil microbial community induced by cropping sequence affect soil properties and crop yield. *Frontiers in Microbiology* **14**, 1095688 (2023).

Kumar, S. & Hasty, J. Stability, robustness, and containment: preparing synthetic biology for real-world deployment. *Current Opinion in Biotechnology* **79**, 102880 (2023).

Fuhrman, J. A., Cram, J. A. & Needham, D. M. Marine microbial community dynamics and their ecological interpretation. *Nature Reviews Microbiology* **13**, 133-146 (2015).

Khazaei, T. *et al.* Metabolic multistability and hysteresis in a model aerobe-anaerobe microbiome community. *Science Advances* **6**, eaba0353 (2020).

Dubinkina, V., Fridman, Y., Pandey, P. P. & Maslov, S. Multistability and regime shifts in microbial communities explained by competition for essential nutrients. *Elife* **8**, e49720 (2019).

Fujita, H. *et al.* Alternative stable states, nonlinear behavior, and predictability of microbiome dynamics. *Microbiome* **11**, 63 (2023).

Mangal, U. *et al.* Multistability and hysteresis in states of oral microbiota: Is it impacting the dental clinical cohort studies? *Journal of Periodontal Research* **58**, 381-391 (2023).

Zaoli, S. & Grilli, J. A macroecological description of alternative stable states reproduces intra-and inter-host variability of gut microbiome. *Science Advances* **7**, eabj2882 (2021).

Fassarella, M. *et al.* Gut microbiome stability and resilience: elucidating the response to perturbations in order to modulate gut health. *Gut* **70**, 595-605 (2021).

van den Berg, N. I. *et al.* Ecological modelling approaches for predicting emergent properties in microbial communities. *Nature Ecology and Evolution* **6**, 855-865 (2022).

Soucy, S. M., Huang, J. & Gogarten, J. P. Horizontal gene transfer: building the web of life. *Nature Reviews Genetics* **16**, 472-482 (2015).

21 Bordenstein, S. R. & Reznikoff, W. S. Mobile DNA in obligate intracellular bacteria. *Nature Reviews*
Microbiology **3**, 688-699 (2005).
22 Brito, I. L. *et al.* Mobile genes in the human microbiome are structured from global to individual scales.
Nature **535**, 435-439 (2016).
23 Brockhurst, M. A. *et al.* The ecology and evolution of pangenomes. *Current Biology* **29**, R1094-R1103
(2019).
24 Smillie, C. S. *et al.* Ecology drives a global network of gene exchange connecting the human
microbiome. *Nature* **480**, 241-244 (2011).
25 Rankin, D. J., Rocha, E. P. & Brown, S. P. What traits are carried on mobile genetic elements, and why?
Heredity **106**, 1-10 (2011).
26 Werren, J. H. Selfish genetic elements, genetic conflict, and evolutionary innovation. *Proceedings of*
the National Academy of Sciences **108**, 10863-10870 (2011).
27 Gilpin, M. E. & CASE, T. J. Multiple domains of attraction in competition communities. *Nature* **261**, 40-
42 (1976).
28 Zhu, S., Hong, J. & Wang, T. Horizontal gene transfer is predicted to overcome the diversity limit of
competing microbial species. *Nature Communications* **15**, 800, doi:10.1038/s41467-024-45154-w
(2024).
29 Abreu, C. I., Friedman, J., Andersen Woltz, V. L. & Gore, J. Mortality causes universal changes in
microbial community composition. *Nature Communications* **10**, 2120 (2019).
30 Finlay, B. B. & Medzhitov, R. Host-microbe interactions: fulfilling a niche. *Cell Host and Microbe* **1**, 3-
4 (2007).
31 Bonham, K. S., Wolfe, B. E. & Dutton, R. J. Extensive horizontal gene transfer in cheese-associated
bacteria. *Elife* **6**, e22144 (2017).
32 Caro-Quintero, A. & Konstantinidis, K. T. Inter-phylum HGT has shaped the metabolism of many
mesophilic and anaerobic bacteria. *The ISME journal* **9**, 958-967 (2015).
33 Acar Kirit, H., Lagator, M. & Bollback, J. P. Experimental determination of evolutionary barriers to
horizontal gene transfer. *BMC Microbiology* **20**, 1-13 (2020).
34 Gama, J. A., Zilhão, R. & Dionisio, F. Plasmid interactions can improve plasmid persistence in bacterial
populations. *Frontiers in Microbiology* **11**, 2033 (2020).
35 Silva, R. F. *et al.* Pervasive sign epistasis between conjugative plasmids and drug-resistance
chromosomal mutations. *PLoS Genetics* **7**, e1002181 (2011).
36 San Millan, A., Heilbron, K. & MacLean, R. C. Positive epistasis between co-infecting plasmids promotes
plasmid survival in bacterial populations. *The ISME Journal* **8**, 601-612 (2014).
37 Leibold, M. A. *et al.* The metacommunity concept: a framework for multi-scale community ecology.
Ecology Letters **7**, 601-613 (2004).
38 Miller, E. T., Svanbäck, R. & Bohannan, B. J. Microbiomes as metacommunities: understanding host-
associated microbes through metacommunity ecology. *Trends in Ecology and Evolution* **33**, 926-935
(2018).
39 Serván, C. A., Capitán, J. A., Grilli, J., Morrison, K. E. & Allesina, S. Coexistence of many species in random
ecosystems. *Nature Ecology and Evolution* **2**, 1237-1242 (2018).

40   Grilli, J. *et al.* Feasibility and coexistence of large ecological communities. *Nature Communications* **8**,
14389 (2017).
41   Liu, L. *et al.* The human microbiome: a hot spot of microbial horizontal gene transfer. *Genomics* **100**,
265-270 (2012).
42   Lipsitch, M., Colijn, C., Cohen, T., Hanage, W. P. & Fraser, C. No coexistence for free: neutral null models
for multistrain pathogens. *Epidemics* **1**, 2-13 (2009).
43   Igler, C., Huisman, J. S., Siedentop, B., Bonhoeffer, S. & Lehtinen, S. Plasmid co-infection: linking
biological mechanisms to ecological and evolutionary dynamics. *Philosophical Transactions of the*
Royal Society B **377**, 20200478 (2022).
44   Ravel, J. *et al.* Vaginal microbiome of reproductive-age women. *Proceedings of the National Academy*
of Sciences **108**, 4680-4687 (2011).
45   Ding, T. & Schloss, P. D. Dynamics and associations of microbial community types across the human
body. *Nature* **509**, 357-360 (2014).
46   Lahti, L., Salojärvi, J., Salonen, A., Scheffer, M. & De Vos, W. M. Tipping elements in the human intestinal
ecosystem. *Nature Communications* **5**, 4344 (2014).
47   Kootte, R. S. *et al.* Improvement of insulin sensitivity after lean donor feces in metabolic syndrome is
driven by baseline intestinal microbiota composition. *Cell Metabolism* **26**, 611-619. e616 (2017).
48   Canfora, E. E., Meex, R. C., Venema, K. & Blaak, E. E. Gut microbial metabolites in obesity, NAFLD and
T2DM. *Nature Reviews Endocrinology* **15**, 261-273 (2019).
49   Moustafa, A. *et al.* Genetic risk, dysbiosis, and treatment stratification using host genome and gut
microbiome in inflammatory bowel disease. *Clinical Translational Gastroenterology* **9**, e132 (2018).
50   Getino, M. & de la Cruz, F. Natural and artificial strategies to control the conjugative transmission of
plasmids. *Microbiology Spectrum* **6**, 10.1128/microbiolspec. mtbp-0015-2016 (2018).
51   Fernandez-Lopez, R. *et al.* Unsaturated fatty acids are inhibitors of bacterial conjugation. *Microbiology*
151, 3517-3526 (2005).
52   Getino, M. *et al.* Synthetic fatty acids prevent plasmid-mediated horizontal gene transfer. *mBio* **6**,
10.1128/mbio. 01032-01015 (2015).
53   Ripoll-Rozada, J. *et al.* Type IV traffic ATPase TrwD as molecular target to inhibit bacterial conjugation.
Molecular Microbiology **100**, 912-921 (2016).
54   Palencia-Gándara, C. *et al.* Conjugation inhibitors effectively prevent plasmid transmission in natural
environments. *MBio* **12**, 10.1128/mbio. 01277-01221 (2021).
55   Kosterlitz, O. *et al.* Estimating the transfer rates of bacterial plasmids with an adapted Luria–Delbrück
fluctuation analysis. *PLoS Biology* **20**, e3001732 (2022).
56   Lopatkin, A. J. *et al.* Persistence and reversal of plasmid-mediated antibiotic resistance. *Nature*
Communications **8**, 1689 (2017).
57   Dieterich, W., Schink, M. & Zopf, Y. Microbiota in the gastrointestinal tract. *Medical Sciences* **6**, 116
(2018).
58   Flemming, H.-C. & Wuertz, S. Bacteria and archaea on Earth and their abundance in biofilms. *Nature*
Reviews Microbiology **17**, 247-260 (2019).
797

Code Availability

All the codes and source data associated with the numerical simulations and analysis of this

manuscript are available at the Github repository:

(https://github.com/twang1993/MultistabilityByHGT).

Acknowledgements

This work is supported by Shenzhen Institute of Synthetic Biology Scientific Research

Program (Grant No. HSE199011086). The funder had no role in study design, data collection and

analysis, the decision to publish, or the preparation of the manuscript.

Declaration of interests

The authors declare no competing interests.

810
811