Dachsous and Fat coordinately repress the Dachs-Dlish-Approximated

complex to control growth

Hitoshi Matakatsu* and Richard G. Fehon*

Department of Molecular Genetics and Cell Biology

University of Chicago

Chicago, IL 60637, USA

*Co-corresponding authors

Primary corresponding author

Richard G. Fehon

Department of Molecular Genetics and Cell Biology

University of Chicago

Chicago, IL 60637, USA

Phone: 773-702-5694

e-mail; rfehon@uchicago.edu

Abstract

Two protocadherins, Dachsous (Ds) and Fat (Ft), regulate organ growth *in Drosophila* via the

Hippo pathway. Ds and Ft bind heterotypically to regulate the abundance and subcellular

localization of a 'core complex' consisting of Dachs, Dlish and Approximated. This complex

localizes to the junctional cortex where it promotes growth by repressing the pathway kinase

Warts. Ds is believed to promote growth by recruiting and stabilizing the core complex at the

junctional cortex, while Ft represses growth by promoting degradation of core complex

components.

Here, we examine the functions of intracellular domains of Ds and Ft and their relationship to the

core complex. While Ds promotes accumulation of the core complex proteins in cortical puncta,

it is not required for core complex assembly. Indeed, the core complex assembles maximally in

the absence of both Ds and Ft. Furthermore, while Ds promotes growth in the presence of Ft, it

represses growth in the absence of Ft by removing the core complex from the junctional cortex.

Ft similarly recruits core complex components, however it normally promotes their degradation.

Our findings reveal that Ds and Ft constrain tissue growth by repressing the default 'on' state of

the core complex.

Introduction

The ability to precisely control the subcellular localization of proteins, particularly at the cell cortex, is critical to the function of diverse intra- and intercellular signaling pathways. This property is well demonstrated by the Hippo pathway, which controls organ growth during development (Zheng and Pan, 2019; Karaman and Halder, 2018; Boggiano and Fehon, 2012). Hippo pathway kinases, including Tao, Hippo and Warts (Wts), function in a cascade to promote phosphorylation of the transcriptional co-factor Yorkie (Yki), resulting in Yki repression and decreased transcription of its pathway targets. While the kinases are normally diffusely cytoplasmic, previous studies have revealed that their assembly and activity are controlled by cortically localized protein complexes (Boggiano and Fehon, 2012; Karaman and Halder, 2018). Expanded, a FERM-domain protein, associates with the transmembrane protein Crumbs at the junctional cortex to assemble and activate the core kinases (Ling et al., 2010; Chen et al., 2010; Robinson et al., 2010). In parallel, Merlin (another FERM domain protein) and Kibra together assemble the same kinases at the apical medial cortex (Su et al., 2017).

Perhaps the most enigmatic cortically localized protein complex controlling Hippo pathway output is the Dachsous (Ds)-Fat (Ft) system. Ds and Ft are giant protocadherins that regulate planar cell polarity, proximo-distal patterning and organ growth (Mahoney et al., 1991; Clark et al., 1995; Bryant et al., 1988; Adler et al., 1998; reviewed in Blair and McNeill, 2018; Strutt and Strutt, 2021). Ft and Ds bind extracellularly via heterotypic interaction (Matakatsu and Blair, 2004; Ma et al., 2003; Strutt and Strutt., 2002). Opposing proximal-distal expression gradients of Ds and Four-jointed (Fj), a Golgi resident kinase that modulates Ds-Ft binding (Ma et al., 2003; Zeidler et al., 2000; Brittle et al., 2010; Simon et al., 2010), result in a planar polarized

subcellular distribution of Ds on distal junctions and Ft on proximal junctions in wing discs

(Brittle et al., 2012; Bosveld et al., 2012; Ambegaonkar et al., 2012).

Ft and Ds are thought to control tissue growth primarily via an atypical myosin, Dachs at the

junctional cortex (Mao et al., 2006; Rogulja et al., 2008). Dachs promotes growth by inhibiting

Wts function, by degrading Wts and/or regulating Wts conformational state (Cho et al., 2006;

Vrabioiu and Struhl., 2015). Two additional proteins, <u>D</u>achs <u>L</u>igand with <u>SH</u>3 domains (Dlish,

aka Vamana) and a DHHC palmitoyltransferase Approximated (App) are required for Dachs

function and localization to the junctional cortex (Matakatsu and Blair, 2008; Matakatsu et al.,

2017; Zhang et al., 2016; Misra and Irvine, 2016). Dlish forms a complex with Dachs, Ft and Ds

when expressed in S2 cells, and Dlish and Dachs are interdependent for cortical localization *in*

vivo (Zhang et al., 2016; Misra and Irvine, 2016). App palmitoylates both Dlish and Ft

(Matakatsu et al., 2017; Zhang et al., 2016). Palmitoylation often increases the affinity of a -

protein for the plasma membrane (Linder and Deschenes, 2007), leading to the suggestion that

App promotes Dlish-dependent Dachs localization to the junctional cortex (Zhang et al., 2016).

Since the accumulation of Dachs at the cortex is influenced by Ds-Ft planar polarity, it has been

proposed that Ds recruits Dachs to the distal edge to promote growth while Ft promotes Dachs

degradation at the proximal edge to suppress growth (Rogulja et al., 2008; Bosveld et al., 2012;

Brittle et al., 2012). However, this model seems inconsistent with the observation that *ds ft*

double mutants have a much stronger overgrowth phenotype than either single mutant alone,

which instead suggests that Ft and Ds work synergistically to repress growth (Matakatsu and

Blair., 2006).

To understand how Ds-Ft signaling regulates the abundance, localization and function of

Dachs, Dlish and App (hereafter called the core complex) at the junctional cortex, we explored

the functions of the intracellular domains of Ds and Ft. We find that members of the core

complex, in particular App, depend on the Ds ICD to accumulate in cortical puncta and promote

tissue growth. In contrast, in the absence of its ligand Ft, Ds (via its ICD) promotes

mislocalization and inactivation of the core complex. Strikingly, loss of both Ds and Ft allows

the core complex to accumulate to very high levels at the junctional cortex, indicating that

neither Ds nor Ft is required for its cortical localization. Instead, our results suggest that Ds and

Ft organize core complex proteins within the junctional cortex into discrete puncta with opposing

consequences: Ft promotes degradation of the core complex whereas Ds protects it from

degradation. Together these findings reveal previously unrecognized mechanisms by which Ds

and Ft control the subcellular localization and abundance of Dachs, Dlish and App at the

junctional cortex and thereby constrain tissue growth.

Results

Ft and Ds act synergistically to repress Yki activity

Current models propose that Ds and Ft function antagonistically. However, mutation of either

gene causes overgrowth of wing discs (Fig. 1A, B and F) and combining mutations in both genes

produces much greater overgrowth (Fig. 1G, Bryant et al., 1988; Matakatsu and Blair, 2006).

Together, these results suggest that rather than functioning antagonistically Ds and Ft function

synergistically to repress tissue growth.

To determine if Ds and Ft synergistically repress growth via the Hippo pathway, we compared

the effect of *ds ft* double mutants to *ft* single mutant cells on the *ban3-GFP, a* reporter of Yki

activity (Matakatsu and Blair, 2012). We observed that expression of *ban3-GFP* is upregulated

to a greater extent in *ds ft* than in *ft* cells (Fig. 1H, I). Thus, Ds and Ft synergistically repress

growth by repressing Yki activity.

Ft and Ds restrict tissue growth by regulating the abundance of Dachs at the junctional

cortex

The finding that Ft and Ds function synergistically to restrict growth seems at odds with the

model that Ds promotes Dachs accumulation at the junctional cortex while Ft promotes its

degradation. To explore this paradox, we examined Dachs localization and abundance in *ds ft*

double mutants relative to *ds* or *ft* single mutants. Compared to wild type, we found that

Dachs:GFP accumulated at slightly greater levels in *ds* mutant tissue (Fig. 1J) and was even

more abundant in *ft* mutant tissue (Fig. 1J). Additionally, we noticed that in *ft* mutant cells Dachs

was distributed in puncta across the apical cortex except in mutant cells at the clone borders (Fig.

1J). We infer this effect might occur because Ds protein in *ft* mutant cells at the edge of clones

interacts with Ft and is tightly localized at the boundary with adjacent wild-type cells, while in

the middle of *ft* clones Ds is mislocalized across the apical cortex (Strutt and Strutt, 2002; Ma et

al., 2003; see below).

If Ds is necessary to recruit Dachs to the junctional cortex, then loss of both Ds and Ft

should result in decreased Dachs cortical accumulation in comparison to loss of just Ft alone.

Remarkably, we instead found that *ds ft* cells accumulated Dachs to even higher levels than those

lacking Ft alone (Fig. 1K). Furthermore, the non-junctional Dachs puncta we observed in *ft*

mutant cells were absent in *ds ft* double mutant cells (Fig. 1K). These results clearly indicate that

neither Ds nor Ft is required for cortical localization of Dachs. Additionally, the increased level

of cortical Dachs and absence of non-junctional Dachs puncta in *ds ft* double mutant cells

suggests that Ds can promote removal of Dachs from the junctional cortex. Given that *ds ft*

double mutant discs are significantly more overgrown than *ft* mutants alone (Fig. 1F, G,

Matakatsu and Blair, 2006), the non-junctional Dachs we observed in *ft* mutant cells likely does

not promote growth.

To determine how loss of Ft and/or Ds affects Dachs abundance, we compared Dachs

protein levels in *ds* null, *ft* null and *ds ft* null double mutant imaginal tissue using immunoblots

(Fig. 1L, M). While *ds* mutant tissues displayed slightly higher levels of Dachs, presumably

because loss of Ds causes decreased Ft activity (Sopko et al., 2009; Feng et al., 2009) Dachs

abundance was significantly greater in *ft* mutant tissue than either *ds* or wild-type tissues.

However, in *ds ft* double mutant tissues, Dachs levels were similar to *ft* mutants (Fig. 1L, M),

suggesting that increased cortical Dachs in *ds ft* double mutants relative to *ft* mutants (Fig. 1K)

reflects changes in localization rather than abundance. These results also indicate that increased

junctional accumulation of Dachs is critical to growth. Consistent with this idea, loss of one copy

of *dachs* strongly suppressed *ds ft* overgrowth (Fig. 1E, G, Fig. S1A) and homozygous loss of

dachs was epistatic to it (Fig. 1C, D, G).

The Dachsous intracellular domain regulates growth in a context-dependent manner

Previous studies have shown that in the absence of Ft, Ds becomes mislocalized from the

junctions and is much more diffusely distributed across the apical membrane (Strutt and Strutt,

2002; Ma et al., 2003; Fig. 1K). Because Ds and Dachs are known to physically interact

(Bosveld et al., 2012), mislocalized Ds might recruit Dachs away from the junctional cortex,

thereby reducing growth. Careful examination of Ds and Dachs in *ft* mutant cells revealed that

both displayed non-junctional puncta that often co-localized (Fig. 1N), consistent with the

possibility that Ds can remove Dachs from the junctional cortex, perhaps through endocytosis.

To better understand how Ds affects Dachs subcellular localization, we next focused on the

intracellular domain of Ds, which has been proposed to form a complex with Dachs and Dlish

(Bosveld et al., 2012 and 2016; Zhang et al., 2016; Misra and Irvine, 2016). To dissect the

function of Ds, we used CRISPR-Cas9 to replace the Ds ICD with GFP (Ds$^{\Delta ICD:GFP}$; Fig. 2A).

Ds$^{\Delta ICD:GFP}$ expression levels and localization were indistinguishable from wild-type Ds (Fig. 2B,

C and Fig. S1B, C). Ft also was normally localized in $ds^{\Delta ICD:GFP}$ cells, suggesting that Ds$^{\Delta ICD:GFP}$

binds normally with Ft (Fig. 2D). In marked contrast to ds null wings and wing discs, which are

slightly overgrown, $ds^{\Delta ICD:GFP}$ wings and wing discs were smaller than normal (Fig. 2I-K). This

result is surprising given our evidence that Ds can repress growth by removing Dachs from the

junctional cortex, but likely reflects the complex nature of Ds function in growth control: in the

presence of Ft, Ds functions both to promote growth by binding and protecting Dachs from Ft-

mediated degradation and to reduce growth by binding and activating Ft. If so, then loss of ft

should be epistatic to $ds^{\Delta ICD:GFP}$. Indeed, the $ds^{\Delta ICD}$ ft double mutant combination produced

severe overgrowth similar to that of ds ft double mutant combination (Fig. 2M, compare with ds

ft double mutant in Fig. 1G). Taken together, our results indicate that the function of the Ds ICD

in growth control is context-dependent: in the presence of Ft the Ds ICD promotes Dachs activity

while in the absence of Ft the Ds ICD represses Dachs activity.

The intracellular domain of Dachsous organizes Dachs into cortical puncta.

How does the Ds ICD regulate growth? Dachs and Dlish are known to co-localize with Ds in

puncta at the junctional cortex in wing discs and to associate with the Ds ICD *in vitro* (Bosveld

et al., 2012 and 2016; Brittle et al., 2012; Zhang et al., 2016; Misra and Irvine, 2016). Consistent

with these studies, we found a strong correlation between wild-type Ds and Dachs in puncta at

the junctional cortex (Fig. 3A, B). However, in $ds^{\Delta ICD}$ mutant clones, Dachs lost its punctate

appearance and was irregularly localized at the junctional cortex (Fig. 3C). Moreover, unlike

wild-type Ds, Ds$^{\Delta ICD:GFP}$ did not strongly colocalize with Dachs (Fig. 3D-E).

In the absence of ft, Dachs colocalizes with Ds in non-junctional puncta (Fig. 1N). To ask if

the Ds ICD is necessary for this effect, we examined Dachs in $ds^{\Delta ICD}$ ft double mutant tissue. We

found that non-junctional Dachs puncta were dramatically reduced and instead Dachs

accumulated strongly at the junctional cortex, similar to ds ft double mutants (Fig. 3F). Given

that $ds^{\Delta ICD}$ ft and ds ft double mutants display similar overgrowth phenotypes (Figs. 1G and 2M),

these results suggest that the Ds ICD associates with Dachs, thereby influencing Dachs'

localization and growth-promoting function at junctional cortex.

Dachsous regulates Approximated localization at the junctional cortex

App encodes a multi-pass transmembrane palmitoyltransferase that is thought to regulate both Ft

and Dlish through post-translational palmitoylation (Matakatsu et al., 2017a; Zhang et al., 2016).

App is also localized to the junctions (Matakatsu and Blair, 2008), but what determines its

localization is unclear. To visualize App, we tagged it at the C-terminus with YFP (App:YFP) or

in the extracellular domain with V5 (App:V5) using CRISPR-Cas9 (Fig. S1D-F). Both tagged

alleles are homozygous viable, indicating the tags do not interfere with App function. App:YFP

localized to junctional puncta in a similar pattern to Dachs and Ds and was predominantly

localized at proximodistal junctions in wing discs (Fig. 4A-D), reminiscent of Dachs, Dlish and

Ds planar polarized distribution (Brittle et al., 2012; Ambegaokar et al., 2012; Misra and Irvine,

2016). In fact, App colocalized with Dachs, Ds and Dlish (Fig. 4A-D and Fig. S1G, see below).

We next explored App's relationship with Ft and Ds. In *ft* mutant cells, App displayed

decreased junctional accumulation but retained its punctate appearance, similar to Ds (Fig. 4E).

In *ds* mutant cells, App displayed junctional localization and lacked distinct puncta (Fig. 4F).

Notably, App distribution differed significantly from both Dachs and Dlish, which displayed

increased accumulation at the junctional cortex in either *ft* or *ds* mutant cells (Fig. 1J, Bosveld et

al., 2012; Brittle et al., 2012; Zhang et al., 2016; Misra and Irvine, 2016). These observations

suggest that Dachs and Dlish are regulated differently from App and that App is more directly

dependent upon Ds for junctional localization.

Previous studies have shown that due to their heterophilic interactions, Ft and Ds

accumulate at the boundary of either *ds* or *ft* mutant clones (Strutt and Strutt, 2002; Ma et al.,

2003). We found that App also accumulated at these boundaries (Fig. 4E, F), raising the

possibility that Ft or Ds recruits App. To determine this, we used *ft* RNAi in the posterior

compartment of the wing (driven by *hh-Gal4*) to create polarized accumulation of Ds on the

posterior side of the boundary and Ft on the anterior (wild-type) side. Conversely, *ds* RNAi in

the posterior caused Ft to accumulate on the posterior side of the boundary and Ds to accumulate

on the anterior (wild-type) side. As we observed in somatic mosaic clones (Fig. 4E, F), App

accumulated at the boundary between wild-type cells and either *ft* (Fig. 4G) or *ds* knockdown

tissue (Fig. 4I). To determine on which side of the boundary App accumulated, we used double

RNAi to deplete both App and either Ft or Ds. Co-depletion of Ft and App in the posterior

strongly reduced accumulation of App at the boundary (Fig. 4H), indicating that App

accumulates with Ds in the posterior (Ft-depleted) cells at the boundary. Conversely, co-

depletion of Ds and App did not affect App accumulation at the boundary (Fig. 4J), indicating

that when Ds is depleted App accumulates in adjacent wild-type cells at the boundary. In short,

these results show that Ds recruits App to these boundaries while Ft does not.

Three further observations suggest that the Ds ICD interacts with App. First, App puncta did

not form at the junctional cortex in $ds^{\Delta ICD}$ mutant clones (Fig. 4K). Second, App co-

immunoprecipitated with the Ds ICD in S2 cells (Fig. 4L). Third, when we ectopically expressed

Ds in the posterior compartment we found that as with Dachs, this resulted in repolarization and

stabilization of App in cells just anterior to the compartment boundary (Fig. 4M, Bosveld et al.,

2012; Brittle et al., 2012; Ambegaonkar et al., 2012). Taken together, our results are consistent

with the idea that the Ds ICD coordinately recruits and stabilizes Dachs and App (and possibly

Dlish) in puncta at the junctional cortex.

Mapping functional domains in the Ds ICD.

We next mapped functional domains in the Ds ICD using a combination of cell culture and *in*

vivo experiments. To identify binding domains in the Ds ICD, we used a previously described

approach to polarize protein domains at the cortex of cultured S2 cells (Johnston et al., 2009;

Bosveld et al., 2016; Tokamov et al., 2023). Specifically, we fused the full-length or truncated

Ds ICD to the extracellular domain of the homophilic adhesion protein Echinoid (Ed), which

causes cells to aggregate and accumulate the fusion protein at points of contact. Using this

approach, we asked which domains in the Ds ICD are necessary to recruit core complex proteins

(Fig. 5A). In control cells, the Ed extracellular domain fused to Flag (Ed:Flag) accumulated at

cell-cell contacts but co-transfected core complex proteins instead localized to the cytoplasm or

to cytoplasmic foci (Fig. S2A). In contrast, Ed fused to Ds ICD (Ed:Ds:Flag) recruited core

complex proteins to cell contacts (Fig. 5B, C).

To determine the essential domains for core complex recruitment by Ds ICD, we first

focused on two non-overlapping domains we denote A and B (Fig. 5A), because each contains

conserved motifs (CM1-3) found in mammalian Ds (Hulpiau and Roy, 2009). Core complex

proteins were recruited to cell contacts by domain B (Ed:Ds ΔA:Flag; Fig. 5D), but not by

domain A (Ed:Ds ΔB:Flag; Fig. 5E). We then defined a smaller region within B (domain D) that

includes CM2 and is necessary for recruitment (Fig. 5A, F, Fig. S2B-E).

To examine the functional significance of these domains *in vivo*, we generated fly lines

lacking these portions using CRISPR-Cas9 (Fig. 5G-S). Interestingly, $ds^{\Delta A:Flag}$, $ds^{\Delta B:Flag}$, $ds^{\Delta C:Flag}$

and $ds^{\Delta D:Flag}$ all led to undergrowth in the wing that was milder than that of $ds^{\Delta ICD:GFP}$ (Fig. 5G-

L, M). Additionally, we noted that $ds^{\Delta B:Flag}$, $ds^{\Delta C:Flag}$ and $ds^{\Delta D:Flag}$ displayed defects in leg tarsal

segmentation commonly observed in Ft signaling mutants while $ds^{\Delta A:Flag}$ did not (Fig. 5N-S).

Thus, while both domain A and B are involved in growth control, domain B alone is sufficient

for the Ds-Ft signaling functions that pattern the leg.

To determine if results from our S2 cell experiments reflect *in vivo* functions, we asked if

Ds deletions recruit the core complex to junctional puncta (Fig. 5T-X). The expression pattern,

protein level and subcellular distribution of all Ds mutant proteins were indistinguishable from

endogenous Ds or Ds:Flag (Fig. S3A-D). While Dachs colocalized with Ds:Flag and DsΔA:Flag

at junctional puncta (Fig. 5T-U), it did not colocalize with DsΔB:Flag and DsΔD:Flag (Fig. 5V-

W). Moreover, App is dramatically reduced in $ds^{\Delta B:Flag}$ and $ds^{\Delta D:Flag}$ (Fig. 5X and data not

shown). Taken together, our results suggest that domain D is responsible for core complex

recruitment to the junctional cortex while domain A might control growth in parallel through

other effectors (see discussion). Consistent with this idea, animals carrying the heteroallelic

combinations $ds^{\Delta A:Flag}/ds^{\Delta B:Flag}$ or $ds^{\Delta A:Flag}/ds^{\Delta D:Flag}$ showed normal tissue growth in the wing and

leg patterning (Fig. S3E-J).

The Fat intra- and extracellular domains have opposing functions.

In the absence of Ft, we propose that Ds, which is no longer anchored junctionally, represses

growth by removing Dachs from the junctional cortex. If so, then restoring Ds junctional

localization, without restoring Ft's ability to degrade Dachs, should strongly enhance tissue

growth. To do this, we deleted the Ft ICD using CRISPR-Cas9 ($ft^{\Delta ICD}$; Fig. S4A). Interestingly,

we observed much greater overgrowth in $ft^{\Delta ICD}$ wing discs compared to ft null mutant tissues

(Fig. 6A-C). Consistent with the idea that Ft$^{\Delta ICD}$ promotes growth by holding Ds (and Dachs) at

the junctional cortex, we found that in $ft^{\Delta ICD}$ mutant cells Ds was localized in an almost normal

pattern (Fig. 6D), in striking contrast to Ds in ft null mutant tissue (Fig. 1K, 4E). Additionally,

although Dachs was mislocalized in non-junnctional puncta in ft null mutant cells (Fig. 1J, K),

Dachs was exclusively localized to the junctional cortex in $ft^{\Delta ICD}$ mutant cells (Fig. 6D). App

appeared normally localized at the junctional cortex in $ft^{\Delta ICD}$ mutant cells (Fig. 6E), in striking

contrast to $ds^{\Delta ICD}$, ds null mutant cells (Fig. 4F, K).

To further identify functions of the Ft ICD, we examined the effects of mutations that

severely diminish activity of the Ft ICD without affecting its extracellular interactions with Ds.

ft^{sum} and ft^{61} previously were identified as point mutations in the Ft ICD that cause overgrowth

(Bossuyt et al., 2014; Bosch et al., 2014). Examining ft^{sum} and ft^{61} mutant clones, we found

increased junctional accumulation of both Dachs and App in a punctate pattern (Fig. 6F and Fig.

S4B, C). These observations are in contrast to $ft^{\Delta ICD}$ mutant clones, where App levels were

unaffected (Fig. 6E) and Dachs was evenly distributed along the junctional cortex (compare Fig.

6D-F). Clones of dco^3 (discs-overgrown) cells, which also decrease activity of the Ft ICD by

blocking its phosphorylation (Sopko et al., 2009; Feng and Irvine, 2009), had a similar effect on

App (Fig. 6G). Interestingly, Dco fails to promote phosphorylation of Ft^{sum} in S2 cells (Fig.

S4D), suggesting that phosphorylation of Ft decreases core complex accumulation at junctional

puncta.

While the Ft ICD clearly has a role for reducing core protein levels, there is evidence that it

recruits core proteins to the junction before degrading them. In fact, the Ft ICD forms a complex

with core complex proteins including Dlish and App in S2 cells (Matakatsu et al., 2017; Zhang et

al., 2016; Misra and Irvine, 2016). To further examine this possibility, we made use of the planar

polarized distribution of Ft and Ds in the imaginal epithelium, where Ft accumulates primarily at

the proximal junctions of cells and Ds accumulates primarily at the distal edge. By examining

mosaic clones of cells expressing tagged forms of Ds, Ft, Dachs or App, we could discern at

which junction, distal or proximal, these proteins accumulate in either wild-type or dco^3

backgrounds. As previously suggested, we found that Ds and Dachs accumulated at the distal

junctions of wild-type imaginal epithelial cells (Fig. 6H), while Ft accumulated on the proximal

side (Fig. 6J-K, Mao et al., 2006; Rogulja et al., 2008; Brittle et al., 2012; Bosveld et al., 2012).

Additionally, although App colocalized with Ds at distal junctions (Fig. 6 I), Ft did not

colocalize with App on proximal junctions suggesting that App localized distally with the rest of

the core complex (Fig. 6K). In contrast, when similar clones were generated in the dco^3 mutant

background, Dachs and App colocalized with Ft at the proximal junctions (Fig. 6L, M). Taken

together these observations suggest that like Ds, Ft recruits App and Dachs to the junctional

cortex but prevents their retention by promoting their degradation.

Mapping functional domains in the Ft ICD.

To further examine the functions of the Ft ICD, we employed the Echinoid S2 cell assay to

examine the ability of Ed:Ft ICD fusions to recruit core complex components to the cell cortex

(Fig. 7A). Just as we observed with Ds ICD fusions, Ed:Ft:Flag strongly recruited Dachs, App

and Dlish (Fig. 7A). Recruitment of Dachs did not appear to depend on other core complex

proteins, as RNAi-mediated depletion of Dlish and App did not affect Dachs recruitment

(Fig.7B). Similarly, we found that Dachs co-immunoprecipitated with the Ft ICD and that

depletion of App and Dlish did not affect their co-immunoprecipitation (Fig. 7C). These results

suggest that while the Ft ICD can recruit Dachs, Dlish and App to the cortex of S2 cells, it likely

cannot promote their degradation, presumably because other proteins essential for this process

are not expressed or are inactive in S2 cells.

　　　Previous studies have revealed the domains in the Ft ICD responsible for PCP and growth

control *in vivo* (Fig. 7D, Matakatsu et al., 2012). To better understand how different Ft ICD

domain function, we next examined their role in core complex recruitment (Fig. 7 D-H).

Ed:FtΔPH-C:Flag, which lacks most of the ICD but retains a domain known to function in planar

cell polarity (Matakatsu and Blair, 2012), failed to recruit any of the core complex proteins (Fig.

7E). To narrow down regions necessary for recruitment, we generated two smaller deletions, one

containing the Hippo N and C domains previously shown to be sufficient for growth control

(Ed:FtΔ6-C:Flag) and one that deletes those domains but retains regions that modulate Hippo

pathway function (Ed:FtΔN-6:Flag; Matakatsu and Blair, 2012; see Fig. 7D). Interestingly, we

found that either region (referred to as the M and C domains, Fig. 7D) was sufficient to recruit

core complex proteins, consistent with previous work showing that both domains co-

immunoprecipitate with App and Dlish in S2 cells (Fig. 7F-H, Zhang et al. 2016; Misra and

Irvine, 2016).

The relationship between Dachs, Dlish and App

The different effects of loss of Ft and Ds on Dachs, Dlish and App led us to further explore the

relationship between these proteins. In the simplest case, these three proteins act together in a

complex to promote growth. Consistent with this idea, Dachs interacts with Dlish *in vitro*,

ectopic Dachs expression strongly stabilizes Dlish at the junctional cortex, and vice versa (Zhang

et al., 2016; Misra and Irvine, 2016). Additionally, loss of App disrupts cortical localization of

Dachs and Dlish (Matakatsu and Blair, 2008; Matakatsu at el., 2017; Zhang et al., 2016; Misra

and Irvine, 2016). However, the relationship between Dachs or Dlish and App at the junctional

cortex is less well studied. To address this, we expressed Dachs or Dlish and found that either

promotes accumulation of App at the junctional cortex (Fig. 8A, C and Fig. S5A). In contrast,

Dachs and Dlish levels changed only slightly in response to ectopic App expression, though they

did have a more punctate appearance (Fig. 8B, C, Fig. S5B). Thus, while ectopic Dachs or Dlish

can promote App accumulation, ectopic App has little effect on Dachs or Dlish.

Although Ft and Ds affect accumulation of Dachs, App and Dlish at the junctional cortex,

these proteins accumulate at the highest levels in the complete absence of *ds* and *ft* (Fig. 8D, E).

Moreover, loss of *dachs* is epistatic to the *ds ft* overgrowth phenotype (Fig. 1D) and loss of either

dlish or *app* partly suppresses this phenotype (Fig. S5C-I), indicating that these proteins can

function together, independent of Ds and Ft, to promote growth. Therefore, we reasoned that it

should be easier to understand how these proteins regulate their cortical accumulation and

promote growth in *ds ft* mutant background.

We started by examining Dlish and App in *ds ft dachs* triple mutant cells. In this

background, Dlish was severely reduced from apical cortex (Fig. 8F), while App appeared more

diffuse but still largely junctional (Fig. 8G). Loss of Dlish blocked the strong accumulation of

Dachs normally seen in *ds ft* mutant cells (Fig. 8H). Taken together, these data suggest that

Dachs and Dlish are strongly interdependent for their localization to the junctional cortex, while

App can localize junctionally independent of Dachs and Dlish. However, we note that in *ds ft*

dachs triple mutant cells App abundance is reduced relative to *ds ft* cells (compare Fig. 8E and

G), suggesting that levels of Dachs and Dlish can affect App.

To explore the role of App in junctional recruitment of Dachs and Dlish we next examined

these proteins in *ds ft app* triple mutant cells, either by removing *ds* and *ft* in an *app* mutant

background (Fig. 8I) or by removing *app* in *ds ft* mutant background (Fig. 8J). Loss of *app* in *ds*

ft background showed that App promotes accumulation of Dachs and Dlish at the junctional

cortex in the absence of Ft and Ds (Fig. 8J), while loss of *ds* and *ft* in an *app* mutant background

clearly showed that App is not absolutely required for Dachs and Dlish accumulation (Fig. 8 I).

These observations are consistent with previous evidence that App palmitoylates Dlish *in vitro*

and promotes junctional localization of Dlish as well as Dachs *in vivo*, since palmitoylation can

promote association of cytoplasmic proteins with the plasma membrane (Zhang et al., 2016). To

test this more directly, we examined the effect of app^{DHHS}, a CRISPR engineered allele that

blocks the palmitoyltransferase activity of App (Matakatsu et al., 2017), on Dachs and Dlish

accumulation and found it to be indistinguishable from an *app* null allele (Fig. 8K). These results

suggest that App's primary role is to promote the accumulation of Dachs and Dlish at the

junctional cortex through its enzymatic function rather than a structural one.

To examine interactions between core complex components, we modified the Echinoid S2

cell assay to work with any GFP-tagged protein by fusing an anti-GFP nanobody (Caussinus et

al., 2012) to the cytoplasmic domain of Echinoid (pMT-Ed:Flag:vhh4) or to a myristoylation

sequence (UAS-myr:Flag:vhh4). Ed:Flag:vhh4 expressed in S2 cells promoted cell aggregation

and strongly accumulated at cell contacts, as shown previously for other Ed protein fusions

(Johnston et al., 2009), and recruited co-transfected GFP (Fig. S5J). Ed:Flag:vhh4 strongly

recruited Dachs:GFP to sites of cell contact, which in turn recruited co-transfected App and

Dlish (Fig. 8L). This result was confirmed using myr:Flag:vhh4 (Fig. S5K). Additionally, Dlish

and App could recruit other core complex components when they were tagged with GFP (Fig.

S5L-O) in S2 cells. Thus, interactions among Dachs, Dlish and App help core complex

formation in S2 cells.

Discussion

In this study we have examined functional interactions between the intracellular domains of Ds

and Ft and their effectors in growth control, a 'core complex' consisting of Dachs, Dlish and

App. Previous work has suggested that while Ds functions to recruit core complex components

to the cell cortex, thereby promoting growth, Ft promotes degradation of Dachs, thereby

repressing growth (Mao et al., 2006; Bosveld et al., 2012; Brittle et al., 2012). The results

presented here revise this model in several important ways. First, we show that the core complex

requires neither Ft nor Ds to associate with the junctional cortex and drive growth. Rather, Ft and

Ds function coordinately to precisely restrict the amount and distribution of core complex

components at the junctional cortex. Second, we show that in the absence of Ft, Ds functions to

repress growth, apparently by removing core complex components from the junctional cortex.

This finding is surprising and suggests that in some tissue contexts Ds could play a significant

role in restricting tissue growth. Third we demonstrate that Ft also can recruit core complex

components to the junctional cortex and that the stability and/or duration of these interactions is

controlled by phosphorylation of the Ft ICD. Our structure/function analysis of Ft and Ds shows

that they interact with core complex components via specific regions of their ICDs and that

different regions of the Ds ICD function in parallel to promote growth. Taken together, our work

reveals the underlying mechanisms by which Ds-Ft signaling represses a default 'on' state that is

driven by the inherent ability of core complex components to localize to the junctional cortex

and promote growth.

The Dachs-Dlish-App core complex localizes at the junctional cortex to promote growth

independent of Ft and Ds

Previous studies have found a strong correlation between the ability of Dachs to localize to

the junctional cortex and its ability to promote tissue growth (Mao et al., 2006). The results

presented here strongly support this model. In particular, we find that although overall Dachs

abundance is indistinguishable between *ft* mutant tissues and *ds ft* mutant tissues, Dachs

abundance at the junctional cortex and overall tissue growth are much greater in the latter than in

the former. Additionally, the finding that Dachs, Dlish and App all accumulate at the junctional

cortex to the greatest extent in *ds ft* double mutant tissues demonstrates that core complex

proteins accumulate at the junctional cortex independently of Ft and Ds.

To better understand how the core complex associates with the cell cortex, we examined

interdependencies between these proteins in the absence of either Ft or Ds. Dachs and Dlish are

highly interdependent-removing one results in almost complete loss of the other from the cell

cortex, consistent with previous studies showing that Dachs and Dlish form a complex (Zhang et

al., 2016; Misra and Irvine, 2016). App has also been shown to form a complex with Dachs and

Dlish and we showed here that Dachs and Dlish can recruit App in S2 cells, but its relationship

with these proteins at the cortex appears less direct. For example, while ectopic expression of

Dachs or Dlish causes increased accumulation of App, ectopic expression of App has little effect

on either Dach or Dlish. Additionally, loss of App reduces but does not prevent increased

accumulation of Dachs and Dlish caused by absence of Ft and Ds. These observations are

consistent with a model where App promotes the interaction of Dachs and Dlish with the cell

cortex but does not contribute structural support for this interaction. Consistent with this idea,

App can palmitoylate Dlish (Zhang et al., 2016) and we found that a mutation that specifically

affects App enzymatic activity (app^{DHHS}) affects Dachs and Dlish recruitment similar to an app

null allele. Nonetheless, we consider App to be a member of the core complex because 1) co-IP

and the Ed recruitment assays in S2 cells demonstrate that all three proteins can form a complex

and 2) loss of Dachs severely diminishes App accumulation at the cell cortex (Matakatsu et al.

2017).

Ds effects on growth are context dependent

Previous studies have highlighted the importance of Ds in stabilizing Dachs accumulation at

the junctional cortex to promote growth (Bosveld et al., 2012; Brittle et al., 2012). While our

findings are consistent with this model, they also show that in some contexts Ds can function to

remove Dachs from the junctional cortex and as a result inhibit growth. This aspect of Ds

function is clearly observed in the absence of Ft. As previously described, we observed that loss

of Ft results in dramatic mislocalization of Ds, suggesting that Ds-Ft heterophilic interactions

stabilize both proteins at the junctional cortex. We observed Ds and Dachs in non-junctional

punctae that appeared to be in an endocytic pathway in the absence of *ft*. These punctae were the

consequence of Ds being unable to bind Ft on adjacent cells rather than a cell autonomous effect

of loss of Ft because *ft* mutant cells at the borders of clones, where Ds could interact with Ft

expressed on adjacent wild-type cells, did not display non-junctional Ds or Dachs. Additionally,

these punctae were absent in in $ds^{\Delta ICD}$ *ft* cells indicating that interaction with the Ds ICD is

necessary to remove Dachs from the junctional cortex.What is most intriguing about these

observations is that they provide a cell biological basis for the observation that while Ds

promotes growth in the presence of Ft, it represses growth in the absence of Ft. While Ft is

believed to play the major role in repressing Dachs activity and *ft* mutants display significant

overgrowth, *ds ft* double mutants display much greater overgrowth. These observations raise the

still unanswered question of whether, in some contexts, Ds might function to repress growth,

independent of its role in activating Ft, during normal development. In relation to this it is

interesting that the opposing gradients of Ds and Fj expression in the wing result in less stable

Ds-Ft binding in the proximal wing. Further studies will be required to elucidate to what extent

Ds represses growth independent of Ft in development.

The Ft ICD can recruit the core complex to the cell cortex

Although Ft is well recognized to promote Dachs degradation, the underlying mechanistic

basis of this function is poorly understood. Using the Echinoid S2 cell assay, we found that the

Ft ICD functions similarly to the Ds ICD in having the ability to recruit core complex

components to the cell cortex, consistent with previous co-immunoprecipitation experiments

(Matakatsu et al., 2017; Zhang et al., 2016; Misra and Irvine, 2016). We further found that while

the Ft ICD can recruit Dachs in the absence of App and Dlish, deletion of either the M or C

domain prevents recruitment of Dachs unless App and Dlish are present (Fig. 7 F, G and Fig

S4E-G). This result suggests that multiple Ft ICD domains could be involved in forming a

complex between Ft and the core complex. Using endogenously tagged alleles and surface

labeling protocols we found that in wing discs under normal conditions core complex

components accumulate with Ds, as has been suggested previously, but do not accumulate with

Ft. However, in the absence of Dco, which phosphorylates the Ft ICD and promotes its ability to

degrade Dachs, core complex components do accumulate with Ft at cell contacts. Thus, our

results suggest that Ft degrades Dachs by first recruiting the core complex in a similar manner to

Ds.

Functions of the Ds intracellular domain

Using CRISPR we have performed an extensive structure/function analysis of the Ds ICD.

Removal of the Ds ICD has no observable effect on the abundance or localization of either Ds or

Ft, but does dramatically affect the core complex. Consistent with the idea that the Ds ICD

promotes core complex activity, $ds^{\Delta ICD}$ mutants display marked undergrowth. Interestingly both

Dachs and App lost their normal punctate appearance in $ds^{\Delta ICD}$ clones, suggesting a key role for

Ds in organizing these puncta and that they normally promote Dachs function in promoting

growth.

Fine scale structure/function studies mapped the domains responsible for growth control

and interaction with the core complex. Previous studies identified three motifs (CM1-3) in the

ICD that are conserved between *Drosophila* Dachsous and its vertebrate orthologs Dachsous 1

and Dachsous 2 (Hulpiau and van Roy, 2009). Using the Echinoid S2 cell assay and CRISPR

deletion mutants *in vivo*, we found that domain D is necessary for core complex recruitment.

Additionally, deletion of domain D leads to proximo-distal patterning defects and undergrowth,

similar to mutation of core complex components. Domain D contains the highly conserved CM2

motif (Hulpiau and van Roy, 2009). In zebrafish, tetratrico peptide repeat containing protein 28

(Ttc-28) was shown to interact with the CM2 motif in Dachsous-1b (Chen et al., 2018). Although

its functions have not been studied, a Ttc-28 homologue (CG43163) exists in *Drosophila*.

Interestingly, while other regions of the Ds ICD are not necessary for core complex recruitment,

they also display growth defects, suggesting that these regions might facilitate core complex

recruitment or interact with other proteins such as Riquiqui, Lowfat and Myo31DF that have

been implicated in Ds-Ft signaling (Degoutin et al., 2013; Mao et al., 2009; González-Morales et

al., 2015).

Concluding remarks

A striking feature of Ft, Ds and core complex localization that is emphasized by our imaging

of endogenously tagged components is that all of these proteins predominantly accumulate in

distinct puncta along the junctional cortex. Previous experiments have shown that the mobility of

Ft and Ds in these puncta is dependent on their interaction with one another (Hale et al., 2015),

suggesting that each punctum consists of Ds and Ft accumulated in trans on adjacent cells. We

show in addition that all components of the core complex localize to these puncta in association

with Ds. Furthermore, their formation is not an inherent property of the core complex but rather

is driven by Ft and Ds because in *ds ft* mutant cells cortical Dachs, Dlish and App all appear

much more uniformly distributed along the junctional cortex. It is still not known how these

puncta form, but our observations that 1) App localization is essentially normal in $ft^{\Delta ICD}$ mutants

while Dachs is uniformly distributed at the junctional cortex (Fig. 6 D-E) and 2) App is severely

mislocalized in $ds^{\Delta ICD}$ cells (Fig. 4 K) suggests that App is recruited to puncta by Ds. A possible

model consistent with our observations is that App lateral movement within the plasma

membrane is restricted through interaction with the Ds ICD where it then promotes accumulation

of Dachs and Dlish (Fig. 9). When Ds is bound to Ft in trans these interactions result in stable

puncta that promote growth ('ON' state), while in the absence of Ft binding Ds promotes

removal of the core complex from the junctional cortex ('OFF' state). The Ft ICD similarly

recruits these proteins but their interaction is more transient due to Ft-mediated degradation of

Dachs at junctional cortex ('OFF' state). Further studies will be required to test this model, but it

provides a framework for understanding the molecular and functional interactions between Ds,

Ft and the core complex in regulating organ growth.

Materials and methods

Fly genetics

To induce mitotic clones, larvae were incubated at 37 ℃ for 40-120 min in a water bath (Xu and

Rubin, 1993). The following stocks were used:

vasa-cas9 (Gratz et al., 2013; Sebo et al., 2014), ds^{UAO71}(Adler et al., 1996), ds^{05142}(BL11394),

$dachs^{GC13}$, $dlish^{Y003}$ (Wang et al., 2019), app^{12-3}, app^{DHHS}, ft^{fd} , ft^{G-rv}, ft^{sum} (Bossuyt et al., 2014),

ft^{61} (Bosch et al., 2014), $ft^{\Delta ICD}$, ds^{UAO71} ft^{G-rv}, ds^{UAO71} ft^{fd}, *hh-gal4, ban3-GFP, $app^{:YFP}$, $app^{:V5}$,*

dachs:GFP (Bosveld et al., 2012), *ds:GFP* (Brittle., 2012), $ds^{\Delta ICD:GFP}$, $ds^{\Delta ICD}$, $ds^{:Flag}$, $ds^{\Delta A:Flag}$,

$ds^{\Delta B:Flag}$, $ds^{\Delta C:Flag}$, $ds^{\Delta D:Flag}$, *ft:GFP* (VDRC 318477), *V5:ft*, *HA:ds* (Ambegaonkar et al., 2012),

UAS-dlish:Flag, UAS-dachs:V5 (Mao et al., 2006), *UAS-ds, UAS-app, ft RNAi* (VDRC 9396), *ds*

RNAi (VDRC 36219), *app RNAi* (BL 35280), *y w hs-FLP; FRT 2A, y w hs-FLP; ubi-RFP FRT*

2A, *y w hs-FLP; ubi-GFP FRT 2A, y w hs-FLP; ubi-GFP FRT 40a, y w hs-FLP; ubi-RFP FRT*

40a, TKO.GS00382 (BL79393) and *w^{1118}*.

Molecular cloning

To construct pMT-*ed:ds:Flag* and pMT-*ed:ft:Flag*, the pMT plasmid backbone and Echinoid

extracellular domain was amplified from pMT-*ed:GFP:apkC* (Johnston et al., 2009) with iProof

high-Fidelity DNA Polymerase (BIO-RAD). The intracellular and transmembrane domain of *ds*

and *ft* were amplified from pUAST-*dsTMICD:Flag* (Zhang et al., 2016) or *Ft STI-4:FVH* (Feng

and Irvine, 2009). DNA fragments were assembled with Gibson assembly. Deletions of these

constructs were constructed using the Q5 site-directed mutagenesis kit according to the

manufacturer protocol (NEB). For construction of pMT-*ed:Flag:vhh4*, the GFP nanobody

sequence was amplified from pUAST-attB-NSlmb-vhhGFP4 (Caussinus et al., 2012, Addgene

35577) and fused to the Ed ECD as just described. All resulting plasmids were verified with

DNA sequencing. Detailed information is available by request.

Immunostaining

All imaging was done on fixed tissues or S2 cells. The following primary antibodies were used:

Rat anti-Dachs (1:20,000; Matakatsu et al., 2017), Guinea pig anti-App (1:10,000; Matakatsu

and Blair, 2008), Rabbit anti-Dlish (1:1,000; Zhang et al., 2016), Rabbit anti-Ds (1:1,000; Strutt

and Strutt, 2002), Rat or Guinea pig anti-Ft (1:1,000; Sopko et al., 2009), Rat anti-*D*ECad

(DCAD2, 1:1,000; DHSB; Oda et al, 1994) , Rat anti-Ci (2A1, 1:10; Motzny and Holmgren,

1995), Rabbit anti-HA (1:1,000; Abcam), Rabbit anti-HA (1:1,000; Y-11, Santa Cruz), iFluor

555 or iFluor 647 conjugated Mouse anti-V5 (1:100, GenScript), Mouse anti-Flag (1:20,000;

M2, Millipore Sigma), Mouse anti-β-galactosidase (JIE7, 1:1,000; DHSB) and Alexa Fluor 647

Phalloidin (1:1,000; Thermo Fisher). Fluorescent secondary antibodies were obtained from

Jackson laboratory or Invitrogen. After staining, wing discs were mounted in ProLong Diamond

Antifade Mountant (Thermo Fisher). Images were taken using an LSM 880 confocal microscope

(Zeiss, using Zen software), using Plan Apochromat 20x/0.8 na, Plan Apochromat 40x/1.4 na

and Plan Apochromat 63x/1.4 na objectives. The acquired images were processed with Image J.

Apical z-stacks were combined to make maximal projections using ImageJ.

App:V5 extracellular staining was performed as previously described (Hale et al., 2015). Wing

discs were dissected at 5.5 h after puparium formation and incubated in iFluor 555 or iFluor 647

conjugated Mouse anti-V5 in dissection medium for 30 min on ice. After washing with

dissection medium, wing discs were fixed with 2% formaldehyde in PBS for 10 min at room

temperature, washed with PBS, and mounted. Images for whole wing discs were taken using tile

scanning. Individual scanned images were aligned and combined into single images with

Photoshop (Adobe) or Affinity designer (Serif).

In vitro **studies**

S2 cells were transfected according to Han (1996). pUAST constructs were co-transfected with

pAWGal4 (kindly provided Y. Hiromi). To induce metallothionein promoter expression, $CuSO_4$

was added to culture medium at a final concentration of 0.7 mM 24 h after transfection. S2 cells

expressing pMT Ed constructs were allowed to aggregate as described previously (Matakatsu

and Blair, 2004). After cells were agitated on a rotator at 70 rpm for 30-60 min, cells were

attached on poly-L-lysine coated cover slips and fixed with 2 % formaldehyde in PBS for 10 min

at room temperature. Cells were stained according to standard procedures.

Western blotting and immunoprecipitation

Western blotting and immunoprecipitation are performed as previously described (Matakatsu et

al., 2017). For immunoprecipitation of HA or Flag tagged proteins, anti-HA magnetic beads

(Pierce) and anti-FLAG M2 agarose beads (Millipore Sigma) were used. Quantification of Dachs

protein on immunoblots was performed as previously described (Matakatsu et al., 2017). The

following primary antibodies were used for visualization of proteins: Rat anti-Dachs (1:10,000;

Matakatsu et al., 2017), Mouse anti- **α**Tubulin (1:20,000; Millipore Sigma), Guinea pig anti-App

(1:10,000; Matakatsu and Blair, 2008), Mouse anti-Flag M2 (1:20,000; Millipore Sigma), Mouse

anti-V5 (1:10,000; Genscript), Anti-V5 tag antibody [SV5-P-K](1:5,000; Abcam, ab206558) and

Rabbit anti-GFP (1:2,000; kindly provided Dr. M. Glotzer). Blots were imaged using a LI-COR

Biosciences Odyssey CLx. Efficiency of depletion with *app* and *dlish* dsRNA was confirmed by

immunoblot. All experiments were performed at least three times and reproduced consistent

results.

Gene editing with the CRISPR-Cas9 system

The target sequence for CRISPR-Cas9 editing was designed according to flyCRISPR Optimal

Target Finder (https://flycrispr.org/target-finder/). Target sequences were cloned into pU6-BbsI-

chiRNA (Gratz et al., 2013; Addgene Plasmid 45946). The resulting plasmids were injected with

or without donor plasmid into *vasa-Cas9* expressing embryos (Sebo et al., 2014, Gratz et al.,

2013). Injections were performed by Genetivision.

To replace the Ds ICD with EGFP, EGFP was cloned between 5' and 3' homology arms into

pBluescript II KS using Gibson assembly. The resulting plasmids and a pair of gRNA plasmids

(pU6 target 4 for 5' and pU6 target 1 for 3') were co-injected. To obtain *ds* ICD deletions (Fig.

5A), the same pair of gRNA plasmids and appropriate donor plasmids containing Ds ICD

deletions were co-injected.

To tag App with YFP, PCR amplified YFP sequence was cloned between 5' and 3' homology

arms into pBluescript II KS (Fig. S1E). The resulting donor plasmid and gRNA plasmids (pU6

app target 2 and pU6 *app* target 3) were co-injected as before. The internal V5 tagging for extra

cellular domain of App is summarized in Fig. S1F. gRNA plasmid (pU6 *app* V5 gRNA) and

single stranded DNA oligonucleotides (*app* 3x V5) were co-injected as described. To make indel

mutants for *ft*$^{\Delta ICD}$, a gRNA plasmid (pU6 *ft* 5' target 5) was injected as described (Fig. S4A). The

resulting indel mutations caused premature stop codons after the transmembrane domain. After

evaluation with anti-Ds staining, one mutant (24-3-5) was selected for further analysis.

After screening by genomic PCR with flanking primers, the isogenized lines were sequenced to

determine the exact CRISPR event for all gene editing.

To generate *ds*$^{\Delta ICD}$ that lacks GFP fluorescence, the GFP coding sequence in *ds*$^{\Delta ICD:GFP}$ was

disrupted by crossing with flies expressing *vasa-Cas9* and a *gRNA* for GFP (BL79393).

Image Quantification

Wing from >20 females for each genotype were mounted on a slide glass in Permount mounting

medium (Fisher). Imaging was performed using a Zeiss Axioplan 2 microscope, using an EC

Plan-Neo 5x objective. The images were taken using EOS REBEL T2i camera. The areas of

wing blade were manually measured using ImageJ.

To quantify expression levels for App:YFP, Dachs:GFP and anti-Dlish staining in Fig. 8C, the

mean gray value for a 25 x 25 µm square in the anterior (wild type) and posterior

(overexpression) compartments in the proximal area of the wing disc (N>11) was determined

using ImageJ.

Statistical analysis

Minimum and Maximum are shown as bars in all box plots. All data points for box plots are

shown as dots except Fig. 5M. Graphs were generated using GraphPad Prism 10. Statistical

significances were evaluated using unpaired two-tailed t tests. Line scans were performed with

ImageJ and quantified with Graph Pad Prism10.

Acknowledgements

637
We thank Drs. S. Blair, H. McNeill, D. Strutt, K. Irvine, Y. Bellaïche, G. Halder, I. Hariharan, J.

Price, the Bloomington and Vienna Stock Centers, and the Developmental Studies Hybridoma

Bank for reagents. We also thank M. Wang, W. J. Desrosiers for help with some experiments,

and S. Blair, S. Tokamov and A. Smith for critical reading of the manuscript.

This research was funded by NIH research grant GM134128 to R. Fehon.

The authors declare no competing financial interests.

Author contribution: Conceptualization and methodology, H. Matakatsu and R. G. Fehon;

Investigation, H. Matakatsu; Interpretation of data, H. Matakatsu and R. G. Fehon; Original

draft, H. Matakatsu and R. G. Fehon; Writing, review and editing, H. Matakatsu and R. G.

Fehon. Funding acquisition, R. G. Fehon, Project administration and Supervision, R. G. Fehon.

References

Adler, P.N., Charlton, J., Liu, J., 1998. Mutations in the cadherin superfamily member gene *dachsous* cause

a tissue polarity phenotype by altering *frizzled* signaling. Development 125, 959–968.

https://doi.org/10.1242/dev.125.5.959

Ambegaonkar, A.A., G. Pan, M. Mani, Y. Feng, and K.D. Irvine. 2012. Propagation of Dachsous-Fat

Planar Cell Polarity. *Curr Biol*. 22:1302–1308. doi:10.1016/j.cub.2012.05.049.

Blair, S., and H. McNeill. 2018. Big roles for Fat cadherins. *Curr Opin Cell Biol*. 51:73–80.

doi:10.1016/j.ceb.2017.11.006.

Boggiano, J.C., and R.G. Fehon. 2012. Growth Control by Committee: Intercellular Junctions, Cell Polarity,

and the Cytoskeleton Regulate Hippo Signaling. *Dev. Cell*. 22:695–702. doi:10.1016/j.devcel.2012.03.013.

Bosch, J.A., T.M. Sumabat, Y. Hafezi, B.J. Pellock, K.D. Gandhi, and I.K. Hariharan. 2014. The

Drosophila F-box protein Fbxl7 binds to the protocadherin Fat and regulates Dachs localization and Hippo

signaling. *Elife*. 3:e03383. doi:10.7554/elife.03383.

Bossuyt, W., C.-L. Chen, Q. Chen, M. Sudol, H. McNeill, D. Pan, A. Kopp, and G. Halder. 2014. An

evolutionary shift in the regulation of the Hippo pathway between mice and flies. *Oncogene*. 33:1218–1228.

doi:10.1038/onc.2013.82.

Bosveld, F., I. Bonnet, B. Guirao, S. Tlili, Z. Wang, A. Petitalot, R. Marchand, P.-L. Bardet, P. Marcq,

F. Graner, and Y. Bellaïche. 2012. Mechanical Control of Morphogenesis by Fat/Dachsous/Four-Jointed

Planar Cell Polarity Pathway. *Science*. 336:724–727. doi:10.1126/science.1221071.

Bosveld, F., B. Guirao, Z. Wang, M. Rivière, I. Bonnet, F. Graner, and Y. Bellaïche. 2016. Modulation of

junction tension by tumor suppressors and proto-oncogenes regulates cell-cell contacts. *Development*.

143:623–634. doi:10.1242/dev.127993.

Brittle, A.L., A. Repiso, J. Casal, P.A. Lawrence, and D. Strutt. 2010. Four-Jointed Modulates Growth and

Planar Polarity by Reducing the Affinity of Dachsous for Fat. *Curr Biol*. 20:803–810.

doi:10.1016/j.cub.2010.03.056.

Brittle, A., C. Thomas, and D. Strutt. 2012. Planar Polarity Specification through Asymmetric Subcellular

Localization of Fat and Dachsous. *Curr Biol*. 22:907–914. doi:10.1016/j.cub.2012.03.053.

Bryant, P.J., B. Huettner, L.I. Held, J. Ryerse, and J. Szidonya. 1988. Mutations at the *fat* locus interfere

with cell proliferation control and epithelial morphogenesis in *Drosophila*. *Dev Biol*. 129:541–554.

doi:10.1016/0012-1606(88)90399-5.

Caussinus, E., O. Kanca, and M. Affolter. 2012. Fluorescent fusion protein knockout mediated by anti-GFP

nanobody. *Nat Struct Mol Biol*. 19:117–121. doi:10.1038/nsmb.2180.

Chen, C.-L., K.M. Gajewski, F. Hamaratoglu, W. Bossuyt, L. Sansores-Garcia, C. Tao, and G. Halder.

2010. The apical-basal cell polarity determinant Crumbs regulates Hippo signaling in *Drosophila*. *Proc. Natl.*

Acad. Sci. 107:15810–15815. doi:10.1073/pnas.1004060107.

Chen, J., G.D. Castelvecchi, N. Li-Villarreal, B. Raught, A.M. Krezel, H. McNeill, and L. Solnica-

Krezel. 2018. Atypical Cadherin Dachsous1b Interacts with Ttc28 and Aurora B to Control Microtubule

Dynamics in Embryonic Cleavages. *Dev. Cell*. 45:376-391.e5. doi:10.1016/j.devcel.2018.04.009.

Cho, E., Y. Feng, C. Rauskolb, S. Maitra, R. Fehon, and K.D. Irvine. 2006. Delineation of a Fat tumor

suppressor pathway. *Nat Genet*. 38:1142–1150. doi:10.1038/ng1887.

Clark, H.F., D. Brentrup, K. Schneitz, A. Bieber, C. Goodman, and M. Noll. 1995. Dachsous encodes a

member of the cadherin superfamily that controls imaginal disc morphogenesis in *Drosophila*. *Gene Dev*.

9:1530–1542. doi:10.1101/gad.9.12.1530.

Degoutin, J.L., C.C. Milton, E. Yu, M. Tipping, F. Bosveld, L. Yang, Y. Bellaiche, A.

Veraksa, and K.F. Harvey. 2013. Riquiqui and Minibrain are regulators of the Hippo pathway

downstream of Dachsous. *Nat Cell Biol*. 15:1176–1185. doi:10.1038/ncb2829.

Feng, Y., and K.D. Irvine. 2009. Processing and phosphorylation of the Fat receptor. *Proc National Acad Sci*.

106:11989–11994. doi:10.1073/pnas.0811540106.

González-Morales, N., C. Géminard, G. Lebreton, D. Cerezo, J.-B. Coutelis, and S. Noselli. 2015. The

Atypical Cadherin Dachsous Controls Left-Right Asymmetry in *Drosophila*. *Dev Cell*. 33:675–689.

doi:10.1016/j.devcel.2015.04.026.

Gratz, S.J., A.M. Cummings, J.N. Nguyen, D.C. Hamm, L.K. Donohue, M.M. Harrison, J. Wildonger,

and K.M. O'Connor-Giles. 2013. Genome Engineering of *Drosophila* with the CRISPR RNA-Guided Cas9

Nuclease. *Genetics*. 194:1029–1035. doi:10.1534/genetics.113.152710.

Hale, R., A.L. Brittle, K.H. Fisher, N.A.M. Monk, and D. Strutt. 2015. Cellular interpretation of the long-

range gradient of Four-jointed activity in the *Drosophila* wing. *Elife*. 4:e05789. doi:10.7554/elife.05789.

Han, K., 1996. An Efficient DDAB-Mediated Transfection of *Drosophila* S2 Cells. Nucleic Acids Res. 24,

4362–4363. https://doi.org/10.1093/nar/24.21.4362

Hulpiau, P., and F. van Roy. 2009. Molecular evolution of the cadherin superfamily. *Int. J. Biochem. Cell*

Biol. 41:349–369. doi: 10.1016/j.biocel.2008.09.027.

Johnston, C.A., K. Hirono, K.E. Prehoda, and C.Q. Doe. 2009. Identification of an Aurora-A/PinsLINKER/

Dlg Spindle Orientation Pathway using Induced Cell Polarity in S2 Cells. *Cell.* 138:1150–1163.

doi:10.1016/j.cell.2009.07.041.

Karaman, R., and G. Halder. 2018. Cell Junctions in Hippo Signaling. *Cold Spring Harb. Perspect.*

Biol. 10:a028753. doi:10.1101/cshperspect.a028753.

Linder, M.E., and R.J. Deschenes. 2007. Palmitoylation: policing protein stability and traffic. *Nat. Rev. Mol.*

Cell Biol. 8:74–84. doi:10.1038/nrm2084.

Ling, C., Y. Zheng, F. Yin, J. Yu, J. Huang, Y. Hong, S. Wu, and D. Pan. 2010. The apical transmembrane

protein Crumbs functions as a tumor suppressor that regulates Hippo signaling by binding to Expanded. *Proc.*

Natl. Acad. Sci. 107:10532–10537. doi:10.1073/pnas.1004279107.

Ma, D., C. Yang, H. McNeill, M.A. Simon, and J.D. Axelrod. 2003. Fidelity in planar cell polarity

signalling. *Nature*. 421:543–547. doi:10.1038/nature01366.

Mao, Y., B. Kucuk, and K.D. Irvine. 2009. *Drosophila lowfat,* a novel modulator of Fat

signaling. *Development*. 136:3223–3233. doi:10.1242/dev.036152.

Mao, Y., C. Rauskolb, E. Cho, W.-L. Hu, H. Hayter, G. Minihan, F.N. Katz, and K.D. Irvine. 2006.

Dachs: an unconventional myosin that functions downstream of Fat to regulate growth, affinity and gene

expression in *Drosophila*. *Development*. 133:2539–2551. doi:10.1242/dev.02427.

Mahoney, P.A., U. Weber, P. Onofrechuk, H. Biessmann, P.J. Bryant, and C.S. Goodman. 1991. The fat

tumor suppressor gene in *Drosophila* encodes a novel member of the cadherin gene superfamily. *Cell*. 67:853–

868. doi:10.1016/0092-8674(91)90359-7.

Matakatsu, H., and S.S. Blair. 2004. Interactions between Fat and Dachsous and the regulation of planar cell

polarity in the *Drosophila* wing. *Development*. 131:3785–3794. doi:10.1242/dev.01254.

Matakatsu, H., and S.S. Blair. 2006. Separating the adhesive and signaling functions of the Fat and Dachsous

protocadherins. *Development*. 133:2315–2324. doi:10.1242/dev.02401.

Matakatsu, H., and S.S. Blair. 2008. The DHHC Palmitoyltransferase Approximated Regulates Fat Signaling

and Dachs Localization and Activity. *Curr Biol*. 18:1390–1395. doi:10.1016/j.cub.2008.07.067.

Matakatsu, H., and S.S. Blair. 2012. Separating planar cell polarity and Hippo pathway activities of the

protocadherins Fat and Dachsous. *Development*. 139:1498–1508. doi:10.1242/dev.070367.

Matakatsu, H., S.S. Blair, and R.G. Fehon. 2017. The palmitoyltransferase Approximated promotes growth

via the Hippo pathway by palmitoylation of Fat. *J Cell Biol*. 216:265–277. doi:10.1083/jcb.201609094.

Misra, J.R., and K.D. Irvine. 2016. Vamana Couples Fat Signaling to the Hippo Pathway. *Dev Cell*. 39:254–

266. doi:10.1016/j.devcel.2016.09.017.

Motzny, C.K., and R. Holmgren. 1995. The *Drosophila* cubitus interruptus protein and its role in the

wingless and hedgehog signal transduction pathways. *Mech. Dev.* 52:137–150. doi:10.1016/0925-

4773(95)00397-j.

Oda, H., T. Uemura, Y. Harada, Y. Iwai, and M. Takeichi. 1994. A Drosophila Homolog of Cadherin

Associated with Armadillo and Essential for Embryonic Cell-Cell Adhesion. *Dev Biol*. 165:716–726.

doi:10.1006/dbio.1994.1287.

Robinson, B.S., J. Huang, Y. Hong, and K.H. Moberg. 2010. Crumbs Regulates Salvador/Warts/Hippo

Signaling in *Drosophila* via the FERM-Domain Protein Expanded. *Curr. Biol.* 20:582–590.

doi:10.1016/j.cub.2010.03.019.

Rogulja, D., C. Rauskolb, and K.D. Irvine. 2008. Morphogen Control of Wing Growth through the Fat

Signaling Pathway. *Dev Cell*. 15:309–321. doi: 10.1016/j.devcel.2008.06.003.

Sebo, Z.L., H.B. Lee, Y. Peng, and Y. Guo. 2014. A simplified and efficient germline-specific CRISPR/Cas9

system for *Drosophila* genomic engineering. *Fly*. 8:52–57. doi:10.4161/fly.26828.

Simon, M.A., A. Xu, H.O. Ishikawa, and K.D. Irvine. 2010. Modulation of Fat:Dachsous Binding by the

Cadherin Domain Kinase Four-Jointed. *Curr Biol*. 20:811–817. doi:10.1016/j.cub.2010.04.016.

Sopko, R., E. Silva, L. Clayton, L. Gardano, M. Barrios-Rodiles, J. Wrana, X. Varelas, N.I. Arbouzova,

S. Shaw, S. Saburi, H. Matakatsu, S. Blair, and H. McNeill. 2009. Phosphorylation of the Tumor

Suppressor Fat Is Regulated by Its Ligand Dachsous and the Kinase Discs Overgrown. *Curr Biol*. 19:1112–

1117. doi:10.1016/j.cub.2009.05.049.

Strutt, H., and D. Strutt. 2002. Nonautonomous Planar Polarity Patterning in *Drosophila* Dishevelled-

Independent Functions of Frizzled. *Dev Cell*. 3:851–863. doi:10.1016/s1534-5807(02)00363-5.

Strutt, H., and D. Strutt. 2021. How do the Fat–Dachsous and core planar polarity pathways act together and

independently to coordinate polarized cell behaviours? *Open Biol*. 11:200356. doi:10.1098/rsob.200356.

Su, T., M.Z. Ludwig, J. Xu, and R.G. Fehon. 2017. Kibra and Merlin Activate the Hippo Pathway Spatially

Distinct from and Independent of Expanded. *Dev. Cell*. 40:478-490.e3. doi:10.1016/j.devcel.2017.02.004.

Tokamov, S.A., N. Nouri, A. Rich, S. Buiter, M. Glotzer, and R.G. Fehon. 2023. Apical polarity and

actomyosin dynamics control Kibra subcellular localization and function in *Drosophila* Hippo signaling. *Dev.*

cell. 58:1864-1879.e4. doi:10.1016/j.devcel.2023.08.029.

Vrabioiu, A.M., and G. Struhl. 2015. Fat/Dachsous Signaling Promotes *Drosophila* Wing Growth by

Regulating the Conformational State of the NDR Kinase Warts. *Dev Cell*. 35:737–749.

doi:10.1016/j.devcel.2015.11.027.

Wang, X., Y. Zhang, and S.S. Blair. 2019. Fat-regulated adaptor protein Dlish binds the growth suppressor

Expanded and controls its stability and ubiquitination. *Proc National Acad Sci*. 116:201811891.

doi:10.1073/pnas.1811891116.

Xu, T., and G.M. Rubin. 1993. Analysis of genetic mosaics in developing and adult *Drosophila* tissues. *Dev*

Camb Engl. 117:1223–37. doi:10.1242/dev.117.4.1223.

Zeidler, M.P., N. Perrimon, and D.I. Strutt. 2000. Multiple Roles for four-jointed in Planar Polarity and

Limb Patterning. *Dev Biol*. 228:181–196. doi:10.1006/dbio.2000.9940.

Zhang, Y., X. Wang, H. Matakatsu, R. Fehon, and S.S. Blair. 2016. The novel SH3 domain protein

Dlish/CG10933 mediates fat signaling in Drosophila by binding and regulating Dachs. *Elife*. 5:e16624.

doi:10.7554/elife.16624.

Zheng, Y., and D. Pan. 2019. The Hippo Signaling Pathway in Development and Disease. *Dev. Cell*. 50:264–

282. doi: 10.1016/j.devcel.2019.06.003.

Figures legends

Figure 1. Both Ft and Ds suppress tissue growth by regulating localization and abundance

of Dachs at the junctional cortex.

(A-G) Representative wing discs stained with phalloidin showing the growth defects of the

following genotypes: wild-type (A), ds^{UAO71} (B), $dachs^{GC13}$ (C), ds^{UAO71} ft^{G-rv} $dachs^{GC13}/ds^{UAO71}$

ft^{fd} $dachs^{GC13}$ (D), $ds^{UAO71}ft^{G-rv}$ $dachs^{GC13}/ds^{UAO71}$ ft^{fd} (E), ft^{G-rv}/ft^{fd} (F) and $ds^{UAO71}ft^{G-rv}/ds^{UAO71}ft^{fd}$

(G). (H, I) Expression of *ban3-GFP*, a Hippo pathway reporter, compared between ft^{G-rv}/ft^{fd} (or

ds^{UAO71} ft^{G-rv}/ft^{fd}) cells and ds^{UAO71} ft^{Gr-v} cells in the wing disc. *ban3-GFP* is more highly

expressed in ds^{UAO71} ft^{G-rv} cells than in *ft* mutant cells. *ds ft* clones are marked by the absence of

Ds staining. (I) XZ section indicated by the blue dashed line in H. (J) Comparison of Dachs:GFP

level and subcellular localization in wild type, ds^{UAO71} and ft^{fd} mutant cells. In *ds* mutant cells

Dachs:GFP levels were slightly increased and primarily localized to the junctional cortex (J and

J'). In *ft* mutant cells, Dachs:GFP levels increased more than in *ds* cells and localized both at the

junctional cortex and to non-junctional puncta (J and J"). (K, K') In comparison to *ft* cells, *ds ft*

mutant cells displayed even more Dachs:GFP at the junctional cortex but lacked the non-

junctional puncta. (L, M) Immunoblot analysis showing Dachs levels in imaginal tissue from

wild-type, ds^{UAO71}(N=5), ft^{G-rv}/ft^{fd} (N=16) and $ds^{UAO71} ft^{G-rv}/ds^{UAO71} ft^{fd}$ (N=9). Dachs abundance

increases slightly in *ds* but is much greater in *ft* and *ds ft* imaginal tissue. (M) Relative Dachs

levels normalized to α-Tubulin. **** P<0.0001, *** 0.0001<P<0.001, ns not significant, (two-

tailed unpaired t-test between each genotype). (N) Localization of Ds and Dachs:GFP in the

absence of Ft. Dachs:GFP was highly localized to the junctional cortex and colocalized with Ds

in non-junctional puncta. Scale bars: (A) 100 µm, (H-K, N) 5 µm

Figure 2. The Ds ICD has context-dependent effects on tissue growth.

(A) Structure of $Ds^{\Delta ICD:GFP}$. The intracellular domain was replaced with GFP (green) using

CRISPR-Cas9. ICD, intracellular domain (red line) and TM, transmembrane domain (sky blue).

(B and C) Localization of $Ds^{:GFP}$ in a ds^{GFP} homozygote (B) and $Ds^{\Delta ICD:GFP}$ in a $ds^{\Delta ICD:GFP}$

homozygote (C). Ds:GFP and $Ds^{\Delta ICD:GFP}$ localize at the junctional cortex. Adherens junctions are

marked by *D*E-Cad staining. (D) Ft localization is normal in $ds^{\Delta ICD:GFP}$ mutant cells. Mutant

clones are marked with the absence of nuclear GFP. (E-G) Representative adult wings in

indicated genotypes. The $ds^{\Delta ICD:GFP}$ wing displays undergrowth (F). (H) Quantification of wing

size in wild-type, $ds^{\Delta ICD:GFP}$ and $ds^{\Delta ICD}$ (GFP-) mutants. **** P<0.0001, ns not significant, (two-

tailed unpaired t-test between each genotype). (I-M) Growth of wing discs with the indicated *ds*

and *ft* alleles. Genotypes: wild type (I), $ds^{\Delta ICD:GFP}$ (J), ds^{UAO71} (K), ft^{G-rv}/ft^{fd} (L) and $ds^{\Delta ICD:GFP} ft^{G-}$

$^{rv}/ds^{\Delta ICD:GFP} ft^{fd}$ (M). Although loss of the Ds ICD leads to undergrowth (J), in combination with

loss of *ft* mutant it causes severe overgrowth (M). Scale bars, (B-C) 1 µm, (D) 5 µm, (E) 100 µm

and (I-M) 10 µm.

Figure 3. The Ds ICD influences Dachs localization.

(A) Ds:GFP and Dachs colocalize at junctional puncta (green arrows) in wing discs. (B) Fluorescence intensity of Dachs and Ds:GFP along the yellow line in A showing strong colocalization. (C) In $ds^{\Delta ICD}$ clones, Dachs:GFP loses its punctate appearance and becomes more uniformly distributed along the junctional cortex (the absence of nuclear RFP serves as clonal marker). (D) Localization of Dachs and Ds$^{\Delta ICD:GFP}$ in a $ds^{\Delta ICD:GFP}$ homozygote. Dachs fails to colocalize with Ds$^{\Delta ICD:GFP}$ (green arrows indicate Ds$^{\Delta ICD:GFP}$ puncta). (E) Fluorescence intensity of Dachs and Ds$^{\Delta ICD:GFP}$ along the yellow line in D showing weak colocalization in comparison to B (Pearson's correlation coefficient r=0.135 vs r=0.795). (F-F") Dachs:GFP accumulates strongly and almost exclusively at the junctional cortex in $ds^{\Delta ICD}ft$ mutant clones. Note the absence of non-junctional puncta of Dachs:GFP here in comparison to ft mutant cells in Figure 1J". Scale bars, (A, C", D, F") 1 µm (C, F) 5 µm.

Figure 4. Ds and Ft regulate App at cell junctions.

(A-D) Ds, Dachs and App colocalize at junctional puncta. Representative image (A) and line scan quantification of fluorescence intensity (B) showing colocalization of App:YFP and Dachs:GFP. Representative image (C) and line scan quantification of fluorescence intensity (D) showing colocalization of Ds:GFP and App:YFP. (E) Loss of ft causes mislocalization of Ds and App:YFP from junctional puncta to a more uniform distribution across the apical surface (clonal marker not shown). Note that App:YFP accumulates at the boundary between wild-type and ft mutant cells and appears reduced in the cytoplasm of ft cells at the edge of the clone (yellow dots). (F) The localization of App:YFP is altered in ds mutant cells. App:YFP is more diffuse in the absence of Ds and accumulates at the clone boundary (clonal marker not shown). Note that

cytoplasmic App:YFP is reduced in *ds* cells adjacent to the boundary (yellow dots). (G, H)

App:YFP accumulates at the boundary between wild-type and *ft* RNAi cells (G) but fails to

accumulate at the boundary in *ft*, *app* double RNAi cells (H), indicating that App in *ft* depleted

cells is recruited to contacts with wild-type cells. (I, J) App:YFP accumulates at the boundary

between wild-type and *ds* RNAi cells (I) and at the boundary between wild-type and *ds, app*

double RNAi cells (J), indicating that App in *ds* depleted cells is not recruited to contacts with

wild-type cells. (K) Junctional punctae of App:YFP are much less apparent in $ds^{\Delta ICD}$ clones. (L)

The Ds ICD co-immunoprecipitates App in S2 cells. (M) Ectopic expression of Ds in the

posterior compartment causes Dachs and App:YFP to colocalize in a planar polarized fashion for

2-3 cell diameters outside the Ds overexpression domain. Scale bars, 5 µm.

Figure 5. Mapping Ds functional domains.

(A) Structure of the Ds ICD and the deletions generated for the S2 cell recruitment assay and

CRISPR mutations *in vivo*. Full-length Ds ICD or Ds ICD deletions (ΔA,ΔB, ΔC and ΔD) were

fused to the Ed extracellular domain for S2 cell experiments. The same domains are deleted with

CRISPR-Cas9 *in vivo*. Abbreviations: Conserved motif 1-3 (CM1-3, red boxes): transmembrane

domain (TM, sky blue box). (B) Ed:Ds:Flag recruits Dachs:GFP, App and Dlish:HA to cell

contacts (yellow arrows). Ed homophilic interactions promote accumulation of the Ed fusion

proteins at cell-cell contacts. (C-F) Recruitment of Dachs:GFP by Ds ICD and its deletions.

Ed:Ds:Flag and Ed:DsΔA:Flag recruit Dachs:GFP to cell contacts (yellow arrows), but

Ed:DsΔB:Flag and Ed:DsΔD:Flag fail to recruit Dachs:GFP. (G-L) Adult wings of the indicated

genotypes. (M) Quantification of wing size in wild type and *ds* ICD deletion mutants. ****

P<0.0001, ns not significant, (two-tailed unpaired t-test between each genotype). (N-S) Tarsal

segments in legs from flies of the indicated genotypes. Numbers in the top right corner indicate

the percentage of animals with normal tarsal segments. (T-W) Dachs localization in wing discs

of the indicated genotypes. Dachs colocalizes with Ds in $ds^{:Flag}$ (T) and $ds^{\Delta A:Flag}$ (U), but they

colocalize less in $ds^{\Delta B:Flag}$ (V) and $ds^{\Delta D:Flag}$ (W). Yellow circles indicate Ds puncta. (X) App

puncta, stained extracellularly with a pulse-chase protocol (see Methods) are lost in $ds^{\Delta D:Flag}$

clones. Scale bars, (B, C, X) 5 µm, (T) 1µm, (G, N) 100 µm.

Figure 6. Effects of the Ft ICD on Dachs and App.

(A-C) Representative wing discs of the following genotypes stained with phalloidin: wild-type

(A), ft^{G-rv}/ft^{fd} (B) and $ft^{\Delta ICD}$ (C). Note that $ft^{\Delta ICD}$ has a more severe overgrowth phenotype than ft

null mutants. (D) Localization of Dachs and Ds in $ft^{\Delta ICD}$ mutant cells. Dachs:GFP expression is

elevated and uniformly distributed around at the junctional cortex, while Ds is only slightly

elevated and localizes more normally at the junctions. Note that Dachs does not display the non-

junctional puncta that are seen in ft null mutant cells (Fig. 1 J"). (E) Localization and abundance

of App:YFP is not altered in $ft^{\Delta ICD}$ mutant cells. (F) Dachs and App:YFP levels are elevated but

localization to junctional puncta does not change in ft^{sum} mutant cells. (G) App:YFP abundance

also is elevated at junctional puncta in dco^3 mutant cells. (H-M) Surface labeling of cell clones

expressing HA:Ds (H, I), App:V5 (I, K, M) or V5:Ft (J, L) together with the indicated tagged

proteins. All clones shown border non-expressing sister clones. (H) Dachs:GFP puncta

colocalize with HA:Ds at clone boundaries. (I) Similarly, HA:Ds in puncta colocalizes with

App:V5. (J) In contrast, V5:Ft puncta do not colocalize with Dachs:GFP at clone boundaries

(yellow ellipses). (K) Similarly, Ft:GFP puncta do not co-stain for App:V5 at clone boundaries

(yellow ellipses). Additionally, Ft:GFP fails to co-localize with App:V5 puncta at clone

boundaries on the opposite side of cells (red ellipses). (L) In contrast to wild type, in a *dco³*

mutant background V5:Ft and Dachs:GFP colocalize in puncta at clone boundaries (yellow

circles). (M) Similarly, in a *dco³* mutant background App:V5 and Ft:GFP extensively colocalize

in puncta at clone boundaries (yellow ellipses). Scale bars, 100 µm (A), 5 µm (D-M)

Figure 7. Interactions between the Ft ICD and core complex proteins in S2 cells.

(A) Recruitment of Dachs:V5, App:GFP and Dlish:HA by Ed:Ft:Flag in S2 cells. All three

proteins accumulate with the Ft ICD at cell contacts (yellow arrows). (B) Depletion of *app* and

dlish by RNAi does not affect the ability of the Ft ICD to recruit Dachs:GFP (yellow arrows).

(C) Co-immunoprecipitation experiments similarly show that the Ft ICD forms a complex with

Dachs in S2 cells even when App and Dlish are depleted. (D) Structure cartoons of Ft ICD

deletions designed to identify regions responsible for core complex recruitment in S2 cells.

Functional domains and App and Dlish association sites from previous studies are also shown.

(E-G) S2 cell assays for recruitment of core complex proteins by different Ft ICD deletions. (E)

Ed:FtΔPH-C:Flag fails to recruit core complex proteins to cell contacts. Ed:FtΔ6-C:Flag (F) as

well as Ed:FtΔN-6:Flag (G) recruit Dachs:GFP and Dlish:HA (yellow arrows). (H) Co-

immunoprecipitation experiments confirm the results in E-G. Scale bars, 5 µm.

Figure 8. Dachs, App and Dlish form a core complex to promote growth.

(A) Ectopic Dachs expression causes increased App abundance. (B) Conversely, ectopic App

expression does not increase Dachs abundance but does result in a more punctate appearance at

the junctional cortex. (C) Comparison of apical protein levels of Dachs, Dlish and App in

response to ectopic expression of other core complex components. Y axis shows the mean

intensity ratio of the posterior compartment (ectopic expression) to the anterior compartment

(normal cells). **** P<0.0001, ** 0.001<P<0.01, * 0.01<P<0.05, (two-tailed unpaired t-test

between each genotype).

(D-K) Analysis of core complex components at the cell cortex in the absence of Ds and Ft. (D)

Dachs and Dlish levels are highly elevated at the junctional cortex in *ds ft* mutant cells. (E)

App:YFP abundance also is increased at junctions together with Dachs in *ds ft* clones. (F) In

marked contrast to *ds ft* mutant cells (D''), Dlish is strongly reduced in *ds ft dachs* mutant cells.

(G) App:YFP is less punctate, but is junctionally localized in *ds ft dachs* mutant cells. (H) In the

absence of *dlish*, Dachs abundance and localization are unaffected by loss of *ds* and *ft*. (I) In

contrast, loss of *ds* and *ft* still causes increased Dachs and Dlish abundance at the junctional

cortex in the absence of *app*, although the effect is smaller than when App is present (J). (K)

Mutation in App catalytic domain has a similar effect on Dachs and Dlish accumulation as *app*

null alleles (compare J and K). (L) Dachs:GFP localized to cell contacts also recruits App and

Dlish in S2 cells. Scale bars, 5 µm

Figure 9. Regulation of junctional localization and abundance of the core complex by Ds

and Ft.

Membrane localized App is recruited by Ds to form junctional puncta. App promotes localization

of Dlish and Dachs to the junctional cortex by palmitoylation of Dlish. These proteins form a

core complex that is stabilized at the junctional cortex through mutual interaction. In a similar

manner, Ft also recruits the core complex. However activated Ft promotes degradation of the

core complex thereby preventing formation of puncta.

Fig. 1

Fig. 2

A Ds

cadherin repeats TM ICD

N C

Ds^ΔICD:GFP

GFP

B DECad Ds:GFP

B' Ds:GFP

C DECad Ds^ΔICD:GFP

C' Ds^ΔICD:GFP

D *wt* ds^ΔICD:GFP Ft ubi-GFP

D' *wt* ds^ΔICD:GFP Ft

E *wt*

F *ds^ΔICD:GFP*

G *ds^UAO71/ds^05142*

H

relative wing size

**** ns ****

wildtype ds^ΔICD:GFP ds^ΔICD (GFP-)

I *wt*

J *ds^ΔICD:GFP*

K *ds^UAO71*

L *ft*

M *ds^ΔICD:GFP ft*

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

A +Ed:Ft:Flag
Dachs:V5
App:GFP
Dlish:HA

A' Dachs:V5

A'' App:GFP

A''' Dlish:HA

B +app,dlish dsRNA
Ed:Ft:Flag
Dachs:GFP

B' Dachs:GFP

B'' Ed:Ft:Flag

C

FtΔECD:HA		+	+	+	+
app RNAi			+		+
dlish RNAi				+	+
lacZ RNAi		+	+	+	
Dachs:V5	+	+	+	+	+
GFP	+				

IP:HA — V5, HA

input — V5, HA

D

TM — PCP — PH — Hippo N — Hippo C — Su(DN) — C

App
Dlish

ΔECD or ICD — 5147 — HA or Flag

Δ1-C — PQRPD⁴⁷⁰⁶

ΔPH-C — YDLEN⁴⁷⁵¹

Δ6-C — AQQTS⁴⁹²⁰

ΔN-6 — YRFRG⁴⁶¹³ — Q⁴⁹¹⁶AQQT

M — C

E Dlish (HA)
Dachs:GFP
Ed:FtΔPH-C:Flag

E' Dachs:GFP

E'' Dlish:HA

E''' Ed:FtΔPH-C:Flag

F Dlish:HA
Dachs:GFP
Ed:FtΔ6-C:Flag

F' Dachs:GFP

F'' Dlish:HA

F''' Ed:FtΔ6-C:Flag

G Dlish:HA
Dachs:GFP
Ed:FtΔN-6:Flag

G' Dachs:GFP

G'' Dlish:HA

G''' Ed:FtΔN-6:Flag

H

FtΔECD:HA		+			
FtΔECDΔ1-C:HA			+		
FtΔECDΔ6-C:HA				+	
FtΔECDΔN-6:HA					+
Dachs:V5	+	+	+	+	+
GFP	+				

IP:HA — V5, HA

input — V5

Fig. 8

Fig. 9

Off

On

On

Dlish   Dachs

App

Ds

Ft

Off

P P

Fig. S1

Fig. S1. (A) Comparison of Dachs levels between $ds^{UAO71}\ ft^{G-rv}\ dachs^{GC13}/ds^{UAO71}\ ft^{fd}$ + and $ds^{UAO71}\ ft^{fd}$. Dachs's level at junctional cortex in $ds^{UAO71}\ ft^{G-rv}\ dachs^{GC13}/ds^{UAO71}\ ft^{fd}$ + is lower than $ds^{UAO71}\ ft^{fd}$ cells. Dahs staining is absent in $ds^{UAO71}\ ft^{G-rv}\ dachs^{GC13}$ triple mutant clones (white asterisk). Expression and localization of Ds:GFP in $ds^{:GFP}$ homozygote (B) and DsΔICD:GFP in $ds^{ΔICD:GFP}$ homozygote (C) in wing discs. Adherens junctions are marked by anti-DECad staining. Wing discs in $ds^{ΔICD:GFP}$ homozygotes are smaller than $ds^{:GFP}$ homozygotes. (D) App tagging by CRISPR-Cas9 gene editinng. YFP was tagged at C-terminus. 3xV5 was inserted at extra cellular domain between transmembrane domains 3 (TM 3) and 4 (TM 4). (E) C-terminus tagging with YFP for *app* using CRISPR-Cas9. Target sequence (red line) and donor DNA are shown. HAL (blue line), left homology arm; HAR (light green line), right homology arm.
(F) Internal V5 tagging for *app* using CRISPR-Cas9. 3xV5, target sequence (red underline), and PAM site (red letters) are shown. 3x V5 epitope tag was inserted after Histidine at 221. (G) Colocalization of App, Dlish, and Dachs at junctions. App, Dlish, and Dachs are colocalized at junctional puncta (yellow arrows). Scale bars, 10 µm (A, B), 1 µm (G).

Fig. S2

Fig. S2. (A) Ed:Flag did not recruit GFP tagged Dachs and App to cell-cell contacts in S2 cells. Ed:Flag is localized at cell-cell contacts. Dachs:GFP and App are distributed to cytoplasm and cytoplasmic foci. (B) High homology between Ds and Dchs1 (*H. sapiens*) in domain D. Identical (Black) and similar (gray) amino acids are shaded. (C) Constructs for Ed extracellular domain fused to Ds ICD or its deletions for the S2 cell recruitment assay. (D) Ed:DsΔE:Flag recruits Dachs:GFP to cell-cell contacts. (E) Ed:Ds D:Flag recruits Dachs:GFP to cell-cell contacts. Scale bars, 1 μm.

Fig. S3

Fig. S3.
(A-D) Anti-Ds staining in $ds^{\Delta A:Flag}$, $ds^{\Delta B:Flag}$, $ds^{\Delta C:Flag}$ and $ds^{\Delta D:Flag}$ clones. Mutant clones are labeled with the absence of GFP. The level and distribution of Ds did not change in *ds* deletion mutants. (E-G) Wings in $ds^{\Delta A:Flag}/ds^{\Delta B:Flag}$ (E), $ds^{\Delta A:Flag}/ds^{\Delta D:Flag}$ (F) and $ds^{\Delta B:Flag}/ds^{\Delta D:Flag}$ (G).
$ds^{\Delta A:Flag}/ds^{\Delta B:Flag}$ and $ds^{\Delta A:Flag}/ds^{\Delta D:Flag}$ wings are almost normal size (E, F). $ds^{\Delta B:Flag}/ds^{\Delta D:Flag}$ wings display slight undergrowth (G). (H-J) Leg phenotype in $ds^{\Delta A:Flag}/ds^{\Delta B:Flag}$ (H), $ds^{\Delta A:Flag}/ds^{\Delta D:Flag}$ (I) and $ds^{\Delta B:Flag}/ds^{\Delta D:Flag}$ (J). Junctions of tarsal segments are marked with black lines. $ds^{\Delta A:Flag}/ds^{\Delta B:Flag}$ (H) and $ds^{\Delta A:Flag}/ds^{\Delta D:Flag}$ (I) legs display normal segmentation. $ds^{\Delta B:Flag}/ds^{\Delta D:Flag}$ legs (J) are fused at T2 and T3 (yellow arrow). Scale bars, 5 μm (A), 100 μm (E, H).
.

Fig. S4

A

signal sequence | cadherin repeats | EGF | LG | TM

TM

ft 5'target 5   PAM

wt
ATC TAT CGA TTC CGC GGA AAG CAG GAG AAG ATT GGT AGT TTG AGT
I Y R F R G^{4613}K Q E K I G S L S

ft$^{\Delta ICD}$
ATC TAT CGA TTC CGC GGA GG AGA AGA TTG GTA GTT TGA
I Y R F R G R R R L V V Stop

B *ft^{61}*   *wt*
App:YFP
ubi-RFP

B' *wt*
ft^{61}
App:YFP

C *ft^{61}*
wt
Dachs:GFP
ubi-RFP

C' *ft^{61}*
wt
Dachs:GFP

D

FtΔECD:V5		+	+	+		
FtΔECD^{sum}:V5					+	+	+
Dco WT			+			+	
Dco K/R				+			+
GFP:FL	+						

IP:V5
Blot:V5

— phos
— Non-phos

+*app, dlish* dsRNA

E Ed:Ft:Flag / Dachs:GFP
E' Dachs:GFP
E'' Ed:Ft:FL

F Ed:Ft Δ6-C:Flag / Dachs:GFP
F' Dachs:GFP
F'' Ed:Ft Δ6-C:FL

G Ed:Ft ΔN-6:Flag / Dachs:GFP
G' Dachs:GFP
G'' Ed:Ft ΔN-6:FL

Fig. S4. (A) CRISPR-Cas9 induced *ft$^{\Delta ICD}$* mutant. The target sequence (green line), PAM site (green box) and transmembrane domain (blue letters) are shown. The resulting 4 bp deletion led to a premature stop codon after transmembrane domain. The altered amino acids are shown in red letters. (B) App:YFP is increased in junctional puncta in *ft^{61}* mutant cells. (C) Dachs:GFP is increased at the junctional puncta in *ft^{61}* mutant clones. Mutant clones are marked by absence of ubi-RFP. (D) Phosphorylation of wild-type Ft and Ft^{sum} by Dco in S2 cells. Dco WT promotes phosphorylation of Ft. The Ft^{sum} is less phosphorylated by Dco. (E) Ed:Ft:Flag recruits Dachs:GFP to cell-cell contacts in *app* and *dlish* depleted cells. (F) Ed:FtΔ6-C:Flag fails to recruit Dachs:GFP in *app* and *dlish* depleted cells. (G) Ed: FtΔN-6:Flag fails to recruit Dachs:GFP in *app* and *dlish* depleted cells. Scale bar, 5 μm (B, C) 1 μm (E).

Fig. S5

Fig. S5. (A) App:YFP levels in *hh>dlish:Flag* at the junctional cortex. Dlish:Flag was expressed in the posterior compartment. App:YFP is increased with Dlish overexpression. (B) Dlish and Dachs:GFP in *hh>app.* Dlish and Dachs:GFP are enriched at junctional puncta with App overexpression. (C-I) Representative wing discs stained with phalloidin showing growth defects of the following genotypes: wild type (C), *app^12-3* (D), *ds^UAO71 ft^fd; app^12-3/ds^UAO71 ft^G-rv; app^12-3* (E), *dlish^Y003* (F), *ds^UAO71 ft^fd dlish^Y003/ ds^UAO71 ft^G-rv dlish^Y003* (G), *ds^UAO71 ft^fd dlish^Y003; app^12-3/ ds^UAO71 ft^G-rv dlish^Y003; app^12-3*(H) and *ds^UAO71 ft^fd/ ds^UAO71 ft^G-rv*(I). Overgrowth in *ds ft* is suppressed with *app* or *dlish* mutant. (J) Recruitment of cytoplasmic GFP by Ed:Flag:vhh4. GFP is recruited to cell-cell contacts (yellow arrows). (K) Membrane tethering GFP nanobody recruits Dachs:GFP, App and Dlish to cell cortex. (L) Ed:Flag:vhh4 recruits App:GFP, Dachs and Dlish to cell-cell contacts (yellow arrow). (M) Ed:Flag:vhh4 recruits Dlish:GFP, App and Dachs to cell-cell contacts (yellow arrows). (N) Membrane tethering GFP nanobody recruits App:GFP, Dachs:V5 as well as Dlish:HA to cell cortex. (O) myr:Flag:vhh4 recruits Dlish:GFP, App and Dachs:V5 to cell cortex. Scale bars, 5 μm (A, L, M), 100 μm (C), 1 μm (J, K, N, O)