Predicting the diversity of photosynthetic light-harvesting using thermodynamics and machine learning

Callum Gray[1], Samir Chitnavis[1], Tamara Lucy Buja[1] and Christopher D. P. Duffy[1]

1 Digital Environment Research Institute, Queen Mary University of London, Empire House, 67-75 New Road, London E1 1HH

* callum.gray@qmul.ac.uk, c.duffy@qmul.ac.uk

Abstract

Oxygenic photosynthesis is responsible for nearly all biomass production on Earth, and may have been a prerequisite for establishing a complex biosphere rich in multicellular life. Life on Earth has evolved to perform photosynthesis in a wide range of light environments, but with a common basic architecture of a light-harvesting *antenna* system coupled to a photochemical reaction centre. Using a generalized thermodynamic model of light-harvesting, coupled with an evolutionary algorithm, we predict the type of light-harvesting structures that might evolve in light of different intensities and spectral profiles. We reproduce qualitatively the pigment composition, linear absorption profile and structural topology of the antenna systems of multiple types of oxygenic photoautotrophs, suggesting that the same physical principles underlie the development of distinct antenna structures in various light environments. Finally we apply our model to representative light environments that would exist on Earth-like exoplanets, predicting that both oxygenic and anoxygenic photosynthesis could evolve around low mass stars, though the latter would seem to work better around the coolest M-dwarfs. We see this as an interesting first step toward a general evolutionary model of basic biological processes and proof that it is meaningful to hypothesize on the nature of biology beyond Earth.

Introduction

With the discovery of nearly 100 potentially-habitable exoplanets [1], the question of whether they do harbour life and what it might be like warrants serious consideration [2]. In which case, how do we meaningfully *hypothesize* (and not merely speculate) on the forms life might take in environments very different to Earth? Regardless of environment, evolution must be constrained by the laws of physics and chemistry; the question is then whether it is possible to develop a model that predicts the type of biological processes that may evolve under a given set of external conditions. Such a model would have be *necessarily* broad and coarse-grained, presupposing no (or very little) fine molecular detail, but still capturing the essential physics of basic biological functions. In essence the problem becomes one of defining elementary biological processes and determining the basic thermodynamic constraints on their function.

The first assumption we make is that life, life, wherever it evolves, will require an energy source. A common assumption in astrobiology research is that this will be the parent star, with the irradiant flux on the plant surface coupled to the biosphere by some form of photosynthesis [3–7]. In its most basic form, this involves using absorbed light energy to oxidize some chemical substrate to produce reducing agents and chemical energy which are then used in generating biomass. Of course, photosynthesis is certainly not the only mechanism of biomass production, with *chemo*-synthetic organisms oxidizing high-energy compounds from geochemical sources [8, 9]. Indeed, recent work has suggested that the last universal common ancestor (LUCA) on Earth was not photosynthetic [10, 11]. Regardless, in this work we will exclusively consider photosynthesis. Firstly, stellar light is a universal energy and one that will remain relatively constant over the life of the star. Secondly, the light environment on *known* exoplanets is relatively well-characterized, while conditions on the planets' surface (beyond basic parameters such as temperature and planet density) are not. Thirdly, the presence of photosynthesis on an exoplanet may present the best chances of remote detection. While chemo-autotrophs may have been the first forms of life on Earth, our biosphere (at least on the surface) is dominated by photo-autotrophs and these have left two characteristic signatures: an oxygenated atmosphere and planet surface covered in coloured pigment molecules such as chlorophyll. The presence of oxygen in an exoplanet

atmosphere could be detectable as absorption lines in the light passing through its atmosphere as it transits, making it a priority bio-signature [12–14]. On Earth, the global coverage of chlorophyll generates the *vegetation red edge* (VRE) at $\sim$ 700-750 nm in the surface-integrated reflectance spectrum (first observed by the Galileo probe [15]). The potential of measuring a similar reflectance edge is also widely discussed in th context of upcoming missions which will directly image the surface of nearby exoplanets [5,16,17]. We therefore argue not that all biospheres *must* be founded upon some form of photosynthesis, but that any biosphere in general could be, and that photosynthesis represents a promising candidate for the detection of extraterrestrial life if it exists.

The second assumption we make is that the molecular details of photosynthesis will depend on the intensity and spectral quality of incident light. On Earth almost all of the primary production is via *oxygenic* photosynthesis, in which light is used to strip electrons from chlorophyll a to produce energy and chemical reductants [18]. The electrons are replenished by the oxidation of water. The redox chemistry of this last step presents a second (energetic) constraint: photons of wavelength longer than about 750 nm are insufficiently energetic to power water oxidation, a widely discussed phenomenon known as the *red limit* of oxygenic photosynthesis [19,20]. While larger and warmer stars such as our Sun emit strongly in the visible range (400-700 nm), providing a significant amount of usable light for oxygenic photosynthesis, cooler and smaller M-dwarf stars have limited emission in this range and higher emission at longer wavelengths, with their absorption maxima reaching out to around 1 μm. At first glance, then, oxygenic photosynthesis would appear to be unfavourable around such stars. This would appear to be a disappointing conclusion, since M-dwarfs are the most common and longest-lived stars in the universe [21] and have a high occurrence rate of Earth-like planets in their habitable zones [22]. Indeed, well-known, nearby candidate systems such as Trappist-1 [23] and Proxima Centauri [24] orbit such stars. Or course, there is the possibility of *anoxygenic* photosynthesis, which involves photo-oxidation of compounds such as sulphides, hydrogen, iron, etc. and uses much redder (800-900 nm) light [25]. Of course on Earth such organisms are single cell and responsible for less than 1% of the primary production of the biosphere (though this can locally increase to $\sim$ 30% in ancient, sulphide rich lakes [26]). Moreover, it has been proposed that an

oxygenated atmosphere is a necessary prerequisite for the evolution of widespread ₆₄
multicellular life (though there is considerable debate [27–31]), meaning a purely ₆₅
anoxygenic biosphere may not produce the widespread land-based vegetation needed to ₆₆
generate a detectable VRE. Fortunately, recent experiments have shown that various ₆₇
oxygenic cyanobacteria, algae, moss and non-vascular plants are able to grow under ₆₈
simulated M-dwarf light [6,32], though vascular plants fare less well, showing signs of ₆₉
the stress-response known as *shade avoidance syndrome* [6]. These are extremely ₇₀
encouraging results as they prove that oxygenic photosynthesis around M-dwarf stars is ₇₁
not automatically precluded by their paltry emission above 700 nm. That said, this ₇₂
does not mean oxygenic photosynthesis would evolve in such conditions, and our recent ₇₃
theoretical work indicated that oxygenic organisms would require larger and more ₇₄
complicated light-harvesting structures than they do on Earth [7]. ₇₅

The third basic assumption we make is that photosynthetic processes in general will ₇₆
evolve some form of *antenna* structure. Essentially all of the various photoautotrophs ₇₇
on Earth adopt this approach, in which a small number of photochemical *reaction* ₇₈
centres (RC) are supplied energy by a much larger antenna system [33]. While there is ₇₉
significant diversity in the structural details of the various antennae which have evolved ₈₀
in different light environments, they function according to the same principles: A large, ₈₁
modular assembly of pigment-protein complexes absorb incident light and very ₈₂
efficiently transfer the resulting excitation energy to the RCs. That said, they tend not ₈₃
to absorb light indiscriminately across the visible spectrum, but rather selectively ₈₄
absorb specific (and sometimes rather narrow) wavelength bands. Previous theoretical ₈₅
work has suggested that selective light-harvesting structures may have developed to ₈₆
balance capturing light against the metabolic cost of synthesising multiple pigment ₈₇
types [34], to absorb light preferentially at the maximum incident photon flux [4,35], or ₈₈
to mitigate fluctuations in the light level in order to maintain the average power ₈₉
throughput of the antenna [36]. In this work we consider a different potential organising ₉₀
principle: balancing photon flux capture, the thermodynamics of energy transfer, ₉₁
spectral overlap of between antenna components, and the metabolic burden of building ₉₂
and maintaining a large, complex antenna. The first consideration, maximizing photon ₉₃
flux capture, is obvious. We assume a photoautotroph, requiring energy and subject to ₉₄
no other constraints, would evolve an antenna which intercepted the largest amount of ₉₅

incident flux. This would imply a huge antenna, contain multiple pigment types to cover as much of the incident spectrum as possible. The thermodynamic constraint reflects the fact that light-harvesting, intercepting photons over a large area and concentrating the resulting energy into a small RC, involves an unfavorable reduction in entropy ($\Delta S < 0$) [37–39]. To counteract this, light-harvesting generally also involves a reduction in *enthalpy* ($\Delta H < 0$) so that the total change in free energy, $\Delta F = \Delta H - T\Delta S$, is negative (or at least not prohibitively positive). In, for example, Photosystem II (PSII) of vascular plants it is sufficient to have the RC slightly lower in energy than the surrounding (and somewhat iso-energetic) antenna [40, 41]. For cyanobacteria, the antenna (known as the phycobilisome) could be described as 'funnel-like', with higher-energy pigments located further away from the RC. Still, an antenna cannot be arbitrarily large and in our previous work we showed that increasing antenna size eventually leads to diminishing returns in light-harvesting as the entropic barrier becomes harder and harder to overcome [7]. 'Spectral overlap' essentially means that the antenna and RC must be energetically resonant otherwise energy transfer would violate the conservation of energy. This is also true for the different pigment types/domains in the antenna. In practice this means one cannot have an arbitrarily steep funnel [7]. Finally, we assume that having a huge antenna will place some metabolic burden on the organism. This could be in the direct sense of having to divert a fraction of photosynthetic output to build and maintain the antenna system, or in a more abstract sense of taking up valuable space in the cell. In this work we create a general antenna model that captures these constraints. We then use a genetic/evolutionary algorithm to optimize this antenna to different light-environments.

Methods

We construct a thermodynamic model of a general light-harvesting antenna. The model parameters (discussed below) are related to overall structural features such as size, shape and pigment composition. The 'input' for a given antenna is an incident spectral flux of light and the model outputs the overall *electron output rate*, ν_e, and the absolute quantum efficiency, ϕ, of the antenna. We then generate random populations of these antennae and apply an evolutionary algorithm to mimic the processes of selection,

reproduction and mutation in order to produce antennae better optimised to the light environments they are subjected to.

Antenna model

Thermodynamics of light-harvesting

We use a relatively simple kinetic model of photon absorption and energy transfer within a given light-harvesting structure based on that previously reported in [7], with a diagram of an example antenna shown in Fig. 1A.

The model contains one generalized RC containing N_r pigments plus exactly one trap state $N_t = 1$ which converts an exciton produced by absorption of a photon into a charge-separated state. The rest of the antenna is composed of n_b identical branches, each made up of n_s light-harvesting complexes (*LHCs*) which we refer to as *subunits*; these in turn contain n_i identical pigments chosen from the set shown in Fig. 1B and described in further detail in Table 1 of the Supplementary Material.

We consider five main processes occurring in the antenna: photon absorption, energy dissipation in the antenna, energy transfer, energy trapping by the RC, and electron output. We do not consider dissipative processes within the RC (such as non-radiative charge recombination), since we are primarily concerned here with the effect of antenna evolution. We do not present a rigorous treatment of charge-separation and transfer in the RC at all and so we assume that dissipative processes simply renormalize (and are already accounted for in) the rae constant for electron output. We make the assumption that an RC would evolve to minimize such waste channels as they ultimately defeat the purpose of the RC. A more detailed model of the combined antenna-RC supersystem will be the focus of our next work.

Photon absorption is described by the excitation rate γ_i as

$$\gamma_i = \int_0^\infty \frac{\lambda}{hc} f_p(\lambda) A_i(\lambda) \, d\lambda \tag{1}$$

where $f_p(\lambda)$ is the incident spectrum (which can be one representing an Earth-like light environment or an extrasolar spectrum, with examples shown in Fig. 1B) and $A_i(\lambda)$ is the absorption spectrum of the pigments in subunit i. λ/hc ensures that the resulting

Fig 1. A.) Thermodynamic model of antenna. Rounded squares represent antenna subunits with the small circles representing pigments. Colour (green, blue) crudely represents absorption wavelength. Light is absorbed by each sub-unit and energy is transferred between them (black arrows), ending at the reaction centre (red) which traps (dark red) the energy and outputs electrons. B.) Altering the light environment from full sunlight (yellow) to deep water (blue) or shaded (red) via filters. C.) Genetic algorithm. Pairs of parents are selected (dashed lines) from the initial population to reproduce (left to middle), after which a mutation step is applied to members of the population at random (middle to right) to form the next generation. D.) Fitted absorption lineshapes of the available pigments for the genetic algorithm to choose from [33, 42, 43]. E.) Example of evolution of the average absorption spectrum from initial population (black) to a converged population (red) under red light illumination.

quantity has dimensions of photons m^{-2} s^{-1}.

Energy dissipation in the antenna is assumed to be due to realtively fast processes such as non-radiative excitation decay and fluorescence, which we combine these into one rate k_{diss}.

Considering the kinetics of energy trapping in and electron output from the reaction centre requires us to describe explicitly whether the trap is already closed (occupied) or open (unoccupied), since this affects which processes are possible. The relevant equations are

$$\left[\frac{d}{dt}P_{n_r,0}(t)\right]_{\text{trap}} = -n_r k_{\text{trap}} P_{n_r,0}(t) \tag{2}$$

$$\left[\frac{d}{dt}P_{n_r,1}(t)\right]_{\text{trap}} = -k_{\text{out}} P_{n_r,1}(t) + (n_r+1) k_{\text{trap}} P_{n_r+1,0}(t) \tag{3}$$

for the open and closed traps respectively. Not that here we define *trapping* as the process of an exciton localized on the RC undergoing charge separation. In the photosynthetic literature 'trapping' can also include the migration of the exciton to the RC, which here is accounted for in our energy transfer terms. In real photosynthetic systems the kinetics of trapping involves multiple steps involving progressively longer timescales and larger distances. The initial charge separation steps are fast (order of 10-100 ps), while later steps involving reduction of electron carriers can be much slower (μ-ms). In our model we class the very earliest charge separation steps as trapping, since it is at this point the energy transfer gives way to electron transfer, with the latter not being rigorously treated in our model. We therefore define $k_{\text{trap}}^{-1} \sim 5$ ps for two reasons. Firstly, it is the approximate timescale of the formation of the radical pair states in PSII [41,44] and secondly, if trapping is much slower then the back transfer of excitation into the antenna becomes the favourable process and the entire system ceases to work effectively.

The timescale for acceptor reduction, k_{out}^{-1} is *much* slower than the other timescales in the model. We will choose $k_{\text{out}}^{-1} \sim 10$ ms as this is approximately the timescale of the H_2O oxidation-quinone reduction cycle of PSII [45]). As previously stated, sever over-simplification of a complex system. There is even some evidence that

the instantaneous state of the RC may also modulate exciton migration in the
antenna [46]. However, as previously stated, this is left for our follow-up work.

Finally we consider energy transfer rates between subunits, beginning with a
fundamental hopping rate k_{hop} which sets an overall timescale. This will depend to
some extent on details of the antenna which we deliberately do not model, but we can
say in general that $k_{\mathrm{hop}} >> k_{\mathrm{diss}}$ in order for light-harvesting to work effectively; we use
a fundamental rate $k_{\mathrm{hop}} = 10\mathrm{ps}$ in this work, which is the typical timescale of energy
transfer between subunits in real photosystems [40]. This base hopping rate is then
modified by the overlap of the spectral lineshapes of the initial and final subunits and
the free energy change using,

$$K_{n_i,n_j}^{n_i-1,n_j+1} = k_{\mathrm{hop}} n_i A_{ij}\, \rho\left(E_i - E_j\right) f\left(\Delta F_{n_i,n_j}^{n_i-1,n_j+1}\right) \tag{4}$$

for hopping from subunit $i \to j$ with initial subunit populations as subscript and final
as superscript. A_{ij} is simply an adjacency term equal to 1 if the two subunits are
connected and 0 otherwise. $\rho(E_i - E_j)$ is the *density of states* which enforces energy
conservation on the process; energy transfer is only possible between subunits if the
pigments in the subunits are at resonance, and the degree of resonance is approximately
captured by the overlap integral,

$$\rho\left(E_i - E_j\right) \sim \int_0^\infty d\lambda \tilde{F}_i\left(\lambda; \lambda_i^p, w_i\right) \tilde{A}_j\left(\lambda; \lambda_j^p, w_j\right) \tag{5}$$

where $\tilde{F}_i$ and $\tilde{A}_j$ are the normalized fluoresence and absorption lineshapes of the donor
i and acceptor j respectively. Finally $f\left(\Delta F_{n_i,n_j}^{n_i-1,n_j+1}\right)$ encodes the thermodynamics of
energy transfer via the free energy change. At equilibrium the rates of forward and
backward energy transfer between any pair of subunits must satisfy the *detailed balance*
condition:

$$\frac{K_{n_i,n_j}^{n_i-1,n_j+1}}{K_{n_i-1,n_j+1}^{n_i,n_j}} = \left(\frac{n_i}{n_j+1}\right) \exp\left(-\frac{\Delta F_{n_i,n_j}^{n_i-1,n_j+1}}{k_B T}\right) \tag{6}$$

$\Delta F_{n_i,n_j}^{n_i-1,n_j+1}$ is the Helmholtz free energy change associated with the transfer process,

$$\Delta F_{n_i,n_j}^{n_i-1,n_j+1} = \Delta H_{n_i,n_j}^{n_i-1,n_j+1} - T\Delta S_{n_i,n_j}^{n_i-1,n_j+1} \tag{7}$$

The first term here ΔH represents the enthalpy change, i.e. the difference in internal energy of the system before and after the transfer process,

$$
\begin{aligned}
\Delta H_{n_i,n_j}^{n_i-1,n_j+1} &= (n_i - 1)\, E_i + (n_j + 1)\, E_j - n_i E_i - n_j E_j \\
&= E_j - E_i \\
&= hc \left(\frac{1}{\lambda_j^p} - \frac{1}{\lambda_i^p} \right)
\end{aligned}
\tag{8}
$$

where $E_{i,j}$ are the excitation energy of the pigments in sub-unit i, j, which is inversely proportional to $\lambda_{i,j}^p$, the peak absorption wavelengths.

The second term ΔS is the entropy change of the process. If we have two coupled LHCs with N_i and N_j identical pigments respectively, of which $n_i \leq N_i$ and $n_j \leq N_j$ are excited, the combined entropy of the two is

$$
S_{n_i,n_j} = k_B \, \ln \left[W(n_i, N_i) W(n_j, N_j) \right]
\tag{9}
$$

where $W(n_i, N_i)$ is the multiplicity,

$$
W(n_i, N_i) = \frac{N_i!}{n_i! \, (N_i - n_i)!}
\tag{10}
$$

If one excitation is transferred from N_i to N_j then the entropy change is,

$$
\Delta S_{n_i,n_j}^{n_i-1,n_j+1} = k_B \, \ln \left(\frac{W(n_i - 1, N_i) W(n_j + 1, N_j)}{W(n_i, N_i) W(n_j, N_j)} \right)
\tag{11}
$$

Implicit in this is the assumption that excitons equilibrate *within* each sub-unit much faster than the typical timescale of energy transfer *between* them, so that they can sample all the available states (ergodicity). We have previously shown that this is true up to around $N_p = 100$ [7], considering the typical equilibration times and hopping rates of real photosystems, and hence set a maximum subunit size of 100 pigments throughout this work; in practice, this limit is never reached.

ΔF is antisymmetric with respect to reversal of the the transfer process,

$$
\Delta F_{n_i,n_j}^{n_i-1,n_j+1} = -\Delta F_{n_i-1,n_j+1}^{n_i,n_j}
\tag{12}
$$

and hence, for non-identical subunits, there will be a thermodynamically favoured

direction of energy transfer. This is encoded into the function

$$
f\left(\Delta F_{n_i,n_j}^{n_i-1,n_j+1}\right) =
\begin{cases}
1 \text{ for } \Delta F_{n_i,n_j}^{n_i-1,n_j+1} \leq 0 \\
\\
\exp\left(-\beta \Delta F_{n_i,n_j}^{n_i-1,n_j+1}\right) \text{ for } \Delta F_{n_i,n_j}^{n_i-1,n_j+1} > 0
\end{cases}
\tag{13}
$$

where $\beta = 1/k_B T$ is the inverse thermodynamic temperature. If the forward transfer

process, $K_{n_i,n_j}^{n_i-1,n_j+1}$ represents a decrease in free energy ($\Delta F < 0$), then the backward

rate is unfavourable ($\Delta F > 0$) and subject to a multiplicative *Boltzmann penalty*, and

vice-versa.

Finally, to construct the set of time-dependent rates which we will solve for, we

assume the *single excitation regime*:

$$
n_i = 0, 1 \tag{14}
$$

$$
n_r = 0, 1 \tag{15}
$$

$$
\langle n_r \rangle + \sum_i \langle n_i \rangle \leq 1 \tag{16}
$$

In words, the entire antenna-plus-RC (hereafter 'photosystem') can contain a

maximum of one excitation at any time, though the trap can be either open or closed.

This is reasonable because k_{diss} and k_{trap} are much faster than γ_i, even in bright light.

This assumption allows us to neglect multi-excitation terms in our model and if we

adopt notation,

$$
P_{0,0,n_t} = P\left(n_1 = 0, n_2 = 0, \ldots, n_i = 0, n_r = 0, n_t\right) \tag{17}
$$

$$
P_{1_i,0,n_t} = P\left(n_1 = 0, n_2 = 0, \ldots, n_i = 1, n_r = 0, n_t\right) \tag{18}
$$

$$
P_{0,1,n_t} = P\left(n_1 = 0, n_2 = 0, \ldots, n_i = 1, n_r = 1, n_t\right) \tag{19}
$$

and so on, then the equations of motion reduce to,

$$\frac{d}{dt}P_{0,0,0}(t) = -\sum_i \gamma_i N_i(1-n_i)P_{0,0,0}(t)$$
$$+ k_{\text{diss}}\left(P_{0,1,0}(t) + \sum_i P_{1_i,0,0}(t)\right) + k_{\text{out}}P_{0,0,1}(t)$$

(20)

$$\frac{d}{dt}P_{0,0,1}(t) = -\left(\sum_i \gamma_i N_i(1-n_i) + k_{\text{out}}\right)P_{0,0,1}(t)$$
$$+ k_{\text{diss}}\left(P_{0,1,1}(t) + \sum_i P_{1_i,0,1}(t)\right) + k_{\text{trap}}P_{0,1,0}(t)$$

(21)

$$\frac{d}{dt}P_{1_i,0,0}(t) = -\left(k_{\text{diss}} + k_{i\to r} + \sum_j k_{i\to j}\right)P_{1_i,0,0}(t)$$
$$+ \gamma_i N_i(1-n_i)P_{0,0,0}(t) + k_{r\to i}P_{0,1,0}(t) + \sum_j k_{j\to i}P_{1_j,0,0}(t)$$
$$+ k_{\text{out}}P_{1_i,0,1}(t)$$

(22)

$$\frac{d}{dt}P_{1_i,0,1}(t) = -\left(k_{\text{diss}} + k_{\text{out}} + k_{i\to r} + \sum_j k_{i\to j}\right)P_{1_i,0,1}(t)$$
$$+ \gamma_i N_i(1-n_i)P_{0,0,1}(t) + k_{r\to i}P_{0,1,1}(t) + \sum_j k_{j\to i}P_{1_j,0,1}(t)$$
$$+ k_{\text{trap}}P_{1_i,1,0}(t)$$

(23)

$$\frac{d}{dt}P_{0,1,0}(t) = -\left(k_{\text{diss}} + k_{\text{trap}} + \sum_i k_{r\to i}\right)P_{0,1,0}(t)$$
$$+ \sum_i k_{i\to r}P_{1_i,0,0}(t) + k_{\text{out}}P_{0,1,1}(t)$$

(24)

$$\frac{d}{dt}P_{0,1,1}(t) = -\left(k_{\text{diss}} + k_{\text{out}} + \sum_i k_{r\to i}\right) P_{0,1,1}(t)$$

$$+ \sum_i k_{i\to r} P_{1_i,0,1}(t)$$

(25)

where the rate constants are defined as

$$k_{i\to j} \equiv K_{1_i,0_j}^{0_i,0_j}.$$

(26)

The thermodynamic quantities also simplify:

$$\Delta H_{1_i,0_j}^{0_i,1_j} \equiv \Delta H_{i\to j} = hc\left(\frac{1}{\lambda_j^p} - \frac{1}{\lambda_i^p}\right)$$

(27)

$$\Delta S_{1_i,0_j}^{0_i,1_j} \equiv \Delta S_{i\to j} = k_B \ln\left(\frac{N_j}{N_i}\right)$$

(28)

Now that all the relevant parameters and rates have been defined we can numerically

solve the equations of motion in the steady state:

$$\frac{d}{dt}P_{n_r,n_t} = 0$$

(29)

subject to the constraints

$$\sum_{n_i,n_r,n_t} P_{n_r,n_t}(t) = 1$$

(30)

$$0 \leq P_{n_r,n_t}(t) \leq 1$$

(31)

to obtain the full set of equilibrium probabilities $\mathcal{P}_{n_r,n_t}$. From these we can obtain the

average occupancies

$$\langle n_t \rangle = \sum_{n_i, n_r} \mathcal{P}_{n_r, 1} \tag{32}$$

$$\langle n_r \rangle = \sum_{n_i, n_r} n_r \left(\mathcal{P}_{n_r, 0} + \mathcal{P}_{n_r, 1} \right) \tag{33}$$

$$\langle n_i \rangle = \sum_{n_i, n_r} n_i \left(\mathcal{P}_{n_r, 0} + \mathcal{P}_{n_r, 1} \right) \tag{34}$$

which we then use to calculate two observable quantities. The first is the *electron output rate*,

$$\nu_e = k_{\text{out}} \langle n_t \rangle \tag{35}$$

which can be measured (indirectly) for oxygenic photo-autotrophs using an oxygen electrode. The second is the *absolute antenna quantum efficiency*,

$$\phi_e = \lim_{\gamma_i \to 0} \left[\frac{\nu_e}{\nu_e + k_{\text{diss}} \left(\langle n_r \rangle + \sum_i \langle n_i \rangle \right)} \right] \tag{36}$$

which is the fraction of captured photons that go on to generate an electron in the limit of low illumination. We note that ϕ_e is similar but not directly comparable to Φ_{PSII}, the PSII quantum efficiency, which is routinely measured using specialized fluorimeters in order to quantify the performance of crop species [47, 48], and essentially measures the efficiency of excitation trapping. We use ν_e and ϕ_e to quantify the performance of our generalized photosystems for a given light input.

Constraints on overall antenna structure

The possible ways in which a collection of different pigments could be arranged into an antenna represents a intractably vast parameter space. We therefore impose some reasonable constraints on the overall geometry. We assume that the RC lies at the centre of our system with some arbitrary number, n_b, of antenna branches radiating outward (see. Fig. 1 **A**). A branch can contain an arbitrary number, n_s, of subunits which each contain an arbitrary number, n_p^i of pigments. We assume that (1) all branches are identical and (2) the pigment states in a subunit are all isoenergetic. The

first constraint is applied purely to ensure computational tractability and is partially justified by the modular/symmetric nature of many real antenna systems. However, highly asymmetric antennae do exist and this should be considered in future work. The second constraint is a reflection of the assumption of a separation of timescales. Energy equilibrates rapidly (we assume essentially instantaneously) over clusters of closely-packed, chemically-similar pigments, while energy transfer between such clusters is assumed to be slower (~ 10 ps). The subunits of our model are not meant to necessarily reflect entire protein complexes, merely clusters of strongly coupled pigments. Similarly, the number of pigments in a subunit does not necessarily equate to the number of actual molecules, but simply the number of *thermodynamically equivalent states*. If one has a cluster of N_p^i pigments with different energies, E_i, then the number of *equivalent* pigment states is

$$n_p^i \sim \left\lceil \sum_{i=1}^{N_p^i} \exp\left(-\frac{E_i}{k_B T}\right) \right\rceil \tag{37}$$

Finally, we set some upper limits on parameters based on computational considerations of memory and simulation time. As mentioned above, we assume $n_p^i \leq 100$ which is justified by our previous work [7]. We also set $n_s \leq 100$ and $n_b \leq 12$, with the latter partially justified by close-packing considerations (you can pack 12 spheres around a central sphere of the same size). However, in practice, none of these limits are ever reached in the evolutionary optimization. Beyond this we impose no other structure on our antenna system, allowing the genetic optimization to choose effective structures. There is nothing to stop the model selecting an entirely isoenergetic antenna or one with entirely random energy landscape (beyond the fact that this is unlikely to work very well).

One should not over-interpret the various antenna structures produced by out model. They are better thought of in terms of pigment/spectral composition and topology rather than strict real-space models.

Excitation transfer between antenna branches

It should be possible in general for energy to be transferred between adjacent branches, and we initially allowed for this possibility in our simulations, but found that it has no

effect whatsoever on the resulting kinetics; this is because the branches are assumed to be identical, and hence at equilibrium the rates of transfer between them are identically equal as well. In a radially-symmetric system such as ours, the radial and lateral hopping of the excitation are independent and therefore antenna efficiency *cannot* be improved (or decreased) by allowing hopping between branches. After verifying this we disabled energy transfer between branches, and do not consider it in any of the work published here. Note that any consideration of non-identical branches or non-equilibrium interactions within the complex would necessarily require radial transfer between branches to be properly accounted for.

Different pigment types available in the model

We use a range of pigment lineshapes which are taken from various organisms in different photosynthetic niches. For each subunit of the antenna the algorithm is free to choose any of the available pigment types when constructing or mutating the antennae. This choice is implemented by taking published absorption and emission lineshapes for each pigment type in solution [33, 42, 43] and fitting them to a set of Gaussian functions. The available choices are shown in Fig. 1D; the individual fits as well as the corresponding fits to fluorescence lineshapes are given in Supplementary Material in Tables S1 and S2 and Figs. S1–S3. These fitted Gaussians are then recalculated for the set of wavelengths given in the input spectra to avoid any interpolation issues and also to allow for future work where it may be necessary to introduce shifting of absorption peaks.

We do not consider excitonic mixing or inhomogenous broadening of these lineshapes since we do not model the microscopic structure of the antenna. Of course, looking at the effects of these mechanisms would be a worthy extension of this work.

The Genetic Algorithm

The genetic algorithm used in this work uses the standard selection, crossover and mutation steps [49–51], as schematically represented in Fig 1**C**. The antenna parameters n_b, n_s, and the full set of n_p^i and pigment types constitute the *genome* of our generalized antenna. A *population* of such genomes is initialized and their *fitness* is

evaluated in terms of the overall electron output, ν_e, that the antenna can generate in the pre-specified light environment. The relative fitness of the different members of the population are then used to determine the inherited properties of the next generation, with *crossover* capturing the gene-mixing process of sexual reproduction. Throughout the evolution we track various quantities averaged over the population, including the average absorption spectrum of the antennae, as shown in Fig. 1E.

Since we allow the size of the antenna and its pigment composition to vary, we then have genomes of arbitrary length which are themselves composed of a mixture of numerical and categorical variables; the implementation of the crossover and mutation algorithms in particular then require some care. We detail the procedures for each of the operators here.

Initialisation of the population

We allow for two different methods of creating an initial population of antennae. The first, which is used throughout most of this work, represents *in situ* evolution of an organism in a given environment. To represent this we consider an initial population of *proto-antennae* all composed of a single branch with a single subunit containing a random number of a random pigment: $n_b = 1, \; n_s = 1, \; n_p \in [1, 100]$.

Alternatively we can model *contingent* evolution as discussed below in the context of antenna evolution in underwater environments. In this case we provide a template for the genetic algorithm to begin from together with a variability parameter. The starting template can be the best/average outcome of a previous simulation or some biologically rationalized model. The variability parameter takes values between 0 and 1 and denotes the fraction of the population which should be mutated using the mutation algorithm described below before the simulation begins. Setting the variability to 0 then provides an initial population of identical antennae.

Selection based on fitness

The fitness function for which the algorithm optimises is given by

$$F = \nu_e - \chi \left(n_b \sum_{i=1}^{n_s} n_p^i \right) \tag{38}$$

where χ is a pre-defined cost parameter. χ represents an abstract metabolic burden associated with building and maintaining a large antenna system, a burden that increases with the size of the antenna. For example, $\chi = 0.02$ would mean that an antenna containing 100 pigments would cost 2 electrons per second, deducted from the total of ν_e produced by the RC. Obviously a large number of processes could contribute to χ, such as the metabolic cost of synthesising proteins and pigments, the cost of supporting non-photosynthetic tissues, the sacrifice of physical space within the cell, etc. It is not intended to be a quantitative measure, but qualitatively reflects the fact that building the photosynthetic machinery would come at some expense. It most likely differs for different types of organism and we would expect, say, χ to be higher for vascular plants than for prokaryotes. Practically, χ prevents the genetic algorithm from selecting huge but very inefficient antenna, as shown in Fig. 2A. Consider a situation where one doubles the size of the antenna for an increase in ν_e of 0.001%. If $\chi = 0$ then this technically reflects an increase in fitness, albeit an extremely marginal one.

Selection on the basis of fitness is performed using stochastic universal sampling via the cumulative probability

$$p_m = \frac{1}{\sum_m F_m} \left(1 - \exp\left(-\frac{F_m}{F_{\max}} \right) \right) \tag{39}$$

where F_m denotes the fitness of the m'th antenna in the population and $F_{\max} = 100 \text{ s}^{-1}$ is the maximum possible fitness. In this way pairs of parents are chosen to reproduce; we use a generational model, where the entire population is replaced after each generation. Note that this represents a high selection pressure, and in particular prevents any antenna whose fitness $F < 0$ from reproducing at all, which we identify with individuals whose antennae are so badly optimised for the light environment that they cannot capture enough energy to survive.

Crossover

The multivariate nature and differing lengths of the genomes in the population make standard procedures such as one- or two-point crossover unfeasible in our case; instead, we perform *intermediate recombination* [49] on our pairs of parent antennae when constructing new child genomes. This is done using a recombination parameter

$d = 0.25$ [52] and the function

$$x_{\text{child}} = x_{p_1} b + x_{p_2}(1 - b) \tag{40}$$

where $x \in n_b, n_s, n_p^i$ and the subscripts p_1 and p_2 indicating parents 1 and 2 respectively. $b = \text{random}(-d, 1 + d)$ is a random number weighting the contribution of parent 1 versus parent 2, and the subsequent value is rounded to integer for integer parameters.

The pigment type associated with a particular antenna subunit is a categorical variable; in this case, for each subunit, the algorithm chooses randomly from the pigment types of the two parents at that subunit. If the number of subunits of the child exceeds that of both its parents, the pigment type of the final subunit is used instead.

Mutation

Again, due to the non-typical nature of the genomes used, it is necessary to define a bespoke mutation operator for our antennae. In order to reduce the number of introduced hyperparameters and avoid imposing any assumptions about which mutations should be more or less common, we choose to use a single mutation rate $r = 0.05$; when a particular antenna is chosen to mutate, we mutate several of its parameters one by one by drawing new values from a Gaussian distribution. In this way we allow all the relevant characteristics of our antennae to mutate without creating a hierarchy of mutations which presuppose a particular structure or set of biosynthetic pathways. The Gaussian distribution from which we draw new values for a given parameter is centred at the current value of the parameter μ_{mutation} and has a width of $\sigma_{\text{mutation}} = 0.1\mu_{\text{mutation}}$; it is truncated where necessary to avoid unphysical values being accepted, for example by setting the number of branches or blocks to zero. The exception to this procedure is again the pigment type of a given subunit; in this case, mutation draws a new pigment type as a random choice from the set of available pigment types with no bias towards those already present in the antenna.

Fig 2. A.) Pictorial representation of light-harvesting structures evolved for different values of the cost χ. Different pigment types represented as different colours; the colours themselves are arbitrary. At very low cost, the antennae become extremely large and contain pigments which are useless for light-harvesting; at too high a cost they fail to evolve at all. B.) Average electron output per second ν_e and antenna efficiency ϕ as a function of χ for antennae evolved under standard AM1.5 spectrum [53] C.) Average absorption spectrum of total population as a function of χ, showing a longer tail and more featured spectrum as the cost decreases. The y-axis here and throughout denotes the intensity of the incident light (grey dashed line) rather than the absorption curves, which are simply normalised by setting their maximum to 1 for readability.

Results

Oxygenic photosynthesis in full sunlight

Fig. 2 shows the results of simulating antenna evolution using the standard AM1.5 sunlight spectrum [53] for a range of values of the cost parameter χ. This corresponds to very bright illumination, equivalent to a photosynthetically active (400-700 nm) photon flux of over 1000 μmol m^2 s^{-1}; note that we neglect the effect of any photoprotective processes such as non-photochemical quenching (NPQ); the precise molecular mechanisms of this process are still under debate [54] and a proper treatment is far beyond the scope of this work. The resulting averaged electron output ν_e and antenna efficiency ϕ in Fig. 2B show very high electron output, up to $80s^{-1}$ at lower

cost values where the antennae are larger, and high efficiencies of 70–90%. The averaged antenna absorption spectra shown in Fig. 2C have absorption peaks at around 650nm, 610nm and 570nm, corresponding to the cyanobacterial pigments allophycocyanin (APC), phycocyanin (PC) and phycoerythrin (PE) respectively, with differing proportions depending on cost. More detail on pigment composition is available in Table 1 and in the Supplementary Material Fig. S4. We note that for the lowest costs we also start to see small peaks appearing in the average absorption spectra at around 700 and even 800 nm, corresponding to the inclusion of pigments such as Chl d and BChl a, along with a decrease in the efficiency of the antennae. These pigments do not match the peaks in incident flux and transfer very little energy to the reaction centre, reflecting the fact that as the cost decreases it is less disadvantageous to build a larger, less efficient antenna. Below $\chi = 0.005$ the population becomes even larger and less efficient with more and more unusual features appearing in the absorption spectra, whereas above $\chi = 0.04$ it can become difficult to evolve any population at all depending on light environment. Hence for the rest of this work we apply bounds $\chi \in [0.005, 0.04]$ and simulate several values within this range.

Oxygenic photosynthesis in terrestrial photosynthetic niches

In Fig. 3 we consider a selection of different terrestrial light environments calculated by filtering standard AM1.5 light from the previous section in different ways: redder light such as that found in the shade (top row) and bluer light as found under varying depths of clean water (middle and bottom rows). Further information about antenna composition and performance is given in Table 1 and Supplementary Material Fig. S5-S8. Each column in Fig. 3 corresponds to a different overall intensity of light. The medium and low light simulations were performed by taking the high intensity filtered light and renormalising the integrated flux to 300 and 50 μmol m^{-2} s^{-1} of photosynthetically active radiation (400 - 700 nm, PAR) respectively. In each case our algorithm predicts structures using pigments commonly found in that environment as well as qualitatively reproducing important features of experimental absorption spectra.

We first simulate red light using a simple filter which blocks light of wavelengths shorter than roughly 600 nm [58], finding a population of antennae primarily composed

Fig 3. Averaged absorption spectra as a function of cost χ for varying light environments (red light, top row; shallower water, middle row; deeper water, bottom row) and light levels (high intensity, left column; medium intensity, middle column; low intensity, right column). The y-axis corresponds to the intensity of the incident spectra shown in grey. Dashed lines are experimental absorption spectra for, respectively, PSII-LHCII in spinach [55] (top row), cyanobacterium *Synecocystis* sp. PCC 6803 [56] (middle row) and marine red alga *Lithothamnion glaciale* [57](bottom row).

of chlorophyll a and b as well as small amounts of allophycocyanin (APC), a cyanobacterial phycobiliprotein whose absorption profile is very similar to Chl b, especially for lower cost values. The resulting absorption spectra are shown in the top row of Fig. 3 along with an experimentally measured absorption spectrum of the combined PSII-LHCII spectrum from spinach [55], showing good agreement in the Q_y region of the spectrum.

We next simulate shallow (≈ 1 m depth) and deeper (≈ 2.5 m) water spectra using the formula for light attenuation by water given in [59]. Note that this is a crude approximation for a true aquatic spectral flux since real terrestrial water also contains varying concentrations of dissolved and particulate organic matter (known as gilvin and trypton in the literature), as well as phytoplankton, which absorb differently to the water itself. However, since the relative concentrations of these are heavily location-dependent on Earth, we choose to neglect them at this stage, meaning that the resulting spectral fluxes are generally bluer and more intense than would be expected at equivalent depth in real marine environments.

For the shallower water spectrum (middle row of Fig. 3) we generally obtain antennae composed of APC, phycocyanin (PC) and different forms of phycoerythrin (PE), with small amounts of Chl b at higher cost values. Varying the cost varies the proportions of these pigments, leading to shifts in the largest absorption peak from 650nm (APC) to 600nm (PC) to 550nm (PE). Additionally, reducing the light level also blueshifts the largest absorption peak by enriching the antennae with first PC and then PE. We note that specifically at low light and at cost $\chi = 0.01$ our algorithm cannot find any viable solutions: this seems to be due to the specific location of the peak in flux, and suggests that as the light intensity decreases and the cost increases the algorithm becomes increasingly sensitive to fine features in the input spectral flux.

In deeper water (bottom row of Fig 3 we see antennae enriched even further with PE, especially at higher cost values and lower light intensities, as our antennae evolve to capture as much flux in the blue region of the spectrum as possible. We note that some cyanobacteria also acclimate to bluer light by reducing their PC content in favour of PE via a process called type III complementary chromatic adaptation (CCA) [60, 61], although they do not stop producing PC entirely. The extreme adaptations of the simulated antennae are possible since they are free to completely overhaul their

Light	Light level	n_b	n_s	Pigment composition
Full sunlight	High	3-4	3-5	APC, PC, Chl b
	Medium	2-6	2-6	APC, PC, Chl b
	Low	2-6	1-6	APC, PC, Chl a
Red light	High	2-4	3-5	Chl a, Chl b, APC
	Medium	2-4	2-6	Chl a, Chl b, APC
	Low	3-5	2-6	Chl a, Chl b
Shallow water	High	2-4	3-5	APC, PC, r-PE, Chl b
	Medium	2-6	2-7	PC, PE, r-PE, APC
	Low	1-5	1-6	PC, r-PE, APC
Deep water	High	2-4	3-6	PC, r-PE, PE, APC
	Medium	2-6	2-7	r-PE, PC, PE, APC
	Low	1-7	1-7	r-PE, PC, PE, APC

Table 1. Details of antenna composition for different light environments and intensities, including branch number, n_b, and length, n_s, and approximate pigment composition.

structure as necessary, and are therefore much more plastic (i.e. they can respond far   471
more easily to the illumination they receive) than any real-world organism. Nonetheless,   472
we do find absorption spectra with similar absorption maxima to real cyanobacteria in   473
shallow water, and reproduce the blue absorption peak at $\approx$570 nm of certain species of   474
red alga such as *Lithothamnion glaciale* which thrive in deep water with PE-enriched   475
antennae [57].   476

Finally, we note that the average number of branches chosen by the algorithm does   477
not seem very sensitive to light input, only to the cost parameter χ. Naively, we might   478
expect a larger number of very short branches to be favoured in general, especially   479
where there is high flux at wavelengths close to that of the RC, but do not find this to   480
be the case.   481

Evolution of far-red adapted antennae   482

Certain species of cyanobacteria were recently shown to be able to acclimate to light   483
environments which are extremely depleted in PAR but rich in Far Red (700 - 750 nm)   484
light (often due to shading by other organisms) [58, 63, 64]. This response is known as   485
FaRLiP (Far Red Light Photoacclimation) and involves the antenna and RC becoming   486
enriched in chlorophyll d and/or f. Moreover, there is also evidence that the RC trap of   487
PSII redshifts from its usual 680 nm to $\sim$ 720 nm, again via subsitution of different   488
chlorophylls. We simulated a progressively Far Red this by applying a sharper and   489
sharper filter to reduce the amount of available visible light at wavelengths shorter 700   490

Fig 4. Averaged absorption spectra as a function of cost compared to the far-red adapted cyanobacteria *Acarychloris marina* [62] for (A) 40% of available visible light below 700nm, (B) 20%, and (C) 5%. A clear shifting of the main absorption peak as the intensity of visible light decreases is seen, corresponding to a shift from Chl *a* and *b* to Chl *d* in the antennae. (D) Electron output as a function of cost and percentage of available light under 700nm. Vertical dashed lines indicate approximately the fraction of shorter-wavelength light at which the antennae switch to using predominantly Chl *d* rather than Chls *a* or *b*, showing a significant decrease in electron output around this point.

nm. For a non-FarLiP RC with absorption at 680 nm, our algorithm has difficulty $\quad$ 491

finding any solutions with positive fitness values, and hence a functional antenna does $\quad$ 492

not evolve at all. This reflects the fact that absorbing light at > 700 nm and $\quad$ 493

transferring energy uphill to a 680 nm RC is profoundly inefficient. $\quad$ 494

However, with a prototypical FarLiP RC absorbing at 720 nm, we find that our $\quad$ 495

antennae evolve successfully except at the very lowest PAR intensities and highest cost $\quad$ 496

vales; we also see a transition from antennae predominantly composed of Chl a and b in $\quad$ 497

Figs. 4A and B to Chl d in Fig. 4C as the amount of PAR decreases, matching the red $\quad$ 498

absorption peak of the far-red adapted cyanobacteria *Acaryochloris marina* [62] at $\quad$ 499

around 700 nm. At around 5% of the PAR intensity, for the highest simulated cost $\quad$ 500

value, we see a change again from a Chl d antenna to a very small Chl f antenna; $\quad$ 501

increasing the cost or decreasing the intensity further prevents the algorithm from being $\quad$ 502

able to find any solutions at all. We also see in Fig. 4D that there is a considerable $\quad$ 503

decline in electron output as the main constituent of the antenna changes from Chl a $\quad$ 504

and b to Chl d to Chl f, without a significant gain in efficiency; further information and $\quad$ 505

details of pigment compositions can be found in Supplementary Material Fig. S9. This $\quad$ 506

is consistent with the fact that even FaRLiP cyanobacteria have restricted growth in Far $\quad$ 507

Red light [65]. We attribute this to the fact that even though average antenna size $\quad$ 508

decreases with increasing cost and decreasing visible light, moving from Chl d or Chl f $\quad$ 509

to the reaction centre involves a smaller enthalpy decrease, and hence reduces the free $\quad$ 510

energy gradient, leading to slower energy transfer between the two. $\quad$ 511

Initial populations: contingent vs. *in-situ* evolution $\quad$ 512

So far all antenna have been assumed to evolve *in situ* from a random population of $\quad$ 513

proto-antennae. This may miss potentially viable antenna configurations if no simple $\quad$ 514

evolutionary path connects them to the proto-antennae. To investigate this, we take a $\quad$ 515

population of "shallow water" antennae composed of APC and PC and let them *adapt* $\quad$ 516

to "deep water" conditions (see Fig. 5C and D), with further detail in Supplementary $\quad$ 517

Material Figs. S10 and S11). We find that our evolved antennae stay close in absorption $\quad$ 518

profile to this initial population for each cost value, with phycobilisome-like antennae $\quad$ 519

composed of either APC or Chl b, PC and PE shown in Fig. 5B, but in significantly $\quad$ 520

Fig 5. Comparison of contingent versus *in situ* evolution of antennae under 2.5m depth of water. A.) Average initial absorption spectrum for *in situ* evolution (solid lines), as a function of cost, compared to the initial spectrum of the unevolved population (dashed line).B.) Averaged final absorption spectra for the contingent (solid lines) populations as a function of cost compared to the initial spectrum (dashed line). C and D.) Diagrams illustrating evolution of *in situ* (C) and pre-adapted (D) populations from left to right, with different coloured blocks representing different pigment types (colours arbitrary).

different proportions compared to the *in-situ* evolution shown in Fig. 5A. The APC antenna core is generally either retained or replaced with Chl *b*, but not removed entirely, and forms a larger part of the overall absorption profile in the population pre-adapted to shallow water. In terms of antenna performance, ν_e and ϕ are slightly higher on average for the *in-situ* evolved population, but essentially identical within the error bounds for the two populations. This can be thought of more generally in terms of maxima of the "fitness surface": there may be multiple different antenna configurations which are roughly equal in performance for a given light environment, illustrating the history-dependence of evolutionary solutions [66].

Oxygenic and anoxygenic photosynthesis in extra-solar light conditions

Given that our model appears qualitatively robust, we apply it to incident fluxes one would expect on the surface of exo-planets orbiting in the habitable zone of M- and

Fig 6. Generated absorption spectra as a function of cost for cool M-dwarf (stellar temperature 2300K) (top row) and larger K-dwarf (stellar temperature 3800K) (middle row) for antennae with anoxygenic (left column) and oxygenic (right column) reaction centres respectively. Our algorithm cannot produce viable oxygenic antennae at high cost values for the coolest M-dwarfs; by contrast the algorithm works well for anoxygenic photosynthesis for all cost values and stellar temperatures. Averaged electron output of the antennae are shown (bottom row) as a function of cost for various stellar temperatures, showing dramatically increasing performance of oxygenic photosynthesis as the stellar temperature increases compared to broadly similar performance for anoxygenic photosynthesis. For detailed pigment compositions and absorption spectra of the intermediate temperatures see Supplementary Material.

Light	RC type	Electron output (s^{-1})	Pigment composition
2300K	Anoxygenic	60-75	BChl b, BChl a
	Oxygenic	$\approx$0-30	Chl b, Chl d
3800K	Anoxygenic	70-80	BChl b, BChl a, Chl d
	Oxygenic	65-75	APC, PC, Chl b, Chl a

Table 2. Details of antenna performance and pigment composition for anoxygenic and oxygenic photosynthesis as a function of stellar temperature.

K-dwarf stars. As in our previous work, these are generated by taking model spectral luminosities for various stellar temperatures [67], and assuming an orbital distance that would allow for liquid water [7,68]. Since recent research has noted the potential feasibility of *anoxygenic* photosynthesis arising around cooler stars [69], we compare our oxygenic model to a crude anoxygenic model with a RC wavelength of 890 nm [70–72]. Fig. 6 shows simulated absorption spectra and respective electron output for oxygenic and anoxygenic models for M- and K-dwarfs. We see that constrained by a 680 nm oxygenic RC, for the coolest M-dwarfs (top row), the antenna performance and even the location of the absorption peak varies significantly with cost–this is because lowering the cost allows larger antennae to evolve to capture more flux. Pigment compositions are detailed in Table. 2 and Supplementary Material Figs. S12–S14. The evolved population of antennae are composed mostly of Chl b at lower cost values, and exclusively Chl d at higher cost.

We note that even at the higher cost values, the low electron output does not necessarily mean oxygenic photosynthesis would be impossible: several species of cyanobacteria have adapted to very low levels of illumination [73], and simulated M-dwarf light does not limit growth of cyanobacteria, even without the FaRLiP response [6].

In contrast, for the warmer K-dwarfs (middle row) we see very reasonable antenna performance regardless of cost; interestingly, instead of exclusively Chl b antennae, we predict a phycobilisome-like composition of predominantly APC and PC with small amounts of chlorophyll mixed in.

For a 890 nm anoxygenic RC, under M-dwarf illumination, the model selects a larger antenna composed of BChl b with an absorption peak $\sim$ 800 nm. The electron output is much higher ($\nu_e \geq 60$ s^{-1}), which casually suggests that anoxygenic photosynthesis may be the better evolutionary choice, though numerous other factors will be important. For

warmer stars (Fig. 5B and 5C) we find antennae again predominantly composed of BChl b, but with BChl a and even Chl d included, as shown by the intermediate peak in the absorption spectra at around 700 nm.

Lastly we note that for the warmer K-dwarf stars ($T = 3800$ K) the performance of the oxygenic photosystem is roughly as good as that predicted for an anoxygenic photosystem, in agreement with recent research showing comparable growth for both vascular plants and desert cyanobacteria (both oxygenic organisms) under simulated K-dwarf light [74] to that in solar light.

Conclusions

Our model, which is *necessarily* coarse-grained, when coupled with a sophisticated evolutionary algorithm, seems to predict (qualitatively) the various oxygenic light-harvesting structures that have evolved in various Earth niches. This includes plant-like chlorophyll a and b antennae in redder light (typically of shaded environments) and cyanobacteria-like phycobilin-based antennae in bluer, sub-marine environments. Using a simple model of a far-red light adapted reaction centre, we show that the model predicts a shift from Chl a and b antennae to Chl d and f analogously to the FaRLiP response [20, 63, 64, 75] as the intensity of visible light decreases and the available light becomes redder and redder.

Applied to the light from M-dwarf stars, it implies that oxygenic photosynthesis, with organisms whose antennae are chlorophyll-like, may still evolve, though with reduced output relative to Earth organisms. This may limit oxygenic photosynthesis to simpler, single-cell organisms that, unlike vascular plants, do not have to support large amounts of non-photosynthetic tissue.

We see these results as a promising first step in producing a universal model of photosynthesis. They suggest that we can, cautiously, hypothesize on the nature of biological processes beyond Earth. That said, the model presented here does have certain shortcomings, consideration of which may help to direct subsequent work.

Firstly, the results obtained from the genetic algorithm seem relatively robust with respect to its main hyperparameter χ, which for a given light environment serves primarily to constrain the size of the antenna and prevent pigments being added which

capture no flux, rather than qualitatively altering the evolved structures. However, χ is a relatively broad and nebulously-defined quantity, and any further work focusing on more specific organisms or light environments may need refined strategies for constraining antenna size, for example by separating it into multiple separate parameters representing different costs. In addition, it does not account for the overall light intensity, and so the range of reasonable χ values decreases as the light intensity does; one obvious potential solution to this is to tie the cost to the integrated flux received by the antenna in some way. Organisms evolving in flux-limited environments may very well be rather finely-tuned, minimizing the overall cost of photosynthesis. How exactly such ideas should be incorporated into this model will require further, careful consideration.

Our genetic algorithm also allows for infinitely plastic behaviour–for example, an organism spontaneously starting to produce completely new kinds of pigments–which is biologically unlikely to say the least. Further work which focuses more specifically on the evolution of photosynthesis in specific terrestrial light environments would need to consider the mutation process more carefully in particular to address this.

Another limitation of the model as it stands concerns the relatively crude treatment of the RC. We use the simplest possible kinetics scheme of a two-step process involving fast primary charge separation, which reflects the transition from exciton transfer to charge transfer, followed by much slower 'electron output'. Of course this neglects the fact that the cycle of charge separation, acceptor reduction and donor oxidation is a multi-step process, on a hierarchy of timescales, that must compete with side processes such as *de-trapping* (reformation of the exciton) and charge recombination. We have focused here exclusively on light absorption and energy transfer by the antenna, as in previous works [34, 36], as effective light-harvesting strategies are a near universal feature of photosynthesis and will be required doubly-required if photoautotrophy is to evolve around M-dwarf stars. We have considered "standard" oxygenic RC energetics corresponding to PSII, a far-red adapted RC and an anoxygenic RC, but without explicitly considering the internal RC processes. A more quantitative description of the combined antenna-RC supersystem, for example to investigate the relative performance of anoxygenic vs. oxygenic photosynthesis or to examine the effect of redshifting the RC energy in more detail, is an obvious extension and will be the focus of further work.

Acknowledgements

CDPD and SC would like to acknowledge support from the BBSRC London Interdisciplinary Biosciences PhD Consortium (LIDo). CDPD and CG would like to acknowledge support from the Leverhulme Trust (RPG-2023-096) for direct funding of this project. The authors would also like to thank Thomas J. Haworth, Andrew R. Ringham, Edward Gillen and Conrad W. Mullineaux for useful discussions.

References

1. (phl upra edu) TPHLUA. Habitable Worlds Catalogue, PHL@UPR Arecibo; 2024.

2. McKay C. Requirements and Limits for Life in the Context of Exoplanets. Proceedings of the National Academy of Sciences of the United States of America. 2014;111. doi:10.1073/pnas.1304212111.

3. Kiang, Nancy. VPL Biological Pigments Database;. https://vplapps.astro.washington.edu/pigments.

4. Kiang NY, Siefert J, Govindjee, Blankenship RE. Spectral Signatures of Photosynthesis. I. Review of Earth Organisms. Astrobiology. 2007;7(1):222–251. doi:10.1089/ast.2006.0105.

5. Lehmer OR, Catling DC, Parenteau MN, Kiang NY, Hoehler TM. The Peak Absorbance Wavelength of Photosynthetic Pigments Around Other Stars From Spectral Optimization. Frontiers in Astronomy and Space Sciences. 2021;8.

6. Battistuzzi M, Cocola L, Liistro E, Claudi R, Poletto L, La Rocca N. Growth and Photosynthetic Efficiency of Microalgae and Plants with Different Levels of Complexity Exposed to a Simulated M-Dwarf Starlight. Life. 2023;13(8):1641. doi:10.3390/life13081641.

7. Chitnavis S, Gray C, Rousouli I, Gillen E, Mullineaux CW, Haworth TJ, et al. Optimizing Photosynthetic Light-Harvesting under Stars: Simple and General Antenna Models. Photosynthesis Research. 2024;162(1):75–92. doi:10.1007/s11120-024-01118-1.

8. Dodd MS, Papineau D, Grenne T, Slack JF, Rittner M, Pirajno F, et al. Evidence for early life in Earth's oldest hydrothermal vent precipitates. Nature. 2017;543(7643):1476–4687. doi:10.1038/nature21377.

9. Ricci F, Greening C. Chemosynthesis: a neglected foundation of marine ecology and biogeochemistry. Trends in microbiology. 2023;.

10. Moody ERR, Álvarez-Carretero S, Mahendrarajah TA, Clark JW, Betts HC, Dombrowski N, et al. The Nature of the Last Universal Common Ancestor and Its Impact on the Early Earth System. Nature Ecology & Evolution. 2024;8(9):1654–1666. doi:10.1038/s41559-024-02461-1.

11. Cardona T, Sánchez-Baracaldo P, Rutherford AW, Larkum AW. Early Archean Origin of Photosystem II. Geobiology. 2019;17(2):127–150. doi:10.1111/gbi.12322.

12. Meadows VS, Reinhard CT, Arney GN, Parenteau MN, Schwieterman EW, Domagal-Goldman SD, et al. Exoplanet biosignatures: understanding oxygen as a biosignature in the context of its environment. Astrobiology. 2018;18(6):630–662.

13. Schwieterman EW, Kiang NY, Parenteau MN, Harman CE, DasSarma S, Fisher TM, et al. Exoplanet Biosignatures: A Review of Remotely Detectable Signs of Life. Astrobiology. 2018;18(6):663–708. doi:10.1089/ast.2017.1729.

14. Fauchez TJ, Villanueva GL, Schwieterman EW, Turbet M, Arney G, Pidhorodetska D, et al. Sensitive probing of exoplanetary oxygen via mid-infrared collisional absorption. Nature Astronomy. 2020;4(4):372–376.

15. Sagan C, Thompson WR, Carlson R, Gurnett D, Hord C. A search for life on Earth from the Galileo spacecraft. Nature. 1993;365:715–721.

16. Seager S, Turner EL, Schafer J, Ford EB. Vegetation's Red Edge: A Possible Spectroscopic Biosignature of Extraterrestrial Plants. Astrobiology. 2005;5(3):372–390. doi:10.1089/ast.2005.5.372.

17. O'Malley-James JT, Kaltenegger L. The Vegetation Red Edge Biosignature Through Time on Earth and Exoplanets. Astrobiology. 2018;18(9):1123–1136. doi:10.1089/ast.2017.1798.

18. Leister D. Enhancing the Light Reactions of Photosynthesis: Strategies, Controversies, and Perspectives. Molecular Plant. 2023;16(1):4–22. doi:10.1016/j.molp.2022.08.005.

19. Rutherford AW, Osyczka A, Rappaport F. Back-Reactions, Short-Circuits, Leaks and Other Energy Wasteful Reactions in Biological Electron Transfer: Redox Tuning to Survive Life in O2. FEBS letters. 2012;586(5):603–616.

20. Nürnberg DJ, Morton J, Santabarbara S, Telfer A, Joliot P, Antonaru LA, et al. Photochemistry beyond the Red Limit in Chlorophyll f–Containing Photosystems. Science. 2018;360(6394):1210–1213. doi:10.1126/science.aar8313.

21. Henry TJ, Jao WC, Subasavage JP, Beaulieu TD, Ianna PA, Costa E, et al. The Solar Neighborhood. XVII. Parallax Results from the CTIOPI 0.9 m Program: 20 New Members of the RECONS 10 Parsec Sample. The Astronomical Journal. 2006;132(6):2360.

22. Dressing CD, Charbonneau D. The Occurrence of Potentially Habitable Planets Orbiting M Dwarfs Estimated from the Full Kepler Dataset and an Empirical Measurement of the Detection Sensitivity. The Astrophysical Journal. 2015;807(1):45.

23. Agol E, Dorn C, Grimm SL, Turbet M, Ducrot E, Delrez L, et al. Refining the Transit-Timing and Photometric Analysis of TRAPPIST-1: Masses, Radii, Densities, Dynamics, and Ephemerides. The Planetary Science Journal. 2021;2(1):1. doi:10.3847/PSJ/abd022.

24. Anglada-Escudé G, Amado PJ, Barnes J, Berdiñas ZM, Butler RP, Coleman GA, et al. A Terrestrial Planet Candidate in a Temperate Orbit around Proxima Centauri. nature. 2016;536(7617):437–440.

25. Blankenship RE, Madigan MT, Bauer CE. Anoxygenic photosynthetic bacteria. vol. 2. Springer Science & Business Media; 2006.

26. Overmann J, Garcia-Pichel F. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F, editors. The Phototrophic Way of Life. Berlin,

Heidelberg: Springer Berlin Heidelberg; 2013. p. 203–257. Available from: [706]
https://doi.org/10.1007/978-3-642-30123-0_51. [707]

27. Wood R, Donoghue PC, Lenton TM, Liu AG, Poulton SW. The origin and rise of [708] complex life: progress requires interdisciplinary integration and hypothesis [709] testing; 2020. [710]

28. Mills DB, Canfield DE. Oxygen and Animal Evolution: Did a Rise of [711] Atmospheric Oxygen "Trigger" the Origin of Animals? BioEssays : news and [712] reviews in molecular, cellular and developmental biology. 2014;36(12):1145–1155. [713]

29. Butterfield N. Oxygen, animals and oceanic ventilation: an alternative view; 2009. [714]

30. Cole DB, Mills DB, Erwin DH, Sperling EA, Porter SM, Reinhard CT, et al. On [715] the co-evolution of surface oxygen levels and animals. Geobiology. [716] 2020;18(3):260–281. [717]

31. Bozdag GO, Libby E, Pineau R, Reinhard CT, Ratcliff WC. Oxygen suppression [718] of macroscopic multicellularity. Nature Communications. 2021;12:2041–1723. [719]

32. Claudi R, Alei E, Battistuzzi M, Cocola L, Erculiani MS, Pozzer AC, et al. [720] Super-Earths, M Dwarfs, and Photosynthetic Organisms: Habitability in the Lab. [721] Life. 2021;11(1):10. doi:10.3390/life11010010. [722]

33. Bryant DA, Canniffe DP. How Nature Designs Light-Harvesting Antenna [723] Systems: Design Principles and Functional Realization in Chlorophototrophic [724] Prokaryotes. Journal of Physics B: Atomic, Molecular and Optical Physics. [725] 2018;51(3):033001. doi:10.1088/1361-6455/aa9c3c. [726]

34. Marosvölgyi MA, van Gorkom HJ. Cost and Color of Photosynthesis. [727] Photosynthesis Research. 2010;103(2):105–109. doi:10.1007/s11120-009-9522-3. [728]

35. Kiang NY, Segura A, Tinetti G, Govindjee, Blankenship RE, Cohen M, et al. [729] Spectral Signatures of Photosynthesis. II. Coevolution with Other Stars And The [730] Atmosphere on Extrasolar Worlds. Astrobiology. 2007;7(1):252–274. [731] doi:10.1089/ast.2006.0108. [732]

36. Arp TB, Kistner-Morris J, Aji V, Cogdell RJ, van Grondelle R, Gabor NM. Quieting a Noisy Antenna Reproduces Photosynthetic Light-Harvesting Spectra. Science. 2020;368(6498):1490–1495. doi:10.1126/science.aba6630.

37. Jennings RC, Engelmann E, Garlaschi F, Casazza AP, Zucchelli G. Photosynthesis and Negative Entropy Production. Biochimica et Biophysica Acta (BBA) - Bioenergetics. 2005;1709(3):251–255. doi:10.1016/j.bbabio.2005.08.004.

38. Broess K, Trinkunas G, van der Weij-de Wit CD, Dekker JP, van Hoek A, van Amerongen H. Excitation Energy Transfer and Charge Separation in Photosystem II Membranes Revisited. Biophysical Journal. 2006;91(10):3776–3786. doi:10.1529/biophysj.106.085068.

39. Dong H, Ghosh A, Kim MB, Li SW, Svidzinsky AA, Zhang Z, et al. On the Thermodynamics of the Difference between Energy Transfer Rate and Heat Engine Efficiency. European Physical Journal Special Topics. 2021;230:867–871. doi:10.1140/epjs/s11734-021-00091-3.

40. Novoderezhkin V, Marin A, van Grondelle R. Intra- and Inter-Monomeric Transfers in the Light Harvesting LHCII Complex: The Redfield–Förster Picture. Physical Chemistry Chemical Physics. 2011;13(38):17093–17103. doi:10.1039/C1CP21079C.

41. Broess K, Trinkunas G, van der Weij-de Wit CD, Dekker JP, van Hoek A, van Amerongen H. Excitation Energy Transfer and Charge Separation in Photosystem II Membranes. Biophysical Journal. 2006;91:3776–3786.

42. Taniguchi M, Du H, Lindsey JS. PhotochemCAD 3: Diverse Modules for Photophysical Calculations with Multiple Spectral Databases. Photochemistry and Photobiology. 2018;94(2):277–289. doi:10.1111/php.12862.

43. Lee PT, Yeh HY, Lung WQC, Huang J, Chen YJ, Chen B, et al. R-Phycoerythrin from Colaconema Formosanum (Rhodophyta), an Anti-Allergic and Collagen Promoting Material for Cosmeceuticals. Applied Sciences. 2021;11(20):9425. doi:10.3390/app11209425.

44. Nguyen HH, Song Y, Maret EL, Silori Y, Willow R, Yocum CF, et al. Charge Separation in the Photosystem II Reaction Center Resolved by Multispectral Two-Dimensional Electronic Spectroscopy. Science Advances. 2023;9(18):eade7190. doi:10.1126/sciadv.ade7190.

45. Oja V, Laisk A. Oxygen Yield from Single Turnover Flashes in Leaves: Non-Photochemical Excitation Quenching and the Number of Active PSII. Biochimica et Biophysica Acta (BBA)-Bioenergetics. 2000;1460(2-3):291–301.

46. van Amerongen H, Chmeliov J. Instantaneous Switching between Different Modes of Non-Photochemical Quenching in Plants. Consequences for Increasing Biomass Production. Biochimica et Biophysica Acta (BBA) - Bioenergetics. 2020;1861(4):148119. doi:10.1016/j.bbabio.2019.148119.

47. Baker NR, Rosenqvist E. Applications of Chlorophyll Fluorescence Can Improve Crop Production Strategies: An Examination of Future Possibilities. Journal of Experimental Botany. 2004;55(403):1607–1621. doi:10.1093/jxb/erh196.

48. Semer J, Navrátil M, Špunda V, Štroch M. Chlorophyll Fluorescence Parameters to Assess Utilization of Excitation Energy in Photosystem II Independently of Changes in Leaf Absorption. Journal of Photochemistry and Photobiology B: Biology. 2019;197:111535. doi:10.1016/j.jphotobiol.2019.111535.

49. Eiben AE, Smith JE. Introduction to Evolutionary Computing. Natural Computing Series. Berlin, Heidelberg: Springer; 2015.

50. Holland JH. Adaptation in Natural and Artificial Systems: An Introductory Analysis with Applications to Biology, Control, and Artificial Intelligence. The MIT Press; 1992.

51. Alhijawi B, Awajan A. Genetic Algorithms: Theory, Genetic Operators, Solutions, and Applications. Evolutionary Intelligence. 2024;17(3):1245–1256. doi:10.1007/s12065-023-00822-6.

52. Mühlenbein H, Schlierkamp-Voosen D. Predictive Models for the Breeder Genetic Algorithm I. Continuous Parameter Optimization. Evolutionary Computation;1(1):25–49. doi:10.1162/evco.1993.1.1.25.

53. National Renewable Energy Laboratory, American Society for Testing and Materials. Reference Air Mass 1.5 Spectra; 2024. https://www.nrel.gov/grid/solar-resource/spectra-am1.5.html.

54. Ruban AV, Wilson S. The Mechanism of Non-Photochemical Quenching in Plants: Localization and Driving Forces. Plant and Cell Physiology. 2021;62(7):1063–1072. doi:10.1093/pcp/pcaa155.

55. Eshaghi S, Andersson B, Barber J. Isolation of a Highly Active PSII-LHCII Supercomplex from Thylakoid Membranes by a Direct Method. FEBS Letters. 1999;446(1):23–26. doi:10.1016/S0014-5793(99)00149-0.

56. Lea-Smith DJ, Bombelli P, Dennis JS, Scott SA, Smith AG, Howe CJ. Phycobilisome-Deficient Strains of Synechocystis Sp. PCC 6803 Have Reduced Size and Require Carbon-Limiting Conditions to Exhibit Enhanced Productivity. Plant Physiology. 2014;165(2):705–714. doi:10.1104/pp.114.237206.

57. Voerman SE, Ruseckas A, Turnbull GA, Samuel IDW, Burdett HL. Red Algae Acclimate to Low Light by Modifying Phycobilisome Composition to Maintain Efficient Light Harvesting. BMC Biology. 2022;20(1):291. doi:10.1186/s12915-022-01480-3.

58. Ho MY, Gan F, Shen G, Zhao C, Bryant DA. Far-Red Light Photoacclimation (FaRLiP) in Synechococcus Sp. PCC 7335: I. Regulation of FaRLiP Gene Expression. Photosynthesis Research. 2017;131(2):173–186. doi:10.1007/s11120-016-0309-z.

59. Stomp M, Huisman J, Stal LJ, Matthijs HCP. Colorful Niches of Phototrophic Microorganisms Shaped by Vibrations of the Water Molecule. The ISME Journal. 2007;1(4):271–282. doi:10.1038/ismej.2007.59.

60. Montgomery BL. Seeing New Light: Recent Insights into the Occurrence and Regulation of Chromatic Acclimation in Cyanobacteria. Current Opinion in Plant Biology. 2017;37:18–23. doi:10.1016/j.pbi.2017.03.009.

61. Kehoe DM, Gutu A. Responding to Color: The Regulation of Complementary Chromatic Adaptation. Annual Review of Plant Biology. 2006;57(Volume 57, 2006):127–150. doi:10.1146/annurev.arplant.57.032905.105215.

62. Hamaguchi T, Kawakami K, Shinzawa-Itoh K, Inoue-Kashino N, Itoh S, Ifuku K, et al. Structure of the Far-Red Light Utilizing Photosystem I of Acaryochloris Marina. Nature Communications. 2021;12(1):2333. doi:10.1038/s41467-021-22502-8.

63. Gan F, Shen G, Bryant DA. Occurrence of Far-Red Light Photoacclimation (FaRLiP) in Diverse Cyanobacteria. Life. 2015;5(1):4–24. doi:10.3390/life5010004.

64. Gisriel CJ, Shen G, Brudvig GW, Bryant DA. Structure of the Antenna Complex Expressed during Far-Red Light Photoacclimation in Synechococcus Sp. PCC 7335. Journal of Biological Chemistry. 2024;300(2). doi:10.1016/j.jbc.2023.105590.

65. Nien TS, Bryant DA, Ho MY. Use of Quartz Sand Columns to Study Far-Red Light Photoacclimation (FaRLiP) in Cyanobacteria. Applied and Environmental Microbiology. 2022;88(13):e00562–22. doi:10.1128/aem.00562-22.

66. Blount ZD, Borland CZ, Lenski RE. Historical Contingency and the Evolution of a Key Innovation in an Experimental Population of Escherichia Coli. Proceedings of the National Academy of Sciences. 2008;105:7899–7906.

67. Husser, T -O , Wende-von Berg, S , Dreizler, S , Homeier, D , Reiners, A , Barman, T , et al. A New Extensive Library of PHOENIX Stellar Atmospheres and Synthetic Spectra. Astronomy and Astrophysics (A&A). 2013;553:A6. doi:10.1051/0004-6361/201219058.

68. Duffy CDP, Canchon G, Haworth TJ, Gillen E, Chitnavis S, Mullineaux CW. Photosynthesis under a Red Sun: Predicting the Absorption Characteristics of an Extraterrestrial Light-Harvesting Antenna. Monthly Notices of the Royal Astronomical Society. 2023;526(2):2265–2277. doi:10.1093/mnras/stad2823.

69. Coelho LF, Kaltenegger L, Zinder S, Philpot W, Price TL, Hamilton TL. Purple Is the New Green: Biopigments and Spectra of Earth-like Purple Worlds.

Monthly Notices of the Royal Astronomical Society. 2024;530(2):1363–1368. doi:10.1093/mnras/stae601.

70. Deisenhofer J, Michel H. The Photosynthetic Reaction Center from the Purple Bacterium ¡i¿rhodopseudomonas Viridis¡/I¿. Science. 1989;245(4925):1463–1473. doi:10.1126/science.245.4925.1463.

71. Leonova MM, Fufina TYu, Vasilieva LG, Shuvalov VA. Structure-Function Investigations of Bacterial Photosynthetic Reaction Centers. Biochemistry (Moscow). 2011;76(13):1465–1483. doi:10.1134/S0006297911130074.

72. Niedringhaus A, Policht VR, Sechrist R, Konar A, Laible PD, Bocian DF, et al. Primary Processes in the Bacterial Reaction Center Probed by Two-Dimensional Electronic Spectroscopy. Proceedings of the National Academy of Sciences. 2018;115(14):3563–3568. doi:10.1073/pnas.1721927115.

73. Pierangelini M, Stojkovic S, Orr PT, Beardall J. Photo-Acclimation to Low Light—Changes from Growth to Antenna Size in the Cyanobacterium Cylindrospermopsis Raciborskii. Harmful Algae. 2015;46:11–17. doi:10.1016/j.hal.2015.04.004.

74. Vilović I, Schulze-Makuch D, Heller R. Observation of Significant Photosynthesis in Garden Cress and Cyanobacteria under Simulated Illumination from a K Dwarf Star. International Journal of Astrobiology. 2024;23:e18. doi:10.1017/S1473550424000132.

75. Ho MY, Soulier NT, Canniffe DP, Shen G, Bryant DA. Light Regulation of Pigment and Photosystem Biosynthesis in Cyanobacteria. Current Opinion in Plant Biology. 2017;37:24–33. doi:10.1016/j.pbi.2017.03.006.