Evolutionary dynamics of pro-inflammatory caspases in primates and
rodents

Authors: Mische Holland[a], Rachel Rutkowski[a], Tera C. Levin[a#]

[a] Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA

#Address correspondence to Tera C. Levin: teralevin@pitt.edu

Abstract

Caspase-1 and its related protease homologs are key players in inflammation and innate
immunity. Because some pro-inflammatory caspases can detect microbial molecules and can
be targets of pathogen antagonists, we hypothesized that some of these genes could be
involved in evolutionary arms races between hosts and pathogens. If so, rapid caspase
evolution may lead to differences in activity and function between primates and rodents and
thus discrepancies between humans and animal model studies. Here, we characterize the
evolutionary history of caspase-1 and its close relatives across 15 primate and 21 rodent
species. While caspase-1 has been retained in all these taxa, we find that it has evolved rapidly
at the sequence level in primates. At the same time, the other members of the caspase-1
subfamily (caspase-4, -5, -11, -12, and CARD16, 17, and 18) each have unique evolutionary
trajectories. Whereas caspase-4 is found across primates, we identified multiple instances of
pseudogenization and gene loss in caspase-5, caspase-11, and the CARDs. Because
caspases-4 and -11 bind and recognize cytosolic LPS in the non-canonical inflammasome
pathway, we expected that these proteins would be most likely to evolve rapidly. Instead, we
found that these two proteins are largely conserved, whereas caspase-4's close paralog,
caspase-5, showed significant indications of positive selection. Caspase-12 is thought to be a
non-functional pseudogene in humans. We find this extends across primates, although most
rodents retain an intact, and likely functional, caspase-12. Finally, unlike rodents, primate
caspases have undergone many, repeated rounds of gene duplication, gene conversion, and
gene loss leading to a highly dynamic pro-inflammatory caspase repertoire. We discuss the
potential implications of this evolutionary history for our understanding of caspase gene
functions.
30

Introduction

Pathogen-interacting genes are among the most rapidly evolving in mammalian
genomes [1]. Because of the strong selective pressures involved, such genes often diversify
through evolutionary arms races, which involve multiple, recurrent changes in host and
pathogen molecules as each protein reciprocally adapts [2]. Common mechanisms of
diversification in arms races include gene family expansion and contraction, gene conversion,
and rapid amino acid turnover, particularly at sites of host-pathogen binding [2–4].
38
Innate immune pathways, including those involved in inflammation, are key defenses
against many human pathogens and deficiencies in these pathways lead to heightened
pathogen susceptibility (e.g. [5]). Over-activation of inflammation can also be dangerous for the
host, both during infection [6] and in autoimmunity [7]. Therefore, we expect natural selection to
act strongly on genes in inflammatory pathways, tuning these proteins to recognize an ever-
changing array of microbial infections while avoiding self-harm. Indeed, some of the proteins
involved in these pathways (e.g. inflammasome proteins NLRPs & CARD8) have been found to
rapidly diversify, with signatures of evolutionary arms races [8–10]. We therefore sought to
examine the evolutionary history and diversity of the pro-inflammatory caspase genes, which
participate in these pathways.
49

Caspase-1 is a protease that is a hub of pro-inflammatory signaling. In the 'canonical'
signaling pathway, multi-protein complexes called inflammasomes assemble in response to
cellular or microbial signals [11]. Caspase-1 is recruited to inflammasomes via its CARD
(caspase activation and recruitment domain), where its oligomerization leads to self-cleavage
and activation. The active caspase-1 protease (made up of p20 and p10 subunits, see Fig. 4 for
domain architectures) then targets for maturation the pro-inflammatory cytokines IL-1beta and
IL-18, as well as gasdermin D, which forms membrane pores and leads to an explosive cell
death known as pyroptosis [7].
58
In addition, there are several caspase-1 homologs (collectively known as the caspase-1
subfamily), which all reside within the same syntenic locus as the caspase-1 gene, *CASP1*, in
primates and rodents [12] (Fig. 1A). One of these genes, rodent *CASP11*, has two homologs in
primates, called *CASP4* and *5*, which have been implicated in 'non-canonical' inflammatory
signaling. (Note: although rodent *CASP11* is clearly the ortholog of primate *CASP4/5* and could
be called *CASP4*, we refer to it here as *CASP11*, as this name is more commonly used in the
biomedical, mouse-centric literature.) In the non-canonical pathway, caspase-4 or -11 directly
binds to cytosolic, bacterial LPS, leading to caspase oligomerization, activation, cleavage of
cytokines and gasdermin D, and inflammatory cell death [13,14]. Thus, the non-canonical
pathway can sense and respond to cytosolic bacterial pathogens through direct caspase
activation. Some caspase-5 mutations have been associated with cancer and it is largely
assumed that it participates in similar non-canonical signaling[14]. However, the molecular
functions of caspase-5 have remained more enigmatic, with mainly *in vitro* evidence supporting
similar biochemical activities to its caspase-4 paralog[13].
73
Other caspase-1 subfamily genes with less characterized roles include caspase-12,
which is generally considered non-functional in humans [15–17], and three *CASP1*-like genes
that contain a CARD as their only protein domain (the CARDs, aka CARD-only proteins, or
COPs). The CARD-only genes are named *CARD16, 17*, and *18* (or alternatively *COP1, Inca*,
and *Iceberg*, respectively) [18–20]. Overexpression studies of the CARDs suggested that they
may act as caspase-1 inhibitors [18,19,21] and *in vitro* studies suggested that CARD16 and 18
promote CASP1 CARD oligomerization whereas CARD17 inhibits it [22]. However, a recent *in*
vivo study of CARDs found that they each inhibit caspase-1 activation by competitively binding
to the CARDs of caspase-1 and inflammasome adapters, blocking caspase-1 recruitment to
inflammasomes [23]. These CARDs apparently act non-redundantly, dampening caspase-1
activation during different stages of inflammasome signaling.
85
Pathogens also have abundant mechanisms to evade and shut down these
inflammatory signaling pathways, including antagonists that act both upstream [24,25] and
downstream [26] of caspase-1 activation, as well as multi-caspase inhibitors that can target
caspase-1, -4, and -5 by forming covalent bonds at the caspase active site [27,28]. Specific
microbial effectors that bind caspase-4 and -11 have also been identified, including the OspC3
effector from *Shigella* that inhibits caspase function through ADP-riboxination[29]. Because the
members of the caspase-1 subfamily are key players in immune defenses, which can directly
bind pathogen molecules and are often antagonized by pathogens, we hypothesized that they

94 may evolve in evolutionary arms races, similar to those that have diversified inflammasome
95 proteins[30].
96

97   Previous studies have identified some hints that the caspase-1 subfamily has been
98 evolutionarily dynamic. For example, instances of gene duplication and loss have been
99 observed in mammals, molluscs, and insects[12,31]. Some of the diversity of caspases has
100 interesting and important functional implications, including the caspase-1/4 fusion proteins in
101 carnivores[32]. Because mice are the most commonly used animal models of pro-inflammatory
102 signaling pathways, it is particularly important to understand the similarities and differences in
103 caspases between primates and rodents. However, prior studies of caspase evolution included
104 at most 4 distantly related rodents and 3 primates[12], limiting our evolutionary resolution in
105 these clades.
106

107   Here, we take an in-depth comparison of evolution of the inflammatory caspase locus
108 across 15 primate and 21 rodent species. We find that the caspase-1 locus is more
109 evolutionarily dynamic in primates than in rodents, with evidence of recurrent gene duplications,
110 pseudogenization of new genes, and gene conversion events. Contrary to our expectations, the
111 signatures of selection were strongest and most evident in primate *CASP1* and *5* as compared
112 to the other caspases, with interesting phenotypic implications. Combined with prior evidence of
113 human segregating polymorphisms[33] and copy number variation[34] within the caspase-1
114 locus, we propose that these genes have not only a history of diversification, but also abundant,
115 ongoing genetic innovation in primates.
116

Results

Differential caspase gene birth and loss in primates and rodents

119   Caspase-1 subfamily genes all reside within a single syntenic locus (Fig. 1A), found in
120 humans on chromosome 11q22.3. To assess the repertoires of these genes across species, we
121 analyzed the locus across 15 primate and 21 rodent genomes. These species were selected
122 based on: (1) sampling taxa across the phylogeny that were not too diverged from our reference
123 human and house mouse species [i.e. *CASP1* dS<0.4], and (2) whether the full, ~350kb locus
124 was well-assembled on a single contig in the genome. Within the locus, we identified caspase
125 genes by searching for exons that had 80% (primate) or 60% (rodent) nucleotide identity to the
126 annotated human or house mouse genes. In cases where diverged exons were not identified,
127 we used synteny and alignments to the reference coding sequences to locate the missing
128 exons.

129   Through this process, we classified every homolog into one of four categories. First, we
130 called genes as "present" if they were found within their syntenic location and had a conserved,
131 in-frame sequence that aligned across its full length to the reference. If our initial searches failed
132 to find the homolog or identified only partial genes, we used blastn to search the Refseq
133 Representative Genomes database of predicted or sequenced transcripts for the homolog. If
134 this search yielded evidence of an in-frame transcript, we then identified each exon in the locus,
135 realigned the coding sequence based on transcript evidence, and called the gene as "present".

Some homologs only had shorter in-frame sequences that did not span the full length of the
reference. We categorized these homologs in the second category, "partial in-frame sequence".
If we identified some traces of the gene but did not find evidence of an in-frame coding
sequence in either the genome or transcript database, we placed them in a third category,
"pseudogenes". Finally, if the gene was undetectable and/or if we found fewer than 70% of the
exons as compared to the reference gene, we categorized the homolog as "absent".

Fig 1. Differential patterns of gene gain and loss within the caspase-1 locus. A) Genes in the caspase-1 locus in human and mouse (not to scale). B) PhyML phylogenetic tree of intact caspase genes demonstrates that orthologs have been correctly named and categorized, despite gene losses. Numbers at nodes are aLRT support values. C-D) Gene presence/ absence across primate (C) and rodent (D) species, with colored Xs showing the history of gene losses. See text for definitions of pseudogene vs. partial sequence vs. gene absence. *CASP1* has been retained across all species examined, as has primate *CASP4*. In contrast, there have been multiple cases of gene degradation and loss of rodent *CASP11*, primate *CASP5*, and the primate *CARD* genes. *CASP12* appears to be retained and functional across most (but not all) rodents, whereas it has been largely degraded and lost in primates.

For all caspases that were present and intact, we confirmed their identities using a
phylogenetic tree, which showed that all genes were correctly categorized into their *CASP1, 4,*
5, 11, or *12* gene families (Fig 1B). As expected based on prior studies [12,31], this analysis
also showed that *CASP4* and *5* are primate-specific paralogs most closely related to the rodent
CASP11 gene. For *CASP12*, although we recovered many rodent homologs, primate
sequences were excluded from the tree, as we recovered no instances where a full-length
primate CASP12 was present. To understand the history of gene gains and losses, we then
assessed which species had intact caspase homologs.

<u>Caspase-1</u>: *CASP1* was present within the syntenic locus and we recovered full-length, well-
aligned coding sequences for all 36 primate and rodent species evaluated (Fig. 1C, 1D). This
pattern of widespread gene retention implies that *CASP1* plays an important, non-redundant
role in organismal fitness across primates and rodents.

<u>Caspase-12</u>: For *CASP12*, we find that the previously reported partial gene loss in humans
[15,16] extends across primates, with multiple lineages experiencing additional gene
degradation over time. Thus, we were not able to extract full length *CASP12* coding sequences
from any of the primate genomes. Neither the SHS>SHG mutation found within humans that
disrupts catalytic activity [15] nor the residue 125 human stop codon polymorphism[17] is
shared between humans and other primates, although each primate species has numerous
CASP12 frameshifts and early stop codons. In contrast, in rodents *CASP12* is present in most
species, with a similar length and domain architecture as the other caspase-1 subfamily genes.
We estimate there have been up to six additional instances of independent, lineage-specific
CASP12 pseudogenization and/or loss events within rodents (Fig. 1D), suggesting that *CASP12*
loss is frequent across multiple mammalian clades. To test if CASP12 is slowly becoming a
pseudogene across rodents as it has in primates, we made an alignment of intact rodent
CASP12 sequences and analyzed it with CODEML (model 0 vs. 0a)[35]. We detected strong
evidence that rodent *CASP12* is under purifying selection (Supp Table X), suggesting that it is
likely a functional gene in most rodent species.

<u>Caspase-4, -5, and 11</u>: For *CASP4, 5*, and *11*, because these proteins are thought to share
similar immune functions, we thought they might exhibit similar patterns of evolution.
Alternatively, because recently duplicated genes often lose one paralog and return to single
copy, we expected *CASP11* might be conserved across rodents, whereas *CASP4* and/or
CASP5 would experience some reciprocal losses in primates. In contrast to both hypotheses,
we found *CASP4* was present and shared across all primate genomes whereas there were
multiple cases of pseudogenization (and some cases of complete gene loss) in rodent *CASP11*
and primate *CASP5* (Fig. 1C & D). Because of these losses, four diverse rodent species, *A.*
sylvaticus, M. auratus, M. orchrogaster, and M. marmota, have *CASP1* as their only caspase-1
subfamily gene.

Within primates, we also found at least two species that lack all three *CARDs* (Fig 1C).
Based on the distribution of *CASP5* and the *CARDs* across the primate tree, we can infer that
these genes originated in the last common ancestor of all primates, ~54 million years ago[36].

However, since then all four of these primate-specific genes have experienced multiple,
repeated instances of gene loss.

Recent, repeated gene conversion and duplications in the primate *CASP1* locus

In addition to the caspase and CARD genes described above, the human locus contains
multiple caspase-like pseudogenes, indicating that there have been additional historical
duplication events followed by gene loss. These include two *CASP1*-like pseudogenes that lie
between the *CARDs* (*Casp1P1* and *P2*) and a *CASP4*-like pseudogene (*Casp4LP*) near
CASP12 (Fig. 2). Inspired by this history of innovation, we next examined the nucleotide
sequence of the locus for evidence of recent duplications or gene conversion events that may
not be reflected in simple gene presence/absence data. To do so, we searched for extended
blocks of similar nucleotide sequence, defining blocks of self-similarity as regions at least 1kb
long that contained at least 50% nucleotide identity to another region within the 350kb locus.

In the human locus, we detected many regions of self-similarity, including those that
duplicated both coding and non-coding sequences (Fig. 2A & B). For example, a ~6kb region of
the *CASP4* locus (including both exons and introns) was 60% identical to the *CASP4LP*
pseudogene and 54% identical to the *CASP5* locus. We found *CASP4LP* loci only within great
apes, suggesting that this locus arose ~13Mya, whereas CASP5 dates back to the last common
ancestor of all primates ~54 Mya[36] (Fig 1). These findings (and those of Fig 3 & 4 below)
suggest that *CASP4* and *CASP5* have experienced recent gene conversion events that
overwrote the sequence of one locus with the other at one or more times during primate
evolution. We found similar blocks of self-similarity in the human locus between *CASP1* and
CARD16, CARD17, and nearby pseudogenes (Fig 2A), as well as blocks that included intronic,
intergenic, or promoter regions (Fig 2B). Although *CARD18* arose via *CASP1* duplication at
approximately the same time as *CARD16 & 17* (Fig. 1), it appears that *CARD18* has not
undergone recent gene conversion, as *CARD18*'s sequence is now too diverged to be identified
in this analysis. Overall, recurrent segmental duplications and gene conversion events within the
primate CASP1 locus appear to be common.

Fig 2. The caspase-1 locus has undergone recent gene conversions and tandem duplications in primates, but not rodents. A-C) Annotated genes (filled colored arrows) and pseudogenes (white arrows with colored outlines) in the human (A-B) and mouse (C) loci. Colored arcs (A) connect regions of extended nucleotide identity between caspase genes or pseudogenes, indicative of recent tandem gene duplications or gene conversions. Gray arcs (B and C) show regions of sequence identity within intronic or intergenic regions.

These events have significantly impacted the sequences, and potentially functions, of
genes in the primate locus, as illustrated by a phylogenetic tree of primate CARD sequences
(Fig 3). Unlike in the full caspase gene tree (Fig 1B), in the CARD tree the *CASP5* branches
emerge out of the *CASP4* clade, consistent with a scenario in which *CASP4* and *5* arose from a
duplication event and diverged from each other, then *CASP5* acquired a *CASP4*-like CARD
sequence via gene conversion, followed by additional divergence. In this tree, we also observed
that *CARD18* was clearly differentiated from other primate caspase-like genes, whereas there
was an intermingling of the *CASP1*, *CARD16*, and *CARD17* sequences (Fig 3B). Specifically,
we saw that the sequences of *CASP1* and *CARD16* were intermixed together in Old world
monkeys, independently intermixed within Great apes, and that the sequence of marmoset
CASP1 was most similar to marmoset *CARD17*. Based on gene synteny and the presence of
CARD16 and *17* across primates, we infer that these genes arose in the last common ancestor
of all primates.

However, in at least
these three instances,
CARD16 and *17* have
undergone recent gene
conversion with *CASP1*.
We predict that such
gene conversion events
would enable CARD16
and CARD17 to retain
strong, homotypic
binding to the CASP1
CARD, despite
divergence in the
CASP1 CARD
sequence over time.

In contrast to the
highly dynamic human
locus, few blocks of self-
similarity exist within the
house mouse locus (Fig.
2C). Indeed, we did not
detect any instances of
extended nucleotide
identity that included
coding regions of any
caspase. We did detect
multiple transposon
insertions within
intergenic and intronic
regions (specifically the

Fig 3. PhyML phylogenetic tree of the CARDs of each primate caspase-1 subfamily gene. Region within the dashed box in A) is shown in detail in panel B), with gene identity indicated by symbols. Although sequences for *CASP4, 5*, and *CARD18* have distinct branches, in several species the *CASP1*, CARD16, and/or *CARD17* sequences are highly similar and intermixed on the tree, indicating recent gene conversion. All illustrated nodes on the PhyML trees had aLRT support statistics >5. Other nodes were collapsed.

Long interspersed nuclear element L1MdMus and ERVL-MaLR family of LTR transposon [37]),
demonstrating that we had the power to detect such duplication events. However, unlike in the
human locus, we did not find any evidence in the mouse genome of ongoing caspase-1
subfamily duplication or gene conversion events.

Finally, within the human population it appears that duplications within the *CASP1* locus
are ongoing. Copy number variation among humans is common, including between the
GRCh38 genome assembly, which we focused on here, and the recent telomere-to-telomere
genome sequence of the CHM13 cell line, which derives from a single human haplotype (T2T-
CHM13) [34]. While segmental duplications tend to get collapsed in most genome assemblies,
the T2T-CHM13 genome is currently unique in its completeness and ability to capture structural
variation. In this haploid genome, the *CASP1* locus was identified as one of the regions with
highly variable gene dosage, with T2T-CHM13 containing an estimated one copy of *CARD16*,
two copies of *CASP12, CARD 17*, and *CARD18*, three copies of *CASP1*, five copies of *CASP5*,
and fourteen copies of *CASP4*. Thus, throughout primate history and within humans, the CASP1
locus appears to be a frequent site of gene duplication and innovation.

Unexpected patterns of positive selection in the caspase-1 subfamily genes

Canonical pro-inflammatory pathways activate caspase-1 downstream of pathogen
recognition. In contrast, caspase-4 and -11 have been found to directly bind multiple types of
pathogen molecules [13,29,38], whereas no pathogen antagonists have yet been identified that
specifically target caspase-1. Because both LPS and microbial effectors tend to be highly
variable and because we expect to see rapid evolution at host-pathogen binding interfaces, we
hypothesized that we would see strong signatures of positive selection within *CASP4* and
CASP11, potentially in the previously-defined LPS-binding residues of the CARD region[13],
whereas these signatures might be weaker or absent in *CASP1* and *CASP5*.

To test this hypothesis, we analyzed evolutionary genomic signatures in alignments of
primate *CASP1, CASP4*, and *CASP5* as well as rodent *CASP1* and *CASP11*. Recurrent gene
conversion events (as discussed above) can cause different parts of a gene to have different
evolutionary histories. To detect if this was the case and identify sites of recombination
breakpoints, we used the program GARD [39]. Consistent with our earlier results, GARD
identified multiple recombination breakpoints across the caspases, with more sites identified in
the primate than in the rodent genes: two recombination breakpoints in primate CASP1, one
breakpoint in CASP4, three breakpoints in CASP5, one breakpoint in rodent CASP1, and none
in rodent CASP11. Thus, many extant caspase genes are evolutionary patchworks, reflecting
multiple gene conversion events.

This sort of recombination history can interfere with tests for positive selection.
Therefore, we divided each alignment into different recombination segments and analyzed each
segment independently. For each, we used multiple algorithms to estimate the dN/dS ratio, i.e.
the rate of non-synonymous to synonymous substitutions. If the dN/dS ratio is significantly
higher than 1, that would indicate that the protein regions and amino acid sites have evolved
under positive selection, with more rapid amino acid changes than expected by neutral

mutation. For these statistical tests of positive selection, we used CODEML and FUBAR [35,40].
Both algorithms are designed to detect positive selection that is pervasive across the
phylogenetic tree and they use different statistical models of molecular evolution. Therefore,
these analyses detect related, but complementary, signatures of positive selection.
299

300 **Table 1. Signatures of selection in caspase genes**

Gene	Clade	Recomb. segment	dN/dS	M7 v. M8 p-value	M7 v. M8a p-value	Highest omega in M8	# of positively selected sites in CODEML	# of positively selected sites in FUBAR
CASP1	Primate	1	0.39854	9.58E-01	9.59E-01	1.12	0	1
		2	**1.27225**	**6.98E-09**	**7.01E-09**	**8.62**	**7**	**12**
		3	**0.6961**	**3.50E-07**	**3.51E-07**	**3.73**	**12**	**14**
CASP4	Primate	1	0.35332	3.69E-01	4.75E-01	2.61	0	2
		2	0.23685	1.86E-01	2.75E-01	6.47	0	0
CASP5	Primate	**1**	**3.22362**	**1.80E-03**	**1.85E-03**	**4.79**	**14**	**1**
		2	0.99276	1.80E-02	1.85E-02	2.33	2	4
		3	0.33048	9.98E-01	9.97E-01	1.09	0	0
		4	1.09378	1.60E-02	1.61E-02	5.27	2	2
CASP1	Rodent	1	0.58534	3.60E-01	6.18E-01	1.29	0	1
		2	0.33853	6.76E-02	1.26E-01	1.78	1	6
CASP11	Rodent	1	0.41356	3.65E-01	7.32E-01	1.37	0	2
CASP12	Rodent	1	0.18069	1.00E+00	1.00E+00	1.00	0	0
		2	0.51008	1.00E+00	1.00E+00	1.00	0	0
		3	0.17643	1.00E+00	1.00E+00	1.00	0	2

301 Recombination segments that showed significant signatures of positive selection in CODEML
302 following Bonferonni correction (p<0.003) are bolded.
303

Contrary to our initial hypothesis, CODEML did not detect significant evidence of positive
selection in primate *CASP4* or rodent *CASP11* (Table 1, Fig 4A & B). Similarly, for these genes
only a few sites were called by FUBAR: two sites in the CARD of *CASP4* and two in the p20
subunit of *CASP11*. None of these sites overlapped with the LPS-binding residues proposed by
Shi et al., 2014[13], where we had expected to observe signatures of positive selection. Of the
FUBAR-identified sites, the most convincing was I156 in the p20 domain of mouse *CASP11*,
which toggled between I, S, T, and N codons across the 10 rodents analyzed. Interestingly, I156
was also the only residue where homologous sites in primate *CASP1* and *CASP5* were also
identified as positively selected (see below). In addition, this site is in proximity to the binding
interface between caspase-4 and -11 to a bacterial effector from *Shigella[29]*, suggesting that
this site may evolve rapidly to evade effector binding. However, with the possible exception of

the I156 residue, we found little to no evidence that *CASP4* or *11* have evolved under positive
selection, with no enrichment at putative LPS binding sites.

Fig 4. Positively selected residues in pro-inflammatory caspases in primates and rodents. **A-E**) Sites of positive selection, as determined by the FUBAR and CODEML algorithms. Arrowheads mark sites passing the statistical threshold for positive selection. Vertical dotted lines separate recombination segments identified by GARD, which were each analyzed independently. The domain architecture of each caspase is shown as colored boxes, with active sites illustrated (yellow *) along with the proposed LPS-binding residues of CASP4 and 11 according to Shi et. al 2014 (orange carats). In four cases, sites of positive selection in primate CASP1 were homologous to sites identified in one or more other caspase alignment (green, blue, red, or purple residue names). All coordinates are named according to the residues in the human or mouse sequences for primate or rodent analyses, respectively. F-G) Each identified site of CASP1 positive selection (red) is illustrated on the predicted AlphaFold structure of human (F) or mouse (G) caspase-1. The CARD, p20, and p10 domains are colored as in (A-E), with linker regions in gray, and active sites in yellow. Many more sites of positive selection were identified in primates than in rodents, but in neither clade do the sites show much co-localization in 3D space.

317
In contrast, we identified significant signatures of positive selection in both primate
CASP1 and *CASP5.* For *CASP5*, we analyzed sequences only in apes and Old world monkeys,
because *CASP5* in New world monkeys appears to be a pseudogene (Fig 4). Nevertheless,
despite the reduced statistical power from fewer sequences, we identified 21 sites under
positive selection between CODEML and FUBAR, 4 of which were called by both programs (Fig
4C). Unlike *CASP4*, *CASP5* has an extended N-terminus. This N-terminal region was enriched
for positive selection hits (56% of total sites identified by both algorithms were located within the
first 12% of the coding sequence length), with CODEML calling an especially large number of
sites. The *CASP5* N-terminal tail also included an insertion within Old world monkeys, further
diversifying this region. When we analyzed the N-terminal region both before and after trimming
out the Old world monkey insertion, CODEML and FUBAR identified many positively selected
sites in both cases, although the identities of the sites shifted. Therefore, while we are confident
that the N-terminal region of *CASP5* is rapidly evolving, the specific sites identified should be
interpreted with caution. While the N-terminus of *CASP5* has not previously been associated
with a known function, the diversification observed here could indicate that this part of the
protein is important for organismal fitness. We also note that *CASP5* has multiple annotated
splice isoforms, some of which exclude exon 2 which encodes most of the N-terminus.
Therefore, although these residues passed our statistical thresholds in one or more algorithms,
it's possible that the elevated rate of amino acid turnover in the N-terminus is due in part to
relaxed selection, as this region is less frequently part of the caspase-5 protein.
338
Even ignoring the *CASP5* N-terminus, we still detected more residues under positive
selection and with stronger support in *CASP5* than in its close paralog *CASP4* or rodent
CASP11 (Fig 4; Table 1). Of particular note, three positively selected sites were homologous
between primate CASP5 and CASP1, including V217, homolog of I156 in caspase-11. Within
humans, the V217 residue of CASP5 is polymorphic at high frequencies (11% of individuals in
Africa and 4% worldwide have L or M at this site) [33], demonstrating that variation at this site
still segregates within human populations, not only across different primate species.
346
Finally, in *CASP1*, we identified 29 sites of positive selection primates (16 called by both
CODEML and FUBAR) and 6 in rodents (1 called by both algorithms (Fig. 4D & E)). These sites
were distributed across the length of the protein, including sites within the CARD, p20, p10
domains, and the flexible linkers connecting these domains. Positively selected residues that
are distant in the primary sequence of a protein will sometimes co-localize in the folded protein,
often overlapping with host-pathogen binding interfaces. To see if the *CASP1* selected residues
potentially form this kind of interface, we examined them on the AlphaFold predicted structures
of primate and rodent caspase-1, but did not observe a large concentration of the residues in 3D
space. The localization was similar when we used experimental caspase-1 crystal structures
(not shown).
357
In summary, we detected little to no evidence of positive selection in *CASP4* and
CASP11, whereas there were strong signatures of selection in *CASP5* and *CASP1*, particularly
in primates. In four cases, sites we identified in primate *CASP1* were homologous to positively

selected sites in other caspases. However, with the exception of the N-terminal tail of *CASP5*,
the positively selected sites were not concentrated or co-localized in a particular region of the
protein.
364
365
Variation in caspase-1 aspartate self-cleavage sites
During caspase protease activation, protein oligomerization induces self-cleavage at
aspartate sites to generate the active enzyme [41]. The removal of the flexible linker between
the p20 and p10 subunits is essential for caspase-1 activation. The aspartate cleavage sites
flanking this linker are strictly conserved across caspase-1, 4, 5, and 11 (Fig 5). However, it has
been open to debate whether the CARD-p20 cleavage is essential for pro-inflammatory caspase
function. When we examined the aspartate site at the N-terminal end of p20, we found it was
conserved across caspase-4, 5, and 11, with the exception of one rodent species, *M. pahari*,
which had an asparagine at the caspase-11 site instead. Across caspase-1 homologs, we found
many more species that had variation in the CARD-p20 cleavage site (Fig 5), often with no
aspartate residues nearby that could serve as alternate cleavage sites. These species included
all of the New world monkeys and four species of rodents, including rats. We suggest that
functional studies of the caspase-1 homologs from these lineages may provide important
information about the mechanisms and activation of pro-inflammatory caspases, particularly the
role of CARD-p20 cleavage.

Fig 5. Caspase-1 aspartate cleavage sites between CARD and p20 domains are not strictly conserved. A) domain architecture of caspase-1, showing locations of aspartate cleavage sites ('D'). B-C) Alignments of the caspase-1 aspartate cleavage sites, with residues that do not match consensus highlighted with colored boxes. While the two C-terminal cleavage sites are highly conserved within primates (B) and rodents (C), the N-terminal site is more highly variable and some species lack the aspartate residue entirely. Arrowheads mark nearby positively selected sites.

Discussion

381   **Discussion**

382   While researchers commonly use mice as model systems for studying human pro-
383   inflammatory caspases, we find a number of differences in the genes and evolutionary
384   dynamics between primates and rodents, with potential implications for caspase functions.
385

First, the caspase-1 locus is more evolutionarily dynamic in primates, with evidence of
recurrent gene duplications, pseudogenization of new genes, and gene conversion events (Fig.
1-3). Such events can create blocks of nucleotide identity within the locus that can facilitate
further rearrangements or recombination, thus perpetuating complex genetic dynamics. The
rapid amino acid turnover, particularly in primate CASP5 and CASP1 (Fig. 4), further
accelerates the evolution of these genes. Combined with prior evidence of human segregating
polymorphisms and copy number variation within the caspase-1 locus, we propose that these
genes have not only a history of diversification, but also abundant, ongoing innovation in
primates. The genetic variation in the caspases is likely to cause differences in pro-inflammatory
immune responses across individual humans (and cell lines) as well as non-human primates,
which won't be reflected in mouse models. Therefore, we urge researchers to carefully assess
and report the caspase genotypes and gene copy number for the organisms and cell lines used
in their experimental studies.
399

Second, our work also provides additional information about differences in gene content.
CASP4 and *5* arose from a primate-specific duplication event, as did the *CARD* genes. We
traced all of these duplication events back to the base of the primate tree, approximately 43
million years ago[36]. The degeneration of primate *CASP12* appears similarly ancient, although
the gene is apparently functional and retained across most rodents. Because each of these
genes lacks 1-to-1 orthologs in rodents and primates, their human functions may not be
reflected in mouse models. *CASP12* is likely to have rodent-specific roles in mediating
inflammatory signaling, just as *CASP4/5* and the *CARDs* will have primate-specific biology.
More surprisingly, we discovered multiple pseudogenization and gene loss events of both
primate *CASP5* and rodent *CASP11*, including four rodent species that apparently have *CASP1*
as their only caspase-1 subfamily gene. Given the proposed importance of *CASP11* for
pathogen defense, we speculate that these four species may have acquired new genes or
alleles that can detect cytoplasmic LPS, thus compensating for *CASP11* loss.
413

The evolutionary signatures have interesting implications for the molecular functions of
the caspase-1 subfamily. For example, we were surprised to see such strong selective
signatures in *CASP5* as compared to its close homologs *CASP4* and *CASP11*, because current
studies have yet to identify *in vivo* phenotypes or molecular functions associated with caspase-5
(nor its unique N-terminal tail), whereas both caspase-4 and -11 are known to mediate important
host-microbe molecular interactions. These evolutionary discrepancies suggest that caspase-4
and -5 may have significant differences in their *in vivo* functions and that caspase-5 plays an
important, undefined, role supporting organismal fitness. For example, caspase-5 may
substitute for caspase-4 in certain contexts, or it may act as a molecular decoy for caspase-4
inhibitors. Although these scenarios will remain speculative until future experimental advances

in the role of caspase-5, the positively-selected residues we identified in Fig. 4C are likely to be
sites that mediate caspase-5-specific activities.
426
We were also surprised to see the widespread signatures of positive selection in primate
CASP1, as compared to the other caspase genes. This pattern suggests that caspase-1 is
under substantial selective pressure at multiple molecular interfaces, and that these pressures
are largely not shared with caspase-4 or -11. (The only exception being at I156 of *CASP11*/
V217 of *CASP5*/ T187 of *CASP1*, which may interact with similar pathogen antagonists).
Analogous patterns of protein evolution (i.e. many positively selected sites distributed
throughout the protein) have been found in examples such as PKR and MxB, which are thought
to be on the 'defense' of evolutionary arms races[2]. In these scenarios, proteins are evolving to
evade pathogen antagonists at multiple sites throughout the protein. Although pathogens are
known to antagonize the caspase-1 signaling pathway both upstream[24,25] and
downstream[26] of caspase-1, we would not expect these known, indirect antagonists to drive
adaptation in caspase-1 itself. Several viruses encode multi-caspase inhibitors that can target
caspase-1, -4, and -5 by forming covalent bonds at the caspase active site [27,28]. However,
as we did not observe an enrichment of positive selection around the caspase-1 active site (Fig.
4D-G), and as the inhibitors are not caspase-1-specific, we also do not believe that these
inhibitors fully explain the patterns of caspase-1 evolution. Instead, we postulate that there are
multiple, unknown pathogen proteins that specifically bind and inhibit caspase-1 at different
molecular interfaces. We further predict that the residues we highlighted here are likely to
mediate variation in binding specificity between caspase-1 and various antagonists as well as
between a given antagonist and different members of the caspase-1 family.
447
Methods

Selection of species and genomes to analyze
To compare the evolutionary histories of primate and rodent caspases, our analyses required at
least 10-15 caspase genomic loci from both the rodent and primate clades[42]. We selected
genomes to use in 2022 and therefore included the latest assembly version available at the time
(Sup Table 1). High quality genome assemblies were included based on the following quality
metrics: low assembly fragmentation (N50>300kbp and L50<2000), and high read coverage
(>40X). Rodent and primate species trees were based on previously published work [42–46]
and the Zoonomia database. To approximate divergence of the species, we used CODEML
model 0 [35] to calculate the pairwise dS between the *CASP1* homologs in the species of
interest and either humans (for primates) or house mouse (for rodents). Species that had a
pairwise dS >0.4 from the reference were considered to be outgroups.
460
Extracting the caspase-1 locus from the assemblies
To identify the *CASP1* locus and *CASP1*-related homologs in each of our genome assemblies,
we first created local blast databases of the collected, high-quality genomes for rodents and
primates using BLAST+ v2.13.0 [47] makeblastbd. We then used the Refseq representative
transcript of house mouse or human *CASP1* as a query in a blastn search on our local
databases with default settings. From the output of possible *CASP1* hits in each genome, we

filtered for strong exon hits using a bitscore>100. The coordinates of the hits also revealed that
each *CASP1*-like gene was contained within a single genomic locus in each species, and none
were located on different chromosomes. We then identified start and end coordinates of each
candidate *CASP1* homolog in each assembly, based on homology to the beginning and end of
the reference query. Using the coordinates of the strongest *CASP1* hit, we then extracted the
CASP1 genomic locus with excess flanking two Mbp for each assembly.

Exon annotations, alignments, phylogenetic trees, and caspase paralog identification
To retrieve accurate *CASP* CDS sequences from the *CASP1* locus in each species, we first
annotated each exon. For each of the *CASP1* homologs and *CARD*-only proteins, we chose the
Refseq representative transcript from human or house mouse as our reference CDS for exon
annotation. In humans *CASP12* is considered catalytically inactive and likely a
pseudogene[12,15]. Because pseudogenes are undergoing degeneration of their coding
sequence, annotation of conserved exons from a pseudogenic reference is challenging.
Therefore, we attempted to perform exon annotation of primate *CASP12* loci using both the
human and house mouse *CASP12* CDS sequences.

We searched for conserved exons of each homolog using both nucleotide similarity and
synteny. To identify exons with high nucleotide identity to the reference (>80% identity to
humans in primates, or >60% identity to house mouse in rodents), we used the Geneious Prime
v2024.0.3 "Transfer Annotations" tool to transfer exon annotations from the reference sequence
to the locus of interest. While this approach identified most exons of interest, some diverged
exons were missed, whereas exons similar to multiple caspase paralogs matched to multiple
regions within the same locus. We therefore relied on gene synteny to assign exons to the
correct *CASP* gene and identify paralogs, even if only part of the gene was present.

To identify missing exons and obtain full CDS sequences for each gene from each species, we
aligned the coding sequences of each gene with the annotated genomic DNA using MAFFT[48]
with automatic direction detection. We then used the exon annotations to guide the manual
trimming of introns across the alignment. We then realigned the trimmed sequences by
translation using Geneious Prime translation alignment tool with the MUSCLEv5[49] plugin. We
confirmed the quality of our manual trimming by comparing our trimmed reference species
CDSs to the publicly available reference CDSs and protein sequences. In some cases, like the
Jerboa *CASP11* and *CASP12*, our manual trimming method initially failed because the intronic
regions were very different sizes compared to the rest of rodent homologs. However, using
predicted CDS of the Jerboa caspases and our exon annotation, we were able to recover full-
length inframe CDS of these genes through manually trimming, independent of the gDNA
alignment. For primate *CASP5*, Old world monkeys had an insertion within the N-terminal tail of
the protein, which we manually trimmed out before further analyses. We generated all
phylogenetic trees using the Geneious Prime tree builder PhyML[50] plugin with GTR model
and aLRT statistics.

Pseudogene and gene loss calls

While *CASP1* was well-conserved in all lineages, other genes in the locus were not. To find
predicted or quantified transcript evidence of each caspase homolog, we performed an NCBI
blast search in Refseq representative genomes with each caspase homolog CDS reference.
The presence of a full-length in-frame transcript from blast results, identical in sequence to our
extracted CDS, was additional evidence of a functional full-length *CASP* homolog. If there was
not a full-length in-frame transcript but an alternative splice isoform from blast results was still
in-frame, we called these homologs as "partial in-frame transcripts". However, this CDS search
strategy was inconclusive if *CASP* homologs did not have any transcript evidence at all, leaving
open the possibilities of gene loss, pseudogenization, or lack of genome annotation.
519
We called events as "gene losses" if there were fewer than 70% exons were identified as
described above. These partial loci were not included in subsequent gene alignments or trees.
To define pseudogenes, we examined the sequences for frameshift indel mutations, exon
truncations, or exon loss. Frameshift indel mutations were differentiated from genome assembly
errors by examining closely related species to see if the same defect was present in multiple,
independent studies/assemblies. See Sup Table 2 for a full description of the presence,
absence, and pseudogene data for each gene in each species.
527
Identification of regions of nucleotide identity within the caspase locus
We identified regions of extended nucleotide identity consistent with recent gene conversion in
the caspase locus of humans (GCF_000001405.40) and *M. musculus* (GCA_000001635.9). To
do so, we extracted a ~350kb region from each genome, centered on the *CASP1* gene. We
then used the Dotplot tool of Geneious Prime to detect any regions of self-similarity, which
appeared as off-diagonal lines in the dotplot. We annotated any DNA regions that were at least
1kb long that contained at least 50% nucleotide identity to another region within the 350kb
locus.
536
Measurements of selection and recombination
To perform selection analyses, we selected *CASP* sequences that fit within at an appropriate
level of divergence. 15 primate assemblies were selected for selection analyses with the most
diverged species from human being marmoset at a dS of 0.13. The rodent Muridaea clade of 12
species was selected for selection analyses with the most diverged clade from house mouse
being spiny mouse at a dS of 0.29. To detect recombination breakpoints, we ran GARD[39] on
each homolog nucleotide alignment with default DataMonkey settings: faster run mode,
universal genetic code, no site-to-site rate variation, two rate classes. Because recombination
can cause false positives in selection analyses, we then split the alignments at recombination
breakpoints and analyzed each recombination segment independently. To detect residues of
positive selection, we ran two softwares: PAML CODEML[35] and FUBAR[40]. We performed
PAML CODEML analysis with models 0, 0a, 1, 2, 7, 8, and 8a. Model 0 estimates a single
dN/dS value across the entire gene. Model 1 partitions sites into two dN/dS estimates, dN/dS =1
or dN/dS<1, whereas model 2 allows for a third bin, dN/dS>1; statistical tests compare models 1
& 2 to assess if there are sites with dN/dS significantly greater than 1. Similarly, model 7
includes many dN/dS bins<1, model 8a includes a bin = 1, and model 8 includes a bin with
dN/dS>1. P-values were obtained by comparing the lnL values between M1 v. 2, M7 v. 8, or

M8a v. 8. We performed FUBAR using the default universal genetic code. Residues of positive
selection were defined as posterior probability >0.95 (CODEML) or posterior probability >0.9
(FUBAR). See Sup Table 3 and Sup Data 1 for more detailed CODEML and FUBAR statistics
and results.
558
559
560 **Figure legends**
561
Figure 1. Differential patterns of gene gain and loss within the caspase-1 locus. A) Genes
in the caspase-1 locus in human and mouse (not to scale). B) PhyML phylogenetic tree of intact
caspase genes demonstrates that orthologs have been correctly named and categorized,
despite gene losses. Numbers at nodes are aLRT support values. C-D) Gene presence/
absence across primate (C) and rodent (D) species, with colored Xs showing the history of gene
losses. See text for definitions of pseudogene vs. partial sequence vs. gene absence. *CASP1*
has been retained across all species examined, as has primate *CASP4*. In contrast, there have
been multiple cases of gene degradation and loss of rodent *CASP11*, primate *CASP5*, and the
primate *CARD* genes. *CASP12* appears to be retained and functional across most (but not all)
rodents, whereas it has been largely degraded and lost in primates.
572
Figure 2. The caspase-1 locus has undergone recent gene conversions and tandem
duplications in primates, but not rodents. A-C) Annotated genes (filled colored arrows) and
pseudogenes (white arrows with colored outlines) in the human (A-B) and mouse (C) loci.
Colored arcs (A) connect regions of extended nucleotide identity between caspase genes or
pseudogenes, indicative of recent tandem gene duplications or gene conversions. Gray arcs (B
and C) show regions of sequence identity within intronic or intergenic regions.
579
Figure 3. PhyML phylogenetic tree of the CARDs of each primate caspase-1 subfamily
gene. Region within the dashed box in A) is shown in detail in panel B), with gene identity
indicated by symbols. Although sequences for CASP4, 5, and CARD18 have distinct branches,
in several species the CASP1, CARD16, and/or CARD17 sequences are highly similar and
intermixed on the tree, indicating recent gene conversion. All illustrated nodes on the PhyML
trees had aLRT support statistics >5. Other nodes were collapsed.
586
Figure 4. Positively selected residues in pro-inflammatory caspases in primates and
rodents. A-E) Sites of positive selection, as determined by the FUBAR and CODEML
algorithms. Arrowheads mark sites passing the statistical threshold for positive selection.
Vertical dotted lines separate recombination segments identified by GARD, which were each
analyzed independently. The domain architecture of each caspase is shown as colored boxes,
with active sites illustrated (yellow *) along with the proposed LPS-binding residues of CASP4
and 11 according to Shi et. al 2014 (orange carats). In four cases, sites of positive selection in
primate CASP1 were homologous to sites identified in one or more other caspase alignment
(green, blue, red, or purple residue names). All coordinates are named according to the
residues in the human or mouse sequences for primate or rodent analyses, respectively. F-G)
Each identified site of CASP1 positive selection (red) is illustrated on the predicted AlphaFold

structure of human (F) or mouse (G) caspase-1. The CARD, p20, and p10 domains are colored
as in (A-E), with linker regions in gray, and active sites in yellow. Many more sites of positive
selection were identified in primates than in rodents, but in neither clade do the sites show much
co-localization in 3D space.
602
Figure 5. Caspase-1 aspartate cleavage sites between CARD and p20 domains are not
strictly conserved. A) domain architecture of caspase-1, showing locations of aspartate
cleavage sites ('D'). B-C) Alignments of the caspase-1 aspartate cleavage sites, with residues
that do not match consensus highlighted with colored boxes. While the two C-terminal cleavage
sites are highly conserved within primates (B) and rodents (C), the N-terminal site is more highly
variable and some species lack the aspartate residue entirely. Arrowheads mark nearby
positively selected sites.
610
611
612 ## **Supplemental files**
613
614 **Supplemental Table 1.** Genome statistics and sources for all primate and rodent species used.
615
Supplemental Table 2. Gene presence, absence, and pseudogene calls in each species.
The "Gene Presence" tab shows for each gene in each primate and rodent species whether we
recovered a sequence from the caspase-1 locus that aligned well with orthologs ("align") and
whether we recovered a blast hit to a sequenced or predicted transcript. 1 = present; 0 =
pseudogene; -1 = absent; * indicates genes with partial in-frame sequences.
The "Pseudogene Evidence" tab lists the SNPs, deletions, and frameshifts in each gene that led
us to call it a pseudogene.
623
Supplemental Table 3. Summary of selection analyses.
The "Species per Gene Analysis" tab indicates which species were included in each alignment
analyzed. The "GARD breakpoints" tab lists the N-terminal and C-terminal end of each GARD
breakpoint, according to the human or house mouse amino acid coordinates. The "PAML Model
Comparisons" tab lists the dN/dS and log likelihood values, and p-values obtained for each
model and statistical test for each caspase recombination segment. M0 vs. M0a tests for dN/dS
not equal to 1 across the whole gene (i.e. deviation from drift). The other comparisons (M1 vs.
M2, M7 vs. M8, M8a vs. M8) are each tests for positive selection, i.e. dN/dS>1 with alternative
statistical models. All p-values are compared to a Bonferroni corrected value of 3.33E-03. The
"FUBAR Summary" tab lists the number of positively selected sites and the highest Bayes factor
per segment.
635
636 **Supplemental Data 1.** Alignments and full results of GARD, CODEML, and FUBAR analyses.
637 We provide the alignments and raw data to maximize reproducibility and data transparency.

References

1. Enard D, Cai L, Gwennap C, Petrov DA. Viruses are a dominant driver of protein adaptation in mammals. Elife. 2016;5. doi:10.7554/eLife.12469

2. Daugherty MD, Malik HS. Rules of engagement: molecular insights from host-virus arms races. Annu Rev Genet. 2012;46: 677–700.

3. McLaughlin RN Jr, Malik HS. Genetic conflicts: the usual suspects and beyond. J Exp Biol. 2017;220: 6–17.

4. Daugherty MD, Zanders SE. Gene conversion generates evolutionary novelty that fuels genetic conflicts. Curr Opin Genet Dev. 2019;58-59: 49–54.

5. Zhu Q, Zheng M, Balakrishnan A, Karki R, Kanneganti T-D. Gasdermin D Promotes AIM2 Inflammasome Activation and Is Required for Host Protection against Francisella novicida. J Immunol. 2018;201: 3662–3668.

6. Wu J, Cai J, Tang Y, Lu B. The noncanonical inflammasome-induced pyroptosis and septic shock. Semin Immunol. 2023;70: 101844.

7. Xu Z, Kombe Kombe AJ, Deng S, Zhang H, Wu S, Ruan J, et al. NLRP inflammasomes in health and disease. Mol Biomed. 2024;5: 14.

8. Tenthorey JL, Kofoed EM, Daugherty MD, Malik HS, Vance RE. Molecular basis for specific recognition of bacterial ligands by NAIP/NLRC4 inflammasomes. Mol Cell. 2014;54: 17–29.

9. Tsu BV, Beierschmitt C, Ryan AP, Agarwal R, Mitchell PS, Daugherty MD. Diverse viral proteases activate the NLRP1 inflammasome. Elife. 2021;10. doi:10.7554/eLife.60609

10. Tsu BV, Agarwal R, Gokhale NS, Kulsuptrakul J, Ryan AP, Fay EJ, et al. Host-specific sensing of coronaviruses and picornaviruses by the CARD8 inflammasome. PLoS Biol. 2023;21: e3002144.

11. Sundaram B, Tweedell RE, Prasanth Kumar S, Kanneganti T-D. The NLR family of innate immune and cell death sensors. Immunity. 2024;57: 674–699.

12. Eckhart L, Ballaun C, Hermann M, VandeBerg JL, Sipos W, Uthman A, et al. Identification of novel mammalian caspases reveals an important role of gene loss in shaping the human caspase repertoire. Mol Biol Evol. 2008;25: 831–841.

13. Shi J, Zhao Y, Wang Y, Gao W, Ding J, Li P, et al. Inflammatory caspases are innate immune receptors for intracellular LPS. Nature. 2014;514: 187–192.

14. Sahoo G, Samal D, Khandayataray P, Murthy MK. A Review on Caspases: Key Regulators of Biological Activities and Apoptosis. Mol Neurobiol. 2023;60: 5805–5837.

15. Fischer H, Koenig U, Eckhart L, Tschachler E. Human caspase 12 has acquired deleterious mutations. Biochem Biophys Res Commun. 2002;293: 722–726.

16. Kachapati K, O'Brien TR, Bergeron J, Zhang M, Dean M. Population distribution of the functional caspase-12 allele. Hum Mutat. 2006;27: 975.

17. Xue Y, Daly A, Yngvadottir B, Liu M, Coop G, Kim Y, et al. Spread of an inactive form of
caspase-12 in humans is due to recent positive selection. Am J Hum Genet. 2006;78: 659–
670.

18. Druilhe A, Srinivasula SM, Razmara M, Ahmad M, Alnemri ES. Regulation of IL-1β
generation by Pseudo-ICE and ICEBERG, two dominant negative caspase recruitment
domain proteins. Cell Death Differ. 2001;8: 649–657.

19. Lamkanfi M, Denecker G, Kalai M, D'hondt K, Meeus A, Declercq W, et al. INCA, a novel
human caspase recruitment domain protein that inhibits interleukin-1beta generation. J Biol
Chem. 2004;279: 51729–51738.

20. Indramohan M, Stehlik C, Dorfleutner A. COPs and POPs Patrol Inflammasome Activation.
J Mol Biol. 2018;430: 153–173.

21. Karasawa T, Kawashima A, Usui F, Kimura H, Shirasuna K, Inoue Y, et al. Oligomerized
CARD16 promotes caspase-1 assembly and IL-1β processing. FEBS Open Bio. 2015;5:
348–356.

22. Lu A, Li Y, Schmidt FI, Yin Q, Chen S, Fu T-M, et al. Molecular basis of caspase-1
polymerization and its inhibition by a new capping mechanism. Nat Struct Mol Biol.
2016;23: 416–425.

23. Devi S, Indramohan M, Jäger E, Carriere J, Chu LH, de Almeida L, et al. CARD-only
proteins regulate in vivo inflammasome responses and ameliorate gout. Cell Rep. 2023;42:
112265.

24. Brodsky IE, Palm NW, Sadanand S, Ryndak MB, Sutterwala FS, Flavell RA, et al. A
Yersinia effector protein promotes virulence by preventing inflammasome recognition of the
type III secretion system. Cell Host Microbe. 2010;7: 376–387.

25. Chung LK, Park YH, Zheng Y, Brodsky IE, Hearing P, Kastner DL, et al. The Yersinia
Virulence Factor YopM Hijacks Host Kinases to Inhibit Type III Effector-Triggered Activation
of the Pyrin Inflammasome. Cell Host Microbe. 2016;20: 296–306.

26. Luchetti G, Roncaioli JL, Chavez RA, Schubert AF, Kofoed EM, Reja R, et al. Shigella
ubiquitin ligase IpaH7.8 targets gasdermin D for degradation to prevent pyroptosis and
enable infection. Cell Host Microbe. 2021;29: 1521–1530.e10.

27. Best SM. Viral subversion of apoptotic enzymes: escape from death row. Annu Rev
Microbiol. 2008;62: 171–192.

28. Kamada S, Funahashi Y, Tsujimoto Y. Caspase-4 and caspase-5, members of the
ICE/CED-3 family of cysteine proteases, are CrmA-inhibitable proteases. Cell Death Differ.
1997;4: 473–478.

29. Li Z, Liu W, Fu J, Cheng S, Xu Y, Wang Z, et al. Shigella evades pyroptosis by arginine
ADP-riboxanation of caspase-11. Nature. 2021;599: 290–295.

30. Castro LK, Daugherty MD. Tripping the wire: sensing of viral protease activity by CARD8
and NLRP1 inflammasomes. Curr Opin Immunol. 2023;83: 102354.

31. Lamkanfi M, Declercq W, Kalai M, Saelens X, Vandenabeele P. Alice in caspase land. A

phylogenetic analysis of caspases from worm to man. Cell Death Differ. 2002;9: 358–361.

32. Devant P, Cao A, Kagan JC. Evolution-inspired redesign of the LPS receptor caspase-4
into an interleukin-1β converting enzyme. Sci Immunol. 2021;6.
doi:10.1126/sciimmunol.abh3567

33. Fairley S, Lowy-Gallego E, Perry E, Flicek P. The International Genome Sample Resource
(IGSR) collection of open human genomic variation resources. Nucleic Acids Res. 2020;48:
D941–D947.

34. Vollger MR, Guitart X, Dishuck PC, Mercuri L, Harvey WT, Gershman A, et al. Segmental
duplications and their variation in a complete human genome. Science. 2022;376:
eabj6965.

35. Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol. 2007;24:
1586–1591.

36. Kumar S, Stecher G, Suleski M, Hedges SB. TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. Mol Biol Evol. 2017;34: 1812–1819.

37. Jurka J. Repbase update: a database and an electronic journal of repetitive elements.
Trends Genet. 2000;16: 418–420.

38. Kobayashi T, Ogawa M, Sanada T, Mimuro H, Kim M, Ashida H, et al. The Shigella OspC3
effector inhibits caspase-4, antagonizes inflammatory cell death, and promotes epithelial
infection. Cell Host Microbe. 2013;13: 570–583.

39. Kosakovsky Pond SL, Posada D, Gravenor MB, Woelk CH, Frost SDW. GARD: a genetic
algorithm for recombination detection. Bioinformatics. 2006;22: 3096–3098.

40. Murrell B, Moola S, Mabona A, Weighill T, Sheward D, Kosakovsky Pond SL, et al. FUBAR:
a fast, unconstrained bayesian approximation for inferring selection. Mol Biol Evol. 2013;30:
1196–1205.

41. Makoni NJ, Nichols MR. The intricate biophysical puzzle of caspase-1 activation. Arch
Biochem Biophys. 2021;699: 108753.

42. McBee RM, Rozmiarek SA, Meyerson NR, Rowley PA, Sawyer SL. The effect of species
representation on the detection of positive selection in primate gene data sets. Mol Biol
Evol. 2015;32: 1091–1096.

43. Menezes AN, Bonvicino CR, Seuánez HN. Identification, classification and evolution of owl
monkeys (Aotus, Illiger 1811). BMC Evol Biol. 2010;10: 248.

44. Molaro A, Malik HS, Bourc'his D. Dynamic Evolution of De Novo DNA Methyltransferases in
Rodent and Primate Genomes. Mol Biol Evol. 2020;37: 1882–1892.

45. Swanson MT, Oliveros CH, Esselstyn JA. A phylogenomic rodent tree reveals the repeated
evolution of masseter architectures. Proc Biol Sci. 2019;286: 20190672.

46. Côrte-Real JV, Baldauf H-M, Melo-Ferreira J, Abrantes J, Esteves PJ. Evolution of
Guanylate Binding Protein (GBP) Genes in Muroid Rodents (Muridae and Cricetidae)
Reveals an Outstanding Pattern of Gain and Loss. Front Immunol. 2022;13: 752186.

47. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
architecture and applications. BMC Bioinformatics. 2009;10: 421.

48. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30: 772–780.

49. Edgar RC. MUSCLE v5 enables improved estimates of phylogenetic tree confidence by
ensemble bootstrapping. bioRxiv. 2021. p. 2021.06.20.449169.
doi:10.1101/2021.06.20.449169

50. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New Algorithms
and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of
PhyML 3.0. Syst Biol. 2010;59: 307–321.