Title

Olfactory combinatorial coding supports risk-reward decision making in *C. elegans*.

Authors

Md Zubayer Hossain Saad[1], William G. Ryan V[2], Chelyan A. Edwards[1], Benjamin N. Szymanski[1], Aryan R. Marri[1], Lilian G. Jerow[1, 3], Robert McCullumsmith[2], and Bruce A. Bamber[1]* (Bruce.Bamber@utoledo.edu)

Affiliations

[1]Department of Biological Sciences, The University of Toledo
[2]Department of Neurosciences and Psychiatry, The University of Toledo
[3]Present address: Neuroscience Graduate Program, University of Cincinnati
*Corresponding Author

Abstract

Olfactory-driven behaviors are essential for animal survival, but mechanisms for decoding olfactory inputs remain poorly understood. We have used whole-network Ca^{++} imaging to study olfactory coding in *Caenorhabditis elegans*. We show that the odorant 1-octanol is encoded combinatorially in the periphery as both an attractant and a repellant. These inputs are integrated centrally, and their relative strengths determine the sensitivity and valence of the behavioral response through modulation of locomotory reversals and speed. The balance of these pathways also dictates the activity of the locomotory command interneurons, which control locomotory reversals. This balance serves as a regulatory node for response modulation, allowing *C. elegans* to weigh opportunities and hazards in its environment when formulating behavioral responses. Thus, an odorant can be encoded simultaneously as inputs of opposite valence, focusing attention on the integration of these inputs in determining perception, response, and plasticity.

Introduction

Animals must detect and respond to chemical signals in the environment in order to find food, avoid hazards, and find mates. Olfactory systems balance two competing priorities to optimally represent the external chemical space. First, they must respond rapidly and robustly to specific compounds which elicit innate responses important for survival. These include predator or parasite odors (eliciting avoidance), and pheromones (eliciting social and/or reproductive responses) (*1-4*). Such stimuli are sensed using narrowly-tuned odorant receptors in the periphery, hard-wired to dedicated circuits for specific behaviors (labeled line coding). Second, olfactory systems must be versatile, capable of detecting and discriminating millions of different environmental odorants. This versatility is achieved using broadly-tuned odorant receptors, which activate in different combinations depending on the chemical structure of the odorant, enabling millions of different odorants to be discriminated using, at most, a few thousand receptors (*5*). These combinations are decoded centrally to identify the odor (combinatorial coding)(*4*). Combinatorial coding has been regarded as the predominant mechanism for olfactory coding (*6, 7*), but examples of labeled line-like olfactory coding are increasingly being found (*8-12*), suggesting that olfactory coding is a hybrid of labeled line and combinatorial mechanisms (*1*). Labeled line coding is simpler, and probably evolved first, with animals responding to specific environmental cues using specific receptors. Combinatorial coding likely emerged later as the narrowly-tuned receptor genes underwent duplication and divergence to diversify ligand binding properties. Presumably, the central decoding circuitry then evolved to take advantage of these new sources of environmental information (*1, 13*). The initial steps in

olfactory coding are relatively well studied, with each olfactory receptor neuron expressing 1-2 olfactory receptors (*14, 15*). Inputs from olfactory receptor neurons expressing the same receptor converge on a single glomerulus in the olfactory bulb (mammals) or antennal lobe (insects), and the glomeruli project to higher processing centers in the brain (*16*). Recent studies are beginning to unravel the neural circuitry by which olfactory inputs generate perception and responses (*17-20*). However, we still lack a clear understanding of how combinatorial inputs are decoded, and how combinatorial pathways and labeled line pathways may functionally interact.

We have turned to *C. elegans* to study the downstream processing of olfactory inputs. The small size of the *C. elegans* nervous system and the availability of techniques for pan-neuronal activity recording now make it possible to study the neural basis for sensory-driven behaviors at whole-network scale in this model (*21-31*). Olfaction in *C. elegans* has classically been viewed as a labeled-line coding system rather than a combinatorial system, where specific subsets of chemosensory neurons respond to attractive odorants and drive positive chemotaxis (AWC, AWA), or respond to repulsive odorants and drive negative chemotaxis (ASH, AWB, ADL) (*32-38*). However, it was recently shown that individual odorants activate multiple classes of olfactory neurons, and individual olfactory neurons respond to a wide range of odorants (*39*). This responsiveness profile is the hallmark of combinatorial coding, even though there are many fewer chemosensory neurons in *C. elegans*, and these neurons must express a much wider repertoire of olfactory receptors than their mammalian counterparts. The striking parallels in olfactory coding between animals as divergent as nematodes, insects, and mammals suggests that evolution may have converged on a few optimal solutions for sensing and responding to external odor landscapes.

We have performed an in-depth analysis of a single odorant, 1-octanol (1-oct), which is encoded combinatorially (*39*), and is well-known to stimulate aversive behavior (*32, 33, 38, 40*). We show that 1-oct stimulation concurrently activates two separate and opposing sensory afferent pathways in the nervous system. These pathways converge to control at least four aspects of locomotion at the behavioral and circuit levels, with the balance between attraction and repulsion emerging as the critical factor determining the overall behavioral output. Furthermore, this coding strategy provides a simple framework to modulate sensory-driven responses to produce context-appropriate behavior.

Results
1-octanol repulsion in microfluidic arenas
1-oct avoidance in *C. elegans* has been intensively studied because it is strongly modulated by nutrition state and monoamine/neuropeptide signaling cascades (*34, 41-45*). Our goal was to investigate the network basis of 1-oct repulsion using the NeuroPAL system (*30*), which allows simultaneous whole-brain Ca^{++} imaging with unambiguous neuronal identification, coupled with quantitative analysis of 1-oct aversive behavior. 1-oct has previously been utilized as a volatile stimulus (*33, 34*), or in a gradient on agar plates (*46, 47*), while NeuroPAL recordings are performed in aqueous solution in microfluidics chips under confocal microscopy (*30*). To better match the conditions for NeuroPAL recordings and behavioral assays, we first investigated 1-oct repulsion in microfluidics arenas (*48-50*), which gave us precise control over the concentration and spatio-temporal characteristics of the 1-oct stimulus. Using a simple two-chamber arena (Fig. 1A), we demonstrated that worms were repelled by a saturating concentration of 1-oct in S-basal (2.2 mM, corresponding to a 3.75×10^{-4} dilution; Fig. 1B-D). In our initial

trials, we added 0.01% w/v xylene cyanol (0.1 mg/mL) to the 1-oct solution to visualize the interface between the stimulus and control buffer. However, as another study reported (*50*), worms were significantly attracted to xylene cyanol introducing an unwanted stimulus into the assays. We therefore optimized methods to maintain a stable interface between odorant and buffer chambers without the need for dye (see Methods). We assayed worms between 20 and 40 minutes after removal from food, during area-restricted search behavior (*51*), and calculated average chemotaxis indices (CIs) between 30 and 40 minutes (see Methods, and Fig. 1D, E). The -0.7 CI observed in the experiment shown represents strong repulsion.

Network-wide response to repulsive olfactory stimulation
To investigate the neuronal basis for the observed 1-oct aversive behavior, we performed network-wide Ca^{++} imaging during 1-oct stimulation, and identified the responding cells by imaging the neuronal landmarks encoded by the NeuroPAL fluorescent reporters (Fig. 2A). Worms were immobilized in 'olfactory' microfluidics chips (*52*) for microscopy. Previous whole-brain studies have shown that certain neurons undergo spontaneous transitions between low and high activity states correlating with spontaneous reversals during foraging, notably the reverse command neurons AVA, AVE, AVD, AVB, and the RIM neuron (*23, 27, 53-58*). We therefore used a recording/stimulus protocol consisting of a 6 min pre-stimulus period, followed by 6 30s applications of 2.2 mM 1-oct, spaced 30s apart, to help distinguish between spontaneous transitions and potential sensory-evoked transitions within this neuronal population. We observed dynamic activity patterns in dozens of neurons before and during 1-oct stimulus (Fig. 2B-D). Consistent with previous studies, AVA, along with AVE, AVD, RIM, AIB, and AIZ showed spontaneous state transitions, with the AVB forward command interneuron and the AIY interneuron showing mirror image patterns (i.e. AVB/AIY inactivated when AVA activated and vice versa). Typically, 3-4 transitions were observed in a 12 minute recording. These transitions took place during both the pre-stimulus and stimulus periods (Fig. 2D, 3A), and showed no obvious relationship with 1-oct application or withdrawal.

Sensory signals were also prominent, in both amphid and non-amphid chemosensory neurons. ASH and BAG showed strong consistent responses correlating with stimulus onset ('ON' responses). AWB also showed an ON response but also had high levels of spontaneous activity. ADL showed a strong ON response to the initial stimulus but was strongly desensitized in subsequent applications (Fig. 2D and 3B). ASH, AWB, and ADL were previously identified as the main neurons mediating 1-oct aversive responses (*33, 34*). 1-oct responses were also observed in AWA, ASE, ASK, ASJ, and URX (Fig. 2D, 3B), consistent with the observation of Lin et al, showing that 1-oct activates sensory neurons promiscuously (*39*). Significantly, we observed a strong 1-oct response in AWC neurons (both AWC^OFF and AWC^ON; Fig 2-D, 3B). AWC activity decreased upon 1-oct addition and rebounded upon 1-oct withdrawal, in a typical 'OFF' neuron pattern (*59*), opposite to ASH (Fig. 2B-D, 3C). This was surprising, first, because AWC is usually considered an attractive neuron, not expected to respond to a repulsive odorant, and second, because the observed AWC pattern predicts that AWC should drive attraction to 1-oct (*38, 46, 59, 60*). 1-oct has thus far only been described as a repellent for *C. elegans*, with no attractive properties (*32, 34, 38, 47*). Strong sensory-driven activity was also observed in AIA and SAA interneurons, suggesting these two neurons may receive substantial inputs from the sensory neurons (Fig. 2D, 3C). Apart from AIA, the other 'first layer' interneurons AIB, AIY, and AIZ did not show detectable sensory-driven activity (Fig. 3A), despite their extensive anatomical connections with sensory neurons, suggesting

these interneurons may be receiving their principal functional inputs from elsewhere in the circuit.

To better understand how sensory inputs may pattern global network activity patterns, we performed correlation analysis on selected data sets in which we were able to identify 12 key neurons with well characterized roles in sensory motor coupling (*38, 51, 59*). To detect the sensory signature, we compared neuronal correlation patterns in the first 6 minute interval (prior to stimulation), and the second 6 minute interval (where stimulus was applied and removed for alternating 30s periods) (Fig. 3D, E). We observed the expected increases and decreases in correlation among the sensory neurons (i.e. ASH and BAG became correlated, and anti-correlated with AWC). We did not observe any increased correlation between the motor command interneurons and sensory neurons, consistent with the absence of a consistent relationship between state transitions and 1-oct application. However, the degree of activity cross-correlation among AVA, AVE, AVD, AIB, and RIM was increased during the stimulus period, and AIB also showed increased correlation with ASH. These observations suggest that sensory information was penetrating to the locomotory command circuit level even if there was no obvious sensory entrainment of the principal neurons in those circuits (Fig. 3D, E).

Finally, we observed several neurons with activity patterns which did not correspond to the locomotory command group or the sensory-dependent group, which included SMDV, RIA, and RMDD (Fig. 2D), which are generally associated with turning behavior (*61-63*).

The balance of simultaneous repulsion and attraction shapes the chemotactic response

The observation that 1-oct application and withdrawal cycles activate ASH and AWC in alternation predicts that these two neurons generate antagonistic inputs into the locomotory circuitry. Specifically, ASH activity varies in phase with 1-oct application, predicted to stimulate in-phase reversal consistent with a repulsive chemotactic response. AWC activity varies out of phase with 1-oct application, predicted to inhibit in-phase reversals and stimulate out of phase reversals, consistent with an attractive chemotactic response. This puzzling activity pattern led us to hypothesize that 1-oct is encoded combinatorially, simultaneously as a repellant (by ASH) and an attractant (by AWC), and these two afferent pathways balance one another during the application-withdrawal cycle, and mutually antagonizing sensory entrainment of the reverse command neurons and potentially attenuating 1-oct behavioral responses.

To test this hypothesis, we selectively inhibited repulsive inputs to unmask latent effects of the attractive inputs and vice versa. To manipulate repulsive inputs, we reduced the 1-oct concentration. Recently, Lin et. al. (2023) reported 1-oct concentration-response relationships for ASH and AWC. At sub-saturating concentrations, ASH responses were highly concentration dependent while AWC 'ON' responses were relatively insensitive to concentration. First, we repeated the observations of Lin et al., extending the concentration range to saturation (2.2mM, shown to be repellent in microfluidics-based assays; Fig. 1), recording ASH responses, and both the AWC 'ON' and 'OFF' responses. We serially stimulated worms with 0.22 mM, 0.66 mM, and 2.2 mM 1-oct. ASH responses showed significant concentration-dependent increases, whereas AWC 'ON' and 'OFF' responses did not (Fig. 4A-C). Behaviorally, after 3- and 10-fold dilution, the 1-oct response switched from repulsion to attraction (Fig. 4D), while 1.4 to 2.0-fold dilutions attenuated chemotactic responses (Fig. 4D), potentially reflecting a more even balance

between the two signaling pathways. This result suggests that ASH-mediated repulsive inputs dominate AWC-mediated attractive inputs at saturating concentrations, causing overall repulsion, but as ASH input strength decreases relative to AWC in progressively more dilute 1-oct, AWC attractive inputs come to dominate, causing attraction.

We then used a genetic approach to manipulate attractive inputs. Sensory signal transduction in AWC but not ASH is eliminated in *tax-4* mutants, which lack a cGMP-gated ion channel downstream of sensory receptors (*38, 64-66*). Loss of TAX-4 converted 1-oct back to a repellent at the lower concentrations (Fig. 4C), indicating that TAX-4 signaling is clearly contributing to 1-oct chemotactic responses in this context. Similarly, at saturating concentrations, we observed a trend toward increased repulsion, although this observation did not rise to statistical significance because of high variability in the CI values. Overall, these experiments support the idea that 1-oct is encoded as multiple parallel inputs with opposite valence, operating concurrently. The ASH inputs promote repulsion while the AWC inputs promote attraction, and the balance of their relative strengths dictates the final chemotactic outcome (Fig. 4E).

To understand the interaction of these co-activated afferent pathways at the level of locomotory decision making, we closely examined the behavior of worms as they encountered the interface between 1-oct solution and control buffer in microfluidic devices. We began using the microfluidics 'stripe' chip (*49, 50*), containing a central zone of 2.2 mM 1-oct flanked by zones containing control buffer. We used automated tracking (*67*) to quantify behavior upon encountering zone boundaries (i.e. reversals vs successful transits; Fig. 5A, B). Surprisingly, for worms entering the 1-oct zone, entry transits outnumbered entry reversals by nearly 2:1 (Fig. 5B, C). In *osm-9* mutants, in which the 1-oct responses of ASH are defective (*43, 68, 69*), the ratio was skewed significantly toward entry transits, demonstrating that OSM-9 signaling promotes 1-oct avoidance. Interestingly, in *tax-4* mutants, the ratio was strongly skewed toward entry reversals. This result indicates that *tax-4* signaling actively suppresses 1-oct avoidance, consistent with its role in promoting 1-oct attraction (Fig. 5B, C). This attraction-promoting role was also observed for worms exiting the 1-oct zone (Fig. 5B, D). Here, exit reversals strongly outnumber exit transits. These reversals are dependent upon TAX-4, but do not require OSM-9 (Fig. 5D). These observations confirm that TAX-4 mediated attractive signaling significantly shapes the behavioral response to 1-oct, despite the fact that 1-oct is classified as a strong repellent. Most importantly, it highlights how overall chemotactic behavior is reciprocally regulated by two different locomotory transitions (reversals due to increased d[1-oct]/dt and reversals due to decreased d[1-oct]/dt), and how two opposing sensory afferent pathways converge to regulate a single locomotory transition, namely reversals due to increased d[1-oct]/dt, at a single point in time and space (Fig. 5B).

These behavioral data raised a new important question: how can wild-type worms avoid the 1-oct containing zone when their reversal decisions favor entry and retention? We had noticed that contact with 2.2 mM 1-oct caused worms to rapidly and reversibly increase their locomotory activity, suggesting an additional potential mechanism in play. To quantify this effect, we measured the average distance worms moved when we flooded the microfluidics arena with 2.2 mM 1-oct, comparing the distances traveled in a 2-minute interval in control buffer and 1-oct. Worms significantly increased their locomotion in 1-oct, and returned to baseline rates 2-4 minutes after 1-oct withdrawal (Fig. 5E). Next, we created a simple behavioral simulation to determine whether this selective increase in locomotory activity could counteract the observed bias in reversal probabilities to achieve

overall avoidance. The environment was modeled as a line divided into two equal compartments, representing the control buffer and the 2.2 mM 1-oct halves of the arena. Worms were modeled as one-dimensional particles, free to move along the full length of the line. They were allowed to reverse direction at random along the length of the line, but were specified to reverse direction with fixed probabilities at the buffer/octanol interface, depending on which direction they approached from. Finally, the particles moved at different relative rates in the buffer and 1-oct compartments (Fig. 5F). Using our measured entry and exit reversal probabilities as inputs, this simulation showed that relative locomotory speed is a critical factor determining overall chemotaxis behavior, with a 60% speed increase sufficient to convert attraction to repulsion, given our observed interface reversal probabilities (Fig. 5F). This simulation was highly simplified, and did not take into account the persistence of higher locomotion speeds once worms exited 1-oct (Fig. 5E), which could further potentiate repulsion since faster-moving exiting worms will be oriented away from the 1-oct zone, biasing their trajectories toward the further reaches of the buffer zone where they eventually slow. Importantly, these results extend the idea that 1-oct is encoded combinatorially, generating a sensory signal with at least three components: 1) promotion of reversals when encountering increased [1-oct], 2) promotion of reversals when encountering reduced [1-oct], and 3) increase of locomotory speed in the presence of increased [1-oct]. These three modalities engage differentially at different concentrations, and are ultimately integrated to determine the final chemotactic outcome.

Repulsive and attractive signaling competitively entrain the locomotory command neurons

To understand how the repulsive and attractive 1-oct signals affect locomotory reversals at a circuit level, we examined the activity pattern of the AVA reverse command interneuron in relation to 1-oct addition and withdrawal. First, it was necessary to develop a data analysis approach to detect and quantify sensory-driven patterns of AVA activation, given the well-documented patterns of spontaneous AVA activity state transitions (*23, 27, 53-58*) that could potentially obscure a sensory signal. To reduce the impact of these transitions, we averaged at least 6 recordings of AVA activity for each condition tested so that non sensory-associated AVA transitions would tend to average out with larger numbers of recordings, akin to spike-triggered averaging (*70-72*). We then calculated the 'entrainment index' (EI), as a measure of how strongly a response is positively or negatively correlated with 1-oct application, and quantified the probability that an observed correlation is greater than a random association (see Methods), setting a threshold of P<0.05 for non-random association. To validate this test, we analyzed ASH and AWC responses to 1-oct application, which were obviously highly correlated and anti-correlated, respectively. The ASH and AWC EIs showed the expected positive and negative values (Fig. 6A), with P <0.0015, indicating a less than one in six hundred chance that the correlations in neuronal responses with 1-oct applications were random (Fig. 6B). Applying this test to AVA activity, we observed strong inverse correlation with stimulant application at 0.22 mM 1-oct (the lowest tested concentration), and a significant negative EI value (Fig. 6C, D, H). This concentration was attractive, and showed the lowest ASH:AWC activity ratio (Fig. 4), suggesting that AWC may be driving AVA with minimal interference from ASH. At higher [1-oct] (0.66, 2.2 mM), AVA entrainment with 1-oct application was shifted positively to a large extent (compared to 0.22 mM; Fig. 6C, D, H), and was not significantly different from random (P>0.05). At these concentrations, the ASH:AWC activity ratio is significantly increased (Fig. 4), consistent with the idea that a lack of entrainment may reflect interference between AWC and ASH signals impinging on AVA.

To further support the idea that conflicting sensory signals may mask one another at the level of AVA, we measured EIs in TAX-4 animals. At the lowest [1-oct], the strong negative entrainment previously observed was abolished, confirming that negative entrainment is a function of TAX-4 signaling (Fig. 6E, F, H). At the intermediate [1-oct] (0.66 mM), *tax-4* loss of function mutation caused a slight positive shift in the EI, but did not lift it above the threshold for significant entrainment (P>0.05) (Fig. 6E, F, H). However, at the highest [1-oct], loss of TAX-4 positively shifted the EI into the significant range for positive entrainment (Fig. 6E, F, H). Interestingly, all of the calculated AVA EIs (Fig. 6H) correlate well with the overall chemotactic behavior in terms of sign (i.e negative entrainment predicts attraction, positive entrainment predicts repulsion), even where the strength of entrainment with the sensory input did not exceed our set threshold for non-random entrainment (see Discussion). Most importantly, these results imply that the reverse command interneurons under these assay conditions will not necessarily show overt entrainment to sensory stimuli even though they are receiving sensory inputs because sensory stimuli may be encoded as parallel antagonistic afferent inputs which cancel one another out, until the prevailing conditions permit one input to become substantially stronger than the other.

To investigate how the 1-oct sensory signals are transmitted to AVA, we examined the interneuron AIB, which is positioned to be an integrating hub in this system because it receives afferent inputs from ASH and AWC, and many other sensory neurons, and makes strong connections with the reverse and forward command interneurons. AIB also receives inputs from RIM, which provides feedback about the motor state that is used in the regulation of sensory afferent pathways by corollary discharge (*53, 73, 74*). Initially, we did not observe significant entrainment of AIB activity with 2.2 mM 1-oct application (Fig. 6G). A potential reason might be that motor feedback was obscuring the sensory signal. To address this possibility, we subtracted AVA activity, representing the motor state, from the AIB activity. This modified AIB signal showed significant entrainment with 2.2 mM 1-oct application which was positive in sign (Fig. 6 G and H), and greater than AVA's entrainment at the same 1-oct concentration. These observations suggests that AIB serves as a sensory relay for the aversive inputs, but since aversive entrainment in AVA is weaker than that in AIB (presumably due to interference from attractive inputs), the site of integration between repulsive and attractive pathways must be downstream of AIB and either at or upstream of AVA.

Combinatorial olfactory coding facilitates context-dependent behavioral flexibility
What benefit might *C. elegans* gain from encoding 1-oct combinatorially, simultaneously activating attractive and repulsive afferent input pathways? This strategy could create a simple and versatile mechanism to modulate the behavioral output to best match internal and external conditions, such as specific odorant concentration, the presence of other odorants, or interoceptive inputs such as hunger or stress. A combinatorial strategy allows response sensitivity and even valence to be altered by simply adjusting gains or synaptic outputs within each pathway (Fig. 7A), without requiring extensive functional reorganization of the downstream connections, or recruitment of complex cross-inhibitory circuitry. For example, in the laboratory, 1-oct is used as a repellant for *C. elegans*, but its presence in the environment may convey more complex information. 1-oct is toxic to worms (thus representing a risk), which presumably explains why *C. elegans* has evolved to avoid it, at least at higher concentrations. However, 1-oct is naturally found in decaying plant material (*75, 76*), so it may also signal the presence of a bacterial food source for

worms (thus representing a reward as well). In the presence of non-toxic alternatives, other food sources are safer, but in the absence of alternatives, worms may risk toxicity to avoid starvation. Could combinatorial coding support the risk-reward decision making required of *C. elegans* when evaluating environmental 1-oct signals? Consistent with this idea, many attractive food signals, including isoamyl alcohol (IAA), suppress AWC activity (*35, 39, 59*), and could therefore decrease the activity of the attractive afferent pathway to disinhibit repulsion (Fig. 7A, B). We tested this hypothesis by conducting chemotaxis experiments at 0.66 mM 1-oct, normally an attractive concentration in our microfluidics assays, in the presence of isoamyl alcohol (IAA, 10^{-4} dilution, added to both the 1-oct and control buffer sides of the microfluidics arena). Under these conditions, worms were strongly repelled by 1-oct (Fig. 7C). The probability of reversals when entering the 1-oct zone was increased (Fig. 7D), the probability of reversals when exiting the 1-oct zone was decreased (Fig. 7D), and the repulsion-inducing modulation of locomotory speed was also increased (Fig. 7E, F). Interestingly 0.66 mM 1-oct did not cause a speed increase on its own (unlike 2.2 mM, Fig. 5). Acute application of IAA also did not affect speed. However, 0.66 mM 1-oct in the presence of IAA caused a large speed increase (2-fold). Since IAA reduces AWC activity, this IAA effect is consistent with removal of AWC inhibition of 0.66 mM 1-oct-dependent speed modulation. These three quantitative changes in locomotory parameters will each shift the chemotaxis index in the negative direction, converting 1-oct attraction to repulsion.

In network recordings, the presence of IAA greatly increased the sensory entrainment of AVA, positively correlating with stimulus application. This effect suggests that IAA will potentiate 1-oct avoidance by klinokinesis, or in other words, increasing the probability that worms will execute a reversal when they experience positive d[1-oct]/dt. These results provide proof-of-principle that rebalancing the afferent input pathways to reduce attraction and promote repulsion can control chemotaxis in an ethologically relevant manner, and demonstrate the potential utility and flexibility of the combinatorial coding strategy.

Discussion

We have investigated the response of *C. elegans* to the odorant 1-oct, using a combination of microfluidics-based behavioral analysis and whole-network Ca^{++} imaging. Ca^{++} imaging using the NeuroPAL system (*30*) showed that 1-oct activates at least 11 different sensory neurons including neurons associated with chemorepulsion and chemoattraction. 1-oct activates the ASH and AWC sensory neurons, with ON- and OFF- response kinetics, respectively. Paradoxically, this pattern suggests that sensory signals are being generated to simultaneously promote and inhibit locomotory reversals upon both 1-oct onset and offset. Based on this observation, we hypothesized that 1-oct may be encoded via two simultaneous and opposite afferent pathways: a repulsive pathway driven by ASH and possibly other neurons, and an attractive pathway driven by AWC and possibly other neurons, with the ultimate chemotactic outcome determined by the balance between the two. We tested this hypothesis by altering the balance between the pathways. This goal was achievable by simply changing the 1-oct concentration, because the ASH 1-oct response was highly sensitive to concentration while the AWC 1-oct response was not. At the highest 1-oct concentration tested (saturation, 2.2 mM), worms were repelled by 1-oct in microfluidics-based assays where worms were given the choice between control buffer or buffer containing dissolved 1-oct. However, 3-fold and 10-fold dilutions of 1-oct were attractive, presumably because the ASH-driven repulsive pathway was weakened while the AWC-driven attractive pathway remained unchanged. Consistent with this

interpretation, the attraction could be eliminated by mutating TAX-4, a critical component of sensory signal transduction in AWC. Worms achieved these chemotactic outcomes by modulating at least three distinct locomotory transitions (Fig. 8A): reversals stimulated upon entering the 1-oct zone ('entry reversals'), reversals stimulated upon exiting the 1-oct zone ('exit reversals'), and increase of locomotory activity within the 1-oct zone ('speed increase'). Based on numerical simulations, speed increase was the most important factor: reversal patterns at the highest 1-oct concentration clearly favored attraction but increased locomotory activity within the 1-oct zone overcame this bias and caused repulsion. Input from both hypothesized pathways influenced these key locomotory transitions (Fig. 8A). Entry reversals were reduced in *osm-9* mutants, implying they may be stimulated by ASH, but increased in *tax-4* mutants, implying they may be inhibited by AWC. Exit reversals were reduced in *tax-4* mutants, suggesting they are stimulated by AWC (we did not observe reciprocal regulation by OSM-9/ASH). Speed increase, like ASH activity, was dependent on 1-oct concentration, and was observed in wild type worms only at the highest 1-oct concentration. However, speed is probably also under the control of AWC because IAA addition, which suppresses AWC activity, unmasked strong speed increase at a lower 1-oct concentrations. Our findings provide consistent support for the idea that 1-oct is encoded simultaneously as two opposing afferent pathways which reciprocally modulate multiple locomotory parameters to influence overall chemotactic behaviors (Fig. 8A). AWC and ASH are probably the main drivers of these pathways, based on their activity patterns and their central roles in chemoattraction and chemorepulsion (specifically, ASH is the primary sensor for 1-oct (*34*)), but it will be important to fully map the sensory inputs into these pathways using neuron-specific silencing approaches.

Our results show how combinatorial and labeled line coding schemes may functionally interact. 1-oct is encoded combinatorially, in that it activates multiple sensory neurons (our results, and (*39*)), principally ASH and AWC. However, these neurons have long been regarded to represent repulsive and attractive labeled lines, respectively (*33, 35, 38, 77*). Thus, the initial combinatorial representation of 1-oct superimposes on the underlying labeled line architecture to dictate the final chemotactic outcome. The novel finding here is that multiple labeled lines become activated simultaneously in the context of a combinatorially-coded odorant (previous studies have shown that antagonistic sensory inputs can contribute to chemosensory responses in *C. elegans* (*78-81*), but this study is the first to show in detail how conflicting inputs interact at the behavioral and circuit levels). The labeled lines independently control multiple locomotory parameters, in opposite directions, and the conflict is resolved by the locomotory system which is sensitive to the balance of the two pathways.

The compact size of the *C. elegans* nervous system makes it feasible to characterize at a cellular and molecular level where and how these afferent pathways converge and interact. Both pathways ultimately impinge on the AVA locomotory command interneurons, which drive locomotory reversals (*82, 83*). Because AVA sensory entrainment varies smoothly as a function of the relative activity levels of the two pathways, they probably do not strongly cross-inhibit one another, which would likely produce much steeper transitions (*84*). Which neurons are likely to integrate the attractive and repulsive inputs? Based on the connectome and previous functional studies, the AIB interneuron is a logical candidate. AIB is a first layer interneuron that receives extensive inputs from sensory neurons including ASH and AWC, strong synaptic inputs from other first layer interneurons (AIZ, AIA), and electrical and chemical synapses from locomotory command

neurons AVE and RIM. AIB sends projections to the locomotory command neurons including AVA, RIM, and AVB (*61, 85*). Functionally, AIB serves as an integrating hub for multiple sensory inputs including ASH and AWC (*86, 87*). However, AIB does not appear to play the role of an integrating hub upstream of AVA for 1-oct. Prior to sensory stimulation, AIB activity closely followed reverse command interneuron activity, as has been observed previously (*23, 53*). Upon stimulation, AIB still followed the reverse command interneurons, but subtracting the reverse command signal from the AIB excitation pattern revealed that AIB was significantly entrained to the repulsive sensory input. AVA did not show this repulsive entrainment under the same conditions, therefore AIB cannot be the sole locus for integration. One intriguing possibility is that integration takes place in AVA itself, since it receives strong synaptic inputs directly from ASH, and from SAA, which mirrors the activity pattern of AWC during stimulation.

This coding scheme facilitates olfactory plasticity because the balance of the pathways, evaluated centrally, can switch odor valence. This plasticity differs from what has previously been described, where odorant responses are modulated at the periphery through lateral interactions among sensory afferents (*20, 35*) or reprogramming of individual sensory neuron responses (*46*). The broad versatility of this coding scheme can be appreciated by considering combinatorial coding and integration of inputs as analogous to the refraction of light through two prisms (Fig. 8B). First, the chemosensory neuron array in *C. elegans* creates a multi-factorial representation of the sensory stimulus, (i.e. the first prism), differentiating inputs from multiple sensory neurons, each with different downstream behavioral correlates. Downstream, the different components of this representation will eventually converge and become integrated into a coherent behavioral response by the motor command circuitry (i.e. the second prism). Because the sensory neurons have different concentration-response profiles, worms can easily generate differential behaviors dependent on stimulus concentration: as concentration is reduced, the input from the sensory layer will be skewed toward the higher-affinity components (Fig. 8B). In this study, reduced [1-oct] favored the AWC-driven response (higher affinity) over the ASH-driven response (lower affinity), converting the 1-oct response from repulsion to attraction. This architecture also provides straightforward avenues for modulation. Multiple sensory inputs may be integrated if the additional odorants modulate the initial sensory representation, either additively (via ON neurons) or subtractively (via OFF neurons Fig. 8B). Here, the addition of IAA restored repulsion at lower concentrations by reducing the contribution of AWC. In the two prism analogy, IAA is equivalent to a selective optical filter placed in the light path, skewing the overall balance back toward ASH. Other forms of plasticity, such as monoamine and neuropeptide modulation or experience dependent plasticity, may also be easily incorporated into this scheme at the sensory- or interneuron level. In a complex natural environment, where multiple odorants, interoceptive inputs, and experience-dependent plasticity are all acting simultaneously on a limited set of neurons, potential interference among modulatory signals might be avoided by funneling all of the modulatory influences through one major repulsive pathway and one major attractive pathway.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Strain, strain background (*Escherichia coli*)	OP50	Caenorhabditis Genetics Center (CGC)	OP50	
Strain, strain background (*C. elegans*)	Wild type	Caenorhabditis Genetics Center (CGC)	N2	
Strain, strain background (*C. elegans*)	otIs669 V, otIs672 [rab-3::NLS::GCaMP6s + arrd-4:NLS:::GCaMP6s]	Caenorhabditis Genetics Center (CGC)	OH15500	*(30)*
Strain, strain background (*C. elegans*)	otIs669 V, wtfIs145 [rab-3p::his-24::GCaMP6s::unc-54 3' UTR + pBX]	Caenorhabditis Genetics Center (CGC)	AML320	
Strain, strain background (*C. elegans*)	tax-4 (ok3771) III	Caenorhabditis Genetics Center (CGC)	VC3113	
Strain, strain background (C. elegans)	tax-4 (ok3771) III, otIs669 V, wtfIs145 [rab-3p::his-24::GCaMP6s::unc-54 3' UTR + pBX]	This study	FY1041	
Chemical compound, drug	Sylgard 184	Electron Microscopy Sciences	Cat# 24236 – 10	

Chemical compound, drug	Tetramisole hydrochloride	Sigma	CAS# 16595-80-5)	
Software, algorithm	Adobe Premier Pro	Adobe Inc.	https://www.adobe.com/	
Software, algorithm	Multi-Worm Tracker	*Itskovits et. al. (2018)*	https://github.com/itskov/MultiAnimalTrackerSuite	*(67)*
Software, algorithm	Mathematica	Wolfram Research	https://www.wolfram.com/	
Software, algorithm	MATLAB	MathWorks	https://www.mathworks.com	
Software, algorithm	R	Posit PBC	R-Studio v2022.07.1	
Software, algorithm	Fiji/ImageJ	*Schindelin et. al. (2012)*	https://imagej.net/software/fiji/downloads	*(88)*
Software, algorithm	Graphpad	GraphPad Software	Prism 10.2.3	

Software, algorithm	MoCo plugin	*Dubbs et. al. (2016)*	https://github.com/NTCColumbia/moco	*(89)*
Other	Silicon microfluidics molds	S. Lockery and D. Albrecht		
Other	Olfactory chips	Microkosmos	ukosmos.com	*(52)*

Worm maintenance and strains
C. elegans were maintained at 20°C on NGM agar plates and fed *E. coli* OP50 bacteria. All experiments were conducted at 20-22°C (*90, 91*). The following strains were utilized:

N2: *Wild-type.*
OH15500: otIs669 V, otIs672 [rab-3::NLS::GCaMP6s + arrd-4:NLS:::GCaMP6s] (*30*).
AML320: otIs669 V, wtfIs145 [rab-3p::his-24::GCaMP6s::unc-54 3' UTR + pBX].
VC3113: tax-4 (ok3771) III.
CX10: osm-9 (ky10) IV.
FY1041: tax-4 (ok3771) III, otIs669 V, wtfIs145 [rab-3p::his-24::GCaMP6s::unc-54 3' UTR + pBX].

Strains were obtained from the Caenorhabditis Genetics Center (CGC), funded by NIH (P40 OD010440). VC3113 was crossed into AML320 to generate FY1041, with the presence of *tax-4* verified by PCR. Strains were allowed for three generations of growth before experiments.

Microfluidics device fabrication and behavioral assays
All behavioral studies were conducted in microfluidic devices following standard protocols (*48-50*). Devices were fabricated by thoroughly mixing polydimethylsiloxane (PDMS) and a curing agent in a 10:1 ratio by weight (Sylgard 184, Electron Microscopy Sciences Cat. 24236 – 10), degassing the mixture under vacuum, and then pouring it onto a silicon mold at ~5 mm height. PDMS was cured by incubating at 65°C for at least 3 hours. Olfactory chips were obtained from Microkosmos (ukosmos.com).

Chemotaxis: Chemotaxis assays utilized a 'two-chamber' device dividing the arena into equal chambers for stimulus and control buffer (S-basal [100mM NaCl, 5mM g K_2HPO_4, 45mM KH_2PO_4, pH = 6]) (*48*). Worm behavior was recorded using AmScope microscope cameras (MD 500, MU 500-HS, AmScope, Irvine, CA), and videos were processed using Adobe Premier Pro 2024 (Adobe Inc., San Jose, CA). 1-2 day old adults were picked and washed with S-basal and loaded into the microfluidics arena. Animals were acclimatized

in the arena for 5 minutes; assays were performed between 20-40 minutes from starvation onset while they were in area-restricted search foraging mode (*51*). In the original microfluidics protocols (*49, 50*) 0.01% w/v xylene cyanol (0.1mg/ml) dye was used to verify clean separation between different domains within the microfluidics device. However, worms show clear attraction to 0.01% xylene cyanol (data not shown), indicating that the presence of dye introduces an additional olfactory stimulus. Therefore, we modified the protocol to test the flow pattern using dye before and after the assay, but remove the dye during the assay (Fig 1A). In control trials using dye, we verified that flow patterns stay stable for up to 90 minutes once properly established. If the flow pattern in the post-data collection dye test was compromised, data were rejected. Chemotaxis index was calculated between 10-20 minutes of stimulus onset, using the equation: Chemotaxis Index (CI) = (Number of worms in stimulus – Number of worms in buffer)/Total number of worms, positive and negative indices representing attraction and aversion respectively.

Distance traveled: Locomotory distance-traveled assays used the 'two-chamber' device (*48*) with both chambers containing the same solution. Worms were exposed sequentially to buffer (5 mins), stimulus (5 mins), and buffer (10 mins). A complete exchange of solutions takes approximately 1 minute. Data were collected 90 seconds after solution change to ensure complete exchange. Distance traveled per minute was recorded over 2 minutes in buffer (pre-stimulus), in 1-oct solution (stimulus), and in successive 2-minutes intervals after 1-oct withdrawal (post-stimulus). Worm tracks were manually traced and measured using Adobe Premiere Pro 2024 and Graphic for Mac (Autodesk, San Francisco, CA).

Worm locomotion simulation was written in Mathematica using AI-enabled chat (Mathematica, Princeton NJ, code available upon request). In the simulation, 20 one-dimensional dots, representing worms, were allowed to travel a one-dimensional line, representing behavior arena. The line was divided equally into two zones, representing buffer and stimulus, and dots moved freely within the line with random reversal frequency. Dot speed increased in the stimulus zone, as indicated, and reversal frequency at the interface was specified as observed from behavioral assays. 10 simulations were performed and averaged for each speed differential shown in Fig. 5F.

Reversals: Analyses of locomotory reversals were performed using larger 'stripe' chips (*49*), which provided a greater working area and facilitated automated video tracking by reducing the frequency of collisions between worms. These chips were usually configured with 3 zones of equal size: a stripe of 1-oct down the middle flanked on either side by buffer, although for some experiments, the 1-oct was placed on one side. Video recordings of worm behavior were analyzed using the Multi-Worm Tracker (*92*), using MatLab (Mathworks, Natick, MA), with manual analysis of individual tracks to quantify reversals.

Whole-brain calcium imaging
Neuronal activity was examined by measuring calcium transients detected by fluorescence change (GCaMP6s, OH15500 (*30*), AML320 (*93*)). Calcium imaging was conducted as previously described (*23, 52, 53*). Animals were anesthetized with 1 mM tetramisole hydrochloride (Sigma, CAS 16595-80-5) in S-basal buffer and imaged in PDMS olfactory microfluidic devices, which trap the worm in a chemical delivery system and provides inlets to deliver either buffer or stimulus to the nose using laminar flow (*52*). Head-ganglia neurons were recorded using Leica confocal microscope (SP8) at 0.6 - 0.8 Hz rate. Odorants were diluted in S-basal. Each worm was recorded for 12-14 minutes within 20-

40 minutes from starvation onset, starting with 6 minutes of buffer exposure followed by 6-7 30-second pulses of stimulus with 30-second intervals. Calcium imaging was followed by immediate recording of NeuroPAL color markers. For +IAA experiments, IAA delivery to the nose was started after 5 minutes of recording (1 minute before the first stimulus application) and IAA was was present in the stimulus and buffer channels thereafter. For concentration-response of ASH and AWC experiments, three 0.22 mM, two 0.66 mM and two 2.2 mM 1-oct pulses were given serially to the same worm; repeated application produced identical response amplitudes.

Recorded image files were stacked and analyzed using Fiji/ImageJ (*94, 95*) followed by motion correction with 'MoCo' plug-in (*89*). Neuron identity was manually assigned and matched with calcium response using guidelines described by Yemini and Hobert (*30*). Neurons with ambiguous identities were excluded from the analysis (Fig. 2). Individual neuron activity was then normalized and re-sampled, before averaging, by 'Min-Max normalization' and 'Linear interpolation' respectively, using custom R-scripts (*96*).

Data analyses and statistics
Statistical analyses were performed using Prism v10.2.1 (GraphPad Software Inc., La Jolla, CA), R v4.3, R-Studio v2022.07.1 (Posit PBC, Boston, MA), MatLab vR2023A (MathWorks, CA) and Microsoft Excel 2016. Chemotaxis assays were repeated 3-5 times and analyzed by *Student's t-test* (un-paired). Pooled data from distance-traveled assays were analyzed by one-way ANOVA with post-tests. Reversal assays were repeated 3 times, data were pooled, and analyzed using Fisher exact tests.

Entrainment indices (EIs) were calculated using averaged, normalized calcium responses. First, area-under-curve (AUC) values were calculated for all possible 30s epochs by summing the Ca^{++} signal intensities over 30s after first subtracting the corresponding initial Ca^{++} signal intensity. EIs were calculated by adding six AUCs; stimulus-dependent EIs were calculated by adding the six AUCs corresponding to stimulus application intervals, while random EIs were calculated by adding six randomly-chosen AUCs. The significance of stimulus entrainment was estimated as the probability that the stimulus-dependent EI would exceed any given random EI, with probabilities < 0.05 considered significant (i.e. P<0.05). A P value <0.0015 indicated that the experimental EI was larger (or smaller) than all possible random EIs within the dataset. EIs for AVA were calculated using a 10-second lag, across all conditions, because, where obvious sensory entrainment was seen, AVA signals showed consistently slowed responses compared to sensory signals. No lag was incorporated for sensory and first-layer interneurons (ASH, AWC, AIB).

Neuronal activity correlations were calculated using Spearman's rank correlation test and corrected with Fisher's z-transformation before testing statistical significance (*97-99*), using custom R scripts.

Acknowledgments

We would like to thank Shawn Lockery and Dirk Albrecht for invaluable help and materials for the microfluidics behavioral assays, the *C. elegans* Genetics Center for strains, Tomer Avidor-Reiss and Qian Chen for assistance with confocal microscopy,

Funding: The University of Toledo Office of Research provided funding support through the DeArce-Koch Memorial Foundation Award, the Small Award Program, and the Interdisciplinary Research Initiation Award. WGRV was supported by was NIH NIGMS T32-G-RISE grant number 1T32GM144873-01. The funding sources were not involved in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions:
Conceptualization: MZHS, BAB, WGR, RM. Software: BAB. Methodology: MZHS, BAB, CAE, LGJ. Investigation: MZHS, WGR, CAE, BNS, ARM, RM, BAB. Visualization: MZHS, WGR, BAB. Supervision: BAB, RM. Writing (original draft): MZHS, BAB. Funding acquisition BAB, RM, WGRV.

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: All data needed to evaluate the conclusions in the paper are present in the main text. Some analyses were performed in custom-written R and Mathematica code and will be available for download (https://doi.org/10.5281/zenodo.13328278).

References

1. M. N. Andersson, C. Löfstedt, R. D. Newcomb, Insect olfaction and the evolution of receptor tuning. *Frontiers in Ecology and Evolution* **3**, 53 (2015).
2. A. Haverkamp, B. S. Hansson, M. Knaden, Combinatorial codes and labeled lines: how insects use olfactory cues to find and judge food, mates, and oviposition sites in complex environments. *Frontiers in Physiology* **9**, 49 (2018).
3. V. Grabe, S. Sachse, Fundamental principles of the olfactory code. *Biosystems* **164**, 94-101 (2018).
4. K. Mori, H. Sakano, Olfactory circuitry and behavioral decisions. *Annual Review of Physiology* **83**, 231-256 (2021).
5. B. Malnic, J. Hirono, T. Sato, L. B. Buck, Combinatorial receptor codes for odors. *Cell* **96**, 713-723 (1999).
6. K. Touhara, L. B. Vosshall, Sensing odorants and pheromones with chemosensory receptors. *Annual review of physiology* **71**, 307-332 (2009).
7. T. D. Wyatt, Introduction to chemical signaling in vertebrates and invertebrates. (2014).
8. M. C. Stensmyr, H. K. Dweck, A. Farhan, I. Ibba, A. Strutz, L. Mukunda, J. Linz, V. Grabe, K. Steck, S. Lavista-Llanos, A conserved dedicated olfactory circuit for detecting harmful microbes in Drosophila. *Cell* **151**, 1345-1357 (2012).
9. S. A. Ebrahim, H. K. Dweck, J. Stökl, J. E. Hofferberth, F. Trona, K. Weniger, J. Rybak, Y. Seki, M. C. Stensmyr, S. Sachse, Drosophila avoids parasitoids by sensing their semiochemicals via a dedicated olfactory circuit. *PLoS biology* **13**, e1002318 (2015).
10. H.-C. Jiang, S. J. Park, I.-H. Wang, D. M. Bear, A. Nowlan, P. L. Greer, CD20/MS4A1 is a mammalian olfactory receptor expressed in a subset of olfactory sensory neurons that mediates innate avoidance of predators. *Nature Communications* **15**, 3360 (2024).
11. H. Saito, H. Nishizumi, S. Suzuki, H. Matsumoto, N. Ieki, T. Abe, H. Kiyonari, M. Morita, H. Yokota, N. Hirayama, Immobility responses are induced by photoactivation of single glomerular species responsive to fox odour TMT. *Nature communications* **8**, 16011 (2017).
12. N. Horio, K. Murata, K. Yoshikawa, Y. Yoshihara, K. Touhara, Contribution of individual olfactory receptors to odor-induced attractive or aversive behavior in mice. *Nature communications* **10**, 209 (2019).
13. J. Cande, B. Prud'homme, N. Gompel, Smells like evolution: the role of chemoreceptor evolution in behavioral change. *Current opinion in neurobiology* **23**, 152-158 (2013).
14. L. Buck, R. Axel, A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* **65**, 175-187 (1991).
15. P. Mombaerts, F. Wang, C. Dulac, S. K. Chao, A. Nemes, M. Mendelsohn, J. Edmondson, R. Axel, Visualizing an olfactory sensory map. *Cell* **87**, 675-686 (1996).
16. R. M. Blazing, K. M. Franks, Odor coding in piriform cortex: mechanistic insights into distributed coding. *Curr Opin Neurobiol* **64**, 96-102 (2020).
17. A. Pérez-Gómez, K. Bleymehl, B. Stein, M. Pyrski, L. Birnbaumer, S. D. Munger, T. Leinders-Zufall, F. Zufall, P. Chamero, Innate predator odor aversion driven by parallel olfactory subsystems that converge in the ventromedial hypothalamus. *Current Biology* **25**, 1340-1346 (2015).
18. C. Poo, G. Agarwal, N. Bonacchi, Z. F. Mainen, Spatial maps in piriform cortex during olfactory navigation. *Nature* **601**, 595-599 (2022).
19. Y. Chen, X. Chen, B. Baserdem, H. Zhan, Y. Li, M. B. Davis, J. M. Kebschull, A. M. Zador, A. A. Koulakov, D. F. Albeanu, High-throughput sequencing of single neuron projections reveals spatial organization in the olfactory cortex. *Cell* **185**, 4117-4134. e4128 (2022).

20. D. Zocchi, S. Y. Emily, V. Hauser, T. F. O'Connell, E. J. Hong, Parallel encoding of CO2 in attractive and aversive glomeruli by selective lateral signaling between olfactory afferents. *Current Biology* **32**, 4225-4239. e4227 (2022).

21. T. Schrödel, R. Prevedel, K. Aumayr, M. Zimmer, A. Vaziri, Brain-wide 3D imaging of neuronal activity in Caenorhabditis elegans with sculpted light. *Nature methods* **10**, 1013-1020 (2013).

22. R. Prevedel, Y.-G. Yoon, M. Hoffmann, N. Pak, G. Wetzstein, S. Kato, T. Schrödel, R. Raskar, M. Zimmer, E. S. Boyden, Simultaneous whole-animal 3D imaging of neuronal activity using light-field microscopy. *Nature methods* **11**, 727-730 (2014).

23. S. Kato, H. S. Kaplan, T. Schrödel, S. Skora, T. H. Lindsay, E. Yemini, S. Lockery, M. Zimmer, Global brain dynamics embed the motor command sequence of Caenorhabditis elegans. *Cell* **163**, 656-669 (2015).

24. J. P. Nguyen, F. B. Shipley, A. N. Linder, G. S. Plummer, M. Liu, S. U. Setru, J. W. Shaevitz, A. M. Leifer, Whole-brain calcium imaging with cellular resolution in freely behaving Caenorhabditis elegans. *Proceedings of the National Academy of Sciences* **113**, E1074-E1081 (2016).

25. I. Kotera, N. A. Tran, D. Fu, J. H. Kim, J. Byrne Rodgers, W. S. Ryu, Pan-neuronal screening in Caenorhabditis elegans reveals asymmetric dynamics of AWC neurons is critical for thermal avoidance behavior. *elife* **5**, e19021 (2016).

26. V. Venkatachalam, N. Ji, X. Wang, C. Clark, J. K. Mitchell, M. Klein, C. J. Tabone, J. Florman, H. Ji, J. Greenwood, Pan-neuronal imaging in roaming Caenorhabditis elegans. *Proceedings of the National Academy of Sciences* **113**, E1082-E1088 (2016).

27. A. L. Nichols, T. Eichler, R. Latham, M. Zimmer, A global brain state underlies C. elegans sleep behavior. *Science* **356**, eaam6851 (2017).

28. Y. Toyoshima, S. Wu, M. Kanamori, H. Sato, M. S. Jang, S. Oe, Y. Murakami, T. Teramoto, C. Park, Y. Iwasaki, Neuron ID dataset facilitates neuronal annotation for whole-brain activity imaging of C. elegans. *BMC biology* **18**, 1-20 (2020).

29. K. M. Hallinen, R. Dempsey, M. Scholz, X. Yu, A. Linder, F. Randi, A. K. Sharma, J. W. Shaevitz, A. M. Leifer, Decoding locomotion from population neural activity in moving C. elegans. *Elife* **10**, e66135 (2021).

30. E. Yemini, A. Lin, A. Nejatbakhsh, E. Varol, R. Sun, G. E. Mena, A. D. T. Samuel, L. Paninski, V. Venkatachalam, O. Hobert, NeuroPAL: A Multicolor Atlas for Whole-Brain Neuronal Identification in C. elegans. *Cell* **184**, 272-288.e211 (2021).

31. U. Dag, I. Nwabudike, D. Kang, M. A. Gomes, J. Kim, A. A. Atanas, E. Bueno, C. Estrem, S. Pugliese, Z. Wang, Dissecting the functional organization of the C. elegans serotonergic system at whole-brain scale. *Cell* **186**, 2574-2592. e2520 (2023).

32. C. I. Bargmann, E. Hartwieg, H. R. Horvitz, Odorant-selective genes and neurons mediate olfaction in C. elegans. *Cell* **74**, 515-527 (1993).

33. E. R. Troemel, B. E. Kimmel, C. I. Bargmann, Reprogramming chemotaxis responses: sensory neurons define olfactory preferences in C. elegans. *Cell* **91**, 161-169 (1997).

34. M. Y. Chao, H. Komatsu, H. S. Fukuto, H. M. Dionne, A. C. Hart, Feeding status and serotonin rapidly and reversibly modulate a Caenorhabditis elegans chemosensory circuit. *Proc Natl Acad Sci U S A* **101**, 15512-15517 (2004).

35. K. Yoshida, T. Hirotsu, T. Tagawa, S. Oda, T. Wakabayashi, Y. Iino, T. Ishihara, Odour concentration-dependent olfactory preference change in C. elegans. *Nature communications* **3**, 739 (2012).

36. J. S. Greene, M. Dobosiewicz, R. A. Butcher, P. T. McGrath, C. I. Bargmann, Regulatory changes in two chemoreceptor genes contribute to a Caenorhabditis elegans QTL for foraging behavior. *elife* **5**, e21454 (2016).

37. S. Rengarajan, E. A. Hallem, Olfactory circuits and behaviors of nematodes. *Current Opinion in Neurobiology* **41**, 136-148 (2016).
38. D. M. Ferkey, P. Sengupta, N. D. L'Etoile, Chemosensory signal transduction in Caenorhabditis elegans. *Genetics* **217**, (2021).
39. A. Lin, S. Qin, H. Casademunt, M. Wu, W. Hung, G. Cain, N. Z. Tan, R. Valenzuela, L. Lesanpezeshki, V. Venkatachalam, C. Pehlevan, M. Zhen, A. D. T. Samuel, Functional imaging and quantification of multineuronal olfactory responses in C. elegans. *Science Advances* **9**, eade1249 (2023).
40. E. R. Troemel, J. H. Chou, N. D. Dwyer, H. A. Colbert, C. I. Bargmann, Divergent seven transmembrane receptors are candidate chemosensory receptors in C. elegans. *Cell* **83**, 207-218 (1995).
41. R. T. Wragg, V. Hapiak, S. B. Miller, G. P. Harris, J. Gray, P. R. Komuniecki, R. W. Komuniecki, Tyramine and octopamine independently inhibit serotonin-stimulated aversive behaviors in Caenorhabditis elegans through two novel amine receptors. *Journal of Neuroscience* **27**, 13402-13412 (2007).
42. V. Hapiak, P. Summers, A. Ortega, W. J. Law, A. Stein, R. Komuniecki, Neuropeptides amplify and focus the monoaminergic inhibition of nociception in Caenorhabditis elegans. *Journal of Neuroscience* **33**, 14107-14116 (2013).
43. J. A. Zahratka, P. D. Williams, P. J. Summers, R. W. Komuniecki, B. A. Bamber, Serotonin differentially modulates Ca2+ transients and depolarization in a C. elegans nociceptor. *J Neurophysiol* **113**, 1041-1050 (2015).
44. P. D. Williams, J. A. Zahratka, M. Rodenbeck, J. Wanamaker, H. Linzie, B. A. Bamber, Serotonin disinhibits a Caenorhabditis elegans sensory neuron by suppressing Ca2+-dependent negative feedback. *Journal of Neuroscience* **38**, 2069-2080 (2018).
45. A. Debnath, P. D. Williams, B. A. Bamber, Reduced Ca2+ transient amplitudes may signify increased or decreased depolarization depending on the neuromodulatory signaling pathway. *Frontiers in Neuroscience* **16**, 931328 (2022).
46. M. Khan, A. H. Hartmann, M. P. O'Donnell, M. Piccione, A. Pandey, P.-H. Chao, N. D. Dwyer, C. I. Bargmann, P. Sengupta, Context-dependent reversal of odorant preference is driven by inversion of the response in a single sensory neuron type. *PLoS biology* **20**, e3001677 (2022).
47. E. Fryer, S. Guha, L. E. Rogel-Hernandez, T. Logan-Garbisch, H. Farah, E. Rezaei, I. N. Mollhoff, A. L. Nekimken, A. Xu, L. Selin Seyahi, S. Fechner, S. Druckmann, T. R. Clandinin, S. Y. Rhee, M. B. Goodman, An efficient behavioral screening platform classifies natural products and other chemical cues according to their chemosensory valence in C. elegans. *bioRxiv*, (2024).
48. S. R. Lockery, K. J. Lawton, J. C. Doll, S. Faumont, S. M. Coulthard, T. R. Thiele, N. Chronis, K. E. McCormick, M. B. Goodman, B. L. Pruitt, Artificial dirt: microfluidic substrates for nematode neurobiology and behavior. *Journal of neurophysiology* **99**, 3136-3143 (2008).
49. D. R. Albrecht, C. I. Bargmann, High-content behavioral analysis of Caenorhabditis elegans in precise spatiotemporal chemical environments. *Nat Methods* **8**, 599-605 (2011).
50. R. C. Lagoy, D. R. Albrecht, Microfluidic devices for behavioral analysis, microscopy, and neuronal imaging in Caenorhabditis elegans. *C. elegans: Methods and Applications*, 159-179 (2015).
51. J. M. Gray, J. J. Hill, C. I. Bargmann, A circuit for navigation in Caenorhabditis elegans. *Proc Natl Acad Sci U S A* **102**, 3184-3191 (2005).
52. N. Chronis, M. Zimmer, C. I. Bargmann, Microfluidics for in vivo imaging of neuronal and behavioral activity in Caenorhabditis elegans. *Nat Methods* **4**, 727-731 (2007).

53. A. Gordus, N. Pokala, S. Levy, S. W. Flavell, C. I. Bargmann, Feedback from network states generates variability in a probabilistic olfactory circuit. *Cell* **161**, 215-227 (2015).

54. M. Katz, F. Corson, W. Keil, A. Singhal, A. Bae, Y. Lu, Y. Liang, S. Shaham, Glutamate spillover in C. elegans triggers repetitive behavior through presynaptic activation of MGL-2/mGluR5. *Nature Communications* **10**, 1882 (2019).

55. G. S. Wirak, J. Florman, M. J. Alkema, C. W. Connor, C. V. Gabel, Age-associated changes to neuronal dynamics involve a disruption of excitatory/inhibitory balance in C. elegans. *Elife* **11**, e72135 (2022).

56. A. Sordillo, C. I. Bargmann, Behavioral control by depolarized and hyperpolarized states of an integrating neuron. *Elife* **10**, e67723 (2021).

57. Z. Li, J. Zhou, K. A. Wani, T. Yu, E. A. Ronan, B. J. Piggott, J. Liu, X. S. Xu, A C. elegans neuron both promotes and suppresses motor behavior to fine tune motor output. *Frontiers in Molecular Neuroscience* **16**, (2023).

58. J. Meng, T. Ahamed, B. Yu, W. Hung, S. EI Mouridi, Z. Wang, Y. Zhang, Q. Wen, T. Boulin, S. Gao, A tonically active master neuron modulates mutually exclusive motor states at two timescales. *Science Advances* **10**, eadk0002 (2024).

59. S. H. Chalasani, N. Chronis, M. Tsunozaki, J. M. Gray, D. Ramot, M. B. Goodman, C. I. Bargmann, Dissecting a circuit for olfactory behaviour in Caenorhabditis elegans. *Nature* **450**, 63-70 (2007).

60. J. Im Choi, H. K. Lee, H. S. Kim, S. Y. Park, T. Y. Lee, K.-h. Yoon, J. I. Lee, Odor-dependent temporal dynamics in Caenorhabitis elegans adaptation and aversive learning behavior. *PeerJ* **6**, e4956 (2018).

61. J. G. White, E. Southgate, J. N. Thomson, S. Brenner, The structure of the nervous system of the nematode Caenorhabditis elegans. *Philos Trans R Soc Lond B Biol Sci* **314**, 1-340 (1986).

62. J. White, Clues to basis of exploratory behaviour of the C. elegans snout from head somatotropy. *Philos Trans R Soc Lond B Biol Sci* **373**, (2018).

63. M.-H. Ouellette, M. J. Desrochers, I. Gheta, R. Ramos, M. Hendricks, A gate-and-switch model for head orientation behaviors in Caenorhabditis elegans. *Eneuro* **5**, (2018).

64. C. M. Coburn, C. I. Bargmann, A putative cyclic nucleotide–gated channel is required for sensory development and function in C. elegans. *Neuron* **17**, 695-706 (1996).

65. M. Wojtyniak, A. G. Brear, D. M. O'Halloran, P. Sengupta, Cell- and subunit-specific mechanisms of CNG channel ciliary trafficking and localization in C. elegans. *J Cell Sci* **126**, 4381-4395 (2013).

66. S. R. Taylor, G. Santpere, A. Weinreb, A. Barrett, M. B. Reilly, C. Xu, E. Varol, P. Oikonomou, L. Glenwinkel, R. McWhirter, Molecular topography of an entire nervous system. *Cell* **184**, 4329-4347. e4323 (2021).

67. E. Itskovits, A. Levine, E. Cohen, A. Zaslaver, A multi-animal tracker for studying complex behaviors. *BMC biology* **15**, 1-16 (2017).

68. H. A. Colbert, T. L. Smith, C. I. Bargmann, OSM-9, A novel protein with structural similarity to channels, is required for olfaction, mechanosensation, and olfactory adaptation incaenorhabditis elegans. *Journal of Neuroscience* **17**, 8259-8269 (1997).

69. M. J. Ezak, E. Hong, A. Chaparro-Garcia, D. M. Ferkey, Caenorhabditis elegans TRPV channels function in a modality-specific pathway to regulate response to aberrant sensory signaling. *Genetics* **185**, 233-244 (2010).

70. E. De Boer, P. Kuyper, Triggered correlation. *IEEE Transactions on Biomedical Engineering*, 169-179 (1968).

71. H. L. Bryant, J. P. Segundo, Spike initiation by transmembrane current: a white-noise analysis. *The Journal of physiology* **260**, 279-314 (1976).

840   72.   A. G. Davidson, R. O'Dell, V. Chan, M. H. Schieber, Comparing effects in spike-triggered
841   averages of rectified EMG across different behaviors. *J Neurosci Methods* **163**, 283-294
842   (2007).
843   73.   N. Ji, V. Venkatachalam, H. D. Rodgers, W. Hung, T. Kawano, C. M. Clark, M. Lim, M.
844   J. Alkema, M. Zhen, A. D. Samuel, Corollary discharge promotes a sustained motor state
845   in a neural circuit for navigation. *Elife* **10**, e68848 (2021).
846   74.   J. Riedl, C. Fieseler, M. Zimmer, Tyraminergic corollary discharge filters reafferent
847   perception in a chemosensory neuron. *Curr Biol* **32**, 3048-3058.e3046 (2022).
848   75.   S. Bourgou, F. Z. Rahali, I. Ourghemmi, M. Saïdani Tounsi, Changes of peel essential oil
849   composition of four Tunisian citrus during fruit maturation. *ScientificWorldJournal* **2012**,
850   528593 (2012).
851   76.   P. Neelakandan, C.-C. Young, A. Hameed, Y.-N. Wang, K.-N. Chen, F.-T. Shen, Volatile
852   1-octanol of tea (Camellia sinensis L.) fuels cell division and indole-3-acetic acid
853   production in phylloplane isolate Pseudomonas sp. NEEL19. *Scientific Reports* **11**, 2788
854   (2021).
855   77.   M. Tsunozaki, S. H. Chalasani, C. I. Bargmann, A behavioral switch: cGMP and PKC
856   signaling in olfactory neurons reverses odor preference in C. elegans. *Neuron* **59**, 959-971
857   (2008).
858   78.   R. K. Hukema, S. Rademakers, M. P. Dekkers, J. Burghoorn, G. Jansen, Antagonistic
859   sensory cues generate gustatory plasticity in Caenorhabditis elegans. *EMBO J* **25**, 312-322
860   (2006).
861   79.   H. Jang, K. Kim, S. J. Neal, E. Macosko, D. Kim, R. A. Butcher, D. M. Zeiger, C. I.
862   Bargmann, P. Sengupta, Neuromodulatory state and sex specify alternative behaviors
863   through antagonistic synaptic pathways in C. elegans. *Neuron* **75**, 585-592 (2012).
864   80.   T. Murayama, I. N. Maruyama, Decision making in C. elegans chemotaxis to alkaline pH:
865   Competition between two sensory neurons, ASEL and ASH. *communicative & integrative*
866   *Biology* **6**, 1007-1012 (2013).
867   81.   D. D. Ghosh, T. Sanders, S. Hong, L. Y. McCurdy, D. L. Chase, N. Cohen, M. R. Koelle,
868   M. N. Nitabach, Neural architecture of hunger-dependent multisensory decision making in
869   C. elegans. *Neuron* **92**, 1049-1062 (2016).
870   82.   M. Chalfie, J. E. Sulston, J. G. White, E. Southgate, J. N. Thomson, S. Brenner, The
871   neural circuit for touch sensitivity in Caenorhabditis elegans. *Journal of Neuroscience* **5**,
872   956-964 (1985).
873   83.   B. J. Piggott, J. Liu, Z. Feng, S. A. Wescott, X. S. Xu, The neural circuits and synaptic
874   mechanisms underlying motor initiation in C. elegans. *Cell* **147**, 922-933 (2011).
875   84.   B. Grothe, M. Pecka, D. McAlpine, Mechanisms of sound localization in mammals.
876   *Physiological reviews* **90**, 983-1012 (2010).
877   85.   D. Witvliet, B. Mulcahy, J. K. Mitchell, Y. Meirovitch, D. R. Berger, Y. Wu, Y. Liu, W.
878   X. Koh, R. Parvathala, D. Holmyard, Connectomes across development reveal principles
879   of brain maturation. *Nature* **596**, 257-261 (2021).
880   86.   J. C. Campbell, L. F. Polan-Couillard, I. D. Chin-Sang, W. G. Bendena, NPR-9, a galanin-
881   like G-protein coupled receptor, and GLR-1 regulate interneuronal circuitry underlying
882   multisensory integration of environmental cues in Caenorhabditis elegans. *PLoS genetics*
883   **12**, e1006050 (2016).
884   87.   P. J. Summers, R. M. Layne, A. C. Ortega, G. P. Harris, B. A. Bamber, R. W.
885   Komuniecki, Multiple sensory inputs are extensively integrated to modulate nociception in
886   C. elegans. *Journal of Neuroscience* **35**, 10331-10342 (2015).
887   88.   J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S.
888   Preibisch, C. Rueden, S. Saalfeld, B. Schmid, Fiji: an open-source platform for biological-
889   image analysis. *Nature methods* **9**, 676-682 (2012).

89. A. Dubbs, J. Guevara, R. Yuste, moco: Fast Motion Correction for Calcium Imaging.
Front Neuroinform **10**, 6 (2016).
90. S. Brenner, The genetics of Caenorhabditis elegans. *Genetics* **77**, 71-94 (1974).
91. T. Stiernagle, Maintenance of C. elegans. WormBook. The C. elegans research
community. *WormBook*, (2006).
92. E. Itskovits, R. Ruach, A. Kazakov, A. Zaslaver, Concerted pulsatile and graded neural
dynamics enables efficient chemotaxis in C. elegans. *Nat Commun* **9**, 2866 (2018).
93. X. Yu, M. S. Creamer, F. Randi, A. K. Sharma, S. W. Linderman, A. M. Leifer, Fast deep
neural correspondence for tracking and identifying neurons in C. elegans using semi-
synthetic training. *Elife* **10**, (2021).
94. J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S.
Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J. Y. Tinevez, D. J. White, V. Hartenstein,
K. Eliceiri, P. Tomancak, A. Cardona, Fiji: an open-source platform for biological-image
analysis. *Nat Methods* **9**, 676-682 (2012).
95. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* **9**, 671-675 (2012).
96. M. Kuramochi, M. Doi, A Computational Model Based on Multi-Regional Calcium
Imaging Represents the Spatio-Temporal Dynamics in a Caenorhabditis elegans Sensory
Neuron. *PLoS One* **12**, e0168415 (2017).
97. A. S. Mahadevan, U. A. Tooley, M. A. Bertolero, A. P. Mackey, D. S. Bassett, Evaluating
the sensitivity of functional connectivity measures to motion artifact in resting-state fMRI
data. *NeuroImage* **241**, 118408 (2021).
98. R. Liégeois, A. Santos, V. Matta, D. Van De Ville, A. H. Sayed, Revisiting correlation-
based functional connectivity and its relationship with structural connectivity. *Network
Neuroscience* **4**, 1235-1251 (2020).
99. L. Myers, M. J. Sirois, Spearman correlation coefficients, differences between.
Encyclopedia of statistical sciences **12**, (2004).

Figures

chemotaxis index (CI) =

$$\overline{X}_{t=10-20} \left(\frac{\text{\# worms in stimulus - \# worms in control buffer}}{\text{total \# worms}} \right)$$

Fig. 1. 1-oct chemotaxis in microfluidic arenas. (**A**) Microfluidics arenas were
configured with two zones of equal area, one containing a 1-oct solution and the
other containing buffer alone (1-oct zone contains 0.01% xylene cyanol dye for
visualization here; assays are run without dye). (**B, C**) Typical distribution of
worms when first exposed to 1-oct (B), and after 15 min of 2.2 mM 1-oct exposure
(C). (**D**) Time-course of 1-oct chemotaxis. (**E**) Calculation of chemotaxis index,
averaging worm counts every 2 minutes from 10-20 minutes after introduction of
1-oct solution.

Fig. 2. Network recording of 1-oct (2.2 mM) response. (**A**) Representative image of a NeuroPAL worm immobilized in a microfluidics device visualized using the mTagBFP2, CyOFP, and mNeptune2.5 reporters for neuronal identification (*22*). (**B, C**) Pan-neuronal nGCaMP6s signal for the region indicated by the box in (A) during 1-oct stimulation, showing 1-oct offset (B) and onset (C). Specific time points in the recording for (B) and (C) are indicated by the black triangles in (D). Selected sensory neurons and interneurons are indicated by dashed white circles. (**D**) Heat map of activity profiles for 56 identifiable neurons. Raw (i.e. non-normalized) fluorescence values shown. Stimulation time-course shown on top row (white is buffer, orange is 1-oct). Green and orange vertical bars at left indicate first-layer and locomotory command neurons (+RIM), respectively.

Fig. 3. Neuronal activity correlations. (A-C) Neuronal activity patterns from worm shown in Fig - 2. (**A**) Stochastic activity state transitions of locomotory command and associated interneurons, with AIY and AVB showing opposite activity patterns to AVA, AVD, AVE, RIM, AIB, and AIZ. (**B**) Sensory neuron activity patterns. (**C**) Selected neuron pairs highlighting the opposite activity patterns of AWC and ASH (top), and the correlated activity patterns of AWC with AIA (middle) and SAA (bottom). (**D, E**) Activity correlations for 12 selected neurons based on data from 5 worms. (**D**) Graph representation of activity correlations before stimulus (minutes 0-6, left) and during stimulus (minutes 6-12, right). Neurons pairs with the strongest positive correlations are closest to one another and connected by the darkest purple lines; neurons with the strongest negative correlations are furthest apart and connected by the darkest green lines. (**E**) Relationships from (D) presented as a correlation matrix using the same color code.

Fig. 4. 1-oct valence reverses as a function of concentration. (A-C) ASH ON responses (A, C) are concentration dependent, while AWC ON and OFF responses (B, C) are not, over a 10-fold concentration range from 0.22 to 2.2 mM. Representative traces from a single ASH or AWC neuron are shown (A, B, respectively), averages shown in (C) *** P<0.001, ns non-significant, paired ANOVA (Tukey's multiple comparison test). (**D**) For wild type (black points/line), chemotaxis index for 1-oct switches from negative (repulsive) at the highest concentration (2.2 mM) to positive (attractive) at the lowest concentrations (0.22 and 0.66 mM), and insensitivity at intermediate concentrations (1.21 and 1.54 mM). In *tax-4* mutants, 1-oct responses are repulsive at all tested concentrations (red points/line), n = 3-5. (**E**) Results are consistent with a model in which repulsive and attractive afferent pathways (driven by ASH and AWC, respectively) are co-active, and balance to determine overall chemotactic response.

969
Fig. 5. Locomotory reversals and speed modulation stimulated by 1-oct. (**A**) Example of a single worm track traced by the automated tracking software. (**B**) Worms execute reversals probabilistically at the 1-oct-buffer interface dependent on TAX-4 and OSM-9 signaling cascades. (**C, D**) Reversal probabilities at 1-oct-buffer interface for wild-type, *tax-4*, and *osm-9* (see panel B for summary and explanation of terms). (**C**) movement from buffer zone to 1-oct zone ('entries'); (**D**) movement from 1-oct zone to buffer zone ('exits'). (**E**) Modulation of locomotory activity. Distance traveled in 2 minutes prior to, during, and after flooding the entire chip with 1-oct solution. (**F**) Simulation of chemotaxis outcome based on reversal probabilities (from C, D) and locomotory activity modulation (from E). Points shown are averages of 10 simulation trials with SEM error bars. ***, **** in C, D indicate $P<0.001, 0.0001$ Fisher's Exact Test. *, ** in E indicate $P<0.05, 0.01$, ANOVA (Tukey's multiple comparison test).

Fig. 6. Entrainment of neuronal activity by 1-oct stimulation. (**A, B**) Validation of the entrainment index measurement using ASH (A) and AWC (B), showing very strong positive and very strong negative entrainment by 1-oct application, as expected. Gray dots indicate each randomized sum, while red and blue dots indicate the stimulus-specific sums of ASH and AWC, respectively (see Methods). (**C-F**) Superimposed ASH and AVA activity patterns after stimulation by 0.22, 0.66, and 2.2 mM 1-oct in wild type (C, D) and *tax-4* (E, F). Single worm traces (C, E) and averaged traces are shown (D, F, n ranges from 6-21 worms). (**G**) AIB trace (averaged); showing unprocessed (upper panel, purple trace) and AIB after subtracting AVA activity (lower panel, green trace). (**H**) Entrainment indices for all recordings in C through G, with gray dots indicating randomized EIs and the bold dots represent the stimulus-specific EIs for AVA (black), unprocessed AIB (purple), and AVA-subtracted AIB (green). P values in A, B and H represent the probability that a randomized EI would be as great or greater than the stimulus-selected sum, with a significance cut-off of $P<0.05$.

Fig. 7. Balanced repulsion and attraction support context-dependent modulation of sensory responses. (**A, B**) Combinatorial coding-based mechanism for modulating sensory responses. (**A**) At lower 1-oct concentrations, the attractive afferent pathway is relatively strong, resulting in attraction. (**B**) Addition of IAA reduces AWC activity, weakening the attractive pathway relative to the repulsive pathway, and reversing the 1-oct valence. (**C**) Addition of 0.92 mM IAA reverses the valence of 1-oct (0.66 mM) from attractive to repulsive. (**D**) IAA biases reversal behavior (at the 1-oct/buffer interface) toward repulsion. Reversals upon entering 1-oct zone are increased (left side/red), and reversals upon exiting the 1-oct zone are decreased (right side /green). (**E-F**) Effect of IAA on locomotory activity. (**E**) 1-oct (0.66 mM) in the absence of IAA does not cause increased speed. (**F**) 1-oct (0.66 mM) in the presence of IAA causes markedly increased speed. (**G**) Single (upper) and averaged (lower) recordings of ASH and AVA before and after addition of IAA and 1-oct (0.66mM) as shown, demonstrating increased AVA entrainment to stimulus (compare to Fig-6D/middle). Blue bar indicates addition of IAA (in both control buffer and 1-oct solutions). (**H**) AVA entrainment index confirms that the presence of IAA causes AVA to become significantly entrained with the stimulus. ** in C indicates $P<0.01$, *t-test*; *** and **** in D indicate $P<0.001, 0.0001$ respectively, Fisher's Exact Test; *, *** in F indicate $P<0.05$ and 0.001, respectively, ANOVA and Tukey's multiple comparison test.

1022

1023

Fig 8. Combinatorial coding of 1-oct as an attractant and a repellant, simultaneously,

facilitates sensory motor coupling and context-dependent modulation in *C. elegans*.

(**A**) 1-oct simultaneously activates sensory neurons in the periphery that mediate both

repulsion (ASH and other) and attraction (AWC and others), such that repulsive and

attractive afferent pathways become active concurrently. These inputs are then integrated

centrally to control at least three key locomotory parameters (exit reversals, entry

reversals, and locomotory speed), and the activity state of the reverse command neurons.

(**B**) An analogy based on refraction on light through two prisms highlights the potential

versatility of olfactory combinatorial coding in *C. elegans*. An individual odorant activates

many chemosensory neurons to produce multiple distinct afferent inputs propagating

through the downstream circuitry simultaneously, analogous to the first prism separating

white light into its component colors (top panel, left). The motor integration circuity

formulates a locomotory response by integrating the active afferent pathways, analogous

to the second prism recombining the incoming light (top panel, right). Because chemosensory neurons have different odorant affinities, the specific outputs of the sensory array and motor integration circuit may be concentration dependent. In this study, ASH (coded in red) is a low affinity sensor, while AWC (coded in blue) is a high affinity sensor. Reducing the 1-oct concentration biases the output of the sensory array toward AWC (middle panel, left), and the motor integration circuitry uses this information to produce an attractive locomotory response (middle panel, right). This coding strategy produces a simple framework for modulation. The behavioral outcome may be influenced by simply shifting the balance of the afferent pathways, much like an optical filter placed in a light path can selectively eliminate specific wavelengths (lower panel) to change the color of the transmitted light. In this study, IAA was added to the diluted 1-oct solution, inhibiting AWC, such that the afferent pathways were re-balanced toward the ASH end of the spectrum, and repulsion was restored. This 'filtering' effect could, in principle, take place peripherally or centrally, and could subserve modulation due to the presence of other odorant molecules, monoamine/neuropeptide modulators serving as interoceptive signals, and effectors of experience-dependent plasticity.