OPTIMIZING PROTEIN EXPRESSION IN THE ONE-POT PURE SYSTEM: INSIGHTS INTO
REACTION COMPOSITION AND TRANSLATION EFFICIENCY

AUTHORS:

Yan Zhang,[1] Yanping Qiu,[2] Matas Deveikis,[3] Zachary A. Martinez,[1] Tsui-Fen Chou,[2] Paul S.
Freemont,[3] Richard M. Murray[1]

[1] Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA,
91125, USA.
[2] Proteome Exploration Laboratory, California Institute of Technology, Pasadena, CA, 91125, USA.
[3] Department of Infectious Diseases, Faculty of Medicine, Imperial College London, Exhibition
Road, South Kensington, London SW7 2AZ, UK.
11

ABSTRACT

The One-Pot PURE system for *in vitro* protein expression, which results from a co-culture and
one-step purification of 36 essential proteins to support gene transcription and translation, can
significantly improve the accessibility and affordability of PURE systems. However, replicating this
protocol to match the productivity of the traditional PURE system can take considerable time and
effort due to variability in the expression level of individual proteins. In this work, we observed
unstable PURE protein expression in two *E. coli* protein expression strains, M15/pREP4 and
BL21(DE3), and addressed this instability using catabolite repression. We identified key proteins
whose concentration in the One-Pot PURE mixture significantly impacted the reaction's protein
expression capacity. Compared to the original method using two *E. coli* protein expression strains,
we found that consolidating all expression vectors onto one BL21 (DE3) strain led to more uniform
cell growth at the time of protein induction, thereby improving the composition of critical translation
initiation factors in the purified mixture for efficient translation. We also found that variations in
commercial energy solution formulations could compensate for deficiencies in the One-Pot PURE
protein composition. Additionally, our study revealed significant differences in the translation
capacity of commercially available *E. coli* tRNAs, suggesting the potential of optimizing tRNA
composition to improve protein translation. Taken together, this work highlights the intricate
biochemical interplay influencing protein expression capacity in the One-Pot PURE system and
presents strategies to improve its robustness and productivity.

KEYWORDS

PURE system, *in vitro* protein expression, cell-free transcription-translation, synthetic biology

INTRODUCTION

In vitro protein expression harnesses living cells' core gene expression machinery to allow
transcription and translation to occur in a test tube reaction. The open nature of these expression
platforms allows direct manipulation of the reaction environment. Applications of these platforms
span high-throughput screening,[1–3] novel protein modifications,[4,5] interfacing biomolecules with
synthetic materials,[6–9] on-demand biosensing[10–12] and biomanufacturing,[13,14] and understanding
the rules of life by building it from scratch.[15–18]

Nearly all *in vitro* protein expression platforms can be categorized into two classes – crude lysate-
based and reconstituted systems. Crude lysate-based expression systems use extracts from
lysed cells, which retain most of the cell's proteome and metabolic pathways[19,20] and can thus
support gene expression from a wide range of promoters and enable higher protein expression
due to effective energy regeneration.[21] They can also be exceptionally affordable at approximately
3 cents per microliter of reaction.[22] However, the active endogenous metabolism and intact
proteome can divert critical energy sources and metabolites to side reactions, interfering with the
central reactions implemented in these systems.[20]

On the other hand, the PURE (Protein synthesis Using Recombinant Elements) system isolates
the necessary transcription, translation, and energy regeneration proteins to support gene
expression by protein purification and reconstitution. This creates a more defined reaction
environment, enabling the construction of predictive models for protein expression,[23,24] allowing
for facile removal and replacement of reaction enzymes for customization,[5] and maintaining a
minimal reaction proteome toward a self-regenerating synthetic cell.[16] However, traditional PURE
systems are expensive to purchase ($0.64 – 1.00 per microliter of reaction) and laborious to
prepare in-house, posing a significant bottleneck in the wide adoption of PURE systems.

To address these limitations, the One-Pot PURE system developed by Lavickova and Maerkl
offers a simplified alternative.[25] Instead of individually purifying each of the 36 PURE proteins, a
co-culture of the 36 *E. coli* strains, each expressing a single protein, is grown, induced for protein
expression, and purified in a single preparation. This approach significantly streamlines the PURE
system preparation and delivers comparable protein yield to commercial PURE systems at a
fraction of the cost ($0.10 per microliter of reaction).

Despite the promise of this method, replicating the One-Pot PURE protocol to achieve
comparable productivity to conventional PURE systems still takes considerable time and effort.
In this work, we document the sources of variability that result in less productive One-Pot PURE
preparations, such as expression vector instability and sub-optimal PURE protein composition for
effective protein translation. In addition to optimizing the PURE protein composition, we find that
differences in the reaction's biochemical composition can significantly impact the translation rate
and the final protein yield. Taken together, this work improves our current understanding of the
factors influencing protein expression capacity in the One-Pot PURE system to enable its robust
implementation across different labs and applications.

RESULTS AND DISCUSSION

Enhancing protein expression stability for the One-Pot PURE system through glucose-
mediated catabolite repression

We began by replicating the One-Pot PURE system using plasmids deposited on Addgene and
transforming them into the respective *E. coli* M15/pREP4 and BL21(DE3) expression strains.
Proteins in pQE30 and pQE60 expression vectors were transformed into M15/pREP4 competent
cells, and proteins in pET21 and pET15 expression vectors were transformed into BL21 (DE3)
competent cells (**Figure 1A**).

Our initial monoculture protein expression assessment revealed that several protein expression
vectors were unstable. Despite confirming plasmid sequences through whole plasmid sequencing,
we observed a loss of expression in ten proteins after growing and inducing protein expression

from saved glycerol stocks (**Figure 1B, C**). No growth defect was observed for these non-
expressing cells (**Figure SI.III.1**). This phenomenon was also observed in several expression
strains directly obtained from a collaborator, with the loss of protein expression appearing in other
strains, indicating the expression vector instability is a stochastic yet frequent event for PURE
proteins (**Figure SI.III.2**).

The expression vector instability is problematic because a previously validated set of PURE
protein expression strains can spontaneously lose protein expression ability after being saved to,
and grown from, glycerol stocks. Without addressing this, subsequent preparations of the One-
Pot PURE system can exhibit protein dropouts, leading to unproductive batches. Alternatively,
each One-Pot PURE preparation will have to be made from fresh transformations of the 36 protein
expression vectors, which is labor-intensive. Being able to stabilize the growth and expression of
PURE proteins from glycerol stocks can significantly simplify the preparation workflow and
facilitate the dissemination of these strains to other labs.

We hypothesize that leaky background protein expression from $P_{T7-lacO}$ and $P_{T5-lacO-lacO}$ promoters
from BL21(DE3) and M15/pREP4 strains, respectively, can create growth burdens, leading to the
loss of protein expression. This has been observed in another work, where background protein
expression burden led to promoter mutations, resulting in the loss of protein expression.[5] To
reduce background protein expression, we investigated whether catabolite repression, where the
addition of a preferred carbon source (glucose), could upregulate the intracellular concentration
of repressors (LacI) that block secondary carbon source utilization. [26]

With upregulated LacI repressor production and reduced background protein expression from P_{T7-}
$_{lacO}$ and $P_{T5-lacO-lacO}$ promoters, we expect more stable plasmid maintenance and protein
expression. Indeed, our results showed that by supplementing 1% w/v glucose to the growth
media until the final culture, the PURE protein expression was stably maintained even after five
passages from the seed glycerol stock (**Figure 1D**). For strains grown in catabolite repression
media, we also observed improved cell growth and higher protein expression from fresh
transformations of the 10 protein expression plasmids previously found to lose protein expression
(**Figure SI.III.3**). All subsequent One-Pot PURE batches are grown with LB media supplemented
with 1% w/v glucose.

Figure 1 (A) Schematic of protein expression and regulation in *E. coli* strains M15/pREP4 and BL21(DE3) for Dual
Strain One-Pot PURE systems. Each gene of interest is expressed under the control of either the P_{T5} or P_{T7} promoter
followed by lacO operator site(s). In the M15/pREP4 expression strain, the pREP4 plasmid constitutively expresses the
lac inhibitor, LacI, which represses protein expression by binding to the lacO sites. In BL21(DE3), endogenous LacI
represses the production of T7 RNA polymerase (RNAP) and gene expression by binding to the lacO sites. Addition of
the IPTG inducer de-represses LacI inhibition and activates gene expression. **(B)** Schematic of monoculture protein
expression workflow. Cells were grown in 96-well deep-well plates, and protein expression was activated by adding 0.1
mM IPTG inducer. Following cell growth, protein expression was assessed via SDS-PAGE. **(C)** Monoculture protein
expression assessment from cells grown in LB media. Ten strains (designated by *) were found to carry the protein
expression plasmid but lacked the protein expression capability. **(D)** Stable protein expression from cells grown in LB
with 1% w/v glucose after 5 passages, with all PURE proteins visibly expressed.

Offsets in the Dual Strain One-Pot PURE protein composition contribute to low protein production.

After resolving the protein expression issue, we proceeded to replicate the original One-Pot PURE system with BL21(DE3) and M15/pREP4 expression strains (Dual Strain One-Pot PURE) following the established protocol.[27] To assess the system's transcription and translation capacity, we used a reporter plasmid with a malachite green aptamer (MGA) to approximate mRNA concentration and destabilized enhanced green fluorescent protein (deGFP) to report on the protein yield (**Figure 2A**). Surprisingly, while the system produced a high amount of mRNA transcripts of up to 6 μM, the protein expression was capped at approximately 1.5 μM for deGFP (**Figure 2B**) for two batches of the One-Pot PURE system. This protein yield is less than a third of the reported One-Pot PURE reaction yield.[25,27] We also note the high initial malachite green mRNA concentration may come from rapid mRNA transcription that began before the microwell plate was placed into the plate reader.

Given that we have confirmed all strains are expressing the One-Pot PURE proteins with mono-culture expression assays, we hypothesized that there might be poor cell growth or protein expression in the co-culture environment that was not captured by the monoculture assays. Resolving the purified One-Pot PURE protein mixture on a gel, as done in the monoculture assays, would be challenging due to the similarity in molecular weight of many proteins. Therefore, we opted for a targeted proteomics approach to characterize the relative abundance of these proteins.

Using liquid chromatography with tandem mass spectrometry (LC-MS/MS), we mapped the detected peptides to protein abundances. After scaling for their concentrations in reactions, we compared individual proteins' relative abundances in the Dual Strain One-Pot PURE mixture to protein abundances in commercial PURE systems (**Figure 2C**). Our analysis revealed a consistent deficiency in 7 proteins in Dual Strain One-Pot PURE systems compared to the two commercial PURE systems (PURE$_A$ and PURE$_B$). These proteins are alanyl-tRNA synthetase (AlaRS), initiation factor 2 (IF2), initiation factor 3 (IF3), elongation factor-G (EF-G), elongation factor-Tu (EF-Tu), elongation factor-Ts (EF-Ts), and creatine kinase (CK).

Figure 2: (A) Schematic of the reporter plasmid used to assess transcription and translation activity, using mRNA transcript with a malachite green aptamer (MGA) and coding sequences for destabilized enhanced green fluorescent protein (deGFP) expression. The fluorescence was monitored for malachite green dye-bound MGA and folded deGFP. **(B)** The productivity of two batches of Dual Strain One-Pot PURE systems (Dual Strain A and Dual Strain B) in mRNA transcription and protein translation, measured via MGA and deGFP, respectively. The One-Pot PURE protein concentration in the reactions was varied from 1.75 to 4 mg/mL to optimize yield. The plots represent the average of reaction triplicates, with error bars indicating the standard deviations of reaction triplicates. **(C)** Comparison of Dual Strain One-Pot PURE protein abundances to protein abundances in two commercial PURE systems after scaling by their concentrations in the reaction.

Enhancing One-Pot PURE protein composition and expression capacity with a Single-Strain One-Pot PURE system

Since all seven of the consistently deficient proteins are expressed in the *E. coli* M15/pREP4 expression strain, we investigated whether these cells had growth deficiencies. From the monoculture cell growth data, we observed that M15/pREP4 cells exhibited slower growth rates during and after IPTG induction for protein expression (**Figures SI.III 1 and 2**). As a result, we transferred all M15/pREP4-expressed proteins to BL21(DE3) cells to achieve more uniform growth during and after IPTG induction. We note that a caveat in this single-strain version is that the T7 RNAP needs to be removed from the One-Pot PURE co-culture due to the anticipated genetic instability of having a P_{T7} promoter expressing T7 RNAP. As a result, we opted to purify and add T7 RNAP into One-Pot PURE reactions separately (see **MATERIALS AND METHODS**). Alternatively, T7 RNAP could be purchased commercially.

Our results showed that five out of the 16 proteins previously expressed in the M15/pREP4 cells exhibited at least a 30% improvement in expression when grown and expressed in BL21(DE3) cells, as determined by monoculture growth and protein gel assessment (**Figure SI.III 4**). This enhancement is also evident in the One-Pot PURE co-culture. Compared to the Dual Strain system, the purified PURE protein mixture from the Single Strain system has a slightly higher concentration of translation factors (24% and 26% for Dual Strain and Single Strain, respectively, **Figure 3B**). In resolving the individual protein composition using mass spectrometry, we found that translation initiation factors 2 and 3 (IF2 and IF3) and creatine kinase (CK) involved in energy regeneration (**Figure 3A**) are present at significantly higher concentrations in Single Strain One-Pot PURE. An Individual protein composition comparison of Single Strain One-Pot PURE to commercial PURE systems is also provided in **Figure SI.III 5**.

We next compared the productivity of the Dual Strain and Single Strain versions of the One-Pot PURE system. Using only the energy solution from commercial vendors and replacing PURE proteins with One-Pot PURE proteins, we observed consistent high protein production with the Single Strain One-Pot PURE system (**Figure 3C-D**).

Figure 3: (A) Comparison of the relative abundance of individual One-Pot PURE proteins in the Single Strain and Dual Strain systems. Initiation factors 2 and 3 (IF2, IF3) show the most notable increase in composition in the Single Strain One-Pot PURE system. Previously under-expressed aminoacyl tRNA-synthetases (AlaRS and PheRS) and creatine kinase (CK) also increased in composition in the Single Strain One-Pot PURE system. **(B)** Composition of commercial PURE proteins, Dual Strain One-Pot PURE proteins, and Single Strain One-Pot PURE proteins grouped by protein classes: aminoacyl tRNA synthetase (aaRS), translation factors (TL factors), T7 RNAP, energy regeneration (E Regen.). Detected proteins not part of the PURE proteins are categorized as contaminants (Contam.). Ribosomal proteins are excluded from analysis. **(C)** Schematic of the reporter plasmid P_{T7}-UTR1-deGFP expressed in PURE reactions to assess system productivity. **(D)** Productivity of Dual Strain and Single Strain One-Pot PURE reactions assembled with two commercial energy mixes. $PURE_A$ and $PURE_B$ are control reactions with 5 nM PT7-UTR1-deGFP plasmids assembled according to the manufacturer's instructions. $Dual_n$ 2.5 and 5 designate varying batches of Dual Strain One-Pot PURE proteins added at 2.5 and 5 mg/mL. Likewise, $Single_n$ 2.5 and 5 designate varying batches of Single Strain One-Pot PURE proteins added at 2.5 and 5 mg/mL. The bar graphs represent the average terminal protein yield of reaction triplicates, with error bars indicating the standard deviations of reaction triplicates.

High-yield protein expression from One-Pot PURE requires a balance between the rate of protein translation and energy regeneration.

The differences in productivity between the Dual Strain One-Pot PURE system in two commercial PURE energy mixes, referred to as $PURE_A$ and $PURE_B$, are particularly interesting. It appears that the $PURE_B$ energy mix can support more One-Pot PURE proteins in a reaction as a strategy to compensate for the lower concentration of translation factors in the purified mixture (**Figure 3D**). In reactions with the $PURE_A$ and homemade energy mixes, adding more One-Pot PURE proteins to the reaction decreases the final protein yield after an optimal concentration.

Comparing the protein translation rates of both types of One-Pot PURE systems to commercial PURE systems provides valuable insights (**Figure 4** and **Figure SI.III 6-7**). In the $PURE_A$ energy solution, the translation rates of the Dual Strain and Single Strain One-Pot PURE systems behaved as expected (**Figure 4**). Deficiencies in translation proteins in the Dual Strain One-Pot PURE system contributed to a lower translation rate over the course of the reaction. With more protein translation machinery, the Single Strain One-Pot PURE system could match or even exceed the translation rate of commercial $PURE_A$ proteins. However, higher translation rate peaks led to steeper drops in translation rate shortly after, as observed for the Single Strain One-Pot PURE system with the $PURE_A$ energy solution.

In the $PURE_B$ energy solution, when both the Dual Strain and Single Strain One-Pot PURE proteins are added to 5 mg/mL, both systems exhibited comparable protein expression levels despite the deficiency in translation factors found in the Dual Strain One-Pot PURE proteins (**Figure 4**). It is important to note that the contribution to these differences in translation rate comes from the energy solution alone (amino acids, NTPs, tRNA, creatine phosphate, and other co-factors). This result suggests that an optimal preparation of the energy solution could compensate for variations in the One-Pot PURE protein composition.

We also observed that when the translation rate from the One-Pot PURE system exceeded that of commercial PURE proteins, a steep drop in translation rate soon followed (**Figure 4**). This suggests that while faster translation can occur, it can also significantly deplete the available energy molecules (ATP and GTP) critical for translation initiation and elongation reactions. A fast translation rate that surpasses the system's energy regeneration capacity can negatively impact the final protein yield by stalling protein translation, potentially resulting in premature translation termination and lower apparent protein production.

Figure 4: (A) Schematic of One-Pot PURE reaction composition using in-house preparations of Dual Strain and Single Strain One-Pot PURE proteins with commercial energy solution according to manufacturers' protocols and 5 nM of P_{T7}-UTR1-deGFP. In the case of Single Strain One-Pot PURE reactions, 0.5 μM of purified T7 RNAP was added. **(B)** Comparison of protein translation rate for various preparations of One-Pot PURE proteins (added to the reaction at 2.5 or 5.0 mg/mL) in two commercial energy mixes. Plots represent the average of reaction triplicates. Time traces of deGFP production are provided in the **Supplementary Figures SI.III. 6-7**.

E. coli tRNA composition critically influences One-Pot PURE reaction productivity.

With an optimized Single Strain One-Pot PURE system, we proceeded to validate the protein
expression capacity with an in-house prepared energy solution. Two commercial sources of *E.*
coli tRNA were tested, one from *E. coli* strain MRE600 and one from *E. coli* strain W. The energy
solution prepared using *E. coli* MRE600 tRNA exhibited a high protein translation rate and final
protein yield (**Figure 5B**). When we replaced tRNA in the energy solution with *E. coli* Strain W
tRNA, a significant decrease in protein translation rate and yield was observed(**Figure 5C**).
Increasing *E. coli* Strain W tRNA concentration did not increase protein expression.

This finding is significant because the complexity and composition of tRNAs can be overlooked
when preparing the energy solution for PURE reactions. Differences in codon usage among
various *E. coli* strains can lead to unexpected deficiency in tRNA associated with some codons.
Without prior knowledge of the deficient tRNA, it can be difficult to codon-optimize the protein for
effective expression, leading to low protein yield despite having an otherwise productive system.

Figure 5: (A) Schematic of One-Pot PURE reaction composition using in-house prepared energy solutions with
commercial tRNAs from *E. coli* MRE600 and *E. coli* Strain W. Each reaction contains 5 nM of P_{T7}-UTR1-deGFP to
assess protein production. **(B)** Comparison of deGFP protein production and its rate of production for two batches of
Single Strain One-Pot PURE proteins (added to the reaction either at 2.5 or 5.0 mg/mL) in energy solution with 22
A_{260}U/mL *E. coli* MRE600 tRNA. Error bars represent standard deviations of reaction triplicates. Protein translation rate
plots represent the average of reaction triplicates. **(C)** Comparison of deGFP protein production and its rate of
production for two batches of Single Strain One-Pot PURE proteins (added to the reaction at 2.5 mg/mL) in energy
solution with increasing *E. coli* Strain W tRNA activity units (30 – 80 A_{260}U/mL). The plots represent the average of
reaction triplicates, and the error bars represent standard deviations of reaction triplicates. Protein translation rate plots
represent the average of reaction triplicates.

CONCLUSION

In this work, we identified several important factors that affect the expression capacity of the One-Pot PURE system. We improved the stability of One-Pot PURE protein expression through glucose catabolite repression. We consolidated all protein expression vectors from the original dual strain design to one *E. coli* BL21(DE3) strain, increasing the expression levels of critical translation initiation and energy regeneration proteins and, consequently, the productivity of the One-Pot PURE reaction. Our characterization of both dual and single-strain One-Pot PURE proteins in two commercial PURE energy solutions revealed that optimizing biochemical compositions could overcome deficiencies in translation capacity. Lastly, our in-house energy mixture preparation revealed the critical role of tRNA composition in determining the translation capacity of PURE reactions.

These findings showcase that the gene expression capacity of the PURE system is influenced by both protein and biochemical factors. At the protein level, we found that translation initiation factors need to be at a higher concentration in the One-Pot PURE protein mixture to support effective protein production (**Figure 3**). This suggests the possibility of compensating for a sub-optimal One-Pot PURE protein preparation with a purified mixture of these critical translational proteins.

At the biochemical level, we found that energy mix composition significantly impacts the One-Pot PURE translation rate and the final protein yield. Using the PURE$_A$ energy mix, the reaction proceeded with a slower protein translation rate and took over 6 hours to stabilize to the final deGFP concentration (**Figure SI.III 6**). This suggests that there may be a translation rate-limiting step imposed by the PURE$_A$ energy mix, and this limitation can make One-Pot PURE proteins with sub-optimal protein composition proceed even more slowly, resulting in lower protein yield. On the other hand, the PURE$_B$ energy mix supports a faster translation rate overall, with reactions completed in under 3 hours (**Figure SI.III 7**). This suggests the rate-limiting step observed in the PURE$_A$ energy mix may be removed in PURE$_B$ to deliver a high protein yield with a shorter reaction lifetime. Interestingly, the reaction bottleneck comes from the PURE protein concentration. When more PURE proteins were added to the PURE$_B$ energy mix, a faster translation rate and a higher protein yield were observed.

Additionally, we show that commercial preparations of *E. coli* tRNA sourced from strain MRE600 and strain W significantly impact effective protein translation (**Figure 5**). Although MRE600 is a highly preferred source for *E. coli* tRNA due to the strain's low RNase I activity,[28] there is no apparent reason behind tRNA obtained from strain W (a safe and fast-growing strain utilizing sucrose as an industrially preferred carbon source)[29] exhibiting sub-optimal translation capacity. This highlights a critical limitation and source of variability in most cell-free systems (lysate-based and PURE), which is that *E. coli* tRNA is also a complex mixture and exhibits differential concentration profiles across growth stages and host strains.[30,31] For robust control over both One-Pot PURE reaction and potentially lysate-based cell-free expression systems, a promising approach can be developing an optimal tRNA composition.[32]

Taken together, this work seeks to obtain a comprehensive understanding of the factors influencing the gene expression capacity of the One-Pot PURE system. By addressing both protein and biochemical composition, our findings offer valuable insights into optimizing not only One-Pot PURE systems but also extending to lysate-based cell-free protein synthesis, paving the way for more efficient and reliable applications in synthetic biology and biotechnology.

MATERIALS AND METHODS

Escherichia coli strains and plasmids

E. coli BL21(DE3) (Thermo Scientific) and M15/pREP4 (Qiagen) strains were used for protein
313
expression. **Supplementary Information I** contains the Addgene catalog numbers and plasmid
314
sequences for Dual-Strain One-Pot. To establish the Single Strain One-Pot, all proteins were
315
amplified via Gibson Assembly and placed onto the pET21a backbone. All sequences of cloned
316
proteins transplanted onto pET21a are provided in **Supplementary Information I**.
317

Sequence-verified plasmids (Primordium Labs, now Plasmidsaurus) were transformed into their
318
respective BL21(DE3) and M15/pREP4 expression strains and plated on Luria Broth (LB) Agar
319
plates with 100 μg/mL carbenicillin and 1% glucose. An inoculation loop was used to scrape
320
multiple colonies to inoculate overnight culture in LB with 100 μg/mL carbenicillin and 1% glucose
321
to obtain the population expression average. To facilitate inoculation of subsequent final cultures,
322
250 μL of overnight cultures were saved into frozen glycerol stocks with 250 μL of 50% glycerol
323
in 1.3 mL 96 well plates (NEST Biotech). The glycerol stock plate was sealed with an aluminum
324
film and stored at -80 °C.
325

Sequences of plasmid DNA used for reporter protein expression are provided in **Supplementary**
326
Information I. All plasmid DNA was prepared using the NucleoBond Xtra Midi kit (Macherey-
327
Nagel) and eluted in nuclease-free water.
328

Monoculture Expression assay for One-Pot PURE proteins

The protocol was adapted from Grasemann *et al.*[27] with a few modifications. An overnight culture
330
of One-Pot PURE protein expression strains was inoculated either from freshly transformed cells
331
grown on LB agar plates with carbenicillin (without or with 1% w/v glucose to exert catabolite
332
repression) or from frozen glycerol stocks using a cryo-replicator (Enzyscreen, CR1000). The
333
overnight culture was grown in 96-well deep-well plates sealed with a Breathe-Easier membrane
334
(Diversified Biotech) at 37 °C and 1000 rpm.
335

The next morning, a fresh starter culture was prepared using 4 μL of the overnight cells and 396
336
μL of LB with carbenicillin. OD600 was monitored at the time of 0.1 mM IPTG induction to initiate
337
protein expression after 2-2.5 hrs of growth and 3 hours after protein expression. 10 μL of cells
338
expressing One-Pot PURE proteins were mixed with Lammeli Buffer (Bio-Rad) and boiled at 95
339
°C for 10 minutes, and 10 μL of the mixture was loaded onto Bolt™ 4-12% Bis-Tris Protein Gels
340
(Invitrogen) along with protein ladders (SeeBlue™ Plus2 Pre-stained Protein Standard,
341
Invitrogen). Samples were run at 150 kV, 400 mAmp for 30 minutes with NuPAGE MES SDS
342
buffer (Invitrogen).
343

The gel was stained with SimplyBlue Safe Stain (Invitrogen) for 1 hour and de-stained in water
344
overnight before imaging on a ChemiDoc Imager (Bio-Rad) or a GelDoc Go Imager (Bio-Rad).
345
The obtained protein gel images were analyzed using the accompanying software Image Lab 6.1
346
for Mac (Bio-Rad).
347

One-Pot PURE reaction assembly and data acquisition

Detailed methods describing the preparation of One-Pot PURE protein mixture and energy
solution are provided in **Supplementary Information II**.

All One-Pot reactions were assembled on ice using low protein-binding microcentrifuge tubes
(Eppendorf). Each reaction is 5 μL in total volume and consists of 2 μM *E. coli* Ribosome (NEB),
5 nM reporter protein expression plasmids (either P_{T7}-MGA-UTR1-deGFP or P_{T7}-UTR1-deGFP),
2.5x Energy Mix (125 mM HEPES-KOH pH 7.6, 250 mM Potassium L-Glutamate, 5 mM ATP and
GTP at pH 7.5, 2.5 mM UTP and CTP at pH 7.5, either 56 A_{260}U/mL for *E. coli* MRE600 tRNA or
75 – 200 A_{260}U/mL for *E. coli* W tRNA, 50 mM Creatine Phosphate, 2.5 mM TCEP, 50 μM Folinic
Acid, 5 mM Spermidine, and 0.75 mM amino acids mixture at pH 7.5), 1.75 – 5 mg/mL of One-
Pot PURE proteins and 0.5 – 1 μM T7 RNAP (only for Single Strain One-Pot). For reactions using
malachite green aptamer (MGA) fluorescence as a proxy for mRNA transcripts, 10 μM of
malachite green dye was added to the reaction.

Following assembly, 5 μL of each reaction was pipetted onto a low-volume 384-well microplate
with a non-binding surface (Corning), and the reaction trajectory was measured every 5 minutes
using the BioTek Synergy Plate Reader at 37 ˚C. Fluorescence for deGFP was measured at
485/515 nm excitation/emission wavelengths at Gain 61, and fluorescence for malachite green
aptamer was measured at 610/650 nm excitation/emission wavelengths at Gain 150. The
fluorescence data for MGA and deGFP were calibrated to protein concentration data using
synthesized MGA RNA and deGFP protein standards, provided in **Supplementary Information
III**, **Figure SI.III.8**.

Acquisition and analysis of proteomics data

A brief summary of LC-MS/MS workflow is provided below. Detailed methods are provided in
Supplementary Information II.

LC-MS/MS analyses of the digested peptides were performed on an EASY-nLC 1200 (Thermo
Fisher Scientific) coupled to a Q Exactive HF hybrid quadrupole-Orbitrap mass spectrometer
(Thermo Fisher Scientific). MS2 fragmentation spectra were searched with Proteome Discoverer
SEQUEST (version 2.5; Thermo Scientific, Waltham, MA) against an *in silico* tryptic digested *E.*
coli BL21 (DE3) and *E. coli* M15 database. The relative abundance of parental peptides was
calculated by integrating the area under the curve of the MS1 peaks.

The proteomics data presented in this manuscript are from the same batch of LC-MS/MS run for
results to be comparable. The relative protein abundances for each sample are first normalized
by scaling the total sample protein abundance with the average total protein abundance for all
samples. The proteins involved in the PURE system were extracted from the entire proteome
dataset and scaled by their composition in PURE reactions (manufacturer's recommended
compositions for PURE$_A$ and PURE$_B$, 4 mg/mL for Dual Strain One-Pot PURE batches A and B
in **Figure 2**, and 2.5 mg/mL for all Dual- and Single Strain One-Pot PURE batches A and B in
Figure 3)

AUTHOR CONTRIBUTIONS

Conceptualization: Y.Z., R.M.M., P.S.F., T.C.; *Investigation:* Y.Z., Y.Q., M.D.; *Analysis:* Y.Z., Y.Q., Z.A.M.; *Writing – Original Draft:* Y.Z.; *Writing – Review and Editing:* Y.Z, M.D., R.M.M., P.S.F., Y.Q., T.C., Z.A.M.; *Visualization:* Y.Z., Z.A.M.; *Supervision:* R.M.M., P.S.F., T.C.

FUNDING

R.M.M. and Y.Z. are supported by the National Science Foundation, award MCB-2152267, and Schmidt Futures, grant number 2023-3-2-1. Y.Z. is also supported by the Caltech Presidential Postdoctoral Fellowship. P.S.F and M.D. are supported by the Engineering and the Physical Sciences Research Council (EPSRC), award EP/T013788/1. M.D. is also supported by the EPSRC Centres for Doctoral Training in Biodesign Engineering at Imperial College London.

ACKNOWLEDGEMENT

We sincerely thank the Murray Lab members, particularly Dr. Zoila Jurado and Miryong (Miki) Yun, who contributed to the early setup and discussions of the project. Dr. Jurado also prepared the P_{T7}-MGA-UTR1-deGFP plasmid [33] used in this work. We also thank Dr. Jurado, Manisha Kapasiawala, and Dr. John Marken for their review and feedback on this manuscript. We thank Prof. Lulu Qian for access to the ChemiDoc Imager used for protein gel image acquisition.

CONFLICT OF INTEREST

R.M.M. has a financial stake in Tierra Biosciences, a private company that uses bacterial lysate-based cell-free technologies for protein expression and screening.

REFERENCES

(1) Hunt, A. C.; Vögeli, B.; Hassan, A. O.; Guerrero, L.; Kightlinger, W.; Yoesep, D. J.; Krüger, A.; DeWinter, M.; Diamond, M. S.; Karim, A. S.; Jewett, M. C. A Rapid Cell-Free Expression and Screening Platform for Antibody Discovery. *Nat. Commun.* **2023**, *14* (1), 3897. https://doi.org/10.1038/s41467-023-38965-w.

(2) Swank, Z.; Maerkl, S. J. CFPU: A Cell-Free Processing Unit for High-Throughput, Automated In Vitro Circuit Characterization in Steady-State Conditions. *BioDesign Res.* **2021**, *2021*, 2968181. https://doi.org/10.34133/2021/2968181.

(3) Tabuchi, T.; Yokobayashi, Y. High-Throughput Screening of Cell-Free Riboswitches by Fluorescence-Activated Droplet Sorting. *Nucleic Acids Res.* **2022**, *50* (6), 3535–3550. https://doi.org/10.1093/nar/gkac152.

(4) Martin, R. W.; Des Soye, B. J.; Kwon, Y.-C.; Kay, J.; Davis, R. G.; Thomas, P. M.; Majewska, N. I.; Chen, C. X.; Marcum, R. D.; Weiss, M. G.; Stoddart, A. E.; Amiram, M.; Ranji Charna, A. K.; Patel, J. R.; Isaacs, F. J.; Kelleher, N. L.; Hong, S. H.; Jewett, M. C. Cell-Free Protein Synthesis from Genomically Recoded Bacteria Enables Multisite Incorporation of Noncanonical Amino Acids. *Nat. Commun.* **2018**, *9* (1), 1203. https://doi.org/10.1038/s41467-018-03469-5.

(5) Ngo, P. H. T.; Ishida, S.; Busogi, B. B.; Do, H.; Ledesma, M. A.; Kar, S.; Ellington, A. Changes in Coding and Efficiency through Modular Modifications to a One Pot PURE System for In Vitro Transcription and Translation. *ACS Synth. Biol.* **2023**, *12* (12), 3771–3777. https://doi.org/10.1021/acssynbio.3c00461.

(6) Zhang, Y.; Kojima, T.; Kim, G.-A.; McNerney, M. P.; Takayama, S.; Styczynski, M. P. Protocell Arrays for Simultaneous Detection of Diverse Analytes. *Nat. Commun.* **2021**, *12* (1), 5724. https://doi.org/10.1038/s41467-021-25989-3.

(7) Manzer, Z. A.; Ghosh, S.; Jacobs, M. L.; Krishnan, S.; Zipfel, W. R.; Piñeros, M.; Kamat, N. P.; Daniel, S. Cell-Free Synthesis of a Transmembrane Mechanosensitive Channel Protein into a Hybrid-Supported Lipid Bilayer. *ACS Appl. Bio Mater.* **2021**, *4* (4), 3101–3112. https://doi.org/10.1021/acsabm.0c01482.

(8) Thakur, M.; Breger, J. C.; Susumu, K.; Oh, E.; Spangler, J. R.; Medintz, I. L.; Walper, S. A.; Ellis, G. A. Self-Assembled Nanoparticle-Enzyme Aggregates Enhance Functional Protein Production in Pure Transcription-Translation Systems. *PLOS ONE* **2022**, *17* (3), e0265274. https://doi.org/10.1371/journal.pone.0265274.

(9) Luo, S.; Adam, D.; Giaveri, S.; Barthel, S.; Cestellos-Blanco, S.; Hege, D.; Paczia, N.; Castañeda-Losada, L.; Klose, M.; Arndt, F.; Heider, J.; Erb, T. J. ATP Production from Electricity with a New-to-Nature Electrobiological Module. *Joule* **2023**, *7* (8), 1745–1758. https://doi.org/10.1016/j.joule.2023.07.012.

(10) Jung, J. K.; Alam, K. K.; Verosloff, M. S.; Capdevila, D. A.; Desmau, M.; Clauer, P. R.; Lee, J. W.; Nguyen, P. Q.; Pastén, P. A.; Matiasek, S. J.; Gaillard, J.-F.; Giedroc, D. P.; Collins, J. J.; Lucks, J. B. Cell-Free Biosensors for Rapid Detection of Water Contaminants. *Nat. Biotechnol.* **2020**, *38* (12), 1451–1459. https://doi.org/10.1038/s41587-020-0571-7.

(11) Zhang, Y.; Steppe, P. L.; Kazman, M. W.; Styczynski, M. P. Point-of-Care Analyte Quantification and Digital Readout via Lysate-Based Cell-Free Biosensors Interfaced with Personal Glucose Monitors. *ACS Synth. Biol.* **2021**, *10* (11), 2862–2869. https://doi.org/10.1021/acssynbio.1c00282.

(12) Karlikow, M.; da Silva, S. J. R.; Guo, Y.; Cicek, S.; Krokovsky, L.; Homme, P.; Xiong, Y.; Xu, T.; Calderón-Peláez, M.-A.; Camacho-Ortega, S.; Ma, D.; de Magalhães, J. J. F.; Souza, B. N. R. F.; de Albuquerque Cabral, D. G.; Jaenes, K.; Sutyrina, P.; Ferrante, T.; Benitez, A. D.; Nipaz, V.; Ponce, P.; Rackus, D. G.; Collins, J. J.; Paiva, M.; Castellanos, J. E.; Cevallos,

V.; Green, A. A.; Ayres, C.; Pena, L.; Pardee, K. Field Validation of the Performance of Paper-
Based Tests for the Detection of the Zika and Chikungunya Viruses in Serum Samples. *Nat.*
Biomed. Eng. **2022**, *6* (3), 246–256. https://doi.org/10.1038/s41551-022-00850-0.

(13)   Pardee, K.; Slomovic, S.; Nguyen, P. Q.; Lee, J. W.; Donghia, N.; Burrill, D.; Ferrante, T.;
McSorley, F. R.; Furuta, Y.; Vernet, A.; Lewandowski, M.; Boddy, C. N.; Joshi, N. S.; Collins,
J. J. Portable, On-Demand Biomolecular Manufacturing. *Cell* **2016**, *167* (1), 248-259.e12.
https://doi.org/10.1016/j.cell.2016.09.013.

(14)   Stark, J. C.; Jaroentomeechai, T.; Moeller, T. D.; Hershewe, J. M.; Warfel, K. F.; Moricz,
B. S.; Martini, A. M.; Dubner, R. S.; Hsu, K. J.; Stevenson, T. C.; Jones, B. D.; DeLisa, M. P.;
Jewett, M. C. On-Demand Biomanufacturing of Protective Conjugate Vaccines. *Sci. Adv.*
2021, *7* (6), eabe9444. https://doi.org/10.1126/sciadv.abe9444.

(15)   Shin, J.; Jardine, P.; Noireaux, V. Genome Replication, Synthesis, and Assembly of the
Bacteriophage T7 in a Single Cell-Free Reaction. *ACS Synth. Biol.* **2012**, *1* (9), 408–413.
https://doi.org/10.1021/sb300049p.

(16)   Lavickova, B.; Laohakunakorn, N.; Maerkl, S. J. A Partially Self-Regenerating Synthetic
Cell. *Nat. Commun.* **2020**, *11* (1), 6340. https://doi.org/10.1038/s41467-020-20180-6.

(17)   Godino, E.; López, J. N.; Zarguit, I.; Doerr, A.; Jimenez, M.; Rivas, G.; Danelon, C. Cell-
Free Biogenesis of Bacterial Division Proto-Rings That Can Constrict Liposomes. *Commun.*
Biol. **2020**, *3* (1), 1–11. https://doi.org/10.1038/s42003-020-01258-9.

(18)   Okauchi, H.; Ichihashi, N. Continuous Cell-Free Replication and Evolution of Artificial
Genomic DNA in a Compartmentalized Gene Expression System. *ACS Synth. Biol.* **2021**, *10*
(12), 3507–3517. https://doi.org/10.1021/acssynbio.1c00430.

(19)   Garenne, D.; Beisel, C. L.; Noireaux, V. Characterization of the All-E. Coli Transcription-
Translation System myTXTL by Mass Spectrometry. *Rapid Commun. Mass Spectrom.* **2019**,
33 (11), 1036–1048. https://doi.org/10.1002/rcm.8438.

(20)   Miguez, A. M.; Zhang, Y.; Piorino, F.; Styczynski, M. P. Metabolic Dynamics in Escherichia
Coli-Based Cell-Free Systems. *ACS Synth. Biol.* **2021**, *10* (9), 2252–2265.
https://doi.org/10.1021/acssynbio.1c00167.

(21)   Garamella, J.; Marshall, R.; Rustad, M.; Noireaux, V. The All E. Coli TX-TL Toolbox 2.0:
A Platform for Cell-Free Synthetic Biology. *ACS Synth. Biol.* **2016**, *5* (4), 344–355.
https://doi.org/10.1021/acssynbio.5b00296.

(22)   Sun, Z. Z.; Hayes, C. A.; Shin, J.; Caschera, F.; Murray, R. M.; Noireaux, V. Protocols for
Implementing an Escherichia Coli Based TX-TL Cell-Free Expression System for Synthetic
Biology. *JoVE J. Vis. Exp.* **2013**, No. 79, e50762. https://doi.org/10.3791/50762.

(23)   Matsuura, T.; Tanimura, N.; Hosoda, K.; Yomo, T.; Shimizu, Y. Reaction Dynamics
Analysis of a Reconstituted Escherichia Coli Protein Translation System by Computational
Modeling. *Proc. Natl. Acad. Sci.* **2017**, *114* (8), E1336–E1344.
https://doi.org/10.1073/pnas.1615351114.

(24)   Jurado, Z.; Pandey, A.; Murray, R. M. A Chemical Reaction Network Model of PURE.
bioRxiv August 14, 2023, p 2023.08.14.553301. https://doi.org/10.1101/2023.08.14.553301.

(25)   Lavickova, B.; Maerkl, S. J. A Simple, Robust, and Low-Cost Method To Produce the
PURE Cell-Free System. *ACS Synth. Biol.* **2019**, *8* (2), 455–462.
https://doi.org/10.1021/acssynbio.8b00427.

(26)   Stülke, J.; Hillen, W. Carbon Catabolite Repression in Bacteria. *Curr. Opin. Microbiol.*
1999, *2* (2), 195–201. https://doi.org/10.1016/S1369-5274(99)80034-4.

(27)   Grasemann, L.; Lavickova, B.; Elizondo-Cantú, M. C.; Maerkl, S. J. OnePot PURE Cell-
Free System. *JoVE J. Vis. Exp.* **2021**, No. 172, e62625. https://doi.org/10.3791/62625.

(28)   Kurylo, C. M.; Alexander, N.; Dass, R. A.; Parks, M. M.; Altman, R. A.; Vincent, C. T.;
Mason, C. E.; Blanchard, S. C. Genome Sequence and Analysis of Escherichia Coli MRE600,

a Colicinogenic, Nonmotile Strain That Lacks RNase I and the Type I Methyltransferase, EcoKI. *Genome Biol. Evol.* **2016**, *8* (3), 742–752. https://doi.org/10.1093/gbe/evw008.

(29)   Archer, C. T.; Kim, J. F.; Jeong, H.; Park, J. H.; Vickers, C. E.; Lee, S. Y.; Nielsen, L. K. The Genome Sequence of E. Coli W (ATCC 9637): Comparative Genome Analysis and an Improved Genome-Scale Reconstruction of E. Coli. *BMC Genomics* **2011**, *12* (1), 9. https://doi.org/10.1186/1471-2164-12-9.

(30)   Dong, H.; Nilsson, L.; Kurland, C. G. Co-Variation of tRNA Abundance and Codon Usage in Escherichia Coli at Different Growth Rates. *J. Mol. Biol.* **1996**, *260* (5), 649–663. https://doi.org/10.1006/jmbi.1996.0428.

(31)   Wei, Y.; Silke, J. R.; Xia, X. An Improved Estimation of tRNA Expression to Better Elucidate the Coevolution between tRNA Abundance and Codon Usage in Bacteria. *Sci. Rep.* **2019**, *9* (1), 3184. https://doi.org/10.1038/s41598-019-39369-x.

(32)   Maheshwari, A. J.; Calles, J.; Waterton, S. K.; Endy, D. Engineering tRNA Abundances for Synthetic Cellular Systems. *Nat. Commun.* **2023**, *14* (1), 4594. https://doi.org/10.1038/s41467-023-40199-9.

(33)   Jurado, Z.; Murray, R. M. Impact of Chemical Dynamics of Commercial PURE Systems on Malachite Green Aptamer Fluorescence. bioRxiv March 26, 2024, p 2024.03.15.585317. https://doi.org/10.1101/2024.03.15.585317.