ER-GUARD: an evolutionarily conserved antioxidant defense system at ER

membranes

Zhijian Ji[1], Taruna Pandey[1], Henry de Belly[1, 2], Bingying Wang[1], Orion D. Weiner[1, 2],

Yao Tang[3], Shouhong Guang[3], Thomas D. Goddard[4], Dengke K. Ma[1, 5, 6, *]

[1] Cardiovascular Research Institute, University of California, San Francisco, San

Francisco, California, USA

[2] Department of Biochemistry and Biophysics, University of California San Francisco,

San Francisco, California, USA

[3] School of Life Sciences, Division of Life Sciences and Medicine, University of Science

and Technology of China, Hefei, Anhui, China

[4] Department of Pharmaceutical Chemistry, University of California San Francisco, San

Francisco, California, USA

[5] Department of Physiology, University of California, San Francisco, San Francisco,

California, USA

[6] Innovative Genomics Institute, University of California, Berkeley, California, USA

* Correspondence: Dengke.Ma@ucsf.edu (D.K.M.)

Abstract

Oxidative protein folding in the endoplasmic reticulum (ER) is essential for all eukaryotic cells yet generates hydrogen peroxide (H_2O_2), a reactive oxygen species (ROS). The ER-transmembrane protein that provides reducing equivalents to ER and guards the cytosol for antioxidant defense remains unidentified. Here we combine AlphaFold2-based and functional reporter screens in *C. elegans* to identify a previously uncharacterized and evolutionarily conserved protein ERGU-1 that fulfills these roles. Deleting *C. elegans* ERGU-1 causes excessive H_2O_2 and transcriptional gene up-regulation through SKN-1, homolog of mammalian antioxidant master regulator NRF2. ERGU-1 deficiency also impairs organismal reproduction and behaviors. Both *C. elegans* and human ERGU-1 proteins localize to ER membranes and form network reticulum structures. We name this system ER-GUARD, Endoplasmic Reticulum Guardian Aegis of Redox Defense. Human and *Drosophila* homologs of ERGU-1 can rescue *C. elegans* mutant phenotypes, demonstrating evolutionarily ancient and conserved functions. Together, our results reveal an ER-membrane-specific protein machinery and defense-net system ER-GUARD for peroxide detoxification and suggest a previously unknown but conserved pathway for antioxidant defense in animal cells.

Introduction

In eukaryotic cells, the endoplasmic reticulum (ER) plays a critical role in protein folding, a process essential for cell function but one that generates the reactive oxygen species hydrogen peroxide (H_2O_2)(*1–3*). The ER-residing oxidase Ero1α stoichiometrically produces one molecule of H_2O_2 per disulfide bond introduced to nascent secreted or membrane proteins. Accordingly, the H_2O_2 concentration in the ER lumen is estimated to be approximately 700 nM, whereas cytosolic H_2O_2 concentration is estimated to be 2.2 nM at steady levels (*4–6*). Low cytosolic H_2O_2 levels are maintained by various H_2O_2-buffering systems and enzymes, including H_2O_2-degrading catalases. Known families of catalase localize to peroxisomes, cytosol and mitochondria, but not ER(*7–10*). While H_2O_2 can cross membranes by slow diffusion or aquaporin-facilitated transport(*11*), it remains unknown how ER membranes are guarded against the exceptionally high levels of ER H_2O_2, whose derivatives (e.g. hydroxyl radicals via Fenton reactions) can chemically attack and damage membrane lipids and proteins.

In bacteria, a family of transmembrane proteins (DsbD/ScsB)(*12*) provide reducing equivalents for correct disulfide bond formation and peroxide reduction in the periplasm, which is considered the bacterial equivalent of eukaryotic ER. DsbD and ScsB carry multi-transmembrane segments and employ a series of cysteine pairs with varying redox potentials to relay electron transfer from cytosol to periplasm(*13*). Mammalian counterparts of DsbD and ScsB are proposed to exist yet remain unidentified(*14*). Our study addresses this gap in knowledge by identifying a previously uncharacterized and

evolutionarily conserved protein *Y87G2A.13* (named as ERGU-1, <u>ER</u> <u>gu</u>ardian of

oxidative stress) as the long-sought ER membrane-localized antioxidant protein. ERGU-

1 protects against elevated H_2O_2 and ensures proper organismal functions in *C.*

elegans. Furthermore, we reveal the structural and functional conservation of ERGU-1

homologs across various animal species, indicating an ancient and fundamental role for

the ER-GUARD system in maintaining cellular redox homeostasis.

Discovery of ERGU-1

We first performed BLASTP search in an attempt to identify potential eukaryotic

sequence homologs of DsbD and ScsB. Such amino acid sequence similarity-based

queries yielded no apparent broadly conserved eukaryotic homologs, even after

adjusting sensitivity and specificity of the search (**Fig. S1A**). We reasoned that

DsbD/ScsB and potential eukaryotic counterparts might not share primary protein

sequence similarities owing to distant evolutionary divergence. Nevertheless, eukaryotic

counterparts of DsbD/ScsB could exhibit structural and functional, rather than protein

sequence features, common for DsbD/ScsB and known families of transmembrane

oxidoreductases. These features include multi-transmembrane segments to facilitate

intramembrane electron transfer, closely spaced cysteine clusters in different

transmembrane segments within close proximity (5-10 Angstroms), and broad

evolutionary conservation in eukaryotic or multicellular organisms. Based on these

features, we sought to leverage the availability of predicted protein structures by

AlphaFold2 for nearly the entire *C. elegans* proteome(*15*) and filter through DsbD/ScsB-

like candidates for subsequent functional screens and validation in *C. elegans*.

We thus conducted a computational search of the *C. elegans* proteome to identify

DsbD/ScsB-like candidates. Using the *C. elegans* UniProt reference proteome (version

26, 19827 genes), we found all genes with at least 4 annotated transmembrane helices

(3177 genes) and further filtered to those with at least two transmembrane helices each

containing at least two cysteines separated by at most 2 intervening residues (sequence

patterns CC, CXC, or CXXC). For the 190 genes meeting these criteria, we examined

AlphaFold2 Database-predicted structures and selected those in which cysteine pairs in

two helices were no more than 10 Angstroms apart (SG to SG atom distance),

producing 53 candidate genes (**Fig. 1A, Fig. S1**). We implemented the computational

screen using Python scripts (**Supplementary Data file**) in UCSF ChimeraX(*16*).

Among the 53 candidates with AlphaFold2-predicted structures passing the above

screening criteria, we focused on the non-GPCR category with 11 hits for functional

validation in *C. elegans* (**Fig. 1A, 1B**). We performed RNA interference (RNAi) against

genes encoding these 11 hits and tested if any can activate the transcriptional *gst-*

4p::GFP reporter, which has been previously used to monitor excess oxidative stress

and elevated levels of peroxide in *C. elegans*(*17*). We found that RNAi against one

gene *Y87G2A.13* strongly activated *gst-4p::GFP* under standard laboratory conditions

without exogenous oxidative stress, while RNAi against other genes or empty vector

control did not (**Fig. 1B**). Based on the T777T vector backbone with improved RNAi

specificity and efficacy(*18*), we designed two independent RNAi constructs targeting

different coding regions of *Y87G2A.13* and obtained similar results (**Fig. 1C, 1D**). In

addition, we used CRISPR-mediated genetic deletion to generate a null allele of

Y87G2A.13 and observed similar constitutive *gst-4p::GFP* activation in *Y87G2A.13* null

mutants (**Fig. 1C, 1E**). The transcription factor SKN-1 is the *C. elegans* orthologue of

NRF2, master regulator of antioxidant responses in mammals, and mediates *gst-*

4p::GFP activation upon a variety of oxidative stresses(*19–21*). We found that RNAi

against *skn-1* abolished the constitutive *gst-4p::GFP* activation in *Y87G2A.13* null

mutants (**Fig. 1F**). N-acetyl-cysteine (NAC), a commonly used precursor for glutathione-

based antioxidants and ROS/H_2O_2 scavenger in *C. elegans*(*22–24*), also strongly

suppressed *gst-4p::GFP* activation in *Y87G2A.13* null mutants (**Fig. 1G**). Taken

together, these results show that reduced expression or genetic deficiency of

Y87G2A.13, one of the computationally identified ERGU candidates, constitutively

activates an oxidative stress response reporter. We hereafter refer to the Y87G2A.13

protein as ERGU-1, given additional lines of evidence below.

Molecular and organismal roles of ERGU-1 in antioxidant defense in *C. elegans*

As an intramembrane oxidoreductase, ScsB transfers electrons via cysteine pairs as

redox couples to reduce excess protein disulfide bonds and peroxides in bacterial

periplasm(*12*). We thus assessed similar roles of ERGU-1 in ER protein-folding stress

and H_2O_2 reduction, using the *hsp-4p::GFP* transcriptional reporter and a specific

fluorescent sensor roGFP2-Orp1 for H_2O_2, respectively(*17, 25*). We found that *ergu-1*

genetic deletion or RNAi caused constitutive activation of *hsp-4p::GFP* in the absence

of exogenous ER or protein folding stresses (**Fig. 2A**). Using *C. elegans* strains

expressing roGFP2-Orp1 to specifically monitor cytosolic levels of H_2O_2 through

ratiometric dual-color fluorescence (**Fig. 2B**), we found that *ergu-1* genetic deletion or

RNAi caused markedly higher ratios of sensor oxidation versus reduction, indicating

elevated H_2O_2 levels. An independent assay based on non-fluorescent compound

dichlorofluorescein (DCFH) that reacts with H_2O_2 to produce fluorescent 2',7'-

dichlorofluorescein (DCF) showed similarly higher levels of peroxide stress in *ergu-1*

mutants (**Fig. 2C**). Furthermore, we assayed effects of *ergu-1* null, protein-truncating

mutants or RNAi on a comprehensive panel of stress-responding reporters, cell type-

specific markers, and organelle membrane lipid sensors(*26–28*), but did not observe

apparent phenotypic consequences, except oxidative stress-related reporters *gst-*

*4*p::GFP, roGFP2-Orp1 and Grx1-roGFP2 (**Fig. S2 and S3**). These results suggest

ERGU-1 normally defends against specific peroxide-related oxidative stress, consistent

with its structural features and functional reporter regulation observed.

The AlphaFold2-predicted structure of ERGU-1 reveals two clusters of cysteine pairs in

its two close transmembrane segments (**Fig. 1B, 2D**). To examine the importance of

such cysteine pairs for the antioxidant function of ERGU-1, we obtained CRISPR-

mediated knock-in alleles converting cysteine to alanine at each site, and crossed such

cysteine mutants separately with reporter strains of *gst-4p::GFP*. We found that these

cysteine mutants caused elevated levels of *gst-4p::GFP* (**Fig. 2E, 2F**). In addition, we

found that the *ergu-1* null mutant phenotypes were rescued by transgenic

extrachromosomal arrays of wild type but not cysteine mutants of *ergu-1* (**Fig. 2G**).

These results suggest that the clusters of cysteine pairs of ERGU-1 functionally

contribute to defending against oxidative stress, supporting their roles in relaying

electron transfer across ER membranes to reduce peroxide stress from ER.

We next determined organismal roles of ERGU-1. Morphologically, the null and cysteine

knock-in *ergu-1* mutants appear grossly normal. Previous studies have implicated roles

of SKN-1 activation in several long-lived genetic mutants or by oxidative stress-induced

hormesis in promoting longevity(*19*). Unexpectedly, we found that the *ergu-1* null

mutants, with control or *skn-1* RNAi, exhibited largely normal lifespans (**Fig. S2**) under

standard laboratory conditions at 20 $^{\circ}$C fed with OP50 on FuDR-NGM plates. However,

assay quantification of reproductive capacity revealed a markedly reduced brood size of

ergu-1 null mutants (**Fig. 2H**). Consistently, the *ergu-1* cysteine mutants exhibited

similar reduction in brood sizes (**Fig. 2H**). In addition, we measured the locomotion

speed change of young adult hermaphrodites upon exogenous H_2O_2 and found that

ergu-1 null mutants exhibited a markedly occluded locomotion-slowing response to

H_2O_2 (**Fig. 2I**). These results indicate that ERGU-1 is critical for maintaining normal

organismal functions in reproduction and behavioral responses to H_2O_2.

Subcellular localization of *C. elegans* ERGU-1 at ER membrane

To elucidate the tissue distribution and subcellular localization of ERGU-1, we

constructed both a transcriptional (*ergu-1p*::GFP, fusing the *ergu-1* promoter to the

green fluorescent protein GFP) and a translational reporter (*ergu-1*p::*ergu-1*::GFP,

fusing GFP to the C-terminus of ERGU-1 under the control of its native promoter) (**Fig.**

3A). We microinjected the constructs and integrated the transgenic extrachromosomal

arrays to the genome at low copy number to ensure the faithful recapitulation of ERGU-

1's endogenous expression pattern. Both the transcriptional and translational reporters

similarly revealed *ergu-1* expression in major metabolic tissues, including the intestine,

body wall muscles, and the spermatheca (**Fig. 3B, 3C**). As expected, RNAi against

ergu-1 abolished GFP signals from the *ergu-1*p::*ergu-1*::GFP transgene (**Fig. S2A**).

Notably, the tissues expressing *ergu-1* also exhibit high metabolic activities, generating

elevated levels of H_2O_2 during protein folding, as indicated by roGFP2-Orp1 (**Fig. 2B**).

To definitively establish ERGU-1's subcellular localization, we crossed the translational

reporter with a previously well-characterized ER membrane marker SEL-1(1-

79)::mCherry::HDEL(*28*, *29*). High-resolution dual-fluorescence confocal microscopy

and line-scanning analysis revealed marked co-localization of *ergu-1*p::*ergu-1*::GFP and

SEL-1(1-79)::mCherry::HDEL in a net structure-like reticulum pattern (**Fig. 3D-I**).

ERGU-1::GFP showed particularly prominent intensities in anterior and posterior

intestinal cells, consistent with effects we observed with roGFP2-Orp1 (**Fig. 2B**). The

ER membrane localization aligns remarkably well with ERGU-1's proposed functional

role in mitigating the detrimental effects of H_2O_2 produced during protein folding within

the ER lumen. Furthermore, biochemical purification of *ergu-1*p::*ergu-1*::GFP using

GFP-trap affinity chromatography revealed a redox-sensitive oligomeric organization of

ERGU-1 (**Fig. 2J**), with monomers being prevalent under reducing conditions. Given its

strategic ER membrane localization and antioxidant function, ERGU-1 defines this

system as ER-GUARD, the Endoplasmic Reticulum Guardian Aegis of Redox Defense.

Evolutionarily conserved features of ERGU-1 family proteins

We next investigated the evolutionary conservation of ERGU-1. First, we constructed a

multiple sequence alignment and a maximum likelihood-based phylogenetic tree

encompassing ERGU-1 and its homologs from diverse animal species (**Fig. 4A**). This

analysis revealed a high degree of sequence conservation across the ERGU-1 protein

family, suggesting a potentially ancient and functionally important role. For instance, the

mouse ERGU-1 homolog Tmem161b and human TMEM161B exhibit 93.6% amino acid

percent identities, while *C. elegans* ERGU-1 and human TMEM161B exhibit 27.9%

amino acid percent identities. To experimentally test evolutionary conservation of the

ERGU-1 protein family, we sought to functionally complement the *C. elegans ergu-1*

mutant phenotype with homologs from other species. We expressed the *ergu-1*

homologs *emei* and *TMEM161B* from *Drosophila* and humans, respectively, under the

control of endogenous *C. elegans ergu-1* promoter, in the *ergu-1* mutant background

carrying the *gst-4p::GFP* reporter (**Fig. 4B**). Remarkably, we found that transgenic

expression of these homologs rescued the constitutively activated *gst-4p::GFP*

phenotypes, indicating normalized oxidative stress in *C. elegans ergu-1* mutants (**Fig.**

4C). This successful complementation across species supports the functional

conservation of ERGU-1 and its homologs within the ER-GUARD system.

To additionally explore the functional conservation of ERGU-1 homologs, we focused on

TMEM161B, which appears more closely related to ERGU-1 than its paralog

TMEM161A (**Fig. 4A**). To mirror our *C. elegans* studies, we first examined TMEM161B's

subcellular localization in mammalian cells. We expressed V5-tagged TMEM161B

cDNA under the control of a CMV promoter in U2OS human osteosarcoma cells.

Immunofluorescence staining using an anti-V5 antibody revealed a striking

colocalization with established ER organelle markers (**Fig. 4D**). Furthermore, to assess

the potential antioxidant function of TMEM161B as *C. elegans* ERGU-1, we employed

V5-tagged TMEM161B cDNA expressed in HEK293 human embryonic kidney cells.

With the H_2O_2 sensor roGFP2-Orp1, we found that cDNA expression of human

TMEM161B led to strong suppression of exogenous H_2O_2-induced oxidation of roGFP2-

Orp1 (**Fig. 4E**). We also used the latest AlphaFold3 webserver(*30*) to model various

ligand interaction with TMEM161B and ERGU-1, and identified haem-binding pockets

coordinated by a highly conserved tyrosine residue (Y454) (**Fig. 4F, 4G**). We

experimentally confirmed that biochemically purified TMEM161B, but not Y454 mutants,

bound to haem directly, while functionally the Y454 mutant ERGU-1 or TMEM161B

failed to rescue *ergu-1* null mutants (**Fig. 4H, 4I**). In addition to this highly conserved

tyrosine residue, AlphaFold2-predicted structures indicate that animal homologs of

ERGU-1 (e.g. from *C. elegans*, *Drosophila*, mice and humans) all contain closely

spaced cysteine pairs that are suitable for relaying electron transfer in coordination with

haem (**Fig. S4A**). These findings indicate that human TMEM161B recapitulates the key

features of *C. elegans* ERGU-1, including ER membrane localization and antioxidant

functions, providing further evidence for the evolutionary conservation of the ER-

GUARD system in antioxidant defense across the animal kingdom (**Fig. S4B**).

Discussion

In this study, we identify ERGU-1, a previously uncharacterized and evolutionarily

conserved protein, in a critical ER-resident antioxidant defense system in animal cells.

We used a combination of computational and functional screening to pinpoint ERGU-1

among the entire UniProt-defined *C. elegans* proteome. Determining the initial list of

ERGU candidates for functional validation exemplifies the power of AlphaFold2, an AI

for protein structure prediction. Notwithstanding known caveats of predicted structures

by AlphaFold2 as compared to experimentally determined ones in accuracy, predicted

structures can guide researchers towards promising avenues for functional studies,

leading to advances in our understanding of fundamental biological processes and

potentially paving the way for new therapeutic strategies(*30–33*). We designed the

filtering criteria for screening ERGU candidates based on empirical knowledge gained

from DsbD/ScsB and membrane oxidoreductases. Similar strategy based on modified

criteria could lead to discovering other protein functions beyond ERGU-1. More broadly,

this approach also highlights the potential of AI in accelerating new biological discovery.

Despite the known generation of H_2O_2 during oxidative protein folding, the identity of the

ER membrane-resident protein responsible for its detoxification has remained elusive.

Our work bridges this gap by identifying ERGU-1 as such a key player, which harbors

functional and structural features characteristic of transmembrane oxidoreductases.

Importantly, genetic deletion of ERGU-1 led to constitutive activation of oxidative stress-

and ER stress-response reporters, elevated cytosolic H_2O_2 levels, and organismal

phenotypes in reproduction and behaviors. These findings strongly suggest ERGU-1's

crucial role in mitigating ER-derived oxidative stress, and mechanistically by transferring

electrons via cysteine pairs across ER membranes, with ER luminal H_2O_2 as a likely

electron acceptor. As loss of ERGU-1 also activated the *hsp-4p::GFP* reporter indicative

of unfolded protein stress, it is plausible that ERGU-1 provides reducing equivalents

from cytosolic electron donors for additional ER substrates other than H_2O_2, e.g.

misfolded proteins and certain peroxide products that ScsB or DscB can act upon to

reduce (*12*, *34*, *35*). However, it is worth noting that neither ScsB nor DsbD is known to

bind to haem, unlike the *C. elegans* and human homologs of ERGU-1. We thus

postulate that eukaryotic ER-GUARD may have evolved additional haem-binding

capacity that likely enables efficient electron transfer together with cysteine pairs.

The subcellular localization of ERGU-1 at the ER membrane is well suited for its

antioxidant function. By residing at the site close to H_2O_2 generation, ERGU-1 may

efficiently neutralize it, safeguarding the ER membrane and nearby cytosol from

oxidative damage. Furthermore, the redox-sensitive nature of the tetrameric ERGU-1

organization suggests a potential regulatory mechanism for ER-GUARD function. Under

more oxidizing conditions, the tetrameric form appears to be more prevalent, potentially

enhancing ERGU-1's antioxidant activity to safeguard against excess peroxide stress.

Interestingly, our findings on ER-GUARD and its proximity to H_2O_2 generation in the ER

align with recent studies suggesting a limited role for mitochondrial ROS in causing

global oxidative stress and nuclear DNA damage(*36*). This highlights the ER as a

potential major source of cellular oxidative stress impacting nuclear membranes and the

genome. By effectively neutralizing ER-derived H_2O_2, ERGU-1 may play a critical role in

shielding the genome from oxidative damage. Furthermore, ribosomal RNA, densely

packed on ER membranes and crucial for protein synthesis, is another likely target of

such oxidative damage. It remains determined whether ERGU-1 protects against

oxidative damage in general or more specifically biases towards certain biomolecules.

In specific tissues (e.g. intestine), excess peroxide stress caused by loss of ERGU-1

may also activate multiple compensatory antioxidant pathways, including SKN-1/NRF2,

and recruit additional mechanisms to safeguard cellular and organismal functions.

The high degree of sequence conservation observed within the ERGU-1 protein family

across diverse animal species underscores its potential evolutionary ancestry and

functional importance. The successful complementation of the *C. elegans ergu-1* mutant

phenotype with homologs from *Drosophila* and humans further strengthens this notion.

Although we did not find any apparent orthologues of ERGU-1 in eukaryotic fungi, our

findings indicate evolutionary conservation of the ER-GUARD system across the animal

kingdom, emphasizing its critical and previously undescribed role in maintaining cellular

redox balance. Mutations in the *Drosophila* orthologue *emei* impair ER calcium

dynamics(*37*), whereas mutations in its vertebrate orthologue *Tmem161B* cause severe

pathological cardiac arrhythmias in mice and brain polymicrogyria in humans(*38–41*).

Our studies suggest that these previously unexplained phenotypic defects might be

mechanistically caused by defects in ER-GUARD functions and redox homeostasis.

Thus, future studies are warranted to investigate the mechanisms of action, redox

regulation of ER-GUARD, and its functional consequences at the cellular and

organismal levels that may be broadly important for human physiology and diseases.

Figures

Fig. 1. Computational and genetic identification of ERGU-1. (**A**) Schematic of

AlphaFold2-assisted computational screens. (**B**) Phenotype-driven functional screens

based on RNAi activation of *gst-4p::GFP* oxidative stress reporters. (**C**) Gene structure

of *ergu-1* showing various alleles. (**D–E**) Representative epifluorescence images

showing *gst-4p::GFP* activation in two independent *ergu-1* RNAi (D) and mutants (E)

(n=15 for each group). (**F–G**) Representative epifluorescence images showing that *gst-*

4p::GFP activation in *ergu-1* null mutants requires ROS and *skn-1* (F), which are

suppressed by NAC dose-dependently (G) (n=10 for each group), with quantification of

gst-4p::GFP fluorescence intensities under conditions indicated. *P* value was

328   determined by an unpaired *t*-test, two-tailed (comparison between two groups) or one-

329   way ANOVA (comparison between multiple groups). Scale bars, 100 µm.

Fig. 2. ERGU-1 defends against H_2O_2 and maintains organismal functions. (**A**)

Representative epifluorescence images showing increased *hsp-4p::GFP* in *ergu-1* null

mutants (n=15 for each group). (**B**) Representative confocal images showing increased

oxidation/reduction ratio of Orp1-roGFP (H_2O_2 indicator) with *ergu-1* RNAi. (**C**) DCFDA

staining showing increased DCF in null mutants (n=15 for each group). (**D**) AlphaFold2-

predicted structure revealing cysteine residues that likely mediate electron transfer

across membranes. (**E**) Representative epifluorescence images showing increased *gst-*

4p::GFP in two different cysteine pair mutants. (**F**) Quantification of *gst-4p*::GFP

fluorescence intensities in WT and two different cysteine pair mutants indicated (n=10

for each group). (**G**) Representative fluorescence images of *gst-4p::GFP* showing no

rescue of *C. elegans ergu-1* mutants by two different cysteine pair mutants of ERGU-1.

(**H**) Quantification of brood sizes in wild type, *ergu-1* null and cysteine mutants. (**I**)

Quantification of average moving speed in wild type, *ergu-1* null and above strain

treated with 10 mM H_2O_2. *P* value was determined by an unpaired *t*-test, two-tailed

(comparison between two groups) or one-way ANOVA (comparison between multiple

groups). Scale bars, 100 μm.

Fig. 3. ERGU-1 constitutes a net-like reticulum structure at ER membranes. (**A**)

Schematic of the *ergu-1*p::ERGU-1::GFP translation reporter. (**B–C**) Representative

low-mag view of *ergu-1*p::GFP (B) and *ergu-1*p::*ergu-1*::GFP (C), respectively, showing

expression in anterior and posterior intestine, body wall muscles and spermatheca.

Scale bars, 100 μm. (**D–G**), Representative high-mag confocal view of *ergu-1*p::ERGU-

1::GFP showing protein localization and expression in anterior (D) and posterior (E)

intestinal cells, body wall muscles (F) and spermatheca (G). (**H**) Representative high-

mag confocal view of ERGU-1::GFP proteins showing co-localization with the ER

membrane marker SEL-1(1-79)::mCherry::HDEL in the intestine, body wall muscles and

spermatheca. (**I**) Line scans showing co-localization. (**J**) Representative SDS-PAGE

western blots with antibodies against GFP showing formation of oligomers (likely dimers

and tetramers based on the molecular weight) by ERGU-1::GFP and reduction to

monomers by 10 mM dithiothreitol (DTT). Scale bars, 100 μm.

Fig. 4. Conserved ER localization and antioxidant roles of animal homologs of

ERGU-1. (**A**) Maximum likelihood-based phylogenetic tree of ERGU-1 family proteins.

(**B**) Schematic of the rescuing transgenic constructs and strategy. (**C**) Representative

fluorescence images of *gst-4p::GFP* showing rescue of *C. elegans ergu-1* mutants by

Drosophila and human homologs (n=10 for each rescue group). Quantification (right).

Scale bars, 100 μm. (**D**) Immunofluorescence images showing that Human

TMEM161A/B localizes to the ER membrane. Scale bars, 10 μm. (**E**) Representative

confocal fluorescence images of roGFP2-Orp1 showing reduced exogenous H_2O_2 by

human TMEM161A/B and quantification. Scale bars, 20 µm. (**F**) Hydrophobic protein

surface representation of ERGU-1 structure, with AlphaFold3-predicted haem binding

pocket labelled by circle. (**G**) Schematic of key residues for haem binding and

interactions. (**H**) Absorption spectrum of purified TMEM161B showing binding to haem.

Haem binding pocket is in high confidence region (90 > plDDT > 70). (**I**) Representative

fluorescence images and quantification of *gst-4p::GFP* showing no rescue of *C. elegans*

ergu-1 mutants by *C. elegans* or human Y454 mutants of ERGU-1 homologs. *P* value

was determined by an unpaired *t*-test, two-tailed (comparison between two groups) or

one-way ANOVA (comparison between multiple groups). Scale bars, 100 µm.

Fig. S1. AlphaFold2-based computational screen for ERGU candidates. (**A**)

BLASTP search yielding no apparent broadly conserved eukaryotic homologs of

DscB/ScsB based on protein sequence similarities, prompting a search for DscB/ScsB

homologs based on functional and structural features using AlphaFold2-predicted

structures. (**B**) Predicted structures of the 53 ERGU candidates identified. (**C**) Table

listing the annotated functions of the 11 non-GPCR ERGU candidates.

Fig. S2. Phenotypic consequences of RNAi or mutant *ergu-1*. (**A**) Representative

fluorescence images of *ergu-1p::ergu-1::GFP* showing abolished GFP signals by *ergu-1*

RNAi. (**B**) Representative fluorescence images of Vitamin B12 stress reporter *acdh-1p::*

GFP showing unaltered GFP signals by *ergu-1* RNAi. (**C**) Representative fluorescence

images of glutathione stress GSSG/GSH reporters showing elevated oxidative stress in

ergu-1 null mutants. (**D**) Representative fluorescence images of *ergu-1p::ergu-1::GFP*

showing reporter up-regulation by oxidative stressors (0.5% NaAsO$_2$). (**E**) Lifespan

curves of wild type and *ergu-1* null mutants with control or *skn-1* RNAi showing

comparable lifespans in wild type and *ergu-1* null mutants and reduction by *skn-1* RNAi.

(**F**) Lifespan curves of wild type and *ergu-1* protein-truncating mutants (*gk840471*) with

control or *skn-1* RNAi comparable lifespans in wild type and *ergu-1(gk840471)* mutants

and reduction by *skn-1* RNAi. (**G**) Quantification of brood sizes in wild type and *ergu-1*

null treated with 10 mM NAC. (**H**) Representative fluorescence images of cytoskeletal

actin ACT-1::GFP reporters showing no apparent alteration by *ergu-1* RNAi. (**I**)

Representative confocal fluorescence images showing no apparent reporter (PVD

neuron-marking GFP) alteration in *ergu-1* null mutants. (**J**) Representative confocal

fluorescence images showing no apparent reporter (neuronal nuclear GCaMP6)

alteration in *ergu-1* null mutants. (**K**) Representative confocal fluorescence images

showing no apparent reporter (secreted MANF-1::GFP) alteration by *ergu-1* RNAi. *P*

value was determined by an unpaired *t*-test, two-tailed (comparison between two

groups). (A**–**H) Scale bars, 100 μm. (I–K) Scale bars, 20 μm.

Fig. S3. Phenotypes of ERGU-1 protein-truncating *ergu-1(gk840471)* mutants.

(**A**) Representative epifluorescence images showing *gst-4p::GFP* activation in *ergu-1(gk840471)* mutants (n=15 for each group). (**B**) Quantification of *gst-4p*::GFP fluorescence intensities in WT and *ergu-1(gk840471)* mutants indicated (n=15 for each group). (**C**) Representative epifluorescence images showing *gst-4p::GFP* activation in *ergu-1(gk840471)* mutants requires *skn-1*. (**D**) Representative confocal fluorescence images showing no apparent reporter (Golgi apparatus-marking Golgi::GFP, apical and endosomal membrane-marking AKT-PH::GFP, yolk organelle-marking VIT-2::GFP and autophagosome-marking SQST-1::mOrange2) alteration in *ergu-1(gk840471)* mutants. *P* value was determined by an unpaired *t*-test, two-tailed (comparison between two groups). (A–C) Scale bars, 100 µm. (D) Scale bars, 20 µm.

Fig. S4. Roles of the conserved ERGU family proteins in anti-oxidant defense. (**A**) AlphaFold2-predicted structures showing cysteine pairs in ERGU homologs from different animal species. (**B**) Schematic model of *C. elegans* ERGU-1 roles in defending against ER peroxide and protein folding stresses and maintaining organismal functions.

Materials and Methods

C. elegans

C. elegans strains were maintained under standard laboratory conditions unless
otherwise specified. The N2 Bristol strain was used as the reference wild type. Feeding
RNAi was performed as previously described(*42*). Transgenic strains were generated
by germline transformation as described(*43*). Transgenic constructs were co-injected (at
10 - 50 ng/µl) with dominant *unc-54p*::mCherry or *myo-2p*::mCherry, and stable
extrachromosomal lines of fluorescent animals were established for UV-mediated
genome integration. Genotypes of strains used are as follows: *ergu-1(ust572, syb9153,*
syb9160), dvIs19 [gst-4p::GFP], zcIs4 [hsp-4::GFP], jrIs10 [unc119(+) rps-0p::roGFP2-
Orp1], dmaIs160 [ergu-1p:: ergu-1::gfp; unc-54p::mCherry], dmaEx701[ergu-
1p::hsTMEM161B::ergu-1utr; myo-2p::mCherry], dmaEx702 [ergu-1p::ergu-1::ergu-1utr;
myo-2p::mCherry], dmaEx704 [ergu-1p::Dm-161:: ergu-1utr; myo-2p::mCherry].

AlphaFold-assisted computational screen

A comprehensive computational screen of the *Caenorhabditis elegans* proteome was
performed to identify potential DsbD/ScsB-like proteins. The reference proteome utilized
for this analysis was the *C. elegans* UniProt reference proteome (version 26),
encompassing 19,827 genes. Initial filtering was conducted to select genes encoding
proteins with a minimum of four annotated transmembrane helices, resulting in a subset
of 3,177 genes. Subsequent filtering criteria focused on the presence of transmembrane

helices containing cysteine residues, specifically targeting helices with at least two

cysteines separated by no more than two intervening residues, corresponding to the

sequence motifs CC, CXC, or CXXC. This filtering narrowed the candidate list to 190

genes. The structural analysis was conducted using predicted protein structures from

the AlphaFold2 Database. For each of the 190 genes, the predicted structures were

examined to determine the spatial proximity of cysteine pairs located within the

transmembrane helices. Specifically, structures in which the sulfur atoms (SG) of

cysteine pairs were within 10 angstroms of each other were identified. This criterion was

met by 53 candidate genes. All computational and structural analyses were executed

using custom Python scripts, which are provided in the Supplementary file. Visualization

and further structural analyses were performed using UCSF ChimeraX software.

RNA interference (RNAi) cloning and screen

RNAi and screen for hits upregulating *gst-4p*::GFP were performed by feeding worms

with E. coli strain HT115 (DE3) expressing double-strand RNA (dsRNA) targeting

endogenous genes. Briefly, dsRNA–expressing bacteria were replicated from the

Ahringer library to LB plates containing 100 μg/ml ampicillin (BP1760-25, Fisher

Scientific) at 37 °C for 16 hrs. Single clone was picked to LB medium containing 100

μg/ml ampicillin at 37 °C for 16 hrs and positive clones (verified by bacteria PCR with

pL4440 forward and pL4440 reverse primers) were spread onto NGM plates containing

100 μg/ml ampicillin and 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG, 420322,

Millopore) for 24 hrs. Developmentally synchronized embryos from bleaching of gravid

adult hermaphrodites were plated on RNAi plates and grown at 20 °C. The worms are

collected for imaging at indicated stages.

To make the *ergu-1* RNAi-1 and RNAi-2 plasmids, RNAi-1 targeting sequences are

PCR- amplified with primers 5'-

GGCCCCCCTCGAGGTCGACAGGAGACGATGCTATAAACTATGCT -3' and 5'-

TCCACCGCGGTGGCGGCCGCCGTGACGTGTTTTCAGCTCG -3' from N2 gDNA and

subcloned into the SalI and NotI sites of a digested T777T vector with Promega T4 DNA

ligase (Promega, M1801). RNAi-2 targeting sequences are PCR- amplified with primers

5'- GGCCCCCCTCGAGGTCGACATGCCAAAATGAATGCGGGC -3' and 5'-

TCCACCGCGGTGGCGGCCGCACGTGGTGATATGCCTGGTG -3'.

Fluorescence microscopy and H_2O_2 sensor imaging

SPE confocal (Leica) and epifluorescence microscopes were used to capture

fluorescence images. Animals were randomly picked at the same stage and treated with

1 mM levamisole in M9 solution (31742-250MG, Sigma-Aldrich), aligned on a 2% agar

pad on a slide for imaging. Identical setting and conditions were used to compare

experimental groups with control. For quantification of GFP fluorescence, animals were

outlined and quantified by measuring gray values using the ImageJ software. The data

were plotted and analyzed by using GraphPad Prism10. For *jrIs10 [unc119(+) rps-*

0p::roGFP2-Orp1] strain, orp1-roGFP2 was excited sequentially at 405 and 488 nm and

emission was recorded at 500-540 nm. Fifteen images were sequentially captured at 1-

micrometer z-intervals and subsequently stacked to form a composite image.

Western blotting

For SDS-PAGE samples, stage-synchronized animals for control and experiment
groups were picked (n = 50) in 60 µl M9 buffer and lysed directly by adding 20 µl of 4x
Laemmli sample buffer (1610747, Bio-Rad) contain 10% of 2-Mercaptoethanol (M6250-
100ML, Sigma (v/v)). Protein extracts were denatured at 95 °C for 10 min and
separated on 10% SDS-PAGE gels (1610156, Bio-Rad) at 80 V for ~40 min followed by
110 V for ~70 min. The proteins were transferred to a nitrocellulose membrane
(1620094, Bio-Rad,) at 25 V for 45 mins by Trans-Blot® Turbo™ Transfer System (Bio-
Rad). The NC membrane was initially blocked with 5% non-fat milk and 2% BSA
(A4503, Sigma (v/v)) in tris buffered saline with 0.1% Tween 20 (93773, Sigma) (TBST)
at room temperature for 1 h. Proteins of interest were detected using antibodies against
GFP (A6455, Invitrogen) or V5 () in cold room for overnight. After three washes of 5 min
each with tris-buffered saline with 0.1% Tween 20, anti-mouse IgG, HRP-linked
Antibody (7076, Cell Signaling Technology) was added at a dilution of 1:5000. For DTT
treatment, worm lysates were treated with 10 mM DTT and boiled at 95 °C for 10 min.

Mammalian cell culture experiments

Human embryonic kidney (HEK) 293T cells and osteosarcoma U2OS cells were
maintained in Dulbecco's modified Eagle's medium with 10% inactive fetal bovine
serum and penicillin-streptomycin (Gibco, Grand Island, 15140122) at 37 °C supplied
with 5% CO_2 in an incubator (Thermo Fisher Scientific) with a humidified atmosphere.
Cells were washed once using PBS and digested with 0.05% trypsin EDTA (Gibco) at
37 °C for routine passage of the cells. All cells were transfected with 3 µl LipoD293™

(SignaGen, SL100668) per 1 µg DNA mixture. HEK293T cells were transfected by the

pLX304-TMEM161A-V5 (DNASU, HsCD00441633) or pLX304-TMEM161B-V5

(DNASU, HsCD00444935) and collected 2 days after transfection. HEK293T and U2OS

cells were transfected by the pHAGE2-TMEM161A-gfp/mCherry or pHAGE2-

TMEM161B-gfp/mCherry and collected for imaging 2 days after transfection. pHAGE2-

TMEM161A/B-GFP/mCherry was constructed by inserting TMEM161A or TMEM161B

into pHAGE2-gfp/mCherry plasmids. TMEM161A was PCR- amplified with the primers

5'- TAATTAAACCTCTAGAGCCACCATGGCGGTCCTCGG -3' and 5'-

CTCACCATAGCTCGAGacccagacccGGAGCCTGCCAAGTGC -3' from pLX304-

TMEM161A-V5 mentioned before. TMEM161B was PCR- amplified with the primers 5'-

TAATTAAACCTCTAGAGCCACCATGGGTGTGATAGGTATACAGC -3' and 5'-

CTCACCATAGCTCGAGacccagacccTGCCACAGTCAGATACTGGT -3' from pLX304-

TMEM161B-V5 mentioned before.

Haem binding assay

Mock plasmid, pLX304-TMEM161B-V5 and pLX304-TMEM161B Y454S-V5 expressing

HEK293T cells were collected for protein extraction by ChromoTek V5-Trap® Agarose

(proteintech, v5ta). Extracted protein were incubated with 40 mM Hemin and washed

three times for 5 min to remove unspecific binding. Samples were analyzed by UV/VIS

spectroscopy and SDS-PAGE.

Quantification of brood size

The brood size assay was carried out according to the standard protocol. (1) Briefly,

single L4 worms of different stains (N2, ergu-1 null, PHX9155, PHX9160) were

individually placed in 60mm Petri plates and kept in the incubator at 20 C. Then, the

worms were transferred to a new OP50 containing plate each day at the same time till

day 6 of the worms. Progenies were scored after 3 days and plotted for each day and

the total sum of progenies. For NAC treatment the plates were supplemented with

10mM of NAC during the entire duration of the assay. Each experiment was repeated 3

times as independent biological replicates with more than 5 animals per group.

DCFDA ROS detection

ROS staining in live worms was carried out as in standard protocol with minor

modifications. Briefly, day 1 worms were washed three times with M9 and then

transferred into 200 µl of M9 buffer containing 10 mM H2DCFDA (carboxy-H2DCFDA

[5-(and-6)-carboxy-2′,7'-dichlorodihydrofluorescein diacetate) in an 1.5 ml Eppendorf

tube and incubated in the dark for 3 hrs. The worms were randomly selected and

treated with 10 mM sodium azide (Sigma-Aldrich) in M9, symmetrically aligned on 2%

agar pads on slides for imaging the oxidized dichlorofluorescin (DCF).

Statistical analysis

Data were analyzed using GraphPad Prism 9.2.0 Software (Graphpad, San Diego, CA)

and presented as means ± S.D. unless otherwise specified, with *P* values calculated by

unpaired two-tailed t-tests (comparisons between two groups), one-way ANOVA

(comparisons across more than two groups) and adjusted with Bonferroni's corrections.

Acknowledgment

We are grateful for the feedback from and discussion with other members of the Ma lab

at UCSF and Drs. P. Sigala, P. R. Ortiz de Montellano, A. Correia, Z. Chen, B.

DeGrado, I. Jain, B. Black and G. Huang. We thank the *Caenorhabditis* Genetics Center

(NIH grant #P40 OD010440), Wormbase.org (NIH grant #U24 HG002223 to P.

Sternberg), Wormatlas.org (NIH grant #OD010943 to D.H. Hall.) and CenGen

(cengen.org) for their immensely helpful resources. The work was supported by NIH

grants (R35GM139618 to D.K.M., R35GM118167 to O.D.W), BARI Investigator Award

(D.K.M.), and UCSF PBBR New Frontier Research Award (D.K.M., O.D.W).

Author contributions

Z.J. and D.K.M. designed, performed and analyzed the *C. elegans* and cell culture

experiments, contributed to project conceptualization and wrote the manuscript. T.P.,

B.W., Y.T. and S.H.G. designed, performed and analyzed the *C. elegans* transgenic

and CRISPR experiments and editing manuscript. H.B. and Z.J. performed and

analyzed the ER imaging experiments. T.D.G conducted the computational screen and

contributed to project conceptualization and wrote the manuscript. D.K.M, O.D.W. and

T.D.G contributed to research materials, funding acquisition and editing manuscript.

Competing interests

The authors declare no competing interests.

References

1. B. P. Tu, J. S. Weissman, Oxidative protein folding in eukaryotes: mechanisms and consequences. *J. Cell Biol.* **164**, 341–346 (2004).

2. B. P. Tu, S. C. Ho-Schleyer, K. J. Travers, J. S. Weissman, Biochemical basis of oxidative protein folding in the endoplasmic reticulum. *Science* **290**, 1571–1574 (2000).

3. B. S. Mamathambika, J. C. Bardwell, Disulfide-linked protein folding pathways. *Annu Rev Cell Dev Biol* **24**, 211–235 (2008).

4. C. Gao, Y. Tian, R. Zhang, J. Jing, X. Zhang, Endoplasmic Reticulum-Directed Ratiometric Fluorescent Probe for Quantitive Detection of Basal H2O2. *Anal Chem* **89**, 12945–12950 (2017).

5. O. Lyublinskaya, F. Antunes, Measuring intracellular concentration of hydrogen peroxide with the use of genetically encoded H2O2 biosensor HyPer. *Redox Biol* **24**, 101200 (2019).

6. H. Sies, Hydrogen peroxide as a central redox signaling molecule in physiological oxidative stress: Oxidative eustress. *Redox Biol* **11**, 613–619 (2017).

7. C. Glorieux, P. B. Calderon, Catalase, a remarkable enzyme: targeting the oldest antioxidant enzyme to find a new cancer treatment approach. *Biol Chem* **398**, 1095–1108 (2017).

8. J. A. Imlay, Cellular defenses against superoxide and hydrogen peroxide. *Annu Rev Biochem* **77**, 755–776 (2008).

9. S. E. Schriner, N. J. Linford, G. M. Martin, P. Treuting, C. E. Ogburn, M. Emond, P. E. Coskun, W. Ladiges, N. Wolf, H. Van Remmen, D. C. Wallace, P. S. Rabinovitch, Extension of murine life span by overexpression of catalase targeted to mitochondria. *Science* **308**, 1909–1911 (2005).

10. R. J, B. N, H. Jm, Disulfide formation in the ER and mitochondria: two solutions to a common process. *Science (New York, N.Y.)* **324** (2009).

11. G. P. Bienert, A. L. B. Møller, K. A. Kristiansen, A. Schulz, I. M. Møller, J. K. Schjoerring, T. P. Jahn, Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *J Biol Chem* **282**, 1183–1192 (2007).

12. S.-H. Cho, D. Parsonage, C. Thurston, R. J. Dutton, L. B. Poole, J.-F. Collet, J. Beckwith, A new family of membrane electron transporters and its substrates, including a new cell envelope peroxiredoxin, reveal a broadened reductive capacity of the oxidative bacterial cell envelope. *mBio* **3**, e00291-11 (2012).

13. A. Rozhkova, R. Glockshuber, Thermodynamic aspects of DsbD-mediated electron
transport. *J Mol Biol* **380**, 783–788 (2008).

14. L. Ellgaard, C. S. Sevier, N. J. Bulleid, How Are Proteins Reduced in the
Endoplasmic Reticulum? *Trends Biochem Sci* **43**, 32–43 (2018).

15. J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K.
Tunyasuvunakool, R. Bates, A. Žídek, A. Potapenko, A. Bridgland, C. Meyer, S. A.
A. Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler,
T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M.
Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K.
Kavukcuoglu, P. Kohli, D. Hassabis, Highly accurate protein structure prediction
with AlphaFold. *Nature* **596**, 583–589 (2021).

16. E. C. Meng, T. D. Goddard, E. F. Pettersen, G. S. Couch, Z. J. Pearson, J. H.
Morris, T. E. Ferrin, UCSF ChimeraX: Tools for structure building and analysis.
Protein Sci **32**, e4792 (2023).

17. B. Leiers, A. Kampkötter, C. G. Grevelding, C. D. Link, T. E. Johnson, K. Henkle-
Dührsen, A stress-responsive glutathione S-transferase confers resistance to
oxidative stress in Caenorhabditis elegans. *Free Radic Biol Med* **34**, 1405–1415
(2003).

18. Á. Sturm, É. Saskoi, K. Tibor, N. Weinhardt, T. Vellai, Highly efficient RNAi and
Cas9-based auto-cloning systems for C. elegans research. *Nucleic Acids Res* **46**,
e105 (2018).

19. T. K. Blackwell, M. J. Steinbaugh, J. M. Hourihan, C. Y. Ewald, M. Isik, SKN-1/Nrf,
stress responses, and aging in Caenorhabditis elegans. *Free Radic Biol Med* **88**,
290–301 (2015).

20. H. Sies, C. Berndt, D. P. Jones, Oxidative Stress. *Annu Rev Biochem* **86**, 715–748
(2017).

21. F. Pa, S. Tf, B. T, B.-Z. R, G. Hk, Z. H, H.-S. R, D. A, SKN-1 regulates stress
resistance downstream of amino catabolism pathways. *iScience* **25** (2022).

22. I. Gusarov, I. Shamovsky, B. Pani, L. Gautier, S. Eremina, O. Katkova-
Zhukotskaya, A. Mironov, A. A. Makarov, E. Nudler, Dietary thiols accelerate aging
of C. elegans. *Nat Commun* **12**, 4336 (2021).

23. C. D. De Magalhaes Filho, B. Henriquez, N. E. Seah, R. M. Evans, L. R. Lapierre,
A. Dillin, Visible light reduces C. elegans longevity. *Nat Commun* **9**, 927 (2018).

24. O. I. Aruoma, B. Halliwell, B. M. Hoey, J. Butler, The antioxidant action of N-
acetylcysteine: its reaction with hydrogen peroxide, hydroxyl radical, superoxide,
and hypochlorous acid. *Free Radic Biol Med* **6**, 593–597 (1989).

25. P. Back, W. H. De Vos, G. G. Depuydt, F. Matthijssens, J. R. Vanfleteren, B. P. Braeckman, Exploring real-time in vivo redox biology of developing and aging Caenorhabditis elegans. *Free Radic Biol Med* **52**, 850–859 (2012).

26. Z. Zhang, M. Bai, G. O. Barbosa, A. Chen, Y. Wei, S. Luo, X. Wang, B. Wang, T. Tsukui, H. Li, D. Sheppard, T. B. Kornberg, D. K. Ma, Broadly conserved roles of TMEM131 family proteins in intracellular collagen assembly and secretory cargo trafficking. *Sci Adv* **6**, eaay7667 (2020).

27. D. K. Ma, R. Vozdek, N. Bhatla, H. R. Horvitz, CYSL-1 interacts with the O2-sensing hydroxylase EGL-9 to promote H2S-modulated hypoxia-induced behavioral plasticity in C. elegans. *Neuron* **73**, 925–940 (2012).

28. C. Wang, B. Wang, T. Pandey, Y. Long, J. Zhang, F. Oh, J. Sima, R. Guo, Y. Liu, C. Zhang, S. Mukherjee, M. Bassik, W. Lin, H. Deng, G. Vale, J. G. McDonald, K. Shen, D. K. Ma, A conserved megaprotein-based molecular bridge critical for lipid trafficking and cold resilience. *Nat Commun* **13**, 6805 (2022).

29. R. W. Klemm, J. P. Norton, R. A. Cole, C. S. Li, S. H. Park, M. M. Crane, L. Li, D. Jin, A. Boye-Doe, T. Y. Liu, Y. Shibata, H. Lu, T. A. Rapoport, R. V. Farese, C. Blackstone, Y. Guo, H. Y. Mak, A conserved role for atlastin GTPases in regulating lipid droplet size. *Cell Rep* **3**, 1465–1475 (2013).

30. J. Abramson, J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L. Willmore, A. J. Ballard, J. Bambrick, S. W. Bodenstein, D. A. Evans, C.-C. Hung, M. O'Neill, D. Reiman, K. Tunyasuvunakool, Z. Wu, A. Žemgulytė, E. Arvaniti, C. Beattie, O. Bertolli, A. Bridgland, A. Cherepanov, M. Congreve, A. I. Cowen-Rivers, A. Cowie, M. Figurnov, F. B. Fuchs, H. Gladman, R. Jain, Y. A. Khan, C. M. R. Low, K. Perlin, A. Potapenko, P. Savy, S. Singh, A. Stecula, A. Thillaisundaram, C. Tong, S. Yakneen, E. D. Zhong, M. Zielinski, A. Žídek, V. Bapst, P. Kohli, M. Jaderberg, D. Hassabis, J. M. Jumper, Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature*, doi: 10.1038/s41586-024-07487-w (2024).

31. L. Aithani, E. Alcaide, S. Bartunov, C. D. O. Cooper, A. S. Doré, T. J. Lane, F. Maclean, P. Rucktooa, R. A. Shaw, S. E. Skerratt, Advancing structural biology through breakthroughs in AI. *Curr Opin Struct Biol* **80**, 102601 (2023).

32. Z. Yang, X. Zeng, Y. Zhao, R. Chen, AlphaFold2 and its applications in the fields of biology and medicine. *Signal Transduct Target Ther* **8**, 115 (2023).

33. S. Mosalaganti, A. Obarska-Kosinska, M. Siggel, R. Taniguchi, B. Turoňová, C. E. Zimmerli, K. Buczak, F. H. Schmidt, E. Margiotta, M.-T. Mackmull, W. J. H. Hagen, G. Hummer, J. Kosinski, M. Beck, AI-based structure prediction empowers integrative structural analysis of human nuclear pores. *Science* **376**, eabm9506 (2022).

34. R. Ushioda, J. Hoseki, K. Araki, G. Jansen, D. Y. Thomas, K. Nagata, ERdj5 is
required as a disulfide reductase for degradation of misfolded proteins in the ER.
Science **321**, 569–572 (2008).

35. I. S. Arts, A. Gennaris, J.-F. Collet, Reducing systems protecting the bacterial cell
envelope from oxidative damage. *FEBS Lett* **589**, 1559–1568 (2015).

36. D. M. K. van Soest, P. E. Polderman, W. T. F. den Toom, J. P. Keijer, M. J. van
Roosmalen, T. M. F. Leyten, J. Lehmann, S. Zwakenberg, S. De Henau, R. van
Boxtel, B. M. T. Burgering, T. B. Dansen, Mitochondrial H2O2 release does not
directly cause damage to chromosomal DNA. *Nat Commun* **15**, 2725 (2024).

37. X. Ma, J.-Y. Lu, A. Moraru, A. A. Teleman, J. Fang, Y. Qiu, P. Liu, T. Xu, A novel
regulator of ER Ca2+ drives Hippo-mediated tumorigenesis. *Oncogene* **39**, 1378–
1387 (2020).

38. C. D. Koopman, J. De Angelis, S. P. Iyer, A. O. Verkerk, J. Da Silva, G. Berecki, A.
Jeanes, G. J. Baillie, S. Paterson, V. Uribe, O. V. Ehrlich, S. D. Robinson, L.
Garric, S. Petrou, C. Simons, I. Vetter, B. M. Hogan, T. P. de Boer, J. Bakkers, K.
A. Smith, The zebrafish grime mutant uncovers an evolutionarily conserved role for
Tmem161b in the control of cardiac rhythm. *Proc Natl Acad Sci U S A* **118**,
e2018220118 (2021).

39. S. K. Akula, A. Y. Chen, J. E. Neil, D. D. Shao, A. Mo, N. K. Hylton, S. DiTroia, V.
S. Ganesh, R. S. Smith, K. O'Kane, R. C. Yeh, J. H. Marciano, S. Kirkham, C. J.
Kenny, J. H. T. Song, M. Al Saffar, F. Millan, D. J. Harris, A. V. Murphy, K. C.
Klemp, S. R. Braddock, H. Brand, I. Wong, M. E. Talkowski, A. O'Donnell-Luria, A.
Lai, R. S. Hill, G. H. Mochida, R. N. Doan, A. J. Barkovich, E. Yang, D. Amrom, E.
Andermann, A. Poduri, C. A. Walsh, Polymicrogyria Genetics Research Network,
Exome Sequencing and the Identification of New Genes and Shared Mechanisms
in Polymicrogyria. *JAMA Neurol* **80**, 980–988 (2023).

40. S. K. Akula, J. H. Marciano, Y. Lim, D. Exposito-Alonso, N. K. Hylton, G. H. Hwang,
J. E. Neil, N. Dominado, R. K. Bunton-Stasyshyn, J. H. T. Song, M. Talukdar, A.
Schmid, L. Teboul, A. Mo, T. Shin, B. Finander, S. G. Beck, R. C. Yeh, A. Otani, X.
Qian, E. M. DeGennaro, F. S. Alkuraya, S. Maddirevula, G. D. Cascino, C.
Giannini, Undiagnosed Diseases Network, L. C. Burrage, J. A. Rosenfield, S.
Ketkar, G. D. Clark, C. Bacino, R. A. Lewis, R. A. Segal, J. F. Bazan, K. A. Smith,
J. A. Golden, G. Cho, C. A. Walsh, TMEM161B regulates cerebral cortical gyration,
Sonic Hedgehog signaling, and ciliary structure in the developing central nervous
system. *Proc Natl Acad Sci U S A* **120**, e2209964120 (2023).

41. L. Wang, C. Heffner, K. I. Vong, C. Barrows, Y.-J. Ha, S. Lee, P. Lara-Gonzalez, I.
Jhamb, D. Van Der Meer, R. Loughnan, N. Parker, D. Sievert, S. Mittal, M. Y. Issa,
O. A. Andreassen, A. Dale, W. B. Dobyns, M. S. Zaki, S. A. Murray, J. G. Gleeson,
TMEM161B modulates radial glial scaffolding in neocortical development. *Proc Natl
Acad Sci U S A* **120**, e2209983120 (2023).

42. R. S. Kamath, J. Ahringer, Genome-wide RNAi screening in Caenorhabditis
elegans. *Methods* **30**, 313–321 (2003).

43. C. C. Mello, J. M. Kramer, D. Stinchcomb, V. Ambros, Efficient gene transfer in
C.elegans: extrachromosomal maintenance and integration of transforming
sequences. *EMBO J.* **10**, 3959–3970 (1991).