AlphaFold2-Guided Functional Screens Reveal a Conserved Antioxidant Protein

at ER Membranes

Zhijian Ji[1], Taruna Pandey[1], Henry de Belly[1, 2], Jingxuan Yao[4], Bingying Wang[1], Orion

D. Weiner[1, 2], Yao Tang[3], Shouhong Guang[3], Shiya Xu[4], Zhiyong Lou[4], Thomas D.

Goddard[5], Dengke K. Ma[1, 6, 7, 8, *]

[1] Cardiovascular Research Institute, University of California San Francisco, San

Francisco, California, USA

[2] Department of Biochemistry and Biophysics, University of California San Francisco,

San Francisco, California, USA

[3] School of Life Sciences, Division of Life Sciences and Medicine, University of Science

and Technology of China, Hefei, Anhui, China

[4] MOE Key Laboratory of Protein Science, School of Medicine, Tsinghua University,

Beijing, China.

[5] Department of Pharmaceutical Chemistry, University of California San Francisco, San

Francisco, California, USA

[6] Department of Physiology, University of California, San Francisco, San Francisco,

California, USA

[7] Innovative Genomics Institute, University of California, Berkeley, California, USA

[8] Lead contact

* Correspondence: Dengke.Ma@ucsf.edu (D.K.M.)

Abstract

Oxidative protein folding in the endoplasmic reticulum (ER) is essential for all eukaryotic cells yet generates hydrogen peroxide (H_2O_2), a reactive oxygen species (ROS). The ER-transmembrane protein that provides reducing equivalents to ER and guards the cytosol for antioxidant defense remains unidentified. Here we combine AlphaFold2-based and functional reporter screens in *C. elegans* to discover a previously uncharacterized and evolutionarily conserved protein ERGU-1 that fulfills these roles. Deleting *C. elegans* ERGU-1 causes excessive H_2O_2 and transcriptional gene up-regulation through SKN-1, homolog of mammalian antioxidant master regulator NRF2. ERGU-1 deficiency also impairs organismal reproduction and behavioral responses to H_2O_2. Both *C. elegans* and human ERGU-1 proteins localize to ER membranes and form network reticulum structures. Human and *Drosophila* homologs of ERGU-1 can rescue *C. elegans* mutant phenotypes, demonstrating evolutionarily ancient and conserved functions. In addition, purified ERGU-1 and human homolog TMEM161B exhibit redox-modulated oligomeric states. Together, our results reveal an ER-membrane-specific protein machinery for peroxide detoxification and suggest a previously unknown and conserved mechanisms for antioxidant defense in animal cells.

Introduction

In eukaryotic cells, the endoplasmic reticulum (ER) plays a critical role in protein folding, a process essential for cell function but one that generates the reactive oxygen species hydrogen peroxide (H_2O_2)[1–3]. The ER-residing oxidase Ero1α stoichiometrically produces one molecule of H_2O_2 per disulfide bond introduced to nascent secreted or membrane proteins. Accordingly, the H_2O_2 concentration in the ER lumen is estimated to be approximately 700 nM, whereas cytosolic H_2O_2 concentration is estimated to be 2.2 nM at steady levels[4–6]. Low cytosolic H_2O_2 levels are maintained by various H_2O_2-buffering systems and enzymes, including H_2O_2-degrading catalases. Known families of catalase localize to peroxisomes, cytosol and mitochondria, but not ER[7–10]. While H_2O_2 can cross membranes by slow diffusion or aquaporin-facilitated transport[11], it remains unknown how ER membranes are guarded against the exceptionally high levels of ER H_2O_2, whose derivatives (e.g. hydroxyl radicals via Fenton reactions) can chemically attack and damage membrane lipids, nucleic acids and proteins.

In bacteria, a family of transmembrane proteins (DsbD/ScsB)[12] provide reducing equivalents for correct disulfide bond formation and peroxide reduction in the periplasm, which is considered the bacterial equivalent of eukaryotic ER. DsbD and ScsB carry multi-transmembrane segments and employ a series of cysteine pairs with differential redox potentials to relay electron transfer from cytosol to periplasm[13]. Mammalian counterparts of DsbD and ScsB are proposed to exist yet remain unidentified[14]. Our study addresses this gap in knowledge by identifying a previously uncharacterized and

evolutionarily conserved *C. elegans* protein *Y87G2A.13* (named as ERGU-1, <u>ER</u>

<u>gu</u>ardian of oxidative stress) as the long-sought ER membrane-localized antioxidant

protein. ERGU-1 protects against elevated H_2O_2 and ensures optimal organismal

functions in *C. elegans*. Furthermore, we reveal the structural and functional

conservation of ERGU-1 homologs across various animal species, indicating an ancient

and fundamental role for ERGU-1 in maintaining cellular redox homeostasis.

Results

Computational and genetic discovery of ERGU-1

We first performed BLASTP search to identify potential eukaryotic sequence homologs

of DsbD and ScsB. Such amino acid sequence similarity-based queries yielded no

apparent broadly conserved eukaryotic homologs, even after adjusting sensitivity and

specificity of the search (Fig. S1). We reasoned that DsbD/ScsB and potential

eukaryotic counterparts might not share primary protein sequence similarities owing to

distant evolutionary divergence. Nevertheless, eukaryotic counterparts of DsbD/ScsB

could exhibit structural and functional, rather than protein sequence features, common

for DsbD/ScsB and known families of transmembrane oxidoreductases. These features

include multi-transmembrane segments to facilitate intramembrane electron transfer,

closely spaced cysteine clusters in different transmembrane segments within proximity

(5-10 Angstroms), and broad evolutionary conservation in eukaryotic or multicellular

organisms. Based on these features, we sought to leverage the availability of predicted

protein structures by AlphaFold2 for nearly the entire *C. elegans* proteome[15] and filter

through DsbD/ScsB-like candidates for subsequent functional screens and validation in

C. elegans.

To identify DsbD/ScsB-like candidates, we conducted a computational search of the *C.*

elegans proteome based on the functional features common for DsbD/ScsB-like

transmembrane oxidoreductases. Using the *C. elegans* UniProt reference proteome as

a starting point (version 26, 19,827 proteins), we found all genes with at least 4

annotated transmembrane helices (3,177 proteins) and further filtered to those with at

least two transmembrane helices, each containing at least two cysteines separated by

at most 2 intervening residues (sequence patterns CC, CXC, or CXXC). For the 190

proteins meeting these criteria, we examined AlphaFold2 Database-predicted structures

and selected those in which cysteine pairs in two helices were no more than 10

Angstroms apart (SG to SG atom distance), producing 53 candidate genes (Fig. 1A and

Fig. S1). We implemented the computational screen using customized Python scripts

(Supplementary Data file) in UCSF ChimeraX[16].

Among the 53 candidates with AlphaFold2-predicted structures passing the above

screening criteria, we focused on the non-GPCR category with 11 hits for functional

validation in *C. elegans* (Fig. 1A, B). We performed RNA interference (RNAi) against

genes encoding these 11 hits and tested if any can activate the transcriptional *gst-*

4p::GFP reporter, which has been previously used to monitor excess oxidative stress

and elevated levels of peroxide in *C. elegans*[17]. We found that RNAi against one gene

Y87G2A.13 strongly activated *gst-4p::GFP* under standard laboratory conditions without

exogenous oxidative stress, while RNAi against other genes or empty vector control did

not (Fig. 1B). Based on the T777T vector backbone with improved RNAi specificity and

efficacy[18], we designed two independent RNAi constructs targeting different coding

regions of *Y87G2A.13* and obtained similar results (Fig. 1C, D). In addition, we used

CRISPR-mediated genetic deletion to generate a null allele of *Y87G2A.13* and

observed similar constitutive *gst-4p*::GFP activation in *Y87G2A.13* null mutants (Fig.

1C, E). We also observed similar *gst-4p*::GFP activation in *ergu-1(gk840471)* mutants

with a protein-truncating mutation (Fig 1C, Fig. S2A, B). The transcription factor SKN-1

is the *C. elegans* ortholog of NRF2, master regulator of antioxidant responses in

mammals, and mediates *gst-4p::GFP* activation upon a variety of oxidative stresses[19–

21]. We found that RNAi against *skn-1* abolished the constitutive *gst-4p::GFP* activation

in *Y87G2A.13* null and protein-truncating mutants (Fig. 1F, Fig. S2C). N-acetyl-cysteine

(NAC), a commonly used precursor for glutathione-based antioxidants and ROS/H_2O_2

scavenger in *C. elegans*[22–24], also strongly suppressed *gst-4p::GFP* activation in

Y87G2A.13 null mutants (Fig. 1G). Taken together, these results show that reduced

expression by RNAi or genetic deficiency of *Y87G2A.13*, one of the computationally

identified ERGU candidates, constitutively activates oxidative stress response in a

manner that involves excess oxidants and activation of SKN-1. We hereafter refer to the

Y87G2A.13 protein as ERGU-1 (<u>ER</u> <u>gu</u>ardian of redox defense), given additional lines

of evidence below.

Molecular and organismal roles of ERGU-1 in antioxidant defense in *C. elegans*

As intramembrane oxidoreductases, DsbD and ScsB transfer electrons via cysteine

pairs as redox couples to reduce excess protein disulfide bonds and peroxides in

bacterial periplasm[12]. We thus assessed similar roles of ERGU-1 in ER protein-folding

stress and H_2O_2 reduction, using the *hsp-4p::GFP* transcriptional reporter and a specific

fluorescent sensor roGFP2-Orp1 for H_2O_2, respectively[17,25]. We found that *ergu-1*

genetic deletion or RNAi caused constitutive activation of *hsp-4p::GFP* in the absence

of exogenous ER or protein folding stresses (Fig. 2A, Fig. S3A). Using *C. elegans*

strains expressing roGFP2-Orp1 to specifically monitor cytosolic levels of H_2O_2 through

ratiometric dual-color fluorescence (Fig. 2B), we found that *ergu-1* genetic deletion or

RNAi caused markedly higher ratios of sensor oxidation versus reduction, indicating

elevated H_2O_2 levels. An independent assay based on non-fluorescent compound

dichlorofluorescein (DCFH) that reacts with H_2O_2 to produce fluorescent 2',7'-

dichlorofluorescein (DCF) showed similarly higher levels of peroxide stress in *ergu-1*

mutants (Fig. 2C). Furthermore, we assayed effects of *ergu-1* null, protein-truncating

mutants or RNAi on a comprehensive panel of stress-responding reporters, cell type-

specific markers, and organelle membrane lipid sensors[26–28], but did not observe

apparent phenotypic consequences, except oxidative stress-related reporters *gst-*

4p::GFP, *hsp-4p*::GFP, roGFP2-Orp1 and Grx1-roGFP2 (Fig. S2D, 3A-3M). These

results suggest ERGU-1 normally defends against specific peroxide-related oxidative

stress, consistent with its structural features and functional reporter regulation observed.

The AlphaFold2-predicted structure of ERGU-1 reveals two clusters of cysteine pairs in

its two close transmembrane segments (Fig. 1B, 3D). To examine the importance of

such cysteine pairs for the antioxidant function of ERGU-1, we obtained CRISPR-

mediated knock-in alleles converting cysteine to alanine at each site, and crossed such

cysteine mutants separately with reporter strains of *gst-4p::GFP*. We found that these

cysteine mutants caused elevated levels of *gst-4p::GFP* (Fig. 2E, F). In addition, we

found that the *ergu-1* null mutant phenotypes were rescued by transgenic

extrachromosomal arrays of wild-type but not cysteine mutants of *ergu-1* (Fig. S4A).

These results suggest that the clusters of cysteine pairs of ERGU-1 functionally

contribute to defending against oxidative stress, supporting their roles in relaying

electron transfer across ER membranes to reduce peroxide stress and excess H_2O_2

from ER.

We next determined organismal roles of ERGU-1. Morphologically, the null and cysteine

knock-in *ergu-1* mutants appear grossly normal. Previous studies have implicated roles

of SKN-1 activation in several long-lived genetic mutants or by oxidative stress-induced

hormesis in promoting longevity[19]. Unexpectedly, we found that the *ergu-1* null mutants,

with control or *skn-1* RNAi, exhibited largely normal lifespans (Fig. S2H) under standard

laboratory conditions at 20 ^{o}C fed with OP50 on FuDR-NGM plates. To test genetic

interaction of *ergu-1* with other known genes implicated in ER-specific redox regulation,

we examined effects of RNAi against genes in the *ero-1, gcs-1, gsr-1, gpx* and *pdi*

families and found that *ergu-1* mutants showed increased oxidative stress responses

compared with wild type upon RNAi treatment (Fig. 2G). Notably, assay quantification of

reproductive capacity revealed a markedly reduced brood size of *ergu-1* null mutants.

Consistently, the *ergu-1* cysteine mutants exhibited similar reduction in brood sizes (Fig.

2H). This reduced brood size phenotype can be partially alleviated upon 10 mM NAC

antioxidant treatment (Fig. 2H). By contrast, treatment with exogenous oxidative

stressor paraquat decreased brood size more severely in *ergu-1* mutants than wild type

(Fig. S3I). In addition, we measured the locomotion speed change of young adult

hermaphrodites upon exogenous H_2O_2 and found that *ergu-1* null mutants exhibited a

markedly occluded locomotion-slowing response to H_2O_2, without apparently affecting

neuronal development, cytoskeletal structure or baseline neuronal activity (Fig. S3J-

S3N). These results indicate that ERGU-1 is critical for maintaining normal specific

organismal functions in reproduction and behavioral responses to H_2O_2.

Subcellular localization of *C. elegans* ERGU-1 at ER membrane

To elucidate the tissue distribution and subcellular localization of ERGU-1, we

constructed both a transcriptional (*ergu-1p*::GFP, fusing the *ergu-1* promoter to the

green fluorescent protein GFP) and a translational reporter (*ergu-1*p::*ergu-1*::GFP,

fusing GFP to the C-terminus of ERGU-1 under the control of its native promoter) (Fig.

3A). We microinjected the constructs and integrated the transgenic extrachromosomal

arrays to the genome at low copy number to ensure the faithful recapitulation of ERGU-

1's endogenous expression pattern. Both the transcriptional and translational reporters

similarly revealed *ergu-1* expression in major metabolic tissues, including the intestine,

body wall muscles, and the spermatheca (Fig. 3B, C). As expected, RNAi against *ergu-*

1 abolished GFP signals from the *ergu-1*p::*ergu-1*::GFP transgene (Fig. S3B). Notably,

the tissues expressing *ergu-1* are also known to exhibit high metabolic activities,

generating elevated levels of H_2O_2 during protein folding, as indicated by roGFP2-Orp1

(Fig. 2B).

To ascertain ERGU-1's subcellular localization, we crossed the translational reporter

with a previously well-characterized ER membrane marker SEL-1(1-

79)::mCherry::HDEL[28,29]. High-resolution dual-fluorescence confocal microscopy and

line-scanning analysis revealed marked co-localization of *ergu-1*p::*ergu-1*::GFP and

SEL-1(1-79)::mCherry::HDEL in a net structure-like reticulum pattern (Fig. 3D-3I, Fig.

S5A). ERGU-1::GFP showed particularly prominent intensities in anterior and posterior

intestinal cells, consistent with effects we observed with roGFP2-Orp1 (Fig. 2B). We

confirmed the ER membrane-specific localization of ERGU-1 by imaging ERGU-1::GFP

alongside RFP markers specific for other organelles, including the mitochondria,

lysosomes, and plasma membrane (Fig. S5B). The ER membrane localization aligns

markedly well with ERGU-1's proposed functional role in mitigating the detrimental

effects of H_2O_2 produced during protein folding within the ER lumen. To address

potential ER membrane-localized changes in oxidative stresses, we generated ER-

proximal redox sensor strain GRX1-roGFP-SPSC-1. We observed markedly higher

oxidation to reduction ratio of roGFP upon *ergu-1* and *nduf-7* (positive control) RNAi

treatment (Fig. 3J, 3K). Furthermore, biochemical purification of *ergu-1*p::*ergu-1*::GFP

using GFP-trap affinity chromatography revealed a redox-sensitive oligomeric

organization of ERGU-1 (Fig. 4F), with monomers being prevalent under reducing

conditions. The ER membrane-specific localization, redox regulation and antioxidant

function support ERGU-1 as a guardian against peroxide stress originating from ER.

Evolutionarily conserved features of ERGU-1 family proteins

We next investigated the evolutionary conservation of ERGU-1. First, we constructed a

multiple sequence alignment and a maximum likelihood-based phylogenetic tree

encompassing ERGU-1 and its homologs from diverse animal species (Fig. 4A). This

analysis revealed a high degree of sequence conservation across the ERGU-1 protein

family, suggesting a potentially ancient and functionally important role. For instance, the

mouse ERGU-1 homolog Tmem161b and human TMEM161B exhibit 93.6% amino acid

percent identities, while *C. elegans* ERGU-1 and human TMEM161B exhibit 27.9%

amino acid percent identities. To experimentally test evolutionary conservation of the

ERGU-1 protein family, we sought to functionally complement the *C. elegans ergu-1*

mutant phenotype with homologs from other species. We expressed the *ergu-1*

homologs *emei* and *TMEM161B* from *Drosophila* and humans, respectively, under the

control of endogenous *C. elegans ergu-1* promoter, in the *ergu-1* mutant background

carrying the *gst-4p::GFP* reporter (Fig. 4B). Remarkably, we found that transgenic

expression of these homologs rescued the constitutively activated *gst-4p::GFP*

phenotypes, indicating normalized oxidative stress in *C. elegans ergu-1* mutants (Fig.

4C). This successful complementation across species supports the functional

conservation of ERGU-1 and its homologs within the ER-GUARD system.

To additionally explore the functional conservation of ERGU-1 homologs, we focused on

TMEM161B, which appears more closely related to ERGU-1 than its paralog

TMEM161A (Fig. 4A). To mirror our *C. elegans* studies, we first examined TMEM161B's

subcellular localization in mammalian cells. We expressed V5-tagged TMEM161B

cDNA under the control of a CMV promoter in HEK293T and U2OS human

osteosarcoma cells. Immunofluorescence staining using an anti-V5 antibody revealed a

striking colocalization with established ER organelle markers (Fig. 4D, Fig. S6F).

Furthermore, to assess the potential antioxidant function of TMEM161B as *C. elegans*

ERGU-1, we employed V5-tagged TMEM161B cDNA expressed in HEK293 human

embryonic kidney cells. With the H_2O_2 sensor roGFP2-Orp1, we found that cDNA

expression of human TMEM161B led to strong suppression of exogenous H_2O_2-induced

oxidation of roGFP2-Orp1 (Fig. 4E, Fig. S6G). We also used the latest AlphaFold3

webserver[30] to model various ligand interaction with TMEM161B and ERGU-1, and

identified heme-binding pockets coordinated by a highly conserved tyrosine residue

(Y454) (Fig. S4B). We experimentally confirmed that biochemically purified TMEM161B,

but not Y454 mutants, bound to heme directly, while functionally the Y454 mutant

ERGU-1 or TMEM161B failed to rescue *ergu-1* null mutants (Fig. S4C). Beyond the

redox-sensitive oligomeric organization of ERGU-1 (Fig. 4F), purified human

TMEM161B exhibited native oligomerization under normal conditions and dissociation

into monomers in a reducing environment, as demonstrated by size exclusion

chromatography (Fig. 4G). Interestingly, cryo-electron microscopy analysis revealed a

filamentary oligomeric form of TMEM161B under non-reducing conditions, while

treatment with DTT induced monomerization (Fig. 4H). These findings indicate that

human TMEM161B recapitulates several key features of *C. elegans* ERGU-1, including

ER membrane localization, antioxidant functions and redox-modulated oligomeric

states, providing further evidence for the evolutionary conservation of ERGU-1 in

antioxidant defense across the animal kingdom.

Discussion

In this study, we identify ERGU-1, a previously uncharacterized and evolutionarily

conserved protein, in a critical ER-resident antioxidant defense system in animal cells.

We used a combination of computational and functional screening to pinpoint ERGU-1

among the entire UniProt-defined *C. elegans* proteome. Determining the initial list of

ERGU candidates for functional validation exemplifies the power of AlphaFold2, an AI

for protein structure prediction. Notwithstanding known caveats of predicted structures

by AlphaFold2 as compared to experimentally determined ones in accuracy, predicted

structures can guide researchers towards promising avenues for functional studies,

leading to advances in our understanding of fundamental biological processes and

potentially paving the way for new therapeutic strategies[30–33]. We designed the filtering

criteria for screening ERGU candidates based on empirical knowledge gained from

DsbD/ScsB and membrane oxidoreductases. Similar strategy based on modified criteria

could lead to discovering other protein functions beyond ERGU-1. More broadly, this

approach also highlights the potential of AI in accelerating new biological discovery.

Despite the known generation of H_2O_2 during oxidative protein folding, the identity of the

ER membrane-resident protein responsible for its detoxification has remained elusive.

Our work bridges this gap by identifying ERGU-1 as such a key player, which harbors

functional and structural features characteristic of transmembrane oxidoreductases.

Importantly, genetic deletion of ERGU-1 led to constitutive activation of oxidative stress-

and ER stress-response reporters, elevated cytosolic H_2O_2 levels, and organismal

phenotypes in reproduction and behaviors. These findings strongly suggest ERGU-1's

crucial role in mitigating ER-derived oxidative stress, and mechanistically by transferring

electrons via cysteine pairs across ER membranes, with ER luminal H_2O_2 as a likely

electron acceptor. As loss of ERGU-1 also activated the *hsp-4p::GFP* reporter indicative

of unfolded protein stress, it is plausible that ERGU-1 provides reducing equivalents

from cytosolic electron donors for additional ER substrates other than H_2O_2, e.g.

misfolded proteins and certain peroxide products that ScsB or DscB can act upon to

reduce[12,34,35]. However, it is worth noting that neither ScsB nor DsbD is known to bind to

heme, unlike the *C. elegans* and human homologs of ERGU-1. We thus postulate that

eukaryotic ER-GUARD may have evolved additional heme-binding capacity that likely

enables efficient electron transfer together with cysteine pairs.

The subcellular localization of ERGU-1 at the ER membrane is well suited for its

antioxidant function. By residing at the site close to H_2O_2 generation, ERGU-1 may

efficiently neutralize it, safeguarding the ER membrane and nearby cytosol from

oxidative damage. Furthermore, the redox-sensitive nature of the tetrameric ERGU-1

organization suggests a potential regulatory mechanism for ER-GUARD function. Under

more oxidizing conditions, the tetrameric form appears to be more prevalent, potentially

enhancing ERGU-1's antioxidant activity to safeguard against excess peroxide stress.

Interestingly, our findings on ER-GUARD and its proximity to H_2O_2 generation in the ER

align with recent studies suggesting a limited role for mitochondrial ROS in causing

global oxidative stress and nuclear DNA damage[36]. This highlights the ER as a potential

major source of cellular oxidative stress impacting nuclear membranes and the genome.

By effectively neutralizing ER-derived H_2O_2, ERGU-1 may play a critical role in shielding

the genome from oxidative damage. Furthermore, ribosomal RNA, densely packed on

ER membranes and crucial for protein synthesis, is another likely target of such

oxidative damage. It remains determined whether ERGU-1 protects against oxidative

damage in general or more specifically biases towards certain biomolecules. In specific

tissues (e.g. intestine), excess peroxide stress caused by loss of ERGU-1 may also

activate multiple compensatory antioxidant pathways, including SKN-1/NRF2, and

recruit additional mechanisms to safeguard cellular and organismal functions.

The high degree of sequence conservation observed within the ERGU-1 protein family

across diverse animal species underscores its potential evolutionary ancestry and

functional importance. The successful complementation of the *C. elegans ergu-1* mutant

phenotype with homologs from *Drosophila* and humans further strengthens this notion.

Although we did not find any apparent orthologues of ERGU-1 in eukaryotic fungi, our

findings indicate evolutionary conservation of the ER-GUARD across the animal

kingdom, emphasizing its critical and previously undescribed role in maintaining cellular

redox balance. Mutations in the *Drosophila* orthologue *emei* impair ER calcium

dynamics[37], whereas mutations in its vertebrate orthologue *Tmem161B* cause severe

pathological cardiac arrhythmias in mice and brain polymicrogyria in humans[38–41]. Our

studies suggest that these previously unexplained phenotypic defects might be

mechanistically caused by defects in ER-GUARD functions and redox homeostasis.

Thus, future studies are warranted to investigate the mechanisms of action, redox

regulation of ER-GUARD, and its functional consequences at the cellular and

organismal levels that may be broadly important for human physiology and diseases.

Figures

Fig. 1. Computational and genetic identification of ERGU-1. (**A**) Schematic of

AlphaFold2-assisted computational screens. (**B**) Phenotype-driven functional screens

based on RNAi activation of *gst-4p::GFP* oxidative stress reporters. (**C**) Gene structure

of *ergu-1* showing various alleles. (**D** and **E**) Representative epifluorescence images

showing *gst-4p::GFP* activation in two independent *ergu-1* RNAi (D) and mutants (E)

(n=15 for each group). (**F** and **G**) Representative epifluorescence images showing that

gst-4p::GFP activation in *ergu-1* null mutants requires ROS and *skn-1* (F), which are

suppressed by NAC dose-dependently (G) (n=10 for each group), with quantification of

gst-4p::GFP fluorescence intensities under conditions indicated. *P* value was

determined by an unpaired *t*-test, two-tailed (comparison between two groups) or one-

way ANOVA (comparison between multiple groups). Scale bars, 100 µm.

Fig. 2. ERGU-1 defends against H₂O₂ and maintains organismal functions. (**A**)

Representative epifluorescence images showing increased *hsp-4p::GFP* in *ergu-1* null

mutants (n=15 for each group). (**B**) Representative confocal images showing increased

oxidation/reduction ratio of Orp1-roGFP (H₂O₂ indicator) with *ergu-1* RNAi. (**C**) DCFDA

staining showing increased DCF in null mutants (n=15 for each group). (**D**) AlphaFold2-

predicted structure revealing cysteine residues that likely mediate electron transfer

across membranes. (**E**) Representative epifluorescence images showing increased *gst-*

4p::GFP in two different cysteine pair mutants. (**F**) Quantification of *gst-4p*::GFP

fluorescence intensities in WT and two different cysteine pair mutants indicated (n=10

for each group). (**G**) Fluorescence intensity of *gst-4p*::GFP in wild type (WT) and *ergu-1*

null mutants under control (Ctrl) and RNAi conditions targeting redox regulators,

including *ero-1*, *pdi-2*, *pdi-6*, *gcs-1*, *gsr-1*, *gpx-1*, *gpx-3*, *gpx-4*, *gpx-5*, *gpx-6*, *gpx-7*, and

manf-1. (**H**) Quantification of brood sizes in wild type, *ergu-1* null and cysteine mutants

(left panel). Comparison of brood sizes between wild-type and mutant strains under

mock and 10 mM NAC treatments (right panel). *P* value was determined by an unpaired

t-test, two-tailed (comparison between two groups) or one-way ANOVA (comparison

between multiple groups). [(A), (B) and (E)] Scale bars, 100 μm. (C) Scale bars, 20 μm.

Fig. 3. ERGU-1 constitutes a net-like reticulum structure at ER membranes. (**A**)

Schematic of the *ergu-1*p::ERGU-1::GFP translation reporter. (**B** and **C**) Representative

low-mag view of *ergu-1*p::GFP (B) and *ergu-1*p::*ergu-1*::GFP (C) respectively, showing

expression in anterior and posterior intestine, body wall muscles and spermatheca.

Scale bars, 100 µm. (**D** to **G**) Representative high-mag confocal view of *ergu-*

*1*p::ERGU-1::GFP showing protein localization and expression in anterior (D) and

posterior (E) intestinal cells, body wall muscles (F) and spermatheca (G). (**H**)

Representative high-mag confocal view of ERGU-1::GFP proteins showing co-

localization with the ER membrane marker SEL-1(1-79)::mCherry::HDEL in the

intestine, body wall muscles and spermatheca. (**I**) Line scans showing co-localization.

Scale bars, 100 µm. (**J**) Schematic design of the ER-proximal H_2O_2 sensor GRX1-

roGFP-SPSC-1. (**K**) Representative ratio metric images and quantification of elevated

H_2O_2 levels in the cytosol near the ER under *ergu-1* and *nduf-7* RNAi treatment, as

indicated by the ER-proximal H_2O_2 sensor. Scale bars, 100 µm.

Fig. 4. Conserved ER localization and antioxidant roles of ERGU-1 homologs. (**A**)

Maximum likelihood-based phylogenetic tree of ERGU-1 family proteins. (**B**) Schematic

representation of transgenic constructs and strategy to rescue *ergu-1* null phenotype.

(**C**) Representative fluorescence images of *gst-4p::GFP* showing rescue of *C. elegans*

ergu-1 mutants by *Drosophila* and human homologs (n=10 for each rescue group).

Quantification (right). Scale bars, 100 μm. (**D**) Immunofluorescence images showing

that Human TMEM161A/B localizes to the ER membrane in 293T cells. Scale bars, 10

μm. (**E**) Reduction of cytosolic H_2O_2 detected by roGFP2-Orp1 in 293T cells

overexpressing human TMEM161A or TMEM161B. (**F**) Representative SDS-PAGE

western blots with antibodies against GFP showing formation of oligomers (likely dimers

and tetramers based on the molecular weight) by ERGU-1::GFP and reduction to

monomers by 10 mM dithiothreitol (DTT). (**G**) Chromatography plot showing reduction

of purified human TMEM161B to smaller size peaks in the presence of 5 mM DTT as

analyzed by SEC. (**H**) Reduction of TMEM161B strep-HA from filamentary forms to

monomers or dimers in 5 mM DTT as shown by cryo-electron microscopy. Scale bars,

10 nm. [(C) and (E)] *P* value was determined by one-way ANOVA.

Fig. S1. AlphaFold2-based computational screen for ERGU candidates. (**A**) BLASTP search yielding no apparent broadly conserved eukaryotic homologs of DscB/ScsB based on protein sequence similarities, prompting a search for DscB/ScsB homologs based on functional and structural features using AlphaFold2-predicted structures. (**B**) Predicted structures of the 53 ERGU candidates identified. (**C**) Table listing the annotated functions of the 11 non-GPCR ERGU candidates.

Fig. S2. Phenotypes of ERGU-1 protein-truncating *ergu-1(gk840471)* mutants. (A)

Representative epifluorescence images showing *gst-4p::GFP* activation in *ergu-1(gk840471)* mutants (n=15 for each group). (**B**) Quantification of *gst-4p*::GFP fluorescence intensities in WT and *ergu-1(gk840471)* mutants indicated (n=15 for each group). (**C**) Representative epifluorescence images showing *gst-4p::*GFP decrease by *ero-1* RNAi and increase by *gcs-1* RNAi (**D**) Representative epifluorescence images showing *gst-4p::GFP* activation in *ergu-1(gk840471)* mutants requires *skn-1.* (**E**) Representative confocal fluorescence images showing no apparent reporter (Golgi apparatus-marking Golgi::GFP, apical and endosomal membrane-marking AKT-PH::GFP, yolk organelle-marking VIT-2::GFP and autophagosome-marking SQST-1::mOrange2) alteration in *ergu-1(gk840471)* mutants. *P* value was determined by an unpaired *t*-test, two-tailed (comparison between two groups). [(A), (B) and (C)] Scale bars, 100 μm. (D) Scale bars, 20 μm.

Fig. S3. Phenotypic consequences of RNAi or mutant *ergu-1*. (A) Representative

epifluorescence images showing increased *hsp-4p::GFP* upon *ergu-1* RNAi treatment.

(**B**) Representative fluorescence images of *ergu-1p*::*ergu-1::GFP* showing abolished

GFP signals by *ergu-1* RNAi. (**C**) Representative fluorescence images of Vitamin B12

stress reporter *acdh-1p*:: *GFP* showing unaltered GFP signals by *ergu-1* RNAi. (**D** and

E) Representative fluorescence images of glutathione stress GSSG/GSH reporters

showing elevated oxidative stress in *ergu-1* null mutants (D) and quantification (n=10)

(E). (**F**) Representative fluorescence images of *ergu-1p*::*ergu-1::GFP* showing reporter

up-regulation by oxidative stressors (0.5% NaAsO$_2$). (**G**) Lifespan curves of wild-type

and *ergu-1* null mutants with control or *skn-1* RNAi showing comparable lifespans in

wild-type and *ergu-1* null mutants and reduction by *skn-1* RNAi. (**H**) Lifespan curves of

wild-type and *ergu-1* protein-truncating mutants (*gk840471*) with control or *skn-1* RNAi

comparable lifespans in wild-type and *ergu-1(gk840471)* mutants and reduction by *skn-*

1 RNAi (n ⩾ 50). (I) Quantification of brood sizes in wild type, *ergu-1* null and cysteine

mutants upon 10 mM paraquat treatment. *P* value was determined by one-way ANOVA

(comparison between multiple groups). (**J**) Quantification of average moving speed in

wild-type, ergu-1 null and above strain treated with 10 mM H2O2. (**K**) Representative

fluorescence images of cytoskeletal actin ACT-1::GFP reporters showing no apparent

alteration by *ergu-1* RNAi. (**L**) Representative confocal fluorescence images showing no

apparent reporter (PVD neuron-marking GFP) alteration in *ergu-1* null mutants. (**M**)

Representative confocal fluorescence images showing no apparent reporter (neuronal

nuclear GCaMP6) alteration in *ergu-1* null mutants. (**N**) Representative confocal

fluorescence images showing no apparent reporter (secreted MANF-1::GFP) alteration

by *ergu-1* RNAi. *P* value was determined by an unpaired *t*-test, two-tailed (comparison

between two groups). [(A) to (J)] Scale bars, 100 µm. [(K) to (M)] Scale bars, 20 µm.

Fig. S4. Importance of key cysteine residues for ERGU-1 functions. (**A**)

Representative fluorescence images of *gst-4p::GFP* showing no rescue of *C. elegans*

ergu-1 mutants by two different cysteine pair mutants of ERGU-1. (**B**) Schematic of key

residues for heme binding and interactions. Heme binding pocket is in high confidence

region (90 > pIDDT > 70). (**C**) Representative fluorescence images and quantification of

gst-4p::GFP showing no rescue of *C. elegans ergu-1* mutants by *C. elegans* or human

Y454 mutants of ERGU-1 homologs. [(A) and (C)] Scale bars, 100 μm.

Fig. S5. Colocalization of ERGU-1::GFP with RFP-marked ER organelle reporters but not non-ER organelle reporters. (A) Representative confocal fluorescence images showing colocalization between ERGU-1::GFP (green) and endoplasmic reticulum-specific reporters (red). Scale bars, 10 μm. (**B**) Representative confocal fluorescence images showing no colocalization between ERGU-1::GFP (green) and lysosome, mitochondria or plasma membrane-specific RFP reporters (red). Arrow indicates the organelles. Scale bars, 5 μm.

Fig. S6. Protein characteristics of human TMEM161B, an ERGU-1 ortholog and evolutionarily conserved member of the ERGU protein family. (**A**) Peak fractions of human TMEM161B-2X Strep-HA protein purified by SEC in a non-reductive environment. (**B**) Peak fractions of TMEM161B-2X Strep-HA protein purified by SEC in a DTT-containing reductive environment. (**C**) TMEM161B-GFP in a non-reductive environment shown by Cryo-EM. (**D**) Peak of TMEM161B-GFP in a non-reductive environment shown by SEC. (**E**) Peak fractions of TMEM161B-GFP protein purified by SEC in a non-reductive environment. (**F**) Immunofluorescence images showing that Human TMEM161A/B localizes to the ER membrane in U2OS cells. Scale bars, 10 μm.

488   (**G**) Representative confocal fluorescence images of roGFP2-Orp1 detecting cytosolic

489   H_2O_2 in 293T cells overexpressing human TMEM161A or B. Scale bars, 20 μm.

 Materials and Methods

C. elegans

C. elegans strains were maintained under standard laboratory conditions unless

otherwise specified. The N2 Bristol strain was used as the reference wild-type. Feeding

RNAi was performed as previously described[42]. Transgenic strains were generated by

germline transformation as described[43]. Transgenic constructs were co-injected (at 10 -

50 ng/µl) with dominant *unc-54p*::mCherry or *myo-2p*::mCherry, and stable

extrachromosomal lines of fluorescent animals were established for UV-mediated

genome integration. Genotypes of strains used are as follows: *ergu-1(ust572, syb9153,*

syb9160), dvIs19 [gst-4p::GFP], zcIs4 [hsp-4::GFP], jrIs10 [unc119(+) rps-0p::roGFP2-

Orp1], dmaIs160 [ergu-1p:: ergu-1::gfp; unc-54p::mCherry], dmaEx701[ergu-

1p::hsTMEM161B::ergu-1utr; myo-2p::mCherry], dmaEx702 [ergu-1p::ergu-1::ergu-1utr;

myo-2p::mCherry], dmaEx704 [ergu-1p::Dm-161:: ergu-1utr; myo-2p::mCherry].

AlphaFold-assisted computational screen

A comprehensive computational screen of the *Caenorhabditis elegans* proteome was

performed to identify potential DsbD/ScsB-like proteins. The reference proteome utilized

for this analysis was the *C. elegans* UniProt reference proteome (version 26),

encompassing 19,827 genes. Initial filtering was conducted to select genes encoding

proteins with a minimum of four annotated transmembrane helices, resulting in a subset

of 3,177 genes. Subsequent filtering criteria focused on the presence of transmembrane

helices containing cysteine residues, specifically targeting helices with at least two

cysteines separated by no more than two intervening residues, corresponding to the

sequence motifs CC, CXC, or CXXC. This filtering narrowed the candidate list to 190

genes. The structural analysis was conducted using predicted protein structures from

the AlphaFold2 Database. For each of the 190 genes, the predicted structures were

examined to determine the spatial proximity of cysteine pairs located within the

transmembrane helices. Specifically, structures in which the sulfur atoms (SG) of

cysteine pairs were within 10 angstroms of each other were identified. This criterion was

met by 53 candidate genes. All computational and structural analyses were executed

using custom Python scripts, which are provided in the Supplementary file. Visualization

and further structural analyses were performed using UCSF ChimeraX software.

RNA interference (RNAi) cloning and screen

RNAi and screen for hits upregulating *gst-4p*::GFP were performed by feeding worms

with E. coli strain HT115 (DE3) expressing double-strand RNA (dsRNA) targeting

endogenous genes. Briefly, dsRNA–expressing bacteria were replicated from the

Ahringer library to LB plates containing 100 µg/ml ampicillin (BP1760-25, Fisher

Scientific) at 37 °C for 16 hrs. Single clone was picked to LB medium containing 100

µg/ml ampicillin at 37 °C for 16 hrs and positive clones (verified by bacteria PCR with

pL4440 forward and pL4440 reverse primers) were spread onto NGM plates containing

100 µg/ml ampicillin and 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG, 420322,

Millopore) for 24 hrs. Developmentally synchronized embryos from bleaching of gravid

adult hermaphrodites were plated on RNAi plates and grown at 20 °C. The worms are

collected for imaging at indicated stages.

To make the *ergu-1* RNAi-1 and RNAi-2 plasmids, RNAi-1 targeting sequences are

PCR- amplified with primers 5'-

GGCCCCCCTCGAGGTCGACAGGAGACGATGCTATAAACTATGCT -3' and 5'-

TCCACCGCGGTGGCGGCCGCCGTGACGTGTTTTCAGCTCG -3' from N2 gDNA and

subcloned into the SalI and NotI sites of a digested T777T vector with Promega T4 DNA

ligase (Promega, M1801). RNAi-2 targeting sequences are PCR- amplified with primers

5'- GGCCCCCCTCGAGGTCGACATGCCAAAATGAATGCGGGC -3' and 5'-

TCCACCGCGGTGGCGGCCGCACGTGGTGATATGCCTGGTG -3'.

Fluorescence microscopy and H$_2$O$_2$ sensor imaging

SPE confocal (Leica) and epifluorescence microscopes were used to capture

fluorescence images. Animals were randomly picked at the same stage and treated with

1 mM levamisole in M9 solution (31742-250MG, Sigma-Aldrich), aligned on a 2% agar

pad on a slide for imaging. Identical setting and conditions were used to compare

experimental groups with control. For quantification of GFP fluorescence, animals were

outlined and quantified by measuring gray values using the ImageJ software. The data

were plotted and analyzed by using GraphPad Prism10. For *jrIs10 [unc119(+) rps-*

0p::roGFP2-Orp1] strain, orp1-roGFP2 was excited sequentially at 405 and 488 nm and

emission was recorded at 500-540 nm. Fifteen images were sequentially captured at 1-

micrometer z-intervals and subsequently stacked to form a composite image.

Western blotting

For SDS-PAGE samples, stage-synchronized animals for control and experiment groups were picked (n = 50) in 60 μl M9 buffer and lysed directly by adding 20 μl of 4x Laemmli sample buffer (1610747, Bio-Rad) contain 10% of 2-Mercaptoethanol (M6250-100ML, Sigma (v/v)). Protein extracts were denatured at 95 °C for 10 min and separated on 10% SDS-PAGE gels (1610156, Bio-Rad) at 80 V for ~40 min followed by 110 V for ~70 min. The proteins were transferred to a nitrocellulose membrane (1620094, Bio-Rad,) at 25 V for 45 mins by Trans-Blot® Turbo™ Transfer System (Bio-Rad). The NC membrane was initially blocked with 5% non-fat milk and 2% BSA (A4503, Sigma (v/v)) in tris buffered saline with 0.1% Tween 20 (93773, Sigma) (TBST) at room temperature for 1 h. Proteins of interest were detected using antibodies against GFP (A6455, Invitrogen) or V5 (13202T, Cell Signaling Technology) in cold room for overnight. After three washes of 5 min each with tris-buffered saline with 0.1% Tween 20, anti-mouse IgG, HRP-linked Antibody (7076, Cell Signaling Technology) was added at a dilution of 1:5000. For DTT treatment, worm lysates were treated with 10 mM DTT and boiled at 95 °C for 10 min.

Mammalian cell culture experiments

Human embryonic kidney (HEK) 293T cells and osteosarcoma U2OS cells were maintained in Dulbecco's modified Eagle's medium with 10% inactive fetal bovine serum and penicillin-streptomycin (Gibco, Grand Island, 15140122) at 37 °C supplied with 5% CO_2 in an incubator (Thermo Fisher Scientific) with a humidified atmosphere. Cells were washed once using PBS and digested with 0.05% trypsin EDTA (Gibco) at

37 °C for routine passage of the cells. All cells were transfected with 3 µl LipoD293™

(SignaGen, SL100668) per 1 µg DNA mixture. HEK293T cells were transfected by the

pLX304-TMEM161A-V5 (DNASU, HsCD00441633) or pLX304-TMEM161B-V5

(DNASU, HsCD00444935) and collected 2 days after transfection. HEK293T and U2OS

cells were transfected by the pHAGE2-TMEM161A-gfp/mCherry or pHAGE2-

TMEM161B-gfp/mCherry and collected for imaging 2 days after transfection. pHAGE2-

TMEM161A/B-GFP/mCherry was constructed by inserting TMEM161A or TMEM161B

into pHAGE2-gfp/mCherry plasmids. TMEM161A was PCR- amplified with the primers

5'- TAATTAAACCTCTAGAGCCACCATGGCGGTCCTCGG -3' and 5'-

CTCACCATAGCTCGAGacccagacccGGAGCCTGCCAAGTGC -3' from pLX304-

TMEM161A-V5 mentioned before. TMEM161B was PCR- amplified with the primers 5'-

TAATTAAACCTCTAGAGCCACCATGGGTGTGATAGGTATACAGC -3' and 5'-

CTCACCATAGCTCGAGacccagacccTGCCACAGTCAGATACTGGT -3' from pLX304-

TMEM161B-V5 mentioned before.

Quantification of brood size

The brood size assay was carried out according to the standard protocol. (1) Briefly,

single L4 worms of different stains (N2, ergu-1 null, PHX9155, PHX9160) were

individually placed in 60mm Petri plates and kept in the incubator at 20 C. Then, the

worms were transferred to a new OP50 containing plate each day at the same time till

day 6 of the worms. Progenies were scored after 3 days and plotted for each day and

the total sum of progenies. For NAC treatment the plates were supplemented with 10

mM of NAC during the entire duration of the assay. For H_2O_2 and paraquat treatment

the plates were supplemented with 10 mM of H_2O_2 and paraquat respectively during the

assay. Each experiment was repeated 3 times as independent biological replicates with

more than 5 animals per group.

Behavioral assay

D1 worms (24 hrs post L4) were transferred to a fresh NGM plate seeded with a small

and thin OP50 bacterial lawn and allowed to settle for at least ten minutes to recover at

room temperature. Moving average speeds, and track lengths of *C. elegans* were

monitored for 10 minutes and were analyzed using WormLab. For H_2O_2 treatment, D1

worms are treated with 10 mM H_2O_2 for 30 min before assay.

Lifespan analysis

For lifespan assays, Animals were cultured under non-starved conditions for at least 2

generations before life span assays. For normal NGM life span assay, stage-

synchronized L4 stage animals (n $\geqslant$ 50) were picked to new NGM plates seeded with

OP50 containing 50 μM 5-fluoro-2′-deoxyuridine (FUDR) to prevent embryo growth at

20 °C incubator. Animals were scored for survival per 24 hrs. Animals failing to respond

to repeated touch of a platinum wire were scored as dead. Three biological replicates

were performed, with population sizes larger than 50 in each trial.

DCFDA ROS detection

ROS staining in live worms was carried out as in standard protocol with minor

modifications. Briefly, day 1 worms were washed three times with M9 and then

transferred into 200 µl of M9 buffer containing 10 mM H2DCFDA (carboxy-H2DCFDA

[5-(and-6)-carboxy-2',7'-dichlorodihydrofluorescein diacetate) in an 1.5 ml Eppendorf

tube and incubated in the dark for 3 hrs. The worms were randomly selected and

treated with 10 mM sodium azide (Sigma-Aldrich) in M9, symmetrically aligned on 2%

agar pads on slides for imaging the oxidized dichlorofluorescin (DCF).

Expression and purification of TMEM161B for cryo-EM

The complementary DNA encoding human TMEM161B was cloned into a modified

pDNA3.1 vector with a C-terminal twin-strep tag and a HA tag. For expression,

Expi293F cells (Thermo Fisher Scientific) cultured in Freestyle293 Expression Medium

were transfected with the vector DNA/polyethylenimine (1 µg DNA per ml culture,

w/w = 1:3) complex at a cell density of ~1.0×10^6 cells per ml and incubated at 37 °C

under 8% CO_2 with agitation at 100 r.p.m for 60 h. Cell pellets were resuspended in

buffer A (25 mM Hepes pH 7.5, 150 mM NaCl, protease inhibitor cocktail) and were

disrupted by sonication. For TMEM161B monomer purification, an extra incubation with

5 mM DTT at room temperature after sonication was required. The cell lysate was then

spun at 150,000× g for 1h to sediment crude membranes. The membrane pellet was

mechanically homogenized in buffer A. The suspension was solubilized in 1% (w/v)

1.0% DDM (Anatrance) and 0.1% CHS (Anatrance) for 60 min at 4 °C. The solubilized

material was centrifuged at 100,000× g for 30 min, and the supernatant was incubated

with Strep-Tactin®XT resins (iba) for 2 h at 4 °C. Resins were then washed with 20

column volumes of buffer B (25 mM Hepes pH 7.5, 0.025% DDM, 0.0025% CHS,

150 mM NaCl). The protein was eluted with buffer B supplied with 50 mM D-biotin,

concentrated, and further purified by gel-filtration chromatography on a Superdex 200

increase column equilibrated with wash buffer B. The peak fractions of protein were

pooled and concentrated to ~5.0 mg/mL. For Cryo-EM observation, 3-μl aliquots of each

sample were applied onto a glow-discharged Quantifoil grid (R1.2/R1.3 300 mesh, Au),

blotted for 4.5–5.5 s in 100% humidity at 4 °C, and plunged into liquid ethane using a

Vitrobot MkIV (Thermo Fisher Scientific). Cryo-EM micrographs were obtained by using

a Talos Arctica G2 microscope (Thermo Fisher Scientific) running at 200 kV and a 300

kV Titan Krios microscope (Thermo Fisher Scientific).

Statistical analysis

Data were analyzed using GraphPad Prism 9.2.0 Software (Graphpad, San Diego, CA)

and presented as means ± S.D. unless otherwise specified, with P values calculated by

unpaired two-tailed t-tests (comparisons between two groups), one-way ANOVA

(comparisons across more than two groups) and adjusted with Bonferroni's corrections.

Box plots are presented as min to max, showing all data points.

Acknowledgment

We are grateful for the feedback from and discussion with other members of the Ma lab

at UCSF and Drs. P. Sigala, P. R. Ortiz de Montellano, A. Correia, Z. Chen, B.

DeGrado, I. Jain, B. Black and G. Huang. We thank the *Caenorhabditis* Genetics Center

(NIH grant #P40 OD010440), Wormbase.org (NIH grant #U24 HG002223 to P.

Sternberg), Wormatlas.org (NIH grant #OD010943 to D.H. Hall.) and CenGen

(cengen.org) for their immensely helpful resources. The work was supported by NIH

Author contributions

Z.J. and D.K.M. designed, performed and analyzed the *C. elegans* and cell culture

experiments, contributed to project conceptualization and wrote the manuscript. T.P.,

B.W., Y.T. and S.H.G. designed, performed and analyzed the *C. elegans* transgenic

and CRISPR experiments and editing manuscript. H.B. and Z.J. performed and

analyzed the ER imaging experiments. T.D.G conducted the computational screen and

contributed to project conceptualization and wrote the manuscript. Z.Y.L., J.X.Y., S.Y.X.

contributed to human TMEM161B biochemical analysis. D.K.M, O.D.W. and T.D.G

contributed to research materials, funding acquisition and editing manuscript.

Competing interests

686 The authors declare no competing interests.

References

1. Tu, B.P., and Weissman, J.S. (2004). Oxidative protein folding in eukaryotes: mechanisms and consequences. J. Cell Biol. *164*, 341–346. https://doi.org/10.1083/jcb.200311055.

2. Tu, B.P., Ho-Schleyer, S.C., Travers, K.J., and Weissman, J.S. (2000). Biochemical basis of oxidative protein folding in the endoplasmic reticulum. Science *290*, 1571–1574. https://doi.org/10.1126/science.290.5496.1571.

3. Mamathambika, B.S., and Bardwell, J.C. (2008). Disulfide-linked protein folding pathways. Annu Rev Cell Dev Biol *24*, 211–235. https://doi.org/10.1146/annurev.cellbio.24.110707.175333.

4. Gao, C., Tian, Y., Zhang, R., Jing, J., and Zhang, X. (2017). Endoplasmic Reticulum-Directed Ratiometric Fluorescent Probe for Quantitive Detection of Basal H2O2. Anal Chem *89*, 12945–12950. https://doi.org/10.1021/acs.analchem.7b03809.

5. Lyublinskaya, O., and Antunes, F. (2019). Measuring intracellular concentration of hydrogen peroxide with the use of genetically encoded H2O2 biosensor HyPer. Redox Biol *24*, 101200. https://doi.org/10.1016/j.redox.2019.101200.

6. Sies, H. (2017). Hydrogen peroxide as a central redox signaling molecule in physiological oxidative stress: Oxidative eustress. Redox Biol *11*, 613–619. https://doi.org/10.1016/j.redox.2016.12.035.

7. Glorieux, C., and Calderon, P.B. (2017). Catalase, a remarkable enzyme: targeting the oldest antioxidant enzyme to find a new cancer treatment approach. Biol Chem *398*, 1095–1108. https://doi.org/10.1515/hsz-2017-0131.

8. Imlay, J.A. (2008). Cellular defenses against superoxide and hydrogen peroxide. Annu Rev Biochem *77*, 755–776. https://doi.org/10.1146/annurev.biochem.77.061606.161055.

9. Schriner, S.E., Linford, N.J., Martin, G.M., Treuting, P., Ogburn, C.E., Emond, M., Coskun, P.E., Ladiges, W., Wolf, N., Van Remmen, H., et al. (2005). Extension of murine life span by overexpression of catalase targeted to mitochondria. Science *308*, 1909–1911. https://doi.org/10.1126/science.1106653.

10. J, R., N, B., and Jm, H. (2009). Disulfide formation in the ER and mitochondria: two solutions to a common process. Science (New York, N.Y.) *324*. https://doi.org/10.1126/science.1170653.

11. Bienert, G.P., Møller, A.L.B., Kristiansen, K.A., Schulz, A., Møller, I.M., Schjoerring, J.K., and Jahn, T.P. (2007). Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. J Biol Chem *282*, 1183–1192. https://doi.org/10.1074/jbc.M603761200.

12. Cho, S.-H., Parsonage, D., Thurston, C., Dutton, R.J., Poole, L.B., Collet, J.-F., and
Beckwith, J. (2012). A new family of membrane electron transporters and its
substrates, including a new cell envelope peroxiredoxin, reveal a broadened
reductive capacity of the oxidative bacterial cell envelope. mBio *3*, e00291-11.
https://doi.org/10.1128/mBio.00291-11.

13. Rozhkova, A., and Glockshuber, R. (2008). Thermodynamic aspects of DsbD-
mediated electron transport. J Mol Biol *380*, 783–788.
https://doi.org/10.1016/j.jmb.2008.05.050.

14. Ellgaard, L., Sevier, C.S., and Bulleid, N.J. (2018). How Are Proteins Reduced in the
Endoplasmic Reticulum? Trends Biochem Sci *43*, 32–43.
https://doi.org/10.1016/j.tibs.2017.10.006.

15. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly
accurate protein structure prediction with AlphaFold. Nature *596*, 583–589.
https://doi.org/10.1038/s41586-021-03819-2.

16. Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris,
J.H., and Ferrin, T.E. (2023). UCSF ChimeraX: Tools for structure building and
analysis. Protein Sci *32*, e4792. https://doi.org/10.1002/pro.4792.

17. Leiers, B., Kampkötter, A., Grevelding, C.G., Link, C.D., Johnson, T.E., and Henkle-
Dührsen, K. (2003). A stress-responsive glutathione S-transferase confers
resistance to oxidative stress in Caenorhabditis elegans. Free Radic Biol Med *34*,
1405–1415. https://doi.org/10.1016/s0891-5849(03)00102-3.

18. Sturm, Á., Saskoi, É., Tibor, K., Weinhardt, N., and Vellai, T. (2018). Highly efficient
RNAi and Cas9-based auto-cloning systems for C. elegans research. Nucleic Acids
Res *46*, e105. https://doi.org/10.1093/nar/gky516.

19. Blackwell, T.K., Steinbaugh, M.J., Hourihan, J.M., Ewald, C.Y., and Isik, M. (2015).
SKN-1/Nrf, stress responses, and aging in Caenorhabditis elegans. Free Radic Biol
Med *88*, 290–301. https://doi.org/10.1016/j.freeradbiomed.2015.06.008.

20. Sies, H., Berndt, C., and Jones, D.P. (2017). Oxidative Stress. Annu Rev Biochem
86, 715–748. https://doi.org/10.1146/annurev-biochem-061516-045037.

21. Pa, F., Tf, S., T, B., R, B.-Z., Hk, G., H, Z., R, H.-S., and A, D. (2022). SKN-1
regulates stress resistance downstream of amino catabolism pathways. iScience *25*.
https://doi.org/10.1016/j.isci.2022.104571.

22. Gusarov, I., Shamovsky, I., Pani, B., Gautier, L., Eremina, S., Katkova-Zhukotskaya,
O., Mironov, A., Makarov, A.A., and Nudler, E. (2021). Dietary thiols accelerate
aging of C. elegans. Nat Commun *12*, 4336. https://doi.org/10.1038/s41467-021-
24634-3.

23. De Magalhaes Filho, C.D., Henriquez, B., Seah, N.E., Evans, R.M., Lapierre, L.R.,
and Dillin, A. (2018). Visible light reduces C. elegans longevity. Nat Commun *9*, 927.
https://doi.org/10.1038/s41467-018-02934-5.

24. Aruoma, O.I., Halliwell, B., Hoey, B.M., and Butler, J. (1989). The antioxidant action
of N-acetylcysteine: its reaction with hydrogen peroxide, hydroxyl radical,
superoxide, and hypochlorous acid. Free Radic Biol Med *6*, 593–597.
https://doi.org/10.1016/0891-5849(89)90066-x.

25. Back, P., De Vos, W.H., Depuydt, G.G., Matthijssens, F., Vanfleteren, J.R., and
Braeckman, B.P. (2012). Exploring real-time in vivo redox biology of developing and
aging Caenorhabditis elegans. Free Radic Biol Med *52*, 850–859.
https://doi.org/10.1016/j.freeradbiomed.2011.11.037.

26. Zhang, Z., Bai, M., Barbosa, G.O., Chen, A., Wei, Y., Luo, S., Wang, X., Wang, B.,
Tsukui, T., Li, H., et al. (2020). Broadly conserved roles of TMEM131 family proteins
in intracellular collagen assembly and secretory cargo trafficking. Sci Adv *6*,
eaay7667. https://doi.org/10.1126/sciadv.aay7667.

27. Ma, D.K., Vozdek, R., Bhatla, N., and Horvitz, H.R. (2012). CYSL-1 interacts with
the O2-sensing hydroxylase EGL-9 to promote H2S-modulated hypoxia-induced
behavioral plasticity in C. elegans. Neuron *73*, 925–940.
https://doi.org/10.1016/j.neuron.2011.12.037.

28. Wang, C., Wang, B., Pandey, T., Long, Y., Zhang, J., Oh, F., Sima, J., Guo, R., Liu,
Y., Zhang, C., et al. (2022). A conserved megaprotein-based molecular bridge
critical for lipid trafficking and cold resilience. Nat Commun *13*, 6805.
https://doi.org/10.1038/s41467-022-34450-y.

29. Klemm, R.W., Norton, J.P., Cole, R.A., Li, C.S., Park, S.H., Crane, M.M., Li, L., Jin,
D., Boye-Doe, A., Liu, T.Y., et al. (2013). A conserved role for atlastin GTPases in
regulating lipid droplet size. Cell Rep *3*, 1465–1475.
https://doi.org/10.1016/j.celrep.2013.04.015.

30. Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger,
O., Willmore, L., Ballard, A.J., Bambrick, J., et al. (2024). Accurate structure
prediction of biomolecular interactions with AlphaFold 3. Nature.
https://doi.org/10.1038/s41586-024-07487-w.

31. Aithani, L., Alcaide, E., Bartunov, S., Cooper, C.D.O., Doré, A.S., Lane, T.J.,
Maclean, F., Rucktooa, P., Shaw, R.A., and Skerratt, S.E. (2023). Advancing
structural biology through breakthroughs in AI. Curr Opin Struct Biol *80*, 102601.
https://doi.org/10.1016/j.sbi.2023.102601.

32. Yang, Z., Zeng, X., Zhao, Y., and Chen, R. (2023). AlphaFold2 and its applications
in the fields of biology and medicine. Signal Transduct Target Ther *8*, 115.
https://doi.org/10.1038/s41392-023-01381-z.

33. Mosalaganti, S., Obarska-Kosinska, A., Siggel, M., Taniguchi, R., Turoňová, B.,
Zimmerli, C.E., Buczak, K., Schmidt, F.H., Margiotta, E., Mackmull, M.-T., et al.
(2022). AI-based structure prediction empowers integrative structural analysis of
human nuclear pores. Science *376*, eabm9506.
https://doi.org/10.1126/science.abm9506.

34. Ushioda, R., Hoseki, J., Araki, K., Jansen, G., Thomas, D.Y., and Nagata, K. (2008).
ERdj5 is required as a disulfide reductase for degradation of misfolded proteins in
the ER. Science *321*, 569–572. https://doi.org/10.1126/science.1159293.

35. Arts, I.S., Gennaris, A., and Collet, J.-F. (2015). Reducing systems protecting the
bacterial cell envelope from oxidative damage. FEBS Lett *589*, 1559–1568.
https://doi.org/10.1016/j.febslet.2015.04.057.

36. van Soest, D.M.K., Polderman, P.E., den Toom, W.T.F., Keijer, J.P., van
Roosmalen, M.J., Leyten, T.M.F., Lehmann, J., Zwakenberg, S., De Henau, S., van
Boxtel, R., et al. (2024). Mitochondrial H_2O_2 release does not directly cause
damage to chromosomal DNA. Nat Commun *15*, 2725.
https://doi.org/10.1038/s41467-024-47008-x.

37. Ma, X., Lu, J.-Y., Moraru, A., Teleman, A.A., Fang, J., Qiu, Y., Liu, P., and Xu, T.
(2020). A novel regulator of ER Ca2+ drives Hippo-mediated tumorigenesis.
Oncogene *39*, 1378–1387. https://doi.org/10.1038/s41388-019-1076-z.

38. Koopman, C.D., De Angelis, J., Iyer, S.P., Verkerk, A.O., Da Silva, J., Berecki, G.,
Jeanes, A., Baillie, G.J., Paterson, S., Uribe, V., et al. (2021). The zebrafish grime
mutant uncovers an evolutionarily conserved role for Tmem161b in the control of
cardiac rhythm. Proc Natl Acad Sci U S A *118*, e2018220118.
https://doi.org/10.1073/pnas.2018220118.

39. Akula, S.K., Chen, A.Y., Neil, J.E., Shao, D.D., Mo, A., Hylton, N.K., DiTroia, S.,
Ganesh, V.S., Smith, R.S., O'Kane, K., et al. (2023). Exome Sequencing and the
Identification of New Genes and Shared Mechanisms in Polymicrogyria. JAMA
Neurol *80*, 980–988. https://doi.org/10.1001/jamaneurol.2023.2363.

40. Akula, S.K., Marciano, J.H., Lim, Y., Exposito-Alonso, D., Hylton, N.K., Hwang, G.H.,
Neil, J.E., Dominado, N., Bunton-Stasyshyn, R.K., Song, J.H.T., et al. (2023).
TMEM161B regulates cerebral cortical gyration, Sonic Hedgehog signaling, and
ciliary structure in the developing central nervous system. Proc Natl Acad Sci U S A
120, e2209964120. https://doi.org/10.1073/pnas.2209964120.

41. Wang, L., Heffner, C., Vong, K.I., Barrows, C., Ha, Y.-J., Lee, S., Lara-Gonzalez, P.,
Jhamb, I., Van Der Meer, D., Loughnan, R., et al. (2023). TMEM161B modulates
radial glial scaffolding in neocortical development. Proc Natl Acad Sci U S A *120*,
e2209983120. https://doi.org/10.1073/pnas.2209983120.

42. Kamath, R.S., and Ahringer, J. (2003). Genome-wide RNAi screening in
837   Caenorhabditis elegans. Methods *30*, 313–321.

838   43. Mello, C.C., Kramer, J.M., Stinchcomb, D., and Ambros, V. (1991). Efficient gene
839   transfer in C.elegans: extrachromosomal maintenance and integration of
840   transforming sequences. EMBO J. *10*, 3959–3970.

841