Perturbing human V1 degrades the fidelity of visual working memory

Mrugank Dake[1] and Clayton E. Curtis[1, 2]

1. Department of Psychology, New York University
2. Center for Neural Science, New York University

Correspondence: clayton.curtis@nyu.edu

Keywords: working memory, transcranial magnetic stimulation, electroencephalography, memory-guided saccade, sensory recruitment, saccade, visual cortex

Abstract

Decades of macaque research established the importance of prefrontal cortex for working memory. Surprisingly, recent human neuroimaging studies demonstrated that the contents of working memory can be decoded from primary visual cortex (V1). However the necessity of this mnemonic information remains unknown and contentious. Here we provide causal evidence that transcranial magnetic stimulation targeting human V1 disrupted the fidelity of visual working memory. Errors increased only for targets remembered in the portion of the visual field disrupted by stimulation. Moreover, concurrently measured electroencephalography confirmed that stimulation disrupted not only memory behavior, but neurophysiological signatures of working memory. These results change the question from whether visual cortex is necessary for working memory to what mechanisms it uses to support memory. Moreover, they point to models in which the mechanisms supporting working memory are distributed across brain regions, including sensory areas that here we show are critical for memory storage.

Introduction

Our ability to maintain information no longer present in the environment, referred to as working memory, acts as a building block supporting most higher forms of cognition [1]. Indeed, impaired working memory can cause a cascade of broad cognitive symptoms in psychiatric disease [2]. Based on pioneering studies, the prefrontal cortex is theorized to store information in working memory in recurrent networks whose neural activity forms a mnemonic bridge between perception and memory-guided behavior [3,4]. There are two major pillars of support for this theory from studies of macaques. One, activity in prefrontal cortical neurons persists during memory delays [5]. Two, lesions to the macaque prefrontal cortex impair working memory performance [6]. More recently, human neuroimaging studies demonstrate that working memory representations may be widely distributed in the brain beyond the prefrontal cortex [7,8], including evidence for persistent neural activity in many cortical [9] and even subcortical areas [10]. Similarly, lesions to human cortical areas, both within and outside of the prefrontal cortex, impact working memory performance [11,12].

The most surprising finding in working memory research, which has now been replicated numerous times, is that the multivariate patterns of neural activity in primary visual cortex, V1, can be used to decode the contents of visual working memory [13,14]. Moreover, patterns in V1 are often the best for decoding the contents of working memory. Therefore, activity in V1 may encode representations needed for storing visual features in working memory [15–17], an idea that has been termed the sensory recruitment model of working memory. However, healthy skepticism exists about the necessity of V1 for working memory, as both macaque neurophysiology and human neuroimaging are mixed as to whether the neural activity in V1 persists during retention intervals [9,18], a hallmark of working memory. Perhaps the information decoded from V1 is not necessary for working memory despite being present. Lesions cannot resolve whether V1 is necessary for working memory because lesions to V1 would simply prevent visual encoding.

To overcome this impasse, here we directly tested whether human V1 is necessary for working memory using transcranial magnetic stimulation (TMS) to perturb neural processing during retention intervals.

Results

We first used fMRI to identify and model the receptive field properties of V1 [19]. Then, we identified the portion of V1 accessible to target with TMS - the dorsal part of the retinotopic map that represents the contralateral lower quadrant of the visual field (Figure 1A). We modeled the electric field induced by TMS [20] to target V1 in each participant (Figure 1B; see individual E-field maps in Supp Figure 1). These models indicated that TMS induced a greater electrical field in the V1 hemisphere targeted (Figure 1C). Although we directed the focus of the TMS pulse to V1, our electric field models suggested that to a lesser extent dorsal V2 may have also been affected.

76   Targeting V1 with TMS induced phosphenes whose spatial extents were drawn by participants.
77   This allowed us to identify the phosphene field and its mirrored location (Figure 1D). Phosphene
78   fields aligned well with the biophysical models we used to estimate the electric fields induced by
79   TMS.
80
In Experiment 1, we measured the accuracy of memory-guided saccades for targets placed within
the phosphene field (i.e., contralesional to TMS), and compared the performance to that when no
TMS was applied and to that when the targets were placed in the hemifield opposite the phosphene
field (i.e., ipsilesional to TMS) (Figure 1E/F). Remarkably, TMS applied to V1 during the middle
of the delay caused larger memory errors, but only for targets within the phosphene field (Figure
2A). Specifically, memory errors were greater for targets inside the phosphene field compared to
when no TMS was applied ($t(14)=2.431$, $p=0.013$, $d=1.30$), and compared to targets in the mirrored
hemifield of the phosphene field ($t(14)=2.506$, $p=0.012$, $d=1.34$). Memory errors were not different
for targets outside the phosphene field during TMS and during no TMS trials ($t(14)=-0.678$,
$p=0.745$, $d=0.36$) confirming the retinotopic specificity of the effect of TMS targeting V1 on
working memory performance.
92
In Experiment 2, we aimed to both test the reproducibility of these results and how the timing of
the stimulation impacts working memory performance. The same participants (n=15) returned and
we again delivered TMS to V1 during the delay period of the memory-guided saccade task (Figure
1F). First, we perfectly replicated our findings from Experiment 1. TMS applied to V1 during the
middle of the delay period caused greater memory errors for targets inside the phosphene field
compared to targets in the mirrored hemifield ($t(14)=2.742$, $p=0.007$, $d=1.47$). Next, we asked how
TMS applied during working memory maintenance compares to when applied during encoding
(Figure 1F). TMS applied to V1 early in the delay (at the offset of the visual target) impaired
memory accuracy for targets inside the phosphene field compared to targets outside it
($t(14)=3.303$, $p=0.003$, $d=1.77$). Although numerically larger, these errors caused by TMS at
encoding were not statistically greater than those caused by TMS during maintenance
($t(14)=1.248$, $p=0.095$, $d=0.67$).
105
Next, since we used memory-guided saccades as our measure of working memory, we explored
what aspects of the saccade metrics were affected by TMS. First, TMS applied during the middle
of the delay ($t(14)=-0.457$, $p=0.655$, $d=0.244$) or during the early delay ($t(14)=0.251$, $p=0.806$,
$d=0.134$) did not impact saccade reaction times (Supp Figure 2A). Similarly, TMS applied during
the middle of the delay ($t(14)=0.321$, $p=0.753$, $d=0.171$) or during the early delay ($t(14)=0.261$,
$p=0.798$, $d=0.140$) had no impact on the peak velocity of saccades (Supp Figure 2B). Therefore
the effects of TMS we observed cannot be explained by a speed-accuracy tradeoff or slowing of
saccades. Second, using regression we found that memory-guided saccade errors were driven by
both radial (i.e., gain; No TMS $t(14)=24.992$, $p=5.1487e-13$, $d=13.359$; TMS outPF $t(14)=19.789$,
$p=1.2420e-11$, $d=10.577$; TMS inPF $t(14)=24.090$, $p=8.5177e-13$, $d=12.877$) and tangential (i.e.,

angular; No TMS $t(14)=7.822$, $p=1.7791e-06$, $d=4.181$; TMS outPF $t(14)=8.236$, $p=9.7491e-07$,
$d=4.402$; TMS inPF $t(14)=7.214$, $p=4.4699e-06$, $d=3.856$) error. Moreover, TMS condition did
not impact the way in which saccade errors were decomposed into these components (radial - No
TMS vs TMS $t(14)=-1.732$, $p=0.105$, $d=0.926$; radial - TMS inPF vs TMS outPF $t(14)=0.651$,
$p=0.525$, $d=0.348$; tangential - No TMS vs TMS $t(14)=0.214$, $p=0.834$, $d=0.114$; tangential - TMS
inPF vs TMS outPF $t(14)=-0.4874$, $p=0.634$, $d=0.261$). Therefore, both radial and tangential
components were equally impacted by TMS. This is important because it suggests that targeting
V1 with TMS caused errors that cannot merely be explained by some systematic motor factors
(e.g., only hypometricity). Instead, they point to induced errors in memory that were highly
specific to targets in the phosphene field.
126
Another possible explanation for our results is that the visual presence of phosphenes acted as
distraction or somehow captured attention. Phosphenes were rarely reported by participants
because of the brighter gray background during the experiment compared to the black used during
phosphene mapping. When they were reported on rare occasions, the intensity of the TMS was
reduced and participants no longer reported seeing phosphenes. Moreover, it is not clear what the
prediction would be for such a visual distraction as behaviorally relevant, but not irrelevant, stimuli
can impact working memory representations in early visual cortex [21,22]. Nonetheless, we performed
a control experiment in which participants were shown simulated phosphenes as three flickers
matching the exact times of the three TMS pulses during the middle of the memory delay (See
Methods). The pseudo phosphenes were low contrast noise patterns created to match subjective
reports from our participants, and were shown at the same exact positions as the real phosphene
fields experimentally defined (Supp Figure 4A). Memory errors were comparable for trials where
pseudo phosphenes were presented on the same or opposite hemifield compared to the target
($t(9)=0.671$, $p=0.5194$, $d=0.447$) (Supp Figure 4B), therefore ruling out that our TMS effects were
caused by rare distracting effects of visible phosphenes. Together, these results demonstrate that
perturbations targeting V1 are reliable, retinotopically specific, and are not likely the cause of
oculomotor disturbances or secondary effects of visual distraction.
144
We used electroencephalography (EEG) in Experiment 1 to measure the impact that TMS targeting
V1 had on the characteristic asymmetry of alpha band topography over occipital cortex electrodes
during working memory [23]. Without TMS, alpha power decreased in the hemisphere contralateral
to the memory target early in the delay, which we quantified with an alpha lateralization index
(αLI) (Figure 2B, top). The lateralized alpha response persisted into the late portion of the delay,
albeit lower in amplitude. On TMS trials, early in the delay the topography of alpha was
indistinguishable from the no TMS trials before stimulation. Following TMS, the asymmetric
alpha response was reduced compared to no TMS (Figure 2B, bottom). Next, we subtracted the
EEG time-frequency responses of the no TMS trials from the TMS trials among the occipital
electrodes over the hemisphere in which stimulation was applied (see *Methods*). Once the TMS
artifact resolved (Supp Figure 3), we found significant reductions in alpha band power for about

500ms and beta band power for about 250ms (Figure 2C). Therefore, consistent with the increase
in behavioral memory errors, TMS also disrupted a prominent EEG signature of working memory.
158
159 **Discussion**
160
In summary, TMS targeting human V1 caused a deficit in working memory that was spatially
restricted to the portion of the visual field perturbed by the electrical field induced by the TMS
pulses. While previous studies testing how visual information is consolidated into working
memory report that TMS to occipital cortex during the early encoding phase of working memory
disrupts memory performance [24–26], we find that TMS during the maintenance phase produced
deficits just as robust as those during encoding. The few other TMS studies of early visual cortex
for working memory have been unable to confirm or refute whether it is necessary. For instance,
TMS to occipital cortex only slowed working memory based decisions when applied after the
delay period [25]. In one study, TMS worsened working memory precision very early in the delay
(100ms), but not later (400ms) or even during the presentation of the visual array [26]. In another
study, TMS had an effect in only 1 out of 6 conditions (high working memory load at 400ms into
the delay) [24]. In yet another study, 10Hz TMS actually improved working memory precision for
gabor orientations [27]. Finally, TMS to human MT disrupted working memory but only for the more
prioritized of two memorized motion stimuli [28]. Although we can only speculate about the various
factors that might underlie the inconsistency of these results, we will highlight a few here. First,
most of the previous studies were designed to mainly test the role of visual cortex in encoding or
early consolidation, consistent with the notion that its role is limited to memory encoding and not
maintenance. Second, methodological issues might have hampered clear interpretations. Here, we
controlled the retinal specificity of the effects by placing stimuli in the phosphene fields during
the early and middle of the memory delay. We applied strong triple pulses of TMS that produced
robust effects. We used measures sensitive to small changes in working memory behavior. We
also used EEG to confirm that our perturbations impacted neural indices associated with working
memory. Bolstered by these experimental procedures, our findings provide direct causal evidence
that early visual cortex may be necessary for working memory, beyond simply the visual encoding
or consolidation of stimuli into memory.
186
As these results have important consequences for theories of working memory, we considered
alternative explanations. For instance, perhaps higher-order areas known to be important for
working memory, like the frontal and parietal cortices, were somehow affected by TMS targeting
V1. We believe that this is unlikely. First, neither V1 or V2 have direct monosynaptic projections
to the dorsolateral prefrontal cortex, area BA46 in the macaque [29]. The vast majority of connections
between V1 and lateral intraparietal area (LIP) or frontal eye field (FEF) are not monosynaptic [29].
For instance, monkey V1 is mainly connected to LIP indirectly through several synapses [30].
Second, PET and fMRI studies of concurrent TMS suggest that distal areas affected are only those
that have monosynaptic connections to the brain target of the TMS [31,32]. Third, electrical
microstimulation of macaque FEF without concurrent visual stimulation does not evoke BOLD

activity in V1 [33]. Fourth, TMS perturbs spiking activity only within a 2mm radius, and synaptic
potentials only within a slightly larger extent in the macaque cortex [34]. This evidence together
suggests that the effects we observed here are due to highly localized effects of TMS to V1 and
perhaps V2d.
201
We must also acknowledge two caveats to our conclusion that V1 is necessary for working
memory. One, while we targeted retinotopically defined V1, perhaps the effects we observed were
due to the propagation of electrical activity to other distant brain areas like the prefrontal cortex.
While by far the strongest electrical effects are localized to brain areas directly targeted by TMS,
one would assume that smaller effects might propagate to areas a single synapse away. Based on
the macaque brain, neither V1 or V2 has direct monosynaptic connections with the dorsolateral
prefrontal cortex, and the overwhelming majority of connections between V1 to LIP and FEF are
not monosynaptic [29,30]. Therefore, if our results were due to indirect effects of TMS on areas
monosynaptically connected to V1, they are most likely limited to areas extensively connected
with V1, such as V2, V3, and V4 and subcortical areas such as the lateral geniculate nucleus.
Indeed, no evidence exists that the indirect effects of TMS on distant brain areas have effects on
behavior. Rather, the effects on neuronal spiking caused by TMS are extremely focal (~2mm) [34]
and single finger contractions can be individuated by TMS displacements of only ~1mm across
cortex [35]. Two, TMS targeting V1 worsened the accuracy of memory-guided saccades, which we
interpreted as an impact on visuospatial working memory. However, other possible cognitive or
motor behaviors may have been affected, and thus masqueraded as a memory deficit. Although
spatial working memory and spatial attention are thought to rely on overlapping neural
mechanisms [36,37], TMS to V1 surprisingly does not affect endogenous covert attention [38].
Additionally, perhaps TMS targeting V1 somehow impacted the oculomotor metrics used to plan
saccades [39,40]. We do not believe that the results are consistent with this possibility as the memory-
guided saccade errors induced by TMS were highly specific to memory targets inside the
contralateral phosphene field. Saccade response times and peak velocity were not affected.
Moreover, the types of memory-guided saccade errors were not systematic; TMS impacted both
the gain and angle of saccade errors (again, only in the contralateral visual field). Nonetheless,
future research should extend our results by testing if more complicated forms of visual working
memory depend on V1, taking into account the difficult challenges in establishing causality [41].
228
Our findings provide compelling support to human neuroimaging studies that consistently report
that the visual properties of memoranda stored in working memory can be decoded from the
patterns of activity in V1 during retention intervals [7,9]. Moreover, they align with
neurophysiological recordings from macaque V1 demonstrating stimulus selective activity during
working memory delays [42–45]. They provide a key piece of evidence that had been missing, but
needed to directly test the sensory recruitment hypothesis of working memory. Indeed, we
demonstrate that the neural mechanisms housed in V1 (and perhaps V2d) are necessary to maintain
accurate working memory representations. Therefore, the question is no longer if visual cortex is

necessary for working memory, but now it is by what mechanisms does it contribute to working
memory? Assuming indirect effects do not account for our results, they are consistent with models
in which the neural mechanisms housed in V1 (and perhaps V2d) play an important role in
maintaining accurate working memory representations. Given the accumulation of evidence from
neuroimaging, electrophysiology, and TMS, we argue that the question should no longer be about
if visual cortex plays an important role in working memory, but instead should be about by what
mechanisms does it contribute to working memory?
244
245

Methods

Participants

All experiments were conducted following the institutional guidelines of New York University and the regulations of the institutional review board (IRB).Twenty-seven neurologically healthy humans (12 female, 15 male; mean age: 28, range: 21-55) were recruited to participate in multi-session TMS studies. All participants had normal to corrected normal vision and were screened for TMS eligibility and excluded from the study if they had any brain-related medical issues or were currently taking certain drugs (e.g. antidepressants, amphetamines, etc.). All participants provided written, informed consent and were compensated $50 for each session, regardless of whether it involved TMS (since the order of TMS was randomized across participants). Three participants found the procedures uncomfortable and withdrew. Five participants were withdrawn because we could not measure their gaze well enough (e.g., poor pupil lock, excessive blinking, lid closures, and dry/watery eyes). Since gaze was our main dependent variable, we could not study these participants. One participant was excluded because they did not follow instructions and made saccades only after the visual feedback was given, and thus not based on memory. Three participants were excluded because we could not reliably evoke phosphenes in the contralateral hemifield. The results, therefore, were based on fifteen participants (7 female, 8 male; mean age: 31, range: 21-55) who completed the full study. Two of the participants were authors and the rest were naive to the purpose of the study.

Rapid Serial Visual Presentation (RSVP) Task and Retinotopic Mapping

Anatomical and functional brain scans for pRF mapping were collected at New York University Center for Brain Imaging (CBI) using a 3T Siemens Prisma MRI scanner using a Siemens 64-channel head/neck radiofrequency coil. Volumes were acquired using a T2*-sensitive echo planar imaging pulse sequence (repetition time (TR), 1300 ms; echo time (TE), 36 ms; flip angle, 66°; 56 slices; 2 mm x 2mm x 2mm voxels). High-resolution T1-weighted images (0.8 mm x 0.8 mm x 0.8 mm voxels) were collected at the end of the session, with the same slice prescriptions as for the functional data, and used for registration, segmentation, and display. Multiple distortion scans (TR 6,690 ms; TE 63.4 ms; flip angle, 90°; 56 slices; 2 mm x 2 mm x 2 mm voxels) were collected during each scanning session.

To define visual field maps, each participant underwent retinotopic mapping in the MRI scanner, following previously established procedures [19]. Participants performed a RSVP task composed of sweeping bars of stimuli that swept across the screen in swift and discrete steps, moving in four possible directions: leftward to rightward, top to bottom, rightward to leftward, or bottom to top. Each bar contained six images of colorful objects. At the start of each block, participants were shown the target object and were instructed to make a button press every time the target object was

present anywhere within the array while maintaining fixation. The participants received online
feedback with the fixation cross turning green indicating their response was accurate. The inter-
stimulus interval was 400 ms and was titrated at the end of each sweep to match participants'
accuracy to 70-80%. The data were monitored online for fixation breaks and blocks where
participants broke fixation were eliminated from pRF model-fitting. Participants performed around
10 runs of the task, with 8 bar sweeps per run.
292
Resulting BOLD time series were fitted with a population receptive field (pRF) model with
compressive spatial summation [46,47], and visual field maps, including V1, were drawn manually
using afni and suma. Specifically, we visualized the polar angle and the eccentricity maps on the
cortical surface, thresholded to include only voxels for which the pRF model explained more than
10% of the variance. These ROIs were then inflated using volumetric data as grid parent and used
for targeting with TMS (as described below).
299
300 **Transcranial Magnetic Stimulation (TMS)**
301
The T1-weighted images of participants were used to reconstruct brain and skin surfaces to localize
targets for TMS using a frameless stereotaxic neuronavigation system (Brainsight, Rogue
Research Inc., CAN) combined with Polaris Spectra (Northern Digital, USA) infrared cameras
that tracked markers positioned on the TMS coil and on the participant's head. TMS was applied
using a MagVenture MagPro X100 stimulator with a MCF-B70 static cooled coil. TMS pulses
were triggered and coupled with EEG using custom MATLAB and Python scripts
(https://github.com/clayspacelab/EEG_TMS_triggers). We identified V1 in each participant from
fMRI-based retinotopy (see *Retinotopic Mapping* above) and overlaid the target on the brain
surface in Brainsight. We initially randomized whether the participant would receive TMS to the
left or right V1. However, if we had difficulties evoking reliable phosphenes we switched to the
other hemisphere. Once we identified a putative target that evoked phosphenes, we then
determined the phosphene threshold and mapped the spatial extent of the phosphene (see
Phosphene Mapping next).
315
316 **Phosphene Mapping**
317
Participants were first screened for their ability to detect phosphenes in a dimly lit room by placing
the TMS coil over retinotopically defined V1 and applying 7 pulses at 20 Hz. The coil was oriented
with the handle away from the interhemispheric fissure to induce currents along the caudal-rostral
axis. We made small changes to the position and angle of the TMS coil until we identified a target
location that reliably induced phosphenes. If phosphenes were inconsistent or weak, we tried
stimulating V1 in the other hemisphere. Next, we then measured the spatial extent and borders of
phosphenes. Participants used a computer mouse to encircle the perceived phosphene on several
trials until consistency was achieved, defined as at least three trials with overlapping phosphenes.

For each participant, we defined their phosphene field (PF), which we derived from the
overlapping area of the phosphene drawings. The PF was the area of visual space consistently
impacted by TMS to V1 at the given coil position. The area of the PF was determined by the
following criteria. First, the centroid was calculated as the center of the overlapping area. Second,
the angular extent spanned the full distance along an arc running through the centroid. Third, the
radial extent was defined as a 2 degree band centered at the centroid. In the experiment, we placed
memory targets inside the area of the defined PF (see *Memory-Guided Saccade Task* below). Once
a reliable PF was identified, we estimated their phosphene threshold of TMS intensity. Starting at
30% of maximum stimulator output we delivered 7 pulses at 20 Hz and recorded if a phosphene
was evoked. We increased the stimulation intensity by 10% until phosphenes were reliably evoked,
then reduced stimulation intensity by 5% to find the threshold defined as reported phosphene in 5
out of 10 trials. In the main experiment, we used a stimulation intensity at 110% of that threshold,
but only applied 3 pulses at 20 Hz. Note that the phosphene thresholds were determined using 7
pulses and in a dark room looking at a screen with a black background. During the main
experiment, participants received 3 pulses looking at a brighter screen with gray background.
Therefore, the phosphenes were rarely perceived. On rare occasions when participants did report
phosphenes, usually noted during the first couple trials of the first block, the intensity was lowered
until participants no longer perceived phosphenes. See below for details of Control Experiment 3,
which was designed to test if the visual perception of phosphenes impact memory.
345
346 **Memory-Guided Saccade Task**
347
348 Experiment 1: TMS during working memory maintenance. Middle Delay.
349
Participants performed a memory-guided saccade (MGS) task. Each block consisted of 40 trials
and began with a fixation cross, followed by a target presented for 500ms inside the PF, or the
mirrored hemifield of the PF, randomized across trials. Participants maintained in working
memory the location of the target over a 4s retention interval while fixating a central circular
fixation stimulus. When the fixation stimulus turned green, they made a saccade, guided by their
memory alone, to the location of the target. After 850ms, the target reappeared in green serving as
feedback. Experiments 1 and 2 were conducted over four days to minimize both fatigue and the
number of TMS pulses participants experienced in a given day. On day 0, participants performed
a 1 hour TMS session of phosphene thresholding, phosphene mapping (described above), and
practice of MGS task. On days 1-3, participants performed 5 blocks (200 trials; 600 TMS pulses)
of the MGS task during which their brain activity was recorded using EEG (see
Electroencephalography below). On 2 of these 3 days, TMS was applied to V1 during the middle
(at 2s) of the 4s long delay (what we refer to as *Middle Delay)*. On the other day, no TMS was
applied. For each participant, we randomized the order of the days on which TMS was applied.
364
Experiment 2: TMS during working memory encoding. Early and Middle Delay.

Each participant returned on another day to complete two sessions of the MGS task, while TMS
was applied to the V1 target area previously identified. EEG was not collected. The experiment
was identical to that described in Experiment 1, with the only difference being that TMS was
applied either at target offset (what we refer to as *Early Delay*) or in the middle of the delay (what
we refer to as *Middle Delay*). The order of sessions was randomized across participants. Each
session lasted 40 minutes and participants performed 5-7 blocks of MGS task (described above),
followed by a mandatory 10-minute break before they performed the next session lasting 40
minutes.

Control Experiment 3: Simulated phosphenes during working memory delay

Ten participants returned for another day of study to complete one session of MGS task where
TMS during the middle of the delay was replaced by simulated phosphenes. Specifically, the visual
appearance of phosphenes was simulated in the phosphene field reported by the participants during
the phosphene mapping task. A white noise mask was added to the phosphene field area, presented
as three brief flickers during the middle of the delay, simulating the three pulses of TMS during
experiments 1 and 2. In order to control for any potential hemispheric differences in performance,
simulated phosphenes were presented either on the same or opposite hemifield compared to the
target. The purpose of this experiment was to test the degree to which perceived phosphenes could
account for memory errors. Each participant completed 10 blocks of MGS in one session that
lasted 75 minutes and participants were compensated $15 for their time.

Eye-tracking and Visual Display

Participants were seated in a dim-lit room with the head positioned on a chin-rest 56 cm away from
a gamma-calibrated ViewPix monitor (120Hz refresh rate, 1920*1080 resolution) controlled by
an Ubuntu video card. Stimuli were presented using MATLAB and Psychtoolbox [48]. An Eyelink
1000 Desktop Mount (SR Research) was used to record the right eye position at 1000 Hz. For all
participants, a 9-point calibration was performed at the start of the session and repeated whenever
the participant took a break by moving the head from the chin-rest. The data were then
preprocessed using iEye (https://github.com/clayspacelab/iEye), an in-house gaze-analysis
toolbox written in MATLAB. The preprocessing involved multiple steps. First, blinks were
identified as pupil diameter lower than 1 percentile. Gaze data 100 ms before and after blinks were
removed. Next, a Gaussian smoothing kernel was applied to gaze traces to get an estimation of
velocity. This was used to detect saccades, which were defined as gaze transitions with velocities
greater than 30 deg/s, having an amplitude greater than 0.25 degree, and a duration of at least 7.5
ms. The data were then corrected for any instrumental drift by demeaning the gaze traces during
fixation and delay epochs. Lastly, we also performed calibration using the corrective saccades
during the feedback for any instrumental or head-movement-related drift that might have occurred

in a trial. Given eye-tracking was the sole metric of behavior, these extra precautions ensured
quality gaze data. Trials where stimulus timings were off (~0.2%) participants made large errors
(>10dva) (~4.6%), where the saccade reaction time was less than 150ms (likely suggesting
participants started saccade during delay epoch) (~0.54%), and trials where participants blinked
during the saccade or the eye tracker lost saccade traces during experimental setup failures (~8.9%)
were eliminated from analysis.
412
413   **Electroencephalography (EEG)**
414
We used a BrainProducts 64-channel actiChamp system with low-profile active electrodes for
recording EEG activity. Data were sampled at 1000Hz, with the ground placed at the center of
forehead above nasion and Cz was used as the online reference. EEG data were analyzed using
Fieldtrip toolbox [49]. Any trials with errors in timing of the TMS pulse were detected (less than 2
out of 400 trials on average per session) and removed from analysis. Then, the data for each session
were band-pass filtered between 0.5-50 Hz using a two-pass butterworth filter of filter order 4. The
data were then screened for bad channels. Bad channels were identified as ones that exhibited more
than 90 percentile of median variance amongst all channels, were flat (std < 0.01 µV), or were
extremely noisy (std > 100 µV) [50]. The cleaned data were then offline re-referenced to the common
average. Next, owing to long delays in our working memory task, the data were heavily
contaminated with blink artifacts. Independent Component Analysis (ICA) was performed using
the 'fastica' algorithm in Fieldtrip and components exhibiting muscle twitches, eye-movement
artifacts and saccades were carefully examined and eliminated. Bad channels removed earlier were
then interpolated using a weighted interpolation of neighboring electrodes. The data were then
epoched and downsampled to 200Hz. The epoched data were then examined for bad trials which
were then eliminated (less than 0.5% on average). Additionally, time-frequency analysis (TFA)
was conducted to investigate spectral dynamics. The analysis was performed using the wavelet
method. The frequency range was set from 2 to 40 Hz, discretized into 53 linearly spaced frequency
bins. The number of cycles for the wavelet analysis varied linearly with frequency, ranging from
4 to 15 cycles across the specified frequency range. Subsequently, the power values were converted
to decibel (dB) scale. Alpha Lateralization Index (αLI) was computed as the ratio of power
difference between the contralateral and ipsilateral activity for each trial from the occipito-parietal
electrodes (O1, O2, PO3, PO4, PO7, PO8, P1, P2, P3, P4, P5, P6, P7, P8) to the total power across
all occipito-parietal electrodes. $αLI = (P_{contra} - P_{ipsi}) / (P_{contra} + P_{ipsi})$ where, P_{contra} was power in an
electrode on trials where stimulus was present in contralateral hemifield, and P_{ipsi} was power in an
electrode on trials where stimulus was present in ipsilateral hemifield.
441
442   **Electrical field simulations**
443
Simulation of Noninvasive Brain Stimulation (SimNIBS) [20] 4.0 was used to estimate the electric
field induced in the brain based on anatomical scans and the target location chosen for each

participant as described above. Using CHARM-based head-segmentation [51] and target locations,
E-field estimates were computed for each individual brain taking into account conductivities of
different tissue types. The model uses the coil specifications provided by MagVenture to estimate
the magnetic field induced by the coil. For each participant, the exact placement of the TMS coil,
including the rotational angle was input to the model. This allowed us to create biophysical models
of the amplitude and extent of the electric field induced by TMS to V1 for each participant.
452
453 **Statistical Analyses**
454
Data analysis and statistics were performed using Python 3.11.5, Numpy 1.26.2, Scipy 1.11.4[52] for
behavioral statistics and Fieldtrip to perform cluster-based statistical analyses for EEG. No
statistical methods were used to predetermine sample size but we used sample size estimates from
similar previous studies[38]. To ensure a reliable measure of behavior, participants and trials were
eliminated as described in "Participants" and "Eye-tracking and Visual Display" sections.
Participants were carefully screened to be eligible for the study and hence the investigators were
not blinded to the selection of the participants. However, the assignment of perturbation was
randomized across participants and the investigators were blinded prior to the onset of the session.
463
For behavioral data, we randomly shuffled the trial labels for TMS conditions (No TMS, Early
TMS, or Middle TMS) and target location (inside PF or outside PF). For each permutation, we
computed the group average for each condition, thereby generating a null distribution of t-
statistics. The observed t-statistic was then compared against this null distribution to obtain the p-
value. In order to assess the contribution of radial and tangential components to saccade errors, we
first z-scored radial, tangential and absolute errors and performed an ordinary least squares
regression using statsmodel OLS to estimate the contribution of each component to the total error,
separately for each subject and condition. We then performed repeated-measures ANOVA using
statsmodel AnovaRM to assess the differences in the contribution of components to total error
across conditions.
474
475 For time-frequency EEG analyses, we employed a cluster-based permutation approach to perform
476 paired t-tests at each time-frequency bin using *ft_freqstatistics* function in Fieldtrip. The primary
477 alpha level was set to 0.05, while the cluster threshold was set at cluster-alpha of 0.1. Significance
478 of the clusters was assessed using Monte Carlo method with 1000 random permutations.
479
480

481
Data Availability
482
483
484 Source data to reproduce bar plots in all figures are provided as a Source Data file. All data used
485 in this study is made publicly available at https://osf.io/hf5a4/. The published dataset contains
486 preprocessed and de-identified behavioral data across both experiments, preprocessed time-
487 frequency analyzed EEG data, and the TMS-efield simulations and ROIs for each participant. Any
488 raw data can be made available upon request.
489
490 ## Code Availability
491
492 Code used to trigger the TMS and EEG systems is available at
493 https://github.com/clayspacelab/EEG_TMS_triggers. The toolbox iEye used to process eye-
494 tracking data is available at https://github.com/clayspacelab/iEye. And the codes required to
495 replicate figures from data are made available at
496 https://github.com/clayspacelab/Dake_Curtis_TMSV1_2025.
497
498
499

References

1. Conway, A. R. A., Kane, M. J. & Engle, R. W. Working memory capacity and its relation to general intelligence. *Trends Cogn. Sci.* **7**, 547–552 (2003).
2. Goldman-Rakic, P. S. Working memory dysfunction in schizophrenia. *J. Neuropsychiatry Clin. Neurosci.* **6**, 348–357 (1994).
3. Goldman-Rakic, P. S. Cellular basis of working memory. *Neuron* **14**, 477–485 (1995).
4. Fuster, J. The prefrontal cortex—an update time is of the essence. *Neuron* **30**, 319–333 (2001).
5. Funahashi, S., Bruce, C. J. & Goldman-Rakic, P. S. Mnemonic coding of visual space in the monkey's dorsolateral prefrontal cortex. *J. Neurophysiol.* **61**, 331–349 (1989).
6. Funahashi, S., Bruce, C. J. & Goldman-Rakic, P. S. Dorsolateral prefrontal lesions and oculomotor delayed-response performance: evidence for mnemonic 'scotomas'. *J. Neurosci.* **13**, 1479–1497 (1993).
7. Christophel, T. B., Klink, P. C., Spitzer, B., Roelfsema, P. R. & Haynes, J.-D. The Distributed Nature of Working Memory. *Trends Cogn. Sci.* **21**, 111–124 (2017).
8. Sreenivasan, K. K. & D'Esposito, M. The what, where and how of delay activity. *Nat. Rev. Neurosci.* **20**, 466–481 (2019).
9. Curtis, C. E. & Sprague, T. C. Persistent Activity During Working Memory From Front to Back. *Front. Neural Circuits* **15**, 696060 (2021).
10. Rahmati, M., DeSimone, K., Curtis, C. E. & Sreenivasan, K. K. Spatially Specific Working Memory Activity in the Human Superior Colliculus. *J. Neurosci.* **40**, 9487–9495 (2020).
11. Mackey, W. E., Devinsky, O., Doyle, W. K., Meager, M. R. & Curtis, C. E. Human Dorsolateral Prefrontal Cortex Is Not Necessary for Spatial Working Memory. *J. Neurosci.* **36**, 2847–2856 (2016).
12. Mackey, W. E., Devinsky, O., Doyle, W. K., Golfinos, J. G. & Curtis, C. E. Human parietal cortex lesions impact the precision of spatial working memory. *J. Neurophysiol.* **116**, 1049–1054 (2016).
13. Harrison, S. A. & Tong, F. Decoding reveals the contents of visual working memory in early visual areas. *Nature* **458**, 632–635 (2009).
14. Serences, J. T., Ester, E. F., Vogel, E. K. & Awh, E. Stimulus-Specific Delay Activity in Human Primary Visual Cortex. *Psychological Science* vol. 20 207–214 Preprint at https://doi.org/10.1111/j.1467-9280.2009.02276.x (2009).
15. Duan, Z. & Curtis, C. E. Visual working memories are abstractions of percepts. *Elife* **13**, (2024).
16. Woodry, R., Curtis, C. E. & Winawer, J. Feedback scales the spatial tuning of cortical responses during visual memory. *J. Neurosci.* (in press).
17. Rahmati, M., Saber, G. T. & Curtis, C. E. Population Dynamics of Early Visual Cortex during Working Memory. *J. Cogn. Neurosci.* **30**, 219–233 (2018).
18. Leavitt, M. L., Mendoza-Halliday, D. & Martinez-Trujillo, J. C. Sustained Activity Encoding Working Memories: Not Fully Distributed. *Trends in Neurosciences* vol. 40 328–346 Preprint at https://doi.org/10.1016/j.tins.2017.04.004 (2017).
19. Mackey, W. E., Winawer, J. & Curtis, C. E. Visual field map clusters in human frontoparietal cortex. *Elife* **6**, (2017).
20. Thielscher, A., Antunes, A. & Saturnino, G. B. Field modeling for transcranial magnetic stimulation: A useful tool to understand the physiological effects of TMS? in *2015 37th Annual International Conference of the IEEE Engineering in Medicine and Biology Society*

546   *(EMBC)* 222–225 (2015).

21. Hallenbeck, G. E., Sprague, T. C., Rahmati, M., Sreenivasan, K. K. & Curtis, C. E. Working memory representations in visual cortex mediate distraction effects. *Nature Communications* **12**, 4714 (2021).

22. Rademaker, R. L., Chunharas, C. & Serences, J. T. Coexisting representations of sensory and mnemonic information in human visual cortex. *Nat. Neurosci.* **22**, 1336–1344 (2019).

23. Medendorp, W. P. *et al.* Oscillatory activity in human parietal and occipital cortex shows hemispheric lateralization and memory effects in a delayed double-step saccade task. *Cereb. Cortex* **17**, 2364–2374 (2007).

24. van de Ven, V., Jacobs, C. & Sack, A. T. Topographic contribution of early visual cortex to short-term memory consolidation: a transcranial magnetic stimulation study. *J. Neurosci.* **32**, 4–11 (2012).

25. Cattaneo, Z., Vecchi, T., Pascual-Leone, A. & Silvanto, J. Contrasting early visual cortical activation states causally involved in visual imagery and short-term memory. *Eur. J. Neurosci.* **30**, 1393–1400 (2009).

26. van Lamsweerde, A. E. & Johnson, J. S. Assessing the Effect of Early Visual Cortex Transcranial Magnetic Stimulation on Working Memory Consolidation. *J. Cogn. Neurosci.* **29**, 1226–1238 (2017).

27. Rademaker, R. L., van de Ven, V. G., Tong, F. & Sack, A. T. The impact of early visual cortex transcranial magnetic stimulation on visual working memory precision and guess rate. *PLoS One* **12**, e0175230 (2017).

28. Zokaei, N., Manohar, S., Husain, M. & Feredoes, E. Causal evidence for a privileged working memory state in early visual cortex. *J. Neurosci.* **34**, 158–162 (2014).

29. Anderson, J. C., Kennedy, H. & Martin, K. A. C. Pathways of attention: synaptic relationships of frontal eye field to V4, lateral intraparietal cortex, and area 46 in macaque monkey. *J. Neurosci.* **31**, 10872–10881 (2011).

30. Blatt, G. J., Andersen, R. A. & Stoner, G. R. Visual receptive field organization and cortico-cortical connections of the lateral intraparietal area (area LIP) in the macaque. *The Journal of Comparative Neurology* vol. 299 421–445 Preprint at https://doi.org/10.1002/cne.902990404 (1990).

31. Paus, T. *et al.* Transcranial magnetic stimulation during positron emission tomography: a new method for studying connectivity of the human cerebral cortex. *J. Neurosci.* **17**, 3178–3184 (1997).

32. Bestmann, S., Baudewig, J., Siebner, H. R., Rothwell, J. C. & Frahm, J. Functional MRI of the immediate impact of transcranial magnetic stimulation on cortical and subcortical motor circuits. *Eur. J. Neurosci.* **19**, 1950–1962 (2004).

33. Ekstrom, L. B., Roelfsema, P. R., Arsenault, J. T., Bonmassar, G. & Vanduffel, W. Bottom-up dependent gating of frontal signals in early visual cortex. *Science* **321**, 414–417 (2008).

34. Romero, M. C., Davare, M., Armendariz, M. & Janssen, P. Neural effects of transcranial magnetic stimulation at the single-cell level. *Nat. Commun.* **10**, 2642 (2019).

35. Gentner, R. & Classen, J. Modular organization of finger movements by the human central nervous system. *Neuron* **52**, 731–742 (2006).

36. Awh, E. & Jonides, J. Overlapping mechanisms of attention and spatial working memory. **5**, 119–126 (2001).

37. Jerde, T. A., Merriam, E. P., Riggall, A. C., Hedges, J. H. & Curtis, C. E. Prioritized maps of space in human frontoparietal cortex. *J. Neurosci.* **32**, 17382–17390 (2012).

38. Fernández, A., Hanning, N. M. & Carrasco, M. Transcranial magnetic stimulation to frontal but not occipital cortex disrupts endogenous attention. *Proc. Natl. Acad. Sci. U. S. A.* **120**, e2219635120 (2023).

39. Curtis, C. E. & Connolly, J. D. Saccade preparation signals in the human frontal and parietal cortices. *Journal of Neurophysiology* **99**, 133–145 (2008).

40. Miller, L. M., Sun, F. T., Curtis, C. E. & D'Esposito, M. Functional interactions between oculomotor regions during prosaccades and antisaccades. *Hum. Brain Mapp.* **26**, 119–127 (2005).

41. Bergmann, T. O. & Hartwigsen, G. Inferring causality from noninvasive brain stimulation in cognitive neuroscience. *J. Cogn. Neurosci.* **33**, 195–225 (2021).

42. Fuster, J. M. & Jervey, J. P. Inferotemporal neurons distinguish and retain behaviorally relevant features of visual stimuli. *Science* **212**, 952–955 (1981).

43. Supèr, H., Spekreijse, H. & Lamme, V. A. A neural correlate of working memory in the monkey primary visual cortex. *Science* **293**, 120–124 (2001).

44. van Kerkoerle, T., Self, M. W. & Roelfsema, P. R. Layer-specificity in the effects of attention and working memory on activity in primary visual cortex. *Nat. Commun.* **8**, 13804 (2017).

45. Huang, J. *et al.* Neuronal representation of visual working memory content in the primate primary visual cortex. *Sci. Adv.* **10**, eadk3953 (2024).

46. Dumoulin, S. O. & Wandell, B. A. Population receptive field estimates in human visual cortex. *Neuroimage* **39**, 647–660 (2008).

47. Kay, K. N., Winawer, J., Mezer, A. & Wandell, B. A. Compressive spatial summation in human visual cortex. *J. Neurophysiol.* **110**, 481–494 (2013).

48. Brainard, D. H. The Psychophysics Toolbox. *Spat. Vis.* **10**, 433–436 (1997).

49. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput. Intell. Neurosci.* **2011**, 156869 (2011).

50. Komosar, M., Fiedler, P. & Haueisen, J. Bad channel detection in EEG recordings. *Current Directions in Biomedical Engineering* **8**, 257–260 (2022).

51. Puonti, O. *et al.* Accurate and robust whole-head segmentation from magnetic resonance images for individualized head modeling. *Neuroimage* **219**, 117044 (2020).

52. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–272 (2020).

Acknowledgments

We thank the staff of the Center for Brain Imaging (CBI) at New York University (NYU), and Yuyang Xu and Qingqing Yang for help with data collection. We are also grateful to Nathan Tardiff and Zhengang Lu for their comments on drafts of the manuscript. Funded by NIH R01 EY-016407 and R01 EY-033925 to C.E.C.

Author Contributions

M.D. performed the experimental studies, analyzed the data, and wrote the manuscript. C.E.C. conceived the experiments, supervised the work, and wrote the manuscript.

Competing Interests

The authors declare no competing interests.

638
Figure Legends
Figure 1. Working memory task conditions, trial types, and TMS procedures. A. In each
participant, V1 was identified using population receptive field mapping [19]. Dorsal V1, which
represents the contralateral lower visual field (LVF), was accessible to TMS stimulation (asterisk
marks TMS target), while the upper visual field (UVF) is buried deeper and not accessible. The
bottom radial plot depicts the group average visual field coverage of the dorsal V1 accessible to
TMS, whose pRFs are localized to the lower visual field. **B.** The modeled electrical field induced
by TMS targeting the retinotopically-defined left V1 (magenta encircled) for an example
participant. V1 in the hemisphere not perturbed with TMS encircled by an orange line. See Supp
Figure 1 for simulated electrical fields for all participants. Simulated electric field in Volts/meter
(V/m). **C.** Based on the model [20], TMS induced a significantly larger electric field in the stimulated
V1 compared to the V1 in the other hemisphere, (paired-*t* test, two-sided, $t(14)=4.677$, $p=0.00035$,
$d=2.50$, $n=15$ humans). Thin gray lines are individual participants (n=15), bars are means (+/-
SEM) across participants. **D.** In each participant, phosphene fields were defined as the portion of
the visual field in which phosphenes were reliably evoked. TMS to dorsal V1 produced phosphenes
in the contralateral lower visual field. Prior to memory experiments, participants drew with a
mouse the extent of TMS-induced phosphenes (top). During the memory experiments, we placed
visual targets in the phosphene field, and out of it (in the mirrored location) (bottom). **E.**
Participants performed a memory-guided saccade (MGS) task, during which they maintained in
working memory a target location. After a retention interval of 4 seconds, participants generated
a memory-guided saccade. Feedback was then given, followed by an intertrial interval (2-3 s; not
shown). We measured the accuracy of memory-guided saccades for targets in the phosphene field
(experimental condition; magenta). Control conditions included no TMS (blue) and TMS to V1
when targets were in the hemifield opposite to the phosphene field (orange). **F**. Trial schematics
for Experiment 1 and Experiment 2. Data are provided in the Source Data file.
664
Figure 2. Effects of TMS to V1 on working memory A. In Experiment 1, TMS applied to V1
during the middle of the delay period caused an increase in memory errors for targets in the
phosphene field (in), compared to when no TMS was applied and to memory for targets in the
opposite hemifield (out). In Experiment 2, we replicated the results from Experiment 1 by again
showing that TMS to V1 during the middle of the delay caused a specific worsening of memory
for targets in the phosphene field. Moreover, we found similar effects when TMS to V1 was
delivered as soon as the visual target was extinguished (early) and when it was applied later in the
middle of the delay (middle). Bars depict means, with SEM error bars, and circles are individual
participants (n = 15). (* denote two-sided paired *t*-tests $p<0.05$). **B.** EEG collected during
Experiment 1 were used to compute an alpha lateralization index (αLI), sensitive to the topography
of alpha power in electrodes over occipital cortex related to the location of working memory
targets. Without TMS, alpha topography tracked the remembered targets evidenced by a decrease
in alpha power contralateral to the target (Delay 1). This pattern persisted late into the delay period
(Delay 2). Following TMS applied to V1 during the middle of the memory delay (2000 ms), note
how the strength of the lateralized alpha decreased (Delay 2). **C**. EEG time-frequency responses
(TMS minus no TMS) during the working memory delay for occipital electrodes over the
hemisphere being stimulated. Following the TMS evoked artifact (large red broadband increase),
power significantly decreased in alpha and beta bands. Yellow outlines significant clusters of
power differences between TMS minus no TMS (cluster-based permutation two-sided paired t-

test, corrected for multiple comparisons, $p<0.05$). This further explains the effect of TMS on αLI.
Gray lines along the timeline denote the Delay 1 and Delay 2 epochs in B. Source Data are
provided in the Source Data file.
687
688

A.

UVF

LVF

V3d
V2d
V1

*

24°
16°
8°

V1
LVF

B.

TMS
V1

Other
V1

0   V/m   1.5

C.

*

Modeled electric field (V/m)

2.0

1.5

1.0

0.5

0.0

TMS   Other

V1 hemisphere

D.

Phosphene
Drawings

out   in

Phosphene
Field

E.

Target
500 ms

Delay
4 s

Response
850 ms

Target
500 ms

Delay
4 s

Response
850 ms

No TMS

V1 TMS

MGS in Phosphene Field

MGS out Phosphene Field

F.

Exp. 1: No TMS and TMS middle of delay

Target

TMS

MGS

0   1   2   3   4   5   6

Time (s)

Exp. 2: TMS early and middle of delay

TMS

Target

TMS

MGS

0   1   2   3   4   5   6

Time (s)

A. Exp. 1   Exp. 2

Memory Error (deg)

No TMS / Middle delay (in / out)

Condition

Early delay (in / out) / Middle delay (in / out)

Condition

B. Delay 1 500-1500ms   Delay 2 3000-4000ms

No TMS

V1 TMS

$-.05$ $.05$

$$\alpha_{LI} = \frac{(P_{contra} - P_{ipsi})}{(P_{contra} + P_{ipsi})}$$

C.

Frequency (Hz)

Power (dB)

Target   Delay 1   TMS   Delay 2

Time (s)