Cortical synaptic vulnerabilities revealed in a α-synuclein aggregation model of Parkinson's disease

Saroj Sah [1,7], Andrew D. Sauerbeck [2,7], Jyoti Gupta [3,7], Dayana Pérez-Acuña [1,7], Jacob E. Reiber [5,7], Dreson L. Russell [5,7], Thomas M. Goralski [6,7], Michael X. Henderson [6,7], Laura A. Volpicelli-Daley [5,7], Michael J. Higley [3,4,7], Terrance T. Kummer [2,7] and Thomas Biederer [1,3,7] *

[1] Department of Neurology, Yale School of Medicine, New Haven, CT 06510, USA
[2] Department of Neurology, Washington University School of Medicine, St. Louis, MO 63110, USA
[3] Department of Neuroscience, Yale School of Medicine, New Haven, CT 06510, USA
[4] Department of Biomedical Engineering, Yale School of Medicine, New Haven, CT 06510, USA
[5] University of Alabama at Birmingham, Birmingham, AL 35294, USA
[6] Department of Neurodegenerative Science, Van Andel Institute, Grand Rapids, MI 49503
[7] Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD

* Correspondence should be addressed to Thomas Biederer, thomas.biederer@yale.edu

Running Title: Cortical synaptopathy in Parkinson's disease
Keywords: Parkinson's disease; α-synuclein; cortex; synapse; excitation/inhibition imbalance

Abstract

Cognitive impairment is a frequent non-motor symptom in Parkinson's disease, and cortical Lewy pathology is strongly associated with cognitive decline. Synaptic pathology has been observed in the PD cortex, but the extent of synaptic vulnerabilities and their temporal and spatial relationship to pathology remains unclear. We employed high-resolution imaging to analyze synaptic abnormalities in layer 5 of the secondary motor cortex. We used striatal injections of α-synuclein pre-formed fibrils as a model to cause the progressive pathological aggregation of endogenous α-synuclein. We find that cortical α-synuclein pathology results in the progressive loss of excitatory synapses, followed by a reduction in inhibitory postsynaptic sites. Synapse loss is most pronounced in areas with high pathology. Additionally, we observed ultrastructural changes in the remaining excitatory synaptic loci, including smaller synaptic vesicles. Consistent with these results, gene ontology analysis of synaptic genes exhibiting altered expression in pathological neurons supported pre- and post-synaptic changes, including in synapse organizing pathways. Our results demonstrate that α-synuclein aggregation in the cortex is linked to molecular and structural alterations that disrupt synaptic connectivity and provide insights into the progressive PD-relevant vulnerability of cortical synapses.

Introduction

Parkinson's disease (PD) features both motor and non-motor impairments (Poewe et al., 2017). Up to 80% of PD patients develop cognitive changes, with a 6-fold increased risk of progression to dementia (Aarsland et al., 2021; Buter et al., 2008; Hely et al., 2008). Cortical Lewy pathology correlates with cognitive decline in PD (Braak et al., 2006; Irwin et al., 2017; Irwin et al., 2012; Ryman et al., 2021). Deficits in cognition have already been observed at the early stages of the disease, with a spectrum of abnormalities such as impaired working memory, visuospatial functions, and language (Caviness et al., 2011). The underlying molecular and cellular aberrations in the neocortex are mainly unknown.

A significant hallmark of the disease is the abnormal aggregation of the protein α-synuclein that is endogenously enriched at presynaptic sites into Lewy bodies and Lewy neurites. Lewy bodies and neurites are present across multiple brain regions beyond the substantia nigra, including limbic and cortical regions, and their abundance strongly correlates with cognitive decline and psychiatric symptoms in PD (Irwin et al., 2012; Schapira et al., 2017). Cortical dysfunction and morphological alterations have been described in PD (Bruck et al., 2005; Caviness et al., 2011; Li et al., 2022), supporting the relevance of neocortical circuit dysfunction in cognition. Propagation of misfolded α-synuclein is a key pathological event in PD, where abnormally aggregated α-synuclein species serve as templates that seed the conversion of monomeric α-synuclein into inclusions as the disease progresses (Braak et al., 2003; Recasens et al., 2014). This phenomenon has been modeled and reproduced *in vivo* using pre-formed fibrils (PFFs) comprised of *in vitro* assembled β-sheet-rich α-synuclein. Once injected into the brain, PFFs recapitulate aspects of PD Lewy body pathology (Henderson et al., 2019), causing dysregulation in multiple cellular and molecular pathways (Goralski et al., 2024) and leading to neuroinflammation (Earls et al., 2019; Iba et al., 2022), metabolic impairment (Creed et al., 2022; Lu et al., 2022) and circuit dysfunction (Blumenstock et al., 2021). This can be analyzed to measure the progression of pathology between anatomically connected neuronal populations, including in cortical neurons projecting to the striatum (Henderson et al., 2019; Luk et al., 2012).

α-Synuclein is enriched in the presynaptic terminal, and the three members of this protein family shape synaptic architecture (Vargas et al., 2017). This suggested that synapses could be

vulnerable sites for α-synuclein pathology and synaptopathy could be among the first pathological processes in multiple neurodegenerative diseases, including PD (Schirinzi et al., 2016). Indeed, brain imaging studies have shown a loss of synaptic markers in PD patients (Matuskey et al., 2020), and dendritic spine loss has been reported in other synucleinopathies (Kramer and Schulz-Schaeffer, 2007). Post-mortem studies of PD, PDD, and Dementia with Lewy Bodies show reduced synaptic marker density in the cortex that is correlated with the extent of Lewy pathology (Frigerio et al., 2024; Thomsen et al., 2021). It has been proposed that a reduction in synaptic vesicle proteins in the prefrontal cortex could be a potential cause of dementia in patients with PD (Holmes et al., 2024). Indeed, in experimental models, it has been reported that α-synuclein aggregation impairs excitatory neuron transmission and connectivity (Blumenstock et al., 2017; Chen et al., 1996; Froula et al., 2018; Volpicelli-Daley et al., 2011).

Here, we aim to define synaptic aberrations in PD to understand alterations in cortical connectivity that can underlie cognitive impairments. As an approach, we use intrastriatal PFF injection to address the extent to which cortical synapses are a cellular locus for dysfunction in PD. We analyze layer 5 of the secondary motor cortex, as it includes intratelencephalic (IT) neurons that project to the striatum as a subpopulation of neurons that preferentially develops α-synuclein aggregates (Goralski et al., 2024), including spine and synapse loss and exhibit dendrite malformation (Blumenstock et al., 2017; Pérez-Acuña et al., 2023). Tracking the gradual increase in α-synuclein pathology, we measured a reduction of synapses proximal to areas with high pathology. SEQUIN high-resolution imaging determined that cortical pathology impacts excitatory synapses after a lag period, and they were then progressively lost. Structural changes were observed in the remaining excitatory synapses, including physical separation of synapses and enlargement of pathology-bearing presynaptic sites. Using EM, we observed reduced synaptic vesicle volumes and shortened postsynaptic sites. After excitatory synapse pathology appeared, inhibitory postsynaptic sites were lost. These pathologies matched our synaptic gene ontology analysis of gene expression alterations in pathology-bearing neurons conserved between PD brain and mice injected with α-synuclein PFFs (Goralski et al., 2024). We additionally determined that cortical α-synuclein pathology in this PFF model results in substantial aberrations of excitatory transmission. Together, this work delineates synapse-type

75   vulnerabilities in the cortex that result from α-synuclein pathology on molecular, cellular, and

76   functional levels.

Results

Progressive excitatory synapse loss in the prefrontal cortex is associated with α–synuclein pathology. Both human and PFF mouse models have revealed that excitatory neurons are most likely to bear a higher abundance of α– synuclein inclusions in layer 5 of the prefrontal cortex. Furthermore, it has been validated in the PFF mouse model that α-synuclein-inclusion-positive excitatory neurons have downregulated differentially expressed genes (DGEs) for VGlut1 and Homer (Goralski et al., 2024). Thus, we analyzed the distribution of excitatory synapses by measuring the immunoreactivity of VGluT1 and Homer using SEQUIN in the layer 5 prefrontal cortex of mice injected into the dorsolateral striatum with PFFs. Monomer-injected mice served as controls.

Mice exhibited α–synuclein inclusions in the prefrontal cortex 1 month post-injection (MPI) of PFFs in the striatum as reported, which plateaued at 3 months, while no inclusions appeared after monomer injection, which served as control (Luk et al., 2012) (Figure 1A and Supplementary Fig 1). SEQUIN identifies synaptic loci by recognizing separations of pre- and post-synaptic markers consistent with ultrastructural parameters and utilizes an enhanced image resolution methodology (Sauerbeck et al., 2020). Staining for the excitatory pre- and post-synaptic markers VGluT1 and Homer followed by SEQUIN analysis showed no apparent change in the density of pre-and post-synaptic pairs containing both VGluT1/Homer in striatally PFF-injected mice compared with monomer-injected mice at 1 MPI (Fig.1B). Interestingly, we identified that the abundance of paired excitatory VGluT1/Homer synaptic loci was lower in the prefrontal cortex 3 months after striatal PFF injection compared to monomer-injected mice (monomer: 0.086 ± 0.003 / μm^3 vs PFF: 0.076 ± 0.003 / μm^3, p = 0.042) (Fig.1C). The decrease in excitatory synaptic loci persisted at 6 MPI, with a reduction in density (monomer: 0.066 ± 0.002 / μm^3 vs PFF: 0.060 ± 0.002 / μm^3, p=0.018) (Fig.1D)

Fewer total excitatory specializations at 3 months post-PFF injection. Analysis of the SEQUIN imaging data by Imaris showed no change in the density of total VGluT1 and Homer puncta at 1 MPI, irrespective of whether they were within the synaptic loci (Fig.2A). The loss of synapses at

3 MPI was coupled with a decreased number of total VGluT1 puncta and Homer puncta (Fig.2B).

No changes in the VGluT1 and Homer puncta were observed at 6 MPI (Fig.2C). This could indicate

that a compensatory improvement in the alignment of synaptic markers at later stages of

α–synuclein pathology.

A reduction in cortical inhibitory postsynaptic sites follows excitatory synapse loss. The extent

of the vulnerability of different synapse subtypes to α–synuclein pathology is unknown. We

performed SEQUIN analysis to study whether inhibitory synapses are affected by synuclein

inclusion in the prefrontal cortex of PFF-injected mice compared to control mice injected with

monomers. Staining for inhibitory presynaptic VGAT and inhibitory postsynaptic Gephyrin

revealed no significant changes in the density of synaptically paired or total VGAT or Gephyrin at

3 MPI (Fig.3A). However, the number of postsynaptic gephyrin puncta was reduced at 6 MPI

(Fig.3B). This impact on inhibitory synaptic markers was therefore subsequent to the loss of

excitatory synapses and may reflect a homeostatic balancing mechanisms to partially restore

excitation/inhibition balance rather than direct synaptic pathology caused by a-synuclein

aggregates.

Localization of phospho-α-synuclein aggregates relative to synaptic loci after PFF injection. We

next addressed to what extent synapses are marked by a-synuclein pathology. One month after

either monomer or PFF injection into the striatum, the synaptic proteins PSD-95 and synapsin, as

well as phosphorylated α-synuclein, were imaged at enhanced resolution in 3-dimensional

volumes of the fronal cortex (Fig.4A). Co-staining with an antibody against pS129 phospho-α-

synuclein that marks the aggregated pathological form of this protein was detected in close

association with apparent synaptic loci (Fig.4B). To quantify synaptic localization of phospho-α-

synuclein, SEQUIN was employed to identify synaptic loci and measure the amount of phospho-

α-synuclein associated with each locus (within a 200 nm field centered on the centroid of each

SEQUIN-identified synaptic locus) (Fig.4C). Compared to monomer treatment, an elevation in

synapse associated phospho-α-synuclein was observed following PFF treatment (Fig.4D, PFF-

effect, p=0.1 Mann-Whitney U test.). Regardless of the general level of pathology, all areas

assessed exhibited an increase in synapse-associated phospho-α-synuclein relative to

parenchymal levels, indicative of specific synaptic enrichment (Fig.4D). The ratio between

synapse-associated and parenchymal phospho-α-synuclein tended to be higher as the overall

level of pathology increased (Fig.4E, p=0.0786 Kruskal-Wallis on group average).

Mesoscale analysis of the impact of phospho-a-synuclein pathology on synaptic density after

PFF injection. We aimed to complement the high-resolution analysis of synaptic puncta with a

mesoscale quantification of how α-synuclein pathology impacts synaptic distribution. Cortical

mesoscans of synaptic proteins and phospho-α-synuclein were collected from animals 1 MPI of

either monomer or PFF (Fig.4F-H). For both the monomer and PFF treatment groups, average

heatmaps of synaptic density and synapse-associated phospho-α-synuclein were generated

(Fig.4I-L). Areas of elevated synapse-specific phospho-α-synuclein in these heatmaps were

associated with an overall reduction in synaptic density, a trend that was also apparent on

quantitative analysis (Fig.4M, note the modest group sizes and model variability preclude

drawing firm conclusions based on statistical analysis: p=0.4 Mann-Whitney U test, Fig.4N p= 0.1

Mann-Whitney U test). To explore these effects further, within animal analysis of synaptic density

and synapse-associated phospho-α-synuclein levels were performed (Fig.4O-Q). In two of the

three PFF-treated subjects, a significant, negative correlation was detected between phospho-α-

synuclein level and synaptic density (Fig.4O,P), while a third animal exhibited a positive

correlation (Fig.4Q). These data provide further evidence of synapse loss in response to

synuclein-related pathology.

Synapse-associated phospho-α-synuclein and synaptic density are associated with proximity

to Lewy neurites. We next addressed the spatial relationship of cortical phospho-a-synuclein

aggregates formed after striatal PFF injection and synaptic aberrations. At 1 MPI of PFF, synapses

were identified using SEQUIN (Fig.5A). In these same areas, phospho-a-synuclein was imaged and

used to identify Lewy neurites (Fig.5A-B). The distance to the closest Lewy neurite edge was

measured for each synaptic centroid (Fig.5C). A significant association was observed between the

proximity of a synaptic locus to a Lewy neurite and the enrichment of phospho-a-synuclein at the

same synaptic locus (Fig.5D). To gain insight into the implications of this effect, the density of synaptic loci with 800 μm of the perimeter of Lewy neurites with varying degrees of pathology (as assessed by total intra-neuritic phospho-α-synuclein levels) was quantified. This analysis revealed a significant, negative association between synaptic density and Lewy neurite pathology (Fig.5E). These results suggest that Lewy neurites locally impact synapses, leading to their loss.

Analysis of dysregulated synaptic pathways in $\pi\eta o\sigma\pi\eta o-\alpha-\sigma\psi\nu\upsilon\chi\lambda\epsilon\iota\nu$ bearing neurons using SYNGO database. We further analyzed gene enrichment of synaptic pathways in pS129 phospho-α-synuclein bearing neurons from previous transcriptomic results (Goralski et al., 2024) by using SynGO, a database for synaptic ontology (Koopmans et al., 2019). After comparing the LAMDA (Lewy-associated molecular dysfunction from aggregates) signature from 3 months after injection mice, against the 1602 annotated genes in the SYNGO database, we found that 1306 genes were identified in this transcriptomic analysis. Among these, 469 were differentially expressed genes, with 418 of them downregulated and 51 upregulated in phospho-α-synuclein positive neurons (Fig.6A and Supplemental Fig.2). Enrichment analysis from genes downregulated showed that by cellular component, genes from postsynaptic density and synaptic vesicles were overrepresented (Fig.6B). Biological processes from these DEGs corresponded to the synaptic organization, presynaptic vesicle cycle, and processes in the post-synapse (Fig.6C). Markers used for our synaptic density analyses, vGlut1 and Homer1, were downregulated. In the case of the markers vGAT and Gephrin, we did not find a decrease in their expression, which was expected for vGAT as pathological neurons are excitatory.

For upregulated genes, cellular component enrichment, similar to downregulated genes, included terms annotated in synaptic specialization and postsynaptic membranes (Fig.4D). Biological processes also included synaptic organization, processes in the post-synapse, and synaptic signaling. We noted enrichment of GPC6 (glypican 6) and TNM2 (teneurin 2), regulators of neurotransmitter receptor localization for postsynaptic specialization and synapse maintenance, suggesting that neurons with pathology might activate response mechanisms to promote and sustain synaptic activity.

A separation of excitatory pre- and post-synaptic specializations is accompanied by ultrastructural changes. Based on our SYNGO analysis of DEGs, we hypothesized that synuclein inclusions may also contribute to nanostructure aberrations in the synapses. Using SEQUIN, we measured the average pre- and post-synaptic separation for VGluT1 and Homer at 1 MPI and at 3 MPI, the timepoint when we observed a decrease in excitatory synapses. The Gaussian distribution of the pre- and post-synaptic separation for synaptic loci showed no change at 1 MPI but revealed an increased mean separation for VGluT1/Homer pairs in the prefrontal cortex at 3 MPI of PFF fibrils compared to monomers in mice (Fig.7A,B).

Our findings were further supported by ultrastructural analysis using transmission electron microscopy, which demonstrated a reduction in the size of synaptic vesicles in PFF-injected mice compared to control mice (Fig.7C). These nanostructure changes could potentially contribute to the increased separation between pre- and post-synaptic proteins in excitatory synapses positive to VGluT1/Homer.

Discussion

The abnormal accumulation of α−synuclein into inclusions, known as Lewy bodies, is a major hallmark of PD and responsible for dopaminergic neuronal loss. These inclusions and neurites are widespread across brain regions beyond the substantia nigra, including limbic and cortical regions. Their presence strongly correlates with cognitive decline in PD (Irwin et al., 2012; Schapira et al., 2017). PD is associated with cortical dysfunction and morphological alterations (Bruck et al., 2005; Caviness et al., 2011; Li et al., 2022), highlighting the relevance of neocortical circuit dysfunction in cognitive impairment. A reduction in the presynaptic marker synaptophysin in the pyramidal layer of the neocortex is considered a significant factor in the development of dementia in Parkinson's disease (Zhan et al., 1993). A recent study confirmed the loss of synaptic vesicle glycoprotein 2A in the cortex in both Parkinson's disease dementia and dementia with Lewy bodies, supporting that pathological changes and subsequent synapse loss extend beyond the substantia nigra to affect the cortex, contributing to cognitive impairment (Andersen et al., 2021). However, the specificity of synapse loss and the nanostructural aberrations contributing to mechanobiological changes contributing with cognitive deficits remain poorly investigated. The prefrontal cortex is considered the essential brain region that is affected by PD cognitive impairment (Cai et al., 2021). To address these pathophysiological aspects of synapse-specific aberrations in the prefrontal cortex, we injected pre-formed fibrils (PFFs) into mice to replicate cortical alpha-synuclein inclusion pathology resembling Lewy bodies and Lewy neurites found in humans, and briefly investigated resulting synaptic aberrations in this region.

We observed α−synuclein inclusions in the secondary motor prefrontal cortex after 1 month of PFF injection in the striatum, although there was no change in synapse loci nor in the number of pre-/postsynaptic proteins positive for VGluT1/Homer. By the 3-month mark, we noted a reduction in excitatory synaptic loci accompanied by decreased levels of both VGluT1 and Homer. This suggests that the loss of excitatory synapses at this time point may be attributed to prolonged exposure to p-Syn inclusions and progressive synuclein pathology. Our data further revealed persistent synapse loss 6 months post-PFF injection, although changes in VGluT1 and Homer puncta were compensated. Analysis using the SYNGO database indicated upregulation of specific genes promoting synaptic protein formation in response to PFF injection, supporting

compensatory upregulation of synaptogenic pathways and the generation of orphan synaptic

sites. These findings underscore the relevance of overall excitatory synapse loss during the later

stages of disease pathology development. The enrichment of pathological phospho-α-synuclein

at synapses may contribute to their widespread loss following PFF injection.

Our in vivo findings align with previously reported in vitro data demonstrating that PFF

treatment compromises excitatory synaptic activity in a time- and dose-dependent manner. The

severity of these deficits correlates with the formation of αSyn pathology in cultured excitatory

hippocampal neurons (Wu et al., 2019). We observed the least pronounced effect on

postsynaptic inhibitory proteins, with no apparent overall loss of inhibitory synapses. This

minimal or absent impact on inhibitory synapses may be due to fewer synuclein inclusions in

inhibitory neurons compared to excitatory neurons in layer 5 of the secondary motor cortex

(Goralski et al., 2024). where we also conducted synaptic aberration analysis. Our results

highlight differential vulnerabilities across synapse types, reflecting varied expression of alpha-

synuclein pathology within specific neuron populations in layer 5 of the secondary motor cortex.

We identified an increased separation between VGluT1 and Homer in the cortex after 3 and

6 months PFF post-injection. This stretched separation indicates a potential shift in the

localization of pre- and postsynaptic proteins or a decrease in network connectivity at synaptic

loci, which can potentially lead to defects in neurotransmission and synapse stability (Frye et al.,

2021). Furthermore, reductions in synaptic vesicle and postsynaptic density sizes at synaptic loci,

as shown by ultrastructural analysis, could also be a cause of these aberrant phenotypes. These

alterations to synaptic architecture may contribute to behavioral deficits (Gcwensa et al., 2024),

including cognitive impairment. The downregulation of pathways involved in synapse

organization, as indicated by analysis of dysregulated synaptic pathways in phospho-α-synuclein

bearing neurons using SYNGO database, in PFF-injected mice could be the mechanistic approach

involved in the nanostructural aberrations we observed within excitatory synapses.

Materials and Methods

Antibodies. Primary antibodies and application notes for immunohistochemistry (IHC) are provided in Table 1. Secondary antibodies, including isotype-specific ones to detect monoclonal primary antibodies, were conjugated to Alexa dyes 405, 488, 568, and 647 (Thermo Fisher) and incubated overnight at 4 °C.

Target	Vendor	Cat. No.	RRID	Host	Clonality	Application IHC
VGluT1	Synaptic Systems	135 304	AB_887878	Guinea Pig	Polyclonal	1:200 (48 h, RT)
Homer	Synaptic Systems	160 003	AB_887730	Rabbit	Polyclonal	1:250 (48 h, RT)
VGAT	Synaptic Systems	131 003	AB_887869	Rabbit	Polyclonal	1:250 (36 h at 4 °C)
Gephyrin	Synaptic Systems	147 111	AB_887719	Mouse	Monoclonal IgG1	1:250 (36 h at 4 °C)

Table 1. Primary and secondary antibody applications.

Animals. Studies were performed using 3-month-old wild-type C57BL/6J mice obtained from Jackson Laboratories (Bar Harbor, ME, USA). Both female and male mice were included in the study unless otherwise stated. All animal procedures undertaken in this study to generate SEQUIN or immunochemistry data were approved by the Yale University Institutional Animal Care & Use Committee and complied with the National Institutes of Health guidelines for the Care and Use of Laboratory Animals.

Preparation of recombinant α-synuclein monomer and preformed fibrils. Monomeric mouse α-synuclein was expressed in E. coli and purified, followed by endotoxin removal using Pierce LAL high-capacity endotoxin removal resin, as previously described (Volpicelli-Daley et al., 2014). The monomer was stored at −80°C. Preformed fibrils (PFFs) at a concentration of 5 mg/mL were produced from monomeric α-synuclein by agitation in 150 mM KCl/50 mM Tris-HCl at 37°C for seven days (Bousset et al., 2013; Stoyka et al., 2020). PFFs in a volume of 22-25 µL were placed in a 1.5 mL sonicator tube (Active Motif, NC0869649) and sonicated for 15 min at an amplitude of 30, with a pulse on and off for 3 sec and 2 sec, respectively. The resulting PFF fragments, ranging from 20 to 70 nm, were verified by dynamic light scattering and stored at -80°C until needed. On the day of injection, the sonicated PFFs were re-sonicated for 15 min at an amplitude of 40, with a pulse on and off for 3 s, respectively, at 10°C and used within 8 h. The PFFs were kept at room temperature during the injections. In contrast, the monomer was kept on ice during the injections to prevent aggregation.

Stereotaxic injections of recombinant proteins. Mice were deeply anesthetized using isoflurane vaporizer (Harvard Apparatus Mobile Anesthesia System, model 750239) and placed on a

stereotactic frame (Kopf, model 900 LS). Thereafter, mice were unilaterally injected with 2 µL (5
mg/mL) of either PFFs or monomer (as a control) into the right forebrain at coordinates +0.2 mm
relative to Bregma, +2.0 mm from midline, and -2.6 mm beneath the dura to target the dorsal
striatum. The injection was performed using a Nanoject III microinjector (Drummond Scientific)
at a rate of 20 nL/s over 100 cycles with a delay of 8 s between cycles. The needle remained
undisturbed for 10 min after the protein delivery inside the dorsal striatum and was gradually
withdrawn to prevent backflow or leakage into the cortex.
299
Tissue processing for SEQUIN synapse quantification. SEQUIN enables enhanced resolution of
synapses, allowing to measure pre- and post-synaptic pairs on a 3D mesoscale (Sauerbeck et al.,
2020). SEQUIN multiscale imaging was used to quantify synaptic loci in layer 5 of the prefrontal
cortex and characterize synaptic nanostructural properties at the individual and population
levels, as previously described (Reitz et al., 2021). In brief, mice injected with either monomer or
PFFs, 1, 3, or 6 months post-injection, were deeply anesthetized by intraperitoneal ketamine (100
mg/kg) and xylazine (10 mg/kg) cocktail injection and transcardially perfused using a 27G needle
at a rate of 6 ml/min with ice-cold 1x Phosphate Buffered Saline (PBS) for 4 min, followed by ice-
cold 4% formaldehyde (PFA) in PBS (pH 7.6) for 4 min. The brains were then extracted and post-
fixed for 24 h at 4°C in 4% PFA in PBS and stored in PBS at 4°C in scintillation vials protected from
light. Fixed brains were sectioned coronally into 40 µm-thick slices using a vibratome (Leica
VT1000 S). Brain slices were stored in 24-well plates in cryoprotectant buffer (PBS containing 30%
ethylene glycol, 30% sucrose, and 0.02% sodium azide) at −20°C.
Slices from the prefrontal cortex were washed 3 times for 5 min each in PBS, then blocked in
20% normal goat serum (Jackson ImmunoResearch) in PBS for 1 h at room temperature (RT).
Slices were then immunolabeled with primary antibodies and washed 3 times for 5 min each in
PBS, followed by Alexa Fluor-labeled secondary antibodies prepared in PBS containing 10% NGS
and 0.3% Triton X-100 in the dark (details on antibodies used and conditions are listed in Table
1). The slices were washed 3 times for 5 min each in PBS to remove the unbound secondary
antibodies, counterstained with DAPI in PBS for 10 min, rinsed 3 times for 5 min each in PBS,
transferred to charged microscopic glass slides, dried at RT, and rinsed in distilled water to
remove PBS residues. The slides were dried at RT and embedded in mounting media freshly
prepared by mixing Tris-MWL 4-88 (Electron Microscopy Sciences #17977-150, Hatfield, PA, USA)
with AF300 (Electron Microscopy Sciences #17977-25) in a 9:1 ratio with 1.5H coverglass. The
coverglass used for SEQUIN was stain-free, thoroughly cleaned with 100% ethanol, rinsed with
distilled water, and air-dried prior to coverslipping. The slides were cured at RT for 3 days.
326
SEQUIN imaging of synapses and image analysis. Images were obtained, processed, and
analyzed as previously described (Reitz et al., 2021) using a Zeiss LSM 900 Airyscan Microscope
(Carl Zeiss Microscopy) with 63x (numerical aperture 1.4) oil immersion objectives (6-8 images
per animal from 3 slices) and processed in 3D Airyscan. The channel alignment of the Airyscan-
processed images was inspected and corrected if required. Pre- and post-synaptic puncta were
detected as spots using Imaris 3D visualization software (Bitplane), and individual spots and
synaptic locations were quantified through a custom nearest neighbor analysis in MATLAB. The
code is available on GitHub: https://github.com/KummerLab/SEQUIN.

To quantify synapse density, the frequency distribution of nearest-neighbor pre- and post-
synaptic protein pairs was generated for the top 20% intensity of pre- and post-synaptic puncta.
The frequency distribution of nearest neighbor pairs, which forms a Gaussian distribution and
excludes all random non-synaptic pairings, was used to calculate the synaptic locus density. The
mean value calculated at the peak of the Gaussian distribution provided the average pre-post
synaptic separation. The number of pre- and posts-synaptic puncta, including those randomly
allocated and not associated with synaptic loci, irrespective of whether they were within synaptic
loci, was quantified from the spot analysis using Imaris 3D visualization software.

GeoMx statistics and reproducibility.
Quality Control. In the mouse GeoMx experiment, 124 segments were collected in total from 11
mice, and 21 of these segments failed Quality Control (QC) analysis and were removed from the
dataset. One segment was removed for sequence stitching < 80%, and aligned sequence reads <
75%. Five additional segments were removed for aligned sequence reads < 75% alone, and four
segments were removed for sequence saturation < 50%. An additional 11 segments were
removed for < 3% of genes being detected above the Limit of Quantification (LOQ). Of the 20175
gene targets contained in the GeoMx Mouse Whole Transcriptome Atlas, 9035 were included in
the downstream analyses following QC analysis. 1 gene target was removed as a global outlier
(Grubbs test, P<0.01), and 7 were removed as local outliers (Grubbs test, P<0.01). 211 genes were
detected below the limit of quantification and were removed. Of the remaining gene targets,
9,031 gene targets were detected above the LOQ in at least 10% of segments, and therefore were
selected for further analysis. An additional 4 gene targets were identified as interesting *a priori*
and were retained in the study for additional analysis. Detailed QC parameters are described in
our   mouse   analysis   https://github.com/Goralsth/Spatial-transcriptomics-reveals-molecular-
[dysfunction-associated-with-cortical-Lewy-pathology (DOI: 10.5281/zenodo.10732492)](https://github.com/Goralsth/Spatial-transcriptomics-reveals-molecular-dysfunction-associated-with-cortical-Lewy-pathology) Sample
size was determined based on pilot data. No statistical method was used to determine sample
size. Experiments did not involve multiple experimental conditions.
In the human GeoMx experiment, 238 segments were collected in total from 8 human
PD/PDD/DLB cingulate cortex samples (n=6 male, n=2 female). The Human GeoMx data was
assessed on the same parameters as the mouse tissue for Quality Control (QC). In total 63
segments failed QC and were removed from the dataset. 33 samples were removed for low
sequence saturation, and 6 segments were removed for low area. An additional 24 segments
were removed for < 5% of genes being detected above the Limit of Quantification (LOQ). Of the
18,815 gene targets present in the GeoMx Human Whole Transcriptome Atlas, 8,601 gene targets
were included in the downstream analyses following QC. 9 gene targets were removed as local
outliers (Grubbs test, P<0.01). 138 genes were detected below the limit of detection and were
removed. Of the remaining gene targets, 8,595 gene targets were detected in at least 10% of
segments. 6 gene targets were identified as interesting *a priori* and were retained in the study
for additional analysis. Detailed QC parameters are described in our human analysis pipeline
https://github.com/Goralsth/Spatial-transcriptomics-reveals-molecular-dysfunction-associated-
[with-cortical-Lewy-pathology (DOI: 10.5281/zenodo.10732492)](https://github.com/Goralsth/Spatial-transcriptomics-reveals-molecular-dysfunction-associated-with-cortical-Lewy-pathology). Sample size was determined
based on pilot data. No statistical method was used to determine sample size. Experiments did
not involve multiple experimental conditions.

Normalization. To account for systematic variation between AOI's, we normalized each count by
the 75th percentile (Q3) of expression for each AOI. To ensure normalization and Quality Control
(QC) procedures were accurate, we compared each AOI's Q3 value to its limit of quantification.
Normalization and QC procedures were determined to be adequate for robust downstream
analysis.

Differential Expression. Differential expression of gene targets in the GeoMx experiments was
determined via a mixed effects linear regression model. The regression model implemented was
as follows: (1) *y*= gene ~ comparison of interest * variable to control + (1/individual). This model
robustly tests for differential gene expression for comparisons of interest (ex. phospho-α-
synuclein negative vs. phospho-α-synuclein positive brain region differences, etc.) while
controlling for confounding variables and multiple sampling of the same samples.

Gene Set Enrichment Analysis. Gene Set Enrichment Analysis was implemented using the GSVA
(Hanzelmann et al., 2013) (RRID:SCR_021058) package in R to determine disrupted pathways
between neurons positive and negative for phospho-α-synuclein immunoreactivity. The
minimum pathway size was set to 5 genes, and the maximum pathway size was set to 500 genes.
We utilized Kegg Brite pathways to determine which transcriptional pathways are differentially
expressed.

Cell-type deconvolution. We used nanoString's cell-type deconvolution algorithm, SpatialDecon
(Danaher et al., 2022), to determine the cell-type proportions in our spatial transcriptomics data.
We determined cell-type proportions in our data via comparison of our data to single-cell atlas
data from the Allen Brain Institute via *SpatialDecon*'s log normal regression. To determine
statistical significance of differences in cell type proportions between inclusion bearing
(phospho-α-synuclein immunopositive) and non-inclusion bearing (phospho-α-synuclein
immunonegative) segments, we utilized the following equation:
(2) *y*= cell-type proportion ~ inclusion status group + (1/individual)
403
404 **GeoMx data availability.** The raw sequencing files and associated metadata for both mouse and
405 human experiments generated in this study have been deposited in the Sequence Read Archive
406 (SRA): https://www.ncbi.nlm.nih.gov/sra/PRJNA1082339
407 The pre-quality control and supporting data for all experiments are available at the following
408 Zenodo repository: https://zenodo.org/records/10729767 [95] (DOI: 10.5281/zenodo.10729767)
409
410 **GeoMX code availability.** Code used to generate the analyze gene expression and cell detection
411 is available on GitHub: https://github.com/Goralsth/Spatial-transcriptomics-reveals-molecular-
412 dysfunction-associated-with-cortical-Lewy-pathology [96] (DOI: 10.5281/zenodo.10732492).
413
SynGO enrichment analysis. The expert-curated SynGO gene ontology database (Koopmans et
al., 2019) was used to characterize the GeoMX-derived synaptic transmission pathways enriched
in differentially expressed genes from phospho-α-synuclein bearing neurons available from
Goralski et al. studies. The list of DEGs was loaded against the 1602 annotated genes (SynGO
release 1.2 '20231201'). For downregulated genes (enriched in phospho-α-synuclein
immunonegative neurons), 418 genes were mapped (397 genes had a cellular component

annotation, and 299 had a biological process annotation). For upregulated genes, 51 were mapped where 46 had a cellular component annotation, and 34 had a biological processes annotation. Additionally, 40 and 18 cellular component terms, and 82 and 15 biological processes were significantly enriched at 1% FDR for downregulated and upregulated genes respectively.

References

Aarsland, D., Batzu, L., Halliday, G.M., Geurtsen, G.J., Ballard, C., Ray Chaudhuri, K., and Weintraub, D. (2021). Parkinson disease-associated cognitive impairment. Nat Rev Dis Primers 7, 47.

Andersen, K.B., Hansen, A.K., Damholdt, M.F., Horsager, J., Skjaerbaek, C., Gottrup, H., Klit, H., Schacht, A.C., Danielsen, E.H., Brooks, D.J., and Borghammer, P. (2021). Reduced Synaptic Density in Patients with Lewy Body Dementia: An [(11) C]UCB-J PET Imaging Study. Mov Disord 36, 2057-2065.

Blumenstock, S., Rodrigues, E.F., Peters, F., Blazquez-Llorca, L., Schmidt, F., Giese, A., and Herms, J. (2017). Seeding and transgenic overexpression of alpha-synuclein triggers dendritic spine pathology in the neocortex. EMBO Mol Med 9, 716-731.

Blumenstock, S., Sun, F., Klaus, C., Marinkovic, P., Sgobio, C., Paeger, L., Liebscher, S., and Herms, J. (2021). Cortical circuit dysfunction in a mouse model of alpha-synucleinopathy in vivo. Brain Commun 3, fcab273.

Bousset, L., Pieri, L., Ruiz-Arlandis, G., Gath, J., Jensen, P.H., Habenstein, B., Madiona, K., Olieric, V., Bockmann, A., Meier, B.H., and Melki, R. (2013). Structural and functional characterization of two alpha-synuclein strains. Nat Commun 4, 2575.

Braak, H., Del Tredici, K., Rub, U., de Vos, R.A., Jansen Steur, E.N., and Braak, E. (2003). Staging of brain pathology related to sporadic Parkinson's disease. Neurobiol Aging 24, 197-211.

Braak, H., Rub, U., and Del Tredici, K. (2006). Cognitive decline correlates with neuropathological stage in Parkinson's disease. J Neurol Sci 248, 255-258.

Bruck, A., Aalto, S., Nurmi, E., Bergman, J., and Rinne, J.O. (2005). Cortical 6-[18F]fluoro-L-dopa uptake and frontal cognitive functions in early Parkinson's disease. Neurobiol Aging 26, 891-898.

Buter, T.C., van den Hout, A., Matthews, F.E., Larsen, J.P., Brayne, C., and Aarsland, D. (2008). Dementia and survival in Parkinson disease: a 12-year population study. Neurology 70, 1017-1022.

Cai, M., Dang, G., Su, X., Zhu, L., Shi, X., Che, S., Lan, X., Luo, X., and Guo, Y. (2021). Identifying Mild Cognitive Impairment in Parkinson's Disease With Electroencephalogram Functional Connectivity. Front Aging Neurosci 13, 701499.

Caviness, J.N., Lue, L.F., Beach, T.G., Hentz, J.G., Adler, C.H., Sue, L., Sadeghi, R., Driver-Dunckley, E., Evidente, V.G., Sabbagh, M.N., *et al.* (2011). Parkinson's disease, cortical dysfunction, and alpha-synuclein. Mov Disord 26, 1436-1442.

Chen, C., Kim, J.J., Thompson, R.F., and Tonegawa, S. (1996). Hippocampal lesions impair contextual fear conditioning in two strains of mice. Behav Neurosci 110, 1177-1180.

Creed, R.B., Memon, A.A., Komaragiri, S.P., Barodia, S.K., and Goldberg, M.S. (2022). Analysis of hemisphere-dependent effects of unilateral intrastriatal injection of alpha-synuclein pre-formed fibrils on mitochondrial protein levels, dynamics, and function. Acta Neuropathol Commun 10, 78.

Danaher, P., Kim, Y., Nelson, B., Griswold, M., Yang, Z., Piazza, E., and Beechem, J.M. (2022). Advances in mixed cell deconvolution enable quantification of cell types in spatial transcriptomic data. Nat Commun 13, 385.

Earls, R.H., Menees, K.B., Chung, J., Barber, J., Gutekunst, C.A., Hazim, M.G., and Lee, J.K. (2019). Intrastriatal injection of preformed alpha-synuclein fibrils alters central and peripheral immune cell profiles in non-transgenic mice. J Neuroinflammation 16, 250.

Frigerio, I., Bouwman, M.M.A., Noordermeer, R., Podobnik, E., Popovic, M., Timmermans, E., Rozemuller, A.J.M., van de Berg, W.D.J., and Jonkman, L.E. (2024). Regional differences in synaptic degeneration are linked to alpha-synuclein burden and axonal damage in Parkinson's disease and dementia with Lewy bodies. Acta Neuropathol Commun 12, 4.

Froula, J.M., Henderson, B.W., Gonzalez, J.C., Vaden, J.H., McLean, J.W., Wu, Y., Banumurthy, G., Overstreet-Wadiche, L., Herskowitz, J.H., and Volpicelli-Daley, L.A. (2018). alpha-Synuclein fibril-induced paradoxical structural and functional defects in hippocampal neurons. Acta Neuropathol Commun 6, 35.

Frye, H.E., Izumi, Y., Harris, A.N., Williams, S.B., Trousdale, C.R., Sun, M.Y., Sauerbeck, A.D., Kummer, T.T., Mennerick, S., Zorumski, C.F., *et al.* (2021). Sex Differences in the Role of CNIH3 on Spatial Memory and Synaptic Plasticity. Biol Psychiatry 90, 766-780.

Gcwensa, N.Z., Russell, D.L., Long, K.Y., Brzozowski, C.F., Liu, X., Gamble, K.L., Cowell, R.M., and Volpicelli-Daley, L.A. (2024). Cortico-amygdala synaptic structural abnormalities produced by templated aggregation of alpha-synuclein. bioRxiv.

Goralski, T.M., Meyerdirk, L., Breton, L., Brasseur, L., Kurgat, K., DeWeerd, D., Turner, L., Becker, K., Adams, M., Newhouse, D.J., and Henderson, M.X. (2024). Spatial transcriptomics reveals molecular dysfunction associated with cortical Lewy pathology. Nature Communications 15.

Hänzelmann, S., Castelo, R., and Guinney, J. (2013). GSVA: gene set variation analysis for microarray and RNA-seq data. BMC Bioinformatics 14, 7.

Hely, M.A., Reid, W.G., Adena, M.A., Halliday, G.M., and Morris, J.G. (2008). The Sydney multicenter study of Parkinson's disease: the inevitability of dementia at 20 years. Mov Disord 23, 837-844.

Henderson, M.X., Cornblath, E.J., Darwich, A., Zhang, B., Brown, H., Gathagan, R.J., Sandler, R.M., Bassett, D.S., Trojanowski, J.Q., and Lee, V.M.Y. (2019). Spread of alpha-synuclein pathology through the brain connectome is modulated by selective vulnerability and predicted by network analysis. Nat Neurosci 22, 1248-1257.

Holmes, S.E., Honhar, P., Tinaz, S., Naganawa, M., Hilmer, A.T., Gallezot, J.D., Dias, M., Yang, Y., Toyonaga, T., Esterlis, I., *et al.* (2024). Synaptic loss and its association with symptom severity in Parkinson's disease. NPJ Parkinsons Dis 10, 42.

Iba, M., McDevitt, R.A., Kim, C., Roy, R., Sarantopoulou, D., Tommer, E., Siegars, B., Sallin, M., Kwon, S., Sen, J.M., *et al.* (2022). Aging exacerbates the brain inflammatory micro-environment contributing to alpha-synuclein pathology and functional deficits in a mouse model of DLB/PD. Mol Neurodegener 17, 60.

Irwin, D.J., Grossman, M., Weintraub, D., Hurtig, H.I., Duda, J.E., Xie, S.X., Lee, E.B., Van Deerlin,
V.M., Lopez, O.L., Kofler, J.K*., et al.* (2017). Neuropathological and genetic correlates of survival
and dementia onset in synucleinopathies: a retrospective analysis. Lancet Neurol 16, 55-65.

Irwin, D.J., White, M.T., Toledo, J.B., Xie, S.X., Robinson, J.L., Van Deerlin, V., Lee, V.M., Leverenz,
J.B., Montine, T.J., Duda, J.E*., et al.* (2012). Neuropathologic substrates of Parkinson disease
dementia. Ann Neurol 72, 587-598.

Koopmans, F., van Nierop, P., Andres-Alonso, M., Byrnes, A., Cijsouw, T., Coba, M.P., Cornelisse,
L.N., Farrell, R.J., Goldschmidt, H.L., Howrigan, D.P*., et al.* (2019). SynGO: an evidence-based,
expert-curated knowledge base for the synapse. Neuron 103, 217-234 e214.

Kramer, M.L., and Schulz-Schaeffer, W.J. (2007). Presynaptic alpha-synuclein aggregates, not
Lewy bodies, cause neurodegeneration in dementia with Lewy bodies. J Neurosci 27, 1405-1410.

Li, J., Zhang, Y., Huang, Z., Jiang, Y., Ren, Z., Liu, D., Zhang, J., La Piana, R., and Chen, Y. (2022).
Cortical and subcortical morphological alterations in motor subtypes of Parkinson's disease. NPJ
Parkinsons Dis 8, 167.

Lu, J., Wang, C., Cheng, X., Wang, R., Yan, X., He, P., Chen, H., and Yu, Z. (2022). A breakdown in
microglial metabolic reprogramming causes internalization dysfunction of alpha-synuclein in a
mouse model of Parkinson's disease. J Neuroinflammation 19, 113.

Luk, K.C., Kehm, V., Carroll, J., Zhang, B., O'Brien, P., Trojanowski, J.Q., and Lee, V.M. (2012).
Pathological alpha-synuclein transmission initiates Parkinson-like neurodegeneration in
nontransgenic mice. Science 338, 949-953.

Matuskey, D., Tinaz, S., Wilcox, K.C., Naganawa, M., Toyonaga, T., Dias, M., Henry, S., Pittman, B.,
Ropchan, J., Nabulsi, N*., et al.* (2020). Synaptic Changes in Parkinson Disease Assessed with in
vivo Imaging. Ann Neurol 87, 329-338.

Pérez-Acuña, D., Shin, S.J., Rhee, K.H., Kim, S.J., and Lee, S.J. (2023). alpha-Synuclein propagation
leads to synaptic abnormalities in the cortex through microglial synapse phagocytosis. Mol Brain
16, 72.

Poewe, W., Seppi, K., Tanner, C.M., Halliday, G.M., Brundin, P., Volkmann, J., Schrag, A.E., and
Lang, A.E. (2017). Parkinson disease. Nat Rev Dis Primers 3, 17013.

Recasens, A., Dehay, B., Bove, J., Carballo-Carbajal, I., Dovero, S., Perez-Villalba, A., Fernagut,
P.O., Blesa, J., Parent, A., Perier, C*., et al.* (2014). Lewy body extracts from Parkinson disease
brains trigger alpha-synuclein pathology and neurodegeneration in mice and monkeys. Ann
Neurol 75, 351-362.

Reitz, S.J., Sauerbeck, A.D., and Kummer, T.T. (2021). SEQUIN: An imaging and analysis platform
for quantification and characterization of synaptic structures in mouse. STAR Protoc 2, 100268.

Ryman, S.G., Yutsis, M., Tian, L., Henderson, V.W., Montine, T.J., Salmon, D.P., Galasko, D., and
Poston, K.L. (2021). Cognition at Each Stage of Lewy Body Disease with Co-occurring Alzheimer's
Disease Pathology. J Alzheimers Dis 80, 1243-1256.

Sauerbeck, A.D., Gangolli, M., Reitz, S.J., Salyards, M.H., Kim, S.H., Hemingway, C., Gratuze, M.,
Makkapati, T., Kerschensteiner, M., Holtzman, D.M.*, et al.* (2020). SEQUIN multiscale imaging of
mammalian central synapses reveals loss of synaptic connectivity resulting from diffuse
traumatic brain injury. Neuron 107, 257-273 e255.

Schapira, A.H.V., Chaudhuri, K.R., and Jenner, P. (2017). Non-motor features of Parkinson disease.
Nat Rev Neurosci 18, 435-450.

Schirinzi, T., Madeo, G., Martella, G., Maltese, M., Picconi, B., Calabresi, P., and Pisani, A. (2016).
Early synaptic dysfunction in Parkinson's disease: Insights from animal models. Mov Disord 31,
802-813.

Stoyka, L.E., Arrant, A.E., Thrasher, D.R., Russell, D.L., Freire, J., Mahoney, C.L., Narayanan, A.,
Dib, A.G., Standaert, D.G., and Volpicelli-Daley, L.A. (2020). Behavioral defects associated with
amygdala and cortical dysfunction in mice with seeded alpha-synuclein inclusions. Neurobiol Dis
134, 104708.

Thomsen, M.B., Jacobsen, J., Lillethorup, T.P., Schacht, A.C., Simonsen, M., Romero-Ramos, M.,
Brooks, D.J., and Landau, A.M. (2021). In vivo imaging of synaptic SV2A protein density in healthy
and striatal-lesioned rats with [11C]UCB-J PET. J Cereb Blood Flow Metab 41, 819-830.

Vargas, K.J., Schrod, N., Davis, T., Fernandez-Busnadiego, R., Taguchi, Y.V., Laugks, U., Lucic, V.,
and Chandra, S.S. (2017). Synucleins Have Multiple Effects on Presynaptic Architecture. Cell Rep
18, 161-173.

Volpicelli-Daley, L.A., Luk, K.C., and Lee, V.M. (2014). Addition of exogenous alpha-synuclein
preformed fibrils to primary neuronal cultures to seed recruitment of endogenous alpha-
synuclein to Lewy body and Lewy neurite-like aggregates. Nat Protoc 9, 2135-2146.

Volpicelli-Daley, L.A., Luk, K.C., Patel, T.P., Tanik, S.A., Riddle, D.M., Stieber, A., Meaney, D.F.,
Trojanowski, J.Q., and Lee, V.M. (2011). Exogenous alpha-synuclein fibrils induce Lewy body
pathology leading to synaptic dysfunction and neuron death. Neuron 72, 57-71.

Wu, Q., Takano, H., Riddle, D.M., Trojanowski, J.Q., Coulter, D.A., and Lee, V.M. (2019). alpha-
Synuclein (alphaSyn) Preformed Fibrils Induce Endogenous alphaSyn Aggregation, Compromise
Synaptic Activity and Enhance Synapse Loss in Cultured Excitatory Hippocampal Neurons. J
Neurosci 39, 5080-5094.

Zhan, S.S., Beyreuther, K., and Schmitt, H.P. (1993). Quantitative assessment of the
synaptophysin immuno-reactivity of the cortical neuropil in various neurodegenerative disorders
with dementia. Dementia 4, 66-74.

Figure Legends

Figure 1. Progressive loss of cortical excitatory synapses in the PFF mouse model stratingat 3 months post-PFF injection.
(A) Model of striatal PFF injection to assess the impact of cortical pathology.
(B-D) Left, 3D reconstructed images of excitatory presynaptic VGluT1 (green) and excitatory postsynaptic Homer (red) in the cortex after (B) 1 month, (C) 3 months, and (D) 6 months of monomer vs. PFF injection. Right, differences in paired synaptic loci density are denoted by the p-value (Mann-Whitney test) for PFF vs. monomer injection. VGluT1 and Homer-positive puncta densities were analyzed based on puncta intensities and are shown as mean ± SEM. B,C, n = 3 animals/group, 6-8 slices per animal. D, n = 6 animals/group, 6-8 slices per animal.

Figure 2. Reduction of excitatory synaptic specializations.
Images as in Figure 1 were analyzed for the density of individual excitatory pre- and post-synaptic specializations, including those scored as paired and isolated. No reduction was observed after 1 MPI (A), but excitatory pre- and post-synaptic specializations were less abundant at 3 MPI (B). No change was detected at 6 MPI.

Figure 3. Inhibitory postsynaptic Gephyrin is reduced in the cortex at 6 months post-injection.
(A, B) vGAT- and gephyrin-positive puncta densities were analyzed based on puncta intensities after (A) 3 months and (B) 6 months post-injection. Densities for vGAT, gephyrin, and synaptic loci are shown as mean ± SEM. (A) n = 3 animals/group, 7-9 slices per animal). (B) n = 4 animals/group, 7-9 slices per animal.

Figure 4. Phospho-α-synuclein accumulates at synaptic loci and is associated with synapse loss.
(A) Representative images of 3D imaging fields for PSD-95, synapsin, and phospho-α-synuclein.
(B) High magnification 2D projection 600nm thick showing overlap of pre- and post-synaptic proteins with phospho-α-synuclein.
(C) Schematic of methodology for measuring phospho-α-synuclein in 200 nm fields centered on synaptic loci.
(D) Quantification of synapse-associated and parenchymal phospho-α-synuclein from synuclein monomer and synuclein PFF treated animals across all tiles and regions of low and high relative pathology. Dark symbols are individual image tiles and red symbols are medians for each animal.
(E) Ratio of synaptic associated versus parenchymal phospho-α-synuclein staining in areas with low, mid, and high total phospho-α-synuclein pathology. Dark symbols are individual image tiles and red symbols are the median for each animal.
(F-H) Representative image of area of tilescan performed 1-month post monomer or PFF injection (PFF-injected animal depicted). Maximum intensity projection of mesoscan from monomer (G) and PFF (H) injected animals. Images correspond to boxed area in (F).
(I-J) Heatmaps of synapse-associated phospho-α-synuclein and synaptic density 1-month after injection of synuclein PFFs.

(K-L) Heatmaps of synapse-associated phospho-α-synuclein and synaptic density 1-month after
injection of synuclein monomers. Boxed region in (J) used for analysis in (M-Q). Group-wise
analysis of synaptic density (M) and parenchymal phospho-α-synuclein (N) between monomer
and PFF injected animals. Statistics performed on red symbols representative of group averages
and not individual tiles.
(O-Q) Within animal correlations of synaptic density with synapse-associated phospho-α-
synuclein (subjects 1-3 left to right). Spearman's ρ -0.50, -0.65, 0.81 subject 1, 2, and 3,
respectively. Scale bars: (A):5 μm, (B):425 nm, (F): 1000 μm, (G,H): 250 μm, (I-L): 250 μm.
624

Figure 5. Lewy neurite pathology influences the density and phospho-α-synuclein content of
local synapses.
(A) Representative image of pSD-95, synapsin, and phospho-α-synuclein 1 month following
synuclein PFF injection.
(B) Masking of Lewy neurites for measurement of distance between synaptic loci and neurite
borders.
(C) Schematic corresponding to boxed region in (B) showing distance measurement from
synaptic centroids and borders of Lewy neurite masks.
(D) Quantification of synaptic phospho-α-synuclein enrichment as a function of distance to a
Lewy neurite.
(E) The extent of pathology in Lewy neurites (phospho-α-synuclein intensity) is anticorrelated
with the density of synaptic loci in its vicinity (within 800 nm of a neurite edge) (Spearman's ρ -
0.92).

Figure 6. Dysregulation of synaptic pathways in phospho-α-synuclein bearing neurons.
(A) Volcano plot of SynGO annotated genes differentially expressed in phospho-α-synuclein
immunopositive neurons.
(B, D) Enrichment of synaptic pathways in phospho-α-synuclein pathology bearing neurons.
Sunburst plots illustrate cellular components and biological processes from downregulated and
upregulated DEGs.
(C) Bar plots illustrates top enriched GO terms within categories: Synaptic Organization, Process
in the presynapse and Process in the postsynapse
646

Figure 7. Apparent increase in pre-post synaptic separation with ultrastructural changes in the
excitatory synapse as pathology progresses.
(A) Apparent increase in pre-post synaptic separation for VGluT1/Homer-positive synaptic loci
was analyzed by determining the mean distance from Gaussian distribution of synaptic loci after
PFF vs. monomer injection at 3 months post-injection (n = 3 animals/group, 6-8 slices per animal,
*p, Mann-Whitney test compared to monomer-injected mice).
(B) Ultrastructural analysis of synaptic vesicle size in PFF-injected mice compared to control mice
injected with PBS using transmission electron microscopy.
655

A

PFF

birth · 3 months · 1 MPI · 3 MPI · 6 MPI

B *No excitatory synapse density change 1 MPI*

monomer control · 5 μm

PFF · 5 μm

n.s.

synaptic loci / μm³

monomer · PFF

C *Excitatory synapse loss 3 MPI*

monomer control · 5 μm

PFF · 5 μm

p = 0.025

synaptic loci / μm³

monomer · PFF

D *Excitatory synapse loss 6 MPI*

monomer control · 5 μm

PFF · 5 μm

p = 0.007

synaptic loci / μm³

monomer · PFF

presynaptic vGluT1/postsynaptic Homer/pSyn

658   **Figure 1**

A *No change excitatory pre- and post-synaptic sites 1 MPI*

B *Loss excitatory pre- and post-synaptic sites synapse loss 3 MPI*

C *No change excitatory pre- and post-synaptic sites 6 MPI*

660   **Figure 2**

A *No inhibtory synaptic site change 3 MPI*

monomer control

PFF

B *Inhibitory postsynaptic site loss 6 MPI*

monomer control

PFF

presynaptic VGAT/postsynaptic Gephyrin/pSyn

Figure 3

664 **Figure 4**

667   **Figure 5**

669 **Figure 6**

A *No pathology synaptic spacing 1 MPI*

PFF

0.5 µm

vGluT1/Homer/pSyn — pre-/post-synaptic separation [µm]

monomer
PFF

B *Synaptic separation 3 MPI*

PFF

vGluT1/Homer/pSyn — pre-/post-synaptic separation [µm]

p < 0.0001

pre-/post-synaptic separation (µm)

monomer   PFF

C *Ultrastructural pathology of excitatory synapses*

PBS   PFF

D

p < 0.0001

synaptic vesicle, area (nm²)

PBS   PFF

671   **Figure 7**