STHD: probabilistic cell typing of single Spots in whole Transcriptome spatial

data with High Definition

Chuhanwen Sun[1]*, Yi Zhang[1-4]*#

1. Department of Neurosurgery, Duke University

2. Department of Biostatistics and Bioinformatics, Duke University

3. Brain Tumor Omics Program, The Preston Robert Tisch Brain Tumor Center, Duke University

4. Duke Cancer Institute

*These authors contributed equally.

#Corresponding author: Yi Zhang, Ph.D. yi.zhang@duke.edu

Abstract

Recent spatial transcriptomics (ST) technologies have enabled single- and sub-cellular

resolution profiling of gene expression across the whole transcriptome. However, the transition

to high-definition ST significantly increased data sparsity and dimensionality, posing

computational challenges in identifying cell types, deciphering neighborhood structure, and

detecting differential expression - all are crucial steps to study normal and disease ST samples.

Here we present STHD, a novel machine learning method for probabilistic cell typing of single

spots in whole-transcriptome, high-resolution ST data. Unlike the current binning-aggregation-

deconvolution strategy, STHD directly models gene expression at single-spot level to infer cell

type identities without cell segmentation or spot aggregation. STHD addresses sparsity by

modeling count statistics, incorporating neighbor similarities, and leveraging reference single-

cell RNA-seq data. We show in VisiumHD data that STHD accurately predicts cell type identities

at single-spot level, which achieves precise segmentation of both global tissue architecture and

local multicellular neighborhoods. The high-resolution labels facilitate various downstream

analyses, including cell type-stratified bin aggregation, spatial compositional comparisons, and

cell type-specific differential expression analyses. Moreover, STHD labels further reveal

frontlines of inter-cell type interactions at immune hubs in cancer samples. STHD is scalable

and generalizable across diverse samples, tissues, and diseases, facilitating genome-wide

analyses in various spatial organization contexts. Overall, computational modeling of individual

spots with STHD facilitates discoveries in cellular interactions and molecular mechanisms in

whole-genome spatial technologies with high resolution. STHD is available at

https://github.com/yi-zhang/STHD.

Keywords

Machine learning, spatial transcriptomics, high-resolution, tumor microenvironment, cell-cell

interactions

Background

Spatial transcriptomics (ST) technologies have enabled gene expression profiling in the native

spatial context of tissues. These technologies provide a systematic approach to profile cell type

distribution, regional variations, and cellular neighborhoods across normal and disease

samples. Current ST technologies are based on next-generation sequencing or in situ

hybridization imaging, in short, seq-ST and image-ST[1]. Often, seq-ST approaches cover

towards whole-transcriptome with compromised resolution or depth (e.g. 10X Visium, Slide-

seq[2], Stereo-seq[3]), whereas image-based approaches can provide sub-cellular resolution

measurement of hundreds of genes in a designed panel (e.g. 10X Xenium, STARmap[4],

MERFISH[5]). The technical tradeoff between resolution and throughput determined that seq-

ST is generally suitable for hypothesis generation while image-ST for mechanism validation[6].

Emerging ST technologies have rapidly advanced with image-ST increasing gene coverage and

seq-ST enhancing resolution to sub-cellular scale. Notable technologies that scale to whole-

transcriptome and provide single- or sub-cellular resolution include HDST(7) (2um spot), Seq-

Scope(8) (0.5-0.8um resolution), Stereo-seq(3) (spot size about 0.22um and center-to-center

distance 0.5 or 0.715um), and Open-ST(3) (center-to-center distance 0.6um), *etc*. Recently,

VisiumHD from 10X Genomics is further available for formalin-fixed paraffin-embedded (FFPE)

samples to profile whole transcriptome in spots of size 2x2um, a sub-cellular size for most cell

types and tissue contexts(9). Despite these technological advancements, the analysis of this

data type presents significant computational challenges. The read coverage in each high-

resolution spot is often highly sparse, and it is usually required to bin adjacent spots at a chosen

size. For example, aggregation of 4x4 spots into 8x8um bins is usually recommended in

VisiumHD sample processing. Similarly, typical Stereo-seq analyses aggregate data by binning

50 spots(3,10). The binning process computationally merges spots for sufficient read depth in

downstream analytical tasks(11). However, this approach creates bins spanning multiple single

cells, potentially mixing distinct cell types and thus compromising resolution. To address this,

computational deconvolution of cell proportions is further needed, similar to the scenario in low-

resolution seq-ST using tools like RCTD(12), CARD(13), Cell2Location(14), *etc*. Furthermore,

the vast number of million spots or bins in high-resolution ST data poses additional challenges

in computational efficiency.

We here developed STHD (probabilistic cell typing of single <u>S</u>pots in whole <u>T</u>ranscriptome

spatial data with <u>H</u>igh <u>D</u>efinition), a machine learning method designed for cell typing of

individual spots in high-resolution, whole-transcriptome spatial data. STHD reverses the

traditional approach by first inferring cell type identities directly at the raw spot level, without

requiring cell segmentation or bin aggregation. This is followed by cell type-specific bin

aggregation as needed for downstream genome-wide analyses. To tackle high dimensionality

and data sparsity, the STHD model employs a neighbor-augmented loss. It leverages cell type-

specific gene expression from reference single-cell RNA-seq data, constructs a statistical model

on spot gene counts, and employs regularization from neighbor similarity. The output includes

cell type labels and probabilities for each individual spot. The STHD labels further facilitate

multiple downstream analyses, including cell type-specific compositional comparison, cell type-

specific differential expression, global tissue architecture segmentation, local cellular niche

identification, cell type-augmented binning, and cell-cell interactions analyses. We demonstrated

application of STHD to a colon cancer VisiumHD sample to uncover spatially heterogenous

cancer cell expression programs and identifying immune infiltrate hubs with diverse immune

functions and interactions. Moreover, we demonstrated that STHD is scalable and generalizable

across different tissues, providing both local subcellular classification and global tissue

segmentation.

Results

STHD efficiently models high-resolution spots using a machine learning model with

neighbor regularization.

An overview of STHD model is illustrated in **Fig.1a**. STHD infers latent cell type identities of

each spot using a loss function that simultaneously optimizes two components: likelihood of

spot gene counts and similarity with neighbor spots (**Fig.1a**, **Methods, Supplementary File 1**).

The first part optimizes log likelihood of spot gene counts following Poisson distribution with the

latent cell type variable, where parameters are based on normalized gene expression derived

from a reference scRNA-seq dataset with cell type annotation from a matched disease or

system. The second part denoises sparsity using cross entropy loss based on neighborhood

similarity in cell type probabilities. The contribution from neighbors is controlled by the neighbor

parameter β, which can be tuned with automatic stopping criterion to achieve balance between

statistical likelihood and neighborhood similarity. For each high-resolution spot, STHD outputs

cell type probabilities and labels based on Maximum a Posterior (MAP). The spot cell types

enable multiple downstream analytical tasks, including cell type-stratified bin aggregation, cell

type-specific compositional and differential gene expression analyses, and cell-cell interaction

analyses. STHD implements fast optimization enabled by gradient derivation, numba(15), and

Adam(16) optimizer (full derivation of loss gradient in **Supplementary File 1**). STHD also

includes local low-count detection to filter cell and tissue gaps (**Methods, Supplementary**

Fig.1). To be compatible with the millions of spots in one sample, we built the STHD pipeline

that parallelizes patch-level inference followed by whole-sample integration (**Supplementary**

Fig.2). The pipeline is thus adaptive to various computing resources without requiring GPU for

optimization. Finally, to enable close examination of cellular neighborhoods, we developed

STHDviewer for interactive and scalable visualization of spots in the entire sample

(**Supplementary Fig.2**).

Different from current standards that start from binning, we propose in STHD that the analyses

of whole-transcriptome, high-resolution ST data can begin with modeling the sub-cellular HD

spots. This strategy offers advantages compared to alternative approaches. First, direct

unsupervised clustering of spot-level gene expression results in false grouping of spots due to

extreme sparsity (**Fig.1b**, left). Second, the current bin-aggregation approach can increase

depth, but it mixes cell types and compromise resolution (**Fig.1b**, middle). Lastly, segmenting

cell areas based on the hematoxylin and eosin (H&E) histological images may provide precise

nucleus boundaries, but whole-cell segmentation has remained a challenging task(17) (**Fig.1b**,

right).

STHD infers cell type identities at the spot level, outperforms other inference tools, and

guides cell type-specific binning.

We first demonstrated STHD using the publicly available human VisiumHD sample from 10X

Genomics, where a 6.5x6.5mm area of human colorectal tumor was profiled by 8,726,600 HD

spots covering 18,085 genes. We constructed the normalized gene expression reference from

an independent human colon cancer atlas study that profiled 43,113 genes from 370,115 single

cells annotated into 98 fine-grained cell types from GSE178341(18) (**Supplementary Fig.3**).

The reference gene profile construction step of STHD selected 4618 cell type-informative genes

(**Methods, Supplementary Fig.3**). **Fig.2a** illustrated STHD label for a tumor-containing patch in

the colorectal cancer sample containing 22,336 spots in a 300x300um area or 1100x1100 pixels

in the full-resolution H&E image. First, we observed that STHD-predicted cell type labels

collectively captured the compartmentalization of colon crypt-like structure with tumor-like

epithelial cell type signatures (**Fig.2a**). In detail, stem-like proliferating cells were positioned

inside, circled by the abundant enterocyte cells with scattered tumor stem and transit amplifying

(TA)-like cells(18,19). The intercrypt regions were composed of abundant endothelial and

fibroblast cells, with diverse immune cell populations such as plasma cells, monocytes,

macrophages, dendritic cells, and T cells (**Fig.2a**). Interestingly, the cell type probabilities

displayed high confidence for abundant epithelial spots as well as for rarer cells like

macrophages and plasma IgG B cells (**Fig.2b**, **Supplementary Fig.4**). By default, spots with

maximum posterior probability below 0.8 were assigned as ambiguous. The ambiguous spots

mainly located at boundaries of different cell types and tissue structures and could represent

cell type mixtures or cell gaps (**Fig.2a, b**). In comparison, firstly, clustering of spot-level gene

expression vaguely reflected colon crypt structure with mingled cell identities (**Fig.2c,**

Supplementary Fig.5a). This contradicts the expectation of consistent cell type labels within

each cell, given that the raw spots (2x2um) are much smaller than a single cell (about 10um in

diameter). Secondly, clustering of gene expression of bins aggregated from 4x4 spots (8x8um

bins) within the patch resulted in less noise in crypt center and intercrypt regions (**Fig.2d** left**,**

Supplementary Fig.5b). Moreover, similar bin grouping from the whole-sample clustering result

could clearly distinguish the stemness center, enterocytes, and intercrypt regions, but the

boundaries are assigned as one separated cluster that mixed in intercrypt immune cells (**Fig.2d**

right). Further deconvolution on the aggregated bins using RCTD separated epithelial structure

and immune cells but demonstrated a loss of spot-level resolution (**Fig.2e**).

Since high-resolution ST data lacks ground truth of spot labels, we evaluated the performance

from aspects of histopathology, comparison to other methods, simulation, and marker gene

expression. First, the spatial distribution of STHD cell type labels aligns with cell organization

morphology based on H&E staining (**Fig.2f**). Second, we compared to cell type deconvolution

on 8x8um bins by RCTD doublet mode or full mode using the same reference human colon

cancer scRNA-seq reference (**Methods**). RCTD also resolved cell type proportions that match

the colon crypt organization and the distribution of STHD labels for enterocytes and Stem/TA-

like proliferating cells (**Fig.2e**, **Supplementary Fig.6a, b**). At the raw 2x2um spot level, we next

tested computational methods that were able to label cell types by reference-based

deconvolution like RCTD(12) and CARD(13), Cell2Location(14), Celloscope(20),

Stereoscope(21), or by neighborhood-augmented clustering like Banksy(22). We noted that the

common gene count filters in these tools need to be removed to avoid filtering of most spots; for

example, 20436 of the 22336 spots are removed by RCTD default UMI cutoff 100

(**Supplementary Fig.6c**). At single-spot level, RCTD, CARD, Cell2Location, and Stereoscope

can vaguely reveal the crypt structure while with highly inter-mingled cell types from different

lineages (**Fig.2g**, **Supplementary Fig.7a-e**). The unsupervised Banksy efficiently separated

epithelial crypt and intercrypt fibroblast/endothelial structures, but mixed the rarer immune cells

inside the fibroblast/endothelial cluster and called the boundary as a separate cluster

(**Supplementary Fig.7f**). Moreover, STHD has highest computational efficiency among all

methods, which enables efficient inference on the entire sample (**Supplementary Fig.8**).

Since binned outputs are commonly used in VisiumHD, we noted that the STHD labels at the

subcellular 2x2um resolution can further guide cell type-specific binning in cell-size resolution,

by aggregating adjacent spots of the same cell type, thereby improving gene coverage while

preventing cell type mixing. With spot-level cell types, an appropriate bin size can be chosen,

allowing separation of distinct cell types within the same bin areas for cell type-specific gene

expression profiles. **Fig.2h** showcased STHD-guided binning with two commonly used bin

sizes, 8x8um (single-cell size) and 16x16um; each cell type-specific bin was positioned at

centroid of the covered spots of corresponding cell type. Thus, tumor epithelial cells and

intercrypt endothelial, fibroblasts and immune cells were clearly separated on the STHD-guided

binning spatial maps (**Fig.2h**, **Supplementary Fig.5c, d**).

STHD achieves high performance in benchmark simulations and preserves cell type-

specific expression profiles.

To quantitatively assess performance of STHD, we simulated comprehensive whole-

transcriptome, high-resolution ST data *in silico* (**Methods, Supplementary Fig.9**). As shown in

Fig.2i, we simulated 500 cells with heterogeneous cell sizes and cell types on 22,500 spots,

each with a true cell type label and gene counts following expression levels from the same

human colon cancer reference. STHD achieved average area under the curve (AUC) 0.9977

with overall accuracy 93.97% for this multi-class classification task with 98 categories (**Fig.2i,**

Supplementary Fig.9). STHD outperformed other popular spot-level cell type annotation

approaches when tested in repeated simulations, achieving highest accuracy, AUC, and

percentage of retained spots (**Methods, Fig.2j**). Next, we systematically investigated cell type-

specific expression of marker genes based on STHD prediction across the entire colon cancer

sample. Despite spot sparsity (**Supplementary Fig.5e, Supplementary Fig.3d**), the STHD

labels displayed similar cell type-specific expression patterns of marker genes as in scRNA-seq

reference data in normal epithelial cells (**Fig.2k**, left two panels, **Supplementary Fig.10a**) and

all other cell lineages: tumor epithelial cells (**Supplementary Fig.10b**), myeloid cells

(**Supplementary Fig.11a**), B and plasma cells (**Supplementary Fig.11b**), CD8T, CD4T, and

other T cells (**Supplementary Fig.12, Supplementary Fig.13a**), fibroblasts (**Supplementary**

Fig.13b), and endothelial cells and pericytes (**Supplementary Fig.14**). STHD-guided binning

effectively increased marker gene coverage and meanwhile retained cell-type specificity, shown

in two example bin sizes: 8x8um bins (**Fig.2k**, third panel, **Supplementary Figs.10-14**) and

16x16um bins (**Fig.2k**, fourth panel, **Supplementary Figs.10-14**). We also observed that the

traditional binning approach that aggregated 8x8um bins yielded 311,396 bins with single cell

type and 169,188 bins with more than one cell type, creating one third of 8x8um grid bins with

mixed cell type signals.

STHD enables scalable spatial inference and interactive exploration of cellular niches

and immune hubs, facilitating genome-wide and cell type-specific spatial comparisons.

The STHD pipeline enabled scalable inference across the entire ST sample through parallelized

inference of spatial patches followed by whole-sample integration (**Supplementary Fig.2c**). The

parameter tuning was thus performed for the colon cancer VisiumHD sample to balance

between statistical likelihood and neighborhood similarity (**Supplementary Fig.15**). To facilitate

close examination of spot identities and scalable investigation of cellular neighborhoods, we

developed STHDviewer, an interactive and scalable visualization functionality capable of

rendering millions of spots in a webpage (**Fig.3a, Supplementary Fig.16**). As demonstrated in

Fig.3a, STHD prediction automatically achieved precise segmentation of global tissue

architecture and local multicellular neighborhoods. Specifically, fibroblasts separated regions of

normal-like area with colon gland morphology (named Epi-Normal-like, **Supplementary Fig.16-**

17) and tumor epithelial areas at the margins (Epi-Tumor-left, Epi-Tumor-top, Epi-Tumor-right)

compared to tumor in deposit (Epi-Tumor-inside)(23,24) (**Fig.3b**). Across tumor regions of

interest (ROI), STHD labels delineated compositional heterogeneity of cell lineages and types

(**Fig.3c**). Notably, spatial heterogeneity of immune cell populations was also observed; for

instance, the Epi-Tumor-inside region contained macrophages(25,26) but few T cells compared

to tumor regions at the margin, indicating distinct "hot" verses "cold" T cell infiltration spatial

patterns within the same sample (**Fig.3c** top, **Supplementary Fig.18**).

Besides cell type compositional comparisons, STHD facilitates genome-wide and cell type-

specific differential expression (DE) analyses across spatial regions. Comparing different tumor

ROI, epithelial cells in Epi-Tumor-inside expressed highest stemness gene *LGR5*(27), whereas

epithelial cells in Epi-Tumor-top highly express *CCL20*(28) and enrich pathways like TNF-alpha

signaling (**Fig.3d,e, Supplementary Fig.19-20**). Since the Epi-Tumor-inside region enrich

macrophages, we compared differentially expressed genes specific to macrophage spots and

observed that macrophage infiltrating inside tumor expressed higher extracellular matrix gene

SPP1 and lower metalloelastase *MMP12*, cytokine *CXCL8*, and antigen presentation gene *B2M*,

indicating that macrophage states segregated spatially in the same tumor(26,29,30)(**Fig.3f**,

Supplementary Fig.21). The cell type-specific DE analyses can also be performed on the level

of STHD-guided bins with higher depth, which resulted in a higher number of significant DE

genes with well correlated effects (**Fig.3f**, **Supplementary Fig.20-21**).

We next demonstrated STHD in analyzing the less abundant immune cells after using

STHDviewer to identify immune cell hubs at different locations in the colon cancer VisiumHD

sample. An advantage of STHD is that rarer cell types and fine-grained cellular neighborhoods

are maintained on top of the denoising effect for those abundant cell types forming continuous

tissue architecture; an example of immune-endothelial cell interactions was demonstrated in

Supplementary Fig.22. **Fig.4a** highlighted two T cell-rich areas marked in **Fig.3b**. The

Immune-Trich-left region contained CD4T and CD8T cells with abundant macrophages and

C1Q-high dendritic cells (DC)(31), while the Immune-Trich-right area contained CD4T, CD8T

cells, mature DCs enriched in immunoregulatory molecules (mregDC)(32), and Plasma IgG B

cells (**Fig.4b**). In neighborhood enrichment analyses, each area displayed unique co-occurrence

patterns among immune cell types; for example, T cells in Immune-Trich-left area frequently

interacted with macrophage and DCs, while T cells in Immune-Trich-right area co-occurred with

DCs (**Fig.4a,c**). Utilizing cell type-specific DE analysis, we observed that T cells in the Immune-

Trich-left region expresses higher cytotoxicity genes like GZMB comparing to T cells in the

Immune-Trich-right region, which can be associated with the immunosuppressive function of

mregDCs in tumor(33–35) (**Supplementary Fig.23**).

STHD identifies precise cell-cell interactions frontlines in immune hubs and reveals

direct ligand-receptor communication activities.

Cell-cell communications between cell types in disease microenvironment could drive disease

progression like tumorigenesis(36,37). Molecular interactions between cells types in single-cell

transcriptomics analyses are traditionally inferred by evaluating averaged expression of

communicating gene pairs, which lacks support on direct and physical interactions(38). With the

sub-cellular resolution spots labeled, STHD can pinpoint the precise frontlines of inter-cell type

interactions. As shown in **Fig.4d**, we highlighted the frontlines of T cell-macrophage interaction

and T cell-DC cell interaction in each of the T cell-rich regions. We hypothesized that cell-cell

communications mediated by ligand-receptor signaling near-frontline could be stronger than that

in far-frontline spots. For a given cell type pair, we identified spots at boundary of two cell types

based on posterior probabilities and considered STHD-labeled spots within 4 spots (comparable

to single cell size) in distance as near-frontline spots (**Methods, Fig.4d**). We calculated ligand-

receptor interaction scores (LRIS)[21,22] between the near-frontline spots and compared with

background interaction scores from far-frontline spots, conditioning on the same cell type pair.

As shown in **Fig.4e**, near-frontline spots (grouped as 3,4) harbored higher LRIS activities

compared to the far-frontline spots (grouped as 1,2) in both cell type pairs of T-Macrophage

(**Fig.4d,e** top) and T-DC interactions (**Fig.4d,e** bottom). Specifically, the T-Macrophage

communications in Immune-Trich-left were largely driven by *CD74-IFNG* interaction, while the

T-DC communications in Immune-Trich-right were driven by *CCL17*/*CCL19-CCR7* interactions

(**Fig.3k**), which indicated the immunoregulatory effect of mature DCs on T cells(39).

Multi-sample analyses of human colon cancer identified sample-specific immune

enrichment patterns.

As data accumulate for whole-genome spatial transcriptomics with high-definition, we next

extended STHD to additional VisiumHD samples. Oliveira et al. recently profiled multiple

colorectal cancer samples and adjacent normal colon tissues using VisiumHD(9). We applied

STHD to analyze two cancer samples (P1, P5) and one adjacent normal colon sample (P3), in

addition to the previously released sample (P2) included in this study. **Fig.5a** illustrated STHD-

inferred cell type identities for colon cancer sample P1 containing 8,115,201 spots at 2x2um

spot resolution. The colon cancer sample P1 exhibits a similar tumor organization to the colon

cancer sample P2, with segmented tumor regions containing colon cancer epithelial cells

interspersed with fibroblasts, and adjacent colon tissue enriched with goblet cells. Notably, the

colon cancer sample P1 features a prominent tertiary lymphoid structure (TLS), characterized

by abundant germinal center B cells and surrounding T cells (**Fig.5b** top). The TLS-like regions

was also validated by B cell markers (*CD19*, *MS4A1*) and the TLS-associated T cell cytokine

CXCL13 (**Fig.5b** bottom)(40). In contrast, other colon cancer samples P2 and P5 lack TLS-like

structures (**Fig.2, Fig.5c** left). STHD also provided detailed insights into the normal colon tissue

from sample P3, revealing goblet cells, an adjacent lymphoid structure, and a myofibroblast

region (**Fig.5c** right). Overall, STHD captures heterogenous cellular compositions across

multiple samples highlighting distinct immune cell profiles (**Fig.5d**).

STHD is generalizable to diverse samples across tissues, diseases, and organisms.

We then extended the application of STHD to normal tissues from both human and mouse

model systems. Using the STHD pipeline with default parameters, we analyzed three VisiumHD

samples from 10X Genomics including human pancreas, mouse small intestine, and mouse

brain (**Methods**). For the adult human pancreas sample, we constructed cell type-specific gene

expression estimation using reference single-cell RNA-seq data from Baron et al.(41), which

included 12,000 pancreatic cells from 4 adult donors profiled by inDrop and annotated 14 cell

types. STHD inference, based on this reference, demonstrated the pancreas's abundant acinar

cells and embedded pancreatic islets - organized cell structure harboring endocrine cells

(**Supplementary Fig.24a**, left). We observed that marker gene expression levels matched

expected patterns for various cell types, such as beta cells (*IAPP*, *INS*), alpha cells (*GCG*,

TTR), epsilon cells (*GHRL*), and pancreatic stellate cells (*RGS5*) (**Supplementary Fig.24a**,

right)(41). For the mouse small intestine, we utilized a reference scRNA-seq profile from the

Mouse Cell Atlas 2.0(42) (**Methods**). STHD inference of spot identities collectively reveals the

tissue structure of the small intestine (**Fig.5e**). Specifically, we highlighted the mesenteric lymph

node with immune cells and the layered intestine-epithelium organizing enterocytes, epithelial

cells, goblet cells, Paneth cells, and fibroblasts (**Fig.5f,g** top row), with matched expression of

cell type-specific gene markers(42) observed at 8x8um bin-aggregated level (**Fig.5f,g** bottom

row). Finally, we mapped the mouse brain VisiumHD data to scRNA-seq data based on Allen

Brain Atlas(43) (**Methods**). STHD revealed brain layer-specific cell types for each high-

resolution spot and differentiated known cortical layers and hippocampus regions

(**Supplementary Fig.24b**). Overall, we observe a high degree of spatial correspondence to

tissue structures collectively by STHD spot-level labels, demonstrating STHD's generalizability

across diverse tissues and disease samples.

Discussion

As spatial transcriptomics technologies advance towards whole-genome coverage at single-

and sub-cellular resolution, the development of computational methods that offer both local

precision and global scalability is crucial for enabling comprehensive genome-wide analyses

across spatial regions, samples, and conditions. However, the enhanced dimension and

resolution introduces challenges such as data sparsity. To address this issue, existing analytical

approaches often rely on hard aggregation of adjacent spots, which compromises resolution

and leads to the mixing of distinct cell types. We presented STHD, a machine learning method

designed to model cell types at the single-spot level for high-definition, genome-wide spatial

transcriptomics data. Unlike traditional methods, STHD achieves soft aggregation of cell type-

specific gene expression information contributed by neighboring spots, effectively addressing

the computational difficulties posed by sparse data through a neighbor-augmented Expectation

Maximization (EM) model. Our benchmarking simulations and analyses of recently available

public VisiumHD samples demonstrate STHD's accuracy, precision, and efficiency. With

genome-wide gene expression profiles, STHD's spot-level cell type labels enabled a range of

downstream analyses, including the study of global tissue architecture, local cellular

neighborhoods, cell type-specific differential expression, cell type-augmented binning, and cell-

cell interactions. The STHD pipeline and viewer are implemented to provid efficient modeling

and interactive examination, scalable to the entire sample containing millions of spots.

Moreover, we have shown STHD's generalizability across different samples, tissues, and

disease contexts.

A tradeoff has existed in ST technologies between resolution and throughput. For seq-ST

technologies, which typically cover a high number of genes, the limitation in spot resolution

often results in the mixing of gene expression signals from different cell types within a single

spot (e.g. Visium's 55um spot diameter). Previous computational efforts have aimed to estimate

cell type proportions using spot-level gene expression decomposition, such as RCTD(12),

Cell2Location(14), CARD(13), *etc*. As seq-ST technology progresses towards subcellular spot

size with genome-wide coverage, we adopted a different assumption: most spots could be

attributed to a single cell type. This assumption reduces model complexity and allows for

estimation based on sparse readouts. Notably, STHD's efficient optimization enabled analyses

of an entire VisiumHD sample with millions of spots using around ten CPU hours without

requirement GPU resource. In both normal and tumor tissue samples, we observed that this

assumption held for most spots, except for a subset classified as ambiguous. STHD detected

these ambiguous spots based on the multinomial cell type probability vector, where lack of a

dominant cell type (default posterior cutoff 0.8) indicates ambiguity. Comparison to histology

reveal that these ambiguous spots often correspond to regions like cell gaps, certain non-

nucleus cytoplasm areas, and mostly inter-cell type boundaries. Consequently, these cell type

boundary spots can be utilized in identifying inter-cell type transitions at the highest possible

resolution; also, STHD's estimated cell type probabilities could implicate spot-level

compositions.

The data sparsity in gene counts presents a significant challenge in subcellular spot modeling.

For instance, in the colon cancer sample P2, the average count depth of a 10X VisiumHD spot

was 1/262 of that of a single cell from 10X scRNA-seq. Consequently, this sparsity often results

in the exclusion of most spots in existing spot-level methods or generation of noisy cell type

labels when each HD spot was treated independently. To tackle sparsity, STHD incorporates a

soft regularization of the loss function based on cross-entropy among neighboring spots' cell

type probability vectors. This approach leverages the neighbor similarities in spatial data, as

seen in previous methods like Banksy, which utilized Gabor filter to incorporate structured

neighborhood contribution in unsupervised tasks like tissue segmentation and spatial

clustering(22). Our results show that STHD labels not only captures the continuous zonation of

those abundant cell types but also identifies rarer cell types, such as immune cells in tissue and

tumor samples. The recovered continuity in the spot identities further enabled cell type-guided

binning, which maintains cell type-specific marker expression and enhances gene coverage for

downstream differential expression analyses. In a recent study, Bin2Cell also avoids traditional

binning methods but operates on the raw VisiumHD spots by pooling spots overlapping the

same nucleus identified from H&E image(44). In another recent study, TopACT also models

neighbor aggregation, where the range of multiscale aggregation was optimized for cell type

classification(45). Similar to the neighbor parameters in Banksy and multiscale design in

TopACT, STHD allows for tunning of neighborhood aggregation effect: the β parameter and the

iteration step parameter control the loss contribution from neighbors, and the neighbor size

parameter can control the range of spatial similarity aggregation, making it adaptable to various

cell sizes and tissue types.

Despite the advancements, several challenges remain in analyzing high-dimensional data with

subcellular resolution. First, the lack of group truth for spot-level cell identities in tissue ST

samples complicates the evaluation of method performance. Our spatial simulations, which

created spots with true cell type labels matching scRNA atlases, provide a starting point for

such evaluations, but future integration with other spatial technologies (e.g., image-ST or

proteomics) could offer additional ground truth data. The STHD model is also generalizable to

other high-resolution, high-dimensional spatial data platforms like Stereo-seq and Xenium 5K,

etc. Second, the availability and quality of the reference scRNA-seq data can limit the

applicability of STHD, as it relies on these references for cell type identification but does not

discover new cell types, a common feature also existed for spot deconvolution methods. In

cases where detailed and cell type annotation are lacking or cell type profiles are highly

correlated, STHD label can provide an initial labeling, STHD can still provide initial labeling,

allowing for further discovery of transcriptional variations within each cell type through clustering

of STHD-guided bins. We anticipate expanding of STHD's capabilities for semi-supervised cell

type identification. Third, the issue of molecular diffusion has been found to exist across ST

methods(46), which may interfere with STHD prediction particularly in spots with weak marker

gene signals. Computational tools to correct spot diffusion, such as SpotClean(47) for low-

resolution data, have been developed, but there remains a need for methods compatible with

millions of sparse spots. Additionally, molecular diffusion could also blur the tissue boundaries,

as observed in some regions outside H&E images containing gene counts. This also imply that

integrating histopathology properties can be informative in modeling the high-resolution spatial

transcriptomics sample, as done in recent studies like iStar, which infers spatial gene

expression from H&E images(48).

Looking ahead, we believe that the combination of individual subcellular spot modeling,

neighborhood information sharing, and multi-modality neighbor information incorporation,

including H&E images, will be key to overcoming the resolution-throughput tradeoff in emerging

spatial transcriptomics technologies. This rationale fully harnesses the potential of recent

advancements in spatial technology and enables the discovery of spatially organized cellular

heterogeneity in samples. The algorithm design of STHD is thus generalizable in other platforms

of high-dimensional spatial transcriptomics profiling with subcellular resolution. In summary,

STHD offers probabilistic subcellular spot-level predictions that facilitate precise and scalable

analyses, with a software toolkit that is generalizable across various tissues and diseases,

providing new insights into cellular and molecular landscapes.

Conclusions

We have presented STHD, a novel machine learning method for probabilistic cell typing of

single spots in whole-transcriptome, high-resolution ST data. Unlike the current binning-

aggregation-deconvolution strategy, STHD directly models gene expression at single-spot level

to infer cell type identities without cell segmentation or spot aggregation. STHD addresses

sparsity by modeling count statistics, incorporating neighbor similarities, and leveraging

reference single-cell RNA-seq data. We show that STHD accurately predicts cell type identities

at single-spot level in simulated benchmarking and biological spatial samples, where STHD

achieves precise segmentation of both global tissue architecture and local multicellular

neighborhoods. The high-resolution STHD labels facilitate various downstream analyses,

including cell type-stratified bin aggregation, spatial compositional comparisons, and cell type-

specific differential expression analyses. Moreover, STHD labels further reveal frontlines of

inter-cell type interactions at immune hubs in cancer samples. Finally, STHD is efficient,

scalable, and generalizable across diverse samples, tissues, and diseases. Overall, STHD will

facilitate biomarker and mechanistic discoveries in spatial heterogeneity, cellular interactions,

and molecular mechanisms in various tissue and disease contexts.

Methods

STHD algorithm.

The STHD model is a machine learning method that combine modeling of per-spot gene counts

based on latent cell type identities with regularizing similarity among neighboring spots in terms

of cell type probability distribution. The overall loss function $\mathcal{L}$ composes of negative log

likelihood $-LL$ that models gene count of all spots and the cross entropy loss CE measuring

similarity of neighboring spots. The amount of contribution from neighbors is controlled by the

parameter β. The cell type identities at every single spot are thus estimated by Expectation

Maximization.

$$\mathcal{L} = -LL + \beta CE$$

For the negative log likelihood part, we model counts of each gene at each spot to follow

Poisson distribution. Let the normalized average gene expression level for gene g within the

latent cell type t is λ_g^t, read depth of spot a is d_a, and spot a's probability of belonging to cell

type t is $P_a(t)$. Thus, the count of gene g at spot a: $n_g^a \sim Poission(d_a \lambda_g^t)$.

$$LL = log \prod_a \prod_g P(n_g^a)$$

465
$$= \sum_a \sum_g logP(n_g^a)$$

466
$$= \sum_a \sum_g log \, E_{P_a(t)}P(n_g^a|t)$$

467
$$\geq \sum_a \sum_g E_{P_a(t)} \, logP(n_g^a|t)$$

468
$$= \sum_a \sum_t P_a(t) \sum_g \left(n_g^a log(d_a\lambda_g^t) - d_a\lambda_g^t\right) + const$$

469 Considering high sparsity of single spots, the second part of the loss function is related to cross

470 entropy loss between spot a and nearest neighboring spots $a' \in NN(a)$ in terms of distribution

471 of cell type probability.

472
$$CE = -\sum_a \sum_{a' \in NN(a)} \sum_t \left(P_a(t)logP_{a'}(t)\right)$$

473 Note that one constraint is $\sum_t P_a(t) = 1$. In calculation, we use the following for inherent

474 incorporation of constraint: $P_a(t) := \frac{e^{w_a^t}}{\sum_{s \in T} e^{w_a^s}}$, where T is the full set of cell types. We thus

475 optimize $P_a(t)$ or w_a^t to minimize the total loss:

476
$$Loss = -\sum_a \sum_t P_a(t) \sum_g \left(n_g^a log(d_a\lambda_g^t) - d_a\lambda_g^t\right) - \beta \sum_a \sum_{a' \in NN(a)} \sum_t \left(P_a(t)logP_{a'}(t)\right)$$

477 More details of the model and derivation of explicit form of gradient for fast optimization is

478 available separately in **Supplementary File 1**. In short, we derived that the explicit form of

479 gradient of the loss function $\mathcal{L}$ at $w_{\tilde{a}}^{\tilde{t}}$ for spot $\tilde{a}$ and cell type $\tilde{t}$ as:

480
$$\frac{\partial \mathcal{L}}{\partial w_{\tilde{a}}^{\tilde{t}}} = -\frac{\partial LL}{\partial w_{\tilde{a}}^{\tilde{t}}} + \beta \frac{\partial CE}{\partial w_{\tilde{a}}^{\tilde{t}}}$$

481
$$= -P_{\tilde{a}}(\tilde{t})\mathcal{F}(\tilde{a}, \tilde{t}) + \sum_t P_{\tilde{a}}(t)P_{\tilde{a}}(\tilde{t})\mathcal{F}(\tilde{a}, t)$$

482
$$-\beta \left[P_{\tilde{a}}(\tilde{t}) \sum_{a' \in NN(\tilde{a})} \left(logP_{a'}(\tilde{t}) - \sum_t \left(P_{\tilde{a}}(t)logP_{a'}(t)\right) \right) - \sum_{a \neq \tilde{a}} \mathbb{I}_{a' \in NN(a)} \left(P_a(\tilde{t}) - P_{\tilde{a}}(\tilde{t})\right) \right]$$

483 Where

$$\mathcal{F}(a,t) = \sum_g \left(n_g^a log\left(d_a \lambda_g^t \right) - d_a \lambda_g^t \right)$$

$$P_a(t) = \frac{e^{w_a^t}}{\sum_{s \in T} e^{w_a^s}}$$

STHD thus optimizes $w_{\tilde{a}}^{\tilde{t}}$ and the probability of spot $\tilde{a}$ having the identity of cell type $\tilde{t}$: $P_{\tilde{a}}(\tilde{t})$,

using Adam(16). We implemented iterative updates using numba(15) on sparse matrix to

achieve high computational efficiency and scalable optimization.

Construct normalized gene expression profile from reference single-cell atlas dataset.

The average normalized gene expression λ_g^t can be estimated from a pre-defined single-cell

RNA-seq reference dataset. A reference single-cell dataset shall match the tissue or disease

origin of the high resolution, whole-transcriptome spatial transcriptomics data. It shall also

contain cell type annotation that is usually based on multi-sample harmonization and

unsupervised clustering of single cells into various cell type or groups. The granularity of cell

type annotation directly influence the number of predicted cell type and downstream analyses.

For each cell type, average normalized gene expression profiles are estimated with $\lambda_g^t = \frac{n_g^t}{\sum_g n_g^t}$

so that $\sum_g \lambda_g^t = 1$. Specifically for human colon cancer, we downloaded the raw counts file of the

single-cell RNA-seq data from the human colon cancer atlas from Pelka et.al(18) through

GSE178341. The data contains 370115 tumor and adjacent normal single cells from 62 primary

treatment-naïve colorectal cancer patients, with 43113 genes profiled. Quality control of single

cells was performed by filtering cells with minimum 100 counts and 100 genes, filtering genes

with minimum 3 cells, and removing cells with mitochondrial read ratio larger than 27.7%, which

is mean plus one standard deviation of the mitochondrial read ratio across all cells. After the

above quality control, 303358 cells and 31,873 genes remain. The cell annotations were

obtained from the study that provided 98 fine-grained cell types and 7 cell lineages. We used

the 98 high-granularity cell type annotation throughout this study. To select genes informative in

distinguishing the 98 cell types, we followed the calculation of cell type expression profile as in

RCTD(12). First, we calculated the estimated cell type expression profiles λ_g^t for all genes and

cell types. We then selected genes with average expression profile above 0.000125, which is

0.0625 counts per 500, and at least 0.5 log-fold-change compared to the average expression

across all cell types, using log2 scale. Moreover, we removed the mitochondrial genes and

ribosome genes in the selection. The selection resulted in 4618 genes that are cell type-

informative, for each we calculated λ_g^t in all cell types. Note that STHD uses the 4618 genes to

infer cell type identity of spots, and later can consider whole-transcriptome for downstream

analyses.

STHD hyperparameter tuning.

To select hyperparameters that balance between statistical likelihood and neighborhood

similarity, we comprehensively tested hyperparameter tuning and provided an automatic

stopping criterion and optional tuning tutorial. The main parameters in STHD includes the

neighborhood parameter β, which controls the contribution of cross entropy loss among local

neighbors to the total loss, the optimization steps n_iteration, and the learning rate that can be

fixed in practice. We provided a tutorial on parameter tuning based on performance in random

sampled patches; for the human colon cancer sample, the optimized parameters are 23 iteration

steps and β=0.1. In detail, for the colon cancer sample, we randomly selected six different

patches in size of 1100x1100 pixels (in full resolution, around 22500 spots). For the optimization

of each patch, we tested a range of β from 0, 0.03, 0.1, 0.3, 1, and 3, each outputting optimal

iteration steps based on the early stopping criterion, where the drop of loss in the past 10 steps

falls within 1%. Since the regularization term from neighborhood contribution β will lower log

likelihood, as β increases, we select the optimal parameter pairs where drop in cross entropy

slows down to around 10% while a high log likelihood is maintained. The optimal iteration steps

when fixing β of 0, 0.03, 0.1, 0.3, 1, 3, is each 24, 23, 22, 23, 23, 24. The optimal iteration steps

is robustly 23.17 on average. Therefore, we chose β=0.1 and n_iteration=23 as optimal

parameters for the whole sample modeling that balance the count-based likelihood and the

contribution from neighbors.

STHD low-count region detection and filtering.

Throughout the spatial transcriptomics data, regions of low gene counts exist, usually arising

from tissue boundaries, tissue gaps, and regions low in DNA/RNA such as dead cells, necrosis,

vessel enriching red blood cells, and collagen rich area. We included an automatic local low-

count region detection and filtering functionality in STHD. By default, STHD computed neighbor

graphs within 2 rings of spots (13 spots for the grid system), aggregated local connected spots,

and detected low-count local regions not exceeding 50 counts, where the model will skip

modeling due to low confidence in cell type inference. Other customized tissue gap and

boundary masks can also be provided to further filter spots.

Assessing model accuracy with simulated spatial transcriptomics data.

To assess the accuracy of STHD model prediction, we simulated the high-dimensional, high-

resolution spatial transcriptomics data *in silico*. We simulated a region with 22500 (150x150)

grid spots and position 500 simulated cells overlapping the region with cell sizes spanning 1-20

spots. Each cell is randomly assigned a cell type from the 98 cell types annotated in colon

cancer atlas, and spots overlapping each whole cell are assigned the corresponding cell type.

To mimic the real-world scenario in gene expression profiles and spot sparsity, we adapted the

same normalized gene expression profile from the same colon cancer atlas data with 98 cell

types. The depth of each spot was simulated following the distribution of exponential decay with

mean read depth 47 matching the colon cancer VisiumHD sample. After simulating counts of

each gene at each spot in the region, we applied STHD with default optimal parameters (β=0.1)

and also STHD no-neighbor model (β=0) to the simulated spatial data, and measured prediction

accuracy, standard error, and area under the receiver operating characteristic curve (ROC), for

this task of multi-class classification of 98 categories at each spot. To compared with other

methods that enable spot-level cell type annotation, we examined six different settings: STHD,

RCTD with default count filter, RCTD with no count filter, CARD with default count filter, CARD

with no count filter, and Cell2Location. The inclusion of both default parameter and adjusted

count filtering for RCTD and CARD is due to the fact that most spots will be removed if following

default parameters, where accuracy measurements were based on non-filtered spots. We

evaluated the methods in ten simulation replicates and assessed performance metrics in mean

accuracy, AUC, and percentage of retained spots.

Comparison with other spot cell typing methods.

To benchmark the STHD results at 2x2 μm resolution, we compared STHD to six methods that

can output cell typing, including the deconvolution-based cell typing tools RCTD(12), CARD(13),

Cell2Location(14), Celloscope(20), Stereoscope(21), and the unsupervised method Banksy(22).

Because of computational efficiency and data size limit of the existing methods, we used the colon

cancer tumor region as in Fig.2a to perform benchmarking, which contains 22,336 cells and

18,085 genes. For methods requiring the reference scRNA-seq data as input, we used the same

human colorectal cancer scRNA-seq datasets (GSE178341) with 98 cell types. Since many

methods in comparison requires a maximum number of cells from reference scRNAseq dataset

– for example, CARD requires below 50000 cells - we down-sampled the reference data to include

maximum 600 cells for each cell type, resulting in a total of 49,351 cells and 41,195 genes. For

running RCTD on single spots, we used RCTD doublet mode, testing both the default minimum

UMI 100 and also by removing the minimum UMI requirement due to loss of most raw spots. For

testing CARD, we followed guidelines from https://github.com/YMa-lab/CARD. Similarly, to avoid

most spots being filtered, the minimum spot count and minimum gene numbers were set to 0.

Cell2location and Stereoscope were ran with default parameters according to the tutorials

https://cell2location.readthedocs.io/en/latest/notebooks/cell2location_tutorial.html,   and

https://github.com/almaan/stereoscope. In Cell2location, by default, the reference data was

filtered using the following parameters: cell_count_cutoff = 5, cell_percentage_cutoff = 0.03, and

nonz_mean_cutoff = 1.12. The reference single-cell regression model was trained with default

parameter max_epochs = 250, lr = 0.002. The spatial cell2location model was obtained with

default parameters max_epoch = 30,000. In Stereoscope, by default, the reference data was

filtered using a minimum count threshold of min_counts = 10, and all mitochondrial genes were

removed. The reference single-cell model was trained with parameters max_epochs = 100; the

spatial stereoscope model was obtained using parameters max_epochs = 10,000. Finally, for

Celloscope, a binary matrix with prior knowledge about marker genes is required among input,

for which we selected top 200 genes in normalized expression from our informative gene matrix

for each cell type. Another required input is estimated number of cells in each spot, for which we

set to 1 due to sub-cellular-resolution nature of the data. Celloscope was then tested following

guidelines at https://github.com/szczurek-lab/Celloscope. Finally, Banksy was tested with default

parameters   following   the   tutorial   at

https://github.com/prabhakarlab/Banksy_py/blob/main/slideseqv2_analysis.ipynb. For all toolkits,

the weights or cell type abundance for each spot were extracted and normalized as cell

proportions for subsequent visualizations.

Cell type deconvolution in aggregated bins.

Following current standards of binning-deconvolution, we binned the spatial data and aggregated

the bins containing 4x4 spots. We applied RCTD deconvolution of doublet mode on the same

spatial patch as in Fig.2a, following the default filtering and parameters of RCTD as on spacexr

(https://github.com/dmcable/spacexr), and using the same subsampled colon cancer reference

scRNA-seq data as in comparison with other methods. To compare cell proportions across bins,

we counted STHD cell type labels and obtained cell proportions for each 4x4 bin after excluding

filtered and ambiguous spots. We also computed cosine similarity between STHD's spot count-

based proportions and RCTD's deconvoluted proportions and visualize the proportions and

cosine similarity on a spatial plot.

Estimating running time in computational efficiency.

To assess computational efficiencies, different cell typing tools are applied onto the tumor

epithelial patch using a server with 8 CPU cores and maximum 128GB memory, running time

recorded. For tools supporting GPU acceleration (cell2location and stereoscope) the time cost

was evaluated on a single GPU, and the spatial training time was recorded without including

scRNA training time. For computational efficacy, we also estimated the time cost for the entire

sample by linearly scaling running time based on number of spots in patch (22,336 spots) and in

the entire colon cancer sample P2 (8,726,224 spots).

STHD pipeline on an entire VisiumHD sample.

The STHD software and pipeline contains several components to flexibly split the whole sample

into patches for inferences and parallelization. The input/output functionalities (STHDio)

expanded from Squidpy Visium interface to enable cropping of full-resolution image. Second, for

the full-size sample, STHD will automatically patchify the full sample into multiple regions with

overlaps, followed by low-count filtering, per-patch inference, and result aggregation where

probabilities of overlapping spots are averaged. The STHD cell type label results are

determined by Maximum A Posterior (MAP) with a preset cutoff (default 0.8). An optional fast

implementation for STHD-guided binning with a predefined bin size is provided, which outputs

per-gene count for each bin stratified by STHD cell type labels. Finally, the predicted cell labels

are exported to STHDviewer for a rasterized and interactive scatter plot implemented using

Bokeh(49). In STHDviewer, the spot grid of array_row and array_column were used for x-axis

and y-axis for visualization efficiency.

Cell type-specific differential gene expression analysis across spatial regions of interest.

To study regional differences within the cancer sample, we first focused on regions enriching

epithelial cell types and manually separated four different regions and extracted spot barcodes.

Based on STHD predicted labels, we performed differential expression analyses specific for

epithelial cell types. Gene counts were normalized towards the spot count depth (target total

counts 10000) and log-transformed followed by Wilcoxon test in Scanpy. Genes passing

adjusted p-value cutoff 10^{-6} and absolute log2 fold change 0.5 were selected as significant, and

top genes passing log2 fold change 0.5 are exported to pathway enrichment analyses by

GSEApy(50) using EnrichR and a few pathway databases, including GO Molecular Function

2023, GO Biological Process 2023, MSigDB Hallmark 2020, Reactome 2022, WikiPathway

2023 Human, KEGG 2021 Human, and BioPlanet 2019. The FDR cutoff of 0.1 was used to filter

the pathways and top four pathways were visualized for tumor epithelial pathway comparison.

We also tested cell type-specific differential expression using both STHD-labeled spot and

STHD-guided bin aggregates of size 4x4 spots. At STHD-guided bin level, we followed the

same step of normalization and performed differential expression with Wilcoxon with the same

cutoff for epithelial and macrophage comparison. For T cell-specific differential expression

analyses between the two T cell rich regions, the cutoff was set log2 fold change 0.5 and

adjusted p-value cutoff 0.05 due to small number of spots from patches. For differential gene

dotplots, significant genes are sorted by log fold change and the top three genes for the tumor

comparison and top ten genes for the immune comparison were plotted.

Neighborhood interaction enrichment analysis for immune-rich regions in colon cancer

VisiumHD sample.

To investigate the neighborhood interaction patterns across various cell types in the two immune-infiltrated regions, local neighborhoods were calculated using the 4 nearest cells in the grid and considering 6 rings of adjacent spots to get a obtain a local neighborhood structure using squidpy.gr.spatial_neighbors in Squidpy. We used STHD predicted cell type labels and excluded spots in the filtered and ambiguous categories. The neighborhood enrichment Z-scores were calculated using squidpy.gr.nhood_enrichment, which follows CellChat(38) with ligand-receptor databases and permutation of 1000 times. Heatmaps of neighborhood interaction patterns were plotted based on the co-occurrence score of pairwise cell types in each region of interest.

Inter-cell type frontline and cell-cell communication analysis.

For a given pair of cell types, we first defined their precise frontlines to be spots where those two cell types are adjacent to each other. We further restrict frontline spots to be those whose neighbor only contains that pair of cell types based on maximum posterior to eliminate potential noises from other cell types. Considering a single cell is comparable to the size of 4 spots, we assumed spots within 4 spots from the frontlines represented one cell with direct interaction with another cell type. Thus, spots were categorized based on their cell type and proximity to frontline. In the colon cancer sample P2, we examined interactions between T cells and dendritic cells, and T cells and macrophages separately for the two immune regions of interest, Immune-Trich-left and Immune-Trich-right. In the chord plots, for a pair of cell types A and B, the labels were assigned as follows: '1' for far-frontline cell type A, '2' for far-frontline cell type B, '3' for near-frontline cell type A, and '4' for near-frontline cell type B. We then hypothesized that cell-cell communications like ligand-receptor expression levels were higher between group 3 and 4 compared to group 1 and 2. To perform the ligand-receptor interaction analysis, normalized data was applied the squidpy.gr.ligrec function which permuted 1000 times to detect significant ligand-receptor pairs.

Top 10 ligand-receptor pairs different between near-frontline and far-frontline communications

were visualized in the chord diagram showing the mean expression of each ligand-receptor pair

and sender-receiver directions.

STHD on additional colon samples and other human and mouse tissue samples.

We performed STHD analyses on the extended list of human colon samples using the same colon

cancer reference gene estimation, yielding cell type inference on 8,115,201 spots in colon cancer

P1, 8,665,510 spots in colon cancer P5 sample, and 7,791,374 spots in the normal colon sample

P3 using normal cell types. We further extended analyses to the human pancreas sample

containing 6,680,357 spots for 18,085 genes, the mouse intestine sample containing 5,479,447

spots with 19,059 genes, and the mouse brain sample containing 6,296,637 spots with 19,059

genes. All analyses were run using default parameters. To match the tissue system, we used the

non-cancerous colon scRNA for the normal colon sample, normal human adult pancreas scRNA

dataset for the human pancreas sample, the adult Mouse Cell Atlas intestine scRNA dataset for

the mouse small intestine sample, and the Allen Brain Atlas reference for the mouse brain sample.

For constructing reference gene expression profile for human pancreas, we downloaded raw

counts file of single-cell RNA-seq data from the human pancreas atlas by Baron et al.(41) from

GSE84133. The dataset contains 8,569 normal single cells from four donors, profiling 20,124

genes. Quality control involved filtering cells with a minimum of 100 counts and 100 genes,

filtering genes with a minimum of 3 cells, and removing cells with a mitochondrial read ratio

larger than 5%. After quality control, 8,569 cells and 16,358 genes remained. The study

provided 14 cell type annotations. Using STHD default setting, we selected 4,436 cell type-

informative genes based on average expression profile above 0.000125 (0.0625 counts per

500) and at least 0.5 log-fold change, excluding mitochondrial and ribosomal genes. For mouse

intestine tissue, the raw counts file of single-cell RNA-seq data from the mouse cell atlas by Fei

et al.(42) was downloaded from GSE176063. The dataset contains 6,684 normal single cells

from three adult mouse intestine samples, profiling 16,416 genes. Quality control involved

filtering cells with a minimum of 100 counts and 100 genes, filtering genes with a minimum of 3

cells, and removing cells with a mitochondrial read ratio larger than 7.15%. After quality control,

6,102 cells and 15,745 genes remained, with 28 cell types annotated. Similarly, we selected

4,083 cell type-informative genes based on average expression profile above 0.0002 (0.1

counts per 500) and at least 0.5 log-fold change, excluding mitochondrial and ribosomal genes.

For mouse brain reference gene expression construction, the processed single-cell RNA-seq

data from the Allen Brain Atlas was downloaded from the Seurat tutorial website

(https://satijalab.org/seurat/articles/visiumhd_analysis_vignette), which contained 199,993 single

cells and 31,053 genes. Quality control involved filtering cells with a minimum of 100 counts and

100 genes, filtering genes with a minimum of 3 cells, and removing cells with a mitochondrial

read ratio larger than 4.8%. After quality control, 181,250 cells and 264,484 genes remained,

with 41 subclass cell types annotated. Similarly, we selected 5,171 cell type-informative genes

based on average expression profile above 0.000125 (0.0625 counts per 500) and at least 0.5

log-fold change, excluding mitochondrial and ribosomal genes.

Declarations

Availability of data and materials

Software availability.

The current STHD software source code is on GitHub at https://github.com/yi-zhang/STHD.

An interactive STHDviewer of the high-resolution cell type result for the human colon cancer

sample P2 is available at https://yi-zhang-compbio-

lab.github.io/STHDviewer_colon_cancer_hd/STHDviewer_crchd.html.

Data Availability.

All analyses in this paper are based on publicly available data.

Colon cancer reference scRNA-seq atlas: data and cell type annotation from Human Colon

Cancer Atlas is available from Pelka et. al through GSE178341:

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE178341).

Colon cancer VisiumHD data (patient P2) is available at

https://www.10xgenomics.com/datasets/visium-hd-cytassist-gene-expression-libraries-of-

human-crc, named Visium HD Spatial Gene Expression Library, Human Colorectal Cancer

(FFPE), by Space Ranger 3.0.0, 10x Genomics (2024, Mar 25).

Colon cancer VisiumHD data (patient P1, P3, P5) from Oliveira et al.(9) include two colon

cancer and one normal colon, available at https://www.10xgenomics.com/products/visium-hd-

spatial-gene-expression/dataset-human-crc.

The human pancreas VisiumHD data is available at

https://www.10xgenomics.com/datasets/visium-hd-cytassist-gene-expression-libraries-human-

pancreas.

The mouse small intestine VisiumHD data is available at

https://www.10xgenomics.com/datasets/visium-hd-cytassist-gene-expression-libraries-of-

mouse-intestine.

The mouse brain VisiumHD data is available at https://www.10xgenomics.com/datasets/visium-

hd-cytassist-gene-expression-libraries-of-mouse-brain-he.

Competing interests

The authors declare no competing interests.

Funding

C.S. and Y.Z. was supported by start-up funds to Y.Z. from Brain Tumor Omics Program,

Preston Robert Tisch Brain Tumor Center, Department of Neurosurgery and Department of

Biostatistics and Bioinformatics at Duke University School of Medicine.

Authors' contributions

This study was conceived and led by Y.Z. Y.Z. designed the model and algorithm, implemented

the STHD software, led the data analyses, and wrote the manuscript. C.S. analyzed the data

and helped with manuscript writing.

Acknowledgements

We acknowledge 10X Genomics to share VisiumHD datasets publicly available. We

acknowledge authors of the Human Colon Cancer Atlas to make scRNAseq datasets publicly

available.

Figures

Fig.1.

Fig.1. Overview of STHD model. a. STHD (probabilistic Cell Typing of Single Spots in Whole

Transcriptome Spatial Data with High Definition) takes two inputs: whole-transcriptome spatial

data of high resolution, and gene expression profiles by cell type from reference single-cell

dataset. The STHD model is a machine learning method that infers latent cell type identity of

individual spots by simultaneously modeling gene counts and neighbor similarity. STHD outputs

cell type identities for each single spot in formats of cell type labels and posterior probabilities.

The spot-level cell identities can then be used for multiple downstream analyses, including

STHD-guided binning with cell type stratification, spatial comparison of cell type-specific

compositional and differential expression analyses, and cell-cell interaction frontline analyses. **b.**

Alternative approaches to analyze the HD datatype. Left, direct unsupervised clustering of spot-

level gene expression is affected by sparsity. Middle, current binning-deconvolution approach

increases coverage but compromise resolution. Right, ideal cell-level aggregation based on

segmentation.

Fig.2.

Fig.2. Spot-level predictions and benchmarking of a zoomed patch in a colon cancer

VisiumHD sample. a. Example of STHD results of a patch from the human colon cancer

VisiumHD sample, revealing tumor-like epithelial cells and intercryptic immune population,

visualized using the interactive STHDviewer. **b.** Cell type posterior probabilities at spot-level for

the same patch, visualized using Squidpy. Top row, three epithelial cell types; Bottom row, two

immune cell types. **c.** A spatial plot of unsupervised leiden clustering on spot gene expression.

d. Spatial plots of unsupervised leiden clustering of gene expression aggregated by bins in size

of 4x4 spots. Left, clustering bins within patch, right, global cluster group of bins. **e.** Cell type

proportion decomposed by RCTD on bins of size of 4x4 spots, demonstrating two tumor

epithelial classes, enterocytes, and stem-like cells. **f.** The H&E histopathology image in full

resolution of the same crop. **g.** Comparison to other cell typing and clustering methods adaptive

to high-resolution spots. **h.** The STHD predicted labels at spot-level and STHD-guided binning

in bins of 4x4 spots and 8x8 spots, visualized using Squidpy using the same colon cancer cell

type colormap. **i.** Benchmarking with simulated high-resolution spatial data. Left, ground truth

cell type labels; middle, STHD predicted cell type labels; right, receiver operating characteristic

curve (ROC) curve for all cell types. **j.** Benchmarking against other methods for spot annotation

in repeated spatial transcriptomics simulations. **k.** Expression dot plots for marker genes of

normal epithelial cells, at spot level or STHD-guided bin level. Left to right: normal epithelial cell

types in reference human colon cancer sample, normal epithelial cell types based on STHD

predicted spots, normal epithelial cell types from STHD-guided bins of size 4x4 spots, normal

epithelial cell types in STHD-guided bins of size 8x8 spots. The same cell type-specific gene

markers from the original colon cancer study were used. ROC, Receiver operating

characteristic. AUC, area under the curve.

Fig.3.

Fig.3. STHD prediction facilitates segmentation of global structures and cell type-specific

differential analyses across spatial regions. a. Interactive spatial visualization by

STHDviewer of predicted cell type labels for spots in the entire human colon cancer VisiumHD

sample. **b.** Annotation of regions of interest, separating epithelial regions and two T cell-

infiltrated regions. **c.** Cell type proportion bar plots at cell lineage and cell type level for epithelial

regions and whole map. Upper: Proportions of major cell lineages including epithelial cells,

fibroblasts, B cells, myeloid cells, and T cells. Bottom: Proportions of finer cell types of epithelial

lineage. **d.** Spot-level gene expression for epithelial-specific differential genes across regions.

Top three genes by log fold change were shown. **e.** Pathway enrichment of epithelial cell-

specific genes differentially expressed in the Epi-Tumor-top region. **f.** Macrophage-specific

differential genes for the Epi-Tumor-inside region, where differential analyses were performed at

bin level of size 4x4 spots with cell type stratification from STHD. DE, differential expression.

Fig.4.

Fig.4. STHD enabled analyses of local neighborhoods for cellular interactions in tumor-infiltrating immune hubs. a. STHD cell type labels for two T cell-rich regions in the colon cancer sample. **b.** Proportions of immune cell types in the two immune regions. **c.** Cell neighborhood interaction patterns of the two T cell-rich regions showing neighborhood enrichment z-scores where spots within 6 spots are considered neighbors. **d.** T cell and myeloid cells in the two immune-rich regions, with frontlines of inter-cell type interactions. Top, T cell-macrophage frontline in Immune-Trich-left region; Bottom, T cell-dendritic cell frontline in Immune-Trich-right region. Considering a pair of cell types, spots within 4 spots (8um) from the inter-cell type interaction location are highlighted as near-frontline spots, with the rest as far-

frontline spots. **e.** The ligand-receptor communications at near-frontline spots compared to far-

frontline communication scores of the same cell type pair as background, where the ligand-

receptor interaction scores were calculated by squidpy.ligrec. Top 10 ligand-receptor pairs

different between near-frontline and far-frontline communications were visualized where width

represents average expression level of ligand receptor pairs. Spots are classified into four

groups 1,2,3,4 based on their cell type and near- or far-frontline categories.

Fig.5.

Fig.5. STHD is scalable to other samples, revealing subcellular cell type identities and tissue structures. a. STHD spot-level inference of a human colon cancer Visium HD sample showing tumor, colon tissue, and enriched lymphoid structure. **b.** Left, spots (2x2um) labeled as germinal center B cells, with B cell marker expression aggregated per 8x8um bins. Right, spots (2x2um) labeled as T cells, and expression of TLS-specific T cell cytokine *CXCL13* by 8x8um bins. Marker expression plots are illustrated by loupe browser. **c.** Spot-level inference of human colon VisiumHD sample P5 (cancer) and P3 (normal adjacent tissue). **d.** Immune cell-specific composition across all four human colon Visium HD samples. **e.** STHD results for the mouse small intestine VisiumHD sample, with two zoomed-in regions highlighted. **f.** Spots with immune cell identities in the lymph node-like region, showing T cell, *Ms4a1*+ B cell and *Ighd*+ B cells. Top, STHD labels in 2x2um spots, Bottom, marker gene expression in 8x8um bins. **g.** Spots of intestine cells forming layered structure, including intestine enterocyte cell, epithelial cell, goblet cell, Paneth cell, and fibroblasts. Top, STHD labels in 2x2um spots, Bottom, marker gene expression in 8x8um bins. GC, germinal center. CRC, colorectal cancer. NAT, normal adjacent tissue.

References

References

1. Tian L, Chen F, Macosko EZ. The expanding vistas of spatial transcriptomics. Nat Biotechnol.

2023 Jun;41(6):773–82.

2. Stickels RR, Murray E, Kumar P, Li J, Marshall JL, Di Bella DJ, et al. Highly sensitive spatial

transcriptomics at near-cellular resolution with Slide-seqV2. Nat Biotechnol. 2021

Mar;39(3):313–9.

3. Chen A, Liao S, Cheng M, Ma K, Wu L, Lai Y, et al. Spatiotemporal transcriptomic atlas of

mouse organogenesis using DNA nanoball-patterned arrays. Cell. 2022 May

12;185(10):1777-1792.e21.

4. Wang X, Allen WE, Wright MA, Sylwestrak EL, Samusik N, Vesuna S, et al. Three-

dimensional intact-tissue sequencing of single-cell transcriptional states. Science. 2018 Jul

27;361(6400):eaat5691.

5. Moffitt JR, Bambah-Mukku D, Eichhorn SW, Vaughn E, Shekhar K, Perez JD, et al.

Molecular, spatial, and functional single-cell profiling of the hypothalamic preoptic region.

Science. 2018 Nov 16;362(6416):eaau5324.

6. Rao A, Barkley D, França GS, Yanai I. Exploring tissue architecture using spatial

transcriptomics. Nature. 2021 Aug;596(7871):211–20.

7. Vickovic S, Eraslan G, Salmén F, Klughammer J, Stenbeck L, Schapiro D, et al. High-

definition spatial transcriptomics for in situ tissue profiling. Nat Methods. 2019

Oct;16(10):987–90.

8. Cho CS, Xi J, Si Y, Park SR, Hsu JE, Kim M, et al. Microscopic examination of spatial

transcriptome using Seq-Scope. Cell. 2021 Jun 24;184(13):3559-3572.e22.

9. Oliveira MF, Romero JP, Chung M, Williams S, Gottscho AD, Gupta A, et al. Characterization
of immune cell populations in the tumor microenvironment of colorectal cancer using high
definition spatial profiling [Internet]. bioRxiv; 2024 [cited 2024 Jul 28]. p. 2024.06.04.597233.
Available from: https://www.biorxiv.org/content/10.1101/2024.06.04.597233v1

10. Wu L, Yan J, Bai Y, Chen F, Zou X, Xu J, et al. An invasive zone in human liver cancer
identified by Stereo-seq promotes hepatocyte–tumor cell crosstalk, local
immunosuppression and tumor progression. Cell Res. 2023 Aug;33(8):585–603.

11. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al. Dictionary
learning for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol. 2024
Feb;42(2):293–304.

12. Cable DM, Murray E, Zou LS, Goeva A, Macosko EZ, Chen F, et al. Robust decomposition
of cell type mixtures in spatial transcriptomics. Nat Biotechnol. 2022 Apr;40(4):517–26.

13. Ma Y, Zhou X. Spatially informed cell-type deconvolution for spatial transcriptomics. Nat
Biotechnol. 2022 Sep;40(9):1349–59.

14. Kleshchevnikov V, Shmatko A, Dann E, Aivazidis A, King HW, Li T, et al. Cell2location
maps fine-grained cell types in spatial transcriptomics. Nat Biotechnol. 2022 May;40(5):661–
71.

15. Lam SK, Pitrou A, Seibert S. Numba: a LLVM-based Python JIT compiler. In: Proceedings
of the Second Workshop on the LLVM Compiler Infrastructure in HPC [Internet]. New York,
NY, USA: Association for Computing Machinery; 2015 [cited 2024 May 14]. p. 1–6.
(LLVM '15). Available from: https://dl.acm.org/doi/10.1145/2833157.2833162

16. Kingma DP, Ba J. Adam: A Method for Stochastic Optimization [Internet]. arXiv; 2017 [cited

2024 May 12]. Available from: http://arxiv.org/abs/1412.6980

17. Hartman A, Satija R. Comparative analysis of multiplexed in situ gene expression profiling

technologies [Internet]. bioRxiv; 2024 [cited 2024 May 12]. p. 2024.01.11.575135. Available

from: https://www.biorxiv.org/content/10.1101/2024.01.11.575135v2

18. Pelka K, Hofree M, Chen JH, Sarkizova S, Pirl JD, Jorgji V, et al. Spatially organized

multicellular immune hubs in human colorectal cancer. Cell. 2021 Sep 2;184(18):4734-

4752.e20.

19. Merlos-Suárez A, Barriga FM, Jung P, Iglesias M, Céspedes MV, Rossell D, et al. The

intestinal stem cell signature identifies colorectal cancer stem cells and predicts disease

relapse. Cell Stem Cell. 2011 May 6;8(5):511–24.

20. Geras A, Darvish Shafighi S, Domżał K, Filipiuk I, Rączkowska A, Szymczak P, et al.

Celloscope: a probabilistic model for marker-gene-driven cell type deconvolution in spatial

transcriptomics data. Genome Biology. 2023 May 17;24(1):120.

21. Andersson A, Bergenstråhle J, Asp M, Bergenstråhle L, Jurek A, Fernández Navarro J, et

al. Single-cell and spatial transcriptomics enables probabilistic inference of cell type

topography. Commun Biol. 2020 Oct 9;3(1):1–8.

22. Singhal V, Chou N, Lee J, Yue Y, Liu J, Chock WK, et al. BANKSY unifies cell typing and

tissue domain segmentation for scalable spatial omics data analysis. Nat Genet. 2024 Feb

27;1–11.

23. Höppener DJ, Stook JLPL, Galjart B, Nierop PMH, Nagtegaal ID, Vermeulen PB, et al. The

relationship between primary colorectal cancer histology and the histopathological growth

patterns of corresponding liver metastases. BMC Cancer. 2022 Aug 22;22(1):911.

24. Sirinukunwattana K, Pluim JPW, Chen H, Qi X, Heng PA, Guo YB, et al. Gland

segmentation in colon histology images: The glas challenge contest. Med Image Anal. 2017

Jan;35:489–502.

25. Ozato Y, Kojima Y, Kobayashi Y, Hisamatsu Y, Toshima T, Yonemura Y, et al. Spatial and

single-cell transcriptomics decipher the cellular environment containing HLA-G+ cancer cells

and SPP1+ macrophages in colorectal cancer. Cell Rep. 2023 Jan 31;42(1):111929.

26. Zhang Y, Xiang G, Jiang AY, Lynch A, Zeng Z, Wang C, et al. MetaTiME integrates single-

cell gene expression to characterize the meta-components of the tumor immune

microenvironment. Nature Communications. 2023 May 6;14(1):2634.

27. Xu L, Lin W, Wen L, Li G. Lgr5 in cancer biology: functional identification of Lgr5 in cancer

progression and potential opportunities for novel therapy. Stem Cell Res Ther. 2019 Jul

29;10(1):219.

28. Kapur N, Mir H, Clark III CE, Krishnamurti U, Beech DJ, Lillard JW, et al. CCR6 expression

in colon cancer is associated with advanced disease and supports epithelial-to-

mesenchymal transition. Br J Cancer. 2016 Jun;114(12):1343–51.

29. Cheng S, Li Z, Gao R, Xing B, Gao Y, Yang Y, et al. A pan-cancer single-cell transcriptional

atlas of tumor infiltrating myeloid cells. Cell. 2021 Feb 4;184(3):792-809.e23.

30. Matusiak M, Hickey JW, van IJzendoorn DGP, Lu G, Kidzinski L, Zhu S, et al. Spatially

Segregated Macrophage Populations Predict Distinct Outcomes In Colon Cancer. Cancer

Discovery [Internet]. 2024 Mar 28 [cited 2024 Jun 8]; Available from:

https://doi.org/10.1158/2159-8290.CD-23-1300

31. Baruah P, Dumitriu IE, Malik TH, Cook HT, Dyson J, Scott D, et al. C1q enhances IFN-

gamma production by antigen-specific T cells via the CD40 costimulatory pathway on

dendritic cells. Blood. 2009 Apr 9;113(15):3485–93.

32. Maier B, Leader AM, Chen ST, Tung N, Chang C, LeBerichel J, et al. A conserved dendritic-

cell regulatory program limits antitumour immunity. Nature. 2020 Apr;580(7802):257–62.

33. Kvedaraite E, Ginhoux F. Human dendritic cells in cancer. Science Immunology. 2022

Apr;7(70):eabm9409.

34. Puleston DJ, Baixauli F, Sanin DE, Edwards-Hicks J, Villa M, Kabat AM, et al. Polyamine

metabolism is a central determinant of helper T cell lineage fidelity. Cell. 2021 Aug

5;184(16):4186-4202.e20.

35. You S, Li S, Zeng L, Song J, Li Z, Li W, et al. Lymphatic-localized Treg-mregDC crosstalk

limits antigen trafficking and restrains anti-tumor immunity. Cancer Cell [Internet]. 2024 Jul

18 [cited 2024 Jul 31];0(0). Available from: https://www.cell.com/cancer-cell/abstract/S1535-

6108(24)00239-3

36. Wang X, Tokheim C, Gu SS, Wang B, Tang Q, Li Y, et al. In vivo CRISPR screens identify

the E3 ligase Cop1 as a modulator of macrophage infiltration and cancer immunotherapy

target. Cell. 2021 Oct 14;184(21):5357-5374.e22.

37. Burdziak C, Alonso-Curbelo D, Walle T, Reyes J, Barriga FM, Haviv D, et al. Epigenetic

plasticity cooperates with cell-cell interactions to direct pancreatic tumorigenesis. Science.

2023 May 12;380(6645):eadd5327.

38. Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan CH, et al. Inference and
analysis of cell-cell communication using CellChat. Nat Commun. 2021 Feb 17;12(1):1088.

39. Li J, Zhou J, Huang H, Jiang J, Zhang T, Ni C. Mature dendritic cells enriched in
immunoregulatory molecules (mregDCs): A novel population in the tumour
microenvironment and immunotherapy target. Clin Transl Med. 2023 Feb 17;13(2):e1199.

40. Chen Z, Wang X, Jin Z, Li B, Jiang D, Wang Y, et al. Deep learning on tertiary lymphoid
structures in hematoxylin-eosin predicts cancer prognosis and immunotherapy response.
NPJ Precis Oncol. 2024 Mar 22;8(1):73.

41. Baron M, Veres A, Wolock SL, Faust AL, Gaujoux R, Vetere A, et al. A Single-Cell
Transcriptomic Map of the Human and Mouse Pancreas Reveals Inter- and Intra-cell
Population Structure. Cell Syst. 2016 Oct 26;3(4):346-360.e4.

42. Fei L, Chen H, Ma L, E W, Wang R, Fang X, et al. Systematic identification of cell-fate
regulatory programs using a single-cell atlas of mouse development. Nat Genet. 2022
Jul;54(7):1051–61.

43. Yao Z, Velthoven CTJ van, Kunst M, Zhang M, McMillen D, Lee C, et al. A high-resolution
transcriptomic and spatial atlas of cell types in the whole mouse brain [Internet]. bioRxiv;
2023 [cited 2023 Sep 8]. p. 2023.03.06.531121. Available from:
https://www.biorxiv.org/content/10.1101/2023.03.06.531121v1

44. Polański K, Bartolomé-Casado R, Sarropoulos I, Xu C, England N, Jahnsen FL, et al.
Bin2cell reconstructs cells from high resolution Visium HD data [Internet]. bioRxiv; 2024
[cited 2024 Jul 31]. p. 2024.06.19.599766. Available from:
https://www.biorxiv.org/content/10.1101/2024.06.19.599766v1

45. Multiscale topology classifies cells in subcellular spatial transcriptomics | Nature [Internet].

[cited 2024 Aug 12]. Available from: https://www.nature.com/articles/s41586-024-07563-1

46. You Y, Fu Y, Li L, Zhang Z, Jia S, Lu S, et al. Systematic comparison of sequencing-based

spatial transcriptomic methods. Nat Methods. 2024 Jul 4;1–12.

47. Ni Z, Prasad A, Chen S, Halberg RB, Arkin LM, Drolet BA, et al. SpotClean adjusts for spot

swapping in spatial transcriptomics data. Nat Commun. 2022 May 27;13(1):2971.

48. Zhang D, Schroeder A, Yan H, Yang H, Hu J, Lee MYY, et al. Inferring super-resolution

tissue architecture by integrating spatial transcriptomics with histology. Nat Biotechnol. 2024

Jan 2;1–6.

49. Bokeh Development Team. Bokeh: Python library for interactive visualization [Internet].

2018. Available from: https://bokeh.pydata.org/en/latest/

50. Fang Z, Liu X, Peltz G. GSEApy: a comprehensive package for performing gene set

enrichment analysis in Python. Bioinformatics. 2023 Jan 1;39(1):btac757.

1068