Hydathode immunity against the vascular pathogen

Xanthomonas campestris pv. campestris by the

Arabidopsis CNL-type receptor SUT1

Short title:

Hydathode immunity by Arabidopsis CNL-type receptor SUT1

Author list:

Nanne W. Taks[1], Marieke van Hulten[1], Jeroen A. van Splunter-Berg[1], Sayantani

Chatterjee[1], Misha Paauw[1], Sebastian Pfeilmeier[1] and Harrold A. van den Burg[1]

Author affiliations:

1) Molecular Plant Pathology, Swammerdam Institute for Life Sciences, Faculty

of Science, University of Amsterdam, Science Park 904, 1098 XH,

Amsterdam, the Netherlands

*Author for correspondence: H.A. van den Burg

Email: H.A.vandenBurg@uva.nl, Tel +31 20 525 7797

Figures: 4

Supporting figures: 7

Supporting data files: 4

Abstract

Bacterial plant pathogens exploit natural openings, such as pores or wounds, to enter

the plant interior and cause disease. Plants actively guard these openings through

defense mechanisms that have been described extensively for stomates, the most

common points of entry. However, bacteria from the genus *Xanthomonas* have

specialized in that they enter their host via hydathodes—a poorly studied organ at the

leaf margin involved in guttation. While hydathodes can mount an effective immune

response against bacteria, a dedicated perception mechanism still needs to be

discovered. To identify a hydathode-specific immune receptor, we mapped a novel

resistance gene against *X. campestris* pv. campestris (Xcc) in Arabidopsis using an

inoculation procedure that promotes natural entry via hydathodes. Using Recombinant

Inbred Lines (RILs) between susceptible accession Oy-0 and resistant Col-0, a QTL

for resistance was identified on the right arm of Chromosome 5 in Col-0. Combining

this finding with results of a genome-wide association analysis, a single candidate

gene was fine-mapped that encoded a coiled-coil nucleotide-binding leucine-rich

repeat (CNL) immune receptor protein called SUPPRESSOR OF TOPP4 1 (SUT1).

Whereas the ZAR1 immune receptor acts in the vasculature against Xcc, we establish

that SUT1 already restricts Xcc in hydathodes but is ineffective in the vasculature. In

corroboration, we confirm promoter activity of *SUT1* in the epithem tissue within

hydathodes. Altogether, we provide evidence for an NLR that confers hydathode-

specific resistance in Arabidopsis against infection by Xcc.

54 **Author summary**

Black rot disease, caused by the bacterial pathogen *Xanthomonas campestris* pv. campestris (Xcc), is an economically relevant disease of cabbage crops. Xcc is rather unique in that it enters the plant interior through specialized organs at the edge of the leaf. These structures called hydathodes contain water pores and are involved in leaf water regulation. Although we know that hydathodes can mount an immune response against these bacteria, specific immune receptors still need to be discovered. In our search for hydathode resistance mechanisms, we use the model plant *Arabidopsis thaliana* to identify genetic targets that could be translated to cabbage breeding practices. Here, by screening large populations of genetically diverse Arabidopsis plants, we could pinpoint a genetic locus that is involved in hydathode resistance. On this locus, we identified a gene, *SUT1*, that confers resistance against Xcc, restricting early hydathode colonization by the bacteria and reducing subsequent disease symptoms. Interestingly, this resistance is ineffective in later stages of infection when the bacteria colonize the plant vascular system. Therefore, this study provides new insights in hydathode-specific resistance and opens doors for more research on these tissue- or organ-specific resistance mechanisms in plants.

Introduction

Plant pathogenic microbes exploit different strategies to invade their host and induce disease. Foliar bacterial pathogens primarily take advantage of natural openings of

plants, such as pores involved in gas exchange (stomata) or water and nutrient homeostasis (hydathodes). Consequently, plants have evolved a plethora of immune mechanisms to guard these openings, responding to the presence of pathogens [1,2]. Among these entry points, stomata are arguably the most widely studied in the context of plant immunity. Several immune mechanisms have been elucidated over the past years, including sensing of microbe-associated molecular patterns (MAMPs), such as bacterial flagellin [3] or fungal chitin [4]. Upon perception, plants activate a signaling cascade that includes pre-invasive closure of the stomatal pores followed by post-invasive opening [5–7]. While stomata provide access to the leaf apoplast, hydathodes represent an entry point for highly specialized bacterial pathogens that colonize the leaf vasculature [8–10]. The hydathode anatomy is composed of an enclosed extracellular cavity interspersed with specialized epithem cells. The cavity is connected to the exterior via stomata-like openings in the epidermis called water pores [11,12]. Water pores allow water exudation, called guttation, in conditions when root pressure exceeds water evaporation [13]. The cavity is connected to endings of xylem trachea and in part filled with the epithem tissue, which is hypothesized to act as an active filter of the guttation fluid. Interestingly, the water pores do not completely close in response to pathogens [8,14], providing nearly unrestricted access to the hydathode cavity. Several pathogenic *Xanthomonas* spp. are known to invade their hosts via hydathodes [8,9,15]. Despite the absence of pre-invasive immunity, a post-invasive immune response is mounted in hydathodes against both hydathode-adapted and non-adapted bacterial pathogens [16], which restricts bacterial proliferation in the cavity and thereby reduces the number of vascular outbreaks. Intriguingly, although general immune signaling hubs have been confirmed to be involved in this hydathode immune response, the existence of hydathode-specific immune receptors still needs to be demonstrated.

Plant immune receptors can be classified as cell-surface and intracellular receptors [17,18]. While cell-surface pattern recognition receptors (PRRs) generally detect well-conserved microbe-associated molecular patterns (MAMPs), intercellular nucleotide-binding leucine-rich repeat (NLR) receptors respond directly or indirectly to the mode of action of effector proteins. These effector promote virulence upon delivery to the host cell to manipulate cellular processes including the suppression of plant immunity [19]. The plant NLRs can be divided into three subgroups based on the N-terminal

domains: Coiled-coil (CC) NLRs (CNLs), Toll/Interleukin-1 receptor/Resistance (TIR) NLRs (TNLs), and RPW8-like CC domain (RPW8) helper NLRs (RNLs). While some NLRs can activate immune responses by themselves through oligomerization into a so-called resistosome complex [20], others rely on helper NLRs or are part of an NLR network to confer immune signaling [18]. The evolution of plant immune receptor networks and pathogen immune suppressors has shaped the landscape of most plant-pathogen interactions [17].

These immune receptors and pathogen effectors have been studied extensively in *Arabidopsis thaliana* (hereafter Arabidopsis) for several bacterial pathogens, in particular *Pseudomonas syringae* [21], a stomatal pathogen. For hydathode-infecting bacteria, such as *Xanthomonas campestris* pv. campestris (Xcc) [10], only a few resistance mechanisms, for instance NLR-mediated recognition of the Xcc effector XopAC (AvrAC) by HOPZ-ACTIVATED RESISTANCE (ZAR1), have been unraveled [22,23]. However, this latter finding was achieved using inoculation methods that circumvent the natural infection path via hydathodes, thereby overlooking any resistance mechanisms unique to hydathodes. Furthermore, many Arabidopsis accessions, including Col-0, still display a considerable level of resistance against Xcc strains that lack XopAC [16], indicating the presence of additional resistance mechanisms.

Limited research has been conducted on pathogen recognition and defense mechanisms within hydathodes. It is known for example that in cabbage, lignin is deposited in and around the hydathodes in association with Xcc infections [24]. In cauliflower, a necrotic response was detected around hydathodes in response to Xcc [8], indicating some form of a hypersensitive response (HR) that is often associated with plant immunity [25,26]. This response was found to be more pronounced when using an Xcc mutant defective in the type three secretion system (T3SS). The T3SS is responsible for translocating bacterial effector proteins into host cells, suggesting that Xcc type three effectors (T3Es) are involved in immune suppression within hydathodes. Furthermore, a strain in which the *xopAC* gene is deleted induces a stronger necrotic response in cauliflower than the wildtype strain [8], corroborating the idea that hydathodes themselves can already mount an effective immune response upon pathogen recognition when the hydathode cavity is colonized by Xcc.

Additional data from Arabidopsis hydathodes supports this notion, as mutants lacking PRR co-receptors BRI1-ASSOCIATED RECEPTOR KINASE 1 (BAK1) and its close homolog BKK1 (BAK1-LIKE 1, SERK4) are more susceptible to hydathode colonization by bacterial pathogens [16]. In addition, the immune signaling hub formed by the proteins ENHANCED DISEASE SUSCEPTIBILITY 1 (EDS1) – PHYTOALEXIN DEFICIENT 4 (PAD4) – ACTIVATED DISEASE RESISTANCE 1 (ADR1) was shown to be involved in restricting hydathode colonization. On top of that, a formate dehydrogenase (AtFDH1), known to be involved in several abiotic stresses, is linked to restricting hydathodes from being colonized by Xcc [27]. Hence, it becomes apparent that an effective, post-invasive immune response exists in plants against bacteria in hydathodes, yet the causal receptors for this tissue-specific immune recognition remain elusive to this date. To fill this knowledge gap on hydathode immunity, we set out to identify the plant immune receptors in Arabidopsis that mount an effective immune response against Xcc and study if this response is already effective in hydathodes.

Results

A QTL on Chromosome 5 is associated with resistance against Xcc *ΔxopAC*

In a recent study, we showed that the Arabidopsis accession Col-0 shows a high level of resistance to the vascular bacterial pathogen Xcc8004 in natural (guttation-based) infections and that this response was independent of the recognition of the effector XopAC by the ZAR1 resistosome [16]. Furthermore, the study showed that the accession Oy-0 is hypersusceptible to the same bacterial strain. To map the causal gene(s) for this unknown resistance trait(s), we used an existing Arabidopsis population of Recombinant Inbred Lines (RILs) between Oy-0 × Col-0 composed of 500 RILs [28]. In total, we screened 165 RILs of this population, including the two parental lines Oy-0 and Col-0, for resistance using a bioluminescent reporter strain (Tn*5:lux*) of Xcc8004 *ΔxopAC* [16,29]. For each plant, the three most-symptomatic leaves were sampled at 14 days post inoculation (dpi) and bacterial colonization was visualized using light-sensitive films (Fig 1A). The level of resistance was then scored using a luminescence index ordinal scale [29] and normalized to the Oy-0 control for each experiment (Figs 1B and C). Using an Expectation-maximization (EM) model for

a single QTL [30] and LOD threshold set at 2.49 by permutation (n = 1000), three QTLs were identified, that is, one at Chromosome 4 and two on Chromosome 5 (Fig 1D). These QTLs showed an explained variance (EV) of 10, 11 and 32%, respectively. The QTL at the end of the right arm of Chromosome 5 had an LOD score of 14.92 and thus displayed a strong genetic association with the resistance phenotype in the Oy-0 × Col-0 RIL population.

Genome-wide association study (GWAS) identified candidate R-gene *SUT1*

To identify candidate genes in this strong QTL, a GWAS was performed with the same Xcc8004 *ΔxopAC* Tn*5:lux* reporter strain on the Arabidopsis HapMap population, consisting of 360 natural accessions [31]. Disease assays were performed on a subset of 321 lines and disease was scored 14 dpi and normalized to Oy-0 control. GWA analysis was performed on phenotypic data in an online GWA portal, using an imputed SNP dataset of over five million SNPs [32]. Unexpectedly, no SNPs were identified with a significant correlation to the disease phenotype on Chromosome 5. However, several SNPs that were close to the False Discovery Rate (FDR) significance thresholds coincided with the most significant QTL in the Oy-0 × Col-0 population (Fig 1E), indicating an association with disease. These SNPs were located in the coding sequence of the gene At5g63020, which codes for SUPPRESSOR OF TOPP4 (SUT1), a previously characterized CNL involved in plant immunity [33]. Combining the results of the RIL and the GWAS analyses, we concluded that the *SUT1* gene could be involved in resistance against Xcc in Arabidopsis.

Fig 1. Mapping of *SUT1* as a candidate R-gene against Xcc8004 *ΔxopAC* in Arabidopsis Col-0. A) Disease symptoms and bacterial luminescence in the resistant Col-0 and hypersusceptible Oy-0 accessions following spray inoculation with bioluminescent Xcc8004 *ΔxopAC.* Leaves shown are representative of spray inoculation with either Tn*5:lux* or Tn*7:lux* tagged reporter strains. Bacterial spread is visualized on light-sensitive film 14 days post inoculation (dpi). **B)** Histogram depicting the variation in luminescence index score 14 dpi (normalized to Oy-0) for all 165 Oy-0 × Col-0 Recombinant Inbred Lines (RILs) following spray inoculation with Xcc8004 *ΔxopAC* Tn*5:lux*. **C)** Population structure of 165 RILs in the Oy-0 × Col-0 RIL population, ordered by luminescence index score at 14 dpi (normalized to Oy-0) shown

in B). **D)** QTL graph showing LOD scores for each of the 85 markers in the RIL
population (Expectation-maximization algorithm). LOD significance threshold
determined by permutation (n = 1000) at 2.49 **E)** Manhattan plot of Chromosome 5
from a screen of 321 accession present in the HapMap collection. Highlighted SNPs
in orange are located within the coding sequence of *SUT1*. False Discovery Rate
(FDR) thresholds set at $-\log_{10}$ (p) = 8 (Bonferroni) and $-\log_{10}$ (p) = 7.29 (Benjamini-
Hochberg).

SUT1 is involved in hydathode resistance against Xcc

To further investigate the role of SUT1 in the differential immune response between
Oy-0 and Col-0, we aligned the respective *SUT1* alleles and identified seven SNPs in
the *SUT1^Oy-0* coding sequence, of which six are non-synonymous and were predicted
to result in amino acid substitutions (Figs 2A and S1). Five of these predicted
substitutions are in the leucine-rich repeat (LRR) domain, which is known to be a
surface for protein-protein interactions [34] and is often involved in target sensing
and/or ligand binding [20,35]. Notably, the stop codon in *SUT1^Col-0* is replaced by an
Arginine codon in *SUT1^Oy-0* (CGA), leading to a predicted 17-residue peptide extension
of SUT1^Oy-0, which potentially could reach into the binding pocket of the LRR domain
based on the Alphafold2 protein structure prediction (Figs 2A and S2-S3). These
sequence differences between *SUT1^Col-0* and *SUT1^Oy-0* could explain the difference in
disease resistance due to impairment of SUT1^Oy-0 to mount an immune response
against Xcc8004 *ΔxopAC*. To test if a null allele in Col-0 would lead to loss-of-
resistance, we assessed Xcc disease susceptibility in *sut1* T-DNA insertion mutants.
Two T-DNA insertion lines, *sut1-8* and *sut1-9* (Fig 2B), were previously shown to
display a loss of SUT1-mediated autoimmunity in a *topp4-1* background [33], implying
the absence of a functional *SUT1* gene. To confirm a role of SUT1 in Xcc resistance,
we performed spray inoculations assays with Xcc8004 *ΔxopAC* Tn*7:lux* and assessed
the level of resistance both in hydathodes (7 dpi) and late stages of infection (14 dpi).
Compared to the wildtype Col-0 control, both *sut1* T-DNA insertion lines showed an
increased number of hydathodes with severe bacterial infections at 7 dpi (Figs 2C and
D), more pronounced leaf chlorosis (Figs 2C and E) combined with a more advanced
bacterial spread across the infected leaves at 14 dpi (Figs 2C and F). During both the
early and late stages of infection, *sut1* knockouts showed an intermediate
susceptibility phenotype compared to plants of the immunocompromised *bak1-5;bkk1-*

1 (*bb*) and the hypersusceptible Oy-0 accession. This corroborates the identification

of several QTLs with significant LOD scores in the RIL population (Fig 1D) indicative

for the presence of one or more loci involved in disease resistance, in addition to

SUT1. A previous study showed that SUT1 mediates an auto-immune response in a

genetic background that encodes a specific modification in a TYPE 1 PROTEIN

PHOPSHATASE 4 (TOPP4), suggesting that SUT1 guards the TOPP4 protein [33].

To infer if TOPP4 is also important for Xcc resistance in Col-0, a previously

characterized *topp4* knockout line was inoculated with Xcc8004 *ΔxopAC* Tn*7:lux*. A

small but significant increase in the number of heavily infected hydathodes was seen

in the *topp4* knockout line, but no increase in the bacterial spread across the leaf or

leaf chlorosis during later infection stages (Fig S4), suggesting that the SUT1-

mediated immunity to Xcc does not act through guarding TOPP4. These results

indicate that SUT1 restricts Xcc proliferation inside the hydathodes, thereby reducing

Black rot disease symptoms and the systemic colonization of the plant vasculature

during later infection stages.

Fig 2. SUT1 contributes to early hydathode resistance against Xcc8004

***ΔxopAC.* A)** Predicted SUT1 protein model. Arrows depict predicted amino acid

substitutions in Oy-0 vs. Col-0. **B)** *SUT1* gene model depicting Col-0 allele and

approximate sites of the T-DNA insertions in the *sut1-8* and *sut1-9*. **C)** Spray

inoculation of *sut1-8* and *sut1-9* with Xcc8004 *ΔxopAC* Tn*7:lux.* Wildtype background

(Col-0), *bak1-5;bkk1-1* (Col-0) and the accession Oy-0 were included as controls.

Top, visualization of high density hydathode colonization using light sensitive films (7

dpi); *bottom*, visualization of the systemic spread of Xcc and disease symptom

development (14 dpi). **D,E,F)** Number of infected hydathodes 7 dpi (**D**), chlorotic

area 14 dpi (**E**) and bacterial colonization 14 dpi (**F**) per leaf for each line in C)

following spray inoculation with Xcc8004 *ΔxopAC* Tn*7:lux*. In panel D, the circle size

depicts the number (#) of leaves with a certain score (y-axis) from three independent

repetitions. Of note, the *sut1* mutants show an increased number of hydathode

infections 7 dpi, culminating in more pronounced disease symptoms and increased

bacterial spread compared to wildtype control 14 dpi (Col-0). Note that Oy-0 is still

more susceptible than the *sut1* mutants. Significance letters from non-parametric

Kruskal-Wallis test with Dunn's Post-Hoc test, p-value threshold = 0.05.

SUT1 **is active in hydathodes**

As SUT1-mediated resistance occurs in hydathodes, we hypothesized that the *SUT1*
gene is expressed in the hydathode epithem cells. To investigate this, transgenic
reporter lines were generated to study the activity of the *SUT1* promoter by fusing a
2.1 kb DNA fragment upstream of the start codon to the *β-glucuronidase (GUS)*
reporter gene for histochemical analysis. We detected GUS straining throughout the
leaves of 4-week-old T1 transgenic plants (Figs 3A and S5), with strong saturation in
hydathode tissue, indicating expression patterns of *SUT1*. To determine whether this
GUS staining corresponds to true expression in the epithem tissue, we used a second
independent reporter system, *NLS-3×mVENUS* [36]. As reference for epithem cells,
we used the epithem marker line *pPUP1::RCl2A-tdTomato* [16], which shows red
fluorescence at the epithem cell membrane (Fig 3B). Col-0 plants carrying the
pSUT1::NLS-3×mVENUS constructs showed fluorescent mVENUS signal in the nuclei
of epithem cells and throughout the mesophyll tissue (Figs 3C and S6). Taken
together, our data confirm the expression of *SUT1* in the hydathodes.

Fig 3. *SUT1* is expressed in epithem cells within hydathodes. A) Arabidopsis
pSUT1::GUS lines stained with X-gluc show expression of *SUT1* throughout 4-week-
old leaves. Strong saturation of signal can be seen in hydathode tissue (white
arrows). **B)** Arabidopsis epithem marker line *pPUP1::RCl2a-tdTomato* shows
tdTomato localization in epithem cell membrane within hydathodes. Epithem cells
can be distinguished from mesophyll cells in the bright-field image (white dashed
line). **C)** Arabidopsis *pSUT1::NLS3×mVENUS* T$_2$ line shows accumulation of
mVENUS in nuclei of mesophyll and epithem cells. Epithem cells can be
distinguished from mesophyll cells in the bright-field image (white dashed line).
Images were taken of epidermal peels and are representative of three independent
lines (Fig S6).

SUT1-mediated resistance is ineffective when hydathodes are bypassed

306 As SUT1 provides resistance to Xcc already during the early stages of infection, we
307 assessed whether SUT1 also contributes to resistance at later stages of the infection
308 or upon wound infection when the bacteria cause a systemic infection along the leaf
309 vasculature. To this end, clipping inoculations were used to provide the bacteria with
310 direct access to the midvein and some lateral veins near the leaf apex, thereby

bypassing the hydathodes. We quantified both the bacterial spread along the midvein and the relative chlorotic leaf area (as a proxy of disease symptom development) (Fig 4A). Strikingly, the analysis revealed that the SUT1-mediated resistance was lost when the hydathodes were bypassed as shown by similar levels of leaf chlorosis in the different genetic backgrounds (Fig 4B). Moreover, by quantifying the total distance the bacteria had progressed along the midvein, we found that long distance movement and colonization itself appears to be unchanged in *sut1-9* compared to the Col-0 wildtype control (Fig 4C). This implies that SUT1 does not provide a high level of resistance to Xcc when the bacteria are directly introduced into vascular tissue by leaf clipping. This tissue-specific response is different from ZAR1-mediated resistance, which remains effective even when the bacteria are introduced directly into the midvein (Fig S7). Overall, we report here on the identification of an NLR that contributes to resistance against Xcc primarily in hydathodes.

Fig 4. SUT1 resistance against Xcc is ineffective when hydathodes are bypassed during inoculation. A) Disease symptoms and bacterial spread in Col-0 leaves at 7 dpi following clip inoculation with Xcc8004 *ΔxopAC* Tn*7:lux*. Bacterial progression is quantified as the spread of the luminescence signal along the midvein. **B,C)** Chlorotic leaf area (**B**) and bacterial spread along the midvein (**C**) 7 dpi following clip inoculation with Xcc8004 *ΔxopAC* Tn*7:lux*. SUT1 resistance against Xcc is apparently ineffective in the vasculature of Col-0 when the hydathodes are bypassed and the bacteria are introduced using leaf clipping. Significance letters from a two-way ANOVA with Tukey Post-Hoc test, p-value threshold = 0.05.

Discussion

Here, we characterized a role for Arabidopsis SUT1 as a functional immune receptor in natural infections by Xcc. We show that SUT1 acts primarily in hydathodes, restricting Xcc colonization in early stages of infection and thereby reducing the subsequent systemic spread of Xcc and with that the onset of disease symptoms. Interestingly, SUT1-mediated resistance is strongly compromised when the hydathodes are bypassed, indicating that SUT1 confers a first layer of defense in the Xcc disease cycle. This role of SUT1 in hydathodes differs from other reports on Xcc immunity in Arabidopsis, such as the ZAR1 immune response mounted upon recognition of XopAC activity [23] and the AMAR1-mediated hypersensitive response

through interaction with the T3E XopAM [37]. In addition, the SUT1-mediated
resistance mechanism seems to be independent of MAPK signaling cascades, as
knockout lines of several MAPK phosphatases (MKPs) show increased vascular
spread and symptom development when Xcc is directly introduced into the vasculature
[22], while under our conditions *sut1* knockout lines do not. Therefore, the role of SUT1
in plant defense in specialized cells in hydathodes is thus far unique compared to that
of characterized NLRs in Arabidopsis [17,38]. In corroboration, *SUT1* gene is
expressed in the epithem cells within hydathodes.

As *SUT1* encodes an intercellular immune receptor, our data support the notion that
Xcc actively manipulates epithem tissue to gain access to the vascular system of the
plant. Importantly, the causal agent for activation of SUT1-mediated immunity remains
unknown albeit that an indirect activation mechanism of SUT1 through guarding the
TOPP4 phosphatase has been proposed in the context of auto-immunity [33]. While
we did not find evidence for this TOPP4 guard-guardee model in the interaction with
Xcc (Fig S4), the two closest homologs of SUT1 in Arabidopsis, RESISTANT TO P.
SYRINGAE 5 (RPS5) and SUPPRESSOR OF MKK1 MKK2 2 (SUMM2) [33], have
both been described as 'guards' in plant-pathogen interactions [39,40], which
highlights the possibility of a similar function of SUT1. TOPP4 is part of an Arabidopsis
gene family of highly similar protein phosphatases called TOPP1 to TOPP9. This
family is involved in abscisic acid (ABA) signaling and leaf water homeostasis [41,42].
Interestingly, multiple TOPP family members are targeted by the *Pseudomonas*
syringae T3E AvrE during infection to hijack ABA signaling thereby suppressing the
reopening of the stomata upon bacterial entry to induce water soaking of the leaf
apoplast—a phenomenon that supports bacterial colonization of the leaf apoplast [43–
45]. Interestingly, Xcc8004 is known to employ T3E XopAM, which belongs to the
same protein superfamily of conserved bacterial effectors as *Pseudomonas* AvrE [46].
Nonetheless, AvrE superfamily members can exhibit a wide variety of functions [47]
and recent work using Y2H-screens on an Arabidopsis cDNA library did not reveal a
protein-protein interaction between XopAM and individual TOPPs [37]. Although the
elicitor of SUT1-mediated resistance has not been identified, it might be associated
with manipulation of TOPPs and therefore ABA signaling.

How manipulation of ABA signaling might benefit a hydathode pathogen remains
unclear. While *Pseudomonas syringae* actively manipulates the stomatal aperture by

targeting ABA signaling [44,45], hydathode pores are known to not fully close in
response to (a)biotic stresses [8]. Indirect evidence for bacterial manipulation of ABA
signaling in hydathodes comes from the observation that *Pseudomonas* naturally does
not progress towards the leaf vascular tissue yet it can colonize and proliferate in
hydathodes [16]. Besides *Pseudomonas*, members of the *Xanthomonas* genus are
also known to affect ABA levels during infection. For example, transcription activator-
like effector (TALE) Tal8 from *Xanthomonas translucens* pv. undulosa (Xtu), the
causal agent of bacterial leaf streak, induces the expression of a host gene encoding
9-*cis*-epoxycarotenoid dioxygenase (*TaNCED-5BS*), which catalyzes a rate-limiting
step in ABA biosynthesis [48]. Although Xcc8004 does not harbor TALEs [49], some
evidence exists for the manipulation of ABA synthesis in Arabidopsis through induction
of the Arabidopsis homolog of *NCED5* by the T3E XopAH (AvrXccC) and increased
virulence of Xcc when ABA is applied exogenously [50]. Moreover, ABSCISIC ACID-
RESPONSIVE 1 (ABR1), a negative regulator of ABA, is required for resistance
against *Xanthomonas campestris* pv. vesicatoria (Xcv) in pepper [51]. Altogether,
these findings underline the importance of ABA signaling as a target for bacterial
manipulation including many members of the *Xanthomonas* genus, but how ABA plays
a role in Xcc hydathode colonization remains elusive.

When it comes to hydathode colonization, we see an intermediate level of resistance
in the *sut1* knockout lines in the Col-0 background compared to the hypersusceptible
accession Oy-0. This indicates the presence of other resistance mechanisms against
Xcc in hydathodes. Indeed, we observed two additional QTLs on Chromosome 4,
which were not fine mapped in this study. The underlying genes for these QTLs might
encode additional NLRs, receptor-like proteins (RLPs) or receptor-like kinases (RLKs),
some of which are known to be specifically expressed in hydathodes [52]. It seems
that the resistance phenotype seen in Col-0, in absence of ZAR1-mediated resistance,
can still be based on several and possibly disconnected resistance mechanisms of
which SUT1 resistance appears to be a major contributor. The existence of a wide
array of immune receptors in Arabidopsis holds promise for application of these genes
in *Brassica* crops to ensure durable resistance. As was shown in other crop-pathogen
interactions, *R*-gene stacking is less readily overcome by pathogens than the
implementation of single *R*-genes [53–55].

In summary, we identified a CNL-type immune receptor, SUT1, that acts in hydathodes
where it reduces Xcc virulence by limiting the initial hydathode colonization and
thereby the subsequent disease development due to vascular spread of the bacteria.
This hydathode-specific response is different from other known plant immune
responses against vascular pathogens, as the efficacy of the *SUT1* gene is lost when
this initial barrier of the hydathode is bypassed using clip inoculations that mimic
wound damage. How the SUT1-mediated immune response is activated in epithem
tissue will be part of future studies, although SUT1 likely acts as a guard of one or
more TOPPs or other plant phosphatases. Studying host manipulation by Xcc is a
promising model to elucidate the intricate interactions that are happening in
hydathodes in response to bacterial pathogens.

Materials and Methods

Plant growth conditions

Arabidopsis thaliana lines used are listed in S1 Table. For all experiments, seeds were
stratified for three to four days on wet filter paper at 4 °C in the dark, and then sown in
40-pot trays (Desh Plantpak, Tray Danish size 40 cell (8x5)) in slitpots (Pöppelmann
Teku, S 5,5 LB), using potting soil (Jongkind Substrates, Hol80 zaaigrond Nr 1) to
which 500 ml Entonem suspension (Koppert Biological Systems, approximately 1.6 x
10^6 third stage *Steinernema feltiae* nematode larvae) was added per tray to prevent
black fly multiplication in growth chambers. Hereafter, trays were placed in standard
short day (SD) conditions (11/13h light/dark, 22 °C, RH 70%). For the first five days
after sowing, trays were covered with transparent plastic domes to ensure seedling
growth.

Plasmid and transgenic plant generation

Primer sequences and plasmids used are listed in S1 Table. The plasmids to create
the *pSUT1::GUS* and *pSUT1::NLS-3×mVENUS* reporter lines in Arabidopsis were
assembled using the GreenGate system [56]. First, a 2.1 kb fragment upstream of the
start codon of *SUT1* (*AT5G63020*) in Col-0 was synthesized with external *Bsa*I sites
(Eurofins Genomics). Next, the *NLS-3×mVENUS* module was amplified from
pGreenII-229-NLS-3×mVENUS [36] with added *Bsa*I sites and cloned into pGGA000.
Final binary vectors were then assembled in the backbone pGGZ003 with either *GUS*

or *NLS-3×mVENUS* as CDS module using a multisite Goldengate reaction, leading to pGGZ_pSUT1_GUS_tUBQ10_BastaR and pGGZ_SUT1_NLS-3×mVENUS_tUBQ10_BastaR. These plasmids were electroporated into *Agrobacterium tumefaciens* strain GV3101 carrying pSOUP and Arabidopsis Col-0 was transformed using the floral dip method [57]. Several independent transgenic lines were analyzed for each construct.

Xanthomonas disease assays

Spray inoculations with bioluminescent Xcc *ΔxopAC* (Tn*5:lux* or Tn*7:lux*) were performed according to previous studies [29,16]. Briefly, Xcc strains were plated on KADO medium [58] with the appropriate antibiotics and grown for 48-72 hours at 28 °C. Bacteria were collected, washed once with 10 mM $MgSO_4$ and diluted to OD_{600} = 0.1. Silwet L-77 was added to a final concentration of 0.0002% and bacteria were sprayed on 4-to-5-week-old Arabidopsis plants (Preval sprayer, SKU# 0221). Inoculated plants were subjected to two guttation cycles over two days in a climate incubator (Snijders Labs, MC1000) to promote hydathode infection. In these cycles, temperature and relative humidity are raised (25°C, RH 95%) to induce guttation, after which a rapid drop in temperature and humidity (21°C, RH 70%) leads to re-uptake of guttation droplets. Subsequently, plants are placed for an additional 12 days at SD growth conditions (11/13h light/dark, 24-22 °C, RH 70%) in the same incubator. Clipping assays were performed by cutting 5 mm of the apex of each leaf (three leaves per plant) with scissors dipped in bacterial inoculum (OD_{600} = 0.1). Plants were subjected to the same two-day guttation cycle and placed for an additional five days at SD growth conditions in an MC1000 climate chamber.

Detection of luminescent bacteria

For spray inoculations, the three most diseased leaves of each plant were sampled 7 dpi and 14 dpi. Each leaf was glued onto an A3 white paper sheet using a grid and covered with a transparent plastic sheet [29]. Next, a light-sensitive X-ray film (Fuji Super RX) was placed on top and left for exposure overnight. The film was developed the next morning and scanned, alongside the leaves, using an A3 flatbed scanner (Epson Expression 12000XL). For 7 dpi, the obtained images were overlayed in Adobe Photoshop 2023/2024 to count the number of luminescent hydathodes per leaf. At 14 dpi, scans of leaves and light-sensitive film were analyzed using the ScAnalyzer image

analysis script [59] to determine chlorosis and bacterial colonization. For clipping assays, at 7 dpi, leaves were sampled in the same manner. For analysis, chlorosis and bacterial colonization were analyzed with ScAnalyzer. Bacterial progression through the midvein was measured using Fiji (ImageJ).

RIL screen and QTL analysis

The Oy-0 × Col-0 RIL population [28] was received from the INRA Versailles stock center (). From the 470 RILs in the population, 165 (including the parental Oy-0 and Col-0) were selected for disease assays (S2 Table). For analysis, the mean luminescence index score [29] per leaf (n = 18) for each line was normalized to the score of Oy-0 and used as input for QTL mapping in R using the R/qtl package version 1.66 [30]. LOD scores for each of the 85 markers were determined in a single QTL model using the Expectation-maximization (EM) algorithm. LOD thresholds were determined by permutation (n = 1000). Explained variance (EV) per QTL was calculated as $EV(\%) = (1 - 10(-2 * LOD/n) * 100$, where LOD is the LOD score and n is the number of RILs.

GWA screen and analysis

The HapMap seed collection was received from NASC (Borevitz lab; NASC ID, N76309; ABRC ID, CS76309). From the 360 lines in the population, 321 accessions (including the controls Oy-0 and Col-0) were selected for disease assays (S3 Table). The mean luminescence index score [29] per leaf (n = 18) for each line was normalized to Oy-0. As input for GWA analysis, the data was transformed in a binary score in which 0 = <50% and 1 = >50% of Oy-0. These phenotypes were analyzed in the GWA portal [32] using the Imputed Fullsequence Dataset, TAIR 9, containing over five million SNPs [60]. P-values were obtained by applying an accelerated mixed model (AMM) [61]. False Discovery Rate (FDR) thresholds were set at $-\log_{10}$ (p) = 8 (Bonferroni) and $-\log_{10}$ (p) = 7.29 (Benjamini-Hochberg). SNPs with $-\log_{10}$ (p) values > 6 are documented in S3 Table.

GUS staining and imaging

Transgenic GUS lines were grown for 4-5 weeks at standard SD conditions. Rosettes were taken from soil, washed in double distilled water and submerged in GUS staining solution (0.5 mM $K_3Fe(CN)_6$, 0.5 mM $K_4Fe(CN)_6$, 0.1 M $NaPO_4$ (pH 7.0), 10 mM EDTA,

0.1% (v/v) Triton X-100 and 1 mg/ml X-Gluc in double distilled water) [62] in a 50 ml falcon tube. Rosettes were twice vacuum infiltrated for two minutes and then incubated at 37 °C for two hours to avoid overstaining [62]. To destain, rosettes were incubated in 96% ethanol at room temperature for two days. The destaining solution was replaced twice until all chlorophyll had been removed. Leaves were imaged using a stereomicroscope (Leica MZ FLIII + Nikon Digital Sight DS-Fi2).

Imaging fluorescent reporter lines

Fluorescent reporter lines were grown for 4-5 weeks using SD conditions (13D/11L). To enable imaging of exposed hydathode tissue, fully expanded leaves were picked and stripped of the epidermal cell layer [63] using household sellotape. Leaves were then imaged with a digital inverted fluorescence microscope (ThermoFisher EVOS fl, RFP filter for tdTomato, YFP filter for mVENUS).

Protein sequence alignments

Coding sequences from both the Arabidopsis alleles of *SUT1* in Col-0 and Oy-0 were downloaded from Araport11 (Tair10)[64] and translated *in silico*. Predicted protein sequences were then aligned in R with ClustalOmega using the Biostrings (v.2.68.1) and msa (v.1.32.0) packages.

Alphafold2 protein structure predictions

Alphafold2 [65] predictions were performed using ColabFold (v.1.5.5) and MMseqs2 using default settings [66]. Out of the five given models, the model with the highest rank score was used for visualization of the predicted fold. Alphafold2 .pdb files are available in S1 and S2 File.

Quantification and statistical analyses

All data analyses and plotting were done in R (v.4.3.2). For statistical analysis of parametric data, a two-way ANOVA followed by a Tukey post-hoc test was performed to detect differences between groups. For non-parametric data, such as hydathode counts or proportional data, a Kruskal-Wallis test followed by a Dunn's post-hoc test was performed. All experiments were performed in at least three independent repetitions with similar results. Data from repetitions was pooled for visualization and statistical analysis.

Acknowledgements

We would like to thank Ludek Tikovsky and Harold Lemereis for the excellent care of our plants in the greenhouse. We also want to thank Manon Richard for previous work on fluorescent reporter lines in Arabidopsis. Lastly, we want to thank Niko Geldner, Laurent Noël and Gail Preston for sharing materials. This research was supported by the Topsector T&U TKI programs "Stop natural entry" and "Finding the Achilles' heel of Brassica for Black Rot disease" (grants EZ-2012-02 and TU18024 to H.A.v.d.B.) and the breeding companies Bejo Zaden B.V. and Rijk Zwaan.

Competing interests

Authors declare no competing interests.

Supporting information

S1 Fig. Alignment of predicted SUT1 protein sequences of the Arabidopsis accessions Col-0 and Oy-0. Coding sequences of *SUT1^{Col-0}* and *SUT1^{Oy-0}* were obtained from Araport11 and translated *in silico* into amino acid sequences. Alignment depicts six predicted amino acid substitutions in SUT1^{Oy-0} compared to SUT1^{Col-0}, of which one is predicted to result in a 17-residue peptide extension at the C-terminus in SUT1^{Oy-0}.

(TIFF)

S2 Fig. Alphafold2 predictions of SUT1^{Col-0} and SUT1^{Oy-0}. Alphafold2 fold prediction for the rank 1 model of SUT1^{Col-0} (**A**) SUT1^{Oy-0} (**B**), colored by pLDDT score from high (blue) to low (red). pLDDT score for best 5 models is shown in right panel. N- and C-termini show low pLDDT scores, indicating low predictive value.

(TIFF)

S3 Fig. Alphafold2 predictions indicate that SUT1^{Oy-0} C-terminal extension might interfere with its protein function. A) Alphafold2 fold prediction of SUT1^{Col-0}, colored by functional domains (CC, NBS-ARC & LRR). **B)** Alphafold2 fold prediction of SUT1^{Oy-0}, colored by functional domains (CC, NBS-ARC & LRR). Green residues indicate differences at amino acid level compared to SUT1^{Col-0}. C-terminal peptide in SUT1^{Oy-0} is predicted in the LRR pocket. **C)** Superimposed model of SUT1^{Col-0} and SUT1^{Oy-0}. Green residues indicate differences at amino acid level in SUT1^{Oy-0}.

(TIFF)

S4 Fig. SUT1-mediated resistance in Arabidopsis Col-0 does not depend on TOPP4 function. A) Number of infected hydathodes per leaf at 7 dpi following spray inoculation with Xcc8004 *ΔxopAC* Tn*7:lux*. Circles depict number of leaves for each score. The topp4 knockout line shows a slight increase in hydathode infections compared to the Col-0 control. Significance letters from non-parametric Kruskal-Wallis test with Dunn's Post-Hoc test, p-value threshold = 0.05. **B)** Bacterial colonization defined as percentage of total leaf area colonized per leaf following spray inoculation with Xcc8004 *ΔxopAC* Tn*7:lux* (14 dpi). Bacterial colonization was calculated using ScAnalyzer [59]. The *topp4* knockout line does not display increased bacterial spread compared to the wildtype control (Col-0). Significance letters from non-parametric Kruskal-Wallis test with Dunn's Post-Hoc test, p-value threshold = 0.05.

(TIFF)

S5 Fig. *SUT1* is expressed in hydathodes and mesophyll. Two independent homozygous T3 lines of 4-week-old Arabidopsis *pSUT1::GUS* show GUS enzymatic activity in hydathodes (black arrows) and mesophyll tissue throughout the leaf. Leaves were incubated in GUS staining solution for two hours.

(TIFF)

S6 Fig. *SUT1* is expressed in epithem cells within hydathodes. Three independent T2 lines of 4-week-old Arabidopsis *pSUT1::NLS-3×mVENUS* show accumulation of mVENUS in nuclei of mesophyll and epithem cells. Epithem cells can be distinguished from mesophyll cells in the bright-field image (white dashed line).

(TIFF)

S7 Fig. ZAR1-mediated immunity is effective in the leaf vasculature. A) Disease

symptoms and bacterial luminescence in resistant Col-0 and susceptible *zar1-1*

knockout lines, 7 dpi following clip inoculation with Xcc8004 WT Tn*5:lux*. **B,C)**

Chlorotic leaf area (**B**) and bacterial spread along the midvein (**C**) 7 dpi following clip

inoculation with Xcc8004 WT Tn*5:lux* on Arabidopsis Col-0, *bak1-5;bkk1-1*, *zar1-1*,

rks1-1 and Oy-0. ZAR1-mediated resistance against Xcc is effective in the

vasculature of Col-0 and *bak1-5;bkk1-1* when the hydathodes are bypassed and the

bacteria are introduced using leaf clipping. Significance letters from a two-way

ANOVA with Tukey Post-Hoc test, p-value threshold = 0.05.

(TIFF)

S1 Table. List of strains, genotypes and oligonucleotides used in this study.

(XLSX)

S2 Table. Raw data of RIL population genetic structure and disease scoring.

(XLSX)

S3 Table. Raw data on disease scoring of HapMap population for GWAS.

(XLSX)

S1 File. SUT1 Col-0 Alphafold2 fold prediction model 1

(PDB)

S2 File. SUT1 Oy-0 Alphafold2 fold prediction model 1

(PDB)

References

1. Melotto M, Zhang L, Oblessuc PR, He SY. Stomatal defense a decade later.
Plant Physiol. 2017;174(2):561–71.

2. Zhang J, Coaker G, Zhou JM, Dong X. Plant immune mechanisms: from
reductionistic to holistic points of view. Mol Plant. 2020;13(10):1358–78.

3. Zipfel C, Robatzek S, Navarro L, Oakeley EJ, Jones JDG, Felix G, et al. Bacterial
disease resistance in Arabidopsis through flagellin perception. Nature. 2004 Apr
1;428(6984):764–7.

4. Miya A, Albert P, Shinya T, Desaki Y, Ichimura K, Shirasu K, et al. CERK1, a
LysM receptor kinase, is essential for chitin elicitor signaling in Arabidopsis.
Proc Natl Acad Sci. 2007;104(49):19613–8.

5. Liu Z, Hou S, Rodrigues O, Wang P, Luo D, Munemasa S, et al. Phytocytokine
signalling reopens stomata in plant immunity and water loss. Nature. 2022 May
12;605(7909):332–9.

6. Melotto M, Underwood W, Koczan J, Nomura K, He SY. Plant stomata function in
innate immunity against bacterial invasion. Cell. 2006;126(5):969–80.

7. Thor K, Jiang S, Michard E, George J, Scherzer S, Huang S, et al. The calcium-
permeable channel OSCA1. 3 regulates plant stomatal immunity. Nature.
2020;585(7826):569–73.

8. Cerutti A, Jauneau A, Auriac MC, Lauber E, Martinez Y, Chiarenza S, et al.
Immunity at Cauliflower Hydathodes Controls Systemic Infection by
Xanthomonas campestris pv *campestris*. Plant Physiol. 2017 Jun;174(2):700–
16.

9. Gluck-Thaler E, Cerutti A, Perez-Quintero AL, Butchacas J, Roman-Reyna V,
Madhavan VN, et al. Repeated gain and loss of a single gene modulates the
evolution of vascular plant pathogen lifestyles. Sci Adv. 2020 Nov
13;6(46):eabc4516.

10. Hugouvieux V, Barber CE, Daniels MJ. Entry of *Xanthomonas campestris* pv.
campestris into Hydathodes of *Arabidopsis thaliana* Leaves: A System for
Studying Early Infection Events in Bacterial Pathogenesis. Mol Plant-Microbe
Interactions®. 1998 Jun;11(6):537–43.

11. Cerutti A, Jauneau A, Laufs P, Leonhardt N, Schattat MH, Berthomé R, et al.
Mangroves in the leaves: anatomy, physiology, and immunity of epithemal
hydathodes. Annu Rev Phytopathol. 2019;57:91–116.

12. Jauneau A, Cerutti A, Auriac MC, Noël LD. Anatomy of leaf apical hydathodes in
four monocotyledon plants of economic and academic relevance. PLoS One.
2020;15(9):e0232566.

13. Routaboul JM, Bellenot C, Clément G, Citerne S, Remblière C, Charvin M, et al.
Arabidopsis hydathodes are sites of intense auxin metabolism and nutrient
scavenging. bioRxiv; 2022.

14. Bellenot C, Routaboul JM, Laufs P, Noël LD. Hydathodes. Curr Biol.
2022;32(14):R763–4.

15. Jian Y, Gong D, Wang Z, Liu L, He J, Han X, et al. How plants manage
pathogen infection. EMBO Rep. 2024;25(1):31–44.

16. Paauw M, van Hulten M, Chatterjee S, Berg JA, Taks NW, Giesbers M, et al.
Hydathode immunity protects the Arabidopsis leaf vasculature against
colonization by bacterial pathogens. Curr Biol. 2023 Feb;S0960982223000131.

17. Ngou BPM, Ding P, Jones JDG. Thirty years of resistance: Zig-zag through the
plant immune system. Plant Cell. 2022 May 1;34(5):1447–78.

18. Ngou BPM, Jones JDG, Ding P. Plant immune networks. Trends Plant Sci. 2022
Mar;27(3):255–73.

19. Büttner D. Behind the lines–actions of bacterial type III effector proteins in plant
cells. FEMS Microbiol Rev. 2016;40(6):894–937.

20. Wang J, Hu M, Wang J, Qi J, Han Z, Wang G, et al. Reconstitution and
structure of a plant NLR resistosome conferring immunity. Science.
2019;364(6435):eaav5870.

21. Xin XF, Kvitko B, He SY. Pseudomonas syringae: what it takes to be a
pathogen. Nat Rev Microbiol. 2018 May;16(5):316–28.

22. Lin H, Wang M, Chen Y, Nomura K, Hui S, Gui J, et al. An MKP-MAPK protein
phosphorylation cascade controls vascular immunity in plants. Sci Adv.
2022;8(10):eabg8723.

23. Wang G, Roux B, Feng F, Guy E, Li L, Li N, et al. The decoy substrate of a
pathogen effector and a pseudokinase specify pathogen-induced modified-self
recognition and immunity in plants. Cell Host Microbe. 2015;18(3):285–95.

24. Gay PA, Tuzun S. Involvement of a novel peroxidase isozyme and lignification
in hydathodes in resistance to black rot disease in cabbage. Can J Bot. 2000
Sep 1;78(9):1144–9.

25. Lam E, Kato N, Lawton M. Programmed cell death, mitochondria and the plant
hypersensitive response. Nature. 2001 Jun 1;411(6839):848–53.

26. Morel JB, Dangl JL. The hypersensitive response and the induction of cell death
in plants. Cell Death Differ. 1997 Dec 1;4(8):671–83.

27. Marzorati F, Vigani G, Morandini P, Murgia I. Formate dehydrogenase
contributes to the early Arabidopsis thaliana responses against Xanthomonas
campestris pv campestris infection. Physiol Mol Plant Pathol. 2021
Apr;114:101633.

28. Simon M, Loudet O, Durand S, Bérard A, Brunel D, Sennesal FX, et al. QTL
mapping in five new large RIL populations of Arabidopsis thaliana genotyped
with consensus SNP markers. Genetics. 2008;178(2253):64.

29. van Hulten M, Chatterjee S, van den Burg HA. Infection Assay for Xanthomonas
campestris pv. campestris in Arabidopsis thaliana Mimicking Natural Entry via
Hydathodes. In: Plant Innate Immunity. Humana, New York, NY: Springer;
2019. p. 159–85.

30. Broman KW, Wu H, Sen Ś, Churchill GA. R/qtl: QTL mapping in experimental
crosses. bioinformatics. 2003;19(7):889–90.

31.   Weigel D, Mott R. The 1001 genomes project for Arabidopsis thaliana. Genome
697   Biol. 2009;10:1–5.

698   32.   Seren Ü. GWA-Portal: Genome-Wide Association Studies Made Easy. In:
699   Ristova D, Barbez E, editors. Root Development: Methods and Protocols
700   [Internet]. New York, NY: Springer New York; 2018. p. 303–19. Available from:
701   https://doi.org/10.1007/978-1-4939-7747-5_22

702   33.   Yan J, Liu Y, Huang X, Li L, Hu Z, Zhang J, et al. An unreported NB-LRR
703   protein SUT1 is required for the autoimmune response mediated by type one
704   protein phosphatase 4 mutation (topp4-1) in Arabidopsis. Plant J.
705   2019;100(2):357–73.

706   34.   Padmanabhan M, Cournoyer P, Dinesh-Kumar SP. The leucine-rich repeat
707   domain in plant innate immunity: a wealth of possibilities. Cell Microbiol.
708   2009;11(2):191–8.

709   35.   Martin R, Qi T, Zhang H, Liu F, King M, Toth C, et al. Structure of the activated
710   ROQ1 resistosome directly recognizing the pathogen effector XopQ. Science.
711   2020;370(6521):eabd9993.

712   36.   Zhou F, Emonet A, Dénervaud Tendon V, Marhavy P, Wu D, Lahaye T, et al.
713   Co-incidence of Damage and Microbial Patterns Controls Localized Immune
714   Responses in Roots. Cell. 2020 Feb;180(3):440-453.e18.

715   37.   Xie Q, Wei B, Zhan Z, He Q, Wu K, Chen Y, et al. Arabidopsis membrane
716   protein AMAR1 interaction with type III effector XopAM triggers a hypersensitive
717   response. Plant Physiol. 2023 Dec 1;193(4):2768–87.

718   38.   Van de Weyer AL, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, Witek K, et
719   al. A species-wide inventory of NLR genes and alleles in Arabidopsis thaliana.
720   Cell. 2019;178(5):1260-1272. e14.

721   39.   Pottinger SE, Innes RW. RPS5-mediated disease resistance: fundamental
722   insights and translational applications. Annu Rev Phytopathol. 2020;58:139–60.

723   40.   Zhang Z, Wu Y, Gao M, Zhang J, Kong Q, Liu Y, et al. Disruption of PAMP-
724   induced MAP kinase cascade by a Pseudomonas syringae effector activates
725   plant immunity mediated by the NB-LRR protein SUMM2. Cell Host Microbe.
726   2012;11(3):253–63.

727   41.   Hou YJ, Zhu Y, Wang P, Zhao Y, Xie S, Batelli G, et al. Type one protein
728   phosphatase 1 and its regulatory protein inhibitor 2 negatively regulate ABA
729   signaling. PLoS Genet. 2016;12(3):e1005835.

730   42.   Zhang J, Qin Q, Nan X, Guo Z, Liu Y, Jadoon S, et al. Role of protein
731   phosphatase1 regulatory subunit3 in mediating the abscisic acid response.
732   Plant Physiol. 2020;184(3):1317–32.

733   43.   Hu Y, Ding Y, Cai B, Qin X, Wu J, Yuan M, et al. Bacterial effectors manipulate
734   plant abscisic acid signaling for creation of an aqueous apoplast. Cell Host
735   Microbe. 2022;30(4):518-529. e6.

44. Roussin-Léveillée C, Lajeunesse G, St-Amand M, Veerapen VP, Silva-Martins
G, Nomura K, et al. Evolutionarily conserved bacterial effectors hijack abscisic
acid signaling to induce an aqueous environment in the apoplast. Cell Host
Microbe. 2022 Apr;30(4):489-501.e4.

45. Xin XF, Nomura K, Aung K, Velásquez AC, Yao J, Boutrot F, et al. Bacteria
establish an aqueous living space in plants crucial for virulence. Nature. 2016
Nov;539(7630):524–9.

46. Degrave A, Siamer S, Boureau T, Barny MA. The AvrE superfamily: ancestral
type III effectors involved in suppression of pathogen-associated molecular
pattern-triggered immunity: The AvrE superfamily of type III effectors. Mol Plant
Pathol. 2015 Oct;16(8):899–905.

47. Nomura K, Andreazza F, Cheng J, Dong K, Zhou P, He SY. Bacterial pathogens
deliver water-and solute-permeable channels to plant cells. Nature.
2023;621(7979):586–91.

48. Peng Z, Hu Y, Zhang J, Huguet-Tapia JC, Block AK, Park S, et al.
Xanthomonas translucens commandeers the host rate-limiting step in ABA
biosynthesis for disease susceptibility. Proc Natl Acad Sci. 2019 Oct
15;116(42):20938–46.

49. Denancé N, Szurek B, Doyle EL, Lauber E, Fontaine-Bodin L, Carrère S, et al.
Two ancestral genes shaped the Xanthomonas campestris TAL effector gene
repertoire. New Phytol. 2018;219(1):391–407.

50. Ho YP, Tan CM, Li MY, Lin H, Deng WL, Yang JY. The AvrB_AvrC domain of
AvrXccC of Xanthomonas campestris pv. campestris is required to elicit plant
defense responses and manipulate ABA homeostasis. Mol Plant Microbe
Interact. 2013;26(4):419–30.

51. Choi DS, Hwang BK. Proteomics and functional analyses of pepper abscisic
acid–responsive 1 (ABR1), which is involved in cell death and defense
signaling. Plant Cell. 2011;23(2):823–42.

52. Wu Y, Xun Q, Guo Y, Zhang J, Cheng K, Shi T, et al. Genome-wide expression
pattern analyses of the Arabidopsis leucine-rich repeat receptor-like kinases.
Mol Plant. 2016;9(2):289–300.

53. Shehryar K, Khan RS, Iqbal A, Hussain SA, Imdad S, Bibi A, et al. Transgene
stacking as effective tool for enhanced disease resistance in plants. Mol
Biotechnol. 2020;62:1–7.

54. Zhang J, Hewitt TC, Boshoff WH, Dundas I, Upadhyaya N, Li J, et al. A
recombined Sr26 and Sr61 disease resistance gene stack in wheat encodes
unrelated NLR genes. Nat Commun. 2021;12(1):3378.

55. Zhu S, Li Y, Vossen JH, Visser RG, Jacobsen E. Functional stacking of three
resistance genes against Phytophthora infestans in potato. Transgenic Res.
2012;21:89–99.

56. Lampropoulos A, Sutikovic Z, Wenzl C, Maegele I, Lohmann JU, Forner J.
GreenGate-a novel, versatile, and efficient cloning system for plant
transgenesis. PloS One. 2013;8(12):e83043.

57. Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J. 1998;16(6):735–43.

58. Kado CI, Heskett MG. Selective media for isolation of agrobacterium,
Corynebacterium, Erwinia, Pseudomonas and Xanthomonas. Phytopathology.
1970;60(6):969–76.

59. Paauw M, Hardeman G, Taks NW, Lambalk L, Berg JA, Pfeilmeier S, et al.
ScAnalyzer: an image processing tool to monitor plant disease symptoms and
pathogen spread in Arabidopsis thaliana leaves. Plant Methods. 2024 May
31;20(1):80.

60. Atwell S, Huang YS, Vilhjálmsson BJ, Willems G, Horton M, Li Y, et al.
Genome-wide association study of 107 phenotypes in Arabidopsis thaliana
inbred lines. Nature. 2010;465(7298):627–31.

61. Zhang Z, Ersoz E, Lai CQ, Todhunter RJ, Tiwari HK, Gore MA, et al. Mixed
linear model approach adapted for genome-wide association studies. Nat
Genet. 2010;42(4):355–60.

62. Asai S, Cevik V, Jones JD, Shirasu K. Cell-specific RNA profiling reveals host
genes expressed in Arabidopsis cells haustoriated by downy mildew. Plant
Physiol. 2023;193(1):259–70.

63. Wu FH, Shen SC, Lee LY, Lee SH, Chan MT, Lin CS. Tape-Arabidopsis
Sandwich - a simpler Arabidopsis protoplast isolation method. Plant Methods.
2009 Nov 24;5(1):16.

64. Cheng CY, Krishnakumar V, Chan AP, Thibaud-Nissen F, Schobel S, Town CD.
Araport11: a complete reannotation of the Arabidopsis thaliana reference
genome. Plant J. 2017 Feb 1;89(4):789–804.

65. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al.
Highly accurate protein structure prediction with AlphaFold. Nature. 2021 Aug
1;596(7873):583–9.

66. Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M.
ColabFold: making protein folding accessible to all. Nat Methods. 2022 Jun
1;19(6):679–82.

Fig 1

Fig 2

Fig 3

Fig 4