1

Title: **Age-related MICOS Complex Dysregulation Impairs**

Mitochondrial 3D Architecture and Metabolic Homeostasis in the Liver

Short title: **MICOS impairment in aged liver mitochondria**

Sepiso K. Masenga[1,2,3*], Alexandria Murphy[4*], Prasanna Venkhatesh[7*], Zer Vue[1], Ashlesha Kadam[5], Andrea G. Marshall[1,6], Benjamin Rodriguez[1], Estevão Scudese[1], Brenita Jenkins[4], Amber Crabtree[1], Praveena Prasad[4], Edgar Garza-Lopez[5,6], Han Le[1], Ky'Era V. Actkins[7], Elma Zaganjor[1], Nelson Wandira[1,3], Jeremiah Afolabi[3], Prasanna Katti[7], Chantell Evans[8], Young Do Koo[9], Dhanendra Tomar[10], Mark A. Phillips[11], David Hubert[11], Chandravanu Dash[12,13,14], Pooja Jadiya[10], Olujimi A. Ajijola[15], Magdalene Ameka[16], Okwute M. Ochayi[17], Eric Wang[18], Quinton Smith[18], Ronald McMillan[3], Annet Kirabo[1,3,], André Kinder [19], Tyne W. Miller-Fleming[3], Bret Mobley[20], Julia D. Berry[20], Nathan Winn[1,21], Vernat Exil[22], Anita M. Quintana[23], Kit Neikirk[1], Jenny Schafer[24], Sean Schaffer[24], Oleg Kovtun[25], Mohd Mabood Khan[1,3,26], Calixto Pablo Hernandez Perez[1,27], Margaret Mungai[1], Melanie R. McReynolds[#3], Antentor Hinton, Jr[#^1]

Author Affiliations
1. Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, 37232, USA.
2. Department of Physiological Sciences, Pathology and Microbiology, HAND research group, Mulungushi University School of Medicine and Health Sciences, Livingstone, Zambia.
3. Department of Medicine, Division of Genetic Medicine, Vanderbilt University Medical Center, Nashville, TN 37232, USA
4. Department of Biochemistry and Molecular Biology, The Huck Institute of the Life Sciences, Pennsylvania State University, State College, PA 16801
5. Department of Internal Medicine, Section of Cardiovascular Medicine, Wake Forest University School of Medicine, Winston-Salem, NC 27157 USA
6. Department of Biomedical Sciences, School of Graduate Studies, Meharry Medical College, Nashville, TN 37208-3501, USA.
7. Department of Biology, Indian Institute of Science Education and Research (IISER) Tirupati, AP, 517619, India.
8. Department of Cell Biology, Duke University School of Medicine, Durham, NC, 27708, USA.
9. Division of Endocrinology, Diabetes and Metabolism, Department of Medicine, David Geffen School of Medicine and UCLA Health, University of California-Los Angeles, Los Angeles, CA, 90095, USA
10. Department of Internal Medicine, Section of Cardiovascular Medicine, Wake Forest University School of Medicine, Winston-Salem, NC 27157 USA
11. Department of Integrative Biology, Oregon State University, Corvallis, OR, 97331, USA.
12. The Center for AIDS Health Disparities Research, Meharry Medical College, Nashville, TN.

2

13. Department of Microbiology, Immunology, and Physiology, Meharry Medical
College, Nashville, TN.
14. Department of Biochemistry, Cancer Biology, Pharmacology and
Neuroscience, Meharry Medical College, Nashville, TN
15. UCLA Cardiac Arrhythmia Center, University of California, Los Angeles, CA,
USA.
16. Department of Medical Microbiology and Immunology, KAVI-ICR, University of
Nairobi, Nairobi, Kenya.
17. Department of Physiology, Faculty of Basic Medical Sciences, Baze University
Abuja, Nigeria.
18. Department of Chemical and Biomolecular Engineering, University of
California, Irvine, CA, 92697, USA.
19. Artur Sá Earp Neto University Center - UNIFASE-FMP, Petrópolis Medical
School, Brazil.
20. Department of Pathology, Vanderbilt University Medical Center, Nashville, TN,
37232, USA
21. Department of Internal Medicine, Section of Gerontology and Geriatric
Medicine, Sticht Center for Healthy Aging and Alzheimer's Prevention, Wake
Forest University School of Medicine, Winston-Salem, NC
22. Department of Pediatrics, Division of Cardiology, St. Louis University School of
Medicine, Missouri, St. Louis, USA.
23. Department of Biological Sciences, Border Biomedical Research Center, The
University of Texas at El Paso, El Paso, Texas, 79968, USA
24. Department of Cell and Developmental Biology, Vanderbilt University,
Nashville, TN, 37232, USA
25. Department of Chemistry, Vanderbilt University, Nashville, TN, 37232, USA.
26. Department of Medicine, Division of Clinical Pharmacology, Vanderbilt
University Medical Center, Nashville, TN 37232, USA.
27. Department of Medicine, Division of Cardiology, Vanderbilt University Medical
Center, Nashville, TN 37232, USA.
73
74 *These authors share co-first authorship.
75 #These authors share co-senior authorship.
76
77 ^Corresponding author. Address: 2222 Pierce Avenue, Robinson Research Building
78 750, Nashville, TN, 37212, USA; Tel.: 319-383-3095 (A.H Hinton).
79 *E-mail address*: antentor.o.hinton.jr@Vanderbilt.Edu

80
81 **Keywords:** Aging; 3D Structure; Mitochondria; Metabolism; MICOS Complex; Liver
82 Disease; *CHCHD6; CHCHD3; OPA1*
83

ABSTRACT:

Background & Aims: Aging is associated with a significant decline in mitochondrial

function in the liver, leading to an increased risk of liver disease. This study examines

age-related changes in the mitochondrial structure of human and murine livers using a

combination of Serial Block-Face Scanning Electron Microscopy (SBF-SEM) and mass

spectrometry approaches.

Methods: This study integrates mitochondrial structure analysis in a murine model with

an analysis of liver architecture, lipogenesis, and genetically regulated gene expression

in human cohorts. We explored the Mitochondrial Contact Site and Cristae Organizing

System (MICOS) complex using SBF-SEM, three-dimensional reconstruction with Amira

software, and mass spectrometry techniques.

Results: Aging leads to a reduction in mitochondrial size and complexity, resulting in

changes in the metabolomic and lipidomic profiles of murine liver cells that are

comparable to those observed in aged human samples. We find that genetically

modeled expression of MICOS complex genes *OPA1* and *CHCHD3* is associated with

chronic liver disease phenotypes within a large biobank population. Furthermore, we

observed dysregulated mitochondrial calcium handling and increased oxidative stress

due to the disruption of the MICOS complex.

Conclusion: Our study highlights the age-associated decline in mitochondrial

complexity and metabolic regulation within the aging murine liver and the human

population. We have identified that these changes are partially attributable to the age-

related loss of the MICOS complex.

108   Impact and implications

109 This study offers new insights into the changes to mitochondrial ultrastructure that occur

110 during aging. Using SBF-SEM, the quantification of young and aged murine

111 mitochondrial structure was performed, which had previously been an underexplored

112 avenue for measuring mitochondrial changes. The discovery of mitochondrial

113 ultrastructural changes, in conjunction with measurements of age-associated metabolic

114 alterations and gene association data, provides a model for how changes in MICOS

115 expression may modulate age-related impairment of hepatic mitochondria. These

116 results provide a new model by which changes in MICOS protein expression may both

117 cause and be a potential therapeutic target for age-related impairment in hepatic

118 function.

119

120   Highlights

121 • Decreased modeled expression of *CHCHD3* in individuals of European genetic

122   ancestry is linked to liver transplant and cirrhosis, while decreased modeled

123   expression of *OPA1* in individuals of African genetic ancestry is associated with

124   chronic liver disease and cirrhosis.

125 • Aging alters liver lipid accumulation, MICOS mRNA levels, and disease markers.

126 • Aging reduces the volume and complexity of murine liver ultrastructure.

127 • Aging and diet significantly alter the MICOS complex in mice.

128 • Knockdown of *Mic60* and *Chchd6* lowers Ca^{2+} uptake, retention, and induces

129   oxidative stress in HepG2 cells.

130 ## Introduction
131

132   Mitochondria, which are essential for energy production, calcium regulation, and

133 various metabolic pathways, undergo dynamic structural changes through fusion and

134 fission to maintain their function.[1-5] Aging disrupts these dynamics, impairing

135 mitochondrial ultrastructure and contributing to various diseases.[6-8] Traditional 2D

136 imaging (e.g. TEM) reveals cristae alterations, but 3D techniques (e.g. SBF-SEM)

137 uncover novel morphologies (e.g., donut-shaped mitochondria, nanotunnels), which

138 remain underexplored in aging tissues.[9-16] In many organs and experimental models,

139 researchers have not yet performed 3D reconstruction, creating a gap in our

140 understanding of 3D mitochondrial structural changes and their molecular regulators

141 across the aging process in different tissues.

142   The liver, a metabolic hub critical for detoxification, lipid/glucose homeostasis,

143 and hormone metabolism, experiences age-related declines in mitochondrial function,

144 lipid accumulation, and increased susceptibility to hepatic diseases (e.g., cirrhosis,

145 steatohepatitis, and metabolic dysfunction-associated steatotic liver disease (MASLD),

146 among other hepatic conditions.[17,18] These pathologies correlate with mitochondrial

147 structural defects, including volume loss, cristae disorganization, and dysregulated

148 dynamics governed by proteins such as MFN1/2, OPA1, and DRP1.[1,19-23] Additionally,

149 the mitochondria contact site and cristae organizing system (MICOS) complex, which is

150 vital for cristae architecture,[24] plays a role in age-related dysfunction.[25-28] Here, we

151 integrate 3D mitochondrial reconstruction, metabolomics, and murine models (juvenile

152 vs. aged) to investigate how aging reshapes hepatic mitochondria. We explore MICOS

153 dysregulation, lipid-driven structural changes, and their consequences on calcium/ROS

homeostasis, linking these to transcriptional and metabolic shifts in human aging livers.

This work bridges the gaps in understanding 3D mitochondrial remodeling and its role in

hepatic aging.

Materials and methods
Testing genetically modeled gene expression of MICOS genes with liver disease

phenotypes in BioVU.

BioVU is a biorepository of matched genotype and de-identified electronic health record

(EHR) data curated by the Vanderbilt Institute for Clinical and Translational Research at

Vanderbilt University Medical Center. Details regarding the BioVU program, including

oversight, patient engagement, and ethical considerations, have been previously

discussed.[29,30] Briefly, this opt-in program collects leftover blood samples from routine

patient care visits at clinics across Tennessee, which are then processed for

genotyping. BioVU currently houses samples for 344,467 individuals, with ongoing

sample collection.

Genotype data for 94,474 individuals were generated on the Illumina Multi-Ethnic

Genotyping Array (MEGA[EX]) as previously described.[31] Quality control procedures

included filtering for SNPs and individual call rates, sex discrepancies, and excessive

heterozygosity. We used principal component analysis to identify individuals of

European and African genetic ancestry, using the 1000 Genomes populations as

reference.[32,33] After imputation using the Michigan Imputation server and the Haplotype

Reference Consortium (HRC) reference panel, genotype data underwent additional

quality control procedures, including filtering for imputation quality, minor allele

frequency, and Hardy-Weinberg Equilibrium.[34,35] After removing genetically related

individuals, we identified 65,363 individuals of European genetic ancestry and 12,313

individuals of African genetic ancestry for analysis. We calculated genetically regulated

gene expression (GREX) in BioVU individuals using models built from Genotype-Tissue

Expression (GTEx) version 8 project data, which includes genotype and matched

transcriptome data for 838 donors across 49 distinct tissues.[36] The best-performing

GREX models based on the highest predictive performance (r^2) from the JTI (Joint-

Tissue Imputation) approach, as implemented in the TIGAR tool, were selected for each

gene.[37-39]

To assess the relationship between *CHCHD3, CHCHD6*, and *OPA1* GREX and liver

disease, we extracted liver disease case status from the de-identified EHR information

of BioVU participants. Specifically, we tested three phenotypes or phecodes mapped

from ICD9/10 (International Classification of Diseases, 9th and 10th editions) billing

codes: cirrhosis of liver without mention of alcohol (571.51), chronic liver disease and

cirrhosis (571), and liver replaced by transplant (573.2). These phenotypes were

selected because we were able to identify at least 500 cases within the BioVU

population. Previous studies have described the mapping of phecodes to ICD9/10

codes, and researchers can access these mappings through the PheWAS package in R

(version   0.99.5-2   and   3.6.0,   respectively)   or   online

(https://phewascatalog.org/phewas/#home).[40,41] We required individuals to have at least

two documented instances of the liver disease codes on unique dates within the

medical record to be considered a case. We defined controls as individuals with no liver

disease codes in their medical records. We used logistic regression analysis to examine

the association between *CHCHD3*, *CHCHD6*, and *OPA1* GREX (predictor variables)

200 and the status of liver disease (outcome variable). Covariates included in the regression

201 models were principal components (PC1-10), sex, current age, median age of the

202 medical record, and genotype batch in the ancestry-stratified analyses. We applied the

203 most stringent global Bonferroni p-value by adjusting the p-value for all gene-tissue

204 GREX pairs tested (n=80) and all phenotypes tested (n=3, p=0.05/240). Because the

205 GREX models across tissues are highly correlated, we also generated a less stringent

206 within-tissue Bonferroni p-value adjustment, correcting the p-value for the number of

207 genes tested (n=3) and the number of phenotypes tested (n=3, p=0.05/9). Nominal

208 significance was considered $p<0.05$.

209 **Sex As a Biological Variable**

210 In this study, sex was considered as a biological variable in the experimental design. All

211 initial experimental procedures and analyses were performed with attention to potential

212 sex-dependent differences. As TEM analyses revealed minimal sex-dependent

213 differences in the assessed parameters, and given the resource-intensive nature of

214 SBF-SEM, subsequent deep ultrastructural and functional analyses were conducted

215 using male murine cohorts. While sex-specific effects on aging and longevity have been

216 reported and may influence mitochondrial morphology, this study focuses on

217 characterizing aging-associated mitochondrial alterations in a male model.

218 **Human Cohort**

219 The Vanderbilt University Institutional Review Board (IRB) approved the collection of

220 human liver tissue under the study title "Mitochondria in Aging and Disease - study of

221 archived and autopsy tissue" (IRB No. 231584). Samples came from participants who

222 provided informed consent to participate in this study. We collected samples from

participants who died from natural causes or accidents without underlying conditions

affecting the liver.

Animal Care and Maintenance

Per protocols previously described,[43] the care and maintenance of the male C57BL/6J

mice conformed to the National Institute of Health's guidelines for the use of laboratory

animals. We used six male mice for this study: three aged 3 months and three aged 2

years. The University of Iowa's Institutional Animal Care and Use Committee (IACUC)

or the University of Washington IACUC approved the housing and feeding of these

mice. We achieved anaesthesia using a mixture of 5% isoflurane and 95% oxygen.

Oil Red O

We cut OCT blocks into 7 µm thick sections, affixed to glass slides, allowed them to

equilibrate to room temperature for 10 minutes, and then stained with Oil Red O

(Sigma-Aldrich) as previously described.[44] We used Student's t-test to assess statistical

significance, ** indicates $p < 0.01$; and *$p < 0.05$.

Bile Acid

From frozen liver tissue, 100 mg of tissue homogenized in 75% ethanol, was incubated

for 2 hours, and then centrifuged at 6000g for 10 minutes. Once prepared, we

measured bile with the Mouse Total Bile Acids Assay Kit (Crystal Chem) according to

the manufacturer's instructions. We used Student's t-test to assess statistical

significance; ** indicates $p < 0.01$, and *indicates $p < 0.05$.

246

247

248

249 **Triglyceride Levels**

250 We measured the triglyceride levels in the liver and in serum collected after a 6⬜hour

251 fast using the EnzyChrom™ Triglyceride Assay Kit (BioAssay Systems), with

252 triglyceride extraction performed using a solution of isopropanol and Triton X⬜100 as

253 described previously.[45,46] We used Student's t-test to assess statistical significance; **

254 indicates $p < 0.01$, and *indicates $p < 0.05$.

255

256 **Quantification of TEM Micrographs and Parameters Using ImageJ**

257 As per established protocols, we fixed samples in a manner to avoid any bias.[47]

258 Following preparation, tissue was embedded in 100% Embed 812/Araldite resin with

259 polymerization at 60 °C overnight. After collecting ultrathin sections (90–100 nm), we

260 post-stained them with lead citrate and imaged them using a JEOL 1400+ at 80 kV,

261 equipped with a GatanOrius 832 camera. The National Institutes of Health

262 (NIH) *ImageJ* software was used for quantification of TEM images, as described

263 previously.[14,48] We used Student's t-test to assess statistical significance; ** indicates p<

264 0.01, and *p< 0.05.

265

266 **SBF-SEM Processing of Mouse Liver Tissue**

267 We performed SBF-SEM according to previously defined protocols.[10,47,48] Anesthesia

268 was induced in male mice using 5% isoflurane. After skin and hair removal, the liver

269 was treated with 2% glutaraldehyde in 100 mM phosphate buffer for 30 minutes. It was

270 then dissected into 1-mm³ cubes and further fixed in a solution containing 2.5%

271 glutaraldehyde, 1% paraformaldehyde, and 120 mM sodium cacodylate for 1 hour.

272 Fixation and subsequent steps were collected onto formvar-coated slot grids (Pella,

273 Redding, CA), stained and imaged as previously described.[10,47,48] This includes tissue

274 washing with 100 mM cacodylate buffer, incubation in a mixture of 3% potassium

275 ferrocyanide and 2% osmium tetroxide, followed by dehydration in an ascending series

276 of acetone concentrations. We then embedded the tissues in Epoxy Taab 812 hard

277 resin. We performed sectioning and imaging of the sample using a VolumeScope 2

278 SEM (Thermo Fisher Scientific, Waltham, MA). Conventional TEM analysis was

279 performed on 300–400 serial sections from each sample, following staining and imaging

280 protocols. Subsequently, analysis via imaging was performed under low-vacuum/water-

281 vapor conditions with a starting energy of 3.0 keV and beam current of 0.10 nA.

282 Sections of 50 nm thickness were cut, allowing for imaging at a spatial resolution of 10

283 nm × 10 nm × 50 nm.

284 **Segmentation and Quantification of 3D SBF-SEM Images Using Amira**

285 SBF-SEM images were manually segmented in Amira to perform 3D reconstruction, as

286 described previously[10]. We used and analyzed between 300 and 400 slices, and the

287 individual performing the analysis remained blinded to all other details. We collected a

288 total of 250 mitochondria across from each of three mice for each quantification. For 3D

289 reconstruction of liver cells, 10 cells and a total of approximately 200 mitochondria are

290 used, which meets field standards for fulfilling power requirements to determine

291 significant differences between groups. Quantification of 3D structures was performed

292 using the Amira software, which included built-in parameters or previously described

293 measurements.[10] We performed Mann–Whitney tests for statistical analysis, and

294 significance values indicate **P ≤ 0.01, ****P ≤ 0.001, and ns, not significant.

295 **LCMS Methods for Metabolomics**

296 Frozen tissues were weighed, ground with a liquid nitrogen in a cryomill (Retsch) at 25

297 Hz for 45 seconds, before extracting tissues 40:40:20 acetonitrile: methanol: water

298 +0.5% FA +15% NH4HCO3.[49] with a volume of 40mL solvent per 1mg of tissue,

299 vortexed for 15 seconds, and incubated on dry ice for 10 minutes. We then centrifuged

300 the tissue samples at 16,000 g for 30 minutes. The supernatants were transferred to

301 new Eppendorf tubes and then centrifuged again at 16,000 g for 25 minutes to remove

302 any residual debris before analysis.

303

304 Extracts were analyzed within 24 hours by liquid chromatography coupled to a mass

305 spectrometer (LC-MS). The LC–MS method was based on hydrophilic interaction

306 chromatography (HILIC) coupled to the Orbitrap Exploris 240 mass spectrometer

307 (Thermo Scientific).[50] The LC separation was performed on a xBridge BEH Amide

308 column (2.1 x 150 mm, 3.5 µm particle size, Waters, Milford, MA). Solvent A is 95%: 5%

309 H_2O: acetonitrile with 20 mM ammonium acetate and 20mM ammonium hydroxide, and

310 solvent B is 90%: 10% acetonitrile: H_2O with 20 mM ammonium acetate and 20mM

311 ammonium hydroxide. The gradient was 0 min, 90% B; 2 min, 90% B; 3 min, 75% B; 5

312 min, 75% B; 6 min, 75% B; 7 min, 75% B; 8 min, 70% B; 9 min, 70% B; 10 min, 50% B;

313 12 min, 50% B; 13 min, 25% B; 14min, 25% B; 16 min, 0% B; 18 min, 0% B; 20 min, 0%

314 B; 21 min, 90% B; 25 min, 90% B. The following parameters were maintained during the

315 LC analysis: flow rate 150 mL/min, column temperature 25 °C, injection volume 5 μL

316 and autosampler temperature was 5 °C. For the detection of metabolites, the mass

317 spectrometer was operated in both negative and positive ion mode. The following

318 parameters were maintained during the MS analysis: resolution of 180,000 at m/z 200,

319 automatic gain control (AGC) target at 3×10^6, maximum injection time of 30 ms and

320 scan range of m/z 70-1000. Raw LC/MS data were converted to mzXML format using

321 the command line "msconvert" utility.[51] Data were analyzed using the EL-MAVEN

322 software version 12. Student's t-test was used to test for statistical significance, **

323 indicates p< 0.01; and *p< 0.05.

324

325 **LCMS Methods for Lipidomic Profiling**

326 Tissue homogenization and extraction for lipids: Tissues were homogenized using a

327 Retsch CryoMill. The homogenate was mixed with 1 mL of Extraction Buffer containing

328 IPA/H2O/Ethyl Acetate (30:10:60, v/v/v) and Avanti Lipidomix Internal Standard (diluted

329 1:1000) (Avanti Polar Lipids, Inc. Alabaster, AL). Samples were vortexed and

330 transferred to bead mill tubes for homogenization using a VWR Bead Mill at 6000 g for

331 30 seconds, repeated twice. The samples were then sonicated for 5 minutes and

332 centrifuged at 15,000 g for 5 minutes at 4°C. The upper phase was transferred to a new

333 tube and kept at 4°C. To re-extract the tissues, an additional 1 mL of Extraction Buffer

334 (30:10:60, v/v/v) was added to the tube containing the tissue pellet. The samples were

335 vortexed, homogenized, sonicated, and centrifuged as described earlier. The

336 supernatants from both extractions were combined, and the organic phase was dried

337 under liquid nitrogen gas.

338

339 Sample reconstitution for lipids: The dried samples were reconstituted in 300 µL of

340 Solvent A (IPA/ACN/H2O, 45:35:20, v/v/v). After brief vortexing, the samples were

341 sonicated for 7 minutes and centrifuged at 15,000 g for 10 minutes at 4°C. The

342 supernatants were transferred to clean tubes and centrifuged again for 5 minutes at

343 15,000 g at 4°C to remove any remaining particulates. For LC-MS lipidomic analysis, 60

344 µL of the sample extracts were transferred to mass spectrometry vials.

345

346 LC-MS analysis for lipids: Sample analysis was performed within 36 hours after

347 extraction using a Vanquish UHPLC system coupled with an Orbitrap Exploris 240™

348 mass spectrometer equipped with a H-ESI™ ion source (all Thermo Fisher Scientific). A

349 Waters (Milford, MA) CSH C18 column (1.0 × 150 mm × 1.7 µm particle size) was used.

350 Solvent A consisted of Acetonitrile (ACN): H_2O (60:40; v/v) with 10 mM Ammonium

351 formate and 0.1% formic acid, while solvent B contained Iso-propyl alcohol (IPA): ACN

352 (95:5; v/v) with 10 mM Ammonium formate and 0.1% formic acid. The mobile phase

353 flow rate was set at 0.11 mL/min, and the column temperature was maintained at 65 °C.

354 The gradient for solvent B was as follows: 0 min 15% (B), 0–2 min 30% (B), 2–2.5 min

355 48% (B), 2.5–11 min 82% (B), 11–11.01 min 99% (B), 11.01–12.95 min 99% (B),

356 12.95–13 min 15% (B), and 13–15 min 15% (B). Ion source spray voltages were set at

357 4,000 V and 3,000 V in positive and negative mode, respectively. Full scan mass

358 spectrometry was conducted with a scan range from 200 to 1000 m/z, and AcquireX

359 mode was utilized with a stepped collision energy of 30% with a 5% spread for fragment

360 ion MS/MS scan. Student's t-test was used to test for statistical significance, ** indicates

361 $p < 0.01$; and *$p < 0.05$.

362

363 **RNA Extraction**

364 RT-qPCR was used to determine the change in mRNA levels in both human and mouse

365 samples. In the human cohort, the two groups were a "young" cohort (18-55 years) and

366 an "old" cohort (>60 years). In the mouse cohort, the "young" group consisted of mice

367 aged 3 months, and the "old" group consisted of mice aged 2 years. Five samples were

368 run for each experimental group for the human samples, and three samples were run

369 for each experimental group for the mouse samples. For both experimental groups,

370 comprising both human and mouse samples, at least three samples were necessary for

371 a statistical test to determine significant differences.

372 Murine liver tissue was collected via microdissection, while the human tissue was

373 collected by macrodissection. Collected samples were immediately placed in liquid

374 nitrogen to freeze and were stored in -80°C for 3 months prior to initiation of reverse

375 transcription and qPCR. To isolate RNA, we first used the TRIzol reagent (Invitrogen)

376 protocol on its user guide, with further purification of isolated RNA being performed with

377 the RNeasy Micro kit and protocol (Qiagen Inc), including DNase treatment of RNA

378 samples with kit-included DNase I. RNA concentration was determined by measuring

379 absorbance at 260 nm and 280 nm using a NanoDrop 1000 spectrophotometer

380 (NanoDrop products, Wilmington, DE, USA). The determination of significant DNA

381 contamination was assessed by the A280/A260 ratio, with a ratio of 1.8-2.0 being

382 necessary for RNA samples to be considered sufficiently pure for downstream

383 applications.

384 **RT-qPCR**

385 Reverse transcription was performed with the High-Capacity cDNA Reverse

386 Transcription Kit (Applied Biosciences, Carlsbad, CA). A total reaction volume of 20 µL,

387 containing 1 µg of RNA, was used. The kit-included MultiScribe™ Reverse

388 Transcriptase (2.5 U/µL) was used as the reverse transcriptase. We used the thermal

389 cycle recommended by the kit's protocol: 25°C for 10 minutes, 37°C for 120 minutes,

390 85°C for 5 minutes, and then held at 4°C.

391 Quantitative PCR (qPCR) was then performed using SYBR Green (Life Technologies,

392 Carlsbad, CA).[52] For qPCR, a 20 µL reaction volume containing 50 ng of cDNA was

393 loaded into each well of a 384-well plate for nuclear gene expression measurement,

394 with the reaction carried out on an ABI Prism 7900HT system (Applied Biosystems). For

395 qPCR for mtDNA, five nanograms of DNA were used for the quantification of

396 mitochondrial (Cox1) and nuclear (Rpl13a) DNA markers with Rpl13a. The reaction was

397 performed using primers (Table 1) at a concentration of 500 nM and with dNTP

398 provided by 2X SYBR® Green PCR Master Mix diluted to 1X (Life Technologies). The

399 DNA polymerase used was kit-included AmpliTaq Gold® DNA polymerase, diluted by

400 the recommended factor from 2X to 1X. The SYBR® Green PCR Master Mix also

401 included the buffer necessary for the protocol. Thermal cycling parameters were

402 adapted from previous qPCR protocol[45] with the following cycle: 1 cycle at 95°C for 10

403 min; 40 cycles of 95°C for 15 s; 59°C for 15 s, 72°C for 30 s, and 78°C for 10 s; 1 cycle

404   of 95°C for 15 s; 1 cycle of 60°C for 15 s; and one cycle of 95°C for 15 s. GAPDH

405   normalization was used to present the data as fold changes. qPCR primers used were

406   from previously published sequences,[45] as detailed in **Table 1**.

407   Specificity for fragments was determined by matching amplicon sizes, as determined by

408   gels with positive control samples. For no template controls (NTCs), detection of DNA

409   amplification remained negative after 40 cycles. Quantification of Ct for runs was

410   determined by fitting to a 6-point standard curve of log DNA dilution vs. Ct (y-intercept:

411   38.664; slope: -3.57; r^2=.96; PCR efficiency: 90.5%). Cq variation at the lower limits was

412   minimal, confirming the assay's accuracy. The limit of detection was determined by the

413   melting curve.

414   qPCR analysis was performed using the RQ Manager software on the ABI Prism

415   7900HT system (Applied Biosystems), which was utilized for qPCR. The Automatic C_T

416   setting on RQ Manager was used to determine the C_T for samples, with outliers

417   determined by using the C_T vs. Well Position View plot on the RQ Manager software.

418   GAPDH was used as the sole reference gene due to its widespread use as a reference

419   gene across multiple protocols[53]. A comparison of results across treatments was

420   performed by normalizing target gene expression to GAPDH. Intra-assay variation for

421   samples of the same treatment, run on the same plate, was measured to ensure

422   sufficient repeatability in the experiment. For each reverse transcription and qPCR

423   experiment, there were 3 technical replicates. To determine significance, Mann–

424   Whitney tests were performed with an α of 0.05, and the results were visualized in

425   GraphPad Prism 10.2.3 (La Jolla, CA, USA).

426   **Western Blotting**

Western blotting was performed as previously described.[54] Briefly, following

homogenization and lysis in RIPA lysis buffer (1% NP40, 150 mM NaCl, 25 mM Tris

base, 0.5% sodium deoxycholate, 0.1% SDS, 1% phosphatase inhibitor cocktails #2

(Sigma P5726-1ML) and #3 (Sigma P0044-1ML), and one complete protease inhibitor

tablet (Sigma 04693159001)), protein concentration in 3-month and 2-year tissue

lysates were quantified using a BCA Assay (Thermo Scientific VLBL00GD2). Equal

amounts of proteins were run on 4%–20% Tris-glycine gels (Invitrogen

WXP42012BOX). Protein was then transferred to a nitrocellulose membrane (Li-Cor

926-31092) and incubated overnight at 4°C with gentle agitation in the presence of

primary antibodies: Mic60/mitofilin (Abcam ab110329), SAM50 (Proteintech 20824-1-

AP), or tubulin (Novus NB100-690). Secondary antibodies [1:10,000; donkey anti-

mouse IgG (H+L) (Invitrogen A32789) and donkey anti-rabbit IgG (H+L)

(Invitrogen A32802)] were incubated with the membrane at room temperature for 1 h.

Using the Li-Cor Odyssey CLx infrared imaging system, blots were imaged. We

performed the student's t-test for statistical significance, with ** indicating $p < 0.01$ and

*indicating $p < 0.05$.

Confocal mCherry-Mito-7 Labeling

To label the mitochondria of cardiac fibroblasts, the mCherry-Mito-7 plasmid was

transfected into the cells using a transfection reagent according to the manufacturer's

instructions[55] and as previously described.[80] Briefly, following plasmid and transfection

reagent dilution in Opti-MEM medium and incubation at room temperature for 20

minutes, the dilution was added to the culture medium of the cells, which were

incubated for 24–48 hours to allow expression of the mCherry-Mito-7 protein.

Localization in fibroblasts was visualized using a Leica SP8 Confocal Microscope. Cells

were washed with PBS, fixed with 4% paraformaldehyde for 10 minutes, and mounted

with DAPI-containing mounting medium. Fluorescent signals were observed using

appropriate filters and recorded with a digital camera.[31,36-40,56-59]

Knockdown of *MIC60* and *CHCHD6* in HepG2 cells.

The transfection of *MIC60* and *CHCHD6* siRNAs into HepG2 cells was carried out using

Lipofectamine RNAiMax (Invitrogen) in accordance with the manufacturer's instructions

and as previously described.[60] Authentication of cell line determined by confirming

proper morphology by microscopy. Following a 48-hour incubation period, the cells were

utilized for mitochondrial calcium ($_mCa^{2+)}$ and ROS measurements.

Measurement of mitochondrial calcium uptake and retention capacity in HepG2

cells.

The $_mCa^{2+}$ uptake and retention capacity in HepG2 cells were assessed using a

ratiometric Ca^{2+} sensor Fura-FF, as detailed earlier,[61] with slight modifications. In brief,

cells ($2.5x10^6$) were washed with Ca^{2+}/Mg^{2+}-free DPBS (GIBCO), permeabilized in

intracellular medium (ICM: 120 mM KCl, 10 mM NaCl, 1 mM KH_2PO_4, 20 mM

HEPES-Tris, pH 7.2), and supplemented with thapsigargin and digitoxin. Fura-FF (1

µM) was added at the 0s time point, and fluorescence emissions at 340- and 380 nm

excitation/ and emission at 510 nm were monitored using a multi-wavelength excitation

dual-wavelength emission fluorimeter (Delta RAM, PTI). To assess the $_mCa^{2+}$ uptake, a

bolus of 5 µM Ca^{2+} and the mitochondrial uncoupler FCCP (10 µM) were added at

specified time points with continuous stirring at 37°C. To assess mCa^{2+} retention

capacity, following baseline recordings, a series of Ca^{2+} boluses (5 µM) were

introduced at specified time points. Upon reaching a steady state, 10 µM FCCP was

added to collapse the Δψm and release matrix free-Ca^{2+}. The number of Ca^{2+} boluses

taken up by cells was counted to calculate mitochondrial CRC. For all statistical tests, a

one-way ANOVA statistical test was performed with Dunnett's multiple comparisons

test. N = 5 -10 for all calcium experiments, each indicated by dots, as run in triplicate.

Significance values indicate **$P ≤ 0.01$ and ****$P ≤ 0.0001$.

Evaluation of ROS levels

Approximately 0.2 million HepG2 cells were plated in 35 mm dishes. The next day,

MIC60 and CHCHD6 siRNAs were transfected using Lipofectamine RNAiMax

(Invitrogen) according to the manufacturer's instructions. After incubation for 30 hrs.,

cells were co-stained for 30 minutes at 37°C with two different dyes for ROS detection:

MitoBright ROS Deep Red (10 µM, Dojindo Laboratories) for mitochondrial superoxide,

and DCFDA (10 µM, Invitrogen) for intracellular total ROS. Following incubation with

staining dyes, cells were washed three times with 1X HBSS, and ROS analysis was

performed using a confocal microscope (FV4000, Olympus Life Science).

For mitochondrial H_2O_2 assessment, cells were stained with MitoPY1 (5 µM, Bio-

Techne) for 45 min at 37°C. Cells were then washed with 1x HBSS and imaged using a

confocal microscope (FV4000, Olympus Life Science). ImageJ was used for quantifying

fluorescence intensities. For all statistical tests, one-way ANOVA statistical test was

496 performed with Dunnett's multiple comparisons test. N = 9 -13 for all oxidative stress

497 experiments, each indicated by dots, as run in triplicates. Significance values indicate

498 **P ≤ 0.01 and ****P ≤ 0.0001.

499

500 **Data Analysis**

501 GraphPad Prism 10.2.3 (La Jolla, CA, USA) was used for all statistical analyses. All

502 experiments involving SBF-SEM and TEM data had at least three independent

503 experiments. Statistics were not handled by those conducting the experiments. For all

504 experiments, at least three samples were used for each experimental group to enable

505 statistical testing. The black bars represent the standard error of the mean. For all

506 analyses, one-way ANOVA was performed with tests against each independent group

507 and significance was assessed using Fisher's protected least significant difference

508 (LSD) test. *, **, ***, **** were set to show significant difference, denoting $p < 0.05$, $p <$

509 0.01, $p < 0.001$, and $p < 0.0001$, respectively.

510 **Results**
511

512 **Genetically Regulated MICOS Gene Expression Associated with Liver Disease**

513 **Across Ancestries**

514 GREX models derived from GTEx v8 were applied to the BioVU cohort, enabling

515 systematic association testing between predicted expression levels of MICOS genes

516 responsible for cristae stabilization, formation, remodeling and cell survival, and liver

517 disease phenotypes across various tissues (Figure 1A). To examine the relationship

518 between MICOS complex genes *CHCHD3 (MIC19), CHCHD6 (MIC25), and OPA1* and

519 liver disease, we leveraged Vanderbilt University Medical Center's EHR-linked biobank,

BioVU (**Table 2**). We used logistic regression analysis to test whether the GREX of

CHCHD3, CHCHD6, and OPA1 was associated with liver disease phenotypes extracted

from the EHR (Figure 1B-E). Although no associations met a strict global Bonferroni-

corrected p-value threshold, we found significant associations that met the less stringent

within-tissue Bonferroni-corrected p-value (see Methods). In individuals of European

genetic ancestry (EUR), we identified significant associations between decreased

GREX of *CHCHD3* in three tissues with liver transplant and liver cirrhosis without

mention of alcohol (p < 0.0056, Figure 1B). In individuals of African genetic ancestry

(AFR), we identified significant associations between decreased GREX of *OPA1* in five

tissues and chronic liver disease and cirrhosis (p <0.0056, Figure 1E).

Aging is Associated with Coordinated Suppression of MICOS Genes and

Activation of ER Stress Pathways

To further investigate whether the MICOS genes that regulate mitochondrial

bioenergetics, ER stress, and apoptosis differ with aging, we performed qPCR in both

human (Figure 2A-I) and mouse (Figure 2J-M) and compared the results between

young and old. In the human cohort (n = 5/group), the expression of MICOS proteins

Mic19, Mic25, And Mic60 (Figure 2A, B, C) was reduced in the older compared to the

younger samples. Genes associated with regulating mitochondrial dynamics and

tethering to the ER, such as *BIP* (Figure 2D), *PERK* (Figure 2E), *XBP1* (Figure 2F),

CHOP (Figure 2G), *IRE1* (Figure 2H), and *ATF4* (Figure 2I), had increased expression.

In the murine model (n = 3/group), we observed reduced expression of MICOS proteins

including, *Opa1, Mitofilin, Chchd3, and Chchd6* (the murine ortholog of *MIC60*), in old

mice compared with young mice (Figure 2J-M).

Loss of MICOS Components Disrupts Calcium Homeostasis and Promotes

Cellular Stress in HepG2 Cells

While the reduction in expression of *Mic19, Mic25*, and *Mic60* highlights compromised

mitochondrial bioenergetics, and the increased expression of Bip, Xbp1, Ire1, Atf4 and

Chop may suggest proteostatic and metabolic stress. We sought to demonstrate and

investigate the role of the MICOS complex in age-related changes by modeling MICOS

loss *in vitro*. To investigate the role of *MIC60* and *CHCHD6* in mitochondrial calcium

($_mCa^{2+}$) homeostasis, we monitored $_mCa^{2+}$ uptake and $_mCa^{2+}$ retention capacity (CRC) in

human-derived HepG2 cells with *MIC60* and *CHCHD6* knockdown (Figure 3). Raw

mitochondrial calcium uptake traces showed reduced calcium uptake dynamics in

permeabilized HepG2 cells following *MIC60* or *CHCHD6* knockdown compared with

scramble-siRNA controls (Figure 3A and B). Furthermore, to determine if the altered

MICOS and cristae structure contributes to impaired CRC and is involved in

mitochondrial permeability transition pore opening, we measured the CRC in *MIC60* and

CHCHD6-knockdown cells. Knockdown of Mic60 as well as Chchd6 sensitized cells to

mPTP opening, as indicated by a sudden Ca^{2+} release at a lower cumulative Ca^{2+} load

compared to scramble-siRNA controls (Figure 3C). We also found that knocking of

MIC60 and *CHCHD6* in HepG2 cells significantly reduced $_mCa^{2+}$ retention capacity

565 (Figure 3D). Quantitative analysis confirmed a significant reduction in CRC in both

566 knockdown conditions relative to scramble-siRNA–treated cells (Figure 3D). Immunoblot

567 analysis verified efficient siRNA-mediated knockdown of Chchd6 (Figure 3E) and Mic60

568 (Figure 3F) in HepG2 cells. These data suggest that altered MICOS complex and

569 cristae disorganization increase the susceptibility of HepG2 cells to $_mCa^{2+}$ dysregulation

570 and Ca^{2+}-induced cell death.

571

572 Since Ca^{2+} directly affects oxidative stress signaling and the generation of reactive

573 oxygen species (ROS),[81] we wanted to determine if MICOS influences ROS production.

574 We evaluated total ROS, mitochondrial superoxide, and H_2O_2 production in *MIC60* and

575 *CHCHD6* knockdown cells (Figures 3G and H). Mitochondrial H_2O_2 content, as

576 measured by Mitochondria Peroxy Yellow 1 (Figure 3G), increased following *MIC60* and

577 *CHCHD6* knockdown when quantified by both plate-reader-based (Figure 3I) and

578 microscopy-based (Figure 3J) ROS quantification. Silencing *MIC60* and *CHCHD6* in

579 HepG2 cells significantly increased mitochondrial superoxide production and more

580 general intracellular ROS, detected by MitoBright Deep Red (Figure 3K) and DCFDA

581 (Figure 3L), respectively. These findings indicate that suppression of *MIC60* and

582 *CHCHD6* disrupts mitochondrial ROS homeostasis, demonstrating that the MICOS

583 complex is associated with oxidative stress.

584

585 **Aging is Associated with Broad Alterations in Liver Metabolism, Including**

586 **Changes in Energy Utilization and Storage**

587 Due to observed changes in liver gene expression and function during aging, we

588 proceeded to analyze alterations in small biomolecules that regulate energy

589 metabolism, cellular homeostasis, and storage. Our investigation involved global

590 metabolic profiling of young (n = 4) and aged (n = 4) liver tissues, revealing significant

591 changes in aged mouse livers (Figure 4A and B). Principal component analysis showed

592 a clear separation between the metabolic profiles of young and aged livers (Figure 4C).

593 We noted an accumulation of metabolites related to Vitamin A metabolism, specifically

594 retinoic acid and retinol (Figures 4D and E). Our metabolomics analysis also uncovered

595 an age-related decrease in tricarboxylic acid (TCA) cycle intermediates: succinate and

596 malate, which could be consistent with increasing cataplerosis associated with aging for

597 biosynthetic use or impaired activity (Figures 4Fand G). Furthermore, we observed

598 decreases in nucleotide monophosphates involved in purine and pyrimidine metabolism,

599 including GMP, CMP, UMP, and AMP (Figures 4H-K). The synthesis of these

600 nucleotides, which involves mitochondrial processes, is influenced by age-related

601 mitochondrial dysfunction, thereby affecting the overall nucleotide biosynthesis

602 pathway, which supports our observations[66]. Our results additionally confirmed existing

603 literature on dysregulated nicotinamide adenine dinucleotide (NAD^+) metabolism in the

604 aging liver (Figure 4M).[63] Significant depletions in tissue ADP, NAD^+, NADP, and NMN

605 pools were detected (Figure 4L-O). In summary, these findings collectively support the

606 presence of altered energy metabolism and cellular homeostasis in the metabolically

607 active liver, resulting from mitochondrial dysfunction and impairment.

608

609 **Aging Causes Alterations in Liver Lipid Accumulation in humans and mice**

610 Based on the evidence of an altered metabolomic profile in aged livers, we also

611 hypothesized that lipid metabolism is altered in aged livers. Damage to the liver's tissue

612 and alterations in hepatic metabolism during aging might result from abnormal lipid

613 levels in the liver.[18] First, we utilized magnetic resonance imaging (MRI) to determine

614 how lipid content changes during aging in human samples. By enrolling female and

615 male participants (n = 10 per group) (Figures 5A–D'), we created a "young" cohort

616 consisting of individuals from 18 to 55 years old and an "old" cohort of individuals older

617 than 60 years old. For both sexes, the percentage of lipid in the liver was higher in the

618 old compared to the young (Figures 5E and F). Specifically, when combining the male

619 and female cohorts (Figure 5G), the mean fat percentage was higher in the older group

620 (5.93 $\pm$ 6.11% SD) compared to the younger group (1.41 $\pm$ 1.44% SD). Additionally, the

621 75% percentile of fat percentage was higher, at 10.5% in the aged cohort, as compared

622 to 2.10% in the young cohort. Notably, Grade 1 of fat fraction classification, representing

623 mild hepatic steatosis, has a fat percentile cutoff of 6.5% or higher.[42]

624

625 To further assess the intricate changes in the lipid profile, we used murine livers. We

626 used Oil Red O, a neutral lipid stain, to visualize lipids in 3-month and 2-year murine

627 liver tissue (Figures 6A-D). We saw an increased amount of Oil Red O in the 2-year

628 cohort (n=10 per group), signifying an increase in lipid amount across the aging process

629 in the liver, with a more than 6-fold increase in the number of lipid droplets in 2-year

630 samples (Figure 6E; 3-months: 12.2 $\pm$ 6.55 SD; 2-years: 77.5 $\pm$ 20.4 SD). An uptick of

631 lipid droplets in hepatocytes is indicative of MASLD.[65] To further explore markers of

632 potential disease progression and dystrophy outside of lipid accumulation, we also

633 looked at liver weight, which was decreased in aged samples when normalized to body

634 weight (Figure 6F; 3-months: 4.56 $\pm$ 0.329% SD; 2-years: 3.93 $\pm$ 0.376% SD). Since

635 mtDNA content reduction is another hallmark of MASLD, we also measured mtDNA

636 content and observed an age-related decline (Figure 6G; 3 months: 1.00 (normalized

637 mean); 2 years: 0.805 ± 0.119 SD). Additionally, abnormal liver function and disease

638 states increase bile acid concentrations, which the liver normally metabolizes.[67] We

639 looked at the concentration of bile acid, but no significant difference was noted (Figure

640 6H; 3 months: 0.871 $\pm$ 0.138 µMol/µg SD; 2 years: 0.89 $\pm$ 0.15 µMol/µg SD). Finally,

641 central to the pathology and a key marker of MASLD is high triglycerides.[68,69] We

642 observed that aging murine samples had significantly elevated triglycerides in the liver

643 (Figure 6I; 3 months: 0.30 ± 0.10 mmol/L SD; 2 years: 0.748 ± 0.15 mmol/L SD), which

644 was more drastic than the general increase in serum triglycerides (Figure 6J; 3 months:

645 0.211 ± 0.100 mmol/L SD; 2 years: 0.379 ± 0.118 mmol/L SD). TEM analysis revealed

646 increased lipid droplet accumulation in livers from 2-year-old mice compared with 3-

647 month-old controls (Figure 6K and L). Aged samples had significantly increased both

648 the number of lipid droplets per unit area and the total lipid droplet area within TEM

649 images (Figure 6M and N), indicating enhanced hepatic lipid deposition with age. These

650 results support the observation of an age-related progression toward liver disease

651 through the accumulation of hallmarks associated with MASLD.

652

653 Lipidomic profiling of both young and aged liver tissues unveiled age-related changes in

654 lipid classes and chain lengths (Figure 6O-R). Lipid class enrichment heatmap and

655 volcano plot analyses highlighted distinct age-dependent shifts in lipid abundance

656 (Figures 6O and P). In the aging liver, we observed significant alterations in triglyceride

657 oligomers (TGO), triglycerides (TG), ceramides (Cer), and acylcarnitines (CAR) relative

658 to other lipid classes (Figure 6Q). Aging also significantly altered lipid chain-length

659 distributions in the liver (Figure 6R), suggesting changes in membrane composition that

660 may influence membrane integrity, fluidity, and function. Together, these findings align

661 with the altered metabolomic and lipogenic remodeling associated with age-related

662 changes in mitochondrial structure and function.

663

664 **Ultrastructural Changes in Murine Liver Reveal Aging is Associated with Lower**

665 **Mitochondrial Volume and Complexity**

666 Next, we sought to determine whether these age-related changes in liver mass, disease

667 markers, and metabolomic and lipidomic profiles are correlated with atypical

668 mitochondrial structures in the liver. For these studies, we utilized aged C57BL/6J mice

669 at 2 age points, 3 months, and 2 years, which are generally understood to be good

670 representations of "young" and "old" aged points akin to aging displayed in humans.[74]

671 To begin with, we used TEM to measure changes in mitochondrial morphology in aged

672 mice (Figures 7A-D). TEM images revealed an increase in lipid droplets in the livers of

673 aged individuals of both sexes, which appeared less circular. TEM analysis revealed

674 that the mitochondrial count increased while the average mitochondrial area decreased

675 with aging (Figures 7E-H). Analysis by two-way ANOVA (Age versus Sex) revealed a

676 significant main effect of age on mitochondrial count (Figure 7I) and area (Figure 7J),

677 but no significant interaction with sex, indicating similar age-related trends in both males

678 and females. In aged male and female murine samples, we found a significant decrease

679 in cristae score (a semi-quantitative measure of cristae integrity[14]) in both sexes

680 (Figures 7K and L). A two-way ANOVA confirmed a main effect of age (Figure 7M).

681 Together, these results highlight that cristae integrity is lower in aged liver tissue

682 samples, indicating a loss of ATP production efficiency. Given these findings and the

683 resource-intensive nature of 3D imaging, we focused the subsequent SBF-SEM

684 analysis on male mice (Figure 7N).

685

686 With the 2D mitochondrial differences observed, we utilized 3D reconstruction of SBF-

687 SEM images to further investigate changes in mitochondrial structure. For each age

688 cohort, we analyzed approximately 250 mitochondria from each of the three mice

689 surveyed (Figure 7O), totaling around 750 mitochondria per age point. SBF-SEM

690 allowed for a resolution of 10 nm in the x- and y- axes, with 50 nm m intervals in the z-

691 axis, to image a total of 300 slices. Of these, 50 slices were used for analysis. These

692 slices underwent manual contour segmentation (Figure 7P) for the 3D reconstruction of

693 mitochondria (Figure 7Q). Displayed first in both age cohorts are representatives of

694 each orthoslice (Figure 7R and S), which allows for the identification of mitochondria

695 (Figures 7R' and S'). From there, we manually segmented each ortho slice in Amira to

696 reconstruct mitochondria. Additionally, for better visualization of each individually

697 colored mitochondria, the ortho slice may be removed (Figures 7R'' and S''). Once

698 modeled, the dimensions of the 3D reconstructions can be quantified accurately. We

699 found that in comparing the young and aged mice, mitochondrial size decreased

700 significantly in all metrics (Figures 7T-X). Specifically, the perimeter, which represents

701 the sum of all external distances in the mitochondria, was approximately 20% lower in

aged samples (Figure 7T; 3 months: 10,761 ± 1,560 μm SD; 2 years: 8,054 ± 450 μm

SD). Mitochondrial surface area, a metric of the outer mitochondrial membrane (OMM)

area, was decreased by nearly 50% in aged samples (Figure 7U; 3 months: 7.66 ±

0.808 μm² SD; 2 years: 4.76 ± 0.511 μm² SD). Finally, mitochondrial volume, which

represents the total of all internal pixels within the 3D mitochondrial reconstruction, was

approximately 30% lower in aged male murine liver samples (Figure 7V; 3 months: 1.01

± 0.182 μm³ SD; 2 years: 0.696 ± 0.107 μm³ SD). Losses in mitochondrial volume may

indicate a decrease in internal volume for ATP synthesis,[75] but mitochondrial roles

extend beyond their energetics, including their ability to interface with the endoplasmic

reticulum (ER) to modulate calcium homeostasis.[76] Thus, it is equally important to look

at their morphology and capacity to form contact sites with the ER.

To aid in visualizing mitochondria, we present mitochondria 3D reconstructions from

transverse (Figures 7W and X) and longitudinal (Figures 7W' and X') viewpoints. To

further verify this change across aging, we examined sphericity and the mitochondrial

complexity index (MCI),[77] which are analogous measures of mitochondrial morphology.

Sphericity (calculated as $\frac{\pi^{\frac{1}{3}}(6*Volume)^{\frac{2}{3}}}{Surface\ Area}$), exhibited an approximate 20% higher mean

value (Figure 7Y; 3 months: 0.638 ± 0.0211 SD; 2 years: 0.785 ± 0.0257 SD). MCI

(calculated as Surface Area^3 /16 π ^{2}volume^2) conversely showed a decrease in the

aged cohort (Figure 7Z; 3 months: 3.22 ± 0.240 SD; 2 years: 1.64 ± 0.236 SD).

Together, these findings validate decreased mitochondrial complexity in aged murine

liver samples, with mitochondria adopting a more spherical morphology. To better

visualize these changes, we organized mitochondria by their volume using a method

known as mito-typing, which allows for comparison across different sizes (Figure FF

and GG). In the three-month age cohort (Figure 5FF), we observed the typical

morphology of mitochondria, characterized by a compact or tubular structure. However,

we also observed some diversity in mitochondrial structure, such as elongation,

branching, and other structures that prioritize surface area over volume. In contrast, the

2-year sample showed much less heterogeneity and mitochondrial structures were

mostly presented as tubular or compact (Figure 7GG). Three-dimensional SBF-SEM

analysis revealed significant age-associated changes in hepatic mitochondrial

morphology. Compared with mitochondria from 3-month-old mice, those from 2-year-old

mice exhibited reduced mitochondrial perimeter (7AA), area (7BB), volume (7CC),

increased sphericity (7DD), and a decreased mitochondrial complexity index (Figure

7EE). Mito-otyping analysis further revealed that mitochondria from young liver tissue

exhibited more elongated and morphologically diverse structures, whereas those from

aged liver tissue appeared more spherical and less complex (Figure 7FF-GG). We

based these analyses on approximately 1,500 mitochondria from three mice per age

group. For all these metrics, while intra-sample heterogeneity is prevalent, there is

minimal inter-sample heterogeneity and no outliers.

Dietary and Metabolic Interplay with MICOS

Beyond aging, we wanted to further investigate the impact of a high-fat diet (HFD) on

mitochondrial ultrastructure. We performed TEM analysis in a 20-week-old murine

cohort subjected to a low-fat diet (LFD) and an HFD (Figures 8A-B'). Qualitatively,

examination of TEM images revealed more lipid droplets with less circularity in HFD

samples. Looking at mitochondria, although mitochondrial count, when normalized to

the μm cell area, was lower in HFD samples (Figure 8C; LFD: 0.210 ± 0.114 SD; HFD:

0.128 ± 0.0972 SD; not significant). Similar to aging, HFD samples exhibit a significantly

higher average mitochondrion area (Figure 8E; LFD: 0.570 ± 0.401 μm² SD; HFD: 0.923

± 1.12 μm² SD). When considering these two quantifications in tandem, we also

calculated the percentage of mitochondrial area related to the total cell area, which

showed no significant difference (Figure 8D; LFD: 11.8 ± 5.59 μm² SD; HFD: 11.7 ±

5.10 μm² SD). When examining the changes in mitochondrial shape with an HFD, we

observed that mitochondria had a higher circularity, indicative of potentially less

complexity, compared to those with an LFD (Figure 8F; LFD: 0.781 ± 0.149 SD; HFD:

0.897 ± 0.0748 SD). Finally, examining the cristae score, similar to aging samples, HFD

cohorts had a significantly lower cristae score (Figure 8G; LFD: 3.74 ± 0.441 SD; HFD:

2.04 ± 0.770 SD). Together, these findings show that an HFD can parallel age-related

changes, and that an HFD exhibit smaller mitochondria with aberrant cristae. In

addition, HFD induced significant metabolic alterations, including increased serum

cholesterol (Figure 8J), serum triglycerides (Figure 8K), serum free fatty acids (Figure

8L), fat mass (Figure 8M), liver weight (Figure 8N), body weight (Figure 8O), serum

insulin (Figure 8P) and OGTT values (Figure 8Q).

Discussion
766
767

768   **Genetic and Clinical Associations of MICOS Complex Genes with Liver Disease**

769   First, we wanted to demonstrate a link between liver diseases and MICOS genes. Our

770   study identified significant associations between genetically regulated expression

771   (GREX) of MICOS complex genes and liver disease phenotypes within a biobank

772 population. We observed that decreased GREX of *CHCHD3* (*MIC19*) in individuals of

773 European ancestry correlated with liver cirrhosis and transplant status. In contrast,

774 reduced *Opa1* expression in individuals of African ancestry was linked to chronic liver

775 disease and cirrhosis. These findings highlight ancestry-specific genetic susceptibility to

776 hepatic pathologies, potentially reflecting differences in genetic architecture or

777 environmental interactions. However, the limitations of EHR-based phenotyping, such

778 as incomplete data and diagnostic variability, underscore the need for replication in

779 diverse cohorts. Future studies integrating longitudinal data and multi-omics approaches

780 could elucidate how MICOS dysregulation interacts with aging and environmental

781 factors to drive disease progression.

782 Next, we demonstrated that aging is associated with reduced expression of specific

783 mitochondrial genes (*MIC19, MIC25,* and *MIC60*) and increased expression of *BIP,*

784 *XBP1, IRE1,* and *ATF4*. The reduced expression of *MIC19, MIC25,* and *MIC60* has

785 implications for mitochondrial structure and aging.[54,60] As core components of the

786 MICOS, *MIC19, MIC25*, and *MIC60* play a crucial role in maintaining cristae

787 architecture, regulating oxidative phosphorylation, mitochondrial dynamics and

788 bioenergetic function. [22,2478] Reduced mRNA expression in older individuals is

789 associated with loss of cristae integrity, which consequently reduces the surface area of

790 respiratory complexes, thereby reducing mitochondrial efficiency & ATP production, and

791 increasing susceptibility to oxidative damage from ROS.[16,54,60,109] Thus, reduced *MIC19,*

792 *MIC25* and *MIC60* likely reflects mitochondrial structural decay and reduced

793 bioenergetic capacity with age.[114,115]

794

795 In aging cells, the upregulation of *BIP, IRE1, XBP1, ATF4,* and *CHOP* reflects a

796 persistent activation of the endoplasmic reticulum (ER) stress response and the

797 integrated stress response (ISR) as part of attempts to maintain cellular proteostasis

798 under increasing proteotoxic and metabolic stress.[96,103] *BIP* (also known as *GRP78*)

799 acts as a key ER chaperone that dissociates from stress sensors upon accumulation of

800 misfolded proteins, initiating the unfolded protein response (UPR) and signaling

801 adaptive stress pathways.[96] *IRE1* activation promotes the unconventional splicing of

802 *XBP1* mRNA, resulting in the production of a potent transcription factor that enhances

803 the expression of chaperones and ER quality control machinery.[103] In contrast, *ATF4* is

804 induced downstream of PERK signaling to regulate stress-responsive genes.[96] Under

805 prolonged or unresolved stress, *CHOP* expression increases, shifting the UPR from an

806 adaptive to a pro-apoptotic signaling pathway by downregulating anti-apoptotic factors

807 and linking to mitochondrial pathways of cell death.[96,103] Collectively, elevated

808 expression of these factors in aged tissues indicates chronic ER stress and an

809 overwhelmed proteostasis network, which can disrupt ER–mitochondrial

810 communication, exacerbate mitochondrial dysfunction, and contribute to age-related

811 decline and susceptibility to disease.[82,83]

812

813 To validate and assess the importance of the MICOS genes in age-related changes, we

814 knocked down *MIC60* and *CHCHD6*. We found that their deficiency impaired Ca^{2+}

815 homeostasis, cell survival and induced oxidative stress in HepG2 cells.

816

817 **MICOS Complex Dysregulation Drives Mitochondrial Dysfunction**

Previous studies have established that the MICOS complex is critical for mitochondrial

dynamics,[78] and our group have previously investigated the effect of aging on the

MICOS complex in kidney tubular cells.[60] As reported in our study, aging in murine liver

was associated with a significant decline in the mRNA expression of key MICOS

components and the associated protein OPA1. While *Opa1* interacts with the MICOS

complex, it does not require *Opa1* to form cristae junctions, and loss of Opa*1* does not

negatively affect MICOS components.[84] This suggests that the loss of the MICOS

complex across aging occurs in an *Opa1*-independent manner. This parallels our

previous findings in skeletal muscle, the heart, and the kidney.[54,60,80]

Calcium influx in mitochondria can reflect cell viability.[97,98] Mitochondrial mediation of

calcium can impact various cell processes, including apoptosis, signaling, and ATP

production.[99] In some cases, elevated mitochondrial calcium uptake can occur

antecedent to mitochondrial swelling, which results in a pathway often leading to

apoptosis.[100] Aging-associated declines in MICOS components (*MIC60, CHCHD3,*

CHCHD6) correlated with disrupted calcium handling and elevated oxidative stress in

HepG2 cells. Knockdown of *MIC60* or *CHCHD6* impaired mitochondrial calcium

retention and increased ROS production, suggesting that MICOS integrity is vital for

redox homeostasis and apoptosis regulation.[99,101] This provides a plausible disease link

as ER calcium release promotes mitochondrial dysfunction, inducing oxidative stress

and hepatotoxicity,[102] with broader implications in MASLD.[103,104] Since Miro clusters

both interact with the MICOS complex and regulate MERCs,[105] Miro represents a

potential future mechanistic avenue through which the MICOS complex contributes to

MERC tethering.

Researchers have proposed that the mitochondrial redox state is a principal moderator

of mitochondrial function in liver disease.[106] Recent studies have continued to highlight

a link between ROS and mitochondrial dynamics.[107] Lipids can aid in stimulating the

production of ROS.[108] Additionally, many theories of aging propose that mitochondria

lose function because ROS byproducts accumulate during aging.[109] There remains

controversy about whether ROS are generated in the liver. While some studies report

that superoxides arise during liver aging,[110] other studies conversely report that the

aging human liver does not produce superoxides.[111] Future studies should explore how

ROS contributes to the observed 3D mitochondrial phenotypes, as oxidative stress may

be a regulator of the phenotypes we observed. It is noteworthy that the loss of *CHCHD6*

and *MIC60* leads to oxidative stress. This suggests that during aging, the liver undergo

a vicious cycle wherein abnormal mitochondrial structures generate more harmful

byproducts. These byproducts, in turn, worsen mitochondrial structural dysfunction,

contributing to age-related oxidative stress. These findings align with studies linking

cristae disorganization to mPTP sensitization and ATP synthase dysfunction.[102,103] The

vicious cycle of MICOS loss, oxidative stress, and metabolic dysfunction may accelerate

age-related hepatic decline, particularly under lipid-rich conditions that exacerbate ER-

mitochondrial uncoupling.[114,115]

Hepatic Metabolic Reprogramming as a Feature of Aging

Our integrated metabolomic and lipidomic profiling of aged liver tissue reveals a

profound and coordinated rewiring of hepatic metabolism, centering on mitochondrial

864 dysfunction as a key driver of age-related physiological decline. The clear separation of

865 young and aged metabolic signatures, along with the accumulation of vitamin A

866 metabolites, suggests altered nutrient handling and redox signaling in aging

867 hepatocytes.[17,18,62] Concomitant reductions in tricarboxylic acid cycle intermediates,

868 including succinate and malate, indicate impaired mitochondrial oxidative metabolism

869 and/or increased cataplerotic flux, reflecting diminished bioenergetic capacity with

870 age.[3,5,109,110] In parallel, significant depletion of nucleotide monophosphates and NAD^+-

871 related metabolites highlights a decline in mitochondrial-dependent biosynthetic

872 pathways and redox homeostasis, processes essential for energy production, cellular

873 repair, and metabolic flexibility.[49,63,64] Given the central role of mitochondria in

874 coordinating these pathways, the observed metabolite changes collectively indicate

875 compromised mitochondrial function as a key driver of altered energy metabolism,

876 reduced anabolic capacity, and disrupted cellular homeostasis in the aging liver, thereby

877 reinforcing mitochondrial dysfunction as a fundamental contributor to age-associated

878 metabolic decline.[8,115]

879

880   Building upon the profound metabolic dysregulation observed, our findings

881 demonstrate that age-related mitochondrial impairment in the liver has direct

882 pathological consequences, culminating in overt lipid accumulation and a progression

883 towards metabolic dysfunction-associated steatotic liver disease (MASLD).[20,66,116] This

884 transition is evident in both humans and mice, where advanced age is correlated with a

885 significantly increased hepatic lipid content, exceeding established steatosis thresholds

886 in human cohorts.[17,20,42] In murine models, this manifests as a dramatic increase in lipid

droplets and triglyceride concentration, accompanied by hallmark features of disease

progression, such as a reduced liver-to-body weight ratio and decreased mtDNA

content.[44,66,124] Crucially, lipidomic profiling reveals that aging is not characterized by a

uniform increase in lipids, but rather by a specific remodeling of the hepatic lipid

landscape, including significant alterations in triglycerides, ceramides, and

acylcarnitines.[68,69,70,71] This specific signature links the compromised energy metabolism

and NAD^+ depletion which impair fatty acid oxidation, to a lipogenic shift and the

accumulation of lipid species known to promote insulin resistance and cellular

stress.[66,69,102,108] Thus, age-related mitochondrial dysfunction appears to be a central

driver that shifts the liver from a state of metabolic flexibility to one of lipid-laden

inflexibility, significantly increasing the risk of serious hepatic and systemic metabolic

disease. [114,115,116]

Age-Related Mitochondrial Structural Remodeling in the Liver

Using SBF-SEM, we demonstrated that aging reduces mitochondrial volume, perimeter,

and complexity in the murine liver, paralleling 2D TEM findings of cristae disorganization

and fragmentation. These structural declines are associated with impaired metabolic

and lipidomic profiles, indicating compromised bioenergetic efficiency and oxidative

capacity. A limitation of our study is the lack of direct measurement of ATP synthesis

rates, which would have more directly linked structure to function; this is an important

area for future investigation. Notably, aged mitochondria exhibited spherical

morphologies with reduced contact site potential, which may disrupt mitochondria-ER

interactions critical for lipid homeostasis and calcium signaling.[82-84] The structure of

mitochondria remains relevant in aging as structural decline may decrease

mitochondrial function, thereby impairing certain liver roles. While megamitochondria, a

hallmark of advanced liver disease,[85-87] were absent in our aged cohort, this may reflect

a pre-diseased state where compensatory mechanisms, such as upregulated *Drp1*,

remain functional.[86,88,89] The observed structural changes likely precede overt

pathology, underscoring the role of mitochondrial integrity in maintaining hepatic

resilience during aging.

Beyond mitochondrial 3D structure, past studies have shown that fatty acid

accumulation promote endoplasmic reticulum stress and liver injury in rodent models.[90]

Due to the increase of lipid droplets in both aging and high fat diet treatment in TEM

images, there may be an increase in mitochondria-lipid droplet contact sites

(MLDCs).[76,91] Beyond this, lipid droplet-endoplasmic reticulum contacts are also

understood to help form lipid droplets and perform metabolism.[65] Especially relevant is

that MLDCs serve as a place for fatty acid homeostasis.[91] Additionally, mitochondria

endoplasmic reticulum contact sites (MERCs) importantly play a role in lipid

homeostasis and synthesis.[92] MERCs are understood to regulate ER stress,[93] so

increases in fatty acids, triggered by lipid droplet formation, can lead to increased

MERC formation in the aged model. Conversely, we qualitatively observed a decrease

in structures consistent with wrappER, a mitochondria-ER contact known to maintain

lipid flux through a tri-organelle contact site that involves peroxisomes.[94] These

wrappER sites, which contain sites of adhesion, regulate very-low-density lipoproteins.[95]

The age-related loss of mitochondrial complexity may impair the ability and relative

surface area mitochondria in murine liver samples must form contact sites, including

933 MERCs, thus interfering with functions including lipid homeostasis. Indeed, past reviews

934 have suggested targeting MERCs in MASLD due to the role of contact sites in glucose

935 and lipid metabolism.[96] However, in the future, a more rigorous analysis of age-related

936 changes in contact sites within liver tissue is necessary.

937

938 **Dietary and Metabolic Interplay with MICOS in Aging**

939 Our investigation into the impact of a high-fat diet reveals that nutritional stress

940 recapitulates and accelerates the same fundamental mitochondrial pathology observed

941 in aging, directly linking dietary patterns to premature cellular aging and metabolic

942 dysfunction. The HFD-induced mitochondrial remodeling characterized by enlarged yet

943 simplified organelles with significantly disordered cristae architecture mirrors the

944 ultrastructural decay seen in aged livers, suggesting a common endpoint of bioenergetic

945 compromise. This loss of cristae integrity is particularly critical, as it directly impairs the

946 capacity for oxidative phosphorylation. While the total mitochondrial area per cell was

947 maintained, the shift towards fewer, larger, and more circular organelles points to an

948 imbalance in mitochondrial dynamics, likely hindering their ability to network, traffic, and

949 undergo efficient quality control. These structural deficits manifest in severe systemic

950 metabolic alterations, including elevated serum lipids, insulin resistance, and hepatic

951 steatosis. Therefore, HFD does not merely alter metabolism passively but actively

952 drives a pathological transformation of mitochondrial structure that underpins the

953 development of metabolic disease, positioning poor dietary quality as a potent

954 accelerator of the mitochondrial dysfunction typically associated with advancing age.

955

We know that diet induced obesity (DIO) are linked with many liver complications. For

example, oxidative stress, inflammation, and lipogenesis are factors that may be linked

to DIO and exacerbate the onset of MASLD.[116] Additionally, DIO is associated with

metabolic dysregulation of the liver.[117] Elevated retinol and ceramide levels in aged liver

paralleled HFD-associated lipotoxicity, implicating disrupted β-oxidation, impaired

insulin sensitivity and peroxisomal signaling.[17,118] The enhanced supply of vitamin A and

high-fat consumption in diet-induced obese mice is linked to the production of

bisretinoid.[119] Interestingly, obesity within hepatocytes has also been shown to increase

MAMs (i.e., MERC-isolated biochemical fractions), which confers mitochondrial

dysfunction.[120,121] Thus, HFDs can play a role in metabolic defects in the liver tissue,

although this requires further investigation in the future.

An HFD can lead to pathological alterations and damage to the ultrastructure of the

mitochondria, as well as downregulation of MERCs regulators, *MFN2* and *OPA1*[122]

which parallels the TEM ultrastructural changes we observed. Additionally, studies have

shown that *Mic19/Chchd3* is required for normal liver function. In the liver, deletion of

Mic19/Chchd3 reduces ER-mitochondrial contacts, disrupts mitochondrial lipid

metabolism, disorganizes mitochondrial cristae, and triggers an unfolded protein stress

response in mouse hepatocytes, resulting in impairments of liver mitochondrial fatty acid

β-oxidation and lipid metabolism.[123] Thus, while MICOS components like *MIC19* and

CHCHD3 are essential for ER-mitochondrial tethering and lipid metabolism; their age-

related decline may impair compensatory responses to dietary stress. Notably, HFD and

aging both reduce mitochondrial plasticity, exacerbating steatosis and insulin

resistance, key drivers of MASLD progression.[124,125]

Conclusions

Our integrative analysis reveals that age-related disruption of the MICOS complex

contributes to mitochondrial structural and functional decline in the liver, promoting

metabolic dysfunction and increasing disease susceptibility. By linking 3D ultrastructural

changes, genetic associations, and cellular mechanisms, we establish MICOS as a

critical node in the process of hepatic aging. A limitation of this study is the primary use

of a male murine model for deep ultrastructural and functional validation; future work

should explore sex-specific differences in these processes. Deleting the MICOS

complex in HepG2 cells impairs calcium uptake and increases oxidative stress,

underscoring the importance of this complex. A high-fat diet exacerbates lipid synthesis,

catabolism, and mitochondrial changes in aging livers, increasing susceptibility to liver

disease.[17,116] This study demonstrates that impaired mitochondrial structure, calcium

influx, ROS production, metabolic alterations, and MICOS dysfunctions contribute to

liver diseases. These findings highlight potential therapeutic targets, such as enhancing

cristae integrity or modulating MERCs, to mitigate age-related liver pathologies. Future

studies examining tissue-specific MICOS dynamics across diverse populations and

dietary conditions will further elucidate its role in aging and metabolic diseases.

Abbreviations
TEM, transmission electron microscopy; SBF-SEM, serial block-face scanning electron
microscopy; MASLD, metabolic dysfunction-associated steatotic liver disease; MICOS,
mitochondria contact site and cristae organizing system; GREX, genetically regulated

gene expression; MRI, magnetic resonance imaging; TCA, tricarboxylic acid; NAD^+,
nicotinamide adenine nucleotide; TGO, triglyceride oligomer; TG, triglycerides; Cer,
ceramide; CAR, acylcarnitine; ER, endoplasmic reticulum; MCI, mitochondrial
complexity index; ROS, reactive oxygen species; MLDC, mitochondria-lipid contact site;
HFD, high-fat diet; DIO, diet induced obesity; MAM, mitochondria-associated
membrane; MERC, mitochondria endoplasmic reticulum contact site; OMM, Outer
Mitochondrial Membrane.

Financial support
The Synthetic Derivative and BioVU projects at VUMC are supported by numerous
sources: including the NIH funded Shared Instrumentation Grant S10OD017985 and
S10RR025141; CTSA grants UL1TR002243, UL1TR000445, and UL1RR024975 from
the National Center for Advancing Translational Sciences. Its contents are solely the
responsibility of the authors and do not necessarily represent official views of the
National Center for Advancing Translational Sciences or the National Institutes of
Health. Genomic data are also supported by investigator-led projects that include
U01HG004798, R01NS032830, RC2GM092618, P50GM115305, U01HG006378,
U19HL065962, R01HD074711; and additional funding sources listed at
https://victr.vumc.org/biovu-funding/. Other funding sources include 2D43TW009744
and R21TW012635 (SKM and AK), **T**he Howard Hughes Medical Institute Hanna H.
Gray Fellows Program Faculty Phase (Grant# GT15655 awarded to MRM) and the
Burroughs Welcome Fund PDEP Transition to Faculty (Grant# 1022604 awarded to
MRM).

This project was funded by the National Institute of Health (NIH) NIDDK T-32, number
DK007563 entitled Multidisciplinary Training in Molecular Endocrinology to Z.V.;
National Institute of Health (NIH) NIDDK T-32, number DK007563 entitled
Multidisciplinary Training in Molecular Endocrinology to A.C.; NSF MCB #2011577I to
S.A.M.; The UNCF/Bristol-Myers Squibb E.E. Just Faculty Fund, Career Award at the
Scientific Interface (CASI Award) from Burroughs Welcome Fund (BWF) ID #
1021868.01, BWF Ad-hoc Award, NIH Small Research Pilot Subaward to
5R25HL106365-12 from the National Institutes of Health PRIDE Program, DK020593,
Vanderbilt Diabetes and Research Training Center for DRTC Alzheimer's Disease Pilot
& Feasibility Program. CZI Science Diversity Leadership grant number 2022- 253529
from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community
Foundation to A.H.J. and O.A.A.; and National Institutes of Health grant HD090061 and
the Department of Veterans Affairs Office of Research Award I01 BX005352 to J.G.
Howard Hughes Medical Institute Hanna H. Gray Fellows Program Faculty Phase
(Grant# GT15655 awarded to M.R.M); and Burroughs Wellcome Fund PDEP Transition
to Faculty (Grant# 1022604 awarded to M.R.M). National Institutes of Health Grants:
R21DK119879 (to C.R.W.) and R01DK-133698 (to C.R.W.), American Heart
Association Grants 16SDG27080009 (to C.R.W.) and 24IVPHA1297559
https://doi.org/10.58275/AHA.24IVPHA1297559.pc.gr.193866 (S.K.M) and by an
American Society of Nephrology KidneyCure Transition to Independence Grant (to
C.R.W.). NIH Grants R01HL147818, R03HL155041, and R01HL144941 (A. Kirabo).
NIH Grant R00DK120876 (D.T.), Harold S. Geneen Charitable Trust Awards Program
(D.T.), Alzheimer's Association AARG-NTF-23-1144888 (D.T.). NIH Grant

R00AG065445 (P.J.), Alzheimer's Association 24AARG-D-1191292 (P.J.), Wake ADRC REC and Development grant P30AG072947 (P.J.). American Heart Association Grant 23POST1020344 (A.K.). American Heart Association Grant 23CDA1053072 (M. S.). NIH K01AG062757 to (M.T.S.) ANRF (Anusandhan National Research Foundation), ANRF/ECRG/2024/001042/LS, ANRF/IRG/2024/001777/LS. IISER Tirupati, NFSG (P.K). The BioVU project at VUMC is supported by numerous sources: including the NIH funded Shared Instrumentation Grant S10OD017985 and S10RR025141; CTSA grants UL1TR002243, UL1TR000445, and UL1RR024975 from the National Center for Advancing Translational Sciences. Its contents are solely the responsibility of the authors and do not necessarily represent official views of the National Center for Advancing Translational Sciences. Genomic data are also supported by investigator-led projects that include U01HG004798, R01NS032830, RC2GM092618, P50GM115305, U01HG006378, U19HL065962, R01HD074711; and additional funding sources listed at https://victr.vumc.org/biovu-funding/.23CDA1053072 (M.S.). Its contents are solely the responsibility of the authors and do not necessarily represent the official view of the NIH. The funders had no role in study design, data collection, and analysis, decision to publish, or preparation of the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

Authors' contributions

Sepiso K. Masenga, Alexandria Murphy, Prasanna Venkhatesh contributed equally to this work. Zer Vue, Benjamin Rodriguez, Ashlesha Kadam, Andrea G. Marshall, Estevão Scudese, Brenita Jenkins, Amber Crabtree, Praveena Prasad, Edgar Garza-Lopez, Han Le, Ky'Era V. Actkins, Elma Zaganjor, Nelson Wandira, Jeremiah Afolabi, Prasanna Katti, Chantell Evans, Young Do Koo, Dhanendra Tomar, Mark A. Phillips, David Hubert, Chandravanu Dash, Pooja Jadiya, Olujimi A. Ajijola, Magdalene Ameka, Okwute M. Ochayi, Eric Wang, Quinton Smith, Ronald McMillan, Annet Kirabo, André Kinder, Tyne W. Miller-Fleming, Bret Mobley, Julia D. Berry, Nathan Winn, Vernat Exil, Anita M. Quintana, Kit Neikirk, Jenny Schafer, Sean Schaffer, Oleg Kovtun, Mohd Mabood Khan, Calixto Pablo Hernandez Perez, Margaret Mungai, Melanie R. McReynolds, Antentor Hinton, Jr contributed to the conception, design, data acquisition, analysis, and interpretation of data. Melanie R. McReynolds and Antentor Hinton, Jr. conceived and supervised this study. All authors were involved in drafting and critically revising the manuscript for important intellectual content. All authors approved the final version of the manuscript.

Data availability statement

All data generated or analyzed during this study are included in this published article and its Supplementary information files. Additional data can be requested from the corresponding author.

Acknowledgment

We would like to acknowledge the Huck Institutes' Metabolomics Core Facility (RRID:SCR_023864) for use of the OE 240 LCMS and Sergei Koshkin for helpful

discussions on sample preparation and analysis. We would also like to acknowledge the Huck Institutes' Metabolomics Core Facility (RRID:SCR_023864) for use of the OE 240 LCMS and Drs. Imhoi Koo, Ashley Shay, and Sergei Koshkin for helpful discussions on sample preparation and analysis. We would also like to thank UCLA investigators for gifting us old and young human liver samples.

References

[1] Chan DC. Fusion and fission: interlinked processes critical for mitochondrial health. Annual Review of Genetics 2012;46:265–287.

[2] Li M, Wang L, Wang Y, et al. Mitochondrial fusion via OPA1 and MFN1 supports liver tumor cell metabolism and growth. Cells 2020;9(1):121.

[3] Glancy B, Kim Y, Katti P, et al. The functional impact of mitochondrial structure across subcellular scales. Frontier Physiology 2020;11. https://doi.org/10.3389/fphys.2020.541040

[4] McBride HM, Neuspiel M, Wasiak S. Mitochondria: more than just a powerhouse. Current Biology 2006;16(14):R551-560.

[5] Picard M, Taivassalo T, Gouspillou G, et al. Mitochondria: isolation, structure and function. Journal of Physiology 2011;589(18):4413–4421.

[6] Seo AY, Joseph AM, Dutta D, et al. New insights into the role of mitochondria in aging: mitochondrial dynamics and more. Journal of Cell Science 2010;123(15):2533–2542.

[7] Bratic A, Larsson N-G. The role of mitochondria in aging. Journal of Clinical Investigation 2013;123(3):951–957.

[8] Haas RH. Mitochondrial dysfunction in aging and diseases of aging. Biology 2019;8(2):48.

[9] Gautier MK, Ginsberg SD. A method for quantification of vesicular compartments within cells using 3D reconstructed confocal z-stacks: Comparison of ImageJ and Imaris to count early endosomes within basal forebrain cholinergic neurons. Journal of Neuroscience Methods 2021;350:109038.

[10] Garza-Lopez E, Vue Z, Katti P, et al. Protocols for generating surfaces and measuring 3D organelle morphology Using Amira. Cells 2022;11(1):65.

[11] Courson JA, Landry PT, Do T, et al. Serial block-face scanning electron microscopy

(SBF-SEM) of biological tissue samples. Journal of Visualized Experiments

2021;(169):10.3791/62045.

[12] Marshall AG, Neirkirk K, Stephens DC, et al. Serial block face-scanning electron

microscopy as a burgeoning technology. Advanced Biology 2023;e2300139.

[13] Russell MRG, Lerner TR, Burden JJ, et al. 3D correlative light and electron

microscopy of cultured cells using serial block-face scanning electron microscopy.

Journal of Cell Science 2017;130(1):278–291.

[14] Lam J, Katti P, Biete M, et al. A universal approach to analyzing transmission

electron microscopy with ImageJ. Cells 2021;10(9):2177.

[15] Collins HE, Kane MS, Litovsky SH, et al. Mitochondrial morphology and mitophagy

in heart diseases: Qualitative and quantitative analyses using transmission electron

microscopy. Frontiers in Aging 2021;2.

https://www.frontiersin.org/articles/10.3389/fragi.2021.670267. Accessed September 8,

2022.

[16] Cogliati S, Enriquez JA, Scorrano L. Mitochondrial cristae: Where beauty meets

functionality. Trends Biochemical Sciences 2016;41(3):261–273.

[17] Gong Z, Tas E, Yakar S, et al. Hepatic lipid metabolism and non-alcoholic fatty liver

disease in aging. Molecular and Cell Endocrinology 2017;455:115–130.

[18] Chung KW. Advances in understanding of the role of lipid metabolism in aging.

Cells 2021;10(4):880.

[19] Abrisch RG, Gumbin SC, Wisniewski BT, et al. Fission and fusion machineries

converge at ER contact sites to regulate mitochondrial morphology. Journal of Cell

Biology 2020;219(4):e201911122.

[20] Kim H, Kisseleva T, Brenner DA. Aging and liver disease. Current Opinion in

Gastroenterology 2015;31(3):184–191.

[21] Chen H, Detmer SA, Ewald AJ, et al. Mitofusins Mfn1 and Mfn2 coordinately

regulate mitochondrial fusion and are essential for embryonic development. Journal of

Cell Biology 2003;160(2):189–200.

[22] Hu C, Shu L, Huang X, et al. OPA1 and MICOS regulate mitochondrial crista

dynamics and formation. Cell Death & Disease 2020;11(10):1–17.

[23] Takeichi Y, Miyazawa T, Sakamoto S, et al. Non-alcoholic fatty liver disease in
mice with hepatocyte-specific deletion of mitochondrial fission factor. Diabetologia
2021;64:2092–2107.

[24] Kozjak-Pavlovic V. The MICOS complex of human mitochondria. Cell and Tissue
Research 2017;367(1):83–93.

[25] Li Z, Shen W, Wu G, et al. The role of SAMM50 in non-alcoholic fatty liver disease:
from genetics to mechanisms. FEBS Open Bio 2021;11(7):1893–1906.

[26] Lionello S, Marzaro G, Martinvalet D. SAM50, a side door to the mitochondria: The
case of cytotoxic proteases. Pharmacological Research 2020;160:105196.

[27] Shao B, Killion M, Oliver A, et al. Ablation of Sam50 is associated with
fragmentation and alterations in metabolism in murine and human myotubes. BioRxiv
The Preprint Server for Biology 2023;2023.05.20.541602.

[28] Shao B, Killion M, Oliver A, et al. Ablation of Sam50 is associated with
fragmentation and alterations in metabolism in murine and human myotubes. Journal of
Cellular Physiology 2024;n/a(n/a). https://doi.org/10.1002/jcp.31293.

[29] Roden DM, Pulley JM, Basford MA, et al. Development of a large-scale de-
identified DNA biobank to enable personalized medicine. Clinical Pharmacology &
Therapeutics 2008;84(3):362–369.

[30] Danciu I, Cowan JD, Basford M, et al. Secondary use of clinical data: the Vanderbilt
approach. Journal of Biomedical Informatics 2014;52:28–35.

[31] Dennis JK, Sealock JM, Straub P, et al. Clinical laboratory test-wide association
scan of polygenic scores identifies biomarkers of complex disease. Genome Medicine
2021;13(1):6.

[32] Price AL, Patterson NJ, Plenge RM, et al. Principal components analysis corrects
for stratification in genome-wide association studies. Nature Genetics 2006;38(8):904–
909.

[33] 1000 Genomes Project Consortium. A global reference for human genetic variation.
Nature 2015;526(7571):68–74.

[34] Das S, Forer L, Schönherr S, et al. Next-generation genotype imputation service
and methods. Nature Genetics 2016;48(10):1284–1287.

[35] the Halotype Reference Consortium. A reference panel of 64,976 haplotypes for

genotype imputation. Nature Genetics 2016;48(10):1279–1283.

[36] GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across

human tissues. Science 2020;369(6509):1318–1330.

[37] Gamazon ER, Wheeler HE, Shah KP, et al. A gene-based association method for

mapping traits using reference transcriptome data. Nature Genetics 2015;47(9):1091–

1098.

[38] Hu Y, Li M, Lu Q, et al. A statistical framework for cross-tissue transcriptome-wide

association analysis. Nature Genetics 2019;51(3):568–576.

[39] Zhou D, Jiang Y, Zhong X, et al. A unified framework for joint-tissue transcriptome-

wide association and Mendelian randomization analysis. Nature Genetics

2020;52(11):1239–1246.

[40] Carroll RJ, Bastarache L, Denny JC. R PheWAS: data analysis and plotting tools

for phenome-wide association studies in the R environment. Bioinformatics

2014;30(16):2375–2376.

[41] Wu P, Gifford A, Meng X, et al. Mapping ICD-10 and ICD-10-CM codes to

Phecodes: Workflow development and initial evaluation. JMIR Medical Informatics

2019;7(4):e14325.

[42] Tang A, Tan J, Sun M, et al. Nonalcoholic fatty liver disease: MR imaging of iver

proton density fat fraction to assess hepatic steatosis. Radiology 2013;267(2):422–431.

[43] Hinton A, Katti P, Mungai M, et al. ATF4 dependent increase in mitochondrial-

endoplasmic reticulum tethering following OPA1 deletion in skeletal muscle. Journal of

Cellular Physiology [published online ahead of print: 2024].

https://doi.org/10.1101/2022.09.12.507669.

[44] Stahl EC, Delgado ER, Alencastro F, et al. Inflammation and ectopic fat deposition

in the aging murine liver is influenced by CCR2. American Journal of Pathology

2020;190(2):372–387.

[45] Pereira RO, Tadinada SM, Zasadny FM, et al. OPA 1 deficiency promotes

secretion of FGF 21 from muscle that prevents obesity and insulin resistance. EMBO

Journal 2017;36(14):2126–2145.

1210 [46] Tam J, Vemuri VK, Liu J, et al. Peripheral CB1 cannabinoid receptor blockade

1211 improves cardiometabolic risk in mouse models of obesity. Journal of Clinical

1212 Investigation 2010;120(8):2953–2966.

1213 [47] Hinton A, Katti P, Christensen TA, et al. A comprehensive approach to sample

1214 preparation for electron microscopy and the assessment of mitochondrial morphology in

1215 tissue and cultured cells. Advanced Biology 2023;e2200202.

1216 [48] Neikirk K, Vue Z, Katti P, et al. Systematic transmission electron microscopy-based

1217 identification and 3D reconstruction of cellular degradation machinery. Advanced

1218 Biology 2023;7(6):2200221.

1219 [49] Lu W, Wang L, Chen L, et al. Extraction and quantitation of nicotinamide adenine

1220 dinucleotide redox cofactors. Antioxidants & Redox Signaling 2018;28(3):167–179.

1221 [50] Wang L, Xing X, Chen L, et al. Peak annotation and verification engine for

1222 untargeted LC–MS metabolomics. Analytical Chemistry 2019;91(3):1838–1846.

1223 [51] Adusumilli R, Mallick P. Data conversion with ProteoWizard msConvert. In: Comai

1224 L, Katz JE, Mallick P, eds. Proteomics: Methods and Protocols New York, NY: Springer;

1225 2017:339–368.

1226 [52] Boudina S, Sena S, Theobald H, et al. Mitochondrial energetics in the heart in

1227 obesity-related diabetes: direct evidence for increased uncoupled respiration and

1228 activation of uncoupling proteins. Diabetes 2007;56(10):2457–2466.

1229 [53] Kozera B, Rapacz M. Reference genes in real-time PCR. Journal of Applied

1230 Genetics, 54(4), 391–406. https://doi.org/10.1007/s13353-013-0173-x

1231 [54] Vue Z, Garza-Lopez E, Neirkirk K, et al. 3D reconstruction of murine mitochondria

1232 reveals changes in structure during aging linked to the MICOS complex. Aging Cell

1233 2023;22(12):e14009.

1234 [55] Olenych SG, Claxton NS, Ottenberg GK, et al. The fluorescent protein color palette.

1235 Current Protocols in Cell Biology 2007;Chapter 21:Unit 21.5.

1236 [56] Denny JC, Ritchie MD, Basford MA, et al. PheWAS: demonstrating the feasibility of

1237 a phenome-wide scan to discover gene-disease associations. Bioinformatics

1238 2010;26(9):1205–1210.

[57] Roden DM, Pulley JM, Basford MA, et al. Development of a large-scale de-

identified DNA biobank to enable personalized medicine. Clinical Pharmacology &

Therapeutics 2008;84(3):362–369.

[58] Danciu I, Cowan JD, Basford MA, et al. Secondary use of clinical data: the

Vanderbilt approach. Journal of Biomedical Informatics 2014;52:28–35.

[59] Barbeira AN, Pividori M, Zheng J, et al. Integrating predicted transcriptome from

multiple tissues improves association detection. PLOS Genetics 2019;15(1):e1007889.

[60] Vue Z, Prasad P, Le H, et al. The MICOS complex regulates mitochondrial

structure and oxidative stress during age-dependent structural deficits in the kidney

[preprint]. 2024;2024.06.09.598108.

[61] Tomar D, Dong Z, Shanmughapriya S, et al. MCUR1 is a scaffold factor for the

MCU complex function and promotes mitochondrial bioenergetics. Cell Reports

2016;15(8):1673–1685.

[62] Surmacki JM, Abramczyk H. Confocal Raman imaging reveals the impact of

retinoids on human breast cancer via monitoring the redox status of cytochrome c.

Scientific Reports 2023;13(1):15049.

[63] McReynolds MR, Chellappa K, Chiles E, et al. NAD+ flux is maintained in aged

mice despite lower tissue concentrations. Cell Systems 2021;12(12):1160-1172.e4.

[64] Goodman RP, Calvo SE, Mootha VK. Spatiotemporal compartmentalization of

hepatic NADH and NADPH metabolism. Journal of Biological Chemistry

2018;293(20):7508–7516.

[65] Ma X, Qian H, Chen A, et al. Perspectives on mitochondria–ER and mitochondria–

lipid droplet contact in hepatocytes and hepatic lipid metabolism. Cells 2021;10(9):2273.

[66] Simões ICM, Fontes A, Pinton P, et al. Mitochondria in non-alcoholic fatty liver

disease. International Journal of Biochemistry & Cell Biology 2018;95:93–99.

[67] Chiang JYL, Ferrell JM. Bile acid metabolism in liver pathobiology. Gene

Expression 2018;18(2):71–87.

[68] Tomizawa M, Kawanabe Y, Shinozaki F, et al. Triglyceride is strongly associated

with nonalcoholic fatty liver disease among markers of hyperlipidemia and diabetes.

Biomedical Reports 2014;2(5):633–636.

[69] Semova I, Biddinger SB. Triglycerides in nonalcoholic fatty liver disease: Guilty until

proven innocent. Trends in Pharmacological Sciences 2021;42(3):183–190.

[70] Ueda N. Ceramide-induced apoptosis in renal tubular cells: a role of mitochondria

and sphingosine-1-phoshate. International Journal of Molecular Sciences

2015;16(3):5076–5124.

[71] Fugio LB, Coeli-Lacchini FB, Leopoldino AM. Sphingolipids and mitochondrial

dynamic. Cells 2020;9(3):581.

[72] Parra V, Eisner V, Chiong M, et al. Changes in mitochondrial dynamics during

ceramide-induced cardiomyocyte early apoptosis. Cardiovascular Research

2008;77(2):387–397.

[73] Longo N, Frigeni M, Pasquali M. Carnitine transport and fatty acid oxidation.

Biochimica et Biophysica Acta 2016;1863(10):2422–2435.

[74] Yanai S, Endo S. Functional aging in male C57BL/6J mice across the life-span: a

systematic behavioral analysis of motor, emotional, and memory function to define an

aging phenotype. Frontiers in Aging Neuroscience 2021;13.

https://www.frontiersin.org/articles/10.3389/fnagi.2021.697621. Accessed February 8,

2023.

[75] Robertson GL, Riffle S, Patel M, et al. DRP1-mediated mitochondrial fission is

essential to maintain cristae morphology and bioenergetics [preprint].

2022;2021.12.31.474637.

[76] Giacomello M, Pellegrini L. The coming of age of the mitochondria–ER contact: a

matter of thickness. Cell Death & Differentiation 2016;23(9):1417–1427.

[77] Vincent AE, White K, Davey T, et al. Quantitative 3D mapping of the human

skeletal muscle mitochondrial network. Cell Reports 2019;26(4):996-1009.e4.

[78] Li H, Ruan Y, Zhang K, et al. Mic60/Mitofilin determines MICOS assembly essential

for mitochondrial dynamics and mtDNA nucleoid organization. Cell Death &

Differentiation 2016;23(3):380–392.

[79] Tezze C, Romanello V, Desbats MA, et al. Age-associated loss of OPA1 in muscle

impacts muscle mass, metabolic homeostasis, systemic inflammation, and epithelial

senescence. Cell Metabolism 2017;25(6):1374-1389.e6.

[80] Vue Z, Neirkirk K, Vang L, et al. Three-dimensional mitochondria reconstructions of

murine cardiac muscle changes in size across aging. American Journal of Physiology-

Heart and Circulatory Physiology [published online ahead of print: August 25, 2023].

https://doi.org/10.1152/ajpheart.00202.2023.

[81] De Nicolo B, Cataldi-Stagetti E, Diquigiovanni C, et al. Calcium and reactive oxygen

species signaling interplays in cardiac physiology and pathologies. Antioxidants

2023;12(2):353.

[82] Chen C, Dong X, Zhang W, et al. Dialogue between mitochondria and endoplasmic

reticulum-potential therapeutic targets for age-related cardiovascular diseases. Frontiers

in Pharmacology 2024;15:1389202.

[83] Jenkins BC, Neirkirk K, Katti P, et al. Mitochondria in disease: changes in shapes

and dynamics. Trends in Biochemical Sciences 2024;49(4):346–360.

[84] Li X, Zhao X, Qin Z, et al. Regulation of calcium homeostasis in endoplasmic

reticulum–mitochondria crosstalk: implications for skeletal muscle atrophy. Cell

Communication and Signaling 2025;23(1):17.

[85] Yamada T, Murata D, Kleiner DE, et al. Prevention and regression of

megamitochondria and steatosis by blocking mitochondrial fusion in the liver. iScience

2022;25(4):103996.

[86] Palma E, Ma X, Riva A, et al. Dynamin-1-like protein inhibition drives

megamitochondria formation as an adaptive response in alcohol-induced hepatotoxicity.

American Journal of Pathology 2019;189(3):580–589.

[87] Tandler B, Hoppel CL. Giant Mitochondria (Megamitochondria). In: Lennarz WJ,

Lane MD, eds. *Encyclopedia of Biological Chemistry (Second Edition)*. Waltham:

Academic Press; 2013:375–376.

[88] Palma E, Riva A, Mudan S, et al. Perturbation of mitochondrial dynamics in

Alcoholic Liver Disease. Journal of World Mitochondria Society 2016;2.

https://doi.org/10.18143/JWMS_V2I2_1948.

[89] Palma E, Riva A, Moreno C, et al. Perturbations in mitochondrial dynamics are

closely involved in the progression of Alcoholic Liver Disease. Alcoholism: Clinical and

Experimental Research [published online ahead of print: 2020].

https://doi.org/10.1111/acer.14299.

[90] Wang D, Wei Y, Pagliassotti MJ. Saturated fatty acids promote endoplasmic

reticulum stress and liver injury in rats with hepatic steatosis. Endocrinology

2006;147(2):943–951.

[91] Renne MF, Hariri H. Lipid droplet-organelle contact sites as hubs for fatty acid

metabolism, trafficking, and metabolic channeling. Frontiers in Cell and Developmental

Biology 2021;9. https://www.frontiersin.org/articles/10.3389/fcell.2021.726261.

Accessed February 12, 2023.

[92] Kim H, Lee S, Jun Y, et al. Structural basis for mitoguardin-2 mediated lipid

transport at ER-mitochondrial membrane contact sites. Nature Communications

2022;13(1):3702.

[93] Marchi S, Patergnani S, Pinton P. The endoplasmic reticulum–mitochondria

connection: One touch, multiple functions. Biochimica et Biophysica Acta BBA -

Bioenergetics 2014;1837(4):461–469.

[94] Ilacqua N, Anastasia I, Raimondi A, et al. A three-organelle complex made by

wrappER contacts with peroxisomes and mitochondria responds to liver lipid flux

changes. Journal of Cell Science 2021;135(5):jcs259091.

[95] Anastasia I, Ilacqua N, Raimondi A, et al. Mitochondria-rough-ER contacts in the

liver regulate systemic lipid homeostasis. Cell Reports 2021;34(11):108873.

[96] Wang J, He W, Tsai PJ, et al. Mutual interaction between endoplasmic reticulum

and mitochondria in nonalcoholic fatty liver disease. Lipids in Health and Disease

2020;19(1):72.

[97] Duchen MR, Szabadkai G. Roles of mitochondria in human disease. Essays in

Biochemistry 2010;47:115–137.

[98] Alonso MT, Villalobos C, Chamero P, et al. Calcium microdomains in mitochondria

and nucleus. Cell Calcium 2006;40(5–6):513–525.

[99] Bertero E, Maack C. Calcium signaling and reactive oxygen species in

mitochondria. Circulation Research 2018;122(10):1460–1478.

[100] Hoek JB, Farber JL, Thomas AP, et al. Calcium ion-dependent signalling and

mitochondrial dysfunction: mitochondrial calcium uptake during hormonal stimulation in

intact liver cells and its implication for the mitochondrial permeability transition.

Biochimica et Biophysica Acta BBA - Molecular Basis of Disease 1995;1271(1):93–102.

[101] Csordás G, Weaver D, Hajnóczky G. Endoplasmic reticulum–mitochondrial

contactology: Structure and signaling functions. Trends in Cell Biology 2018;28(7):523–

540.

[102] Egnatchik RA, Leamy AK, Jacobson DA, et al. ER calcium release promotes

mitochondrial dysfunction and hepatic cell lipotoxicity in response to palmitate overload.

Molecular Metabolism 2014;3(5):544–553.

[103] Kim YS, Kim SG. Endoplasmic reticulum stress and autophagy dysregulation in

alcoholic and non-alcoholic liver diseases. Clinical and Molecular Hepatology

2020;26(4):715–727.

[104] Amaya MJ, Nathanson MH. Calcium Signaling in the liver. Comprehensive

Physiology 2013;3(1):515–539.

[105] Modi S, López-Doménech G, Halff EF, et al. Miro clusters regulate ER-

mitochondria contact sites and link cristae organization to the mitochondrial transport

machinery. Nature Communications 2019;10(1):4399.

[106] Cichoż-Lach H, Michalak A. Oxidative stress as a crucial factor in liver diseases.

World Journal of Gastroenterology 2014;20(25):8082–8091.

[107] Ježek J, Cooper KF, Strich R. Reactive oxygen species and mitochondrial

dynamics: the yin and yang of mitochondrial dysfunction and Cancer progression.

Antioxidants 2018;7(1). https://doi.org/10.3390/antiox7010013.

[108] Li P-L, Gulbins E. Bioactive lipids and redox signaling: Molecular mechanism and

disease pathogenesis. Antioxidants & Redox Signaling 2018;28(10):911–915.

[109] Barja G. The mitochondrial free radical theory of aging. Progress in Molecular

Biology and Translational Science 2014;127:1–27.

[110] Navarro A, Boveris A. Rat brain and liver mitochondria develop oxidative stress

and lose enzymatic activities on aging. American Journal of Physiology-Regulatory,

Integrative, and Comparative Physiology 2004;287(5):R1244–R1249.

[111] Maeso-Díaz R, Ortega-Ribera M, Fernández-Iglesias A, et al. Effects of aging on

liver microcirculatory function and sinusoidal phenotype. Aging Cell 2018;17(6):e12829.

[112] Garone C, Pietra A, Nesci S. From the structural and (dys)function of ATP

synthase to deficiency in age-related diseases. Life 2022;12(3):401.

[113] He B, Yu H, Liu S, et al. Mitochondrial cristae architecture protects against mtDNA
release and inflammation. Cell Reports 2022;41(10):111774.
[114] Amorim JA, Coppotelli G, Rolo AP, et al. Mitochondrial and metabolic dysfunction
in ageing and age-related diseases. Nature Reviews Endocrinology 2022;18(4):243–
258.
[115] Pahal S, Mainali N, Balasubramaniam M, et al. Mitochondria in aging and age-
associated diseases. Mitochondrion 2025;82:102022.
[116] Lian C-Y, Zhai Z-Z, Li Z-F, et al. High fat diet-triggered non-alcoholic fatty liver
disease: A review of proposed mechanisms. Chemico-Biological Interactions
2020;330:109199.
[117] Graham DS, Liu G, Arasteh A, et al. Ability of high fat diet to induce liver pathology
correlates with the level of linoleic acid and Vitamin E in the diet. PLOS One
2023;18(6):e0286726.
[118] Softic S, Meyer JG, Wang G-X, et al. Dietary sugars alter hepatic fatty acid
oxidation via transcriptional and post-translational modifications of mitochondrial
proteins. Cell Metabolism 2019;30 4:735–7534.
[119] Kim HJ, Zhao J, Walwski JL, et al. A high fat diet fosters elevated bisretinoids.
Journal of Biological Chemistry 2023;299(6):104784.
[120] Parlakgül G, Pang S, Artico LL, et al. Spatial mapping of hepatic ER and
mitochondria architecture reveals zonated remodeling in fasting and obesity. Nature
Communications 2024;15(1):3982.
[121] Arruda AP, Pers BM, Parlakgül G, et al. Chronic enrichment of hepatic
endoplasmic reticulum-mitochondria contact leads to mitochondrial dysfunction in
obesity. Nature Medicine 2014;20(12):1427–1435.
[122] Zheng P, Ma W, Gu Y, et al. High-fat diet causes mitochondrial damage and
downregulation of mitofusin-2 and optic atrophy-1 in multiple organs. Journal of Clinical
Biochemistry and Nutrition 2023;73(1):61–76.
[123] Dong J, Chen L, Ye F, et al. Mic19 depletion impairs endoplasmic reticulum-
mitochondrial contacts and mitochondrial lipid metabolism and triggers liver disease.
Nature Communications 2024;15(1):168.

1421 [124] Rector RS, Thyfault JP, Uptergrove GM, et al. Mitochondrial dysfunction precedes

1422 insulin resistance and hepatic steatosis and contributes to the natural history of non-

1423 alcoholic fatty liver disease in an obese rodent model. Journal of Hepatology

1424 2010;52(5):727–736.

1425 [125] Zhang Y, Yang J, Min J, et al. The emerging role of E3 ubiquitin ligases and

1426 deubiquitinases in metabolic dysfunction-associated steatotic liver disease. Journal of

1427 Translational Medicine 2025;23:368.

1428

1429

1430 **Figure legends**

1431 **Graphical Abstract:** Liver aging causes metabolic, lipidomic, and mitochondrial
1432 structural alterations, reflecting age-dependent losses in the MICOS complex. Key
1433 components of the MICOS complex (MIC60, CHCHD3 and CHCHD6) are illustrated.

1434

Figure 1. Assessing the relationship between *CHCHD3, CHCHD6*, and *OPA1*
genetically modeled gene expression and liver diseases in a clinical biobank.

(A) Genetically regulated gene expression (GREX) was modeled from GTEx version 8
data, which includes genotype data linked to measured gene expression across 49
tissues. GREX was calculated within the BioVU population for the MICOS complex
genes CHCHD3, CHCHD6, and OPA1. Liver disease phenotypes were extracted from
the de-identified medical records of BioVU participants. Associations between the
MICOS GREX and liver phenotypes were performed using logistic regression models.
(B-C). Association results for CHCHD3 GREX in (B) 65,363 individuals of European
(EUR) genetic ancestry and (C) 12,313 individuals of African (AFR) genetic ancestry
with three liver disease phenotypes. (D-E) Association results for OPA1 GREX in (D)
65,363 individuals of European (EUR) genetic ancestry and (E) 12,313 individuals of
African (AFR) genetic ancestry with three liver disease phenotypes (phecode 571:
Chronic liver disease and cirrhosis, phecode 571.51: Cirrhosis of liver without mention
of alcohol, phecode 573.2: Liver replaced by transplant). Grey line annotates nominal
significance (p=0.05) and black line annotates the within-tissue Bonferroni-corrected p-
value (p=0.0056). GREX tissue model results are ranked by -log10p value on the y axis
and -log10p values are represented on the x axis. A comprehensive supplementary
table with all association results (effect sizes, standard errors, p-values) for all gene-
tissue-phenotype combinations are found in the supplementary information.

Figure 2: MICOS mRNA fold changes in young and old subjects.

Human mRNA fold change in (A) Mic19, (B) Mic25, (C) Mic60, (D) BIP, (E) PERK, (F)
XBP1, (G) CHOP, (H) IRE1, and (I) ATF4 expressions between young and old human
liver tissue samples (n = 5/group). Murine model mRNA fold change in (J) Opa1, (K)
Mitofilin, (L) Chchd3, and (M) Chchd6 expression in mouse liver tissue between young

(3 months of age) and old (2 years of age) cohorts (n = 3/group). Mann–Whitney tests
were used for statistical analysis. Statistical significance is denoted as ns (not
significant), *p < 0.05, **p < 0.01 and ***p < 0.001.

1463

Figure 3: Loss of *Mic6*0 and *Chchd6* in HepG2 cells results in reduced mCa2+
uptake and calcium retention capacity and oxidative stress.

(A) Raw traces showing mitochondrial calcium uptake in permeabilized *MIC60* and
CHCHD6 knockdown HepG2 cells along with scr-siRNA transected controls. (B)
Percentage change in mCa2+ uptake rate quantified from raw traces. (C) Recordings of
mitochondrial calcium retention capacity in scr-siRNA, *MIC60* siRNA, and *CHCHD6*
siRNA HepG2 cells. The circles indicate the number of calcium boluses taken up by
specific cells. (D) Percentage change in mitochondrial calcium retention capacity
quantified from recordings of mitochondrial calcium retention capacity (CRC). (E)
Immunoblot confirming siRNA-mediated knockdown of *CHCHD6* in HepG2 cells. (F)
Immunoblot confirming siRNA-mediated knockdown of *MIC60* in HepG2 cells. (G) 4′,6-
diamidino-2-phenylindole (DAPI) staining, MitoPY1 (5 μM, 45 min at 37°C magnification
of 60x), and merge channels in scramble-siRNA (control), MIC60-siRNA, and CHCHD6-
siRNA transfected permeabilized HepG2 cells. (H) 4′,6-diamidino-2-phenylindole (DAPI)
staining, MitoBright Deep Red (10 μM, 30 min at 37°C), DCFDA (10 μM, 30 min at
37°C, magnification of 60x), and merge channels in scramble-siRNA (control), MIC60-
siRNA, and CHCHD6-siRNA transfected permeabilized HEK293 cells. (I) Plate reader-
based reactive oxygen species (ROS) quantification. (J) Microscopy-based ROS
quantification of MitoPY1 orange, (K) MitoSox Deep Red, and (L) DCFDA. For all
statistical tests, a one-way ANOVA statistical test was performed with Dunnett's multiple
comparisons test. N=5-10 for all calcium experiments, each indicated by dots, as run in
triplicate. N=9-13 for all oxidative stress experiments, each indicated by dots, as run in
triplicate. Significance values indicate **P ≤ 0.01 and ****P ≤ 0.0001.

1487

1488

1489 **Figure 4. Global metabolomic profiling reveals changes in lipid composition,**
1490 **mitochondrial DNA, and metabolic pathways.**

1491 (A) Metabolomic heatmap of the top 25 most enriched metabolic pathways in 3-month
1492 and 2-year mouse liver tissue. For each tissue and metabolite in the heatmaps, the 2-
1493 year samples were normalized to the median of the 3-month samples and then log2
1494 transformed. (note: p-values were adjusted to correct for multiple comparisons using an
1495 FDR procedure) and log fold changes greater than 1 or less than –1. 3-month, n= 4; 2-
1496 year, n= 4. (B) Pathway impact graph of metabolic pathways in 3-month and 2-year
1497 mouse liver tissue that are affected by aging. A higher impact factor and a larger circular
1498 area are associated with greater fold enrichment of metabolic pathways. More red
1499 coloring of circles is associated with higher statistical significance. (C) PCA chart of
1500 metabolite concentration between 3-month and 2-year-old samples. (D-O)
1501 Concentration of metabolites (D) retinoic acid, (E) retinal, (F) succinate, (G) malate, (H)
1502 GMP, (I) CMP, (J) UMP, (K) AMP, (L) ADP, (M) NAD+, (N) NADP, and (O) NMN in
1503 mouse liver in 3-month (blue) and 2-year-old (pink) samples. For all panels, error bars
1504 indicate SEM, Mann–Whitney tests were used for statistical analysis, and significance
1505 values indicate $*P \leq 0.05$, $**P \leq 0.01$, $***P \leq 0.001$, $****P \leq 0.0001$, and ns indicates
1506 non-significant.
1507 For all panels, error bars indicate SEM, ** indicates p< 0.01; and *p< 0.05, calculated
1508 with Student's t-test.

1509

1510 **Figure 5. Comparative analyses of liver lipid content in young and old humans**

1511 (A-D') Cross-sectional imaging of in-phase (A-D) and out-of-phase (A'-D') liver anatomy
1512 data from (A,A') males under 55 years old (aged 14–53 years old; n = 10), (B,B')
1513 females under 55 years old (aged 17–41 years old; n = 10), (C,C') males over 60 years
1514 old (aged 60–85 years old; n = 10), and (D,D') females over 60 years old (aged 64–96
1515 years old; n = 10). (E-F) Liver lipid percentage in young and old cohort in (E) males and
1516 (F) females and (G) Body Fat%.

1517

1518 **Figure 6. Global lipidomic profiling reveals variations in lipid classes and chain**
1519 **lengths with age in liver tissues.**

1520 (A-D) Oil Red O Staining of liver tissue from 3-month (A-B) and 2-year (C-D) old mice at
1521 (A, C) 20x and (B, D) 40x magnification. (E) Quantification of Oil Red O staining per
1522 area (n=10 per group). (F) Quantification of liver weight relative to total body weight as a
1523 percent (n=10), (G) relative real-time quantitative polymerase chain reaction mtDNA
1524 content (n=8), (H) bile acids concentrations (n=10), (I) liver triglycerides concentration
1525 (n=8), (J) and serum triglycerides concentration (n=8). (K-N) TEM image of mouse liver
1526 tissue from (K) 3-month and (L) 2-year-old mice. (M) Number of lipid droplets per μm^2 of
1527 area in TEM image. (N) Total area of TEM image consisting of lipid droplets in μm^2. (O)
1528 Lipid class enrichment heatmap. (P) Volcano plot labeling significant hits in lipid
1529 expression between 3-month and 2-year-old livers, which have adjusted p-value <0.05
1530 and fold change (+ or -) greater than 1. (Q) Lipid class (R) and lipid chain length
1531 enrichment based on comparison between 3-month and 2-year-old livers. For each
1532 tissue and metabolite in the heatmaps, the 2-year-old samples were normalized to the
1533 median of the 3-month samples and then log2 transformed. Significantly different lipid
1534 classes represented in the figures are those with adjusted p-values < 0.05 (note: p-
1535 values were adjusted to correct for multiple comparisons using an FDR procedure) and
1536 log fold changes greater than one or less than –1. 3-month, n= 4; 2-year-old, n= 4. For
1537 all panels, error bars indicate SEM, ** indicates p< 0.01; and *p< 0.05, calculated with
1538 Student's   t-test.

1539

1540 **Figure 7: TEM and SBF-SEM Shows Changes in Mitochondria Murine Liver**
1541 **Across Aging.**

1542 (A-D) Representative transmission electron micrographs in (A, C) males and (B, D)
1543 females from (A-B) 3-month and (C-D) 2-year murine liver tissue. (E) Mitochondrial
1544 quantifications of male mitochondria number per μm^2 area in TEM image (n=21 images
1545 across 5 mice, 3-months; n=24 images across 5 mice, 2-years) (G) mitochondrial area
1546 quantified in 3 mice, over 3 ROI (n=1309 mitochondria, 3-months; n=1333, 2-years) and
1547 (K) cristae score (n=555, 3-months; n=555, 2-years) in male murine liver TEM images.

(F) Mitochondrial quantifications of the number of female mitochondria per μm^2 unit area

in the TEM image. (n=21, 3-months; n=25, 2-years) (H) mitochondrial area (n=1253, 3-

months; n=1018, 2-years) (I) Combined Mitochondrial quantifications of male and

female mitochondria number per μm^2 unit area in TEM image across aging. (n=21 each

sex, 3-months; n=25 each sex, 2-years) (J) Combined mitochondrial area for males and

females across aging (n=1253 female, n=1309 male, 3-months; n=1018 female, n=1333

males, 2-years) and (L) cristae score (n=684, 3-months; n=684, 2-years) in female

murine liver TEM images. (M) Combined cristae score by sex across aging. (N-Q) Work

through of 3D reconstruction by SBF-SEM. (N) Schematic depicting removal of the liver

from mice. (O) Schematic showing Serial Block Face-Scanning Electron Microscopy

(SBF-SEM) for serial imaging of ortho slices representing different depths of tissue. (P)

Manual contour segmentation of ortho slices was performed to yield (Q) 3-dimensional

(3-D) reconstructions of individually colored mitochondria. (R-S) Representative ortho

slice images from (R) 3-month murine liver tissue and (S) 2-year murine liver tissue. (R'-

S') Representative ortho slice images with 3D reconstructions of mitochondria overlaid

from (R') 3-month murine liver tissue and (S') 2-year murine liver tissue. (R''-S'')

Isolated 3D reconstructions of mitochondria from (R'') 3-month murine liver tissue and

(S'') 2-year murine liver tissue (T-Z) Quantification of 3-D reconstructed mitochondria for

(T) mitochondrial perimeter, (U) area, and (V) volume in 3-month and 2-year-old mice

(W-X) 3D reconstructions of mitochondria displayed from the transverse viewpoint in

(W) 3-month and (X) 2-year murine liver tissue. (W'-X') Representative images of 3D

reconstructions of mitochondria displayed from the longitudinal viewpoint in (W') 3-

month and (X') 2-year murine liver tissue. (Y-EE) Measurement of average

mitochondrial sphericity (Y) and complexity index (Z) in 3-month and 2-year-old murine

liver samples. AA-EE) Measurements of mitochondrial (AA) perimeter, (BB) 3D area,

(CC) volume, (DD) sphericity, and (EE) complexity index in individual mice for 3-month

(blue) and 2-year (pink) mice. Display of individual mitochondria as displayed by mito-

otyping for (FF) 3-month and (GG) 2-year murine liver tissue. For SBF-SEM, a total of

approximately 1500 mitochondria were analyzed, 750 per age group, from 3 mice per

group. For all panels, error bars indicate SEM, Mann–Whitney tests were used for

1578 statistical analysis, and significance values are denoted as follows: **P ≤ 0.01, ****P ≤

1579 0.001, and ns, not significant.

1580 **Figure 8: High-Fat vs. Low-Fat Diet Associated with Changes to Mitochondrial**

1581 **Morphology and Alterations to Physiological Parameters**

1582 (A-B') Representative transmission electron images of murine livers of mice fed (A-A')

1583 low-fat and (B-B') high-fat diets. (C-G) Measurements of (C) number of mitochondria per

1584 μm^2 area, (D) percentage of mitochondrial area normalized to total cell area, (E) total

1585 mitochondrial area, (F) circularity index, and (G) cristae score of mitochondria from low-

1586 fat and high-fat diet murine livers. (H-I) Measurement of (H) number of lipid droplets per

1587 μm^2 area and (I) lipid droplet area in low-fat and high-fat diet murine livers. (J-P)

1588 Measurement of physiological parameters (J) serum cholesterol, (K) serum

1589 triglycerides, (L) serum free fatty acids, (M) total fat mass, (N) liver weight, (O) body

1590 weight, and (P) serum insulin in mice fed low-fat and high-fat diets. (Q) Oral glucose

1591 tolerance test (OGTT) for male mice fed low-fat and high-fat diets. For all panels, error

1592 bars indicate SEM, ** indicates $p < 0.01$; and *$p < 0.05$, calculated with Student's t-test.

Visual Abstract

Mitochondrial cristae morphology is maintained by the MICOS complex

- outer membrane
- inner membrane
- cristae
- MICOS complex
- SAM complex

OPA1
MIC25
MIC60
Para
MIC19
Sam50
35
37

In the liver, aging causes changes in fat %, mitochondrial structure, wrappER, metabolomics, and lipidomics

Young

Old

Age-dependent losses of the MICOS complex contribute to dysregulated calcium and oxidative stress

ROS

Ca²⁺

MICOS Complex

Liver aging

Human data

Animal data

	Male		Female	
Young

 | | | |
| Old | | | | |