Title: MICOS Complex Loss Governs Age-Associated Murine Mitochondrial Architecture and Metabolism in the Liver, While Sam50 Dictates Diet Changes

Authors: Zer Vue[*1], Alexandria Murphy[2*], Han Le[1*], Kit Neikirk[1], Edgar Garza-Lopez[3], Andrea G. Marshall[1], Margaret Mungai[1], Brenita Jenkins[2], Larry Vang[1], Heather K. Beasley[1], Mariaassumpta Ezedimma[1], Sasha Manus[1], Aaron Whiteside[1], Maria Fernanda Forni[4], Chanel Harris[1,5], Amber Crabtree[1], Claude F. Albritton[5,6], Sydney Jamison[6], Mert Demirci[6], Praveena Prasad[2], Ashton Oliver[1], Ky'Era V. Actkins[7], Jianqiang Shao[8], Elma Zaganjor[1], Estevão Scudese[1], Benjamin Rodriguez[1], Alice Koh[1], Izabella Rabago[1], Johnathan Moore[1], Desiree Nguyen[1], Muhammad Aftab[3], Benjamin Kirk[3], Yahang Li[3], Nelson Wandira[6], Taseer Ahmed[6]Ahmed[6]Ahmed[6]Ahmad[6,9], Mohammad Saleem[6] Ashlesha Kadam[10], Prasanna Katti[11,12], Ho-Jin Koh[13], Chantell Evans[14], Young Do Koo[3,15], Eric Wang[16], Quinton Smith[16], Dhanendra Tomar[9], Clintoria R. Williams[17], Mariya T. Sweetwyne[18], Anita M. Quintana[19], Mark A. Phillips[20], David Hubert[20], Annet Kirabo[6,21,22,23], Chandravanu Dash[24], Pooja Jadiya[25], André Kinder[26], Olujimi A. Ajijola[27], Tyne W. Miller-Fleming[7], Melanie R. McReynolds[#2], Antentor Hinton, Jr[#1],

Affiliations:

1. Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, 37232, USA.

2. Department of Biochemistry and Molecular Biology, The Huck Institute of the Life Sciences, Pennsylvania State University, State College, PA 16801

3. Department of Internal Medicine, University of Iowa, Iowa City, IA, 52242, USA.

4. Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT 06520.

5. Department of Biomedical Sciences, School of Graduate Studies, Meharry Medical College, Nashville, TN 37208-3501, USA.

6. Department of Medicine, Vanderbilt University Medical Center, Nashville, TN 37232, USA.

7. Division of Genetic Medicine, Department of Medicine, Vanderbilt University Medical Center, Nashville, Tennessee, United States

8. Central Microscopy Research Facility, University of Iowa, Iowa City, IA, 52242, USA.

9. Department of Pharmacology, College of Pharmacy, University of Sargodha, Sargodha, Punjab,40100, Pakistan

10. Department of Internal Medicine, Section of Cardiovascular Medicine, Wake Forest University School of Medicine, Winston-Salem, NC 27157 USA

11. National Heart, Lung and Blood Institute, National Institutes of Health, 9000 Rockville Pike, Bethesda, MD 20892, USA

12. Department of Biology, Indian Institute of Science Education and Research (IISER) Tirupati, AP, 517619, India

13. Department of Biological Sciences, Tennessee State University, Nashville, TN 37209, USA

14. Department of Cell Biology, Duke University School of Medicine, Durham, NC, 27708, USA

15. Fraternal Order of Eagles Diabetes Research Center, Iowa City, Iowa, USA1

16. Department of Chemical and Biomolecular Engineering, University of California, Irvine, CA, 92697, USA.

17. Department of Neuroscience, Cell Biology and Physiology, Wright State University, Dayton,

OH 45435 USA

18. Department of Laboratory Medicine and Pathology, University of Washington, Seattle, WA,

98195, USA

19. Department of Biological Sciences, Border Biomedical Research Center, The University of

Texas at El Paso, El Paso, Texas, USA

20. Department of Integrative Biology, Oregon State University, Corvallis, OR, 97331, USA.

21. Vanderbilt Center for Immunobiology, Nashville, TN, 37232, USA.

22. Vanderbilt Institute for Infection, Immunology and Inflammation, Nashville, TN, 37232,

USA.

23. Vanderbilt Institute for Global Health, Nashville, TN, 37232, USA.

24. Department of Microbiology, Immunology and Physiology, Meharry Medical College,

Nashville, TN, United States

25. Department of Internal Medicine, Section of Gerontology and Geriatric Medicine, Sticht

Center for Healthy Aging and Alzheimer's Prevention, Wake Forest University School of

Medicine, Winston-Salem, NC

26. Artur Sá Earp Neto University Center – UNIFASE-FMP, Petrópolis Medical School, Brazil

27. UCLA Cardiac Arrhythmia Center, University of California, Los Angeles, CA, USA.

*These authors share co-first authorship.

#These authors share co-senior authorship.

Corresponding Author:

4

Antentor Hinton

Department of Molecular Physiology and Biophysics

Vanderbilt   University

antentor.o.hinton.jr@Vanderbilt.Edu 319-383-3095

ABSTRACT:

The liver, a metabolic hub, undergoes significant declines due to aging, affecting mitochondrial

function and increasing the risk of systemic liver diseases. This study investigates age-related

changes in murine liver mitochondrial structure using Serial Block Face-Scanning Electron

Microscopy (SBF-SEM). We found aging leads to diminished mitochondrial size and

complexity, accompanied by altered metabolomic and lipidomic profiles, reflecting findings in

aged human samples. We explored the Mitochondrial Contact Site and Cristae Organizing

System (MICOS) complex as a regulator of these changes, observing its decline with age, unlike

Sam50, which remains stable. Sam50, part of the sorting and assembly machinery (SAM)

complex, decreases notably in murine models on high-fat diets, correlating with increased

susceptibility to liver diseases in humans. Additionally, we found dysregulated mitochondrial

calcium handling and increased oxidative stress with MICOS complex disruption. Our study

underscores age-related reductions in mitochondrial complexity and metabolic dysregulation in

murine liver aging. Together, we establish that a decrease in mitochondrial complexity and

dysregulated metabolism occur with murine liver aging. While these changes are partially be

regulated by age-related loss of the MICOS complex, the confluence of a murine high-fat diet

can also cause loss of *Sam50*, which contributes to liver diseases.

Keywords: Aging, 3D Structure, Mitochondria, Metabolism, MICOS Complex, Liver Disease

93 **INTRODUCTION:**

94   Mitochondria are required for various purposes including oxidative phosphorylation,

95 calcium homeostasis, and biochemical pathways (1). Mitochondria structure is crucial to many

96 mitochondrial functions (2, 3). Unlike many other organelles, mitochondria have independent

97 fusion and fission dynamics, which allow them to dynamically respond to their environment and

98 adapt a variety of phenotypes beyond the spherical or tubular shapes that are typically associated

99 with mitochondria (2, 4, 5). It is well understood that across aging, changes in morphology and

100 size can impair mitochondrial functions and structures **(6–8)**. Typically, these declines are

101 measured through transmission electron microscopy (TEM), which offers high x- and y-

102 resolutions (9, 10) providing insights into mitochondrial ultrastructure such as cristae, the folds

103 of the inner mitochondrial membrane (11). However, a nascent research avenue is studying 3D

104 phenotypes and exploring novel mitochondria structures (2, 12), such as donut-shaped

105 mitochondria (13, 14), nanotunnels (15, 16), and megamitochondria (17). These structures

106 require imaging techniques, such as serial block face-scanning electron microscopy (SBF-SEM),

107 which allows 3D reconstruction of mitochondria **(18–22)**. For many organs and experimental

108 models, 3D reconstruction has not yet been performed, thus creating a gap in knowledge of 3D

109 mitochondrial structural changes and molecular regulators that are present across the aging

110 process in various tissues. Previously, we have demonstrated changes in mitochondria in other

111 tissues, such as cardiac and skeletal muscle tissue (23, 24). Here, we aim to use 3D

112 reconstruction to understand mitochondrial remodeling in the aging murine liver—which serves

113 as the metabolic hub for the body.

114   The liver is the largest internal organ in vertebrates, performing essential biological functions

115 such as detoxification, metabolism of drugs and nutrients, storage of minerals, and synthesis of

116 plasma (25). The liver facilitates digestion and metabolism, which has an impact on the

117 gastrointestinal and endocrine systems (26). Additionally, the liver generates carrier proteins

118 crucial for growth and reproduction and participates in the metabolism of sex hormones (27).

119 Therefore, studying liver metabolism will provide insights into many crucial biological processes

120 including lipid and glucose homeostasis and detoxification pathways. One of the principal

121 factors that deteriorate the liver is aging (26). After the age of 45, the percentage of mortality

122 linked to liver disease rises significantly in humans (28). With aging comes a progressive

123 expansion of the liver's neural fat and cholesterol volumes and a rise in blood cholesterol, high-

124 density lipoprotein cholesterol, and neutral fat levels (29). Volume alterations, polyploidy nuclei,

125 the accumulation of dense bodies containing lipofuscin, a reduction in smooth endoplasmic

126 reticulum area, and most importantly, a decrease in the quantity and functionality of

127 mitochondria are all associated with aging in liver cells (29). During aging, lipids are also readily

128 accumulated by the liver, raising the risk of cirrhosis, steatohepatitis, and non-alcoholic fatty

129 liver disease (NAFLD), among other hepatic conditions (30, 31). Lipids are necessary for several

130 essential mitochondrial functions, including metabolism and biogenesis, mitochondrial

131 dynamics, oxidative phosphorylation, architecture, and structure of mitochondrial membranes

132 (32). Additionally, the production of energy, regulation of cell death, and bioenergetics are all

133 significantly impacted by the lipid content of the inner mitochondrial membrane (IMM) (32).

134 Thus, as liver cells undergo mitochondrial dysfunction during aging, mitochondrial alterations

135 may contribute to the age-related risk of systematic diseases (25, 33). Thus, within this study, we

136 combine our murine approach of looking at mitochondrial structure with the examination of

137 gross liver structure, lipogenesis, and genetically regulated gene expression in human cohorts.

Age-related changes in mitochondrial function are characterized by dynamic structural

alterations within the mitochondria. Mitochondria are important for generating ATP through the

electron transport chain in their cristae, folds of the IMM (11). Age-related liver disease may

decrease ATP synthesis (34), resulting decreased energy allocation for essential functions.

Beyond this, mitophagy, mitochondria DNA (mtDNA) mutations, disruptions in calcium

homeostasis, and reactive oxygen species (ROS) have been implicated in many liver pathologies,

such as alcohol-related liver disease, non-alcoholic fatty liver disease, chronic hepatitis B and

hepatocellular carcinoma (35, 36). Through mitochondrial quality control mechanisms, oxidative

stress, mtDNA, mitochondrial structure are highly interdependent (37, 38). Notably, changes in

mitochondrial structure are often driven by changes in mitochondrial dynamic proteins (4, 39),

which coordinate fusion and fission processes, allowing mitochondria to increase volume or

repair mtDNA defects through fusion. In contrast, fission can form new mitochondria and

remove damaged mitochondria (40). The organization of proteins belonging to the dynamin-

related GTPase family and their adaptors controls this mitochondrial remodeling (41). The key

regulators of mitochondrial dynamics are mitofusins 1 and 2 (MFN1 and MFN2) and optic

atrophy 1 (OPA1). MFN 1 and 2 regulate outer mitochondrial membrane (OMM) fusion,

whereas OPA1 regulates IMM fusion (42, 43). In contrast, fission is typically associated with

dynamin-related protein 1 (DRP-1). DRP1 redistribution from the cytosol to the OMM is

necessary for mitochondrial fission **(42, 44–46)**. Dysregulation of these finely tuned dynamics

can support pathophysiology; for example, in liver cancer, upregulation of *Mfn1* and *Opa1* play a

role in cancer tumor cell growth mechanisms (5). Conversely, increased *Drp1* and impaired

Mfn1 expression were hallmarks of chronic alcohol exposure in mice; while other models show

Drp1 downregulation upon chronic alcohol exposure (35). Deletion of mitochondrial fission

161 factor, a recruitment factor of DRP1, in hepatocytes, has similarly led to altered mitochondrial

162 morphology concomitant with enhanced susceptibility to NAFLD (47), but conversely, blocking

163 fission can ameliorate hepatic steatosis and oxidative stress in NAFLD (48). This underscores

164 that while traditional fusion and fission dynamic proteins can affect age-related risk to liver

165 diseases, other regulators must be considered.

166   The OMM and IMM include hardwired translocase complexes with a very intricately

167 specialized protein import mechanism that drives mitochondrial dynamics. These translocases

168 operate along five distinct protein import pathways, one of which is through the SAM50 channel

169 (49, 50). *Sam50* is one of the crucial proteins of the outer membrane of the mitochondria.

170 SAM50 is a SAM complex component with a β-barrel domain (50). Although the role of *Sam50*

171 in mitochondria is still unclear, studies showed that Sam50 contributes to the preservation of

172 cristae's structural integrity. *Sam50* maintains the formation of the cristae via interaction with

173 mitofilin and CHCHD6, which are parts of the mitochondrial contact site and cristae organizing

174 system (MICOS) (51). Furthermore, *Sam50* aids in the maintenance of the mitochondrial

175 network (50), establishing and maintaining mitochondrial integrity and mitochondrial

176 metabolism, with its loss conferring volumetric losses in mitochondria structure (52). Recently,

177 *Sam50* has been linked to NAFLD pathogenesis (49), but the interconnection with aging liver

178 mitochondrial structure remains poorly elucidated.

179   In our study, we use human models to explore the impairments of the liver due to aging,

180 investigating changes in the structure and physical appearance of the liver in young and old

181 groups. To establish the tandem roles of MICOS and SAM50, we used two-year-old mice, which

182 represented an elderly model, and three-month-old mice, which represented a juvenile

183 phenotype. We then studied mitochondrial morphological changes in both two-dimensional (2D)

and three-dimensional (3D) which showed age-related losses in volume and complexity. We also

find that metabolomic and lipogenic changes in aging may mediate mitochondrial structural

alterations in the aging liver. We further learned how the MICOS complex is dysregulated in the

aging liver. Notably, deletion of the MICOS genes resulted in impairments in mitochondrial

calcium regulation and ROS production in the liver. Due to the changes observed in

mitochondria function and structure, we also investigated the role of diet-dependent *Sam50*

depletion in the liver. Lastly, we modeled genetically-regulated gene expression (GReX) of

SAMM50 in a biobank of 85,615 participants, and found that SAMM50 GReX was significantly

associated with multiple liver diseases.

Results:

Aging Causes Alterations in Liver Lipid Accumulation and Altered Disease Markers

Damage to the liver's tissue and alterations in hepatic metabolism might result from

abnormal lipid levels in the liver (31). Thus, we utilized magnetic resonance imaging to

determine how lipid content is regulated during aging in human samples. By enrolling female

and male participants (Figures 1A–D), we created a "young" cohort consisting of individuals

from 18 to 55 years old and an "old" cohort of individuals older than 60 years old. For both

sexes, the liver's lipid percentage increased through aging (Figures 1E–F). This age-related

influx in liver validated the progression towards liver diseases in humans. Specifically, when

combining the male and female cohorts together (Figure 1G), the mean fat percent increased

(from a mean of 1.41% $\pm$ 1.44% SD to 5.93% $\pm$ 6.11% SD). Additionally, the 75% percentile of

fat percentage was higher, at 10.5% in the aged cohort, as compared to 2.10% in the young

cohort. Notably, Grade 1 of fat fraction classification, representing mild hepatic steatosis has a

fat percentile cutoff of 6.5% or higher (53). To further validate these results, we switch to our

mouse model and used Oil Red O staining to visualize lipids in 3-month and 2-year murine liver

tissue (Figures 1H-K). Specifically, we saw an increased amount of Oil Red O amount in the 2-

year cohort signifying an increase in lipid amount across the aging process in the liver, with a

more than 6-fold increase in lipid droplets in 2-year samples (Figure 1L; 12.2 $\pm$ 6.55 SD, 3-

months; 77.5 $\pm$ 20.4 SD, 2-years). An uptick of lipid droplets in hepatocytes is indicative of

NAFLD (54). To further explore markers of potential disease progression and dystrophy outside

of lipid accumulation, we also looked at liver mass, which was decreased in aged samples when

normalized to body weight (Figure 1M; 4.56% $\pm$ 0.329% SD, 3-months; 3.93% + 0.376% SD, 2-

years). Since mtDNA content reduction is a hallmark of NAFLD (55), we also showed an age-

dependent loss in mtDNA content (Figure 1N; mean: 1, 3-months; 0.805 $\pm$ 0.119 SD, 2-years).

Additionally, bile acid, which is metabolized in the liver, has increased concentrations with

abnormal liver function and disease states (56). We also looked at the concentration of bile acid,

but no significant difference was noted (Figure 1O; 0.871 µMol/µg $\pm$ 0.138 µMol/µg SD, 3-

months; 0.89 µMol/µg $\pm$ 0.15 µMol/µg SD, 2-years). Finally, central to the pathology and a key

marker of NAFLD is high triglycerides (57, 58). We observed that aging murine samples had

significantly elevated triglycerides in the liver (Figure 1P; 0.30 mmol/L $\pm$ 0.10 mmol/L SD, 3-

months; 0.748 mmol/L $\pm$ 0.15 mmol/L SD, 2-years), which was more drastic than the general

increase in serum triglycerides (Figure 1Q; 0.211 mmol/L $\pm$ 0.100 mmol/L SD, 3-months; 0.379

mmol/L $\pm$ 0.118 mmol/L SD, 2-years). Thus, while we could not confirm aging human and

murine samples had liver disease, these results support the observation of an age-related

progression toward liver disease.

Ultrastructural Changes in Murine Liver Reveals Aging is Associated with Lower Volume

and Complexity

Next, we sought to explain whether these age-related changes in liver mass and disease

markers are correlative with atypical mitochondria structures in the liver. For these studies, we

utilized aged C57BL/6J mice at 2 age points, 3-months and 2-years, which are generally

understood to be good representations of "young" and "old" aged points akin to aging displayed

in humans (59).

To begin with, we used TEM to measure changes in mitochondrial morphology in aged mice

(Figures 2A-D). Qualitatively, our findings of lipid accumulation were confirmed, with aging

also showing lipid droplets that showed less circularity. TEM analysis showed that mitochondria

count increased while the average mitochondrion area decreased with aging (Figures 2E-J). To

compare sex-dependent differences, mitochondrial counts, as normalized to µm cell area, in 3-

month males and females were similar (0.15 ± 0.02 SD, males; 0.174 ± 0.03 SD, females), there

was a slight divergence with aging, although both followed a similar trend (Figures 2E, 2H; 0.56

$\pm$ 0.130 0.15 SD, males; 0.738 ± 0.14 SD, females). This sex-difference which is more

pronounced with age was similarly seen in average mitochondrion area when comparing 3-

month samples ($1.27 \, \mu m^2 \pm 1.34 \, \mu m^2$ SD, males; $1.21 \, \mu m^2 \pm 1.19 \, \mu m^2$ SD, females) with their 2-

year counterparts (Figures 2F, 2I; $0.233 \, \mu m^2 \pm 0.442 \, \mu m^2$ SD, males; $0.558 \, \mu m^2 \pm 0.386 \, \mu m^2$

SD, females). However, in many ways, TEM is inadequate to quantify mitochondria, which exist

in diverse 3D shapes (2). Yet, given the best in-class resolution of TEM (9), it is useful for

measuring subcellular structures of cristae. In male and female murine samples, we found a

decrease in cristae score both (Figures 2G, 2J), which is a semi-qualitative measure (9), that

253   evaluates the relative amount and architectural integrity of cristae. Neither 3-month (3.72 $\pm$ 0.44

254   SD, males; 3.47 $\pm$ 0.62 SD, females) nor 2-year (2.29 $\pm$ 0.71 SD, males; 1.99 $\pm$ 0.87 SD,

255   females) samples showed a significant sex-dependent difference in cristae score. Together, these

256   results highlight that cristae integrity is lower in aged liver tissue samples. Additionally, we

257   performed these studies in both male and female samples, which showed slight sex-dependent

258   differences but were generally minimal with similar trends. Given the time-consuming nature of

259   manual segmentation associated with 3D reconstruction, we proceeded with the study utilizing a

260   male cohort.

261   With the 2D mitochondrial differences observed, we utilized 3D reconstruction of SBF-

262   SEM to further investigate changes in mitochondrial structure. For each age cohort, we analyzed

263   approximately 250 mitochondria from the three mice surveyed (Figure 2K), for a total of around

264   750 mitochondria for each age point. SBF-SEM allowed for 50 µM intervals in the z-axis to be

265   imaged for a total of 300 slices, of which 50 slices were used for analysis. The x- and y-

266   resolution was 10 µM, which allowed for about 50 ortho slices to be reconstructed and quantified

267   for each stack (Figure 2L), which then undergo manual contour segmentation (Figure 2M), for

268   the 3D reconstruction of mitochondria (Figure 2N). Displayed first in both age cohorts is a

269   representative of each orthoslice, which allows for the identification of mitochondria (Figures

270   2O-P). From there, manual segmentation across each ortho slice was performed in the software

271   Amira, for mitochondria to be reconstructed (Figures 2O'-P'). Additionally, for better

272   visualization of each individually colored mitochondria, the ortho slice may be removed (Figures

273   2O''-P''). Once modeled, 3D reconstructions (Videos 1-2) can be quantified. We found that in

274   comparing the young and aged mice, size significantly decreased in all metrics (Figures 2Q-S).

275   Specifically, the perimeter, which represents the total of all external distances in the

mitochondria was approximately 20% lower in aged samples (Figure 2Q; 10761 µm $\pm$ 1560 µm

SD, 3-months; 8054 µm $\pm$ 450 µm SD, 2 year). Three-dimensional area, which represents a

metric of OMM area, was decreased by nearly 50% in aged samples (Figure 2R; 7.66 µm^2 $\pm$

0.808 µm^2 SD, 3-months; 4.76 µm^2 $\pm$ 0.511 µm^2 SD, 2-years). Finally, mitochondrial volume,

which represents the total of all internal pixels within the 3D mitochondrial reconstruction, was

approximately 30% lower in aged male murine liver samples (Figure 2S; 1.01 µm^3 $\pm$ 0.182 µm^3

SD, 3-months; 0.696 µm^3 $\pm$ 0.107 µm^3 SD, 2-years). Losses in mitochondrial volume may

indicate decreased internal volume for ATP synthesis (60), but mitochondrial roles extend

beyond their energetics, including through interfacing with endoplasmic reticulum (ER) to

modulate calcium homeostasis (61). Thus, it is equally important to look at their morphology and

capacity to form compact sites.

To aid in visualizing mitochondria, we present mitochondria 3D reconstructions from

transverse (Figures 3A-B) and longitudinal (Figures 3A'-B') viewpoints (Videos 3-4). To further

verify this change across aging we looked at sphericity and mitochondrial complexity index

(MCI) (62), which are analogous measures of mitochondrial morphology. Sphericity (calculated

as $\frac{\pi^{\frac{1}{3}}(6*Volume)^{\frac{2}{3}}}{Surface\ Area}$), as exhibited an approximate 20% higher mean value (Figure 3C; 0.638 $\pm$

0.0211 SD, 3-months; 0.785 $\pm$ 0.0257 SD, 2-years). MCI (calculated as Surface Area^3 /16 π

^{2}volume^2) expectedly showed an inverse decrease in the aged cohort (Figure 3D; 3.22 $\pm$ 0.240

SD, 3-months; 1.64 $\pm$ 0.236 SD, 2-years). Together, these validate that complexity is decreased

in aged murine liver samples. To better visualize these changes, we organized mitochondria by

their volume, in a method known as mito-otyping, to compare morphology across size (Figure

3E). In the three-month age cohort (Figure 3E), we observed an expected phenotype of normally

sized mitochondria. However, we also observed some diversity in mitochondrial structure, as we

noted elongation, branching, and other structures that prioritize surface area over volume. In

contrast, the 2-year sample showed much less heterogeneity and mitochondrial structures mostly

presented as tubular or compact (Figure 3F). For these metrics, while intra-sample heterogeneity

is plentiful, there is minimal inter-sample heterogeneity or outliers (sFigures 1A-E).

These changes in sphericity and morphology may impact the ability of mitochondria to

form contact sites. Particularly, liver mitochondria are known to exhibit phenotypes in which ER

wraps around them, known as wrappER, to modulate lipid flux **(63–65)**. To investigate this

paradigm, we qualitatively identified wrappER within liver tissue (Figures 3G-H). Qualitatively,

we found that wrappER exists in both 3-month and 2-year murine liver tissue (Figures 3G'-H'),

but in 3-month tissue wrappER contact is much more extensive (Figures 3G"-H"). This suggests

that liver lipid homeostasis may be negatively impacted **(63–65)**. However, to confirm these

changes and further explore the implications of these mitochondrial morphological alterations,

we analyzed metabolism and performed lipidomics in mice.

Global metabolic and lipidomic profiling reveals altered energy metabolism and storage in

the aged liver.

Due to observed changes in mitochondria in the aged liver, we proceeded to analyze

alterations in small biomolecules regulating energy metabolism, cellular homeostasis, and

storage. Our investigation involved global metabolic and lipidomic profiling of young and aged

liver tissues, revealing dynamic features and significant changes in aged mouse livers (Figures

4A-O; sFigures 2A-D). We noted an accumulation of metabolites related to Vitamin A

metabolism, specifically retinoic acid and retinal (Figures 4B-C). While Vitamin A metabolism

must be finely balanced, within breast cancers retinoid changes can be associated with

controlling the redox status of cytochrome c (66). This finding supports the presence of

metabolic shifts, altered cellular signaling, and impaired clearance mechanisms in the aging

liver, all attributed to changes in mitochondrial function. Our metabolomics analysis also

uncovered disruptions in tricarboxylic acid (TCA) cycle intermediates—succinate and malate,

indicating mitochondrial dysfunction and impaired activity (Figures 4D-E). These changes can

impact the production and regulation of these intermediates. Furthermore, we observed decreases

in nucleotide monophosphates involved in purine and pyrimidine metabolism—GMP, CMP,

UMP, AMP (Figures 4F-I). The synthesis of these nucleotides, which involves mitochondrial

processes, is influenced by age-related mitochondrial dysfunction, affecting the overall

nucleotide biosynthesis pathway supporting our observations. Our results additionally confirmed

existing literature on dysregulated nicotinamide adenine dinucleotide (NAD+) metabolism in the

aging liver (67). Significant depletions in tissue NAD+, NADP, NMN pools, and tissue ADP

were detected (Figure 8J-M). Historical liver studies suggest that mitochondrial NAD(H) pools

tend to be oxidized, playing a crucial role in energy homeostasis. Conversely, cytosolic

NADP(H) pools tend to be highly reduced for reductive biosynthesis (68). Our data reveals

disruptions to mitochondrial activity reverses these classical mechanisms. In summary, these

findings collectively support the presence of altered energy metabolism and cellular homeostasis

in the metabolically active liver due to mitochondrial dysfunction and impairment.

Lipidomic profiling of both young and aged liver tissues unveiled age-related changes in

lipid classes and chain lengths (Figure 9N-O; sFigures 2C-D). In the aging liver, significant

alterations were observed in the triglycerides oligomers (TGO), triglycerides (TG), ceramide

(Cer), and acylcarnitine (CAR) lipid classes compared to other lipid groups (Figures 4N-O,

sFigure 2C). These lipid classes, each with distinct roles in liver mitochondria, contribute to

various aspects of cellular metabolism and homeostasis. Disruptions in TGOs and TGs suggest

imbalances in energy storage and release in the aging liver, possibly due to mitochondrial

dysfunction. TGs, being the primary form of stored energy, can participate in mitochondrial beta-

oxidation, providing acetyl-CoA for energy production. Conversely, Cer regulates apoptosis and

modulates mitochondrial membrane permeability (69). Changes in Cer levels can influence

mitochondrial integrity and function (70, 71). CARs play a crucial role in transporting fatty acids

into the mitochondria for beta-oxidation, thus maintaining mitochondrial function (72).

Additionally, significant differences in lipid chain lengths were noted with age in the liver,

impacting membrane integrity, fluidity, and functionality (Figure 4D). These findings align with

altered metabolomic and lipogenic changes attributed to the modified shape and function of

mitochondria with age.

Murine Aging and Diet Exhibit Distinct Changes in the MICOS and SAM Complex

It is well established that the MICOS complex is critical for mitochondrial dynamics (73),

and our group has previously investigated the effect of aging on the MICOS complex in kidney

tubular cells (24), however, it is still unclear how aging affects the MICOS in aging liver. Studies

have shown that *Opa1*, which is epistatic to the MICOS complex and physically interacts with

components of the MICOS complex (43), decreases with age (74). With *Opa1* as a positive

control, we sought to determine if the MICOS complex mRNA expression is also decreased in

murine liver with age. Consistent with previous studies, *Opa1* mRNA decreased by over 50%

between 3 months and 2 years (Figure 5A). Mitofilin also decreased significantly (Figure 5B).

Similarly, *Chchd3* and *Chchd6* also progressively decreased with age but not as much as the

368 decline of other transcripts (Figures 5C-D). While *Opa1* interacts with the MICOS complex, it is

369 not required for the formation of cristae junctions at which the MICOS complex forms, nor does

370 *Opa1* loss negatively affect MICOS components (54). This suggests that the loss of the MICOS

371 complex across aging occurs in an *Opa1*-independent manner. This parallels our previous

372 findings in skeletal muscle, cardiac, and kidney (24, 75, 76).

373   The SAM complex, particularly its β-barrel channel *Sam50*, interacts with the MICOS

374 complex to regulate cristae morphology **(77–80)**. Particularly, overexpression of SAMM50

375 enhances fatty acid oxidation and reduces intracellular lipid accumulation, while polymorphisms

376 in SAMM50 are associated with NAFLD (49), and loss of Sam50 in hepatocytes leads to

377 cardiolipin-dependent mitochondrial membrane remodeling, mtDNA release, and liver injury

378 (81). So, we also looked at SAM50 protein levels across aging. Interestingly, unlike the MICOS

379 complex and *Opa1*, we saw no significant changes in SAM50 levels (Figure 5E). Notably,

380 beyond aging, a key risk factor for NAFLD is a high-fat diet (HFD) which can cause lipid

381 accumulation and dyslipidemia (82). Since past studies have shown that overexpression of

382 SAMM50 can mitigate lipid accumulation (49), we focused on how an HFD also affects these

383 regulators. To do this, we subjected male mice to an HFD, with fat constituting 60% of their diet,

384 and a littermate cohort to a low-fat diet (LFD), with fat constituting only 5% of their calories, for

385 a total of 12 weeks, with mice sampled at 20 weeks. Contrary to our age-related findings, we

386 showed that while MIC60, a key component of the MICOS complex, remains unchanged,

387 SAM50 shows a decrease with an HFD. Given the role of SAMM50 in mitigating lipid

388 accumulation (49), this loss may increase susceptibility to lipid accumulation and dyslipidemia

389 that is a hallmark of HFD (82).

390   To further investigate the impact of an HFD on mitochondrial ultrastructure, we

391   performed TEM analysis in a 20-week-old murine cohort subjected to an LFD and HFD (Figures

392   5G-H''). Qualitatively, we observed more lipid droplets with less circularity in HFD samples.

393   Looking at mitochondria, although mitochondrial count, when normalized to the µm cell area,

394   was fewer in HFD samples (Figure 5I; 0.210 ± 0.114 SD, LFD; 0.128 ± 0.0972 SD, HFD), this

395   change was not significant. Similar to aging, HFD samples exhibit a significantly higher average

396   mitochondrion area (Figure 5J; 0.570 µm^2 ± 0.401 µm^2 SD, LFD; 0.923 µm^2 ± 1.12 µm^2 SD,

397   HFD). When considering these two quantifications in tandem, we also calculated the percentage

398   of mitochondrial area related to the total cell area, which showed no significant difference

399   (Figure 5K; 11.8 µm^2 ± 5.59 µm^2 SD, LFD; 11.7 µm^2 ± 5.10 µm^2 SD, HFD). When looking at

400   how mitochondrial shape changes with a HFD, we observed that mitochondrial had a higher

401   circularity, indicative of potentially less complexity, with a HFD (Figure 5L; 0.781 ± 0.149 SD,

402   LFD; 0.897 ± 0.0748 SD, HFD). Finally, looking at cristae score, similar to aging samples, HFD

403   cohorts had a significantly lower cristae score (Figure 5M; 3.74 ± 0.441 SD, LFD; 2.04 ± 0.770

404   SD, HFD). Together, these findings show that a HFD can parallel age-related changes, and that

405   HFD exhibit smaller mitochondria with aberrant cristae due, in part, to a loss of SAM50.

406   Previously, we have performed a rigorous 3D analysis of mitochondrial structure and

407   networking in *Sam50*-deficient human and murine myotubes, showing that the deletion of *Sam50*

408   results in smaller and less complex mitochondria (52). To further investigate SAM50's role in

409   mitochondrial structure, we knocked out *Sam50* in fibroblasts and used mito-mCherry confocal

410   to investigate structure (Figures 5N-O''). Our results further validate our previous findings,

411   showing that Sam50 deletion causes impairments in mitochondrial length (Figure 5P; 4.14 µm $\pm$

412 1.57 µm SD, WT; 2.59 µm $+ 0.539$ µm SD, *Sam50* KO). Additionally, when comparing the

percentage of fragmented mitochondria to tubular mitochondria, there was a significantly higher

rate of fragmented mitochondria in *Sam50* KO than in WT. (Figure 5Q; 3.33% $\pm$ 2.08% SD, WT;

57.3% + 18.0% SD, *Sam50* KO). Notably, these findings parallel our findings in HFD of reduced

mitochondrial size, indicating that loss of *Sam50* may dictate the negative bioenergetics caused

by an HFD (83, 84).

Clinical relevance of Sam50 in liver pathologies

Previously, we found that *Sam50* is responsible for establishing and maintaining

mitochondrial integrity, cristae structure, and mitochondrial metabolism (52). To identify the

clinical implications of dysregulated *SAMM50* gene expression in humans, we modeled

genetically regulated gene expression (GReX) of *SAMM50* across 85,615 individuals in the

Vanderbilt biobank, BioVU (Table 1). Cross-tissue *SAMM50* GReX was calculated using the

genetic information for each individual, combined with *SAMM50* gene expression models

developed from the GTEX dataset (see Methods, sFigure3). The relationship between *SAMM50*

GReX and clinical phenotypes (n=1,704) and laboratory values (n=326) were then evaluated by

logistic and linear regression models, respectively. In 70,440 BioVU individuals of European

ancestry, we found that *SAMM50* GReX is significantly associated with multiple liver disease

diagnoses, including: chronic liver disease and cirrhosis, alcoholic liver damage, liver transplant,

acute gastritis, and esophageal bleeding (p < 2.93427E-05, Figure 5R, SFile1). Within the 15,175

BioVU individuals of African ancestry, we did not find any significant associations between

SAMM50 GReX and the clinical phenotypes tested; however, we did see a nominally significant

association with acute gastritis (p=0.00142, SFile3). We did not find any significant associations

435 between SAMM50 GReX and clinical lab values within the BioVU individuals of European or

436 African ancestry (SFile2, SFile4).

437

438 **Knockdown of MIC60 and CHCHD6 reduces Ca^{2+} uptake and retention capacity in HepG2**

439 **cells**

440　　　While this underscores how diet changes may contribute to liver diseases, we also wanted

441 to establish how the MICOS complex contributes to age-related changes. To unravel the

442 involvement of *MIC60* and *CHCHD6* in mitochondrial calcium ($_mCa^{2+}$) homeostasis, we

443 monitored $_mCa^{2+}$ uptake and $_mCa^{2+}$ retention capacity in *MIC60* and *CHCHD6* knockdown

444 human-derived HepG2 cells. Both *MIC60* and *CHCHD6* knockdown cells show reduced $_mCa^{2+}$

445 uptake (Figure 6A). Furthermore, to determine if the altered MICOS and cristae structure

446 contributes to impaired $_mCa^{2+}$ retention capacity and involved in mitochondrial permeability

447 transition pore opening, we measured the $_mCa^{2+}$ retention capacity in *MIC60* and *CHCHD6*

448 knockdown cells. *MIC60* as well as *CHCHD6* knockdown cells underwent rapid permeability

449 transition opening as compared to control cells (Figure 6B). We also found that $_mCa^{2+}$ retention

450 capacity is significantly reduced in both *MIC60* and *CHCHD6* knockdown HepG2 cells (Figure

451 6C). These data suggest that altered MICOS complex and cristae disorganization increase the

452 susceptibility of HepG2 cells to $_mCa^{2+}$ dysregulation and Ca^{2+}-induced cell death.

453

454 **Knockdown of MIC60 and CHCHD6 induces oxidative stress**

455 Ca^{2+} directly affects oxidative stress signaling and reactive oxygen species (ROS) generation

456 (85). To determine if MICOS affects ROS production, we evaluated total ROS, mitochondrial

457 superoxide, and H$_2$O$_2$ production in *MIC60* and *CHCHD6* knockdown cells (Figures 6G-H).

458 Mitochondrial H_2O_2 content, as measured by Mitochondria peroxy yellow 1, increased following

459 *MIC60* and *CHCHD6* knockdown when quantified by both plate-reader based (Figure 6I) and

460 microscopy-based (Figure 6J) ROS quantification. Silencing *MIC60* and *CHCHD6* in HepG2

461 cells significantly increased mitochondrial superoxide production and more general intracellular

462 ROS, detected by MitoBright Deep Red (Figure 6K) and DCFDA (Figure 6L), respectively.

463 These findings indicate that suppression of *MIC60* and *CHCHD6* disrupts mitochondrial ROS

464 homeostasis, demonstrating that the MICOS complex is associated with oxidative stress.

465

466 **Discussion:**

467

468 *Age-Related Changes in Liver Mitochondria Structure*

469 Liver mitochondria have been imaged using 3D in rodent models since the early 1990s (86).

470 Previously, in rat liver, 3D reconstruction revealed that cristae showed high variability in

471 structure (87). 3D reconstruction also showed that there are extensive mitochondria endoplasmic

472 reticulum contact sites (MERCs) in liver mitochondria [35]. Additionally, MERCs are sites for

473 various purposes, such as calcium homeostasis and lipid homeostasis (88). In mice with high

474 alcohol consumption, alcohol increased mitochondrial volume while conversely reducing the

475 networking of mitochondria (89). Notably, the formation of mega mitochondria, a novel

476 mitochondria 3D structure, arose in human non-alcoholic fatty liver disease (90). Beyond these

477 3D studies, previous studies using cryo-electron tomography (cryo-ET) have found that

478 mitochondria fragmented and lamellar cristae lost integrity across the aging process (91). While

479 these studies have aided in revealing previously unknown specific mitochondrial structures in the

liver, the same has not yet been done for different mitochondrial structures that may arise across

aging.

Here, we use SBF-SEM to attempt to answer how mitochondria change in an aged liver and

the functional impact of such changes. We found that in aged states, the size and morphology of

liver mitochondria are altered in both 2D and 3D. Using SBF-SEM and Amira for 3D

morphology of mitochondria and comparing 2-year against 3-month, we observed fragmentation

of mitochondria and decreased complexity. Importantly, we found that an aged murine liver has

a decrease in area, perimeter, volume, and complexity in mitochondria. Mitochondria structure

remains relevant in aging as structural decline may decrease mitochondrial function, thus

reducing certain liver roles. For example, mitochondria dysfunction has been linked to an

increased risk of liver damage upon drug treatment (92). Beyond this, in humans and mice with

alcoholic liver disease, it has been suggested that megamitochondria form due to the lack of

Drp1 and serve a protective role (93, 94). This may be due to the downregulation of the

NR4A1/DNA-PKcs/p53 pathway which has been shown to activate Drp1 in other liver diseases

(95). Aging and disease states affect each other; therefore, future experiments must look at the

mitochondria 3D reconstruction in different disease states across aging to see if the confluence of

these factors leads to the unique mitochondria 3D structures. Interestingly, while

megamitochondria are a hallmark of hepatotoxicity and steatosis (94, 96, 97), and aging

exhibited numerous hallmarks of progression towards liver diseases (Figure 1), we did not

observe these megamitochondria in our sample, rather showing mostly smaller mitochondria.

Since the formation of megamitochondria in Alcoholic Liver Disease follows a significant

increase in Drp1 expression and mitochondrial fragmentation (98), it may be that our aged

502   sample represents a pre-diseased state that can subsequently shift to megamitochondria as a

503   compensatory response to reduce alcohol-induced toxicity (94, 98, 99).

504   Beyond mitochondrial 3D structure, past studies have shown that fatty acids promote

505   endoplasmic reticulum stress and liver injury in rodent models (100). Due to the increase of lipid

506   droplets in both aging and HFD in TEM images, there may be an increase in mitochondria-lipid

507   droplet contact sites (MLDCs) (61, 101). Beyond this, lipid droplet-endoplasmic reticulum

508   contacts are also understood to help form lipid droplets and perform metabolism (54). Especially

509   relevant is that MLDCs serve as a place for fatty acid homeostasis (101). Additionally, MERCs

510   importantly play a role in lipid homeostasis and synthesis (102). MERCs are understood to

511   regulate ER stress (103), so increases in fatty acids, triggered by lipid droplet formation, can

512   increase MERC formation in the aged model. Conversely, we qualitatively showed a decrease in

513   wrappER, which is known to maintain lipid flux through a 3-organelle contact site that involves

514   peroxisomes (63). These wrappER sites, which contain sites of adhesion, regulate very-low-

515   density lipoproteins (104). The age-related loss of mitochondrial complexity may impair the

516   ability and relative surface area mitochondria in murine liver samples must form contact sites,

517   including MERCs, thus interfering with functions including lipid homeostasis. Indeed, past

518   reviews have suggested targeting MERCs in NAFLD due to the role of contact sites in glucose

519   and lipid metabolism (105). However, in the future, a more rigorous analysis of age-related

520   changes in contact sites within liver tissue is necessary.

521

522   *Sam50 as a Regulator of Liver Disease*

523   Due to the observed dysfunctions in mitochondrial function and structure, we

524   investigated the role of *SAM50*, part of the SAM required for β-barrel protein assembly, in the

liver (80). There is evidence that the IMM proteins MIC60 (Mitofilin) and CHCHD3 were part

of a protein complex that includes the SAM components, known as the mitochondrial

intermembrane space bridging (MIB) complex (80, 106). The preservation of cristae depends on

interactions between the MIB complex's OMM and IMM components. The destabilization of the

MIB complex led to a decrease in the respiratory chain complex assembly. SAM50 depletion

over time affects the protein contents of all large respiratory complexes containing subunits

encoded by the mitochondria, suggesting a link between SAM50 and cristae structural integrity,

respiratory complex assembly, and mitochondrial DNA (mtDNA) maintenance (107). Our

previous study have shown that *Sam50* plays a significant role in the maintenance of both the

mitochondrial network and the structure of cristae and MICOS, especially in the context of liver

injury (52). Mice with *Sam50* deletion or liver-specific deletion produced by acetaminophen, an

efficient antipyretic and analgesic, experience mtDNA release, which activates the cGAS-STING

pathway and causes liver inflammation (81). Here, using phenome-wide and clinical lab-wide

scans for *SAMM50* GReX in a medical biobank, we saw a significant association between

SAMM50 and multiple liver diseases including nonalcoholic or alcoholic liver disease and

cirrhosis. This suggested that the expression of *SAMM50* as a transcription factor may directly or

indirectly contribute to mitochondrial defects of the liver.

To corroborate the clinical consequences of *SAMM50* GReX in BioVU participants, we

queried publicly available databases including: PheWeb (108), Biobank Japan PheWeb (109),

and the FinnGen web browser (110). PheWeb reports genome-wide association results from the

UK Biobank population (n > 400,000). The top phenotype associations for the *SAMM50* locus

within PheWeb included: Chronic liver disease and cirrhosis (p=2.8 X 10^{-24}, cases=2,954), Other

chronic nonalcoholic liver disease (p=1.2e-18, cases =1,687), Alcoholic liver damage (p=1.4 X

548 10^{-11}, cases=836), Injury to other and unspecified nerves (p=2.2 X 10^{-10}, cases=269), Portal

549   hypertension (p=1.7 X 10^{-9}, cases=527), and Liver abscess and sequelae of chronic liver disease

550   (p=4.1 X 10^{-9}, cases=974). BioBank Japan PheWeb similarly reports genome-wide association

551   results from the BioBank Japan Project (n > 260,000). The top phenotype associations for the

552   *SAMM50* locus in BioBank Japan PheWeb included Cirrhosis (p=2.8 X 10^{-14}, cases=2,551), as

553   well as Aspartate transaminase (p=3.3e-87, cases=150,068), Alanine aminotransferase (p=3.0 X

554 10^{-71}, cases=150,545), and platelet count (p=6.5 X 10^{-38}, cases=148,623). The FinnGen web

555   browser reports genome-wide association studies from the FinnGen data (n=342,499). Top

556   results from FinnGen included several phenotypes related to liver diseases including cirrhosis

557   (p=9.6 X 10^{-54}, cases=3,548), diseases of liver (p=4.9e-49, cases=9,548), and nonalcoholic fatty

558   liver disease (p=5.7 X 10^{-40}, cases=1,908). This underscores the therapeutic potential of targeting

559   *SAMM50*.

560

561   *The MICOS Complex as a Regulator of Aging:*

562   In this study, we demonstrated the disruptions in many aspects of mitochondria structure

563   in both 2D and 3D analysis of mitochondria in aging liver. We also showed MICOS impairment

564   in the aging liver. Therefore, we investigated whether MICOS regulates mitochondrial calcium

565   influx and oxidative stress. Calcium influx in mitochondria can reflect cell viability (111, 112).

566   Mitochondria mediation of calcium can impact many cell processes, such as apoptosis, signaling,

567   and ATP production (113). In some cases, elevated mitochondria calcium uptake can occur

568   antecedent to mitochondrial swelling, which results in a pathway often leading to apoptosis

569   (114). However, there was a decline in calcium uptake as *CHCHD6* and *MIC60* were disrupted

570   (Figure 6), suggestive of impaired cellular calcium signaling, which has previously been linked

571 with ER stress and dysregulation of MERCs (113, 115). This provides a plausible disease link as

572 ER calcium release promotes mitochondrial dysfunction, inducing oxidative stress and

573 hepatotoxicity (116), with broader implications in NAFLD (117, 118). Since Miro clusters both

574 interact with the MICOS complex, as well as regulate MERCs (119), Miro represents a potential

575 future mechanistic avenue through which MICOS complex contributes to MERC tethering.

576   Mitochondria redox state has been proposed as a principal moderator of mitochondrial

577 function in liver disease (120). Recent studies have continued to highlight a link between ROS

578 and mitochondrial dynamics (38). Lipids can aid in stimulating the production of ROS (121).

579 Additionally, many theories regarding aging have been formulated around the basis that

580 mitochondria lose function due to ROS byproducts which accumulate across aging (122). There

581 remains controversy about whether ROS are generated in the liver, while some studies have

582 found they arise during aging in the liver (123), other studies conversely found that in the aging

583 human liver, no superoxides are produced (124). Future studies should explore how ROS

584 contributes to the observed 3D mitochondrial phenotypes, as oxidative stress may be a regulator

585 of the phenotypes we observed. It is noteworthy that the loss of CHCHD6 and MIC60 leads to

586 oxidative stress (Figure 6). This suggests that during aging, the liver undergo a vicious cycle

587 wherein abnormal mitochondrial structures generate more harmful byproducts. These

588 byproducts, in turn, worsen mitochondrial structural dysfunction, contributing to age-related

589 oxidative stress.

590

591 *The Interaction Effect of the SAM Complex and MICOS Complex in Aging and Diet:*

592   In aged murine liver, decreases in the mtDNA numbers, urea cycle, mitochondrial

593 function, locomotor activity, loss of enzymatic activity, and anti-oxidative stress are well

594 established (123, 125, 126). Mitochondria also show diminished membrane potential across

595 aging in murine liver (127) and declined mitophagy responses (128). Within our study, we

596 looked at *Sam50* in the context of aging. Although we did not observe a change in the protein

597 level of SAM50 in the aging liver, there were defects in SAM50, but not MIC60, protein

598 production during the HFD. Notably, our study contrasts previous quantitative proteomic

599 analyses of liver mitochondria from HFD Diabetic mice, which found that Mitofilin, Sam50, and

600 Chchd3 are all upregulated (129). While more research is warranted to resolve this controversy,

601 our findings suggest that *Sam50* may not regulate mitochondrial machinery during aging

602 directly, but future studies performing an RNA sequencing scan of aged liver tissue, compared to

603 young tissue, will provide some candidates that interact with Sam50.

604 We know that HFDs are linked with many liver complications. For example, oxidative

605 stress, inflammation, and lipogenesis are some of the factors that might be linked to HFD and

606 exacerbate the onset of non-alcoholic fatty liver disease (83). Additionally, HFD is associated

607 with metabolic dysregulation of the liver (130). In this study, we found an increase in retinoic

608 acid and retinal in the aged liver (Figure 4). Notably, there is evidence linking retinol-binding

609 proteins and HFD. The enhanced supply of vitamin A and high-fat consumption in diet-induced

610 obese mice is linked to the production of bisretinoid (131). Interestingly, obesity within

611 hepatocytes has also been shown to increase MAMs (i.e., MERC-isolated biochemical fractions),

612 which confers mitochondrial dysfunction (132, 133). High fat within a diet may further have an

613 additive effect with aging to exacerbate age-related hepatic pathologies. For example, an HFD

614 can alter mitochondrial structure, leading to decreased fatty acid oxidation (134). Within age-

615 related hepatic steatosis, impaired fatty acid oxidation, and other factors that HFD can contribute

616 to such as insulin resistance, are key risk factors (30). Notably, *SAMM50* deficiency causes lipid

617 accumulation and impaired fatty acid oxidation (49). Thus, HFDs can play a role in metabolism

618 defects in the liver tissue, although this must be further investigated in the future.

619 The changes in SAM50 during HFD may be due to several factors. Our lipidomic studies

620 confirm suggested dysregulations of lipids in the aged liver (Figure 4). Our observation of

621 SAM50 decreased in HFD contrasts *in vitro* models of hepatic cellular steatosis, revealing that

622 mitochondrial dysfunction is determined by oxidative stress instead of lipid buildup (135). This

623 finding is novel since *Sam50*-deletion or overexpression, and the broader SAM complex, during

624 HFD in the liver, have not been studied. However, past studies of HFD offer some insight. HFD

625 can lead to pathological alterations and damage to the ultrastructure of the mitochondria and

626 downregulation of MERCs regulators, MFN2 and OPA1 (84), which parallels the TEM

627 ultrastructural changes we observed. Additionally, *Mic19/Chchd3* was shown to be necessary in

628 the liver (136). *Sam50* is closely related to *Mic19*. *Sam50* mediates mitochondrial outer and inner

629 membrane interactions via the Sam50-Mic19-Mic60 axis (79). In the liver, *Mic19/Chchd3*

630 deletion reduces ER-mitochondrial contacts, disrupts mitochondrial lipid metabolism,

631 disorganizes mitochondrial cristae, and causes unfolded protein stress response in mouse

632 hepatocytes, leading to impairments of liver mitochondrial fatty acid β-oxidation and lipid

633 metabolism (136). This may explain our results of declined *Sam50* synthesis during HFD, but it

634 remains unclear why *Mic60* is not affected. Alternatively, linking MICOS and SAM complex

635 changes, in hepatocytes both SAM50 and MIC60 cooperate to bind to cardiolipin, a phospholipid

636 exclusively located in mitochondria (81). ROS also targets cardiolipin, which is associated with

637 mitochondrial dysfunction in NAFLD (137). Cardiolipin inhibition has mitigated non-alcoholic

638 steatohepatitis through NLRP3 inflammasome activation, suggesting a therapeutic role of

639 cardiolipin (138). *Sam50* depletion leads to cardiolipin externalization and subsequent liver

inflammation (81). Interestingly, paralleling our findings of increased lipogenesis with aging

(Figure 1), our lipidomic analysis showed altered lipogenesis across multiple lipid classes, which

has implications extending beyond cardiolipin, since the liver is a principal site for lipid

homeostasis (139). Thus, while aging exhibits a protective mechanism to maintain *Sam50*, the

additional challenge of diet changes may cause loss of integrity in both the SAM complex and

MICOS complex, which results in the buildup of lipids in the liver, raising the risk of cirrhosis,

steatohepatitis, and NAFLD, among other hepatic conditions.

CONCLUSION

To broaden our understanding of age-related changes in male murine hepatic tissue, we have

established how 2D and 3D ultrastructure undergo similar reductions in mitochondrial volume

and connectivity, potentially due to an age-related loss of the MICOS complex. This is an

important age-related regulator of mitochondrial structure and could be linked to functional

changes across aging in the liver. Deletion of the MICOS complex in HepG2 cells results in

impaired calcium uptake and increased oxidative stress, highlighting its significance. We have

also established that *Sam50*, which is genetically associated with liver diseases, is maintained

across aging but lost with a high-fat diet. Since a high-fat diet can exacerbate changes in lipid

synthesis, catabolism, and mitochondrial alterations that already occur in hepatic aging (30, 83),

this suggests the confluence of aging and a high-fat diet in increasing susceptibility to liver

disease, in part due to the loss of both the MIB complex (i.e., MICOS and SAM complex).

Together, this study demonstrates dysfunctions of mitochondrial structure in both 2D and 3D,

mitochondrial calcium influx and ROS production, changes in metabolism and lipid content, and

MICOS impairments in the aging liver, with alterations in *Sam50* expression contributing to liver

diseases.

EXPERIMENTAL PROCEDURES

Sex As a Biological Variable:

In this study, the consideration of sex as a biological variable was integral to our experimental

design. All experimental procedures and analyses were conducted with careful consideration of

sex differences. Since minimal sex-dependent differences were observed in TEM studies, we

proceeded with a male murine model for all other studies. Sex differences in aging-related

changes and longevity can affect relative mitochondrial morphology which is a limitation of our

male model.

Human Cohort

All human samples were obtained from Brazilian cohorts according to the CAEE (Ethics

Appreciation Presentation Certificate) guidelines. Samples from young and old individuals were

collected and analyzed under CAEE number 77570224.2.0000.5281. Fat percentage was

calculated as a function of in-phase and out-phase quantifications via the following calculator:

https://www.ricardoromano.com/esteatoseRM (53).

Animal Care and Maintenance:

Per protocols previously described (140), the care and maintenance of the male C57BL/6J mice

conformed to the National Institute of Healt's guidelines for the use of laboratory animals. The

University of Iow's Institutional Animal Care and Use Committee (IACUC) or University of

Washington IACUC approved the housing and feeding of these mice. Anesthesia was achieved

using a mixture of 5% isoflurane and 95% oxygen.

Oil Red O

OCT blocks were cut into 7 μm thick sections, affixed to glass slides, brought to room

temperature for 10 minutes, then stained with Oil Red O (Sigma-Aldrich), as previously

described (141).

mtDNA Content

As previously described (142), mitochondrial DNA content was quantified using real-time PCR

(RT-PCR) from DNA extracted and purified from gastrocnemius muscle tissue. Five nanograms

of DNA were used for the quantification of mitochondrial (Cox1) and nuclear (β-actin) DNA

markers. Mitochondrial DNA content was normalized to the genomic Rpl13a gene using specific

primers for Cox1 and Rpl13a.

Bile Acid

From frozen liver tissue, 100 mg of tissue homogenized in 75% ethanol was incubated for 2

hours and then centrifuged at 6000g for 10 minutes. Once prepared, bile was measured with

Mouse Total Bile Acids Assay Kit (Crystal Chem), per manufacturer's instruction.

Triglyceride Levels

As previously described (142, 143), triglyceride levels were measured in the liver and in serum

collected after a 6□h fast using the EnzyChrom™ Triglyceride Assay Kit (BioAssay Systems),

with triglycerides extraction using a solution of isopropanol and Triton X□100.

Quantification of TEM Micrographs and Parameters Using ImageJ

Samples were fixed in a manner to avoid any bias, per established protocols (144). Following

preparation, tissue was embedded in 100% Embed 812/Araldite resin with polymerization at 60

°C overnight. After ultrathin sections (90–100 nm) were collected, they were post-stained with

lead citrate and imaged (JEOL 1400+ at 80 kV, equipped with a GatanOrius 832 camera). The

National Institutes of Health (NIH) *ImageJ* software was used for quantification of TEM images,

as described previously (9, 145).

Serial Block-Face Scanning Electron Microscope (SBF-SEM) Processing of Mouse Muscle

Fibers

SBF-SEM was performed according to previously defined protocols (19, 144, 145). Anesthesia

was induced in male mice using 5% isoflurane. Post skin and hair removal, the liver was treated

with 2% glutaraldehyde in 100 mM phosphate buffer for 30 minutes, dissected into 1-mm³

cubes, and further fixed in a solution containing 2.5% glutaraldehyde, 1% paraformaldehyde, and

120 mM sodium cacodylate for 1 hour.

Fixation and subsequent steps collected onto formvar-coated slot grids (Pella, Redding CA),

stained and imaged as previously described (19, 144, 145). This includes tissue washing with

100 mM cacodylate buffer, incubation in a mixture of 3% potassium ferrocyanide and 2%

osmium tetroxide, followed by dehydration in an ascending series of acetone concentrations. The

tissues were then embedded in Epoxy Taab 812 hard resin. Sectioning and imaging of sample

was performed using a VolumeScope 2 SEM (Thermo Fisher Scientific, Waltham, MA).

Conventional TEM analysis was performed on 300–400 serial sections from each sample,

following staining and imaging protocols. Subsequently, analyzed, via imaging was performed

under low vacuum/water vapor conditions with a starting energy of 3.0 keV and beam current of

0.10 nA. Sections of 50 nm thickness were cut allowing for imaging at 10 nm $\times$ 10 nm $\times$ 50 nm

spatial resolution.

Segmentation and Quantification of 3D SBF-SEM Images Using Amira

SBF-SEM images were manually segmented in Amira to perform 3D reconstruction, as

described previously (19). 300-400 slices were used and analyzed by a blind individual. 250 total

mitochondria across from 3 mice were collected for each quantification. For 3D reconstruction of

cardiomyocytes, 10 cells and a total of about 200 mitochondria. Quantification of 3D structures

was performed using the Amira software with built-in parameters or previously described

measurements (19).

LCMS Methods for Metabolomics:

Frozen tissues were weighed, ground with a liquid nitrogen in a cryomill (Retsch) at 25 Hz for

45 seconds, before extracting tissues 40:40:20 acetonitrile: methanol: water +0.5% FA +15%

NH4HCO3 (146) with a volume of 40mL solvent per 1mg of tissue, vortexed for 15 seconds, and

incubated on dry ice for 10 minutes. Tissue samples were then centrifuged at 16,000 g for 30

752 minutes. The supernatants were transferred to new Eppendorf tubes and then centrifuged again at

753 16,000 g for 25 minutes to remove any residual debris before analysis.

754

755 Extracts were analyzed within 24 hours by liquid chromatography coupled to a mass

756 spectrometer (LC-MS). The LC–MS method was based on hydrophilic interaction

757 chromatography (HILIC) coupled to the Orbitrap Exploris 240 mass spectrometer (Thermo

758 Scientific) (147). The LC separation was performed on a xBridge BEH Amide column (2.1 x 150

759 mm, 3.5 μm particle size, Waters, Milford, MA). Solvent A is 95%: 5% H2O: acetonitrile with

760 20 mM ammonium acetate and 20mM ammonium hydroxide, and solvent B is 90%: 10%

761 acetonitrile: H2O with 20 mM ammonium acetate and 20mM ammonium hydroxide. The

762 gradient was 0 min, 90% B; 2 min, 90% B; 3 min, 75% B; 5 min, 75% B; 6 min, 75% B; 7 min,

763 75% B; 8 min, 70% B; 9 min, 70% B; 10 min, 50% B; 12 min, 50% B; 13 min, 25% B; 14min,

764 25% B; 16 min, 0% B; 18 min, 0% B; 20 min, 0% B; 21 min, 90% B; 25 min, 90% B. The

765 following parameters were maintained during the LC analysis: flow rate 150 mL/min, column

766 temperature 25 °C, injection volume 5 μL and autosampler temperature was 5 °C. For the

767 detection of metabolites, the mass spectrometer was operated in both negative and positive ion

768 mode. The following parameters were maintained during the MS analysis: resolution of 180,000

769 at m/z 200, automatic gain control (AGC) target at 3e6, maximum injection time of 30 ms and

770 scan range of m/z 70-1000. Raw LC/MS data were converted to mzXML format using the

771 command line "msconvert" utility (148). Data were analyzed via the EL-MAVEN software

772 version 12.

773

774 *LCMS Methods for Lipidomic Profiling*

Tissue homogenization and extraction for lipids: Tissues were homogenized using a Retsch

CryoMill. The homogenate was mixed with 1 mL of Extraction Buffer containing

IPA/H2O/Ethyl Acetate (30:10:60, v/v/v) and Avanti Lipidomix Internal Standard (diluted

1:1000) (Avanti Polar Lipids, Inc. Alabaster, AL). Samples were vortexed and transferred to

bead mill tubes for homogenization using a VWR Bead Mill at 6000 g for 30 seconds, repeated

twice. The samples were then sonicated for 5 minutes and centrifuged at 15,000 g for 5 minutes

at 4°C. The upper phase was transferred to a new tube and kept at 4°C. To re-extract the tissues,

another 1 mL of Extraction Buffer (30:10:60, v/v/v) was added to the tissue pellet-containing

tube. The samples were vortexed, homogenized, sonicated, and centrifuged as described earlier.

The supernatants from both extractions were combined, and the organic phase was dried under

liquid nitrogen gas.

Sample reconstitution for lipids: The dried samples were reconstituted in 300 µL of Solvent A

(IPA/ACN/H2O, 45:35:20, v/v/v). After brief vortexing, the samples were sonicated for 7

minutes and centrifuged at 15,000 g for 10 minutes at 4°C. The supernatants were transferred to

clean tubes and centrifuged again for 5 minutes at 15,000 g at 4°C to remove any remaining

particulates. For LC-MS lipidomic analysis, 60 µL of the sample extracts were transferred to

mass spectrometry vials.

LC-MS analysis for lipids: Sample analysis was performed within 36 hours after extraction using

a Vanquish UHPLC system coupled with an Orbitrap Exploris 240™ mass spectrometer

equipped with a H-ESI™ ion source (all Thermo Fisher Scientific). A Waters (Milford, MA)

CSH C18 column (1.0 × 150 mm × 1.7 µm particle size) was used. Solvent A consisted of

ACN:H$_2$O (60:40; v/v) with 10 mM Ammonium formate and 0.1% formic acid, while solvent B

contained IPA:ACN (95:5; v/v) with 10 mM Ammonium formate and 0.1% formic acid. The

mobile phase flow rate was set at 0.11 mL/min, and the column temperature was maintained at

65 °C. The gradient for solvent B was as follows: 0 min 15% (B), 0–2 min 30% (B), 2–2.5 min

48% (B), 2.5–11 min 82% (B), 11–11.01 min 99% (B), 11.01–12.95 min 99% (B), 12.95–13 min

15% (B), and 13–15 min 15% (B). Ion source spray voltages were set at 4,000 V and 3,000 V in

positive and negative mode, respectively. Full scan mass spectrometry was conducted with a

scan range from 200 to 1000 m/z, and AcquireX mode was utilized with a stepped collision

energy of 30% with a 5% spread for fragment ion MS/MS scan.

RNA Extraction and RT-qPCR

Using TRIzol reagent (Invitrogen), total RNA was isolated from tissues and further purified with

the rNeasy kit (Qiagen Inc). RNA concentration was determined by measuring absorbance at 260

nm and 280 nm using a NanoDrop 1000 spectrophotometer (NanoDrop products, Wilmington,

DE, USA). Approximately 1 µg of RNA was reverse-transcribed using a High-Capacity cDNA

Reverse Transcription Kit (Applied Biosciences, Carlsbad CA). Quantitative PCR (qPCR) was

then performed using SYBR Green (Life Technologies, Carlsbad, CA) (149). For qPCR, 50 ng

of cDNA was loaded into each well of a 384-well plate, with the reaction carried out on an ABI

Prism 7900HT system (Applied Biosystems) with the following cycle: 1 cycle at 95°C for 10

min; 40 cycles of 95°C for 15 s; 59°C for 15 s, 72°C for 30 s, and 78°C for 10 s; 1 cycle of 95°C

for 15 s; 1 cycle of 60°C for 15 s; and one cycle of 95°C for 15 s. GAPDH normalization was

used to present the data as fold changes. qPCR primers used were from previously published

sequences (142), as detailed in Table 1.

821 Table 1: qPCR Primers Used

Gene	Primers	
Opa1	Forward	5'-ACCAGGAGACTGTGTCAA-3'
	Reverse	5'-TCTTCAAATAAACGCAGAGGTG-3'
Chchd3	Forward	5'-GAAAAGAATCCAGGCCCTTCCACGCGC-3'
	Reverse	5'-CAGTGCCTAGCACTTGGCACAACCAGGAA-3'
Chchd6	Forward	5'-CTCAGCATGGACCTGGTAGGCACTGGGC-3'
	Reverse	5'-GCCTCAATTCCCACATGGAGAAAGTGGC-3'
Mitofilin	Forward	5'-CCTCCGGCAGTGTTCACCTAGTAACCCCTT-3'
	Reverse	5'-TCGCCCGTCGACCTTCAGCACTGAAAACCTAT-3'

822

823

824 *Western Blotting*

825 Western blotting was performed as previously described (76). Briefly, following RIPA lysis

826 buffer (1% NP40, 150 mM NaCl, 25 mM Tris base, 0.5% sodium deoxycholate, 0.1% SDS,

827 1% phosphatase inhibitor cocktails #2 (Sigma P5726-1ML) and #3 (Sigma P0044-1ML), and

828 one cOmplete protease inhibitor tablet (Sigma 04693159001)), 3-month and 2-year tissue

829 samples were quantified using a BCA Assay (Thermo Scientific VLBL00GD2). Equal amounts

830 of proteins were run on 4%–20% Tris-glycine gels (Invitrogen WXP42012BOX). Protein was

831 then transferred to a nitrocellulose membrane (Li-Cor 926-31092) that was incubated with

832 primary antibodies overnight at 4°C: Mic60/mitofilin (Abcam ab110329), SAM50 (Proteintech

833 20824-1-AP), or tubulin (Novus NB100-690). Secondary antibodies [1:10,000; donkey anti-

834 mouse IgG (H+L) (Invitrogen A32789) and donkey anti-rabbit IgG (H+L) (Invitrogen

A32802)] were incubated with the membrane at room temperature for 1 h. Using the Li-Cor

Odyssey CLx infrared imaging system, blots were imaged.

Confocal mCherry-Mito-7 Labeling

To label the mitochondria of cardiac fibroblasts, the mCherry-Mito-7 plasmid was transfected

into the cells using a transfection reagent according to the manufacturer's instructions (150) and

as previously described (75). Briefly, following plasmid and transfection reagent dilution in

Opti-MEM medium and incubation at room temperature for 20 minutes, the dilution was added

to the culture medium of the cells, which were incubated for 24–48 hours to allow expression of

the mCherry-Mito-7 protein. Localization in fibroblasts was visualized using a Leica SP8

Confocal Microscope. Cells were washed with PBS, fixed with 4% paraformaldehyde for 10

minutes, and mounted with DAPI-containing mounting medium. Fluorescent signals were

observed using appropriate filters and recorded with a digital camera.

SAMM50 Analyses in Vanderbilt University Medical Center's biobank, BioVU

To examine the clinical correlates of SAMM50 gene expression in a human population, we

leveraged the BioVU biobank at Vanderbilt University Medical Center (151, 152). Genotype

data linked to deidentified electronic health records (EHR) for 85,615 individuals were assessed.

Genotype data were collected on Illumina's Multi-Ethnic Genotyping Array (MEGA) and

underwent quality control procedures as previously described (153). SAMM50 genetically-

regulated gene expression (GReX) was calculated across all genotyped individuals incorporating

training data from GTEX version 8 and utilizing PrediXcan, UTMOST, or JTI models based on

the best performance (highest performance r2 for each gene-tissue pair) **(154–157)**. MultiXcan

was then performed to collapse individual tissue results into a single cross-tissue SAMM50

GReX model (158). The BioVU population was then stratified by genetic ancestry as previously

described and within each group, SAMM50 GReX was tested for association with clinical

phenotypes mapped from ICD9/10 codes and clinical lab values extracted from the EHR using

the previously described QualityLabs pipeline (153, 159, 160). In total, we tested 1,704

phenotypes and 326 labs across 70,440 individuals of European ancestry and 1,361 phenotypes

and 265 labs across 15,175 individuals of African ancestry. PheWAS analyses included logistic

regressions for any mapped phenotype with at least 50 cases, requiring at least 2 instances of an

ICD9/10 code on unique dates to be labeled as a case. LabWAS analyses included linear

regressions for any lab values that met the QualityLabs QC criteria and had median

measurements from at least 50 individuals. Covariates for both PheWAS and LabWAS included:

principal components (PCs) 1-10 for genetic ancestry, sex, age, median age of medical record,

and genotyping batch.

Knockdown of MIC60 and CHCHD6 in HepG2 cells.

The transfection of MIC60 and CHCHD6 siRNAs into HepG2 cells was carried out using

Lipofectamine RNAiMax (Invitrogen) in accordance with the manufacturer's instructions and as

previously described (24). Following a 48-hour incubation period, the cells were utilized for

mitochondrial calcium ($_mCa^{2+)}$ and ROS measurements.

Measurement of mitochondrial calcium uptake and retention capacity in HepG2 cells.

$_mCa^{2+}$ uptake retention capacity in HepG2 cells were assessed using a ratiometric Ca^{2+} sensor

Fura-FF, as detailed earlier (161), with slight modifications. In brief, cells (2.5x106) were

881   washed with $Ca^{2+/Mg2+}$-free DPBS (GIBCO), permeabilized in intracellular medium (ICM:

882   120 mM KCl, 10 mM NaCl, 1 mM KH2PO4, 20 mM HEPES-Tris, pH 7.2), and

883   supplemented with thapsigargin and succinate. Fura-FF (1 μM) was added at the 0 s time point,

884   and fluorescence emissions at 340- and 380-nm ex/510-nm em were monitored using a multi-

885   wavelength excitation dual-wavelength emission fluorimeter (Delta RAM, PTI). To assess the

886 $_mCa^{2+}$ uptake, a bolus of 5 μM Ca^{2+} and the mitochondrial uncoupler FCCP (10 μM) were added

887   at specified time points with continuous stirring at 37°C. To assess mCa^{2+} retention capacity,

888   following baseline recordings, a series of Ca^{2+} boluses (5 μM) were introduced at specified time

889   points. Upon reaching a steady state, 10 μM FCCP was added to collapse the $\Delta\psi m$ and release

890   matrix free-Ca^{2+}. The number of Ca^{2+} boluses taken up by cells was counted to calculate

891   mitochondrial CRC.

892

893   *Evaluation of ROS levels*

894   ~0.2 million HepG2 cells were plated in 35 mm dishes. The next day, MIC60 and CHCHD6

895   siRNAs were transfected using Lipofectamine RNAiMax (Invitrogen) according to the

896   manufacturing instructions. After incubation for 30 hrs, cells were co-stained for 30 min at 37°C

897   with two different dyes for ROS detection: MitoBright ROS Deep Red (10 μM, Dojindo

898   Laboratories) for mitochondrial superoxide, and DCFDA (10 µM, Invitrogen) for intracellular

899   total ROS. Following the incubation with staining dyes, cells were washed three times with 1X

900   HBSS and ROS analysis was done using a confocal microscope (FV4000, Olympus Life

901   Science).

902   For mitochondrial H_2O_2 assessment, cells were stained with MitoPY1 (5 µM, Bio-Techne) for 45

903   min at 37°C. Cells were then washed with 1x HBSS and imaged using a confocal microscope

(FV4000, Olympus Life Science). ImageJ was used for the quantification of fluorescence

intensities.

Data Analysis

GraphPad Prism 10.2.3 (La Jolla, CA, USA), was used for all statistical analysis. All

experiments involving SBF-SEM and TEM data had at least three independent experiments.

Statistics were not handled by those conducting the experiments. The black bars represent the

standard error of the mean. For all analysis, one-way ANOVA was performed with tests against

each independent group and significance was assessed using Fisher's protected least significant

difference (LSD) test. *, **, ***, **** were set to show significant difference, denoting $p <$

0.05, $p < 0.01$, $p < 0.001$, and $p < 0.0001$, respectively.

AUTHOR CONTRIBUTIONS:

Zer Vue, Alexandria Murphy, and Han Le contributed equally to this work. Zer Vue, Han Le, Kit

Neikirk, Edgar Garza-Lopez, Andrea G. Marshall, Margaret Mungai, Brenita Jenkins, Larry

Vang, Heather K. Beasley, Mariaassumpta Ezedimma, Sasha Manus, Aaron Whiteside, Chanel

Harris, Amber Crabtree, Claude F. Albritton, Sydney Jamison, Mert Demirci, Ashton Oliver,

Ky'Era V. Actkins, Elma Zaganjor, Estevão Scudese, Benjamin Rodriguez, Alice Koh, Izabella

Rabago, Johnathan Moore, Desiree Nguyen, Muhammad Aftab, Benjamin Kirk, Yahang Li,

Nelson Wandira, Taseer Ahmed, Mohammad Saleem, Ashlesha Kadam, Prasanna Katti, Ho-Jin

Koh, Chantell Evans, Young Do Koo, Eric Wang, Quinton Smith, Dhanendra Tomar, Clintoria

R. Williams, Mariya Sweetwyne, Anita M. Quintana, Mark A. Phillips, David Hubert, Annet

Kirabo, Chandravanu Dash, Pooja Jadiya, André Kinder, Olujimi A. Ajijola, Tyne W. Miller-

Fleming, Melanie R. McReynolds, and Antentor Hinton, Jr., contributed to the conception,

design, data acquisition, analysis, and interpretation of data. Melanie R. McReynolds, and

Antentor Hinton, Jr. conceived and supervised this study. All authors were involved in drafting

and critically revising the manuscript for important intellectual content. All authors approved the

final version of the manuscript.

ACKNOWLEDGEMENTS

All authors have no competing interests.

This project was funded by the National Institute of Health (NIH) NIDDK T-32, number

DK007563 entitled Multidisciplinary Training in Molecular Endocrinology to Z.V.; National

Institute of Health (NIH) NIDDK T-32, number DK007563 entitled Multidisciplinary Training

in Molecular Endocrinology to A.C.; NSF MCB #2011577I to S.A.M.; The UNCF/Bristol-Myers

Squibb E.E. Just Faculty Fund, Career Award at the Scientific Interface (CASI Award) from

Burroughs Welcome Fund (BWF) ID # 1021868.01, BWF Ad-hoc Award, NIH Small Research

Pilot Subaward to 5R25HL106365-12 from the National Institutes of Health PRIDE Program,

DK020593, Vanderbilt Diabetes and Research Training Center for DRTC Alzheimer's Disease

Pilot & Feasibility Program. CZI Science Diversity Leadership grant number 2022- 253529 from

the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation

to A.H.J. and O.A.A.; and National Institutes of Health grant HD090061 and the Department of

Veterans Affairs Office of Research Award I01 BX005352 to J.G. Howard Hughes Medical

Institute Hanna H. Gray Fellows Program Faculty Phase (Grant# GT15655 awarded to M.R.M);

and Burroughs Wellcome Fund PDEP Transition to Faculty (Grant# 1022604 awarded to

M.R.M). National Institutes of Health Grants: R21DK119879 (to C.R.W.) and R01DK-133698

(to C.R.W.), American Heart Association Grant 16SDG27080009 (to C.R.W.) and by an

American Society of Nephrology KidneyCure Transition to Independence Grant (to C.R.W.).

NIH Grants R01HL147818, R03HL155041, and R01HL144941 (A. Kirabo). NIH Grant

R00DK120876 (D.T.), Harold S. Geneen Charitable Trust Awards Program (D.T.), Alzheimer's

Association AARG-NTF-23-1144888 (D.T.). NIH Grant R00AG065445 (P.J.), Alzheimer's

Association 24AARG-D-1191292 (P.J.), Wake ADRC REC and Development grant

P30AG072947 (P.J.). American Heart Association Grant 23POST1020344 (A.K.). American

Heart Association Grant 23CDA1053072 (M. S.). NIH K01AG062757 to (M.T.S.) The BioVU

project at VUMC is supported by numerous sources: including the NIH funded Shared

Instrumentation Grant S10OD017985 and S10RR025141; CTSA grants UL1TR002243,

UL1TR000445, and UL1RR024975 from the National Center for Advancing Translational

Sciences. Its contents are solely the responsibility of the authors and do not necessarily represent

official views of the National Center for Advancing Translational Sciences. Genomic data are

also supported by investigator-led projects that include U01HG004798, R01NS032830,

RC2GM092618, P50GM115305, U01HG006378, U19HL065962, R01HD074711; and

additional funding sources listed at https://victr.vumc.org/biovu-funding/.23CDA1053072

(M.S.). Its contents are solely the responsibility of the authors and do not necessarily represent

the official view of the NIH. The funders had no role in study design, data collection, and

analysis, decision to publish, or preparation of the manuscript. We would also like to

acknowledge the Huck Institutes' Metabolomics Core Facility (RRID:SCR_023864) for use of

the OE 240 LCMS and Drs. Imhoi Koo, Ashley Shay, and Sergei Koshkin for helpful

discussions on sample preparation and analysis. We would also like to thank UCLA investigators

for gifting us old and young human liver samples.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

References:

1. McBride HM, Neuspiel M, Wasiak S. Mitochondria: more than just a powerhouse. *Curr*

***Biol CB*. 2006;16(14):R551-560.**

2. Glancy B, et al. The Functional Impact of Mitochondrial Structure Across Subcellular

Scales. *Front Physiol*. 2020;11.

https://www.frontiersin.org/articles/10.3389/fphys.2020.541040. Accessed December 16,

2022.

3. Picard M, et al. Mitochondria: isolation, structure and function. *J Physiol*.

2011;589(18):4413–4421.

4. Chan DC. Fusion and fission: interlinked processes critical for mitochondrial health.

***Annu Rev Genet*. 2012;46:265–287.**

5. Li M, et al. Mitochondrial Fusion Via OPA1 and MFN1 Supports Liver Tumor Cell

Metabolism and Growth. *Cells*. 2020;9(1):121.

6. Seo AY, et al. New insights into the role of mitochondria in aging: mitochondrial

dynamics and more. *J Cell Sci*. 2010;123(15):2533–2542.

7. Bratic A, Larsson N-G. The role of mitochondria in aging. *J Clin Invest*.

2013;123(3):951–957.

8. Haas RH. Mitochondrial Dysfunction in Aging and Diseases of Aging. *Biology*.

2019;8(2):48.

9. Lam J, et al. A Universal Approach to Analyzing Transmission Electron Microscopy

with ImageJ. *Cells*. 2021;10(9):2177.

10. Collins HE, et al. Mitochondrial Morphology and Mitophagy in Heart Diseases:

Qualitative and Quantitative Analyses Using Transmission Electron Microscopy. *Front*

Aging. 2021;2. https://www.frontiersin.org/articles/10.3389/fragi.2021.670267. Accessed

September 8, 2022.

11. Cogliati S, Enriquez JA, Scorrano L. Mitochondrial cristae: where beauty meets

functionality. *Trends Biochem Sci*. 2016;41(3):261–273.

12. Jenkins BC, et al. Mitochondria in disease: changes in shapes and dynamics. *Trends*

Biochem Sci. 2024;49(4):346–360.

13. Long Q, et al. Modeling of Mitochondrial Donut Formation. *Biophys J*. 2015;109:892–9.

14. Liu X, Hajnóczky G. Altered fusion dynamics underlie unique morphological changes

in mitochondria during hypoxia–reoxygenation stress. *Cell Death Differ*. 2011;18(10):1561–

1572.

15. Vincent AE, et al. Mitochondrial Nanotunnels. *Trends Cell Biol*. 2017;27(11):787–799.

16. Tyumentsev MA, et al. Mitochondria with Morphology Characteristic for Alzheimer's

Disease Patients Are Found in the Brain of OXYS Rats. *Biochem Mosc*. 2018;83(9):1083–

1088.

17. Wakabayashi T. Megamitochondria formation □ physiology and pathology. *J Cell Mol*

Med. 2002;6(4):497–538.

18. Gautier MK, Ginsberg SD. A method for quantification of vesicular compartments

within cells using 3D reconstructed confocal z-stacks: Comparison of ImageJ and Imaris to

count early endosomes within basal forebrain cholinergic neurons. *J Neurosci Methods*.

2021;350:109038.

19. Garza-Lopez E, et al. Protocols for Generating Surfaces and Measuring 3D Organelle

Morphology Using Amira. *Cells*. 2022;11(1):65.

20. Courson JA, et al. Serial Block-Face Scanning Electron Microscopy (SBF-SEM) of

Biological Tissue Samples. *J Vis Exp JoVE*. 2021;(169):10.3791/62045.

21. Marshall AG, et al. Serial Block Face-Scanning Electron Microscopy as a Burgeoning

Technology. *Adv Biol*. 2023;e2300139.

22. Russell MRG, et al. 3D correlative light and electron microscopy of cultured cells using

serial blockface scanning electron microscopy. *J Cell Sci*. 2017;130(1):278–291.

23. Vue Z, et al. 3D reconstruction of murine mitochondria reveals changes in structure

during aging linked to the MICOS complex. *Aging Cell*. 2023;22(12):e14009.

24. Vue Z, et al. The MICOS Complex Regulates Mitochondrial Structure and Oxidative

Stress During Age-Dependent Structural Deficits in the Kidney [preprint].

2024;2024.06.09.598108.

25. Gagliano N, Grizzi F, Annoni G. Mechanisms of Aging and Liver Functions. *Dig Dis*.

2007;25(2):118–123.

26. Hunt NJ, et al. Hallmarks of Aging in the Liver. *Comput Struct Biotechnol J*.

2019;17:1151–1161.

27. Kalra A, et al. Physiology, Liver. *StatPearls*. Treasure Island (FL): StatPearls

Publishing; 2024.

28. Reddy SR, et al. Outcomes of Young Patients With Alcoholic Cirrhosis After First

Hospitalization for Cirrhosis: A Carilion Clinic Experience. *Cureus*;13(7):e16695.

29. Kim H, Kisseleva T, Brenner DA. Aging and liver disease. *Curr Opin Gastroenterol*.

2015;31(3):184–191.

30. Gong Z, et al. Hepatic lipid metabolism and non-alcoholic fatty liver disease in aging.

Mol Cell Endocrinol. 2017;455:115–130.

31. Chung KW. Advances in Understanding of the Role of Lipid Metabolism in Aging.

Cells. 2021;10(4):880.

32. Hoffmann JJ, Becker T. Crosstalk between Mitochondrial Protein Import and Lipids.

Int J Mol Sci. 2022;23(9):5274.

33. Chen G, Kroemer G, Kepp O. Mitophagy: An Emerging Role in Aging and Age-

Associated Diseases. *Front Cell Dev Biol*. 2020;8.

https://www.frontiersin.org/articles/10.3389/fcell.2020.00200. Accessed December 21, 2022.

34. Nishikawa T, et al. A switch in the source of ATP production and a loss in capacity to

perform glycolysis are hallmarks of hepatocyte failure in advance liver disease. *J Hepatol*.

2014;60(6):1203–1211.

35. Middleton P, Vergis N. Mitochondrial dysfunction and liver disease: role, relevance,

and potential for therapeutic modulation. *Ther Adv Gastroenterol*.

2021;14:17562848211031394.

36. Wang D, et al. Research progress on the mitochondrial mechanism of age-related non-

alcoholic fatty liver. *World J Gastroenterol*. 2023;29(13):1982–1993.

37. Shutt TE, McBride HM. Staying cool in difficult times: Mitochondrial dynamics,

quality control and the stress response. *Biochim Biophys Acta BBA - Mol Cell Res*.

2013;1833(2):417–424.

38. Ježek J, Cooper KF, Strich R. Reactive Oxygen Species and Mitochondrial Dynamics:

The Yin and Yang of Mitochondrial Dysfunction and Cancer Progression. *Antioxidants*.

2018;7(1). https://doi.org/10.3390/antiox7010013.

39. Abrisch RG, et al. Fission and fusion machineries converge at ER contact sites to

regulate mitochondrial morphology. *J Cell Biol*. 2020;219(4):e201911122.

40. Youle RJ, van der Bliek AM. Mitochondrial Fission, Fusion, and Stress. *Science*.

2012;337(6098):1062–1065.

41. Belenguer P, Pellegrini L. The dynamin GTPase OPA1: More than mitochondria?

Biochim Biophys Acta BBA - Mol Cell Res. 2013;1833(1):176–183.

42. Chen H, et al. Mitofusins Mfn1 and Mfn2 coordinately regulate mitochondrial fusion

and are essential for embryonic development. *J Cell Biol*. 2003;160(2):189–200.

43. Hu C, et al. OPA1 and MICOS Regulate mitochondrial crista dynamics and formation.

Cell Death Dis. 2020;11(10):1–17.

44. Favaro G, et al. DRP1-mediated mitochondrial shape controls calcium homeostasis and

muscle mass. *Nat Commun*. 2019;10(1):1–17.

45. Anand R, et al. The i-AAA protease YME1L and OMA1 cleave OPA1 to balance

mitochondrial fusion and fission. *J Cell Biol*. 2014;204(6):919–929.

46. Wai T, Langer T. Mitochondrial Dynamics and Metabolic Regulation. *Trends

Endocrinol Metab*. 2016;27(2):105–117.

47. Takeichi Y, et al. Non-alcoholic fatty liver disease in mice with hepatocyte-specific

deletion of mitochondrial fission factor. *Diabetologia*. 2021;64:2092–2107.

48. Galloway C, et al. Decreasing mitochondrial fission alleviates hepatic steatosis in a

murine model of nonalcoholic fatty liver disease. *Am J Physiol Gastrointest Liver Physiol*.

2014;307 6:632–41.

49. Li Z, et al. The role of SAMM50 in non-alcoholic fatty liver disease: from genetics to

mechanisms. *FEBS Open Bio*. 2021;11(7):1893–1906.

50. Lionello S, Marzaro G, Martinvalet D. SAM50, a side door to the mitochondria: The

case of cytotoxic proteases. *Pharmacol Res*. 2020;160:105196.

51. Shao B, et al. Ablation of Sam50 is associated with fragmentation and alterations in

metabolism in murine and human myotubes. *BioRxiv Prepr Serv Biol*.

2023;2023.05.20.541602.

52. Shao B, et al. Ablation of Sam50 is associated with fragmentation and alterations in

metabolism in murine and human myotubes. *J Cell Physiol*. 2024;n/a(n/a).

https://doi.org/10.1002/jcp.31293.

53. Tang A, et al. Nonalcoholic Fatty Liver Disease: MR Imaging of Liver Proton Density

Fat Fraction to Assess Hepatic Steatosis. *Radiology*. 2013;267(2):422–431.

54. Ma X, et al. Perspectives on Mitochondria–ER and Mitochondria–Lipid Droplet

Contact in Hepatocytes and Hepatic Lipid Metabolism. *Cells*. 2021;10(9):2273.

55. Simões ICM, et al. Mitochondria in non-alcoholic fatty liver disease. *Int J Biochem Cell

Biol*. 2018;95:93–99.

56. Chiang JYL, Ferrell JM. Bile Acid Metabolism in Liver Pathobiology. *Gene Expr*.

2018;18(2):71–87.

57. TOMIZAWA M, et al. Triglyceride is strongly associated with nonalcoholic fatty liver

disease among markers of hyperlipidemia and diabetes. *Biomed Rep*. 2014;2(5):633–636.

58. Semova I, Biddinger SB. TRIGLYCERIDES IN NAFLD: GUILTY UNTIL PROVEN

INNOCENT. *Trends Pharmacol Sci*. 2021;42(3):183–190.

59. Yanai S, Endo S. Functional Aging in Male C57BL/6J Mice Across the Life-Span: A

Systematic Behavioral Analysis of Motor, Emotional, and Memory Function to Define an

Aging Phenotype. *Front Aging Neurosci*. 2021;13.

https://www.frontiersin.org/articles/10.3389/fnagi.2021.697621. Accessed February 8, 2023.

60. Robertson GL, et al. DRP1-mediated mitochondrial fission is essential to maintain

cristae morphology and bioenergetics [preprint]. 2022;2021.12.31.474637.

61. Giacomello M, Pellegrini L. The coming of age of the mitochondria–ER contact: a

matter of thickness. *Cell Death Differ*. 2016;23(9):1417–1427.

62. Vincent AE, et al. Quantitative 3D Mapping of the Human Skeletal Muscle

Mitochondrial Network. *Cell Rep*. 2019;26(4):996-1009.e4.

63. Ilacqua N, et al. A three-organelle complex made by wrappER contacts with

peroxisomes and mitochondria responds to liver lipid flux changes. *J Cell Sci*.

2021;135(5):jcs259091.

64. Ilacqua N, Anastasia I, Pellegrini L. Isolation and analysis of fractions enriched in

WrappER-associated mitochondria from mouse liver. *STAR Protoc*. 2021;2(3):100752.

65. Ilacqua N, et al. Expression of Synj2bp in mouse liver regulates the extent of wrappER-

mitochondria contact to maintain hepatic lipid homeostasis. *Biol Direct*. 2022;17(1):37.

1128 66. Surmacki JM, Abramczyk H. Confocal Raman imaging reveals the impact of retinoids

1129 on human breast cancer via monitoring the redox status of cytochrome c. *Sci Rep*.

1130 2023;13(1):15049.

1131 67. McReynolds MR, et al. NAD+ flux is maintained in aged mice despite lower tissue

1132 concentrations. *Cell Syst*. 2021;12(12):1160-1172.e4.

1133 68. Goodman RP, Calvo SE, Mootha VK. Spatiotemporal compartmentalization of hepatic

1134 NADH and NADPH metabolism. *J Biol Chem*. 2018;293(20):7508–7516.

1135 69. Ueda N. Ceramide-Induced Apoptosis in Renal Tubular Cells: A Role of Mitochondria

1136 and Sphingosine-1-Phoshate. *Int J Mol Sci*. 2015;16(3):5076–5124.

1137 70. Fugio LB, Coeli-Lacchini FB, Leopoldino AM. Sphingolipids and Mitochondrial

1138 Dynamic. *Cells*. 2020;9(3):581.

1139 71. Parra V, et al. Changes in mitochondrial dynamics during ceramide-induced

1140 cardiomyocyte early apoptosis. *Cardiovasc Res*. 2008;77(2):387–397.

1141 72. Longo N, Frigeni M, Pasquali M. CARNITINE TRANSPORT AND FATTY ACID

1142 OXIDATION. *Biochim Biophys Acta*. 2016;1863(10):2422–2435.

1143 73. Li H, et al. Mic60/Mitofilin determines MICOS assembly essential for mitochondrial

1144 dynamics and mtDNA nucleoid organization. *Cell Death Differ*. 2016;23(3):380–392.

1145 74. Tezze C, et al. Age-Associated Loss of OPA1 in Muscle Impacts Muscle Mass,

1146 Metabolic Homeostasis, Systemic Inflammation, and Epithelial Senescence. *Cell Metab*.

1147 2017;25(6):1374-1389.e6.

75. Vue Z, et al. Three-Dimensional Mitochondria Reconstructions of Murine Cardiac

Muscle Changes in Size Across Aging. *Am J Physiol-Heart Circ Physiol*. [published online

ahead of print: August 25, 2023]. https://doi.org/10.1152/ajpheart.00202.2023.

76. Vue Z, et al. 3D reconstruction of murine mitochondria reveals changes in structure

during aging linked to the MICOS complex. *Aging Cell*. 2023;22(12):e14009.

77. Körner C, et al. The C-terminal domain of Fcj1 is required for formation of crista

junctions and interacts with the TOB/SAM complex in mitochondria. *Mol Biol Cell*.

2012;23(11):2143–2155.

78. Kozjak V, et al. An Essential Role of Sam50 in the Protein Sorting and Assembly

Machinery of the Mitochondrial Outer Membrane *. *J Biol Chem*. 2003;278(49):48520–

48523.

79. Tang J, et al. Sam50–Mic19–Mic60 axis determines mitochondrial cristae architecture

by mediating mitochondrial outer and inner membrane contact. *Cell Death Differ*.

2020;27(1):146–160.

80. Ding C, et al. Mitofilin and CHCHD6 physically interact with Sam50 to sustain cristae

structure. *Sci Rep*. 2015;5(1):1–11.

81. Chen L, et al. Loss of Sam50 in hepatocytes induces cardiolipin☐dependent

mitochondrial membrane remodeling to trigger mtDNA release and liver injury.

Hepatology. 2022;76(5):1389.

82. Tsuru H, et al. HFD-induced hepatic lipid accumulation and inflammation are

decreased in Factor D deficient mouse. *Sci Rep*. 2020;10:17593.

83. Lian C-Y, et al. High fat diet-triggered non-alcoholic fatty liver disease: A review of

proposed mechanisms. *Chem Biol Interact*. 2020;330:109199.

84. Zheng P, et al. High-fat diet causes mitochondrial damage and downregulation of

mitofusin-2 and optic atrophy-1 in multiple organs. *J Clin Biochem Nutr*. 2023;73(1):61–76.

85. De Nicolo B, et al. Calcium and Reactive Oxygen Species Signaling Interplays in

Cardiac Physiology and Pathologies. *Antioxidants*. 2023;12(2):353.

86. Winslow JL, Hollenberg MJ, Lea PJ. Resolution limit of serial sections for 3D

reconstruction of tubular cristae in rat liver mitochondria. *J Electron Microsc Tech*.

1991;18(3):241–248.

87. Mannella CA. Structure and dynamics of the mitochondrial inner membrane cristae.

Biochim Biophys Acta BBA - Mol Cell Res. 2006;1763(5):542–548.

88. Mannella CA, et al. Electron microscopic tomography of rat-liver mitochondria and

their interactions with the endoplasmic reticulum. *BioFactors*. 1998;8(3–4):225–228.

89. Das S, et al. Mitochondrial morphology and dynamics in hepatocytes from normal and

ethanol-fed rats. *Pflugers Arch*. 2012;464(1):101–109.

90. Shami GJ, et al. Three-dimensional ultrastructure of giant mitochondria in human non-

alcoholic fatty liver disease. *Sci Rep*. 2021;11:3319.

91. Daum B, et al. Age-dependent dissociation of ATP synthase dimers and loss of inner-

membrane cristae in mitochondria. *Proc Natl Acad Sci*. 2013;110(38):15301–15306.

92. Labbe G, Pessayre D, Fromenty B. Drug-induced liver injury through mitochondrial

dysfunction: mechanisms and detection during preclinical safety studies. *Fundam Clin*

Pharmacol. 2008;22(4):335–353.

93. Altamirano J, et al. A histologic scoring system for prognosis of patients with alcoholic

hepatitis. *Gastroenterology*. 2014;146(5):1231-1239.e1–6.

94. Palma E, et al. Dynamin-1-Like Protein Inhibition Drives Megamitochondria

Formation as an Adaptive Response in Alcohol-Induced Hepatotoxicity. *Am J Pathol*.

2019;189(3):580–589.

95. Zhou H, et al. DNA-PKcs promotes alcohol-related liver disease by activating Drp1-

related mitochondrial fission and repressing FUNDC1-required mitophagy. *Signal*

Transduct Target Ther. 2019;4:56.

96. Yamada T, et al. Prevention and regression of megamitochondria and steatosis by

blocking mitochondrial fusion in the liver. *iScience*. 2022;25(4):103996.

97. Tandler B, Hoppel CL. Giant Mitochondria (Megamitochondria). In: Lennarz WJ,

Lane MD, eds. *Encyclopedia of Biological Chemistry (Second Edition)*. Waltham: Academic

Press; 2013:375–376.

98. Palma E, et al. Perturbations in mitochondrial dynamics are closely involved in the

progression of Alcoholic Liver Disease. *Alcohol Clin Exp Res*. [published online ahead of

print: 2020]. https://doi.org/10.1111/acer.14299.

99. Palma E, et al. Perturbation of mitochondrial dynamics in Alcoholic Liver Disease. *J*

World Mitochondria Soc. 2016;2. https://doi.org/10.18143/JWMS_V2I2_1948.

100. Wang D, Wei Y, Pagliassotti MJ. Saturated Fatty Acids Promote Endoplasmic

Reticulum Stress and Liver Injury in Rats with Hepatic Steatosis. *Endocrinology*.

2006;147(2):943–951.

101. Renne MF, Hariri H. Lipid Droplet-Organelle Contact Sites as Hubs for Fatty Acid

Metabolism, Trafficking, and Metabolic Channeling. *Front Cell Dev Biol*. 2021;9.

https://www.frontiersin.org/articles/10.3389/fcell.2021.726261. Accessed February 12, 2023.

102. Kim H, et al. Structural basis for mitoguardin-2 mediated lipid transport at ER-

mitochondrial membrane contact sites. *Nat Commun*. 2022;13(1):3702.

103. Marchi S, Patergnani S, Pinton P. The endoplasmic reticulum–mitochondria

connection: One touch, multiple functions. *Biochim Biophys Acta BBA - Bioenerg*.

2014;1837(4):461–469.

104. Anastasia I, et al. Mitochondria-rough-ER contacts in the liver regulate systemic lipid

homeostasis. *Cell Rep*. 2021;34(11):108873.

105. Wang J, et al. Mutual interaction between endoplasmic reticulum and mitochondria in

nonalcoholic fatty liver disease. *Lipids Health Dis*. 2020;19(1):72.

106. Huynen MA, et al. Evolution and structural organization of the mitochondrial contact

site (MICOS) complex and the mitochondrial intermembrane space bridging (MIB)

complex. *Biochim Biophys Acta BBA-Mol Cell Res*. 2016;1863(1):91–101.

107. Ott C, et al. Sam50 Functions in Mitochondrial Intermembrane Space Bridging and

Biogenesis of Respiratory Complexes. *Mol Cell Biol*. 2012;32(6):1173–1188.

108. Gagliano Taliun SA, et al. Exploring and visualizing large-scale genetic associations by

using PheWeb. *Nat Genet*. 2020;52(6):550–552.

109. Kanai M, et al. Genetic analysis of quantitative traits in the Japanese population links

cell types to complex human diseases. *Nat Genet*. 2018;50(3):390–400.

110. Kurki MI, et al. FinnGen: Unique genetic insights from combining isolated population

and national health register data [preprint]. 2022;2022.03.03.22271360.

111. Duchen MR, Szabadkai G. Roles of mitochondria in human disease. *Essays Biochem*.

2010;47:115–137.

112. Alonso MT, et al. Calcium microdomains in mitochondria and nucleus. *Cell Calcium*.

2006;40(5–6):513–525.

113. Bertero E, Maack C. Calcium Signaling and Reactive Oxygen Species in

Mitochondria. *Circ Res*. 2018;122(10):1460–1478.

114. Hoek JB, et al. Calcium ion-dependent signalling and mitochondrial dysfunction:

mitochondrial calcium uptake during hormonal stimulation in intact liver cells and its

implication for the mitochondrial permeability transition. *Biochim Biophys Acta BBA - Mol*

Basis Dis. 1995;1271(1):93–102.

115. Csordás G, Weaver D, Hajnóczky G. Endoplasmic Reticulum–Mitochondrial

Contactology: Structure and Signaling Functions. *Trends Cell Biol*. 2018;28(7):523–540.

116. Egnatchik RA, et al. ER calcium release promotes mitochondrial dysfunction and

hepatic cell lipotoxicity in response to palmitate overload. *Mol Metab*. 2014;3(5):544–553.

117. Kim YS, Kim SG. Endoplasmic reticulum stress and autophagy dysregulation in

alcoholic and non-alcoholic liver diseases. *Clin Mol Hepatol*. 2020;26(4):715–727.

118. Amaya MJ, Nathanson MH. Calcium Signaling in the Liver. *Compr Physiol*.

2013;3(1):515–539.

119. Modi S, et al. Miro clusters regulate ER-mitochondria contact sites and link cristae

organization to the mitochondrial transport machinery. *Nat Commun*. 2019;10(1):4399.

120. Cichoż-Lach H, Michalak A. Oxidative stress as a crucial factor in liver diseases.

World J Gastroenterol WJG. 2014;20(25):8082–8091.

121. Li P-L, Gulbins E. Bioactive Lipids and Redox Signaling: Molecular Mechanism and

Disease Pathogenesis. *Antioxid Redox Signal*. 2018;28(10):911–915.

122. Barja G. The mitochondrial free radical theory of aging. *Prog Mol Biol Transl Sci*.

2014;127:1–27.

123. Navarro A, Boveris A. Rat brain and liver mitochondria develop oxidative stress and

lose enzymatic activities on aging. *Am J Physiol-Regul Integr Comp Physiol*.

2004;287(5):R1244–R1249.

124. Maeso-Díaz R, et al. Effects of aging on liver microcirculatory function and sinusoidal

phenotype. *Aging Cell*. 2018;17(6):e12829.

125. Nahata M, et al. Decline in Liver Mitochondria Metabolic Function Is Restored by

Hochuekkito Through Sirtuin 1 in Aged Mice With Malnutrition. *Front Physiol*. 2022;13.

https://www.frontiersin.org/articles/10.3389/fphys.2022.848960. Accessed February 12,

2023.

126. Barazzoni R, Short KR, Nair KS. Effects of Aging on Mitochondrial DNA Copy

Number and Cytochromec Oxidase Gene Expression in Rat Skeletal Muscle, Liver, and

Heart *. *J Biol Chem*. 2000;275(5):3343–3347.

127. Serviddio G, et al. Bioenergetics in aging: mitochondrial proton leak in aging rat liver,

kidney and heart. *Redox Rep*. 2007;12(1–2):91–95.

128. Moreira OC, et al. Mitochondrial Function and Mitophagy in the Elderly: Effects of

Exercise. *Oxid Med Cell Longev*. 2017;2017:2012798.

129. Guo Y, et al. Quantitative Proteomic and Functional Analysis of Liver Mitochondria

from High Fat Diet (HFD) Diabetic Mice. *Mol Cell Proteomics MCP*. 2013;12(12):3744–

3758.

130. Graham DS, et al. Ability of high fat diet to induce liver pathology correlates with the

level of linoleic acid and Vitamin E in the diet. *PloS One*. 2023;18(6):e0286726.

131. Kim HJ, et al. A high fat diet fosters elevated bisretinoids. *J Biol Chem*.

2023;299(6):104784.

132. Parlakgül G, et al. Spatial mapping of hepatic ER and mitochondria architecture

reveals zonated remodeling in fasting and obesity. *Nat Commun*. 2024;15(1):3982.

133. Arruda AP, et al. Chronic enrichment of hepatic endoplasmic reticulum-mitochondria

contact leads to mitochondrial dysfunction in obesity. *Nat Med*. 2014;20(12):1427–1435.

134. Softic S, et al. Dietary Sugars Alter Hepatic Fatty Acid Oxidation via Transcriptional

and Post-translational Modifications of Mitochondrial Proteins. *Cell Metab*. 2019;30 4:735–

7534.

135. Lockman KA, et al. Oxidative stress rather than triglyceride accumulation is a

determinant of mitochondrial dysfunction in in vitro models of hepatic cellular steatosis.

Liver Int Off J Int Assoc Study Liver. 2012;32(7):1079–1092.

136. Dong J, et al. Mic19 depletion impairs endoplasmic reticulum-mitochondrial contacts

and mitochondrial lipid metabolism and triggers liver disease. *Nat Commun*.

2024;15(1):168.

137. Paradies G, et al. Oxidative stress, cardiolipin and mitochondrial dysfunction in

nonalcoholic fatty liver disease. *World J Gastroenterol WJG*. 2014;20(39):14205–14218.

138. Liu J, et al. Cardiolipin inhibitor ameliorates the non-alcoholic steatohepatitis through

suppressing NLRP3 inflammasome activation. *Eur Rev Med Pharmacol Sci*.

2019;23(18):8158–8167.

139. Alves-Bezerra M, Cohen DE. Triglyceride metabolism in the liver. *Compr Physiol*.

2017;8(1):1–8.

140. Hinton A, et al. ATF4 Dependent Increase in Mitochondrial-Endoplasmic Reticulum

Tethering Following OPA1 Deletion in Skeletal Muscle. *Journal of Cellular Physiology*.

[published online ahead of print: 2024]. https://doi.org/10.1101/2022.09.12.507669.

141. Stahl EC, et al. Inflammation and Ectopic Fat Deposition in the Aging Murine Liver Is

Influenced by CCR2. *Am J Pathol*. 2020;190(2):372–387.

142. Pereira RO, et al. OPA 1 deficiency promotes secretion of FGF 21 from muscle that

prevents obesity and insulin resistance. *EMBO J*. 2017;36(14):2126–2145.

143. Tam J, et al. Peripheral CB1 cannabinoid receptor blockade improves

cardiometabolic risk in mouse models of obesity. *J Clin Invest*. 2010;120(8):2953–2966.

144. Hinton A, et al. A Comprehensive Approach to Sample Preparation for Electron

Microscopy and the Assessment of Mitochondrial Morphology in Tissue and Cultured

Cells. *Adv Biol*. 2023;e2200202.

145. Neikirk K, et al. Systematic Transmission Electron Microscopy-Based Identification

and 3D Reconstruction of Cellular Degradation Machinery. *Adv Biol*. 2023;7(6):2200221.

146. Lu W, et al. Extraction and Quantitation of Nicotinamide Adenine Dinucleotide Redox

Cofactors. *Antioxid Redox Signal*. 2018;28(3):167–179.

147. Wang L, et al. Peak Annotation and Verification Engine for Untargeted LC–MS

Metabolomics. *Anal Chem*. 2019;91(3):1838–1846.

148. Adusumilli R, Mallick P. Data Conversion with ProteoWizard msConvert. In: Comai

L, Katz JE, Mallick P, eds. *Proteomics: Methods and Protocols*. New York, NY: Springer;

2017:339–368.

149. Boudina S, et al. Mitochondrial energetics in the heart in obesity-related diabetes:

direct evidence for increased uncoupled respiration and activation of uncoupling proteins.

Diabetes. 2007;56(10):2457–2466.

150. Olenych SG, et al. The fluorescent protein color palette. *Curr Protoc Cell Biol*.

2007;Chapter 21:Unit 21.5.

151. Roden DM, et al. Development of a large-scale de-identified DNA biobank to enable

personalized medicine. *Clin Pharmacol Ther*. 2008;84(3):362–369.

152. Danciu I, et al. Secondary use of clinical data: the Vanderbilt approach. *J Biomed

Inform*. 2014;52:28–35.

153. Dennis JK, et al. Clinical laboratory test-wide association scan of polygenic scores

identifies biomarkers of complex disease. *Genome Med*. 2021;13(1):6.

154. GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across

human tissues. *Science*. 2020;369(6509):1318–1330.

155. Gamazon ER, et al. A gene-based association method for mapping traits using

reference transcriptome data. *Nat Genet*. 2015;47(9):1091–1098.

156. Hu Y, et al. A statistical framework for cross-tissue transcriptome-wide association

analysis. *Nat Genet*. 2019;51(3):568–576.

157. Zhou D, et al. A unified framework for joint-tissue transcriptome-wide association and

Mendelian randomization analysis. *Nat Genet*. 2020;52(11):1239–1246.

158. Barbeira AN, et al. Integrating predicted transcriptome from multiple tissues

improves association detection. *PLoS Genet*. 2019;15(1):e1007889.

159. Denny JC, et al. PheWAS: demonstrating the feasibility of a phenome-wide scan to

discover gene-disease associations. *Bioinforma Oxf Engl*. 2010;26(9):1205–1210.

160. Carroll RJ, Bastarache L, Denny JC. R PheWAS: data analysis and plotting tools for

phenome-wide association studies in the R environment. *Bioinforma Oxf Engl*.

2014;30(16):2375–2376.

161. Tomar D, et al. MCUR1 Is a Scaffold Factor for the MCU Complex Function and

Promotes Mitochondrial Bioenergetics. *Cell Rep*. 2016;15(8):1673–1685.

1353

1354

1355

1356 **FIGURE LEGEND:**

1357

1358 **Graphical Abstract:** Liver aging causes metabolic, lipidomic, and mitochondrial structural

1359 alterations, reflecting age-dependent losses in the MICOS complex. Diet-dependent losses of the

1360 SAM complex underlie genetic disease associations and mitochondrial structure.

1361

Figure 1: Comparative analyses of liver lipid content and morphology in young and old

patients. Cross-sectional imaging of in-phase (left) and out-of-phase (right) liver anatomy data

from (**A**) males under 55 years old (aged 14–53 years old; n = 10), (**B**) females under 55 years

old (aged 17–41 years old; n = 10), (**C**) males over 60 years old (aged 60–85 years old; n = 10),

and (**D**) females over 60 years old (aged 64–96 years old; n = 10). (**E**) Computed fat percentage

in males, (**F**) females, and (**G**) combined young and old cohorts. (**H**) Oil Red O staining at 20X

and (**I**) 40X magnification in young and (**J-K**) old samples. (**L**) Quantification of Oil Red O

staining per area in 3-month and 2-year murine liver samples (n=10). (**M**) Quantifications of

liver weight relative to total body weight as a percent (n=10), (**N**) relative reverse transcription-

quantitative polymerase chain reaction mtDNA content (n=8), (**O**) bile acids concentrations

(n=10), (**P**) liver triglycerides concentration (n=8), (**Q**) and serum triglycerides concentration

(n=8). For all panels, error bars indicate SEM, Mann–Whitney tests were used for statistical

analysis, and significance value indicate *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.001, ****P ≤ 0.0001,

and ns indicates non-significant.

1376

1377 **Figure 2: Transmission Electron Microscopy and Serial Block Face-Scanning Electron**

1378 **Microscopy Shows Changes in Mitochondria Murine Liver Across Aging. (A)**

1379 Representative transmission electron micrographs in males and (**B**) females from 3-month and

1380 (**C-D**) 2-year murine liver tissue. (**E**) Mitochondrial quantifications of male mitochondria

1381 number (n=21, 3-months; n=24, 2-years) (**F**) mitochondrial area (n=1309, 3-months; n=333, 2-

1382 years) (**G**) and cristae score (n=555, 3-months; n=555, 2-years)**.** (**H**) Mitochondrial

1383 quantifications of female mitochondria number (n=21, 3-months; n=25, 2-years) (**I**)

1384 mitochondrial area (n=1253, 3-months; n=1018, 2-years) (**J**) and cristae score (n=684, 3-months;

1385 n=684, 2-years)**.** (**K**) Schematic depicting removal of the liver. (**L**) Following embedded fixation,

1386 Serial Block Face-Scanning Electron Microscopy (SBF-SEM) allows for ortho-slice alignment.

1387 (**M**) Manual segmentation of ortho slices was performed to yield (**N**) 3-dimensional (3-D)

1388 reconstructions of individually colored mitochondria. (**O**) Representative ortho slice images from

1389 3-month murine liver tissue and (**P**) 2-year murine liver tissue. (**O'**) Representative ortho slice

1390 images with 3D reconstructions of mitochondria overlaid from 3-month murine liver tissue and

1391 (**P'**) 2-year murine liver tissue. (**O''**) Isolated 3D reconstructions of mitochondria from 3-month

1392 murine liver tissue and (**P''**) 2-year murine liver tissue (**Q**) Mitochondrial quantifications

1393 mitochondrial perimeter, (**R**) area, (**S**) and volume. For SBF-SEM, in total, 3-month samples

1394 both included 750 (n=250, per mouse) for a total of 1500 mitochondria, which were used for

1395 statistical analysis. For all panels, error bars indicate SEM, Mann–Whitney tests were used for

1396 statistical analysis, and significance values indicate **$P \leq 0.01$, ***$P \leq 0.001$, and ns, not

1397 significant.

1398 **Figure 3: Serial Block Face-Scanning Electron Microscopy Shows Changes in**

1399 **Mitochondrial Morphology in Liver Across Aging. (A)** 3D reconstructions of mitochondria

1400 displayed from the transverse viewpoint in 3-month and (**B**) 2-year murine liver tissue. (**A'**)

1401 Representative images of 3D reconstructions of mitochondria displayed from the longitudinal

1402 viewpoint in 3-month and (**B'**) 2-year murine liver tissue. (**C**) Based on these measurements, the

1403 mitochondrial sphericity and (**D**) complexity index was determined. (**E**) Mito-otyping displays

1404 diversity in mitochondrial morphology across each relative volume of mitochondria in 3-month

1405 and (**F**) 2-year murine liver tissue. (**G-H''**) Qualitative identification of wrappER. (**G**)

1406 Orthogonal (ortho) slice from 3-month and (**H**) 2-year samples, with (**G'-H'**) 3D reconstruction

1407 of mitochondria and endoplasmic reticulum overlaid. (**G''-H''**) Isolated 3D reconstruction of

1408 mitochondria, in purple, and endoplasmic reticulum, in blue. For SBF-SEM, in total, 3-month

1409 samples both included 750 (n=250, per mouse) for a total of 1500 mitochondria, which were

1410 used for statistical analysis. For all panels, error bars indicate SEM, Mann–Whitney tests were

1411 used for statistical analysis, and significance value indicate ****$P \leq 0.0001$.

1412

1413

1414 **Figure 4: Global metabolomic and lipidomic profiling revealed metabolic dysregulation**

1415 **and disruptions in lipid classes with age in liver tissues.** (**A**) Metabolomics heatmap showing

1416 the relative abundance of metabolites. (**B-M**) Metabolite pools illustrating the metabolic

1417 pathways that are altered with age in the liver– Vitamin A Metabolism, TCA Cycle, Nucleotide

1418 Metabolism, and NAD+ metabolism. (**N**) Heatmap showing enriched lipid classes based on

1419 comparison between young and old liver tissues. (**O**) Volcano plot labeling significant hits,

1420 which have adjusted p-value <0.05 and fold change (+ or -) greater than 1. For each tissue and

1421 metabolite in the heatmaps, the aged samples were normalized to the median of the young

1422 samples and then log2 transformed. Significantly different lipid classes represented in the figures

1423 are those with adjusted p-values < 0.05 (note: p-values were adjusted to correct for multiple

1424 comparisons using an FDR procedure) and log fold changes greater than 1 or less than −1.

1425 Young, n= 4; aged, n= 4. For all panels, error bars indicate SEM, ** indicates p< 0.01; and *p<

1426 0.05, calculated with Student's t-test.

1427

1428 **Figure 5. Aging Causes MICOS Loss, Diet Changes Affect SAM50 Expression, and SAM50**

1429 **has Genetic Associations with Liver Diseases. (A-D)** Quantitative PCR shows changes in

1430 mRNA transcripts in (**A**) *Opa1*, (**B**) *Mitofilin/Mic60*, (**C**) *Chchd3/Mic19*, and (**D**) *Chchd6/Mic25*

1431 between 3-month and 2-year murine liver samples. (**E**) Western blotting showing SAM50

1432 protein levels, relative to tubulin, in 3-month and 2-year murine liver samples. (**F**) Western

1433 blotting showing MIC60 and SAM50 protein levels, relative to ß-actin, in low-fat diet and high-

1434 fat diet murine liver samples. **(G-G')** Representative transmission electron micrographs from

1435 low-fat diet and (**H-H'**) high-fat diet murine liver tissue. Quantifications of (**I**) mitochondria

1436 number, as normalized to cell area (n=7, low fat diet; n=5, high fat diet), (**J**) individual

1437 mitochondrial area (n=592, low fat diet; n=266, high fat diet), (**K**) the sum of all mitochondrial

1438 area normalized to total cell area (n=7, low fat diet; n=5, high fat diet), (**L**) mitochondrial

1439 circularity index, (n=598, low fat diet; n=271, high fat diet), and (**M**) cristae score (n=425, low

1440 fat diet; n=425, high fat diet). (**N-N'**) Representative confocal fluorescence (using mCherry-

1441 Mito-7) from wildtype and (**O-O'**) *Sam50*-knockout fibroblasts. (**P**) Quantifications of

1442 mitochondrial length. (**Q**) Relative proportion of mitochondria that are either fragmented or

1443 tubular in wildtype and *Sam50* KO fibroblasts, with the white area representing percentage of

1444 fragmentation and colored areas representing percentage of typical tubular. (**R**) Phenome-wide

1445 association study (PheWAS) results for SAMM50 GReX in individuals of European ancestry

(n=70,440). The GReX for SAMM50 was tested for association across 1,704 clinical phenotypes

extracted from the EHR. Association tests were run using logistic regression models, accounting

for genetic ancestry (principle components/PC 1-10), sex, age, median age of medical record,

and genotyping batch. Associations that met the Bonferroni-corrected threshold (red line, p <

2.934272×10^{-5}) are labeled with phenotype name [see SFile 1 for all PheWAS results]. The blue

line represents nominal significance (p = 0.05) For all panels, error bars indicate SEM, Mann–

Whitney tests were used for statistical analysis, and significance value indicate ****$P \leq 0.0001$,

ns indicates non-significant.

1454

Figure 6: Loss of MIC60 and CHCHD6 in HepG2 cells results in reduced mCa2+ uptake

and calcium retention capacity and oxidative stress.

(**A**) Raw traces showing mitochondrial calcium uptake in permeabilized *MIC60* and *CHCHD6*

knockdown HepG2 cells along with scr-siRNA transected controls. (**B**) Percentage change in

mCa2+ uptake rate quantified from raw traces. (**C**) Recordings of mitochondrial calcium

retention capacity in scr-siRNA, *MIC60* siRNA, and *CHCHD6* siRNA HepG2 cells. The circles

indicate the number of calcium boluses taken up by specific cells. (**D**) Percentage change in

mitochondrial calcium retention capacity quantified from recordings of mitochondrial calcium

retention capacity. (**E**) Immunoblot confirming siRNA-mediated knockdown of *CHCHD6* in

HepG2 cells. (**F**) Immunoblot confirming siRNA-mediated knockdown of *MIC60* in HepG2

cells. (**G**) 4′,6-diamidino-2-phenylindole (DAPI) staining, MitoPY1 (5 µM, 45 min at 370 c

magnification of 60x), and merge channels in scramble-siRNA (control), MIC60-siRNA, and

CHCHD6-siRNA transfected permeabilized HepG2 cells. (**H**) 4′,6-diamidino-2-phenylindole

(DAPI) staining, MitoBright Deep Red (10 µM, 30 min at 37^0 c), DCFDA (10 µM, 30 min at 37^0

c, magnification of 60x), and merge channels in scramble-siRNA (control), MIC60-siRNA, and

CHCHD6-siRNA transfected permeabilized HEK293 cells. (**I**) Plate reader-based reactive

oxygen species (ROS) quantification. (**J**) Microscopy-based ROS quantification of MitoPY1

orange, (**K**) MitoSox Deep Red, and (**L**) DCFDA. For all statistical tests, one-way ANOVA

statistical test was performed with Dunnett's multiple comparisons test. N=5-10 for all calcium

experiments, each indicated by dots, as run in triplicates. N=9-13 for all oxidative stress

experiments, each indicated by dots, as run in triplicates. Significance values indicate **P ≤ 0.01

and ****P ≤ 0.0001.

1477

1478 **Tables**:

Genetic Ancestry	European	African
N	70,404	15,175
Sex (%M)	44.10%	38.40%
Current Age	57.04 ± 22.42	46.48 ± 21.39
Median Age of Medical Record	48.43 ± 22.24	38.32 ± 21.41
Visits	61.18 ± 75.14	55.02 ± 80.83

1479

1480 **Table 1: Demographics for BioVU Cohort.** Genetic Ancestry Abbreviations: (EA): European

1481 ancestry, (AA): African ancestry.

1482

1483 **Supplementary Figures:**

1484 **Supplemental Figure 1: Heterogeneity in Mitochondrial Quantification Across Samples (A)**

1485 Distribution of mitochondria for mouse heterogeneity (3 mice, each 250 mitochondria surveyed)

1486 in mitochondrial volume**, (B)** surface area, **(C)** perimeter, **(D)** sphericity, and **(E)** complexity

1487 index in young and old liver tissue.

1488

1489 **Supplemental Figure 2: Global metabolomic and lipidomic profiling revealed metabolic**

1490 **dysregulation and disruptions in lipid classes with age in liver tissues. (A)** Metabolic

1491 pathway analysis and (**B**) PCA plot analysis for metabolomics. **(C)** Lipid class enrichment **(D)**

1492 and lipid chain length enrichment based on comparison between young and old livers.

1493

Supplemental Figure 3: Phenome-wide and clinical lab-wide scan for SAMM50 genetically-

regulated gene expression (GReX) in a medical biobank. Clinical phenotypes and laboratory

values from BioVU participants were extracted from Vanderbilt's de-identified electronic health

record database (n= 85,615, top left panel). Genetically-regulated gene expression for SAMM50

was calculated in BioVU participants using models built from the GTEx version 8 data (top right

panel), which contains genotype data matched to RNA-Seq data from 838 donors across 49

tissues. Imputed gene expression was calculated and tested for association across up to 1,704

phenotypes and 329 clinical lab tests using logistic and linear regression models (bottom panel),

1502 accounting for genetic ancestry (principle components/PC 1-10), sex, age, median age of medical

1503 record and genotyping batch.

1504

1505 **Supplementary Material:**

1506 **Video 1:** 3D structure of 3-month murine liver tissue visualized to show volumetric changes.

1507 **Video 2:** 3D structure of 2-year murine liver tissue visualized to show volumetric changes.

1508 **Video 3:** 3D structure of 3-month murine liver tissue visualized to show complexity changes.

1509 **Video 4:** 3D structure of 2-year murine liver tissue visualized to show complexity changes.

1510 **SFile 1:** Phenome-wide association MultiXCan SAMM50 genetically-regulated gene expression

1511 (GReX) in individuals of European ancestry.

1512 **SFile 2:** Laboratory-wide association MultiXCan SAMM50 genetically-regulated gene

1513 expression (GReX) in individuals of European ancestry.

1514 **SFile 3:** Phenome-wide association MultiXCan SAMM50 genetically-regulated gene expression

1515 (GReX) in individuals of African ancestry.

1516 **SFile 4:** Laboratory-wide association MultiXCan SAMM50 genetically-regulated gene

1517 expression (GReX) in individuals of African ancestry.

1518

1519

1520

1521

1522

Visual Abstract

Mitochondrial Cristae Morphology is Maintained by the MICOS and SAM Complex

Aging Causes Changes in Liver Fat %, Mitochondrial Structure, WrappER, Metabolics, and Lipidomics

Age-Dependent Losses of the MICOS Complex Contribute to Dysregulated Calcium and Oxidative Stress

Diet-Dependent Losses of the SAM Complex Contribute to Dysregulated Structure and Disease Risk

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6