Title: MICOS Complex Loss Governs Age-Associated Murine Mitochondrial Architecture and Metabolism in the Liver, While Sam50 Dictates Diet Changes

Authors: Zer Vue[*1], Alexandria Murphy[2*], Han Le[1*], Kit Neikirk[1], Edgar Garza-Lopez[3], Andrea G. Marshall[1], Margaret Mungai[1], Brenita Jenkins[2], Larry Vang[1], Heather K. Beasley[1], Mariaassumpta Ezedimma[1], Sasha Manus[1], Aaron Whiteside[1], Maria Fernanda Forni[4], Chanel Harris[1,5], Amber Crabtree[1], Claude F. Albritton[5,6], Sydney Jamison[6], Mert Demirci[6], Praveena Prasad[2], Ashton Oliver[1], Ky'Era V. Actkins[7], Jianqiang Shao[8], Elma Zaganjor[1], Estevão Scudese[1], Benjamin Rodriguez[1], Alice Koh[1], Izabella Rabago[1], Johnathan E. Moore[1], Desiree Nguyen[1], Muhammad Aftab[3], Benjamin Kirk[3], Yahang Li[3], Nelson Wandira[6], Taseer Ahmad[6,9], Mohammad Saleem[6] Ashlesha Kadam[10], Prasanna Katti[11,12], Ho-Jin Koh[13], Chantell Evans[14], Young Do Koo[3,15], Eric Wang[16], Quinton Smith[16], Dhanendra Tomar[9], Clintoria R. Williams[17], Mariya T. Sweetwyne[18], Anita M. Quintana[19], Mark A. Phillips[20], David Hubert[20], Annet Kirabo[6,21,22,23], Chandravanu Dash[24], Pooja Jadiya[25], André Kinder[26], Olujimi A. Ajijola[27], Tyne W. Miller-Fleming[7], Melanie R. McReynolds[#2], Antentor Hinton, Jr[#1],

Affiliations:
1. Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, 37232, USA.
2. Department of Biochemistry and Molecular Biology, The Huck Institute of the Life Sciences, Pennsylvania State University, State College, PA 16801
3. Department of Internal Medicine, University of Iowa, Iowa City, IA, 52242, USA.
4. Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT 06520.
5. Department of Biomedical Sciences, School of Graduate Studies, Meharry Medical College, Nashville, TN 37208-3501, USA.
6. Department of Medicine, Vanderbilt University Medical Center, Nashville, TN 37232, USA.
7. Division of Genetic Medicine, Department of Medicine, Vanderbilt University Medical Center, Nashville, Tennessee, United States
8. Central Microscopy Research Facility, University of Iowa, Iowa City, IA, 52242, USA.
9. Department of Pharmacology, College of Pharmacy, University of Sargodha, Sargodha, Punjab,40100, Pakistan
10. Department of Internal Medicine, Section of Cardiovascular Medicine, Wake Forest University School of Medicine, Winston-Salem, NC 27157 USA
11. National Heart, Lung and Blood Institute, National Institutes of Health, 9000 Rockville Pike, Bethesda, MD 20892, USA
12. Department of Biology, Indian Institute of Science Education and Research (IISER) Tirupati, AP, 517619, India
13. Department of Biological Sciences, Tennessee State University, Nashville, TN 37209, USA
14. Department of Cell Biology, Duke University School of Medicine, Durham, NC, 27708, USA
15. Fraternal Order of Eagles Diabetes Research Center, Iowa City, Iowa, USA1
16. Department of Chemical and Biomolecular Engineering, University of California, Irvine, CA, 92697, USA.
17. Department of Neuroscience, Cell Biology and Physiology, Wright State University, Dayton, OH 45435 USA

18. Department of Laboratory Medicine and Pathology, University of Washington, Seattle, WA,
98195, USA
19. Department of Biological Sciences, Border Biomedical Research Center, The University of
Texas at El Paso, El Paso, Texas, USA
20. Department of Integrative Biology, Oregon State University, Corvallis, OR, 97331, USA.
21. Vanderbilt Center for Immunobiology, Nashville, TN, 37232, USA.
22. Vanderbilt Institute for Infection, Immunology and Inflammation, Nashville, TN, 37232,
USA.
23. Vanderbilt Institute for Global Health, Nashville, TN, 37232, USA.
24. Department of Microbiology, Immunology and Physiology, Meharry Medical College,
Nashville, TN, United States
25. Department of Internal Medicine, Section of Gerontology and Geriatric Medicine, Sticht
Center for Healthy Aging and Alzheimer's Prevention, Wake Forest University School of
Medicine, Winston-Salem, NC
26. Artur Sá Earp Neto University Center – UNIFASE-FMP, Petrópolis Medical School, Brazil
27. UCLA Cardiac Arrhythmia Center, University of California, Los Angeles, CA, USA.
63
64   *These authors share co-first authorship.
65   #These authors share co-senior authorship.
66
Corresponding Author:
Antentor Hinton
Department of Molecular Physiology and Biophysics
Vanderbilt   University
antentor.o.hinton.jr@Vanderbilt.Edu 319-383-3095

ABSTRACT:
73

The liver, the largest internal organ and a metabolic hub, undergoes significant declines due to
aging, affecting mitochondrial function and increasing the risk of systemic liver diseases. How
the mitochondrial three-dimensional (3D) structure changes in the liver across aging, and the
biological mechanisms regulating such changes confers remain unclear. In this study, we
employed Serial Block Face-Scanning Electron Microscopy (SBF-SEM) to achieve high-
resolution 3D reconstructions of murine liver mitochondria to observe diverse phenotypes and
structural alterations that occur with age, marked by a reduction in size and complexity. We also
show concomitant metabolomic and lipidomic changes in aged samples. Aged human samples
reflected altered disease risk. To find potential regulators of this change, we examined the
Mitochondrial Contact Site and Cristae Organizing System (MICOS) complex, which plays a
crucial role in maintaining mitochondrial architecture. We observe that the MICOS complex is
lost during aging, but not Sam50. Sam50 is a component of the sorting and assembly machinery
(SAM) complex that acts in tandem with the MICOS complex to modulate cristae morphology.
In murine models subjected to a high-fat diet, there is a marked depletion of the mitochondrial
protein SAM50. This reduction in Sam50 expression may heighten the susceptibility to liver
disease, as our human biobank studies corroborate that Sam50 plays a genetically regulated role
in the predisposition to multiple liver diseases. We further show that changes in mitochondrial
calcium dysregulation and oxidative stress accompany the disruption of the MICOS complex.
Together, we establish that a decrease in mitochondrial complexity and dysregulated metabolism

occur with murine liver aging. While these changes are partially be regulated by age-related loss of the MICOS complex, the confluence of a murine high-fat diet can also cause loss of Sam50, which contributes to liver diseases. In summary, our study reveals potential regulators that affect age-related changes in mitochondrial structure and metabolism, which can be targeted in future therapeutic techniques.

Keywords: Aging, 3D Structure, Mitochondria, Metabolism, MICOS Complex, Liver Disease

4

100   **INTRODUCTION:**
101   Mitochondria are required for various purposes including oxidative phosphorylation,
102   calcium homeostasis, and biochemical pathways (1). Mitochondria structure is crucial to many
103   mitochondrial functions (2, 3). Unlike many other organelles, mitochondria have independent
104   fusion and fission dynamics, which allow them to dynamically respond to their environment and
105   adapt a variety of phenotypes beyond the spherical or tubular shapes that are typically associated
106   with mitochondria (2, 4, 5). It is well understood that across aging, changes in morphology and
107   size can impair mitochondrial functions and structures (6–8). Typically, these declines are
108   measured through transmission electron microscopy (TEM), which offers high x- and y-
109   resolutions (9, 10) providing insights into mitochondrial ultrastructure such as cristae, the folds
110   of the inner mitochondrial membrane (11). However, a nascent research avenue is studying 3D
111   phenotypes and exploring novel mitochondria structures (2, 12), such as donut-shaped
112   mitochondria (13, 14), nanotunnels (15, 16), and megamitochondria (17). These structures
113   require imaging techniques, such as serial block face-scanning electron microscopy (SBF-SEM),
114   which allows 3D reconstruction of mitochondria (18–22). For many organs and experimental
115   models, 3D reconstruction has not yet been performed, thus creating a gap in knowledge of 3D
116   mitochondrial structural changes and molecular regulators that are present across the aging
117   process in various tissues. Previously, we have demonstrated changes in mitochondria in other
118   tissues, such as cardiac and skeletal muscle tissue (23, 24). Here, we aim to use 3D
119   reconstruction to understand mitochondrial remodeling in the aging murine liver—which serves
120   as the metabolic hub for the body.
121   The liver is the largest internal organ in vertebrates, performing essential biological functions
122   such as detoxification, metabolism of drugs and nutrients, storage of minerals, and synthesis of
123   plasma (25). The liver facilitates digestion and metabolism, which has an impact on the
124   gastrointestinal and endocrine systems (26). Additionally, the liver generates carrier proteins
125   crucial for growth and reproduction and participates in the metabolism of sex hormones (27).
126   Therefore, studying liver metabolism will provide insights into many crucial biological processes
127   including lipid and glucose homeostasis and detoxification pathways. One of the principal
128   factors that deteriorate the liver is aging (26). After the age of 45, the percentage of mortality
129   linked to liver disease rises significantly in humans (28). With aging comes a progressive
130   expansion of the liver's neural fat and cholesterol volumes and a rise in blood cholesterol, high-
131   density lipoprotein cholesterol, and neutral fat levels (29). Volume alterations, polyploidy nuclei,
132   the accumulation of dense bodies containing lipofuscin, a reduction in smooth endoplasmic
133   reticulum area, and most importantly, a decrease in the quantity and functionality of
134   mitochondria are all associated with aging in liver cells (29). During aging, lipids are also readily
135   accumulated by the liver, raising the risk of cirrhosis, steatohepatitis, and non-alcoholic fatty
136   liver disease (NAFLD), among other hepatic conditions (30, 31). Lipids are necessary for several
137   essential mitochondrial functions, including metabolism and biogenesis, mitochondrial
138   dynamics, oxidative phosphorylation, architecture, and structure of mitochondrial membranes
139   (32). Additionally, the production of energy, regulation of cell death, and bioenergetics are all
140   significantly impacted by the lipid content of the inner mitochondrial membrane (IMM) (32).
141   Thus, as liver cells undergo mitochondrial dysfunction during aging, mitochondrial alterations
142   may contribute to the age-related risk of systematic diseases (25, 33). Thus, within this study, we
143   combine our murine approach of looking at mitochondrial structure with the examination of
144   gross liver structure, lipogenesis, and genetically regulated gene expression in human cohorts.

Age-related changes in mitochondrial function are characterized by dynamic structural
alterations within the mitochondria. Mitochondria are important for generating ATP through the
electron transport chain in their cristae, folds of the IMM (11). Age-related liver disease may
decrease ATP synthesis (34), resulting decreased energy allocation for essential functions.
Beyond this, mitophagy, mitochondria DNA (mtDNA) mutations, disruptions in calcium
homeostasis, and reactive oxygen species (ROS) have been implicated in many liver pathologies,
such as alcohol-related liver disease, non-alcoholic fatty liver disease, chronic hepatitis B and
hepatocellular carcinoma (35, 36). Through mitochondrial quality control mechanisms, oxidative
stress, mtDNA, mitochondrial structure are highly interdependent (37, 38). Notably, changes in
mitochondrial structure are often driven by changes in mitochondrial dynamic proteins (4, 39),
which coordinate fusion and fission processes, allowing mitochondria to increase volume or
repair mtDNA defects through fusion. In contrast, fission can form new mitochondria and
remove damaged mitochondria (40). The organization of proteins belonging to the dynamin-
related GTPase family and their adaptors controls this mitochondrial remodeling (41). The key
regulators of mitochondrial dynamics are mitofusins 1 and 2 (MFN1 and MFN2) and optic
atrophy 1 (OPA1). MFN 1 and 2 regulate outer mitochondrial membrane (OMM) fusion,
whereas OPA1 regulates IMM fusion (42, 43). In contrast, fission is typically associated with
dynamin-related protein 1 (DRP-1). DRP1 redistribution from the cytosol to the OMM is
necessary for mitochondrial fission (42, 44–46). Dysregulation of these finely tuned dynamics
can support pathophysiology; for example, in liver cancer, upregulation of *Mfn1* and *Opa1* play a
role in cancer tumor cell growth mechanisms (5). Conversely, increased *Drp1* and impaired
Mfn1 expression were hallmarks of chronic alcohol exposure in mice; while other models show
Drp1 downregulation upon chronic alcohol exposure (35). Deletion of mitochondrial fission
factor, a recruitment factor of DRP1, in hepatocytes, has similarly led to altered mitochondrial
morphology concomitant with enhanced susceptibility to NAFLD (47), but conversely, blocking
fission can ameliorate hepatic steatosis and oxidative stress in NAFLD (48). This underscores
that while traditional fusion and fission dynamic proteins can affect age-related risk to liver
diseases, other regulators must be considered.

The OMM and IMM include hardwired translocase complexes with a very intricately
specialized protein import mechanism that drives mitochondrial dynamics. These translocases
operate along five distinct protein import pathways, one of which is through the SAM50 channel
(49, 50). *Sam50* is one of the crucial proteins of the outer membrane of the mitochondria.
SAM50 is a SAM complex component with a β-barrel domain (50). Although the role of *Sam50*
in mitochondria is still unclear, studies showed that Sam50 contributes to the preservation of
cristae's structural integrity. *Sam50* maintains the formation of the cristae via interaction with
mitofilin and CHCHD6, which are parts of the mitochondrial contact site and cristae organizing
system (MICOS) (51). Furthermore, *Sam50* aids in the maintenance of the mitochondrial
network (50), establishing and maintaining mitochondrial integrity and mitochondrial
metabolism, with its loss conferring volumetric losses in mitochondria structure (52). Recently,
Sam50 has been linked to NAFLD pathogenesis (49), but the interconnection with aging liver
mitochondrial structure remains poorly elucidated.

In our study, we use human models to explore the impairments of the liver due to aging,
investigating changes in the structure and physical appearance of the liver in young and old
groups. To establish the tandem roles of MICOS and SAM50, we used two-year-old mice, which
represented an elderly model, and three-month-old mice, which represented a juvenile
phenotype. We then studied mitochondrial morphological changes in both two-dimensional (2D)

and three-dimensional (3D) which showed age-related losses in volume and complexity. We also
find that metabolomic and lipogenic changes in aging may mediate mitochondrial structural
alterations in the aging liver. We further learned how the MICOS complex is dysregulated in the
aging liver. Notably, deletion of the MICOS genes resulted in impairments in mitochondrial
calcium regulation and ROS production in the liver. Due to the changes observed in
mitochondria function and structure, we also investigated the role of diet-dependent *Sam50*
depletion in the liver. Lastly, we modeled genetically-regulated gene expression (GReX) of
SAMM50 in a biobank of 85,615 participants, and found that SAMM50 GReX was significantly
associated with multiple liver diseases.

200
201 **RESULTS:**
202
Aging Causes Alterations in Liver Lipid Accumulation and Altered Disease Markers

Damage to the liver's tissue and alterations in hepatic metabolism might result from
abnormal lipid levels in the liver (31). Thus, we utilized magnetic resonance imaging to
determine how lipid content is regulated during aging in human samples. By enrolling female
and male participants (Figures 1A–D), we created a "young" cohort consisting of individuals
from 18 to 55 years old and an "old" cohort of individuals older than 60 years old. For both
sexes, the liver's lipid percentage increased through aging (Figures 1E–F). This age-related
influx in liver validated the progression towards liver diseases in humans. Specifically, when
combining the male and female cohorts together (Figure 1G), the mean fat percent increased
(from a mean of 1.41% $\pm$ 1.44% SD to 5.93% $\pm$ 6.11% SD). Additionally, the 75% percentile of
fat percentage was higher, at 10.5% in the aged cohort, as compared to 2.10% in the young
cohort. Notably, Grade 1 of fat fraction classification, representing mild hepatic steatosis has a
fat percentile cutoff of 6.5% or higher (53). To further validate these results, we switch to our
mouse model and used Oil Red O staining to visualize lipids in 3-month and 2-year murine liver
tissue (Figures 1H-K). Specifically, we saw an increased amount of Oil Red O amount in the 2-
year cohort signifying an increase in lipid amount across the aging process in the liver, with a
more than 6-fold increase in lipid droplets in 2-year samples (Figure 1L; 12.2 $\pm$ 6.55 SD, 3-
months; 77.5 $\pm$ 20.4 SD, 2-years). An uptick of lipid droplets in hepatocytes is indicative of
NAFLD (54). To further explore markers of potential disease progression and dystrophy outside
of lipid accumulation, we also looked at liver mass, which was decreased in aged samples when
normalized to body weight (Figure 1M; 4.56% $\pm$ 0.329% SD, 3-months; 3.93% + 0.376% SD, 2-
years). Since mtDNA content reduction is a hallmark of NAFLD (55), we also showed an age-
dependent loss in mtDNA content (Figure 1N; mean: 1, 3-months; 0.805 $\pm$ 0.119 SD, 2-years).
Additionally, bile acid, which is metabolized in the liver, has increased concentrations with
abnormal liver function and disease states (56). We also looked at the concentration of bile acid,
but no significant difference was noted (Figure 1O; 0.871 µMol/µg $\pm$ 0.138 µMol/µg SD, 3-
months; 0.89 µMol/µg $\pm$ 0.15 µMol/µg SD, 2-years). Finally, central to the pathology and a key
marker of NAFLD is high triglycerides (57, 58). We observed that aging murine samples had
significantly elevated triglycerides in the liver (Figure 1P; 0.30 mmol/L $\pm$ 0.10 mmol/L SD, 3-
months; 0.748 mmol/L $\pm$ 0.15 mmol/L SD, 2-years), which was more drastic than the general
increase in serum triglycerides (Figure 1Q; 0.211 mmol/L $\pm$ 0.100 mmol/L SD, 3-months; 0.379
mmol/L $\pm$ 0.118 mmol/L SD, 2-years). Thus, while we could not confirm aging human and

235 murine samples had liver disease, these results support the observation of an age-related
236 progression toward liver disease.
237
238 **Ultrastructural Changes in Murine Liver Reveals Aging is Associated with Lower Volume**
239 **and Complexity**
240   Next, we sought to explain whether these age-related changes in liver mass and disease
241 markers are correlative with atypical mitochondria structures in the liver. For these studies, we
242 utilized aged C57BL/6J mice at 2 age points, 3-months and 2-years, which are generally
243 understood to be good representations of "young" and "old" aged points akin to aging displayed
244 in humans (59).
245   To begin with, we used TEM to measure changes in mitochondrial morphology in aged mice
246 (Figures 2A-D). Qualitatively, our findings of lipid accumulation were confirmed, with aging
247 also showing lipid droplets that showed less circularity. TEM analysis showed that mitochondria
248 count increased while the average mitochondrion area decreased with aging (Figures 2E-J). To
249 compare sex-dependent differences, mitochondrial counts, as normalized to µm cell area, in 3-
250 month males and females were similar (0.15 ± 0.02 SD, males; 0.174 ± 0.03 SD, females), there
251 was a slight divergence with aging, although both followed a similar trend (Figures 2E, 2H; 0.56
252 $\pm$ 0.130 0.15 SD, males; 0.738 ± 0.14 SD, females). This sex-difference which is more
253 pronounced with age was similarly seen in average mitochondrion area when comparing 3-
254 month samples ($1.27 \, \mu m^2 \pm 1.34 \, \mu m^2$ SD, males; $1.21 \, \mu m^2 \pm 1.19 \, \mu m^2$ SD, females) with their 2-
255 year counterparts (Figures 2F, 2I; $0.233 \, \mu m^2 \pm 0.442 \, \mu m^2$ SD, males; $0.558 \, \mu m^2 \pm 0.386 \, \mu m^2$
256 SD, females). However, in many ways, TEM is inadequate to quantify mitochondria, which exist
257 in diverse 3D shapes (2). Yet, given the best in-class resolution of TEM (9), it is useful for
258 measuring subcellular structures of cristae. In male and female murine samples, we found a
259 decrease in cristae score both (Figures 2G, 2J), which is a semi-qualitative measure (9), that
260 evaluates the relative amount and architectural integrity of cristae. Neither 3-month (3.72 ± 0.44
261 SD, males; 3.47 ± 0.62 SD, females) nor 2-year (2.29 ± 0.71 SD, males; 1.99 ± 0.87 SD,
262 females) samples showed a significant sex-dependent difference in cristae score. Together, these
263 results highlight that cristae integrity is lower in aged liver tissue samples. Additionally, we
264 performed these studies in both male and female samples, which showed slight sex-dependent
265 differences but were generally minimal with similar trends. Given the time-consuming nature of
266 manual segmentation associated with 3D reconstruction, we proceeded with the study utilizing a
267 male cohort.
268   With the 2D mitochondrial differences observed, we utilized 3D reconstruction of SBF-
269 SEM to further investigate changes in mitochondrial structure. For each age cohort, we analyzed
270 approximately 250 mitochondria from the three mice surveyed (Figure 2K), for a total of around
271 750 mitochondria for each age point. SBF-SEM allowed for 50 µM intervals in the z-axis to be
272 imaged for a total of 300 slices, of which 50 slices were used for analysis. The x- and y-
273 resolution was 10 µM, which allowed for about 50 ortho slices to be reconstructed and quantified
274 for each stack (Figure 2L), which then undergo manual contour segmentation (Figure 2M), for
275 the 3D reconstruction of mitochondria (Figure 2N). Displayed first in both age cohorts is a
276 representative of each orthoslice, which allows for the identification of mitochondria (Figures
277 2O-P). From there, manual segmentation across each ortho slice was performed in the software
278 Amira, for mitochondria to be reconstructed (Figures 2O'-P'). Additionally, for better
279 visualization of each individually colored mitochondria, the ortho slice may be removed (Figures
280 2O''-P''). Once modeled, 3D reconstructions (Videos 1-2) can be quantified. We found that in

comparing the young and aged mice, size significantly decreased in all metrics (Figures 2Q-S).
Specifically, the perimeter, which represents the total of all external distances in the
mitochondria was approximately 20% lower in aged samples (Figure 2Q; 10761 µm $\pm$ 1560 µm
SD, 3-months; 8054 µm $\pm$ 450 µm SD, 2 year). Three-dimensional area, which represents a
metric of OMM area, was decreased by nearly 50% in aged samples (Figure 2R; 7.66 µm^2 $\pm$
0.808 µm^2 SD, 3-months; 4.76 µm^2 $\pm$ 0.511 µm^2 SD, 2-years). Finally, mitochondrial volume,
which represents the total of all internal pixels within the 3D mitochondrial reconstruction, was
approximately 30% lower in aged male murine liver samples (Figure 2S; 1.01 µm^3 $\pm$ 0.182 µm^3
SD, 3-months; 0.696 µm^3 $\pm$ 0.107 µm^3 SD, 2-years). Losses in mitochondrial volume may
indicate decreased internal volume for ATP synthesis (60), but mitochondrial roles extend
beyond their energetics, including through interfacing with endoplasmic reticulum (ER) to
modulate calcium homeostasis (61). Thus, it is equally important to look at their morphology and
capacity to form compact sites.
To aid in visualizing mitochondria, we present mitochondria 3D reconstructions from
transverse (Figures 3A-B) and longitudinal (Figures 3A'-B') viewpoints (Videos 3-4). To further
verify this change across aging we looked at sphericity and mitochondrial complexity index
(MCI) (62), which are analogous measures of mitochondrial morphology. Sphericity (calculated
as $\frac{\pi^{\frac{1}{3}}(6*Volume)^{\frac{2}{3}}}{Surface\ Area}$), as exhibited an approximate 20% higher mean value (Figure 3C; 0.638 $\pm$
0.0211 SD, 3-months; 0.785 $\pm$ 0.0257 SD, 2-years). MCI (calculated as Surface Area^3 /16 π
^{2}volume^2) expectedly showed an inverse decrease in the aged cohort (Figure 3D; 3.22 $\pm$ 0.240
SD, 3-months; 1.64 $\pm$ 0.236 SD, 2-years). Together, these validate that complexity is decreased
in aged murine liver samples. To better visualize these changes, we organized mitochondria by
their volume, in a method known as mito-otyping, to compare morphology across size (Figure
3E). In the three-month age cohort (Figure 3E), we observed an expected phenotype of normally
sized mitochondria. However, we also observed some diversity in mitochondrial structure, as we
noted elongation, branching, and other structures that prioritize surface area over volume. In
contrast, the 2-year sample showed much less heterogeneity and mitochondrial structures mostly
presented as tubular or compact (Figure 3F). For these metrics, while intra-sample heterogeneity
is plentiful, there is minimal inter-sample heterogeneity or outliers (sFigures 1A-E).
These changes in sphericity and morphology may impact the ability of mitochondria to
form contact sites. Particularly, liver mitochondria are known to exhibit phenotypes in which ER
wraps around them, known as wrappER, to modulate lipid flux (63–65). To investigate this
paradigm, we qualitatively identified wrappER within liver tissue (Figures 3G-H). Qualitatively,
we found that wrappER exists in both 3-month and 2-year murine liver tissue (Figures 3G'-H'),
but in 3-month tissue wrappER contact is much more extensive (Figures 3G"-H"). This suggests
that liver lipid homeostasis may be negatively impacted (63–65). However, to confirm these
changes and further explore the implications of these mitochondrial morphological alterations,
we analyzed metabolism and performed lipidomics in mice.
319
Global metabolic and lipidomic profiling reveals altered energy metabolism and storage in
the aged liver.
Due to observed changes in mitochondria in the aged liver, we proceeded to analyze
alterations in small biomolecules regulating energy metabolism, cellular homeostasis, and
storage. Our investigation involved global metabolic and lipidomic profiling of young and aged
liver tissues, revealing dynamic features and significant changes in aged mouse livers (Figures

4A-O; sFigures 2A-D). We noted an accumulation of metabolites related to Vitamin A
metabolism, specifically retinoic acid and retinal (Figures 4B-C). While Vitamin A metabolism
must be finely balanced, within breast cancers retinoid changes can be associated with
controlling the redox status of cytochrome c (66). This finding supports the presence of
metabolic shifts, altered cellular signaling, and impaired clearance mechanisms in the aging
liver, all attributed to changes in mitochondrial function. Our metabolomics analysis also
uncovered disruptions in tricarboxylic acid (TCA) cycle intermediates—succinate and malate,
indicating mitochondrial dysfunction and impaired activity (Figures 4D-E). These changes can
impact the production and regulation of these intermediates. Furthermore, we observed decreases
in nucleotide monophosphates involved in purine and pyrimidine metabolism—GMP, CMP,
UMP, AMP (Figures 4F-I). The synthesis of these nucleotides, which involves mitochondrial
processes, is influenced by age-related mitochondrial dysfunction, affecting the overall
nucleotide biosynthesis pathway supporting our observations. Our results additionally confirmed
existing literature on dysregulated nicotinamide adenine dinucleotide (NAD+) metabolism in the
aging liver (67). Significant depletions in tissue NAD+, NADP, NMN pools, and tissue ADP
were detected (Figure 8J-M). Historical liver studies suggest that mitochondrial NAD(H) pools
tend to be oxidized, playing a crucial role in energy homeostasis. Conversely, cytosolic
NADP(H) pools tend to be highly reduced for reductive biosynthesis (68). Our data reveals
disruptions to mitochondrial activity reverses these classical mechanisms. In summary, these
findings collectively support the presence of altered energy metabolism and cellular homeostasis
in the metabolically active liver due to mitochondrial dysfunction and impairment.
Lipidomic profiling of both young and aged liver tissues unveiled age-related changes in
lipid classes and chain lengths (Figure 9N-O; sFigures 2C-D). In the aging liver, significant
alterations were observed in the triglycerides oligomers (TGO), triglycerides (TG), ceramide
(Cer), and acylcarnitine (CAR) lipid classes compared to other lipid groups (Figures 4N-O,
sFigure 2C). These lipid classes, each with distinct roles in liver mitochondria, contribute to
various aspects of cellular metabolism and homeostasis. Disruptions in TGOs and TGs suggest
imbalances in energy storage and release in the aging liver, possibly due to mitochondrial
dysfunction. TGs, being the primary form of stored energy, can participate in mitochondrial beta-
oxidation, providing acetyl-CoA for energy production. Conversely, Cer regulates apoptosis and
modulates mitochondrial membrane permeability (69). Changes in Cer levels can influence
mitochondrial integrity and function (70, 71). CARs play a crucial role in transporting fatty acids
into the mitochondria for beta-oxidation, thus maintaining mitochondrial function (72).
Additionally, significant differences in lipid chain lengths were noted with age in the liver,
impacting membrane integrity, fluidity, and functionality (Figure 4D). These findings align with
altered metabolomic and lipogenic changes attributed to the modified shape and function of
mitochondria with age.
363
364
Murine Aging and Diet Exhibit Distinct Changes in the MICOS and SAM Complex
It is well established that the MICOS complex is critical for mitochondrial dynamics (73),
and our group has previously investigated the effect of aging on the MICOS complex in kidney
tubular cells (24), however, it is still unclear how aging affects the MICOS in aging liver. Studies
have shown that *Opa1*, which is epistatic to the MICOS complex and physically interacts with
components of the MICOS complex (43), decreases with age (74). With *Opa1* as a positive
control, we sought to determine if the MICOS complex mRNA expression is also decreased in

murine liver with age. Consistent with previous studies, *Opa1* mRNA decreased by over 50%
between 3 months and 2 years (Figure 5A). Mitofilin also decreased significantly (Figure 5B).
Similarly, *Chchd3* and *Chchd6* also progressively decreased with age but not as much as the
decline of other transcripts (Figures 5C-D). While *Opa1* interacts with the MICOS complex, it is
not required for the formation of cristae junctions at which the MICOS complex forms, nor does
Opa1 loss negatively affect MICOS components (54). This suggests that the loss of the MICOS
complex across aging occurs in an *Opa1*-independent manner. This parallels our previous
findings in skeletal muscle, cardiac, and kidney (24, 75, 76).
The SAM complex, particularly its β-barrel channel *Sam50*, interacts with the MICOS
complex to regulate cristae morphology (77–80). Particularly, overexpression of SAMM50
enhances fatty acid oxidation and reduces intracellular lipid accumulation, while polymorphisms
in SAMM50 are associated with NAFLD (49), and loss of Sam50 in hepatocytes leads to
cardiolipin-dependent mitochondrial membrane remodeling, mtDNA release, and liver injury
(81). So, we also looked at SAM50 protein levels across aging. Interestingly, unlike the MICOS
complex and *Opa1*, we saw no significant changes in SAM50 levels (Figure 5E). Notably,
beyond aging, a key risk factor for NAFLD is a high-fat diet (HFD) which can cause lipid
accumulation and dyslipidemia (82). Since past studies have shown that overexpression of
SAMM50 can mitigate lipid accumulation (49), we focused on how an HFD also affects these
regulators. To do this, we subjected male mice to an HFD, with fat constituting 60% of their diet,
and a littermate cohort to a low-fat diet (LFD), with fat constituting only 5% of their calories, for
a total of 12 weeks, with mice sampled at 20 weeks. Contrary to our age-related findings, we
showed that while MIC60, a key component of the MICOS complex, remains unchanged,
SAM50 shows a decrease with an HFD. Given the role of SAMM50 in mitigating lipid
accumulation (49), this loss may increase susceptibility to lipid accumulation and dyslipidemia
that is a hallmark of HFD (82).
To further investigate the impact of an HFD on mitochondrial ultrastructure, we
performed TEM analysis in a 20-week-old murine cohort subjected to an LFD and HFD (Figures
5G-H''). Qualitatively, we observed more lipid droplets with less circularity in HFD samples.
Looking at mitochondria, although mitochondrial count, when normalized to the µm cell area,
was fewer in HFD samples (Figure 5I; 0.210 ± 0.114 SD, LFD; 0.128 ± 0.0972 SD, HFD), this
change was not significant. Similar to aging, HFD samples exhibit a significantly higher average
mitochondrion area (Figure 5J; $0.570 \ \mu m^2 \pm 0.401 \ \mu m^2$ SD, LFD; $0.923 \ \mu m^2 \pm 1.12 \ \mu m^2$ SD,
HFD). When considering these two quantifications in tandem, we also calculated the percentage
of mitochondrial area related to the total cell area, which showed no significant difference
(Figure 5K; $11.8 \ \mu m^2 \pm 5.59 \ \mu m^2$ SD, LFD; $11.7 \ \mu m^2 \pm 5.10 \ \mu m^2$ SD, HFD). When looking at
how mitochondrial shape changes with a HFD, we observed that mitochondrial had a higher
circularity, indicative of potentially less complexity, with a HFD (Figure 5L; 0.781 ± 0.149 SD,
LFD; 0.897 ± 0.0748 SD, HFD). Finally, looking at cristae score, similar to aging samples, HFD
cohorts had a significantly lower cristae score (Figure 5M; 3.74 ± 0.441 SD, LFD; 2.04 ± 0.770
SD, HFD). Together, these findings show that a HFD can parallel age-related changes, and that
HFD exhibit smaller mitochondria with aberrant cristae due, in part, to a loss of SAM50.
Previously, we have performed a rigorous 3D analysis of mitochondrial structure and
networking in *Sam50*-deficient human and murine myotubes, showing that the deletion of *Sam50*
results in smaller and less complex mitochondria (52). To further investigate SAM50's role in
mitochondrial structure, we knocked out *Sam50* in fibroblasts and used mito-mCherry confocal
to investigate structure (Figures 5N-O''). Our results further validate our previous findings,

showing that Sam50 deletion causes impairments in mitochondrial length (Figure 5P; 4.14 µm $\pm$
1.57 µm SD, WT; 2.59 µm + 0.539 µm SD, *Sam50* KO). Additionally, when comparing the
percentage of fragmented mitochondria to tubular mitochondria, there was a significantly higher
rate of fragmented mitochondria in *Sam50* KO than in WT. (Figure 5Q; 3.33% $\pm$ 2.08% SD, WT;
57.3% + 18.0% SD, *Sam50* KO). Notably, these findings parallel our findings in HFD of reduced
mitochondrial size, indicating that loss of *Sam50* may dictate the negative bioenergetics caused
by an HFD (83, 84).

Clinical relevance of Sam50 in liver pathologies
Previously, we found that *Sam50* is responsible for establishing and maintaining
mitochondrial integrity, cristae structure, and mitochondrial metabolism (52). To identify the
clinical implications of dysregulated *SAMM50* gene expression in humans, we modeled
genetically regulated gene expression (GReX) of *SAMM50* across 85,615 individuals in the
Vanderbilt biobank, BioVU (Table 1). Cross-tissue *SAMM50* GReX was calculated using the
genetic information for each individual, combined with *SAMM50* gene expression models
developed from the GTEX dataset (see Methods, sFigure3). The relationship between *SAMM50*
GReX and clinical phenotypes (n=1,704) and laboratory values (n=326) were then evaluated by
logistic and linear regression models, respectively. In 70,440 BioVU individuals of European
ancestry, we found that *SAMM50* GReX is significantly associated with multiple liver disease
diagnoses, including: chronic liver disease and cirrhosis, alcoholic liver damage, liver transplant,
acute gastritis, and esophageal bleeding (p < 2.93427E-05, Figure 5R, SFile1). Within the 15,175
BioVU individuals of African ancestry, we did not find any significant associations between
SAMM50 GReX and the clinical phenotypes tested; however, we did see a nominally significant
association with acute gastritis (p=0.00142, SFile3). We did not find any significant associations
between SAMM50 GReX and clinical lab values within the BioVU individuals of European or
African ancestry (SFile2, SFile4).

Knockdown of MIC60 and CHCHD6 reduces Ca^{2+} uptake and retention capacity in HepG2
cells
While this underscores how diet changes may contribute to liver diseases, we also wanted
to establish how the MICOS complex contributes to age-related changes. To unravel the
involvement of *MIC60* and *CHCHD6* in mitochondrial calcium ($_mCa^{2+}$) homeostasis, we
monitored $_mCa^{2+}$ uptake and $_mCa^{2+}$ retention capacity in *MIC60* and *CHCHD6* knockdown
human-derived HepG2 cells. Both *MIC60* and *CHCHD6* knockdown cells show reduced $_mCa^{2+}$
uptake (Figure 6A). Furthermore, to determine if the altered MICOS and cristae structure
contributes to impaired $_mCa^{2+}$ retention capacity and involved in mitochondrial permeability
transition pore opening, we measured the $_mCa^{2+}$ retention capacity in *MIC60* and *CHCHD6*
knockdown cells. *MIC60* as well as *CHCHD6* knockdown cells underwent rapid permeability
transition opening as compared to control cells (Figure 6B). We also found that $_mCa^{2+}$ retention
capacity is significantly reduced in both *MIC60* and *CHCHD6* knockdown HepG2 cells (Figure
6C). These data suggest that altered MICOS complex and cristae disorganization increase the
susceptibility of HepG2 cells to $_mCa^{2+}$ dysregulation and Ca^{2+}-induced cell death.

Knockdown of MIC60 and CHCHD6 induces oxidative stress
Ca^{2+} directly affects oxidative stress signaling and reactive oxygen species (ROS) generation
(85). To determine if MICOS affects ROS production, we evaluated total ROS, mitochondrial

superoxide, and H_2O_2 production in *MIC60* and *CHCHD6* knockdown cells (Figures 6G-H).
Mitochondrial H_2O_2 content, as measured by Mitochondria peroxy yellow 1, increased following
MIC60 and *CHCHD6* knockdown when quantified by both plate-reader based (Figure 6I) and
microscopy-based (Figure 6J) ROS quantification. Silencing *MIC60* and *CHCHD6* in HepG2
cells significantly increased mitochondrial superoxide production and more general intracellular
ROS, detected by MitoBright Deep Red (Figure 6K) and DCFDA (Figure 6L), respectively.
These findings indicate that suppression of *MIC60* and *CHCHD6* disrupts mitochondrial ROS
homeostasis, demonstrating that the MICOS complex is associated with oxidative stress.
472
473 **Discussion:**
474
Age-Related Changes in Liver Mitochondria Structure
Liver mitochondria have been imaged using 3D in rodent models since the early 1990s (86).
Previously, in rat liver, 3D reconstruction revealed that cristae showed high variability in
structure (87). 3D reconstruction also showed that there are extensive mitochondria endoplasmic
reticulum contact sites (MERCs) in liver mitochondria [35]. Additionally, MERCs are sites for
various purposes, such as calcium homeostasis and lipid homeostasis (88). In mice with high
alcohol consumption, alcohol increased mitochondrial volume while conversely reducing the
networking of mitochondria (89). Notably, the formation of mega mitochondria, a novel
mitochondria 3D structure, arose in human non-alcoholic fatty liver disease (90). Beyond these
3D studies, previous studies using cryo-electron tomography (cryo-ET) have found that
mitochondria fragmented and lamellar cristae lost integrity across the aging process (91). While
these studies have aided in revealing previously unknown specific mitochondrial structures in the
liver, the same has not yet been done for different mitochondrial structures that may arise across
aging.

Here, we use SBF-SEM to attempt to answer how mitochondria change in an aged liver and
the functional impact of such changes. We found that in aged states, the size and morphology of
liver mitochondria are altered in both 2D and 3D. Using SBF-SEM and Amira for 3D
morphology of mitochondria and comparing 2-year against 3-month, we observed fragmentation
of mitochondria and decreased complexity. Importantly, we found that an aged murine liver has
a decrease in area, perimeter, volume, and complexity in mitochondria. Mitochondria structure
remains relevant in aging as structural decline may decrease mitochondrial function, thus
reducing certain liver roles. For example, mitochondria dysfunction has been linked to an
increased risk of liver damage upon drug treatment (92). Beyond this, in humans and mice with
alcoholic liver disease, it has been suggested that megamitochondria form due to the lack of
Drp1 and serve a protective role (93, 94). This may be due to the downregulation of the
NR4A1/DNA-PKcs/p53 pathway which has been shown to activate Drp1 in other liver diseases
(95). Aging and disease states affect each other; therefore, future experiments must look at the
mitochondria 3D reconstruction in different disease states across aging to see if the confluence of
these factors leads to the unique mitochondria 3D structures. Interestingly, while
megamitochondria are a hallmark of hepatotoxicity and steatosis (94, 96, 97), and aging
exhibited numerous hallmarks of progression towards liver diseases (Figure 1), we did not
observe these megamitochondria in our sample, rather showing mostly smaller mitochondria.
Since the formation of megamitochondria in Alcoholic Liver Disease follows a significant
increase in Drp1 expression and mitochondrial fragmentation (98), it may be that our aged

sample represents a pre-diseased state that can subsequently shift to megamitochondria as a
compensatory response to reduce alcohol-induced toxicity (94, 98, 99).

Beyond mitochondrial 3D structure, past studies have shown that fatty acids promote
endoplasmic reticulum stress and liver injury in rodent models (100). Due to the increase of lipid
droplets in both aging and HFD in TEM images, there may be an increase in mitochondria-lipid
droplet contact sites (MLDCs) (61, 101). Beyond this, lipid droplet-endoplasmic reticulum
contacts are also understood to help form lipid droplets and perform metabolism (54). Especially
relevant is that MLDCs serve as a place for fatty acid homeostasis (101). Additionally, MERCs
importantly play a role in lipid homeostasis and synthesis (102). MERCs are understood to
regulate ER stress (103), so increases in fatty acids, triggered by lipid droplet formation, can
increase MERC formation in the aged model. Conversely, we qualitatively showed a decrease in
wrappER, which is known to maintain lipid flux through a 3-organelle contact site that involves
peroxisomes (63). These wrappER sites, which contain sites of adhesion, regulate very-low-
density lipoproteins (104). The age-related loss of mitochondrial complexity may impair the
ability and relative surface area mitochondria in murine liver samples must form contact sites,
including MERCs, thus interfering with functions including lipid homeostasis. Indeed, past
reviews have suggested targeting MERCs in NAFLD due to the role of contact sites in glucose
and lipid metabolism (105). However, in the future, a more rigorous analysis of age-related
changes in contact sites within liver tissue is necessary.

528
Sam50 as a Regulator of Liver Disease

Due to the observed dysfunctions in mitochondrial function and structure, we
investigated the role of *SAM50*, part of the SAM required for β-barrel protein assembly, in the
liver (80). There is evidence that the IMM proteins MIC60 (Mitofilin) and CHCHD3 were part
of a protein complex that includes the SAM components, known as the mitochondrial
intermembrane space bridging (MIB) complex (80, 106). The preservation of cristae depends on
interactions between the MIB complex's OMM and IMM components. The destabilization of the
MIB complex led to a decrease in the respiratory chain complex assembly. SAM50 depletion
over time affects the protein contents of all large respiratory complexes containing subunits
encoded by the mitochondria, suggesting a link between SAM50 and cristae structural integrity,
respiratory complex assembly, and mitochondrial DNA (mtDNA) maintenance (107). Our
previous study have shown that *Sam50* plays a significant role in the maintenance of both the
mitochondrial network and the structure of cristae and MICOS, especially in the context of liver
injury (52). Mice with *Sam50* deletion or liver-specific deletion produced by acetaminophen, an
efficient antipyretic and analgesic, experience mtDNA release, which activates the cGAS-STING
pathway and causes liver inflammation (81). Here, using phenome-wide and clinical lab-wide
scans for *SAMM50* GReX in a medical biobank, we saw a significant association between
SAMM50 and multiple liver diseases including nonalcoholic or alcoholic liver disease and
cirrhosis. This suggested that the expression of *SAMM50* as a transcription factor may directly or
indirectly contribute to mitochondrial defects of the liver.
To corroborate the clinical consequences of *SAMM50* GReX in BioVU participants, we
queried publicly available databases including: PheWeb (108), Biobank Japan PheWeb (109),
and the FinnGen web browser (110). PheWeb reports genome-wide association results from the
UK Biobank population (n > 400,000). The top phenotype associations for the *SAMM50* locus
within PheWeb included: Chronic liver disease and cirrhosis (p=2.8 X 10^{-24}, cases=2,954), Other

chronic nonalcoholic liver disease (p=1.2e-18, cases =1,687), Alcoholic liver damage (p=1.4 X
10^{-11}, cases=836), Injury to other and unspecified nerves (p=2.2 X 10^{-10}, cases=269), Portal
hypertension (p=1.7 X 10^{-9}, cases=527), and Liver abscess and sequelae of chronic liver disease
(p=4.1 X 10^{-9}, cases=974). BioBank Japan PheWeb similarly reports genome-wide association
results from the BioBank Japan Project (n > 260,000). The top phenotype associations for the
SAMM50 locus in BioBank Japan PheWeb included Cirrhosis (p=2.8 X 10^{-14}, cases=2,551), as
well as Aspartate transaminase (p=3.3e-87, cases=150,068), Alanine aminotransferase (p=3.0 X
10^{-71}, cases=150,545), and platelet count (p=6.5 X 10^{-38}, cases=148,623). The FinnGen web
browser reports genome-wide association studies from the FinnGen data (n=342,499). Top
results from FinnGen included several phenotypes related to liver diseases including cirrhosis
(p=9.6 X 10^{-54}, cases=3,548), diseases of liver (p=4.9e-49, cases=9,548), and nonalcoholic fatty
liver disease (p=5.7 X 10^{-40}, cases=1,908). This underscores the therapeutic potential of targeting
SAMM50.

The MICOS Complex as a Regulator of Aging:
In this study, we demonstrated the disruptions in many aspects of mitochondria structure
in both 2D and 3D analysis of mitochondria in aging liver. We also showed MICOS impairment
in the aging liver. Therefore, we investigated whether MICOS regulates mitochondrial calcium
influx and oxidative stress. Calcium influx in mitochondria can reflect cell viability (111, 112).
Mitochondria mediation of calcium can impact many cell processes, such as apoptosis, signaling,
and ATP production (113). In some cases, elevated mitochondria calcium uptake can occur
antecedent to mitochondrial swelling, which results in a pathway often leading to apoptosis
(114). However, there was a decline in calcium uptake as *CHCHD6* and *MIC60* were disrupted
(Figure 6), suggestive of impaired cellular calcium signaling, which has previously been linked
with ER stress and dysregulation of MERCs (113, 115). This provides a plausible disease link as
ER calcium release promotes mitochondrial dysfunction, inducing oxidative stress and
hepatotoxicity (116), with broader implications in NAFLD (117, 118). Since Miro clusters both
interact with the MICOS complex, as well as regulate MERCs (119), Miro represents a potential
future mechanistic avenue through which MICOS complex contributes to MERC tethering.
Mitochondria redox state has been proposed as a principal moderator of mitochondrial
function in liver disease (120). Recent studies have continued to highlight a link between ROS
and mitochondrial dynamics (38). Lipids can aid in stimulating the production of ROS (121).
Additionally, many theories regarding aging have been formulated around the basis that
mitochondria lose function due to ROS byproducts which accumulate across aging (122). There
remains controversy about whether ROS are generated in the liver, while some studies have
found they arise during aging in the liver (123), other studies conversely found that in the aging
human liver, no superoxides are produced (124). Future studies should explore how ROS
contributes to the observed 3D mitochondrial phenotypes, as oxidative stress may be a regulator
of the phenotypes we observed. It is noteworthy that the loss of CHCHD6 and MIC60 leads to
oxidative stress (Figure 6). This suggests that during aging, the liver undergo a vicious cycle
wherein abnormal mitochondrial structures generate more harmful byproducts. These
byproducts, in turn, worsen mitochondrial structural dysfunction, contributing to age-related
oxidative stress.
597
The Interaction Effect of the SAM Complex and MICOS Complex in Aging and Diet:

In aged murine liver, decreases in the mtDNA numbers, urea cycle, mitochondrial
function, locomotor activity, loss of enzymatic activity, and anti-oxidative stress are well
established (123, 125, 126). Mitochondria also show diminished membrane potential across
aging in murine liver (127) and declined mitophagy responses (128). Within our study, we
looked at *Sam50* in the context of aging. Although we did not observe a change in the protein
level of SAM50 in the aging liver, there were defects in SAM50, but not MIC60, protein
production during the HFD. Notably, our study contrasts previous quantitative proteomic
analyses of liver mitochondria from HFD Diabetic mice, which found that Mitofilin, Sam50, and
Chchd3 are all upregulated (129). While more research is warranted to resolve this controversy,
our findings suggest that *Sam50* may not regulate mitochondrial machinery during aging
directly, but future studies performing an RNA sequencing scan of aged liver tissue, compared to
young tissue, will provide some candidates that interact with Sam50.
We know that HFDs are linked with many liver complications. For example, oxidative
stress, inflammation, and lipogenesis are some of the factors that might be linked to HFD and
exacerbate the onset of non-alcoholic fatty liver disease (83). Additionally, HFD is associated
with metabolic dysregulation of the liver (130). In this study, we found an increase in retinoic
acid and retinal in the aged liver (Figure 4). Notably, there is evidence linking retinol-binding
proteins and HFD. The enhanced supply of vitamin A and high-fat consumption in diet-induced
obese mice is linked to the production of bisretinoid (131). Interestingly, obesity within
hepatocytes has also been shown to increase MAMs (i.e., MERC-isolated biochemical fractions),
which confers mitochondrial dysfunction (132, 133). High fat within a diet may further have an
additive effect with aging to exacerbate age-related hepatic pathologies. For example, an HFD
can alter mitochondrial structure, leading to decreased fatty acid oxidation (134). Within age-
related hepatic steatosis, impaired fatty acid oxidation, and other factors that HFD can contribute
to such as insulin resistance, are key risk factors (30). Notably, *SAMM50* deficiency causes lipid
accumulation and impaired fatty acid oxidation (49). Thus, HFDs can play a role in metabolism
defects in the liver tissue, although this must be further investigated in the future.
The changes in SAM50 during HFD may be due to several factors. Our lipidomic studies
confirm suggested dysregulations of lipids in the aged liver (Figure 4). Our observation of
SAM50 decreased in HFD contrasts *in vitro* models of hepatic cellular steatosis, revealing that
mitochondrial dysfunction is determined by oxidative stress instead of lipid buildup (135). This
finding is novel since *Sam50*-deletion or overexpression, and the broader SAM complex, during
HFD in the liver, have not been studied. However, past studies of HFD offer some insight. HFD
can lead to pathological alterations and damage to the ultrastructure of the mitochondria and
downregulation of MERCs regulators, MFN2 and OPA1 (84), which parallels the TEM
ultrastructural changes we observed. Additionally, *Mic19/Chchd3* was shown to be necessary in
the liver (136). *Sam50* is closely related to *Mic19*. *Sam50* mediates mitochondrial outer and inner
membrane interactions via the Sam50-Mic19-Mic60 axis (79). In the liver, *Mic19/Chchd3*
deletion reduces ER-mitochondrial contacts, disrupts mitochondrial lipid metabolism,
disorganizes mitochondrial cristae, and causes unfolded protein stress response in mouse
hepatocytes, leading to impairments of liver mitochondrial fatty acid β-oxidation and lipid
metabolism (136). This may explain our results of declined *Sam50* synthesis during HFD, but it
remains unclear why *Mic60* is not affected. Alternatively, linking MICOS and SAM complex
changes, in hepatocytes both SAM50 and MIC60 cooperate to bind to cardiolipin, a phospholipid
exclusively located in mitochondria (81). ROS also targets cardiolipin, which is associated with
mitochondrial dysfunction in NAFLD (137). Cardiolipin inhibition has mitigated non-alcoholic

steatohepatitis through NLRP3 inflammasome activation, suggesting a therapeutic role of
cardiolipin (138). *Sam50* depletion leads to cardiolipin externalization and subsequent liver
inflammation (81). Interestingly, paralleling our findings of increased lipogenesis with aging
(Figure 1), our lipidomic analysis showed altered lipogenesis across multiple lipid classes, which
has implications extending beyond cardiolipin, since the liver is a principal site for lipid
homeostasis (139). Thus, while aging exhibits a protective mechanism to maintain *Sam50*, the
additional challenge of diet changes may cause loss of integrity in both the SAM complex and
MICOS complex, which results in the buildup of lipids in the liver, raising the risk of cirrhosis,
steatohepatitis, and NAFLD, among other hepatic conditions.
654

CONCLUSION

To broaden our understanding of age-related changes in male murine hepatic tissue, we have
established how 2D and 3D ultrastructure undergo similar reductions in mitochondrial volume
and connectivity, potentially due to an age-related loss of the MICOS complex. This is an
important age-related regulator of mitochondrial structure and could be linked to functional
changes across aging in the liver. Deletion of the MICOS complex in HepG2 cells results in
impaired calcium uptake and increased oxidative stress, highlighting its significance. We have
also established that *Sam50*, which is genetically associated with liver diseases, is maintained
across aging but lost with a high-fat diet. Since a high-fat diet can exacerbate changes in lipid
synthesis, catabolism, and mitochondrial alterations that already occur in hepatic aging (30, 83),
this suggests the confluence of aging and a high-fat diet in increasing susceptibility to liver
disease, in part due to the loss of both the MIB complex (i.e., MICOS and SAM complex).
Together, this study demonstrates dysfunctions of mitochondrial structure in both 2D and 3D,
mitochondrial calcium influx and ROS production, changes in metabolism and lipid content, and
MICOS impairments in the aging liver, with alterations in *Sam50* expression contributing to liver
diseases.
671
672
673 **EXPERIMENTAL PROCEDURES**
674
Sex As a Biological Variable:
In this study, the consideration of sex as a biological variable was integral to our experimental
design. All experimental procedures and analyses were conducted with careful consideration of
sex differences. Since minimal sex-dependent differences were observed in TEM studies, we
proceeded with a male murine model for all other studies. Sex differences in aging-related
changes and longevity can affect relative mitochondrial morphology which is a limitation of our
male model.
682
Human Cohort
All human samples were obtained from Brazilian cohorts according to the CAEE (Ethics
Appreciation Presentation Certificate) guidelines. Samples from young and old individuals were
collected and analyzed under CAEE number 77570224.2.0000.5281. Fat percentage was
calculated as a function of in-phase and out-phase quantifications via the following calculator:
https://www.ricardoromano.com/esteatoseRM (53).
689

Animal Care and Maintenance:
Per protocols previously described (140), the care and maintenance of the male C57BL/6J mice
conformed to the National Institute of Healt's guidelines for the use of laboratory animals. The
University of Iowa's Institutional Animal Care and Use Committee (IACUC) or University of
Washington IACUC approved the housing and feeding of these mice. Anesthesia was achieved
using a mixture of 5% isoflurane and 95% oxygen.

696
697 *Oil Red O*
698 OCT blocks were cut into 7 μm thick sections, affixed to glass slides, brought to room
699 temperature for 10 minutes, then stained with Oil Red O (Sigma-Aldrich), as previously
700 described (141).

701
mtDNA Content
As previously described (142), mitochondrial DNA content was quantified using real-time PCR
(RT-PCR) from DNA extracted and purified from gastrocnemius muscle tissue. Five nanograms
of DNA were used for the quantification of mitochondrial (Cox1) and nuclear (β-actin) DNA
markers. Mitochondrial DNA content was normalized to the genomic Rpl13a gene using specific
primers for Cox1 and Rpl13a.

708
709 *Bile Acid*
710 From frozen liver tissue, 100 mg of tissue homogenized in 75% ethanol was incubated for 2
711 hours and then centrifuged at 6000g for 10 minutes. Once prepared, bile was measured with
712 Mouse Total Bile Acids Assay Kit (Crystal Chem), per manufacturer's instruction.

713
714 *Triglyceride Levels*
715 As previously described (142, 143), triglyceride levels were measured in the liver and in serum
716 collected after a 6□h fast using the EnzyChrom™ Triglyceride Assay Kit (BioAssay Systems),
717 with triglycerides extraction using a solution of isopropanol and Triton X□100.

718
Quantification of TEM Micrographs and Parameters Using ImageJ
Samples were fixed in a manner to avoid any bias, per established protocols (144). Following
preparation, tissue was embedded in 100% Embed 812/Araldite resin with polymerization at 60
°C overnight. After ultrathin sections (90–100 nm) were collected, they were post-stained with
lead citrate and imaged (JEOL 1400+ at 80 kV, equipped with a GatanOrius 832 camera). The
National Institutes of Health (NIH) *ImageJ* software was used for quantification of TEM images,
as described previously (9, 145).

726
Serial Block-Face Scanning Electron Microscope (SBF-SEM) Processing of Mouse Muscle
Fibers
SBF-SEM was performed according to previously defined protocols (19, 144, 145). Anesthesia
was induced in male mice using 5% isoflurane. Post skin and hair removal, the liver was treated
with 2% glutaraldehyde in 100 mM phosphate buffer for 30 minutes, dissected into 1-mm³
cubes, and further fixed in a solution containing 2.5% glutaraldehyde, 1% paraformaldehyde, and
120 mM sodium cacodylate for 1 hour.

Fixation and subsequent steps collected onto formvar-coated slot grids (Pella, Redding CA),
stained and imaged as previously described (19, 144, 145). This includes tissue washing with
100 mM cacodylate buffer, incubation in a mixture of 3% potassium ferrocyanide and 2%
osmium tetroxide, followed by dehydration in an ascending series of acetone concentrations. The
tissues were then embedded in Epoxy Taab 812 hard resin. Sectioning and imaging of sample
was performed using a VolumeScope 2 SEM (Thermo Fisher Scientific, Waltham, MA).
Conventional TEM analysis was performed on 300–400 serial sections from each sample,
following staining and imaging protocols. Subsequently, analyzed, via imaging was performed
under low vacuum/water vapor conditions with a starting energy of 3.0 keV and beam current of
0.10 nA. Sections of 50 nm thickness were cut allowing for imaging at 10 nm $\times$ 10 nm $\times$ 50 nm
spatial resolution.

745
Segmentation and Quantification of 3D SBF-SEM Images Using Amira
SBF-SEM images were manually segmented in Amira to perform 3D reconstruction, as
described previously (19). 300-400 slices were used and analyzed by a blind individual. 250 total
mitochondria across from 3 mice were collected for each quantification. For 3D reconstruction of
cardiomyocytes, 10 cells and a total of about 200 mitochondria. Quantification of 3D structures
was performed using the Amira software with built-in parameters or previously described
measurements (19).

753
LCMS Methods for Metabolomics:
Frozen tissues were weighed, ground with a liquid nitrogen in a cryomill (Retsch) at 25 Hz for
45 seconds, before extracting tissues 40:40:20 acetonitrile: methanol: water +0.5% FA +15%
NH4HCO3 (146) with a volume of 40mL solvent per 1mg of tissue, vortexed for 15 seconds, and
incubated on dry ice for 10 minutes. Tissue samples were then centrifuged at 16,000 g for 30
minutes. The supernatants were transferred to new Eppendorf tubes and then centrifuged again at
16,000 g for 25 minutes to remove any residual debris before analysis.

761
Extracts were analyzed within 24 hours by liquid chromatography coupled to a mass
spectrometer (LC-MS). The LC–MS method was based on hydrophilic interaction
chromatography (HILIC) coupled to the Orbitrap Exploris 240 mass spectrometer (Thermo
Scientific) (147). The LC separation was performed on a xBridge BEH Amide column (2.1 x 150
mm, 3.5 μm particle size, Waters, Milford, MA). Solvent A is 95%: 5% H2O: acetonitrile with
20 mM ammonium acetate and 20mM ammonium hydroxide, and solvent B is 90%: 10%
acetonitrile: H2O with 20 mM ammonium acetate and 20mM ammonium hydroxide. The
gradient was 0 min, 90% B; 2 min, 90% B; 3 min, 75% B; 5 min, 75% B; 6 min, 75% B; 7 min,
75% B; 8 min, 70% B; 9 min, 70% B; 10 min, 50% B; 12 min, 50% B; 13 min, 25% B; 14min,
25% B; 16 min, 0% B; 18 min, 0% B; 20 min, 0% B; 21 min, 90% B; 25 min, 90% B. The
following parameters were maintained during the LC analysis: flow rate 150 mL/min, column
temperature 25 °C, injection volume 5 μL and autosampler temperature was 5 °C. For the
detection of metabolites, the mass spectrometer was operated in both negative and positive ion
mode. The following parameters were maintained during the MS analysis: resolution of 180,000
at m/z 200, automatic gain control (AGC) target at 3e6, maximum injection time of 30 ms and
scan range of m/z 70-1000. Raw LC/MS data were converted to mzXML format using the
command line "msconvert" utility (148). Data were analyzed via the EL-MAVEN software
version 12.

LCMS Methods for Lipidomic Profiling
Tissue homogenization and extraction for lipids: Tissues were homogenized using a Retsch CryoMill. The homogenate was mixed with 1 mL of Extraction Buffer containing IPA/H2O/Ethyl Acetate (30:10:60, v/v/v) and Avanti Lipidomix Internal Standard (diluted 1:1000) (Avanti Polar Lipids, Inc. Alabaster, AL). Samples were vortexed and transferred to bead mill tubes for homogenization using a VWR Bead Mill at 6000 g for 30 seconds, repeated twice. The samples were then sonicated for 5 minutes and centrifuged at 15,000 g for 5 minutes at 4°C. The upper phase was transferred to a new tube and kept at 4°C. To re-extract the tissues, another 1 mL of Extraction Buffer (30:10:60, v/v/v) was added to the tissue pellet-containing tube. The samples were vortexed, homogenized, sonicated, and centrifuged as described earlier. The supernatants from both extractions were combined, and the organic phase was dried under liquid nitrogen gas.

Sample reconstitution for lipids: The dried samples were reconstituted in 300 µL of Solvent A (IPA/ACN/H2O, 45:35:20, v/v/v). After brief vortexing, the samples were sonicated for 7 minutes and centrifuged at 15,000 g for 10 minutes at 4°C. The supernatants were transferred to clean tubes and centrifuged again for 5 minutes at 15,000 g at 4°C to remove any remaining particulates. For LC-MS lipidomic analysis, 60 µL of the sample extracts were transferred to mass spectrometry vials.

LC-MS analysis for lipids: Sample analysis was performed within 36 hours after extraction using a Vanquish UHPLC system coupled with an Orbitrap Exploris 240™ mass spectrometer equipped with a H-ESI™ ion source (all Thermo Fisher Scientific). A Waters (Milford, MA) CSH C18 column (1.0 × 150 mm × 1.7 µm particle size) was used. Solvent A consisted of ACN:H2O (60:40; v/v) with 10 mM Ammonium formate and 0.1% formic acid, while solvent B contained IPA:ACN (95:5; v/v) with 10 mM Ammonium formate and 0.1% formic acid. The mobile phase flow rate was set at 0.11 mL/min, and the column temperature was maintained at 65 °C. The gradient for solvent B was as follows: 0 min 15% (B), 0–2 min 30% (B), 2–2.5 min 48% (B), 2.5–11 min 82% (B), 11–11.01 min 99% (B), 11.01–12.95 min 99% (B), 12.95–13 min 15% (B), and 13–15 min 15% (B). Ion source spray voltages were set at 4,000 V and 3,000 V in positive and negative mode, respectively. Full scan mass spectrometry was conducted with a scan range from 200 to 1000 m/z, and AcquireX mode was utilized with a stepped collision energy of 30% with a 5% spread for fragment ion MS/MS scan.

RNA Extraction and RT-qPCR
Using TRIzol reagent (Invitrogen), total RNA was isolated from tissues and further purified with the rNeasy kit (Qiagen Inc). RNA concentration was determined by measuring absorbance at 260 nm and 280 nm using a NanoDrop 1000 spectrophotometer (NanoDrop products, Wilmington, DE, USA). Approximately 1 µg of RNA was reverse-transcribed using a High-Capacity cDNA Reverse Transcription Kit (Applied Biosciences, Carlsbad CA). Quantitative PCR (qPCR) was then performed using SYBR Green (Life Technologies, Carlsbad, CA) (149). For qPCR, 50 ng of cDNA was loaded into each well of a 384-well plate, with the reaction carried out on an ABI Prism 7900HT system (Applied Biosystems) with the following cycle: 1 cycle at 95°C for 10 min; 40 cycles of 95°C for 15 s; 59°C for 15 s, 72°C for 30 s, and 78°C for 10 s; 1 cycle of 95°C for 15 s; 1 cycle of 60°C for 15 s; and one cycle of 95°C for 15 s. GAPDH normalization was

826 used to present the data as fold changes. qPCR primers used were from previously published
827 sequences (142), as detailed in Table 2.

828
Western Blotting
Western blotting was performed as previously described (76). Briefly, following RIPA lysis
buffer (1% NP40, 150 mM NaCl, 25 mM Tris base, 0.5% sodium deoxycholate, 0.1% SDS,
1% phosphatase inhibitor cocktails #2 (Sigma P5726-1ML) and #3 (Sigma P0044-1ML), and
one cOmplete protease inhibitor tablet (Sigma 04693159001)), 3-month and 2-year tissue
samples were quantified using a BCA Assay (Thermo Scientific VLBL00GD2). Equal amounts
of proteins were run on 4%–20% Tris-glycine gels (Invitrogen WXP42012BOX). Protein was
then transferred to a nitrocellulose membrane (Li-Cor 926-31092) that was incubated with
primary antibodies overnight at 4°C: Mic60/mitofilin (Abcam ab110329), SAM50 (Proteintech
20824-1-AP), or tubulin (Novus NB100-690). Secondary antibodies [1:10,000; donkey anti-
mouse IgG (H + L) (Invitrogen A32789) and donkey anti-rabbit IgG (H + L) (Invitrogen
A32802)] were incubated with the membrane at room temperature for 1 h. Using the Li-Cor
Odyssey CLx infrared imaging system, blots were imaged.

Confocal mCherry-Mito-7 Labeling
To label the mitochondria of cardiac fibroblasts, the mCherry-Mito-7 plasmid was transfected
into the cells using a transfection reagent according to the manufacturer's instructions (150) and
as previously described (75). Briefly, following plasmid and transfection reagent dilution in
Opti-MEM medium and incubation at room temperature for 20 minutes, the dilution was added
to the culture medium of the cells, which were incubated for 24–48 hours to allow expression of
the mCherry-Mito-7 protein. Localization in fibroblasts was visualized using a Leica SP8
Confocal Microscope. Cells were washed with PBS, fixed with 4% paraformaldehyde for 10
minutes, and mounted with DAPI-containing mounting medium. Fluorescent signals were
observed using appropriate filters and recorded with a digital camera.

853
SAMM50 Analyses in Vanderbilt University Medical Center's biobank, BioVU
To examine the clinical correlates of SAMM50 gene expression in a human population, we
leveraged the BioVU biobank at Vanderbilt University Medical Center (151, 152). Genotype
data linked to deidentified electronic health records (EHR) for 85,615 individuals were assessed.
Genotype data were collected on Illumina's Multi-Ethnic Genotyping Array (MEGA) and
underwent quality control procedures as previously described (153). SAMM50 genetically-
regulated gene expression (GReX) was calculated across all genotyped individuals incorporating
training data from GTEX version 8 and utilizing PrediXcan, UTMOST, or JTI models based on
the best performance (highest performance r2 for each gene-tissue pair) **(154–157)**. MultiXcan
was then performed to collapse individual tissue results into a single cross-tissue SAMM50
GReX model (158). The BioVU population was then stratified by genetic ancestry as previously
described and within each group, SAMM50 GReX was tested for association with clinical
phenotypes mapped from ICD9/10 codes and clinical lab values extracted from the EHR using
the previously described QualityLabs pipeline (153, 159, 160). In total, we tested 1,704
phenotypes and 326 labs across 70,440 individuals of European ancestry and 1,361 phenotypes
and 265 labs across 15,175 individuals of African ancestry. PheWAS analyses included logistic
regressions for any mapped phenotype with at least 50 cases, requiring at least 2 instances of an

ICD9/10 code on unique dates to be labeled as a case. LabWAS analyses included linear
regressions for any lab values that met the QualityLabs QC criteria and had median
measurements from at least 50 individuals. Covariates for both PheWAS and LabWAS included:
principal components (PCs) 1-10 for genetic ancestry, sex, age, median age of medical record,
and genotyping batch.
876
877 *Knockdown of MIC60 and CHCHD6 in HepG2 cells.*
878 The transfection of MIC60 and CHCHD6 siRNAs into HepG2 cells was carried out using
879 Lipofectamine RNAiMax (Invitrogen) in accordance with the manufacturer's instructions and as
880 previously described (24). Following a 48-hour incubation period, the cells were utilized for
881 mitochondrial calcium ($_mCa^{2+}$) and ROS measurements.
882
Measurement of mitochondrial calcium uptake and retention capacity in HepG2 cells.
$_mCa^{2+}$ uptake retention capacity in HepG2 cells were assessed using a ratiometric Ca^{2+} sensor
Fura-FF, as detailed earlier (161), with slight modifications. In brief, cells (2.5x106) were
washed with $Ca^{2+/Mg2+}$-free DPBS (GIBCO), permeabilized in intracellular medium (ICM:
120 mM KCl, 10 mM NaCl, 1 mM KH2PO4, 20 mM HEPES-Tris, pH 7.2), and
supplemented with thapsigargin and succinate. Fura-FF (1 μM) was added at the 0 s time point,
and fluorescence emissions at 340- and 380-nm ex/510-nm em were monitored using a multi-
wavelength excitation dual-wavelength emission fluorimeter (Delta RAM, PTI). To assess the
$_mCa^{2+}$ uptake, a bolus of 5 μM Ca^{2+} and the mitochondrial uncoupler FCCP (10 μM) were added
at specified time points with continuous stirring at 37°C. To assess mCa^{2+} retention capacity,
following baseline recordings, a series of Ca^{2+} boluses (5 μM) were introduced at specified time
points. Upon reaching a steady state, 10 μM FCCP was added to collapse the $\Delta\psi m$ and release
matrix free-Ca^{2+}. The number of Ca^{2+} boluses taken up by cells was counted to calculate
mitochondrial CRC.
897
Evaluation of ROS levels
~0.2 million HepG2 cells were plated in 35 mm dishes. The next day, MIC60 and CHCHD6
siRNAs were transfected using Lipofectamine RNAiMax (Invitrogen) according to the
manufacturing instructions. After incubation for 30 hrs, cells were co-stained for 30 min at 37°C
with two different dyes for ROS detection: MitoBright ROS Deep Red (10 μM, Dojindo
Laboratories) for mitochondrial superoxide, and DCFDA (10 μM, Invitrogen) for intracellular
total ROS. Following the incubation with staining dyes, cells were washed three times with 1X
HBSS and ROS analysis was done using a confocal microscope (FV4000, Olympus Life
Science).
For mitochondrial H_2O_2 assessment, cells were stained with MitoPY1 (5 μM, Bio-Techne) for 45
min at 37°C. Cells were then washed with 1x HBSS and imaged using a confocal microscope
(FV4000, Olympus Life Science). ImageJ was used for the quantification of fluorescence
intensities.
911
Data Analysis
GraphPad Prism 10.2.3 (La Jolla, CA, USA), was used for all statistical analysis. All
experiments involving SBF-SEM and TEM data had at least three independent experiments.
Statistics were not handled by those conducting the experiments. The black bars represent the
standard error of the mean. For all analysis, one-way ANOVA was performed with tests against

each independent group and significance was assessed using Fisher's protected least significant
difference (LSD) test. *, **, ***, **** were set to show significant difference, denoting $p <$
0.05, $p < 0.01$, $p < 0.001$, and $p < 0.0001$, respectively.
920
921

AUTHOR CONTRIBUTIONS:
Zer Vue, Alexandria Murphy, and Han Le contributed equally to this work. Zer Vue, Han Le, Kit
Neikirk, Edgar Garza-Lopez, Andrea G. Marshall, Margaret Mungai, Brenita Jenkins, Larry
Vang, Heather K. Beasley, Mariaassumpta Ezedimma, Sasha Manus, Aaron Whiteside, Chanel
Harris, Amber Crabtree, Claude F. Albritton, Sydney Jamison, Mert Demirci, Ashton Oliver,
Ky'Era V. Actkins, Elma Zaganjor, Estevão Scudese, Benjamin Rodriguez, Alice Koh, Izabella
Rabago, Johnathan Moore, Desiree Nguyen, Muhammad Aftab, Benjamin Kirk, Yahang Li,
Nelson Wandira, Taseer Ahmed, Mohammad Saleem, Ashlesha Kadam, Prasanna Katti, Ho-Jin
Koh, Chantell Evans, Young Do Koo, Eric Wang, Quinton Smith, Dhanendra Tomar, Clintoria
R. Williams, Mariya Sweetwyne, Anita M. Quintana, Mark A. Phillips, David Hubert, Annet
Kirabo, Chandravanu Dash, Pooja Jadiya, André Kinder, Olujimi A. Ajijola, Tyne W. Miller-
Fleming, Melanie R. McReynolds, and Antentor Hinton, Jr., contributed to the conception,
design, data acquisition, analysis, and interpretation of data. Melanie R. McReynolds, and
Antentor Hinton, Jr. conceived and supervised this study. All authors were involved in drafting
and critically revising the manuscript for important intellectual content. All authors approved the
final version of the manuscript.
938

ACKNOWLEDGEMENTS
All authors have no competing interests.
This project was funded by the National Institute of Health (NIH) NIDDK T-32, number
DK007563 entitled Multidisciplinary Training in Molecular Endocrinology to Z.V.; National
Institute of Health (NIH) NIDDK T-32, number DK007563 entitled Multidisciplinary Training
in Molecular Endocrinology to A.C.; NSF MCB #2011577I to S.A.M.; The UNCF/Bristol-Myers
Squibb E.E. Just Faculty Fund, Career Award at the Scientific Interface (CASI Award) from
Burroughs Welcome Fund (BWF) ID # 1021868.01, BWF Ad-hoc Award, NIH Small Research
Pilot Subaward to 5R25HL106365-12 from the National Institutes of Health PRIDE Program,
DK020593, Vanderbilt Diabetes and Research Training Center for DRTC Alzheimer's Disease
Pilot & Feasibility Program. CZI Science Diversity Leadership grant number 2022- 253529 from
the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation
to A.H.J. and O.A.A.; and National Institutes of Health grant HD090061 and the Department of
Veterans Affairs Office of Research Award I01 BX005352 to J.G. Howard Hughes Medical
Institute Hanna H. Gray Fellows Program Faculty Phase (Grant# GT15655 awarded to M.R.M);
and Burroughs Wellcome Fund PDEP Transition to Faculty (Grant# 1022604 awarded to
M.R.M). National Institutes of Health Grants: R21DK119879 (to C.R.W.) and R01DK-133698
(to C.R.W.), American Heart Association Grant 16SDG27080009 (to C.R.W.) and by an
American Society of Nephrology KidneyCure Transition to Independence Grant (to C.R.W.).
NIH Grants R01HL147818, R03HL155041, and R01HL144941 (A. Kirabo). NIH Grant
R00DK120876 (D.T.), Harold S. Geneen Charitable Trust Awards Program (D.T.), Alzheimer's
Association AARG-NTF-23-1144888 (D.T.). NIH Grant R00AG065445 (P.J.), Alzheimer's
Association 24AARG-D-1191292 (P.J.), Wake ADRC REC and Development grant
P30AG072947 (P.J.). American Heart Association Grant 23POST1020344 (A.K.). American

Heart Association Grant 23CDA1053072 (M. S.). NIH K01AG062757 to (M.T.S.) The BioVU
project at VUMC is supported by numerous sources: including the NIH funded Shared
Instrumentation Grant S10OD017985 and S10RR025141; CTSA grants UL1TR002243,
UL1TR000445, and UL1RR024975 from the National Center for Advancing Translational
Sciences. Its contents are solely the responsibility of the authors and do not necessarily represent
official views of the National Center for Advancing Translational Sciences. Genomic data are
also supported by investigator-led projects that include U01HG004798, R01NS032830,
RC2GM092618, P50GM115305, U01HG006378, U19HL065962, R01HD074711; and
additional funding sources listed at https://victr.vumc.org/biovu-funding/.23CDA1053072
(M.S.). Its contents are solely the responsibility of the authors and do not necessarily represent
the official view of the NIH. The funders had no role in study design, data collection, and
analysis, decision to publish, or preparation of the manuscript. We would also like to
acknowledge the Huck Institutes' Metabolomics Core Facility (RRID:SCR_023864) for use of
the OE 240 LCMS and Drs. Imhoi Koo, Ashley Shay, and Sergei Koshkin for helpful
discussions on sample preparation and analysis. We would also like to thank UCLA investigators
for gifting us old and young human liver samples.

CONFLICT OF INTEREST
The authors declare that they have no conflict of interest.

References:

1. McBride HM, Neuspiel M, Wasiak S. Mitochondria: more than just a powerhouse. *Curr Biol*
CB. 2006;16(14):R551-560.

2. Glancy B, et al. The Functional Impact of Mitochondrial Structure Across Subcellular Scales.
Front Physiol. 2020;11. https://www.frontiersin.org/articles/10.3389/fphys.2020.541040.
Accessed December 16, 2022.

3. Picard M, et al. Mitochondria: isolation, structure and function. *J Physiol*. 2011;589(18):4413–
4421.

4. Chan DC. Fusion and fission: interlinked processes critical for mitochondrial health. *Annu Rev*
Genet. 2012;46:265–287.

5. Li M, et al. Mitochondrial Fusion Via OPA1 and MFN1 Supports Liver Tumor Cell
Metabolism and Growth. *Cells*. 2020;9(1):121.

6. Seo AY, et al. New insights into the role of mitochondria in aging: mitochondrial dynamics
and more. *J Cell Sci*. 2010;123(15):2533–2542.

7. Bratic A, Larsson N-G. The role of mitochondria in aging. *J Clin Invest*. 2013;123(3):951–
957.

8. Haas RH. Mitochondrial Dysfunction in Aging and Diseases of Aging. *Biology*. 2019;8(2):48.

9. Lam J, et al. A Universal Approach to Analyzing Transmission Electron Microscopy with
ImageJ. *Cells*. 2021;10(9):2177.

10. Collins HE, et al. Mitochondrial Morphology and Mitophagy in Heart Diseases: Qualitative
and Quantitative Analyses Using Transmission Electron Microscopy. *Front Aging*. 2021;2.
https://www.frontiersin.org/articles/10.3389/fragi.2021.670267. Accessed September 8, 2022.

11. Cogliati S, Enriquez JA, Scorrano L. Mitochondrial cristae: where beauty meets
functionality. *Trends Biochem Sci*. 2016;41(3):261–273.

12. Jenkins BC, et al. Mitochondria in disease: changes in shapes and dynamics. *Trends Biochem
Sci*. 2024;49(4):346–360.

13. Long Q, et al. Modeling of Mitochondrial Donut Formation. *Biophys J*. 2015;109:892–9.

14. Liu X, Hajnóczky G. Altered fusion dynamics underlie unique morphological changes in
mitochondria during hypoxia–reoxygenation stress. *Cell Death Differ*. 2011;18(10):1561–1572.

15. Vincent AE, et al. Mitochondrial Nanotunnels. *Trends Cell Biol*. 2017;27(11):787–799.

16. Tyumentsev MA, et al. Mitochondria with Morphology Characteristic for Alzheimer's
Disease Patients Are Found in the Brain of OXYS Rats. *Biochem Mosc*. 2018;83(9):1083–1088.

17. Wakabayashi T. Megamitochondria formation □ physiology and pathology. *J Cell Mol Med*.
2002;6(4):497–538.

18. Gautier MK, Ginsberg SD. A method for quantification of vesicular compartments within
cells using 3D reconstructed confocal z-stacks: Comparison of ImageJ and Imaris to count early
endosomes within basal forebrain cholinergic neurons. *J Neurosci Methods*. 2021;350:109038.

19. Garza-Lopez E, et al. Protocols for Generating Surfaces and Measuring 3D Organelle
Morphology Using Amira. *Cells*. 2022;11(1):65.

20. Courson JA, et al. Serial Block-Face Scanning Electron Microscopy (SBF-SEM) of
Biological Tissue Samples. *J Vis Exp JoVE*. 2021;(169):10.3791/62045.

21. Marshall AG, et al. Serial Block Face-Scanning Electron Microscopy as a Burgeoning
Technology. *Adv Biol*. 2023;e2300139.

22. Russell MRG, et al. 3D correlative light and electron microscopy of cultured cells using
serial blockface scanning electron microscopy. *J Cell Sci*. 2017;130(1):278–291.

23. Vue Z, et al. 3D reconstruction of murine mitochondria reveals changes in structure during
aging linked to the MICOS complex. *Aging Cell*. 2023;22(12):e14009.

24. Vue Z, et al. The MICOS Complex Regulates Mitochondrial Structure and Oxidative Stress
During Age-Dependent Structural Deficits in the Kidney [preprint]. 2024;2024.06.09.598108.

25. Gagliano N, Grizzi F, Annoni G. Mechanisms of Aging and Liver Functions. *Dig Dis*.
2007;25(2):118–123.

26. Hunt NJ, et al. Hallmarks of Aging in the Liver. *Comput Struct Biotechnol J*. 2019;17:1151–
1161.

27. Kalra A, et al. Physiology, Liver. *StatPearls*. Treasure Island (FL): StatPearls Publishing;
2024.

28. Reddy SR, et al. Outcomes of Young Patients With Alcoholic Cirrhosis After First
Hospitalization for Cirrhosis: A Carilion Clinic Experience. *Cureus*;13(7):e16695.

29. Kim H, Kisseleva T, Brenner DA. Aging and liver disease. *Curr Opin Gastroenterol*.
2015;31(3):184–191.

30. Gong Z, et al. Hepatic lipid metabolism and non-alcoholic fatty liver disease in aging. *Mol*
Cell Endocrinol. 2017;455:115–130.

31. Chung KW. Advances in Understanding of the Role of Lipid Metabolism in Aging. *Cells*.
2021;10(4):880.

32. Hoffmann JJ, Becker T. Crosstalk between Mitochondrial Protein Import and Lipids. *Int J*
Mol Sci. 2022;23(9):5274.

33. Chen G, Kroemer G, Kepp O. Mitophagy: An Emerging Role in Aging and Age-Associated
Diseases. *Front Cell Dev Biol*. 2020;8.
https://www.frontiersin.org/articles/10.3389/fcell.2020.00200. Accessed December 21, 2022.

34. Nishikawa T, et al. A switch in the source of ATP production and a loss in capacity to
perform glycolysis are hallmarks of hepatocyte failure in advance liver disease. *J Hepatol*.
2014;60(6):1203–1211.

35. Middleton P, Vergis N. Mitochondrial dysfunction and liver disease: role, relevance, and
potential for therapeutic modulation. *Ther Adv Gastroenterol*. 2021;14:17562848211031394.

36. Wang D, et al. Research progress on the mitochondrial mechanism of age-related non-
alcoholic fatty liver. *World J Gastroenterol*. 2023;29(13):1982–1993.

37. Shutt TE, McBride HM. Staying cool in difficult times: Mitochondrial dynamics, quality
control and the stress response. *Biochim Biophys Acta BBA - Mol Cell Res*. 2013;1833(2):417–
424.

38. Ježek J, Cooper KF, Strich R. Reactive Oxygen Species and Mitochondrial Dynamics: The
Yin and Yang of Mitochondrial Dysfunction and Cancer Progression. *Antioxidants*. 2018;7(1).
https://doi.org/10.3390/antiox7010013.

39. Abrisch RG, et al. Fission and fusion machineries converge at ER contact sites to regulate
mitochondrial morphology. *J Cell Biol*. 2020;219(4):e201911122.

40. Youle RJ, van der Bliek AM. Mitochondrial Fission, Fusion, and Stress. *Science*.
2012;337(6098):1062–1065.

41. Belenguer P, Pellegrini L. The dynamin GTPase OPA1: More than mitochondria? *Biochim*
Biophys Acta BBA - Mol Cell Res. 2013;1833(1):176–183.

42. Chen H, et al. Mitofusins Mfn1 and Mfn2 coordinately regulate mitochondrial fusion and are
essential for embryonic development. *J Cell Biol*. 2003;160(2):189–200.

43. Hu C, et al. OPA1 and MICOS Regulate mitochondrial crista dynamics and formation. *Cell*
Death Dis. 2020;11(10):1–17.

44. Favaro G, et al. DRP1-mediated mitochondrial shape controls calcium homeostasis and
muscle mass. *Nat Commun*. 2019;10(1):1–17.

45. Anand R, et al. The i-AAA protease YME1L and OMA1 cleave OPA1 to balance
mitochondrial fusion and fission. *J Cell Biol*. 2014;204(6):919–929.

46. Wai T, Langer T. Mitochondrial Dynamics and Metabolic Regulation. *Trends Endocrinol*
Metab. 2016;27(2):105–117.

47. Takeichi Y, et al. Non-alcoholic fatty liver disease in mice with hepatocyte-specific deletion
of mitochondrial fission factor. *Diabetologia*. 2021;64:2092–2107.

48. Galloway C, et al. Decreasing mitochondrial fission alleviates hepatic steatosis in a murine
model of nonalcoholic fatty liver disease. *Am J Physiol Gastrointest Liver Physiol*. 2014;307
6:632–41.

49. Li Z, et al. The role of SAMM50 in non-alcoholic fatty liver disease: from genetics to
mechanisms. *FEBS Open Bio*. 2021;11(7):1893–1906.

50. Lionello S, Marzaro G, Martinvalet D. SAM50, a side door to the mitochondria: The case of
cytotoxic proteases. *Pharmacol Res*. 2020;160:105196.

51. Shao B, et al. Ablation of Sam50 is associated with fragmentation and alterations in
metabolism in murine and human myotubes. *BioRxiv Prepr Serv Biol*. 2023;2023.05.20.541602.

52. Shao B, et al. Ablation of Sam50 is associated with fragmentation and alterations in
metabolism in murine and human myotubes. *J Cell Physiol*. 2024;n/a(n/a).
https://doi.org/10.1002/jcp.31293.

53. Tang A, et al. Nonalcoholic Fatty Liver Disease: MR Imaging of Liver Proton Density Fat
Fraction to Assess Hepatic Steatosis. *Radiology*. 2013;267(2):422–431.

54. Ma X, et al. Perspectives on Mitochondria–ER and Mitochondria–Lipid Droplet Contact in
Hepatocytes and Hepatic Lipid Metabolism. *Cells*. 2021;10(9):2273.

55. Simões ICM, et al. Mitochondria in non-alcoholic fatty liver disease. *Int J Biochem Cell Biol*.
2018;95:93–99.

56. Chiang JYL, Ferrell JM. Bile Acid Metabolism in Liver Pathobiology. *Gene Expr*.
2018;18(2):71–87.

57. TOMIZAWA M, et al. Triglyceride is strongly associated with nonalcoholic fatty liver
disease among markers of hyperlipidemia and diabetes. *Biomed Rep*. 2014;2(5):633–636.

58. Semova I, Biddinger SB. TRIGLYCERIDES IN NAFLD: GUILTY UNTIL PROVEN
INNOCENT. *Trends Pharmacol Sci*. 2021;42(3):183–190.

59. Yanai S, Endo S. Functional Aging in Male C57BL/6J Mice Across the Life-Span: A
Systematic Behavioral Analysis of Motor, Emotional, and Memory Function to Define an Aging
Phenotype. *Front Aging Neurosci*. 2021;13.
https://www.frontiersin.org/articles/10.3389/fnagi.2021.697621. Accessed February 8, 2023.

60. Robertson GL, et al. DRP1-mediated mitochondrial fission is essential to maintain cristae
morphology and bioenergetics [preprint]. 2022;2021.12.31.474637.

61. Giacomello M, Pellegrini L. The coming of age of the mitochondria–ER contact: a matter of
thickness. *Cell Death Differ*. 2016;23(9):1417–1427.

62. Vincent AE, et al. Quantitative 3D Mapping of the Human Skeletal Muscle Mitochondrial
Network. *Cell Rep*. 2019;26(4):996-1009.e4.

63. Ilacqua N, et al. A three-organelle complex made by wrappER contacts with peroxisomes
and mitochondria responds to liver lipid flux changes. *J Cell Sci*. 2021;135(5):jcs259091.

64. Ilacqua N, Anastasia I, Pellegrini L. Isolation and analysis of fractions enriched in WrappER-
associated mitochondria from mouse liver. *STAR Protoc*. 2021;2(3):100752.

65. Ilacqua N, et al. Expression of Synj2bp in mouse liver regulates the extent of wrappER-
mitochondria contact to maintain hepatic lipid homeostasis. *Biol Direct*. 2022;17(1):37.

66. Surmacki JM, Abramczyk H. Confocal Raman imaging reveals the impact of retinoids on
human breast cancer via monitoring the redox status of cytochrome c. *Sci Rep*.
2023;13(1):15049.

67. McReynolds MR, et al. NAD+ flux is maintained in aged mice despite lower tissue
concentrations. *Cell Syst*. 2021;12(12):1160-1172.e4.

68. Goodman RP, Calvo SE, Mootha VK. Spatiotemporal compartmentalization of hepatic
NADH and NADPH metabolism. *J Biol Chem*. 2018;293(20):7508–7516.

69. Ueda N. Ceramide-Induced Apoptosis in Renal Tubular Cells: A Role of Mitochondria and
Sphingosine-1-Phoshate. *Int J Mol Sci*. 2015;16(3):5076–5124.

70. Fugio LB, Coeli-Lacchini FB, Leopoldino AM. Sphingolipids and Mitochondrial Dynamic.
Cells. 2020;9(3):581.

71. Parra V, et al. Changes in mitochondrial dynamics during ceramide-induced cardiomyocyte early apoptosis. *Cardiovasc Res*. 2008;77(2):387–397.

72. Longo N, Frigeni M, Pasquali M. CARNITINE TRANSPORT AND FATTY ACID OXIDATION. *Biochim Biophys Acta*. 2016;1863(10):2422–2435.

73. Li H, et al. Mic60/Mitofilin determines MICOS assembly essential for mitochondrial dynamics and mtDNA nucleoid organization. *Cell Death Differ*. 2016;23(3):380–392.

74. Tezze C, et al. Age-Associated Loss of OPA1 in Muscle Impacts Muscle Mass, Metabolic Homeostasis, Systemic Inflammation, and Epithelial Senescence. *Cell Metab*. 2017;25(6):1374-1389.e6.

75. Vue Z, et al. Three-Dimensional Mitochondria Reconstructions of Murine Cardiac Muscle Changes in Size Across Aging. *Am J Physiol-Heart Circ Physiol*. [published online ahead of print: August 25, 2023]. https://doi.org/10.1152/ajpheart.00202.2023.

76. Vue Z, et al. 3D reconstruction of murine mitochondria reveals changes in structure during aging linked to the MICOS complex. *Aging Cell*. 2023;22(12):e14009.

77. Körner C, et al. The C-terminal domain of Fcj1 is required for formation of crista junctions and interacts with the TOB/SAM complex in mitochondria. *Mol Biol Cell*. 2012;23(11):2143–2155.

78. Kozjak V, et al. An Essential Role of Sam50 in the Protein Sorting and Assembly Machinery of the Mitochondrial Outer Membrane *. *J Biol Chem*. 2003;278(49):48520–48523.

79. Tang J, et al. Sam50–Mic19–Mic60 axis determines mitochondrial cristae architecture by mediating mitochondrial outer and inner membrane contact. *Cell Death Differ*. 2020;27(1):146–160.

80. Ding C, et al. Mitofilin and CHCHD6 physically interact with Sam50 to sustain cristae structure. *Sci Rep*. 2015;5(1):1–11.

81. Chen L, et al. Loss of Sam50 in hepatocytes induces cardiolipin☐dependent mitochondrial membrane remodeling to trigger mtDNA release and liver injury. *Hepatology*. 2022;76(5):1389.

82. Tsuru H, et al. HFD-induced hepatic lipid accumulation and inflammation are decreased in Factor D deficient mouse. *Sci Rep*. 2020;10:17593.

83. Lian C-Y, et al. High fat diet-triggered non-alcoholic fatty liver disease: A review of proposed mechanisms. *Chem Biol Interact*. 2020;330:109199.

84. Zheng P, et al. High-fat diet causes mitochondrial damage and downregulation of mitofusin-2 and optic atrophy-1 in multiple organs. *J Clin Biochem Nutr*. 2023;73(1):61–76.

85. De Nicolo B, et al. Calcium and Reactive Oxygen Species Signaling Interplays in Cardiac Physiology and Pathologies. *Antioxidants*. 2023;12(2):353.

86. Winslow JL, Hollenberg MJ, Lea PJ. Resolution limit of serial sections for 3D reconstruction of tubular cristae in rat liver mitochondria. *J Electron Microsc Tech*. 1991;18(3):241–248.

87. Mannella CA. Structure and dynamics of the mitochondrial inner membrane cristae. *Biochim Biophys Acta BBA - Mol Cell Res*. 2006;1763(5):542–548.

88. Mannella CA, et al. Electron microscopic tomography of rat-liver mitochondria and their interactions with the endoplasmic reticulum. *BioFactors*. 1998;8(3–4):225–228.

89. Das S, et al. Mitochondrial morphology and dynamics in hepatocytes from normal and ethanol-fed rats. *Pflugers Arch*. 2012;464(1):101–109.

90. Shami GJ, et al. Three-dimensional ultrastructure of giant mitochondria in human non-alcoholic fatty liver disease. *Sci Rep*. 2021;11:3319.

91. Daum B, et al. Age-dependent dissociation of ATP synthase dimers and loss of inner-membrane cristae in mitochondria. *Proc Natl Acad Sci*. 2013;110(38):15301–15306.

92. Labbe G, Pessayre D, Fromenty B. Drug-induced liver injury through mitochondrial dysfunction: mechanisms and detection during preclinical safety studies. *Fundam Clin Pharmacol*. 2008;22(4):335–353.

93. Altamirano J, et al. A histologic scoring system for prognosis of patients with alcoholic hepatitis. *Gastroenterology*. 2014;146(5):1231-1239.e1–6.

94. Palma E, et al. Dynamin-1-Like Protein Inhibition Drives Megamitochondria Formation as an Adaptive Response in Alcohol-Induced Hepatotoxicity. *Am J Pathol*. 2019;189(3):580–589.

95. Zhou H, et al. DNA-PKcs promotes alcohol-related liver disease by activating Drp1-related mitochondrial fission and repressing FUNDC1-required mitophagy. *Signal Transduct Target Ther*. 2019;4:56.

96. Yamada T, et al. Prevention and regression of megamitochondria and steatosis by blocking mitochondrial fusion in the liver. *iScience*. 2022;25(4):103996.

97. Tandler B, Hoppel CL. Giant Mitochondria (Megamitochondria). In: Lennarz WJ, Lane MD, eds. *Encyclopedia of Biological Chemistry (Second Edition)*. Waltham: Academic Press; 2013:375–376.

98. Palma E, et al. Perturbations in mitochondrial dynamics are closely involved in the progression of Alcoholic Liver Disease. *Alcohol Clin Exp Res*. [published online ahead of print: 2020]. https://doi.org/10.1111/acer.14299.

99. Palma E, et al. Perturbation of mitochondrial dynamics in Alcoholic Liver Disease. *J World Mitochondria Soc*. 2016;2. https://doi.org/10.18143/JWMS_V2I2_1948.

100. Wang D, Wei Y, Pagliassotti MJ. Saturated Fatty Acids Promote Endoplasmic Reticulum Stress and Liver Injury in Rats with Hepatic Steatosis. *Endocrinology*. 2006;147(2):943–951.

1202 101. Renne MF, Hariri H. Lipid Droplet-Organelle Contact Sites as Hubs for Fatty Acid
1203 Metabolism, Trafficking, and Metabolic Channeling. *Front Cell Dev Biol*. 2021;9.
1204 https://www.frontiersin.org/articles/10.3389/fcell.2021.726261. Accessed February 12, 2023.

1205 102. Kim H, et al. Structural basis for mitoguardin-2 mediated lipid transport at ER-
1206 mitochondrial membrane contact sites. *Nat Commun*. 2022;13(1):3702.

1207 103. Marchi S, Patergnani S, Pinton P. The endoplasmic reticulum–mitochondria connection:
1208 One touch, multiple functions. *Biochim Biophys Acta BBA - Bioenerg*. 2014;1837(4):461–469.

1209 104. Anastasia I, et al. Mitochondria-rough-ER contacts in the liver regulate systemic lipid
1210 homeostasis. *Cell Rep*. 2021;34(11):108873.

1211 105. Wang J, et al. Mutual interaction between endoplasmic reticulum and mitochondria in
1212 nonalcoholic fatty liver disease. *Lipids Health Dis*. 2020;19(1):72.

1213 106. Huynen MA, et al. Evolution and structural organization of the mitochondrial contact site
1214 (MICOS) complex and the mitochondrial intermembrane space bridging (MIB) complex.
1215 *Biochim Biophys Acta BBA-Mol Cell Res*. 2016;1863(1):91–101.

1216 107. Ott C, et al. Sam50 Functions in Mitochondrial Intermembrane Space Bridging and
1217 Biogenesis of Respiratory Complexes. *Mol Cell Biol*. 2012;32(6):1173–1188.

1218 108. Gagliano Taliun SA, et al. Exploring and visualizing large-scale genetic associations by
1219 using PheWeb. *Nat Genet*. 2020;52(6):550–552.

1220 109. Kanai M, et al. Genetic analysis of quantitative traits in the Japanese population links cell
1221 types to complex human diseases. *Nat Genet*. 2018;50(3):390–400.

1222 110. Kurki MI, et al. FinnGen: Unique genetic insights from combining isolated population and
1223 national health register data [preprint]. 2022;2022.03.03.22271360.

1224 111. Duchen MR, Szabadkai G. Roles of mitochondria in human disease. *Essays Biochem*.
1225 2010;47:115–137.

1226 112. Alonso MT, et al. Calcium microdomains in mitochondria and nucleus. *Cell Calcium*.
1227 2006;40(5–6):513–525.

1228 113. Bertero E, Maack C. Calcium Signaling and Reactive Oxygen Species in Mitochondria.
1229 *Circ Res*. 2018;122(10):1460–1478.

1230 114. Hoek JB, et al. Calcium ion-dependent signalling and mitochondrial dysfunction:
1231 mitochondrial calcium uptake during hormonal stimulation in intact liver cells and its implication
1232 for the mitochondrial permeability transition. *Biochim Biophys Acta BBA - Mol Basis Dis*.
1233 1995;1271(1):93–102.

1234 115. Csordás G, Weaver D, Hajnóczky G. Endoplasmic Reticulum–Mitochondrial Contactology:
1235 Structure and Signaling Functions. *Trends Cell Biol*. 2018;28(7):523–540.

1236 116. Egnatchik RA, et al. ER calcium release promotes mitochondrial dysfunction and hepatic
1237 cell lipotoxicity in response to palmitate overload. *Mol Metab*. 2014;3(5):544–553.

1238 117. Kim YS, Kim SG. Endoplasmic reticulum stress and autophagy dysregulation in alcoholic
1239 and non-alcoholic liver diseases. *Clin Mol Hepatol*. 2020;26(4):715–727.

1240 118. Amaya MJ, Nathanson MH. Calcium Signaling in the Liver. *Compr Physiol*.
1241 2013;3(1):515–539.

1242 119. Modi S, et al. Miro clusters regulate ER-mitochondria contact sites and link cristae
1243 organization to the mitochondrial transport machinery. *Nat Commun*. 2019;10(1):4399.

1244 120. Cichoż-Lach H, Michalak A. Oxidative stress as a crucial factor in liver diseases. *World J*
1245 *Gastroenterol WJG*. 2014;20(25):8082–8091.

1246 121. Li P-L, Gulbins E. Bioactive Lipids and Redox Signaling: Molecular Mechanism and
1247 Disease Pathogenesis. *Antioxid Redox Signal*. 2018;28(10):911–915.

1248 122. Barja G. The mitochondrial free radical theory of aging. *Prog Mol Biol Transl Sci*.
1249 2014;127:1–27.

1250 123. Navarro A, Boveris A. Rat brain and liver mitochondria develop oxidative stress and lose
1251 enzymatic activities on aging. *Am J Physiol-Regul Integr Comp Physiol*. 2004;287(5):R1244–
1252 R1249.

1253 124. Maeso-Díaz R, et al. Effects of aging on liver microcirculatory function and sinusoidal
1254 phenotype. *Aging Cell*. 2018;17(6):e12829.

1255 125. Nahata M, et al. Decline in Liver Mitochondria Metabolic Function Is Restored by
1256 Hochuekkito Through Sirtuin 1 in Aged Mice With Malnutrition. *Front Physiol*. 2022;13.
1257 https://www.frontiersin.org/articles/10.3389/fphys.2022.848960. Accessed February 12, 2023.

1258 126. Barazzoni R, Short KR, Nair KS. Effects of Aging on Mitochondrial DNA Copy Number
1259 and Cytochromec Oxidase Gene Expression in Rat Skeletal Muscle, Liver, and Heart *. *J Biol*
1260 *Chem*. 2000;275(5):3343–3347.

1261 127. Serviddio G, et al. Bioenergetics in aging: mitochondrial proton leak in aging rat liver,
1262 kidney and heart. *Redox Rep*. 2007;12(1–2):91–95.

1263 128. Moreira OC, et al. Mitochondrial Function and Mitophagy in the Elderly: Effects of
1264 Exercise. *Oxid Med Cell Longev*. 2017;2017:2012798.

1265 129. Guo Y, et al. Quantitative Proteomic and Functional Analysis of Liver Mitochondria from
1266 High Fat Diet (HFD) Diabetic Mice. *Mol Cell Proteomics MCP*. 2013;12(12):3744–3758.

1267 130. Graham DS, et al. Ability of high fat diet to induce liver pathology correlates with the level
1268 of linoleic acid and Vitamin E in the diet. *PloS One*. 2023;18(6):e0286726.

1269 131. Kim HJ, et al. A high fat diet fosters elevated bisretinoids. *J Biol Chem*.
1270 2023;299(6):104784.

1271 132. Parlakgül G, et al. Spatial mapping of hepatic ER and mitochondria architecture reveals
1272 zonated remodeling in fasting and obesity. *Nat Commun*. 2024;15(1):3982.

1273 133. Arruda AP, et al. Chronic enrichment of hepatic endoplasmic reticulum-mitochondria
1274 contact leads to mitochondrial dysfunction in obesity. *Nat Med*. 2014;20(12):1427–1435.

1275 134. Softic S, et al. Dietary Sugars Alter Hepatic Fatty Acid Oxidation via Transcriptional and
1276 Post-translational Modifications of Mitochondrial Proteins. *Cell Metab*. 2019;30 4:735–7534.

1277 135. Lockman KA, et al. Oxidative stress rather than triglyceride accumulation is a determinant
1278 of mitochondrial dysfunction in in vitro models of hepatic cellular steatosis. *Liver Int Off J Int*
1279 *Assoc Study Liver*. 2012;32(7):1079–1092.

1280 136. Dong J, et al. Mic19 depletion impairs endoplasmic reticulum-mitochondrial contacts and
1281 mitochondrial lipid metabolism and triggers liver disease. *Nat Commun*. 2024;15(1):168.

1282 137. Paradies G, et al. Oxidative stress, cardiolipin and mitochondrial dysfunction in
1283 nonalcoholic fatty liver disease. *World J Gastroenterol WJG*. 2014;20(39):14205–14218.

1284 138. Liu J, et al. Cardiolipin inhibitor ameliorates the non-alcoholic steatohepatitis through
1285 suppressing NLRP3 inflammasome activation. *Eur Rev Med Pharmacol Sci*. 2019;23(18):8158–
1286 8167.

1287 139. Alves-Bezerra M, Cohen DE. Triglyceride metabolism in the liver. *Compr Physiol*.
1288 2017;8(1):1–8.

1289 140. Hinton A, et al. ATF4 Dependent Increase in Mitochondrial-Endoplasmic Reticulum
1290 Tethering Following OPA1 Deletion in Skeletal Muscle. *Journal of Cellular Physiology*.
1291 [published online ahead of print: 2024]. https://doi.org/10.1101/2022.09.12.507669.

1292 141. Stahl EC, et al. Inflammation and Ectopic Fat Deposition in the Aging Murine Liver Is
1293 Influenced by CCR2. *Am J Pathol*. 2020;190(2):372–387.

1294 142. Pereira RO, et al. OPA 1 deficiency promotes secretion of FGF 21 from muscle that
1295 prevents obesity and insulin resistance. *EMBO J*. 2017;36(14):2126–2145.

1296 143. Tam J, et al. Peripheral CB1 cannabinoid receptor blockade improves cardiometabolic risk
1297 in mouse models of obesity. *J Clin Invest*. 2010;120(8):2953–2966.

1298 144. Hinton A, et al. A Comprehensive Approach to Sample Preparation for Electron
1299 Microscopy and the Assessment of Mitochondrial Morphology in Tissue and Cultured Cells. *Adv*
1300 *Biol*. 2023;e2200202.

1301 145. Neikirk K, et al. Systematic Transmission Electron Microscopy-Based Identification and 3D
1302 Reconstruction of Cellular Degradation Machinery. *Adv Biol*. 2023;7(6):2200221.

1303 146. Lu W, et al. Extraction and Quantitation of Nicotinamide Adenine Dinucleotide Redox
1304 Cofactors. *Antioxid Redox Signal*. 2018;28(3):167–179.

1305 147. Wang L, et al. Peak Annotation and Verification Engine for Untargeted LC–MS
1306 Metabolomics. *Anal Chem*. 2019;91(3):1838–1846.

1307 148. Adusumilli R, Mallick P. Data Conversion with ProteoWizard msConvert. In: Comai L,
1308 Katz JE, Mallick P, eds. *Proteomics: Methods and Protocols*. New York, NY: Springer;
1309 2017:339–368.

1310 149. Boudina S, et al. Mitochondrial energetics in the heart in obesity-related diabetes: direct
1311 evidence for increased uncoupled respiration and activation of uncoupling proteins. *Diabetes*.
1312 2007;56(10):2457–2466.

1313 150. Olenych SG, et al. The fluorescent protein color palette. *Curr Protoc Cell Biol*.
1314 2007;Chapter 21:Unit 21.5.

1315 151. Roden DM, et al. Development of a large-scale de-identified DNA biobank to enable
1316 personalized medicine. *Clin Pharmacol Ther*. 2008;84(3):362–369.

1317 152. Danciu I, et al. Secondary use of clinical data: the Vanderbilt approach. *J Biomed Inform*.
1318 2014;52:28–35.

1319 153. Dennis JK, et al. Clinical laboratory test-wide association scan of polygenic scores
1320 identifies biomarkers of complex disease. *Genome Med*. 2021;13(1):6.

1321 154. GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across human
1322 tissues. *Science*. 2020;369(6509):1318–1330.

1323 155. Gamazon ER, et al. A gene-based association method for mapping traits using reference
1324 transcriptome data. *Nat Genet*. 2015;47(9):1091–1098.

1325 156. Hu Y, et al. A statistical framework for cross-tissue transcriptome-wide association
1326 analysis. *Nat Genet*. 2019;51(3):568–576.

1327 157. Zhou D, et al. A unified framework for joint-tissue transcriptome-wide association and
1328 Mendelian randomization analysis. *Nat Genet*. 2020;52(11):1239–1246.

1329 158. Barbeira AN, et al. Integrating predicted transcriptome from multiple tissues improves
1330 association detection. *PLoS Genet*. 2019;15(1):e1007889.

1331 159. Denny JC, et al. PheWAS: demonstrating the feasibility of a phenome-wide scan to
1332 discover gene-disease associations. *Bioinforma Oxf Engl*. 2010;26(9):1205–1210.

1333 160. Carroll RJ, Bastarache L, Denny JC. R PheWAS: data analysis and plotting tools for
1334 phenome-wide association studies in the R environment. *Bioinforma Oxf Engl*.
1335 2014;30(16):2375–2376.

1336 161. Tomar D, et al. MCUR1 Is a Scaffold Factor for the MCU Complex Function and Promotes
1337 Mitochondrial Bioenergetics. *Cell Rep*. 2016;15(8):1673–1685.

1338
1339

1340

1341   **FIGURE LEGEND:**

1342

Graphical Abstract: Liver aging causes metabolic, lipidomic, and mitochondrial structural
alterations, reflecting age-dependent losses in the MICOS complex. Diet-dependent losses of the
SAM complex underlie genetic disease associations and mitochondrial structure.

Supplementary Figures:

Supplemental Figure 1: Heterogeneity in Mitochondrial Quantification Across Samples (A)
Distribution of mitochondria for mouse heterogeneity (3 mice, each 250 mitochondria surveyed)
in mitochondrial volume**, (B)** surface area, **(C)** perimeter, **(D)** sphericity, and **(E)** complexity
index in young and old liver tissue.

1351

1352   **Supplemental Figure 2: Global metabolomic and lipidomic profiling revealed metabolic**
1353   **dysregulation and disruptions in lipid classes with age in liver tissues.** **(A)** Metabolic
1354   pathway analysis and **(B)** PCA plot analysis for metabolomics. **(C)** Lipid class enrichment **(D)**
1355   and lipid chain length enrichment based on comparison between young and old livers.

1356

Supplemental Figure 3: Phenome-wide and clinical lab-wide scan for SAMM50 genetically-
regulated gene expression (GReX) in a medical biobank. Clinical phenotypes and laboratory
values from BioVU participants were extracted from Vanderbilt's de-identified electronic health
record database (n= 85,615, top left panel). Genetically-regulated gene expression for SAMM50
was calculated in BioVU participants using models built from the GTEx version 8 data (top right
panel), which contains genotype data matched to RNA-Seq data from 838 donors across 49
tissues. Imputed gene expression was calculated and tested for association across up to 1,704
phenotypes and 329 clinical lab tests using logistic and linear regression models (bottom panel),
accounting for genetic ancestry (principle components/PC 1-10), sex, age, median age of medical
record and genotyping batch.

1367

Supplementary Material:

Video 1: 3D structure of 3-month murine liver tissue visualized to show volumetric changes.

Video 2: 3D structure of 2-year murine liver tissue visualized to show volumetric changes.

Video 3: 3D structure of 3-month murine liver tissue visualized to show complexity changes.

Video 4: 3D structure of 2-year murine liver tissue visualized to show complexity changes.

SFile 1: Phenome-wide association MultiXCan SAMM50 genetically-regulated gene expression
(GReX) in individuals of European ancestry.

1375   **SFile 2:** Laboratory-wide association MultiXCan SAMM50 genetically-regulated gene
1376   expression (GReX) in individuals of European ancestry.

1377   **SFile 3:** Phenome-wide association MultiXCan SAMM50 genetically-regulated gene expression
1378   (GReX) in individuals of African ancestry.

1379   **SFile 4:** Laboratory-wide association MultiXCan SAMM50 genetically-regulated gene
1380   expression (GReX) in individuals of African ancestry.

1381   **Tables**:

Genetic Ancestry	European	African
N	70,404	15,175
Sex (%M)	44.10%	38.40%
Current Age	57.04 ± 22.42	46.48 ± 21.39
Median Age of Medical Record	48.43 ± 22.24	38.32 ± 21.41
Visits	61.18 ± 75.14	55.02 ± 80.83

1382

1383   **Table 1: Demographics for BioVU Cohort.** Genetic Ancestry Abbreviations: (EA): European
1384   ancestry, (AA): African ancestry.

1385

1386

Gene	Primers	
Opa1	Forward	5'-ACCAGGAGACTGTGTCAA-3'
	Reverse	5'-TCTTCAAATAAACGCAGAGGTG-3'
Chchd3	Forward	5'-GAAAAGAATCCAGGCCCTTCCACGCGC-3'
	Reverse	5'-CAGTGCCTAGCACTTGGCACAACCAGGAA-3'
Chchd6	Forward	5'-CTCAGCATGGACCTGGTAGGCACTGGGC-3'
	Reverse	5'-GCCTCAATTCCCACATGGAGAAAGTGGC-3'
Mitofilin	Forward	5'-CCTCCGGCAGTGTTCACCTAGTAACCCCTT-3'
	Reverse	5'-TCGCCCGTCGACCTTCAGCACTGAAAACCTAT-3'

1387
1388 **Table 2**: qPCR Primers Used
1389

1390

Figure 1: Comparative analyses of liver lipid content and morphology in young and old
patients. Cross-sectional imaging of in-phase (left) and out-of-phase (right) liver anatomy data
from (**A**) males under 55 years old (aged 14–53 years old; n = 10), (**B**) females under 55 years
old (aged 17–41 years old; n = 10), (**C**) males over 60 years old (aged 60–85 years old; n = 10),
and (**D**) females over 60 years old (aged 64–96 years old; n = 10). (**E**) Computed fat percentage
in males, (**F**) females, and (**G**) combined young and old cohorts. (**H**) Oil Red O staining at 20X
and (**I**) 40X magnification in young and (**J-K**) old samples. (**L**) Quantification of Oil Red O
staining per area in 3-month and 2-year murine liver samples (n=10). (**M**) Quantifications of
liver weight relative to total body weight as a percent (n=10), (**N**) relative reverse transcription-
quantitative polymerase chain reaction mtDNA content (n=8), (**O**) bile acids concentrations
(n=10), (**P**) liver triglycerides concentration (n=8), (**Q**) and serum triglycerides concentration
(n=8). For all panels, error bars indicate SEM, Mann–Whitney tests were used for statistical
analysis, and significance value indicate *$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$, ****$P \leq 0.0001$,
and ns indicates non-significant.

1405

Figure 2: Transmission Electron Microscopy and Serial Block Face-Scanning Electron
Microscopy Shows Changes in Mitochondria Murine Liver Across Aging. (**A**)
Representative transmission electron micrographs in males and (**B**) females from 3-month and
(**C-D**) 2-year murine liver tissue. (**E**) Mitochondrial quantifications of male mitochondria
number (n=21, 3-months; n=24, 2-years) (**F**) mitochondrial area (n=1309, 3-months; n=333, 2-
years) (**G**) and cristae score (n=555, 3-months; n=555, 2-years). (**H**) Mitochondrial
quantifications of female mitochondria number (n=21, 3-months; n=25, 2-years) (**I**)
mitochondrial area (n=1253, 3-months; n=1018, 2-years) (**J**) and cristae score (n=684, 3-months;
n=684, 2-years). (**K**) Schematic depicting removal of the liver. (**L**) Following embedded fixation,
Serial Block Face-Scanning Electron Microscopy (SBF-SEM) allows for ortho-slice alignment.
(**M**) Manual segmentation of ortho slices was performed to yield (**N**) 3-dimensional (3-D)
reconstructions of individually colored mitochondria. (**O**) Representative ortho slice images from
3-month murine liver tissue and (**P**) 2-year murine liver tissue. (**O'**) Representative ortho slice
images with 3D reconstructions of mitochondria overlaid from 3-month murine liver tissue and
(**P'**) 2-year murine liver tissue. (**O''**) Isolated 3D reconstructions of mitochondria from 3-month
murine liver tissue and (**P''**) 2-year murine liver tissue (**Q**) Mitochondrial quantifications
mitochondrial perimeter, (**R**) area, (**S**) and volume. For SBF-SEM, in total, 3-month samples
both included 750 (n=250, per mouse) for a total of 1500 mitochondria, which were used for
statistical analysis. For all panels, error bars indicate SEM, Mann–Whitney tests were used for
statistical analysis, and significance values indicate **$P \leq 0.01$, ***$P \leq 0.001$, and ns, not
significant.

1427
Figure 3: Serial Block Face-Scanning Electron Microscopy Shows Changes in
Mitochondrial Morphology in Liver Across Aging. (**A**) 3D reconstructions of mitochondria
displayed from the transverse viewpoint in 3-month and (**B**) 2-year murine liver tissue. (**A'**)
Representative images of 3D reconstructions of mitochondria displayed from the longitudinal
viewpoint in 3-month and (**B'**) 2-year murine liver tissue. (**C**) Based on these measurements, the
mitochondrial sphericity and (**D**) complexity index was determined. (**E**) Mito-otyping displays

diversity in mitochondrial morphology across each relative volume of mitochondria in 3-month
and (**F**) 2-year murine liver tissue. (**G-H''**) Qualitative identification of wrappER. (**G**)
Orthogonal (ortho) slice from 3-month and (**H**) 2-year samples, with (**G'-H'**) 3D reconstruction
of mitochondria and endoplasmic reticulum overlaid. (**G''-H''**) Isolated 3D reconstruction of
mitochondria, in purple, and endoplasmic reticulum, in blue. For SBF-SEM, in total, 3-month
samples both included 750 (n=250, per mouse) for a total of 1500 mitochondria, which were
used for statistical analysis. For all panels, error bars indicate SEM, Mann–Whitney tests were
used for statistical analysis, and significance value indicate ****$P \leq 0.0001$.

1442
1443

Figure 4: Global metabolomic and lipidomic profiling revealed metabolic dysregulation
and disruptions in lipid classes with age in liver tissues. (**A**) Metabolomics heatmap showing
the relative abundance of metabolites. (**B-M**) Metabolite pools illustrating the metabolic
pathways that are altered with age in the liver– Vitamin A Metabolism, TCA Cycle, Nucleotide
Metabolism, and NAD+ metabolism. (**N**) Heatmap showing enriched lipid classes based on
comparison between young and old liver tissues. (**O**) Volcano plot labeling significant hits,
which have adjusted p-value <0.05 and fold change (+ or -) greater than 1. For each tissue and
metabolite in the heatmaps, the aged samples were normalized to the median of the young
samples and then log2 transformed. Significantly different lipid classes represented in the figures
are those with adjusted p-values < 0.05 (note: p-values were adjusted to correct for multiple
comparisons using an FDR procedure) and log fold changes greater than 1 or less than −1.
Young, n= 4; aged, n= 4. For all panels, error bars indicate SEM, ** indicates p< 0.01; and *p<
0.05, calculated with Student's t-test.

1457

Figure 5. Aging Causes MICOS Loss, Diet Changes Affect SAM50 Expression, and SAM50
has Genetic Associations with Liver Diseases. (A-D) Quantitative PCR shows changes in
mRNA transcripts in (**A**) *Opa1*, (**B**) *Mitofilin/Mic60*, (**C**) *Chchd3/Mic19*, and (**D**) *Chchd6/Mic25*
between 3-month and 2-year murine liver samples. (**E**) Western blotting showing SAM50
protein levels, relative to tubulin, in 3-month and 2-year murine liver samples. (**F**) Western
blotting showing MIC60 and SAM50 protein levels, relative to ß-actin, in low-fat diet and high-
fat diet murine liver samples. (**G-G'**) Representative transmission electron micrographs from
low-fat diet and (**H-H'**) high-fat diet murine liver tissue. Quantifications of (**I**) mitochondria
number, as normalized to cell area (n=7, low fat diet; n=5, high fat diet), (**J**) individual
mitochondrial area (n=592, low fat diet; n=266, high fat diet), (**K**) the sum of all mitochondrial
area normalized to total cell area (n=7, low fat diet; n=5, high fat diet), (**L**) mitochondrial
circularity index, (n=598, low fat diet; n=271, high fat diet), and (**M**) cristae score (n=425, low
fat diet; n=425, high fat diet). (**N-N'**) Representative confocal fluorescence (using mCherry-
Mito-7) from wildtype and (**O-O'**) *Sam50*-knockout fibroblasts. (**P**) Quantifications of
mitochondrial length. (**Q**) Relative proportion of mitochondria that are either fragmented or
tubular in wildtype and *Sam50* KO fibroblasts, with the white area representing percentage of
fragmentation and colored areas representing percentage of typical tubular. (**R**) Phenome-wide
association study (PheWAS) results for SAMM50 GReX in individuals of European ancestry
(n=70,440). The GReX for SAMM50 was tested for association across 1,704 clinical phenotypes
extracted from the EHR. Association tests were run using logistic regression models, accounting
for genetic ancestry (principle components/PC 1-10), sex, age, median age of medical record,

1479 and genotyping batch. Associations that met the Bonferroni-corrected threshold (red line, p <
1480 2.934272 x 10^{-5}) are labeled with phenotype name [see SFile 1 for all PheWAS results]. The blue
1481 line represents nominal significance (p = 0.05) For all panels, error bars indicate SEM, Mann–
1482 Whitney tests were used for statistical analysis, and significance value indicate ****P ≤ 0.0001,
1483 ns   indicates   non-significant.
1484

1485 **Figure 6: Loss of MIC60 and CHCHD6 in HepG2 cells results in reduced mCa2+ uptake**
1486 **and   calcium   retention   capacity   and   oxidative   stress.**
1487 (**A**) Raw traces showing mitochondrial calcium uptake in permeabilized *MIC60* and *CHCHD6*
1488 knockdown HepG2 cells along with scr-siRNA transected controls. (**B**) Percentage change in
1489 mCa2+ uptake rate quantified from raw traces. (**C**) Recordings of mitochondrial calcium
1490 retention capacity in scr-siRNA, *MIC60* siRNA, and *CHCHD6* siRNA HepG2 cells. The circles
1491 indicate the number of calcium boluses taken up by specific cells. (**D**) Percentage change in
1492 mitochondrial calcium retention capacity quantified from recordings of mitochondrial calcium
1493 retention capacity. (**E**) Immunoblot confirming siRNA-mediated knockdown of *CHCHD6* in
1494 HepG2 cells. (**F**) Immunoblot confirming siRNA-mediated knockdown of *MIC60* in HepG2
1495 cells. (**G**) 4′,6-diamidino-2-phenylindole (DAPI) staining, MitoPY1 (5 µM, 45 min at 370 c
1496 magnification of 60x), and merge channels in scramble-siRNA (control), MIC60-siRNA, and
1497 CHCHD6-siRNA transfected permeabilized HepG2 cells. (**H**) 4′,6-diamidino-2-phenylindole
1498 (DAPI) staining, MitoBright Deep Red (10 µM, 30 min at 37^{0} c), DCFDA (10 µM, 30 min at 37^{0}
1499 c, magnification of 60x), and merge channels in scramble-siRNA (control), MIC60-siRNA, and
1500 CHCHD6-siRNA transfected permeabilized HEK293 cells. (**I**) Plate reader-based reactive
1501 oxygen species (ROS) quantification. (**J**) Microscopy-based ROS quantification of MitoPY1
1502 orange, (**K**) MitoSox Deep Red, and (**L**) DCFDA. For all statistical tests, one-way ANOVA
1503 statistical test was performed with Dunnett's multiple comparisons test. N=5-10 for all calcium
1504 experiments, each indicated by dots, as run in triplicates. N=9-13 for all oxidative stress
1505 experiments, each indicated by dots, as run in triplicates. Significance values indicate **P ≤ 0.01
1506 and ****P ≤ 0.0001.

Figure 1

Figure 2

Figure 3

3 Month

2 Year

Transverse

A   2μm

B

Longitudinal

A'

B'

C   Sphericity   ****
3 month   2 year

D   Complexity Index   ****
3 month   2 year

Mito-otype

E   0.5μm

F

3 Month

2 Year

G

H

G'

H'

G''

H''

Figure 4

Figure 5

Figure 6