Broad diversity of human gut bacteria with traceable strain identifiers to facilitate bulk
deposition

Thomas C. A. Hitch[1], Johannes M. Masson[1,#], Charlie Pauvert[1,#], Johanna Bosch[1],
Selina Nüchtern[1], Nicole Treichel[1], Marko Baloh[1], Soheila Razavi[1], Afrizal Afrizal[1],
Ntana Kousetzi[1], Andrea M. Aguirre[1], David Wylensek[1], Amy Coates[1], Susan A. V.
Jennings[1], Atscharah Panyot[1], Alina Viehof[1], Matthias A. Schmitz[1], Maximilian
Stuhrmann[1], Evelyn C. Deis[1], Kevin Bisdorf[1], Maria D. Chiotelli[2], Artur Lissin[3], Isabel
Schober[3], Julius Witte[3], Thorsten Cramer[4], Thomas Riedel[3,5], Marie Wende[6], Katrin
A. Winter[6], Lena Amend[6], Alessandra Riva[7,8], Stefanie Trinh[9], Laura Mitchell[10],
Jonathan Hartman[11], David Berry[7], Jochen Seitz[12], Lukas C. Bossert[11], Marianne
Grognot[2], Thorsten Allers[10], Till Strowig[5,6,13], Michael Pester[3,14], Birte Abt[3,5], Lorenz
C. Reimer[3], Jörg Overmann[3,5,14], Thomas Clavel[1*]

[1] Functional Microbiome Research Group, Institute of Medical Microbiology, University
Hospital of RWTH Aachen, Germany

[2] Biophysics of Host-Microbe Interactions Research Group, Institute of Medical Microbiology,
University Hospital of RWTH Aachen, Germany

[3] Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures,
Braunschweig, Germany

[4] Molecular Tumor Biology Research Group, Department of General, Visceral, Children and
Transplantation Surgery, University Hospital of RWTH Aachen, Germany

[5] German Centre for Infection Research (DZIF), Partner Site Hannover-Braunschweig,
Braunschweig, Germany

[6] Department of Microbial Immune Regulation, Helmholtz Centre for Infection Research,
Braunschweig, Germany

[7] University of Vienna, Center for Microbiology and Environmental Systems Science,
Department of Microbiology and Ecosystem Science, Vienna, Austria

[8] Chair of Nutrition and Immunology, School of Life Sciences, Technical University Munich,
Freising-Weihenstephan, Germany

[9] Institute of Neuroanatomy, University Hospital of RWTH Aachen, Germany

[10] School of Life Sciences, University of Nottingham, Nottingham, United Kingdom

[11] IT centre, RWTH Aachen, Germany

[12] Clinic for Child and Adolescent Psychiatry, Psychosomatic Medicine and Psychotherapy,
LVR-University Hospital Essen, University of Duisburg-Essen, Germany

[13] Centre for Individualised Infection Medicine (CiiM), a joint venture between the Helmholtz-
Centre for Infection Research (HZI) and the Hannover Medical School (MHH), Hannover,
Germany

[14] Technical University Braunschweig, Braunschweig, Germany

# Authors contributed equally to the manuscript

* Correspondence: tclavel@ukaachen.de

43 Abstract

Numerous bacteria in the human gut microbiome remain unknown and/or have yet to
be cultured. While collections of human gut bacteria have been published, few
strains have been made publicly available. A major hurdle in making strains publicly
available is their deposition to public culture collections. We propose a framework for
the bulk-deposition of strains to culture collections, which removes many of the
barriers previously identified (www.dsmz.de/bulk-deposit). Using this bulk-deposition
system we have created a publicly available collection of human gut isolates. The
Human intestinal Bacteria Collection (HiBC) (www.hibc.rwth-aachen.de) contains
340 strains representing 198 species within 29 families and 7 phyla, of which 29
previously unknown species are taxonomically described and named. These
included two butyrate-producing species of *Faecalibacterium* and new dominant
species associated with health and inflammatory bowel disease, *Ruminococcoides*
intestinale and *Blautia intestinihominis*, respectively. Plasmids were prolific within the
HiBC isolates, with almost half (46%) of strains containing plasmids, with a maximum
of six within a strain. This included a broadly occurring plasmid (pBAC) that exists in
three diverse forms across *Bacteroidales* species. Megaplasmids were identified
within two strains, the pMMCAT megaplasmid is globally present within multiple
Bacteroidales species. This collection of easily searchable and publicly available gut
bacterial isolates will facilitate functional studies of the gut microbiome.

64 Introduction

The cultivation of human gut bacteria has accelerated in the last years [1–4], providing
valuable information on the presence of novel taxa within gut microbiomes. Yet while
these published collections of human gut isolates note the novel taxa within their
collections [1,5], rarely is this novelty made known by describing and validly naming the
taxa [3]. Curating the taxonomic assignment of such collections is essential as often
outdated names are used, or recently described taxa are ignored, causing greater
confusion in the current taxonomic sphere [6–9]. Such curated taxonomy ensures
strains are correctly assigned, allowing strain-level diversity to be studied [10].
Variation between strains of the same species can lead to functional shifts that alter
the association with host dietary habits [11,12]. One-way strains can vary is due to the
presence of mobile genetic elements, which are known to affect the phenotype of the
isolates in which they occur [13]. The study of plasmids within the human gut has been
limited until recently [14,15].

Accessibility to collections of strains from the human gut is essential for the
mechanistic study of microbe-microbe and microbe-host interactions [16–18]. However,
accessibility remains problematic, with few strains being deposited in public culture

collections and even fewer of the proposed novel taxa being validated. For most of
the strains deposited, genomes are available, but their quality is not always high and
metadata such as the source of isolation, cultivation requirements, and full taxonomy
are often lacking. The small number of deposited strains limits future confirmatory or
comparative studies, while the lack of curated metadata reduces the value of the
genomic information. Only the professional acquisition of strains by public collections
can ensure that the high-quality standards required for future work are maintained
and that valuable strain-associated information is gathered and made freely
accessible to researchers and users worldwide. This is particularly relevant for
microbiota research to move beyond associations and towards a more mechanistic
understanding, as exemplified by the study of *Akkermansia muciniphila* [19,20].

The deposit of strains in public collections is a time-demanding and expensive
process, yet necessary to ensure access to pure strains. Once deposited, the
preservation of strains becomes a highly cost-effective means of maintaining the
value of microbial strains over long periods of time [21]. The complete acquisition of
single bacterial strains at culture collections, which includes purity and identity
confirmation, requires several months of time and for fastidious anaerobic strains
such as those isolated from the human intestinal microbiome, is currently
accomplished for only about 20 strains in parallel by one laboratory technician. This
has become a major bottleneck in the creation of publicly accessible collections
emanating from current microbiome research that generates ever increasing
numbers of novel isolates over comparatively short time intervals. A solution that
enables rapid deposition is needed to enhance the time available to process strains.

To provide functional insights into the human gut microbiome based on isolates and
facilitate public access to strains for the scientific community, we sought to create a
bacterial collection and implement unique strain identifiers to facilitate bulk
deposition at an international collection. In this way, all strains are easily traceable
and made publicly available more rapidly. Studying this collection of isolates
provided insight into the prevalence and diversity of plasmids within the human gut
and the presence of megaplasmids. We also highlight health-associated variation
within key genera, including novel species which require further *in vivo* study.

Results

A diverse range of key commensal species isolated from the human gut

The Human intestinal Bacteria Collection (HiBC) consists of 340 strains,
representing 198 species, isolated from human faecal samples (**Figure 1**). It
contains 29 families from across the seven dominant phyla in the human gut:
Bacillota (n = 173 isolates), *Bacteroidota* (n = 95), *Actinomycetota* (n = 46),
Pseudomonadota (n = 22), *Desulfobacterota* (n = 2), *Fusobacteriota* (n = 1), and
Verrucomicrobiota (n = 1). Of the 198 species, 29 are novel taxa that we have
described and named for validation according to both, the SeqCode [6] and the ICNP

[22]. High-quality genomes defined as >90% completeness (99.22 ± 1.10%), <5%
contamination (0.50 ± 0.77%), >10x genome coverage (380.08 ± 293.97), were
generated for all 340 strains. To improve access to the strains, their genomes,
information on their cultivation, and isolation source material, we have created the
HiBC web interface; www.hibc.rwth-aachen.de. The website provides access to the
complete collections of 16S rRNA gene sequences, genomes, plasmids, and
metadata.

Figure 1: Phylogenomic diversity of isolates within HiBC. Tree based on all 340 genomes,
generated using Phylophlan [23]. Phyla are indicated with colours. The *Bacillota* are split due to the
placement of *Fusobacteriota,* which separated strains assigned to 'Bacillota_A' by GTDB. The
potential need for splitting the phylum *Bacillota* is therefore independently supported by the
Phylophlan tree, and GTDB. Blue circles identify strains belonging to the 29 novel species that are
taxonomically described in this paper.

Description of dominant novel taxa within the human gut

There has been a renaissance in isolation of strains from the human gut in the last decade [24–32]. To place the HiBC isolates in the context of prior studies, we analysed previously published work in which phylogenetically diverse isolates were cultured. Across eight large-scale isolation studies and repositories, a total of 12,565 strains from the human gut were reported (**Figure 2a, Supplementary Table 1**). Of these, 8,019 (63.8%) isolates claim to be requestable in the original publications, but only 1,539 (12.2%) have been deposited to a culture collection to ensure long-term availability. However, this includes 1,063 isolates that have only been deposited to China General Microbiological Culture Collection Centre (CGMCC), which prevents the accessibility of risk group 2 organisms, which many of these are, to researchers outside China. This means only 476 bacterial isolates (3.8%) from the human gut are currently accessible to the community. While access to many of the isolates is limited, most isolates have been genomes sequenced (11,498, 91.5%) with most being high-quality (10,893, 86.7%). To understand the diversity captured by each study, we dereplicated the genomes to estimate the number of species cultured by each study (**Figure 2b**). A 33-fold reduction in diversity was found in some studies. Despite the value of capturing variability at the strain level, the limited number of samples used for isolation suggests redundancy within these collections. Therefore, while many studies claim to have a large collection of isolates, this is an overestimate of the true diversity captured.

Next, we assessed the ability of all isolated strains to capture the diversity present in the human gut microbiota (**Figure 2c**). This was achieved by mapping the isolates genomes against a MAG collection for which corresponding relative abundance values were available [33]. The entire landscape of bacteria isolated from the human gut, both from the literature and this study, captured 69.83 ± 18.62 % of an individual's microbiota, while the HiBC alone covered 56.51 ± 20.44 %. The 29 novel taxa described within this work enhanced this coverage by 3.75 ± 3.99 % on average, and represented >10% of the microbiota from 380 people, which peaked at >40% in two samples. Of the 29 novel taxa, 11 species were identified to have a mean relative abundance >0.15% within positive metagenomic samples (**Figure 2d**). The most dominant of these was *Ruminococcoides intestinale* (mean relative abundance = 2.84%; n = 3,023), followed by *R. intestinihominis* (1.19%; n = 285). Although only recently described [34], these results suggest that *Ruminococcoides* is a dominant but understudied genus within the human gut. A second genus of clear ecological importance to the gut is *Lachnospira*, in which we describe two dominant novel species. Interestingly, the two samples in which novel taxa accounted for >40% relative abundance were dominated by either *R. intestinale*, or the novel species *L. rogosae*, identified as a medium priority, HMP most wanted taxa (**Supplementary Results**). Further study of these species is needed to understand

their impact on host health and identify the mechanisms, which is now enabled by access to isolated strains.

To uncover associations of the novel taxa with human health conditions, we studied the ecological occurrence of each protein within *R. intestinale* CLA-AA-H216 (=DSM 117897) (**Figure 2e**), as the most abundant novel taxa, and *Blautia intestinihominis* CLA-AA-H95[T] (=DSM 111354) (**Figure 2f**), as member of an important genus associated with both health and disease conditions. The association of individual proteins with disease states were determined using InvestiGUT[35]. This involves studying the prevalence of each individual protein encoded by the strain's genome, or plasmid, across thousands of metagenomes, then statistically determining if they occur significantly more or less frequently in the gut metagenomes of people with a specific condition or healthy controls. Out of the 2,167 proteins encoded by *R. intestinale* CLA-AA-H216, comparison between Crohn's disease (CD) patients and healthy controls identified a total of 1,902 differentially prevalent proteins, with 1,883 being significantly more prevalent in healthy samples whilst 19 were more prevalent in CD samples. The same pattern was observed with Ulcerative colitis (UC) where a total of 1,715 significantly differentially prevalent proteins were identified, with 1,684 enriched in healthy samples, and 31 enriched in UC samples. Pathway analysis of the health-associated proteins identified multiple anti-inflammatory pathways, including the ABC transporter for spermidine (PotB/C/D), an anti-inflammatory polyamine [36], and biosynthesis of biotin, which has been shown to be immunomodulatory [37,38], ameliorating colitis [39].

The association of *Blautia* spp. with inflammatory bowel diseases is more complex, as some species within this genus have been shown to ameliorate colitis [40], leading to their inclusion within therapeutic products [41]. However, recent studies have suggested some *Blautia* spp. may exacerbate colitis [42]. Given the complex interaction of this genus with inflammatory bowel diseases, we studied the association of the novel species, *B. intestinihominis* CLA-AA-H95[T], with both CD and UC. Out of the 3,906 proteins, comparison between CD patients and healthy controls identified 523 differentially prevalent proteins, with 488 being significantly more prevalent in healthy samples whilst 35 were more prevalent in CD samples. In contrast, comparison between UC patients and healthy controls identified 1,466 proteins, with 16 being significantly more prevalent in healthy samples whilst 1,450 were more prevalent in UC samples. The functionality of the large number of proteins enriched within UC samples was studied further, identifying ABC transporters for branched-chain amino acids (LivF/G/H/K/M) [43], phosphate (PstA/B/C/S) [44], and adenosine (NupA/B/C, BmpA) [45], each of which has been linked to colitis.

These results highlight the importance of describing novel isolates and provide insights into the observed association of their proteins with health or disease.

Figure 2: Ecology of isolated human gut bacteria and their proteins. a. The number of strains and genomes produced by eight major isolation projects, along with HiBC, were compared. Strains were deemed requestable if it was stated as such in the original publication. They were deemed deposited if culture collection identifiers were included in the original paper and were confirmed to exist. Genomes were deemed high-quality if they were >90% complete, and <5% contaminated. The number of strains within each study are stated, while the percentage meeting each criterion are plotted. Red dots highlight datasets which have barriers to their accessibility, *i.e.*, data available upon request or access limited to specific countries. Strain collections: GMbC, Global Microbiome Conservancy [28]; BIO-ML, Broad Institute-OpenBiome Microbiome Library [27]; CGMR, Chinese Gut Microbial Reference [30]; CAMII, Culturomics by Automated Microbiome Imaging and Isolation [31]; hGMB, Human Gut Microbial BioBank [29]; HBC, Human Gastrointestinal Bacterial Collection [26]; HiBC, Human Intestinal Bacteria Collection (this study); IHU, collection of the Institut Hospitalier Universitaire Méditerranée Infection [32]; HMP, Human Microbiome Project at ATCC. **b.** Number of species per isolate collection, either via manual curation (HiBC) or dereplication of the available genomes (ANI values >95 % indicated identical species). **c.** The cumulative relative abundance of gut metagenomes across 4,624 individuals from Leviatan *et al* (2022) covered by all isolated bacteria across studies including the HiBC (Global isolates, dark blue), HiBC alone (green), or the subset represented by the 29 novel taxa described in this work (light blue), which had matches within 4,583 of the samples. **d.** Relative abundance of dominant (mean relative abundance >0.25%) novel taxa across 4,624 individuals, with the number of positive samples stated. Each strain represents a distinct novel species, described in detail in in the protologues at the end of the methods section. **e-f.** Genomic location of proteins significantly differentially prevalent between Crohn's disease (CD) samples and healthy controls (inner ring), or ulcerative colitis (UC) samples and healthy controls (outer ring). The delta-prevalence (prevalence in healthy donors − prevalence in corresponding patients) is shown in blue (more prevalent in healthy controls), red (UC), or mauve (CD). The species, strain, number of proteins predicted within the genome, and those significantly differentially between health conditions are shown within the circle. Only contigs >10 kp were plotted.

<u>Plasmid landscape in the cultured strains and identification of prevalent</u> <u>megaplasmids</u> Plasmids are present in the human gut and vary between geographically distinct populations [13]. However, most analyses have been culture-independent, preventing accurate taxonomic assignment [14,15,46]. Even studies on horizontal gene transfer using isolates have not reconstructed plasmids and accounted for their impact [47]. Analysis of bacterial isolates has uncovered many novel plasmids from non-human primates [48]. As many bacterial genome assembly workflows do not consider plasmids, we developed a genome pipeline that first searches for the presence of plasmids using Recycler [49] and plasmidSPAdes [50], assembles them, and then removes their reads from consideration during genome assembly (see Methods). This resulted in the reconstruction of plasmids from 46% of the strains (155 out of 340), with a total of 266 plasmids (**Figure 3a**). Plasmids were identified in all phyla except *Fusobacteriota*, for which the HiBC includes only a single isolate. Almost half of the plasmid-positive strains contained more than one plasmid (64 out of 155 strains), up to 6 plasmids in a strain of *Phocaeicola vulgatus* and two strains of the novel species *Enterobacter intestinihominis* (**Figure 3b**). Across the 266 plasmids identified in HiBC, 3,697 proteins were predicted, although only 639 (17.28 %) could be functionally annotated. Of these, nine were antibiotic resistance genes, four of which were copies of the *APH(2")-IIIa* aminoglycoside resistance gene found on plasmids of identical size (7,686 bp) in four strains of *Ruthenibacterium lactatiformans* (strains CLA-KB-H110, CLA-KB-H15, CLA-KB-H80, CLA-AA-H80).

Megaplasmids (>100 kp) have been described to occur in *Lactobacillaceae* [51] and *Bifidobacterium breve* [52], but their occurrence in a broader range of commensals from the human gut is unknown. We evaluated the length of the reconstructed plasmids and found the average length was 11.74 ± 17.25 kp (**Figure 3c**). This included the identification of two megaplasmids, one within a strain of the recently described species *Hominifimenecus microfluidus* (strain CLA-JM-H9, = DSM 114605; 125.6 kp) and the other in a strain of *Phocaeicola vulgatus* (strain CLA-AA-H253, = DSM 118718; 103.4 kp). Given the number and size of plasmids a single strain can contain, we assessed their impact on species assignments based on average nucleotide identity, but confirmed that plasmids had only a minor effect on the taxonomic assignment of a genome (**Supplementary Results**).

Figure 3: Plasmid repertoire of human gut bacterial isolates and their association with disease,
and phenotypes. a. Phylum level diversity of plasmid-positive isolates. **b.** Number of plasmids
reconstructed within each isolate. **c.** Length of reconstructed plasmids in log-10 scale; a bar
represents the number; a line represents the distribution. **d**. Network analysis of pBAC plasmid
sequence similarity to each other as determined by MobMess. **e**. Sequence alignment of a
representative from pBAC cluster 1, 2, and 3. Grey lines show regions of >99% similarity. **f**. The
plasmid repertoire purified from 7 strains predicted to contain either a single or multiple pBAC
plasmids, as well as additional plasmids. The arrows indicate bands with a size matching the

reconstructed pBAC plasmids. **g**. Proteins encoded on pBAC plasmids from different strains with
significantly different prevalence between Crohn's disease (CD) patients and healthy controls. Each
point represents a single differentially prevalent protein, coloured based on the pBAC cluster the
protein was encoded on (panel e). **h**. Plasmid map of pMMCAT_H253. The innermost rings represent
the association of proteins differentially prevalent between CD samples and healthy controls (first
ring), or ulcerative colitis (UC) samples and healthy controls (second ring) via InvestiGUT [35]. Proteins
enriched in CD are purple, those enriched in healthy samples are blue, and those enriched in UC are
red. The third ring represents the GC content relative to the average of the entire plasmid. Boxes are
used to represent genes identified on the plasmid in the forward (outer ring) and reverse (inner ring)
strand, with coloured boxes being assigned a COG category, while grey boxes represent COG-
unassigned proteins. The fimbriae loci proteins are indicated in dark grey. Enzymes with a single
restriction site are indicated on the outer ring in orange. **i.** Quantification of cell adhesion from a
pMMCAT-containing strain of *P. vulgatus* (CLA-AA-H253) and its closest relative strain (H-iso) without
the megaplasmid (see methods). Statistics: paired, one-tailed t-tests. **j.** Representative images of cell
adhesion of the two selected strains. The green scale bars represent 50 µm.

Sequence comparison with MobMess[53] identified 168 plasmid clusters, the two
largest of which were specific to the *Bacteroidales* (**Supplementary Figure 1a**). The
largest cluster shared similarities with 1,266 plasmids within the comprehensive
plasmid database, PLSDB [54], but lacked an assigned name, hence we named it
pBAC (plasmid *Bacteroidales*). The second largest plasmid cluster was identified as
the cryptic plasmid, pBI143, which was recently described to be associated with IBD
[14] and occurred in 14 isolates across eight species. Given that pBAC matches 1,266
plasmids within PLSDB, while pBI143 matched only 719, it may be the most prolific
plasmid in the human gut. We therefore investigated pBAC further, observing it has
been most frequently found in the USA but has also been detected in Denmark,
Japan, Ireland, and China (**Supplementary Figure 1b**). Investigation of the
sequence similarity network identified that pBAC occurred in both a shorter (pBAC-1;
~4,220 bp) and longer (pBAC-2; ~5,419 bp) form, with the two cluster being
connected by two 9,494 bp variants (pBAC-3) (**Figure 3d**). Comparison of
representative sequences from each of these three clusters uncovered that both,
pBAC-1 and pBAC-2 shared no similarity, but were identical to regions of pBAC-3
(**Figure 3e**). These results suggest that the different forms of pBAC share a common
origin, implying they can co-occur within the same strain. We identified CLA-KB-
H139 as a strain which was predicted to contain both pBAC-1 and pBAC-2 and
confirmed their existence within the isolate by extracting plasmids from a diverse
range of species predicted to contain different forms of pBAC (**Figure 3f**). This
analysis confirmed that pBAC occurs across the *Bacteroidales*. Given the prevalence
of pBAC within common denizens of the human gut, we aimed to study their
functional potential. All three pBAC clusters encode mobilisation machinery
(MbpA/B/C) [55]. Interestingly, both pBAC-1 and pBAC-2 encode different toxin-
antitoxin pairs, with that in pBAC-1 resembling YoeB and YefM [56], while the pBAC-2
encoded system shared similarity with RelE and Phd [57]. Given their differing protein
content, we hypothesised their association with host health may differ, hence studied
the prevalence of their encoded proteins across disease cohorts (**Figure 3g**). pBAC-

was observed to encode 2-3 proteins that were significantly more prevalent within CD patients, with the most enriched protein being a replication protein, followed by YefM-like and YoeB-like proteins. Conversely, pBAC-2 plasmids encoded 5-6 proteins within significantly lower prevalence in CD patients, with the most being the RelE-like and PhD-like proteins. Toxin-antitoxin systems within *Bacteroidota* plasmids have previously been reported, although pBAC shared no similarity with these described plasmids [58]. These results suggest the pBAC encoded toxin-antitoxin pairs are associated with human health, potentially by altering the fitness of their host strain.

In addition to the highly prevalent pBAC plasmid, we studied the megaplasmid present in *P. vulgatus* (CLA-AA-H253), a prevalent and dominant species in the human gut. The megaplasmid matched 26 plasmids within PLSDB, including multiple designated as pMMCAT (**Supplementary Figure 1a**), a recently proposed large plasmid from *Bacteriodales,* which has yet to be confirmed using sequencing-independent methods [59], but has been shown to impact its host ability to form biofilms [60]. As such, the *P. vulgatus* megaplasmid was designated pMMCAT_H253. While megaplasmids have previously been observed in sequencing data of human gut commensals, their existence is rarely confirmed. PacBio sequencing generated a complete genome for strain CLA-AA-H253, which confirmed the reconstruction of pMMCAT_H253. To provide sequencing-independent validation of pMMCAT_H253, we used pulsed field gel electrophoresis which showed a band of ~100 kb after treatment using XbaI, for which the plasmid had a single restriction site allowing for its linearisation (**Supplementary Figure 2a**). The bands' identity was confirmed using southern blot with primers designed based on the predicted pMMCAT_H253 sequence (**Supplementary Figure 2b**). Out of the 104 proteins encoded on pMMCAT_H253, only 6 could be assigned to a functional category, highlighting the need for further characterisation of these proteins (**Figure 3h**). A locus containing four proteins assigned to 'fimbrillin family proteins' was identified within a region of lower GC content compared to the plasmids average, which may suggest this region represents variable cargo and not the backbone. The presence of 'fimbrillin family proteins' on pMMCAT_H253 was further investigated as fimbriae can facilitate adhesion of cells [61], including to the host epithelium, to enhance colonisation [62]. We therefore studied the ability of CLA-AA-H253, the pMMCAT containing strain, and its closest related strain (H-iso, 98.9% ANI), which lacks pMMCAT, to adhere to plastic wells. CLA-AA-H253 adhered to the plastic significantly more than H-iso after only 4 hours, with 2-fold more bacteria adherent after 30 hours of growth (**Figure 3i**). The adherence of strain CLA-AA-H253 was observed in the wells, while H-iso was adhered in patches across the wells (**Figure 3j**). These results support the need for greater study of pMMCAT as a potential source for phenotype variation within genera known to contribute to host health [64,65].

Enhanced diversity of *Faecalibacterium* species

The genus *Faecalibacterium* currently (January 2025) contains seven validly published and correctly named species. HiBC contains representatives for five of these, and isolates representing two novel species. Initial 16S rRNA gene analysis assigned these novel species as *Faecalibacterium prausnitzii*, a common commensal of the gut microbiota that has been associated with human health conditions ranging from reducing inflammation [66] to improving cognitive function in Alzheimer's disease [67]. Many of these beneficial observations have been attributed to *F. prausnitzii's* ability to produce large amounts of the short-chain fatty acid butyrate [68].

Taxonomically, the genomic diversity of *F. prausnitzii* is proposed to be greater than suggested by 16S rRNA gene sequence analysis [69]. This has led the GTDB to split this species into 12 proposed species. HiBC contains representatives of the not yet described species defined by GTDB-Tk as *F. prausnitzii* A (strain CLA-AA-H175) and *F. prausnitzii* J (strain CLA-AA-H281). This was confirmed using ANI and can be observed in a phylogenomic tree (**Figure 4a**). We therefore propose to name these novel species *Faecalibacterium tardum* (CLA-AA-H175) and *Faecalibacterium intestinale* (CLA-AA-H281). Protologues for these, and the other 27 novel taxa described in this paper are provided at the end of the Methods section. Both novel species were observed in the majority of microbiota studied at similar relative abundances as *F. prausnitzii* (*F. prausnitzii* = 0.24 ± 0.24 %; *F. tardum* = 0.16 ± 0.20 %; *F. intestinale* = 0.14 ± 0.21 %) (**Figure 4b**).

Butyrate production by *Faecalibacterium* spp. is critical for host health, hence we studied variability between the strains in their ability to produce butyrate (**Figure 4; Supplementary Table 2**). Butyrate production by *Faecalibacterium* is achieved via the acetyl-CoA pathway, which is also the dominant pathway for butyrate production in the human gut (**Figure 4c**) [70,71]. While all *Faecalibacterium* strains within HiBC produced butyrate, the amounts varied greatly, with *F. tardum* (strain CLA-AA-H175) producing only 0.8 ± 0.16 mM, whereas strains of *F. longum* and *F. butyricigenerans* produced >10 mM (**Figure 4d**). Given that butyrate production is a conserved phenotype of this genus, we investigated if variation in the acetyl-CoA pathway loci was responsible for the observed variation between isolates (**Figure 4e**). While little variation was observed between most of the strains, *F. tardum* CLA-AA-H175 encoded two truncated copies of the butyl-CoA:acetate-CoA transferase gene. To understand the cause of these truncated genes, we studied their placement in relation to each other and identified they occurred on the same strand but in different frames, with an overlap by 55 bp. Interestingly, the second truncated gene is encoded with 'GTG' as an alternative start codon, which may lead to lower transcription and hence alter butyrate production further [72]. Structural modelling of these proteins using AlphaFold3 identified that the truncated proteins form a complex that share 100% (**Figure 4f**) and 90% (**Figure 4g**) identity with the full protein from *F. prausnitzii* CLA-AA-H222, respectively. Acetate utilisation and butyrate production have been linked to the growth of *Faecalibacterium* spp. [73], hence we considered if the ability of the isolate to grow under the testing conditions alters its metabolism,

and eventually the ability to produce butyrate. We therefore correlated the average
butyrate production of each strain against its average growth, measured by OD600
(**Figure 4h**). A strong significant correlation ($r = 0.76$; $p = 0.01$) was identified
between the growth of a strain and the amount of butyrate it produced. These results
suggest that strains of *Faecalibacterium* vary greatly in their ability to produce
butyrate, either as a result of genetic modification, as in the case of *F. tardum*, or due
to the strains ability to grow *in vitro*. This further implies that strains that grow at high
abundance *in vivo* are likely to produce the most butyrate.

Figure 4: Novel diversity within *Faecalibacterium* and strain-dependent butyrate production. a.
The two novel species of *Faecalibacterium* described within this paper placed within the current
landscape of *Faecalibacterium* spp. with a valid name, along with the type genomes for proposed
divisions of *F. prausnitzii*, as determined by GTDB. The phylogenomic tree was rooted on
Ruminococcus bromii ATCC 27255[T]. Novel species are in bold, and the type strain of *F. prausnitzii* is
underlined. **b.** Relative abundance, and prevalence, of the genus *Faecalibacterium*, and each
Faecalibacterium species represented within HiBC across 4,624 metagenomic samples. **c.** Butyrate
production pathway in *Faecalibacterium* with gene names and KEGG ortholog identifiers when
possible. **d.** Phylogenomic tree of the *Faecalibacterium* strains within HiBC, displaying the ability of
each strain to produce butyrate over a 48 h-period, along with the OD600 that the strain achieved
during the testing period (n = 3 independent batch cultures for each strain; the replicates are shown
with individual boxes). The phylogenomic tree was rooted on *R. bromii* ATCC 27255[T]. **e.** Sequence
comparison of the butyrate production loci across the *Faecalibacterium* strains. Genes are coloured
based on their assignment to each step in the butyrate production pathway in panel c. **f.** AlphaFold3
model of the *But* complex in *F. tardum* CLA-AA-H175 against the full *But* protein in *F. prausnitzii* CLA-
AA-H222. The first CLA-AA-H175 *But* gene is highlighted in yellow in the dashed box, while the
second gene is shown in faded orange. The highlighted protein is indicated in the top right of the
dashed box. **g.** Same as in panel f, but this time the second CLA-AA-H175 *But* gene is highlighted in

446 yellow in the dashed box, while the first gene in faded orange. The highlighted protein is indicated in
447 the top right of the dashed box. **h.** Correlation of the mean OD600 against the mean butyrate
448 production of each strain with a linear regression and its 95% confidence interval.

Traceable strain identifiers facilitate bulk isolates deposition

Published isolates are often not easily accessible to researchers, impeding them
from working with the same quality-controlled strains that are needed to compare
and reproduce data. The main barriers to making isolates publicly available is the
time-demanding process of strain deposition and the lack of unique strain identifiers
for reliable referencing of strains while they are being processed. Researchers
frequently deposit only a small selection of strains (**Figure 2a**), preventing access to
an entire collection that may contain functionally relevant diversity, as highlighted
above with *Faecalibacterium* (**Figure 4**). Current strategies for publishing large
collections of isolates vary, as shown in **Figure 5** (upper box). Publication with a
statement such as "available upon request" without deposition of the strains (case 1)
appears simple and quick. Yet, it should not be accepted anymore, as the strains
and their data are effectively inaccessible and ultimately will be lost over time [74].
Depositing in a culture collection (DSMZ, JCM, CCUG, BCCM, KCCM, etc.) ensures
that metadata quality is maintained and guarantees the long-term accessibility of the
strains for the scientific community. Current best practice is to ensure that all strains
included in a manuscript are deposited and processed at a culture collection before
publication, so that strain catalogue numbers can be issued and included in the
manuscript (case 4) [75,76]. The shortcoming of the latter approach is that deposition
and strain processing for quality checks is time consuming, and the time required
grows linearly with the number of strains to be deposited. The deposition of a
collection such as HiBC is thus a significant undertaking.

Accordingly, a bulk submission system (www.dsmz.de/bulk-deposit) has been
developed by the Leibniz Institute-German Collection of Microorganisms and Cell
Cultures (DSMZ) to address these issues and facilitate large-scale strain deposition,
as shown in **Figure 5** (lower box) (**Supplementary Results; Supplementary Figure
9**). The HiBC presented in this article has been used as a blueprint to develop this
system. After the culture collection has accepted the submission of a researchers
strain collection, it requests the strains in large batches which are deposited in an
iterative process. Once cultures have been received by the DSMZ and passed initial
checks, the system allows for the allocation of stable identifiers in the form of Digital
Object Identifiers (DOIs) provided by the database StrainInfo
(www.straininfo.dsmz.de), where each strain is permanently registered with its
metadata. This step ensures that all data (e.g., publications and sequence data) can
be linked to the corresponding strain entry when the strains become available. Strain
entries are initially displayed with the status 'Deposition in progress'. When the
deposition process is completed (which may take several months) and the strains
become available in the DSMZ catalogue, the status of the strains are updated to
'Published' in StrainInfo. Therefore, this system would accommodate fast publication

without compromising metadata quality, ensuring adherence to good scientific
practices.

Figure 5: Schematic of the proposed system for large-scale strain deposition. Once bacteria
have been isolated and described, there are multiple strategies for publishing the strains. In the upper
box, we have four example strategies, each of which can be applied on a single strain basis, but are
not designed for large-scale submission of strains. The benefits and negatives of each method is
indicated on the right of the methods under four main categories, where a cross indicates a missing
aspect to the strategy. The lower box pictures the main advantage of creating unique strain identifiers
early in the process as a foundation for bulk deposition of many strains (www.dsmz.de/bulk-deposit).
Figure designed in BioRender.

Discussion

Over the last decade, the renewed interest in cultivation has expanded the number
of genomes available for taxa within the human gut. However, the corresponding
strains are rarely accessible to the research community, as highlighted by only 3.8%
of human gut isolates being publicly deposited within a culture collection. The
process of strain deposition within culture collections has previously been highlighted
as a major limiting factor in making strains publicly available [77]. To overcome this, we
initiated a system for bulk deposition of strains. Streamlined registration of strains
before bulk submission ensures their traceability, facilitating publication while
ensuring strains are consistently referenced. Application of this system allowed for
deposition of 340 human gut isolates in the DSMZ, facilitating the naming and
description of 29 novel species, including the type species of three novel genera.
Based on this, the HiBC is the first collection of human gut isolates that is entirely
publicly accessible.

While the new bulk deposition process described here currently relies on manual
steps and separate communication between multiple partners, DSMZ is currently
developing a deposition management system that will largely automate this process.
Similar to the review systems of scientific journals, it will expedite the deposition
process by facilitating the submission of deposition data and the communication
between submitter and curator in a web portal named StrainRegistry
(https://straininfo.dsmz.de/strainregistry).   The   StrainRegistry   will   be   closely
connected to the StrainInfo database described in this manuscript for automated DOI
assignment to identify strains throughout literature and databases. We will also
encourage participation of other culture collections, to allow researchers to submit
their strains to multiple collections at the same time. This will significantly streamline
the deposition process and make it more transparent, while providing the persistent
identifiers necessary to connect strain data at any time.

Large-scale cultivation of human gut bacteria in this study led to the isolation of
many novel species and genera. Whilst previous cultivation studies report many
isolates representing novel diversity, these bacteria are rarely taxonomically
described and named. The discovery of prevalent and important novel species in this
work will facilitate future experiments on the role of bacteria in health and disease.
This included *Ruminococcoides intestinale*, *Blautia intestinihominis*, and the isolation
of multiple strains of *Faecalibacterium*, including species with the potential to reduce
gastrointestinal inflammation [66]. We confirmed that *F. prausnitzii* represents a
diverse group of related species that show reduced 16S rRNA gene sequence
diversity [69]. By naming two of these species and describing the butyrate production
and growth characteristics of seven different species in this genus, we observed the
ability of these strains to produce butyrate was proportional to their ability to grow.
Given these results, further quantification of these strains ability to grow within the
human gut is required to understand their contribution to SCFA levels *in vivo*, and
therefore their potential impact on host health.

The reconstruction of plasmids for almost half of HiBC isolates allows the direct link
of plasmids to strains of various species, information previously lacking from studies
on plasmids from the human gut [13–15]. Of note, *Bacteroidales* strains often contain
multiple plasmids, particularly copies of pBI143 or pBAC. The pBAC plasmid was the
most frequently reconstructed plasmid, and was observed to occur in three forms.
Investigation of these forms uncovered both major forms have differing associations
with host health. The megaplasmid pMMCAT was also identified within a strain of
P. vulgatus, leading to enhanced adhesion to surfaces compared to strains lacking
pMMCAT. By making the reconstructed plasmids, and strains publicly available, we
believe this resource expands the toolbox of methods for genetically modifying non-
model commensal gut microbes.

Metagenomic studies have provided important insights into the taxonomic and
functional diversity present in the human gut, however, they are unable to provide
mechanistic insights. By enabling broad access to human gut isolates in this work,

the functionality of taxa can be experimentally validated, facilitating mechanistic
studies of microbe-host interactions.

Methods

<u>Ethics</u>

The Ethics Committee of the Medical Faculty of RWTH University Aachen permitted
bacterial isolation from human stool under ethical number EK 23-055, EK 316-16,
and EK 194/19. For strains originating from Vienna, isolation was approved by the
University of Vienna ethics committee under ethical number 00161. For strains
originating from Braunschweig, isolation was approved by the Ethics Committee of
Lower Saxony (MHH permit No. 6794, 8629, and 8750). Written informed consent
was signed by all enrolled participants.

<u>Bacterial isolation and cultivation</u>

Human stool was collected in sterile plastic bags and stored in tightly sealed plastic
buckets until further processing. An oxygen scavenger sachet (BD Biosciences; ref.
260683) was added to each bucket before sealing to reduce exposure to oxygen. All
samples were processed in the lab within 24 hours of collection. First the faecal
material was homogenised by manual kneading of the plastic bag. Stool samples
were either mixed 1:5 with anaerobic FMT media [78], distributed to 1 mL aliquots and
stored at -80 °C until further use, or the samples were further processed as
described previously [76]. Hence approximately 5 g of the samples were dissolved in
50 ml of anaerobic PBS supplemented with peptone (0.05% w/v), L-cysteine (0.05%
w/v) and dithiothreitol (DTT) (0.02% w/v) by shaking the glass flask intensely. A
syringe with needle was used to transfer 5 ml of the slurry through a rubber stopper
(previously flamed using ethanol) into a second glass flask containing 45 ml of the
same buffer in an anaerobic atmosphere (89.3% N_2, 6% CO_2, 4.7% H_2) to create a
1:100 dilution of the original sample. The flask was moved into an anaerobic
workstation (MBraun GmbH, Germany) to prepare 2 ml aliquots under anaerobic
conditions, which were mixed with 2 ml of 40% anaerobic glycerol to a final
concentration of 20%, then stored at -80 °C until use.

Two different approaches were used for bacterial isolation. For classical isolation,
the faecal aliquots were thawed inside the anaerobic workstation and diluted with
anoxic PBS (see above) in a tenfold dilution series (10^{-2} to 10^{-6}). 50 µl of each
dilution step were transferred onto different agar plates and spread using a L-
spatula. Single colonies were picked after 1-7 days of incubation at 37 °C under
anaerobic conditions. Bacterial cells were re-streaked at least three times to
guarantee purity of the culture. For the second isolation approach, the single-cell
dispenser b.sight (Cytena GmbH, Germany) was used as described before [79].
Details on culture media preparation and ingredients are provided in the
Supplementary Methods.

The isolated bacteria were first identified using MALDI-TOF MS (Bruker Daltonics,
Bremen, Germany). If a species was not yet present within HiBC, or could not be
reliably identified by MALDI-TOF MS, the bacteria were stored as cryo-stocks and
DNA extracted for genome sequencing. Cryo-stocks were generated by freezing in
glycerol media (end concentration 20%) at -80 °C. All HiBC strains have been
deposited at the Leibniz Institute-German Collection of Microorganisms and Cell
Cultures (DSMZ). In addition, the strains representing novel taxa were deposited at
either the Belgian Coordinated Collections of Microorganisms (BCCM) or the Japan
Collection of Microorganisms (JCM).

Metabolite production analysis

Concentrations of short-chain fatty acids (SCFAs) (acetate, butyrate, propionate,
valerate), branched chain fatty acids (isobutyrate, isovalerate), intermediate
metabolites (ethanol, formate, lactate, 1,2-propandiol, 1-propanol, succinate), as well
as mono- and disaccharides (galactose, glucose, lactose) were measured by high-
performance liquid chromatography (HPLC). The bacterial strains were grown
anaerobically, apart from *Robertmurraya yapensis* CLA-AA-H227 which was grown
under aerobic conditions, in YCFA broth (DSMZ Medium No. 1611) in Hungate tubes
for 48 hours at 37 °C. Triplicate cultures from each strain were measured. Baseline
controls were taken from each sterile Hungate tube before inoculation. Each taken
sample was centrifuged (10,000 x g, 10 min, 4 °C), supernatants were collected and
stored at -80 °C until HPLC measurement. Samples were prepared and measured
(including HPLC settings) as described previously [76]. External standards were used
for concentration determination by comparison of peak retention times (HPLC grade
compounds were purchased from Sigma-Aldrich). Peaks were integrated using the
Chromaster System Manager Software (Version 2.0, Hitachi High-Tech Science
Corporation 2013, 2017). Metabolite concentrations >0.2 mM (limit of detection
(LOD) for citrate, 1-propanol), >0.24 mM (LOD for butyrate, formate, galactose,
glucose, isobutyrate, isovalerate, lactose, valerate), >0.4 mM (ethanol, 1,2-
propandiol) and >0.8 mM (acetate, lactate, propionate, succinate) were considered
for statistical analysis if present in all three replicates. Production and consumption
of metabolites was calculated by subtracting the baseline values from the sample
taken after 48 hours of growth.

Cellular fatty acids (CFAs) determination

Cellular fatty acids were measured at the Leibniz Institute-DSMZ. The strains
representing novel taxa were grown under the conditions indicated in the respective
protologues. Approximately 100 mg (wet weight) of cell biomass was extracted
according to the standard protocol of the Microbial Identification System (MIDI Inc.,
version 6.1; technical note #101). CFAs were analysed by converting them into fatty
acid methyl esters (FAMEs) through saponification, methylation, and extraction. The

resulting FAME mixtures were separated using gas chromatography (GC) and
detected with a flame ionization detector (FID). Subsequent analysis involved the
identification of fatty acids using a GC-MS system (Agilent GC-MS 7000D) as
described by Vieira et al. (2021)[80]. Further derivatization methods were used for
structural elucidation of unidentified compounds. For branched-chain fatty acids,
cyclo-positions, and multiple double bonds, 4,4-dimethyloxazoline (DMOX)
derivatives were analysed [81].

Isolation of genomic DNA

DNA was isolated using a modified protocol according to Godon *et al.* [82]. Frozen cell
pellets were mixed with 600 µl stool DNA stabilizer (Stratec biomedical), thawed, and
transferred into autoclaved 2-ml screw-cap tubes containing 500 mg 0.1 mm-
diameter silica/zirconia beads. Next, 250 µL 4 M guanidine thiocyanate in 0.1 M Tris
(pH 7.5) and 500 µL 5 % N-lauroyl sarcosine in 0.1 M PBS (pH 8.0) were added.
Samples were incubated at 70 °C and 700 rpm for 60 min. A FastPrep® instrument
(MP Biomedicals) fitted with a 24 × 2 ml cooling adaptor filled with dry ice was used
for cell disruption. The program was run 3 times for 40 s at 6.5 M/s. After each run,
the cooling adapter was refilled with dry ice. An amount of 15 mg
Polyvinylpyrrolidone (PVPP) was added and samples were vortexed, followed by 3
min centrifugation at 15.000 x g and 4 °C. Approximately 650 µl of the supernatant
were transferred into a new 2 ml tube, which was centrifuged again for 3 min at
15.000 x g and 4 °C. Subsequently, 500 µl of the supernatant was transferred into a
new 2 ml tube and 50 µg of RNase was added. After 20 minutes at 37 °C and 700
rpm, gDNA was isolated using the NucleoSpin® gDNA Clean-up Kit from Macherey-
Nagel. Isolation was performed according to the manufacturer's protocol. DNA was
eluted from columns twice using 40 µl Elution buffer and concentration was
measured with NanoDrop® (Thermo Scientific). Samples were stored at -20 °C.

Genome library preparation and Illumina Sequencing

Library preparation and sequencing were conducted using the NEBNext Ultra II FS
DNA Library Prep Kit for Illumina with dual index primers and ~300 ng of DNA (NEB
Inc.) on an automation platform (Biomeki5, Beckman Coulter) according to the
manufacturer's instructions. The time used for enzymatic shearing to ca. 500 bp to
be used on a MiSeq run was 10 minutes and to ca. 200 bp to be used on a NextSeq
run was 30 minutes. PCR enrichment of adaptor-ligated DNA was conducted with
five cycles using NEBNext Multiplex Oligos for Illumina (NEB, USA) for paired-end
barcoding. AMPure beads (Beckman Coulter, USA) were used for size selection and
clean-up of adaptor-ligated DNA. Sequencing was performed either at the IZKF Core
Facility Genomics (Uniklinik RWTH Aachen) on a NextSeq platform (Illumina) (PE
150 bp), or on a MiSeq (Illumina) (PE 300 bp) in house.

Sequencing data processing

Genomes were assembled from paired-end Illumina short-reads. Low-quality reads with an expected average quality below 20 over a 5-bases window, containing adapters, and shorter than 50 bases were discarded using Trimmomatic (v0.39) [83]. Reads with phiX sequences were removed using BBduk from the BBtools suite [84]. Plasmids sequences were reported only when inferred from plasmidSPades (v3.15.5) [85] and then from Recycler (v0.7) [49]. Reads free from plasmid sequences were then assembled using SPades (v3.15.5) [50] with default parameters.

Contigs above 500 bp in assemblies were kept for quality evaluation. Following the MIxS specifications [86] as well as requirements from the SeqCode [6], genomes with a coverage above 10X, a completion above 90%, and a contamination below 5% based on CheckM (v1.2.2) [87] estimation using single-copy marker genes were flagged as high-quality draft genomes. Additional quality flags on the assembly included: 1) <100 contigs, longest contig >100 kp, and N50 >25 kp using QUAST (v5.0.2) [88]; 2) detectable 16S and 23S rRNA genes using metaxa2 (v2.2.3) [89]; 3) >18 unique essential tRNAs genes and a detectable 5S rRNA gene sequence using the annotations of bakta (v1.6.1) [90].

The genome assembly pipeline, including the reconstruction of plasmids and quality checks steps, is available as a reproducible workflow using Snakemake (v7.9.0) [91]: www.github.com/clavellab/genome-assembly.

Genome analysis

Taxonomic, functional, and ecological analysis for all genomes was conducted using Protologger (v1.3) [92]. For this, 16S rRNA gene sequences were extracted from each strain's genome using barrnap v0.9 (default settings) (www.github.com/tseemann/barrnap), and the longest 16S rRNA gene sequence from each genome was used as input for Protologger. Phylogenomic trees were generated using Phylophlan v3.0.60 [93] with proteomes predicted using Prodigal v2.6.3 (default settings) [94]. Gut metabolic modules presence was based on the identification of all included KOs within a genome, detected by Kofamscan v1.3.0 [95].

Comparison to published isolate collections

Metadata for previously published collection of microbial isolates from the human gut were obtained from screening literature. To prevent redundancy, we have excluded articles that are a subset of large collections [25], and those that lack strain metadata or genome access [96]. In the case of isolate collections from multiple body sites, the isolation source was checked to be related to the gastrointestinal tract. Of the 12,565 strains identified, 11,498 had genomes and hence could be analysed. Of these, 10,893 had a high-quality genome and were studied. High-quality was determined by completion above 90%, and less than 5% contamination based on CheckM (v1.2.2).

720 Strains were determined to be 'publicly available' if a genome is available and the
721 strain has been deposited within a culture collection, *i.e.*, DSMZ, CGMCC etc. The
722 validity of each published culture accession number was checked to ensure strain
723 availability. Genomes within the collections were dereplicated at the species level
724 based on >95% ANI values via FastANI [101].

Ecological analysis

The occurrence of each strain across 16S rRNA gene amplicon samples was
conducted using Protologger [97]. In essence, 1,000 16S rRNA amplicon samples for
each body site (gut, vagina, skin, lung) were processed by IMNGS [98]. Next, their
operational taxonomic unit (OTU) sequences were compared against the 16S rRNA
gene sequences of the HiBC strains using BLASTN (>97% identity, >80% coverage).
Comparison of HiBC genomes against metagenome assembled genomes (MAGs)
was also conducted using Protologger and its collection of 42,927 MAGs with
curated metadata. The 16S rRNA gene sequences of the Human Microbiome
Projects "most wanted" taxa were obtained from Fodor *et al* (2012) and their priority
taken from the original publication [99]. The most wanted sequences were compared
to the 16S rRNA sequences of the HiBC strains using BLASTN (>97% identity,
>80% coverage) [100]. The prevalence and relative abundance of the HiBC strains
within shotgun metagenomic samples was determined by comparison of the HiBC
strain genomes to the representative genomes from Leviatan *et al* (2022) via
FastANI (>95% ANI) [33,101]. HiBC strains matching to representative genomes were
then connected to the pre-calculated relative abundance of the representative
genomes across 4,623 individuals, published in Leviatan *et al* (2022).

Plasmid analysis

Sequence similarity between the isolates plasmids was determined using MobMess
[15] and clusters were visualised with Cytoscape [102]. Clustering was conducted firstly
based only on the isolate plasmids, and then including the background previously
predicted human gut plasmids. The similarity of each plasmid to known sequences
was determined with the 'mash dist' option of PLSDB (v2023_11_03_v2) [54]. The
ecology of each plasmid was inferred from the geographical location assigned to
each matching plasmid. The sequences of pBAC plasmids were rotating using
Rotate, and then aligned with Easyfig [103]. Only regions of similarity with an identity
>90% were studied.

Pulsed Field Gel Electrophoresis

Cultures were grown overnight in 5 mL BHI media, centrifuged (2 mL for *P. vulgatus*
CLA-AA-H253, 5 mL for *H. microfluidus* CLA-JM-H9), resuspended in 100 µL sterile
PBS and incubated at 45 °C for 10 minutes. Of this, 80 µL was mixed with 320 µL

1% agarose (Bio-Rad Certified Megabase Agarose) at 45 °C, 80 µL transferred to
PFGE plug moulds (Bio-Rad CHEF Disposable Plug Molds, 50-Well) and cooled at 4
°C for 30 minutes. Solidified agarose plugs were transferred to lysis solution inside 2
mL microcentrifuge tubes (20 mM Tris-HCl pH 8.8, 500 mM EDTA pH 8, 1% N-
laurylsarcosine, 1 mg/mL Proteinase K) and incubated with shaking at 250 rpm at 45
°C for 4 h. Lysis solution was replaced with fresh lysis solution + RNAse (10 µg/mL)
and incubated with shaking at 45 °C overnight. Lysis solution was replaced with
wash solution (25 mM Tris-HCl pH 7.5, 100 mM EDTA pH 8) and incubated with
shaking at 45 °C. Plugs were then transferred to fresh 2 mL microcentrifuge tubes
with wash solution containing phenylmethanesulfonyl fluoride (1 mM) and incubated
with shaking at 45 °C for 30 min, twice. Plugs were then transferred to 1 mL wash
solution at stored at 4 °C until use in PFGE. For analysis of intact genomic DNA,
agarose plugs were subjected to 100 Gy of γ radiation using a ^{137}Cs source
(Gammacell 1000), to linearize circular chromosomes [104]. PFGE was performed
using a CHEF Mapper apparatus (Bio-Rad). Intact and XbaI-digested DNA
fragments were separated on a 1.2% agarose gel in 0.5 × TBE at 14 °C, with a
gradient voltage of 6 V/cm, linear ramping, an included angle of 120°, initial and
final switch times of 0.64 s and 1 min 13.22 s, respectively, and a run time of 20 h
46 min.
779
<u>Southern blot</u>
The DNA probe was designed by identifying a 1,428 bp DNA sequence unique to the
pMMCAT_H258 plasmid sequence. Primers sequences were checked against the
P. vulgatus genome sequence using SnapGene (www.snapgene.com) to confirm
specificity to only plasmid DNA. PCR primers were then designed based on the
regions flanking the unique region. *P. vulgatus* total genomic and plasmid DNA was
extracted using the Mericon DNA Bacteria Plus Kit following the manufacturer's
instructions (Qiagen). This DNA was used as a template to amplify the target DNA
sequence using ThermoFisher Scientific Phusion Hot Start 2 DNA polymerase and
the above primer set. The amplified DNA was visualized by gel electrophoresis to
confirm the expected DNA size and the DNA extracted and purified using
ThermoFisher Scientific GeneJET gel extraction kit.
792
The Southern blot gel was stained with 0.5 µg/ml ethidium bromide and visualised,
then acid-nicked in 0.25 M HCl, and subsequently denatured in 1.5 M NaCl, 0.5 M
NaOH. The DNA was transferred onto a GE healthcare Amersham Hybond XL
membrane by vacuum transfer using a Vacugene XL gel blotter (Pharmacia Biotech)
for 1 hour at 40 mBar. The membrane was briefly neutralised in 2x SSPE (20x
SSPE: 3 M NaCl, 230 mM NaH2PO4, 32 mM EDTA, pH 7.4) and the DNA then
crosslinked with 120 mJ/cm2 UV. The membrane was pre-hybridised for 4 h at 65°C
in 6x SSPE, 1% SDS, 5x Denhardt's solution (100x Denhardt's solution: 2% Ficoll
400, 2 % polyvinyl pyrrolidone 360, 2% bovine serum albumin), 200 µg/ml salmon
sperm DNA (Roche, boiled). The DNA probe used 50 ng of DNA template and was
radiolabelled using 0.74 MBq of [α-32P] dCTP (Perkin Elmer) and HiPrime random

priming mix (Roche), then purified on a BioRad P-30 column. The membrane was
hybridised with the radiolabelled DNA probe overnight at 65 °C in 6x SSPE, 1%
SDS, 5% dextran sulphate, 200 µg/ml salmon sperm DNA (Roche, boiled). The
membrane was then washed twice for 30 min at 65 °C in 2x SSPE, 0.5 % SDS, and
twice for 30 min at 65 °C in 0.2x SSPE, 0.5% SDS, before being exposed to a
phosphorimager screen (Fujifilm) for 24 h and then scanned on a GE Healthcare
Typhoon.

Plasmid extraction and visualisation

Frozen cryostocks were plated on mGAM plates and incubated anaerobically at 37
°C for 48 h. A single colony was inoculated into 5 mL of BHI media within a Hungate
tube and incubated for 48 h. 2 mL of culture was transferred into 2 mL
microcentrifuge tubes and centrifuged at 18,000 rcf for 5 min. The medium was
removed and the pellet processed using QIAprep Spin Miniprep kit, following the
manufacturer's recommendations (Qiagen, Cat. No. 27106). Depending on DNA
concentration, 10-20 µL of extracted DNA was run on a 0.5 % agarose gel (Sigma-
Aldrich A9539) at 80 V for 120 min (Bio-Rad Biometra) along with GeneRuler 1 kb
DNA ladder (ThemoFisherScientific SM0312) and stained with a dye (Midori Green
Advance, BulldogBio). The gel was imaged with Bio-Rad GelDoc Go imaging
system.

Crystal violet staining to quantify strain adhesion

Strains of interest were tested in triplicates. For each, one colony was picked and
grown to saturation overnight in BHI. Cultures were diluted back to an OD600nm of
0.1 in fresh BHI, 3 ml were placed per well in 6-well plates (Nunc cell-culture,
treated, flat clear polystyrene plates, Thermo Scientific) and the plates were
incubated in static conditions at 37 °C. For each timepoint, a plate containing the two
strains to be compared were removed from the anaerobic cultivation chamber, rinsed
with water, and dried at 60 °C for 2 h. Each well was incubated with 1 % (w/v) crystal
violet solution then rinsed 3 times with deionised water and air dried. Once all plates
were collected, wells were destained by adding 1 ml of a 1:4 acetone:ethanol
solution. After gentle mixing, at least three technical replicates of 200 µL per well
were placed in wells of a 96-well plate. The optical density of crystal violet staining
present in the destaining solution was measured at 590 nm. Wells filled with 200 µL
of pure destaining solution were used as blank reference. All wells from the same
biological replicate at a given time point were averaged to provide a single point and
used for statistical analysis. Cell counting in Neubauer chambers ensured an initial
cell concentration varying by less than 5 %, and OD600nm measurements in the
supernatant at all timepoints of incubation showed similar OD values over all
biological replicates, confirming that growth rates were not driving the observed 2-
fold difference in attached biomass after 30 h. Representative images of attachment

were obtained after 24 h following the above protocol until the first drying stage. Two
wells were imaged in their centre using phase contrast (Nikon Ti-E inverted
microscope, 40x objective lens).

<u>Protein modelling</u>

Protein sequences were entered into the AlphaFoldServer [105]. Pairwise structure
alignment was conducted using the RCSB PDB webserver with the TM-align method
[106,107].

<u>Website design</u>

Access to taxonomic, cultivation, and genomic information about the HiBC strains is
available at: www.hibc.rwth-aachen.de. This website was created using the Shiny
package from R and the code is available at: https://git.rwth-
aachen.de/clavellab/hibc. The 16S rRNA gene sequence and genome for each strain
can be downloaded directly from this site, both individually for strains of interest via
the research data management platform Coscine, or for the entire collection via the
digital repository Zenodo. Further strain specific metadata provided includes:
isolation conditions, source, and risk group.

<u>Description of novel taxa</u>

The description of novel taxa was based on the analysis provided by Protologger
v1.3 [92], and manually curated into the protologues below. For each isolate, taxonomy
was assigned using the following thresholds: <98.7% 16S rRNA gene sequence
identity (as indication for as-yet undescribed species), <94.5% (undescribed genus),
and <86.5% (undescribed family) [108]. ANI values <95% to separate species [109];
POCP values <50% for distinct genera [110]. Phylogenomic trees were also considered
to make decisions on genus- and family-level delineation [23]. All novel taxa have been
registered with the SeqCode and will be registered with the ICNP.

Description of *Alistipes intestinihominis* sp. nov.

Alistipes intestinihominis (in.tes.ti.ni.ho'mi.nis. L. neut. n. *intestinum*, the intestine; L.
masc. n. *homo*, a human being; N.L. gen. n. *intestinihominis*, of the human intestine).

The genome size is 3.85 Mbp, G+C percentage is 58.29%, with 99.76%
completeness and 0.96% contamination. Strain CLA-KB-H122 was determined to be
a new species based on 16S rRNA gene analysis, with the closest validly named
match being *Alistipes timonensis* (98.17%). Separation from existing *Alistipes*
species was confirmed by ANI comparison, which gave a value of 91.78% to *A.*
timonensis. GTDB-Tk classified strain CLA-KB-H122 as 'Alistipes senegalensis'.
However, the latter name is as yet not valid. Functional analysis showed the strain

has 83 transporters, 17 secretion genes, and predicted utilization of cellulose and
starch along with production of L-glutamate and folate. In total, 395 CAZymes were
identified, with 59 different glycoside hydrolase families and 19 glycoside transferase
families represented. Major (≥5 %) cellular fatty acids after 72 h of growth in DSMZ
medium 1611 included 15:0 ISO FAME (19.4 %), 15:0 FAME (18.9 %), 16:0 ISO
FAME (10.6 %), 17:0 ISO 3OH FAME (8.8 %), 17:0 FAME (7.2 %), and 17:0 3OH
FAME (6.9 %). The type strain, CLA-KB-H122[T] (phylum *Bacteroidota*, family
Rikenellaceae) (=DSM 118481) (StrainInfo: 10.60712/SI-ID414389.1, genome:
GCA_040095975.1), was isolated from human faeces.

Description of *Bifidobacterium hominis* sp. nov.

Bifidobacterium hominis (ho'mi.nis. L. gen. n. *hominis*, of a human being, pertaining
to the human gut habitat, from where the type strain was isolated).

The genome size is 2.03 Mbp, G+C percentage is 55.98%, with 99.77%
completeness and 0.45% contamination. The closest relative to strain CLA-AA-H311
was *Bifidobacterium pseudocatenulatum* (99.07%) based on 16S rRNA gene
analysis. However, ANI comparison identified strain CLA-AA-H311 as a novel
species within the genus *Bifidobacterium*, with an ANI value of 93.18% against
B. pseudocatenulatum. GTDB-Tk classification as 'Bifidobacterium sp002742445'
confirmed the proposition of a novel species. Placement within the genus
Bifidobacterium was confirmed by the presence of fructose-6-phosphate
phosphoketolase (KO1621) [111]. Functional analysis showed the strain has 90
transporters, 20 secretion genes, and predicted utilization of starch and production of
propionate, acetate, and folate. In total, 137 CAZymes were identified, with 28
different glycoside hydrolase families and 11 glycoside transferase families
represented. Major (≥5 %) cellular fatty acids after 24 h of growth in DSMZ medium
1203a included 16:0 FAME (28.8 %), 18:0 FAME (15.3 %), 18:1 CIS 9 DMA (15.1
%), 18:1 CIS 9 FAME (14.3 %), and 14:0 FAME (7.8 %). The type strain, CLA-AA-
H311[T] (phylum *Actinomycetota*, family *Bifidobacteriaceae*) (=DSM 118068, =LMG
33596) (StrainInfo: 10.60712/SI-ID414317.1, genome: GCA_040095915.1), was
isolated from human faeces.

Description of *Blautia aquisgranensis* sp. nov.

Blautia aquisgranensis (a.quis.gra.nen'sis. L. fem. adj. *aquisgranensis,* named after
the German city of Aachen (Latin name *Aquisgranum*) where it was isolated)

The genome size is 3.63 Mbp, G+C percentage is 43.76%, with 99.37%
completeness and 0.32% contamination. It includes two plasmids (37,495 bp; 1,036
bp). The closest relative to strain CLA-JM-H16 was *Blautia intestinalis* (96.11%)
based on 16S rRNA gene analysis. Placement of the strain within *Blautia* was
confirmed based on POCP comparison as values above 50% to multiple *Blautia*
species were obtained. However, comparison to the type species of the genus,

Blautia coccoides, gave a value of 42.36%. This inconsistency was also highlighted
by GTDB-Tk, which classified strain CLA-JM-H16 as 'Blautia_A sp900764225',
suggesting *Blautia* may require splitting into multiple genera in future. As the
separation of *Blautia* would require detailed analysis, which is outside the scope of
this manuscript, we propose strain CLA-JM-H16 as a novel species within *Blautia*. All
three novel species of *Blautia* described within this manuscript were confirmed to
represent distinct species based on ANI comparison (*B. aquisgranensis* Vs. *B.*
caccae, 81.74%; *B. aquisgranensis* Vs. *B. intestinihominis*, 76.95%; *B. caccae* Vs. *B.*
intestinihominis, 78.57%). Functional analysis revealed 157 transporters, 17
secretion genes, and predicted utilization of arbutin, salicin, sucrose, starch, and
production of acetate, propionate, folate, L-glutamate, riboflavin, and cobalamin. In
total, 205 CAZymes were identified, with 40 different glycoside hydrolase families
and 12 glycoside transferase families represented. The type strain, CLA-JM-H16^{T}
(phylum *Bacillota*, family *Lachnospiraceae*) (=DSM 114586, =LMG 33033)
(StrainInfo: 10.60712/SI-ID414368.1, genome: GCA_040096615.1), was isolated
from human faeces.

Description of *Blautia caccae* sp. nov.

Blautia caccae (cac'cae. N.L. fem. n. *cacca*, human ordure, faeces; from Gr. fem. n.
kakkê, human ordure, faeces; N.L. gen. n. *caccae*, of faeces, referring to the source
of isolate).

The genome size is 5.83 Mbp, G+C percentage is 46.73%, with 98.73%
completeness and 0.63% contamination. The closest relative to strain CLA-SR-H028
was *Blautia hominis* (98.66%) based on 16S rRNA gene analysis. ANI comparison
identified CLA-SR-H028 as a novel species within the genus *Blautia*, with all values
being below the species threshold. GTDB-Tk classification as 'Blautia sp001304935'
confirmed the proposition of a novel species within *Blautia*. Functional analysis
showed the strain has 158 transporters, 18 secretion genes, and predicted utilization
of cellobiose, sucrose, starch and production of propionate, acetate, cobalamin, and
folate. In total, 353 CAZymes were identified, with 53 different glycoside hydrolase
families and 15 glycoside transferase families represented. Major (≥5 %) cellular
fatty acids after 24 h of growth in DSMZ medium 1203a included 16:0 FAME (20.2
%), 14:0 FAME (19.8 %), 16:0 DMA (17.5 %), 18:1 CIS 11 DMA (6.8 %), and 14:0
DMA (5.9 %). The type strain, CLA-SR-H028^{T} (phylum *Bacillota*, family
Lachnospiraceae) (=DSM 118556, =LMG 33609) (Straininfo: 10.60712/SI-
ID414428.1, genome: GCA_040095955.1), was isolated from human faeces.

Description of *Blautia intestinihominis* sp. nov.

Blautia intestinihominis (in.tes.ti.ni.ho'mi.nis. L. neut. n. *intestinum*, intestine; L.
masc. n. *homo*, a human being; N.L. gen. n. *intestinihominis*, of the human intestine).

The genome size is 4.1 Mbp, G+C percentage is 43.49%, with 98.73%
completeness and 0.63% contamination. It includes a single plasmid of 22,629 bp.
The isolate was assigned to the species *Blautia obeum* (98.98%) based on 16S
rRNA gene analysis. However, ANI comparison to *B. obeum* clearly identified this
isolate as being a separate species (84.16%). This was confirmed by GTDB-Tk
classification as 'Blautia_A sp000436615', recommending the creation of a novel
species. Functional analysis showed the strain has 155 transporters, 18 secretion
genes, and predicted utilization of sucrose and starch, along with production of L-
glutamate, folate, propionate, and cobalamin. In total, 177 CAZymes were identified.
The type strain, CLA-AA-H95^T (phylum *Bacillota*, family *Lachnospiraceae*) (=DSM
111354, =LMG 33582) (StrainInfo: 10.60712/SI-ID414326.1, genome:
GCA_040096655.1), was isolated from human faeces.

Description of *Coprococcus intestinihominis* sp. nov.

Coprococcus intestinihominis (in.tes.ti.ni.ho'mi.nis. L. neut. n. *intestinum*, intestine; L.
masc. n. *homo*, a human being; N.L. gen. n. *intestinihominis*, of the human intestine).

The genome size is 3.6 Mbp, G+C percentage is 43.29%, with 98.43%
completeness and 2.52% contamination. A single plasmid of 20,255 bp was
detected. The closest relative to strain CLA-AA-H190 was *Coprococcus catus*
(96.76%) based on 16S rRNA gene analysis. ANI comparison identified CLA-AA-
H190 as a novel species within the genus *Coprococcus*, with an ANI value of
90.25% against the closest relative *C. catus*. GTDB-Tk classification as
'Coprococcus_A catus_A' confirmed the proposition of a novel species, but also
suggests that separation of *Coprococcus* into multiple genera could occur in future.
Functional analysis showed the strain has 119 transporters, 18 secretion genes, and
predicted utilization of starch and production of propionate, butyrate, acetate,
cobalamin, and folate. In total, 122 CAZymes were identified, with 19 different
glycoside hydrolase families and 13 glycoside transferase families represented. The
type strain, CLA-AA-H190^T (phylum *Bacillota*, family *Lachnospiraceae*) (=DSM
114688, =LMG 33015) (StrainInfo: 10.60712/SI-ID414125.1, genome:
GCA_040096555.1), was isolated from human faeces.

Description of *Enterobacter intestinihominis* sp. nov.

Enterobacter intestinihominis (in.tes.ti.ni.ho'mi.nis. L. neut. n. *intestinum*, the
intestine; L. masc. n. *homo*, a human being; N.L. gen. n. *intestinihominis*, of the
human intestine).

The genome size is 4.86 Mbp, G+C percentage is 54.88%, with 99.89%
completeness and 0.12% contamination. It contains two plasmids (4,416 bp; 2,494
bp). Strain CLA-AC-H004 was determined to be a strain of *Enterobacter*
quasihormaechei (99.80%) based on 16S rRNA gene analysis. Separation from
existing *Enterobacter* species was confirmed by ANI comparison, which gave a value

of 93.69% to *E. quasihormaechei*. GTDB-Tk classification of strain CLA-AC-H004 as
'Enterobacter hormaechei_A' supports the proposal of a novel species. An ANI value
of 99.01% was obtained when compared to *Enterobacter hormaechei* subsp.
hoffmannii. However, given the separation from *E. hormaechei* we propose it these
strains represent a separate species and not only a sub-species. ANI comparison
also highlighted similarity with *Pedobacter himalayensis* (95.89%), which has been
classified as 'Enterobacter hormaechei_B' within GTDB. However, this suggests
reclassification of *Pedobacter* may be required in future. Functional analysis showed
the strain has 497 transporters, 98 secretion genes, and predicted utilization of
arbutin, salicin, cellobiose, sucrose, and starch along with production of L-glutamate,
biotin, riboflavin, acetate, propionate, and folate. In total, 316 CAZymes were
identified, with 37 different glycoside hydrolase families and 19 glycoside transferase
families represented. Major (≥5 %) cellular fatty acids after 24 h of growth in DSMZ
medium 1203a included 16:0 FAME (38.3 %), 17:0 CYC FAME (18.6 %), 18:1 CIS
11 FAME (9.7 %), 14:0 FAME (9.0 %), 19:0 CYCLO CIS 11,12 FAME (9.0 %), and
14:0 3OH (8.8 %). The type strain, CLA-AC-H004^T (phylum *Pseudomonadota*, family
Enterobacteriaceae) (=DSM 118557, =LMG 33610) (StrainInfo: 10.60712/SI-
ID414328.1, genome: GCA_040096145.1), was isolated from human faeces.

1024

Description of *Enterocloster hominis* sp. nov.

Enterocloster hominis (ho'mi.nis. L. gen. n. *hominis*, of a human being).

The genome size is 6.52 Mbp, G+C percentage is 50.14%, with 99.16%
completeness and 2.53% contamination. It includes a single plasmid of 7,635 bp.
The closest relative to strain CLA-SR-H021 was *Enterocloster aldenensis* (98.44%)
based on 16S rRNA gene analysis. ANI comparison identified CLA-SR-H021 as a
novel species within the genus *Enterocloster*, with all values being below the species
threshold. GTDB-Tk classified CLA-SR-H021 as 'Enterocloster pacaense', a name
derived from the proposed species 'Lachnoclostridium pacaense'. However, the fact
that these two names are not valid supports the proposition of a novel species within
Enterocloster. Functional analysis revealed 118 transporters, 14 secretion genes,
and predicted utilization of cellobiose, starch and production of propionate, acetate,
and folate. In total, 307 CAZymes were identified, with 39 different glycoside
hydrolase families and 14 glycoside transferase families represented. Major (≥5 %)
cellular fatty acids after 24 h of growth in DSMZ medium 1203a included 16:0 FAME
(30.3 %), 16:1 CIS 9 DMA (13.7 %), 16:1 CIS 9 FAME (10.7 %), 14:0 FAME (10.4
%), 16:0 DMA (9.6 %), 18:1 CIS 11 DMA (5.6 %), and 18:1 CIS 11 FAME (5.0 %).
The type strain, CLA-SR-H021^T (phylum *Bacillota*, family *Lachnospiraceae*) (=DSM
118482, =LMG 33606) (StrainInfo: 10.60712/SI-ID414422.1, genome:
GCA_040096035.1), was isolated from human faeces.

1045

Description of *Faecalibacterium intestinale* sp. nov.

Faecalibacterium intestinale (in.tes.ti.na'le. N.L. neut. adj. *intestinale*, pertaining to
the intestine, from where the type strain was isolated).

The genome size is 2.97 Mbp, G+C percentage is 56.43%, with 100.0%
completeness and 0.0% contamination. The isolate was determined to be related to
F. prausnitzii (98.08%) based on 16S rRNA gene analysis. ANI comparison to *F.*
prausnitzii was just below species level assignment (94.46%), and GTDB-Tk
classification as 'Faecalibacterium prausnitzii_J' recommended the creation of a
novel species. Functional analysis showed the strain has 135 transporters, 18
secretion genes, and predicted utilization of starch and production of L-glutamate,
riboflavin, and cobalamin. In total, 159 CAZymes were identified, with 27 different
glycoside hydrolase families and 12 glycoside transferase families represented.
Production of butyrate (4.74 ± 0.30 mM) was confirmed for strain CLA-AA-H281
when grown in YCFA broth (DSMZ Medium No. 1611) in Hungate tubes for 48 h at
37 °C under anaerobic conditions (6% CO_2 and 4,7% H_2 in N_2). Major (≥5 %) cellular
fatty acids after 48 h of growth in DSMZ medium 215 included 16:0 FAME (33.2 %),
18:1 CIS 11 DMA (11.7 %), 18:1 CIS 11 FAME (11.4 %), 14:0 FAME (9.3 %), 16:0
DMA (8.4 %), and 16:1 CIS 9 FAME (6.7 %). The type strain, CLA-AA-H281[T]
(phylum *Bacillota*, family *Oscillospiraceae*) (=DSM 116193, =LMG 33027)
(StrainInfo: 10.60712/SI-ID414306.1, genome: GCA_040096575.1), was isolated
from human faeces.

1067

Description of *Faecalibacterium tardum* sp. nov.

Faecalibacterium tardum (tar'dum. L. neut. adj. tardum, pertaining to its slow
growth).

The genome size is 3.04 Mbp, G+C percentage is 56.28%, with 99.32%
completeness and 0.0% contamination. It contains one plasmid of 14,735 bp in size,
which encodes for vancomycin resistance via the *vanY* gene. The isolate was
determined to be closely related to *Faecalibacterium prausnitzii* (98.70%) based on
16S rRNA gene analysis. ANI comparison to *F. prausnitzii* was on the border of
species level assignment (95.05%), however, GTDB-Tk classification as
'Faecalibacterium prausnitzii_A' recommended the creation of a novel species.
Strain CLA-AA-H175 was confirmed to represent a distinct species to
Faecalibacterium intestinale (CLA-AA-H281, =DSM 116193) (ANI: 94.39%) and
Faecalibacterium faecis (CLA-JM-H7B, =DSM 114587) (ANI: 86.46%), also
described in this paper. Functional analysis showed the strain has 116 transporters,
18 secretion genes, and predicted utilization of starch, and production of L-
glutamate. In total, 177 CAZymes were identified, with 28 different glycoside
hydrolase families and 12 glycoside transferase families represented. Production of
butyrate (0.81 ± 0.16 mM) was confirmed for strain CLA-AA-H175 when grown in
YCFA broth (DSMZ Medium No. 1611) in Hungate tubes for 48 h at 37 °C under
anaerobic conditions (6% CO_2 and 4,7% H_2 in N_2). The type strain, CLA-AA-H175[T]

(phylum *Bacillota*, family *Oscillospiraceae*) (=DSM 116192) (StrainInfo: 10.60712/SI-
ID414281.1, genome: GCA_040096515.1), was isolated from human faeces.

1090

Description of *Faecousia* gen. nov.

Faecousia (Faec.ou'si.a. L. fem. n. *faex*, dregs; N.L. fem. n. *ousia*, an essence; N.L.
fem. n. *Faecousia*, a microbe associated with faeces).

Based on 16S rRNA gene sequence identity, the closest relatives are members of
the genus *Vescimonas* (*Vescimonas fastidiosa*, 93.48%). POCP analysis against
Vescimonas coprocola (44.75%), the type species of this genus, and *V. fastidiosa*
(41.06%) confirmed strain CLA-AA-H192 represents a distinct genus to *Vescimonas*.
GTDB-Tk supported the creation of a novel genus, placing strain CLA-AA-H192
within the proposed genus of "*Candidatus Faecousia*". The type species of this
genus is *Faecousia intestinalis*.

1101

Description of *Faecousia intestinalis* sp. nov.

Faecousia intestinalis (in.tes.ti.na'lis. N.L. fem. adj. *intestinalis*, pertaining to the
intestines, from where the type strain was isolated).

The genome size is 2.98 Mbp, G+C percentage is 58.44%, with 93.29%
completeness and 0.0% contamination. It contains two plasmids (23,094 bp; 3,448
bp). Functional analysis showed the strain has 144 transporters, 26 secretion genes,
and predicted utilization of starch, and production of acetate, propionate, L-
glutamate, and folate. In total, 116 CAZymes were identified, with 19 different
glycoside hydrolase families and 11 glycoside transferase families represented. The
type strain, CLA-AA-H192[T] (phylum *Bacillota*, family *Oscillospiraceae*) (StrainInfo:
10.60712/SI-ID414286.1, genome: GCA_040096185.1), was isolated from human
faeces.

1114

Description of *Flavonifractor hominis* sp. nov.

Flavonifractor hominis (ho'mi.nis. L. gen. n. *hominis*, of a human being).

The genome size is 2.94 Mbp, G+C percentage is 58.64%, with 99.33%
completeness and 0.0% contamination. Strain CLA-AP-H34 was determined to be a
new species based on 16S rRNA gene sequence analysis, with the closest validly
named match being *Flavonifractor plautii* (97.21%). Separation from existing
Flavonifractor species was confirmed by ANI comparison, which gave a value of
82.25% to *F. plautii*. GTDB-Tk classification of strain CLA-AP-H34 as an unknown
species within *Flavonifractor* supports the proposal of a novel species. Functional
analysis showed the strain has 112 transporters, 14 secretion genes, and predicted
utilization of starch and production of L-glutamate, riboflavin, butyrate, and folate. In
total, 140 CAZymes were identified, with 19 different glycoside hydrolase families

and 15 glycoside transferase families represented. Major (≥5 %) cellular fatty acids
after 72 h of growth in DSMZ medium 1611 included 16:0 DMA (25.2 %), 15:0 FAME
(15.7 %), 15:0 DMA (11.4 %), 14:0 FAME (8.9 %), 15:0 ISO FAME (8.4 %), 17:0
DMA (7.5 %), and 16:0 FAME (6.4 %). The type strain, CLA-AP-H34^T (phylum
Bacillota, family *Oscillospiraceae*) (=DSM 118484, =LMG 33602) (StrainInfo:
10.60712/SI-ID414347.1, genome: GCA_040095835.1), was isolated from human
faeces.

1134

Description of *Hominiventricola aquisgranensis* sp. nov.

Hominiventricola aquisgranensis (a.quis.gra.nen'sis. L. masc. adj. *aquisgranensis,*
named after the German city of Aachen, where it was isolated).

The genome size is 3.17 Mbp, G+C percentage is 45.01%, with 99.37%
completeness and 0.27% contamination, including a single plasmid of 7,983 bp. The
closest relative to strain CLA-AA-H78B was *Hominiventricola filiformis* (96.24%)
based on 16S rRNA gene analysis. POCP analysis confirmed genus assignment to
Hominiventricola, with a POCP value of 69.06% to the type strain of the only current
species within this genus, *H. filiformis*. GTDB-Tk classified strain CLA-AA-H78B as
'Choladocola sp003480725' within "*Candidatus* Choladocola". The latter is a
heterosynonym of *Hominiventricola*, a validly published name. ANI comparison
confirmed that strain CLA-AA-H78B represents a novel species as the ANI value to
H. filiformis was 82.98%. Functional analysis showed the strain has 134 transporters,
31 secretion genes, and predicted utilization of cellobiose, starch, arbutin, salicin,
and production of L-glutamate, folate, acetate, propionate, and cobalamin. In total,
134 CAZymes were identified. The type strain, CLA-AA-H78B^T (phylum *Bacillota*,
family *Lachnospiraceae*) (=DSM 111355, =LMG 33583) (StrainInfo: 10.60712/SI-
ID414323.1, genome: GCA_040096225.1), was isolated from human faeces.

1153

Description of *Lachnospira intestinalis* sp. nov.

Lachnospira intestinalis (in.tes.ti.na'lis. N.L. fem. adj. *intestinalis*, pertaining to the
intestines, from where the type strain was isolated).

The genome size is 3.1 Mbp, G+C percentage is 41.75%, with 99.33%
completeness and 0.0% contamination. Strain CLA-AA-H89B was determined to
represent a separate species to its closest relative, *Lachnospira pectinoschiza*
(97.48%), based on 16S rRNA gene analysis. This was confirmed based on ANI
comparison to all close relatives which were below the species threshold. GTDB-Tk
classification of strain CLA-AA-H89B as 'Lachnospira sp000437735' confirmed that
this isolate represents a novel species. Functional analysis showed the strain has
112 transporters, 29 secretion genes, and predicted utilization of starch and
cellulose, along with production of L-glutamate, folate, acetate, propionate, and
riboflavin. Motility was predicted based on the presence of the following genes: *FlhA,*
FlhB, FlgB, FlgC, FlgD, FlgE, FlgF, FlgJ, FlgK, FlgL, FliC, FliD, FliE, FliF, FliG, FliK,

FliM, FliN, MotA, MotB. In total, 162 CAZymes were identified, with 21 different
glycoside hydrolase families and 11 glycoside transferase families represented.
Major (≥5 %) cellular fatty acids after 48 h of growth in DSMZ medium 1611 included
16:0 FAME (32.4 %), 18:1 CIS 11 DMA (20.6 %), 14:0 FAME (7.8 %), 16:0 DMA (6.3
%), 18:1 CIS 11 aldehyde (6.2 %), and 16:1 CIS 9 DMA (5.5 %). The type strain,
CLA-AA-H89B^T (phylum *Bacillota*, family *Lachnospiraceae*) (=DSM 118070)
(StrainInfo: 10.60712/SI-ID414325.1, genome: GCA_040095895.1), was isolated
from human faeces.

1176

Description of *Lachnospira hominis* sp. nov.

Lachnospira hominis (ho'mi.nis. L. gen. masc. n. *hominis*, of a human being).

The genome size is 3.15 Mbp, G+C percentage is 37.05%, with 98.66%
completeness and 0.67% contamination. Strain CLA-JM-H10 was determined to
represent a separate species to its closest relative, *Lachnospira rogosae* (96.37%),
based on 16S rRNA gene analysis. This was confirmed by an ANI of 79.9% between
L. rogosa (CLA-AA-H255), and strain CLA-JM-H10. GTDB-Tk classification of CLA-
JM-H10 as 'Lachnospira sp900316325' confirmed that it represents a novel species.
Functional analysis showed the strain has 109 transporters, 27 secretion genes, and
predicted utilization of starch, and production of L-glutamate, folate, and cobalamin.
Motility was predicted based on detection of the following genes: *FlhA, FlhB, FlgB,*
FlgC, FlgD, FlgE, FlgF, FlgJ, FlgK, FlgL, FliC, FliD, FliE, FliF, FliG, FliK, FliM, FliN,
MotA, MotB. This is consistent with the type species of this genus being motile. In
total, 165 CAZymes were identified, with 22 different glycoside hydrolase families
and 12 glycoside transferase families represented. The type strain, CLA-JM-H10^T
(phylum *Bacillota*, family *Lachnospiraceae*) (=DSM 114599, =LMG 33585)
(StrainInfo: 10.60712/SI-ID414366.1, genome: GCA_040096395.1), was isolated
from human faeces.

1195

Description of *Lachnospira rogosae* sp. nov.

Lachnospira rogosae (ro.go'sae. N.L. gen. masc. n. *rogosae*, of Rogosa).

Strain CLA-AA-H255 was determined to be similar to *Lactobacillus rogosae*
(99.87%) based on 16S rRNA gene analysis. However, the lack of a genome for the
type strain of the latter species, along with the lack of the type strain at any
established culture collection prevented further comparison (Tindall, 2014). GTDB-Tk
classification of CLA-AA-H255 as 'Lachnospira rogosae_A' suggested that it
represents a species within a genus distantly related to *Lactobacillus*. This was
reconfirmed by the same placement of a second strain, CLA-AA-H191, which was
also assigned to the same placeholder by GTDB-Tk (ANI of 98.86% between our two
isolates). Based on these results, we propose *L. rogosae* was previously
misassigned as a member of the genus *Lactobacillus*. To provide a type strain, and
correct its placement, we propose the creation of the species *Lachnospira rogosae*.

Functional analysis showed that strain CLA-AA-H255 has 100 transporters, 25
secretion genes, and predicted utilization of starch, and production of L-glutamate,
folate, and riboflavin. The prediction of motility and acetate production based on
genomic analysis is consisting with the observed phenotype of motility in the original
type strain of *L. ragosae* as stated by Holdeman and Moore (Holdeman and Moore,
1974). In total, 142 CAZymes were identified, including 20 different glycoside
hydrolase families and 12 glycoside transferase families. Ecological analysis of
1,000 human gut 16S rRNA gene amplicon samples identified this strain in 1.20% of
samples with a relative abundance of 0.43 ± 0.78%. The type strain, CLA-AA-H255[T]
(phylum *Bacillota*, family *Lachnospiraceae*) (=DSM 118602, =LMG 33594)
(StrainInfo: 10.60712/SI-ID414300.1, genome: GCA_040096455.1), was isolated
from human faeces.

1221

Description of *Laedolimicola intestinihominis* sp. nov.

Laedolimicola intestinihominis (in.tes.ti.ni.ho'mi.nis. L. neut. n. *intestinum*, intestine;
L. masc. n. *homo*, a human being; N.L. gen. n. *intestinihominis*, of the human
intestine).

The genome size is 3.45 Mbp, G+C percentage is 49.65%, with 99.37%
completeness and 0.16% contamination. It includes a single plasmid of 5,131 bp.
Strain CLA-AA-H132 was determined to represent a separate species to its closest
relative, *Laedolimicola ammoniilytica* (98.38%), based on 16S rRNA gene analysis.
POCP analysis confirmed that strain CLA-AA-H132 belongs to the recently named
genus, *Laedolimicola*, with a POCP value of 75.19% to the type strain of the only
current species within this genus, *L. ammoniilytica*. GTDB-Tk placement as
'Merdisoma sp900553635' suggests that *Laedolimicola* and *Candidatus* Merdisoma
are homonyms and require future reclassification. ANI comparison confirmed that
CLA-AA-H132 represents a novel species, as the ANI value to *L. ammoniilytica* was
only 90.2%. Functional analysis showed the strain has 142 transporters, 21 secretion
genes, and predicted utilization of cellobiose, sucrose, starch, and production of
acetate, propionate, L-glutamate, cobalamin, and folate. In total, 153 CAZymes were
identified, with 21 different glycoside hydrolase families and 16 glycoside transferase
families represented. Major (≥5 %) cellular fatty acids after 24 h of growth in DSMZ
medium 339 included 16:0 FAME (24.8 %), 18:1 CIS 11 DMA (12.8 %), 14:0 FAME
(10.0 %), 16:0 DMA (8.2 %), and 16:0 CIS 9 DMA (8.2 %). The type strain, CLA-AA-
H132[T] (phylum *Bacillota*, family *Lachnospiraceae*) (=DSM 117481, =LMG 33588)
(StrainInfo: 10.60712/SI-ID414271.1, genome: GCA_040096155.1), was isolated
from human faeces.

1246

Description of *Maccoyibacter* gen. nov.

Maccoyibacter (Mac.co.y.i.bac'ter N.L. fem. gen. n. *Maccoyae*, referring to the
immunologist Kathy McCoy; N.L. masc. n. *bacter*, a rod; N.L. masc.

n. *Maccoyibacter*, a rod-shaped bacterium named after the immunologist Kathy
McCoy, for her many scientific contributions in the field of microbe-host interactions).

Based on 16S rRNA gene analysis, the closest species with a valid name are
Roseburia hominis (94.9%) and *Eubacterium oxidoreducens* (94.23%). POCP values
against all close relatives were below the genus delineation threshold of 50%,
including many *Roseburia* spp. (*R. hominis*, *R. faecis*, *R. porci*, *R. intestinalis*), apart
from *R. inulinivorans* (50.83%). GTDB-Tk placement as "UBA11774 sp003507655"
confirmed that strain CLA-AA-H185 represents a novel genus within the family
Lachnospiraceae. The type species is *Maccoyibacter intestinihominis*.

1259

Description of *Maccoyibacter intestinihominis* sp. nov.

Maccoyibacter intestinihominis (in.tes.ti.ni.ho'mi.nis. L. neut. n. *intestinum*, intestine;
L. masc. n. *homo*, a human being; N.L. gen. n. *intestinihominis*, of the human
intestine).

The genome size is 2.85 Mbp, G+C percentage is 41.17%, with 97.41%
completeness and 0.45% contamination. It contains a single plasmid (2,341 bp).
Functional analysis showed the strain has 99 transporters, 35 secretion genes, and
predicted utilization of starch, and production of acetate, propionate, L-glutamate,
cobalamin, folate, and riboflavin. In total, 110 CAZymes were identified, with 17
different glycoside hydrolase families and 13 glycoside transferase families
represented. The type strain, CLA-AA-H185^{T} (phylum *Bacillota*, family
Lachnospiraceae) (=DSM 118601) (StrainInfo: 10.60712/SI-ID414284.1, genome:
GCA_040096355.1), was isolated from human faeces.

1273

Description of *Megasphaera intestinihominis* sp. nov.

Megasphaera intestinihominis (in.tes.ti.ni.ho'mi.nis. L. gen. neut. n. *intestini*, of the
intestine; L. gen. masc. n. *hominis*, of a human being; N.L. gen. masc.
n. *intestinihominis*, of the human intestine).

The genome size is 2.38 Mbp, G+C percentage is 53.59%, with 100.00%
completeness and 0.00% contamination. The closest relative to strain CLA-AA-H81
was *Megasphaera indica* (99.09%) based on 16S rRNA gene analysis. ANI
comparison to all close relatives were below species assignment threshold (highest
to *Megasphaera elsdenii,* 90.78%). GTDB-Tk classification as 'Megasphaera
sp000417505' supports the creation of a novel species. Functional analysis showed
the strain has 133 transporters, 16 secretion genes, and predicted utilization of
starch, and production of butyrate, propionate, L-glutamate, folate, riboflavin, and
cobalamin. Ecological analysis identified 152 matching (MASH distance < 0.05)
MAGs, of which 124 originate from the human gut, suggesting this species is most
commonly observed within this environment. This is supported by ecological analysis
using 16S rRNA gene amplicon datasets which identified it within 19.0% of 1,000

human gut samples, with a relative abundance of 2.14 ± 5.23%. Major (≥5 %)
cellular fatty acids after 24 h of growth in DSMZ medium 215 included 12:0 FAME
(17.1 %), 14:0 3OH (14.8 %), 18:1 CIS 9 FAME (9.6 %), 18:1 CIS 9 DMA (9.3 %),
16:0 FAME (8.0 %), and 16:1 CIS 7 FAME (5.0 %). The type strain, CLA-AA-H81[T]
(phylum *Bacillota*, family *Veillonellaceae*) (=DSM 118069, =LMG 33597) (StrainInfo:
10.60712/SI-ID414324.1, genome: GCA_040096415.1), was isolated from human
faeces.

1297

Description of *Niallia hominis* sp. nov.

Niallia hominis (ho'mi.nis. L. gen. n. *hominis*, of a human being).

The genome size is 4.9 Mbp, G+C percentage is 35.33%, with 92.24%
completeness and 3.20% contamination. Strain CLA-SR-H024 was assigned to
Niallia circulans (100%) based on 16S rRNA gene analysis. However, ANI
suggested the isolate represents a novel species as comparison to all *Niallia* spp.
were below 80%. GTDB-Tk classification of strain CLA-SR-H024 as 'Niallia
sp001076885' confirmed that it represents a novel species. Functional analysis
showed the strain has 244 transporters, 43 secretion genes, and predicted utilization
of arbutin, salicin, cellobiose, starch, and dextran, along with production of L-
glutamate, folate, and cobalamin. In total, 294 CAZymes were identified, with 34
different glycoside hydrolase families and 14 glycoside transferase families
represented. Major (≥5 %) cellular fatty acids after 72 h of growth in DSMZ medium
1203a included 15:0 ANTEISO FAME (29.7 %), 16:0 FAME (16.8 %), 15:0 ISO
FAME (14.9 %), 16:0 ISO FAME (9.8 %), 14:0 FAME (8.4 %), 17:0 ANTEISO FAME
(6.9 %), and 14:0 ISO FAME (5.2 %).The type strain, CLA-SR-H024[T] (phylum
Bacillota, family *Bacillaceae*) (=DSM 118483) (StrainInfo: 10.60712/SI-ID414424.1,
genome: GCA_040095995.1), was isolated from human faeces.

1316

Description of *Peptoniphilus hominis* sp. nov.

Peptoniphilus hominis (ho'mi.nis. L. gen. n. *hominis*, of a human being).

The genome size is 1.99 Mbp, G+C percentage is 33.8%, with 99.3% completeness
and 0.0% contamination. Strain CLA-SR-H025 was assigned to *Peptoniphilus*
gorbachii (99.11%) based on 16S rRNA gene analysis. However, ANI suggested the
isolate represents a novel species as comparison to *P. gorbachii* gave a value of
88.92%. GTDB-Tk classification of strain CLA-SR-H025 as 'Peptoniphilus_A
grossensis' confirmed that it represents a novel species, as the name 'Peptoniphilus
grossensis' was proposed but never validated [112]. Functional analysis showed the
strain has 12 transporters, 1 secretion genes, and no gut metabolic models were
identified within the genome. In total, 59 CAZymes were identified, with 10 different
glycoside hydrolase families and 8 glycoside transferase families represented. The
type strain, CLA-SR-H025[T] (phylum *Bacillota*, family *Peptoniphilaceae*) (=DSM

118555, =LMG 33608) (Straininfo: 10.60712/SI-ID414425.1, genome:
GCA_040096015.1), was isolated from human faeces.

1332

Description of *Pseudoflavonifractor intestinihominis* sp. nov.

Pseudoflavonifractor intestinihominis (in.tes.ti.ni.ho'mi.nis. L. neut. n. *intestinum*, the
intestine; L. masc. n. *homo*, a human being; N.L. gen. n. *intestinihominis*, of the
human intestine).

The genome size is 3.69 Mbp, G+C percentage is 61.54%, with 99.33%
completeness and 0.00% contamination. Strain CLA-AP-H29 was assigned to the
species *Pseudoflavonifractor capillosus* (99.53%) based on 16S rRNA gene
sequence analysis. However, ANI suggested the isolate represents a novel species
as comparison to *P. capillosus* gave a value of 80.96%. GTDB-Tk classified strain
CLA-AP-H29 as 'Pseudoflavonifractor sp944387275', which confirms that it
represents a novel species. Functional analysis predicted the strain has 121
transporters, 14 secretion genes, and predicted utilization of starch, and production
of L-glutamate, folate, and cobalamin. In total, 180 CAZymes were identified, with 25
different glycoside hydrolase families and 15 glycoside transferase families
represented. Major (≥5 %) cellular fatty acids after 48 h of growth in DSMZ medium
215 included 14:0 FAME (33.9 %), 16:0 DMA (33.8 %), 16:0 FAME (9.8 %), and 16:0
ALDE (8.0 %). The type strain, CLA-AP-H29^T (phylum *Bacillota*, family
Oscillospiraceae) (=DSM 118073, =LMG 33601) (Straininfo: 10.60712/SI-
ID414342.1, genome: GCA_040096055.1), was isolated from human faeces.

1352

Description of *Robertmurraya yapensis* sp. nov.

Robertmurraya yapensis (yap'ensis. N.L. fem. adj. *yapensis*, pertaining to Yap
trench, which is the geographical position where the first isolate of this species was
obtained).

The genome size is 4.74 Mbp, G+C percentage is 37.94%, with 98.85%
completeness and 2.13% contamination. It contains a plasmid (2,470 bp). The
closest relative to strain CLA-AA-H227 was *Robertmurraya spiralis* (99.23%) based
on 16S rRNA gene analysis. The highest POCP scores were to members of the
genus *Robertmurraya (Robertmurraya kyonggiensis,* 84.9%; *R. spiralis,* 65.33%).
The placement of strain CLA-AA-H227 within *Robertmurraya* was confirmed by
GTDB-Tk assignment as 'Robertmurraya yapensis', a reclassification of the species
'Bacillus yapensis', although neither name has been validated. ANI comparison
confirmed that CLA-AA-H227 represents a novel species, as values to close
relatives were below the species level threshold. Functional analysis showed the
strain has 246 transporters, 53 secretion genes, and predicted utilization of arbutin,
salicin, cellobiose, starch, and production of butyrate, acetate, propionate, folate,
riboflavin, and cobalamin. In total, 238 CAZymes were identified. The type strain,
CLA-AA-H227^T (phylum *Bacillota*, family *Bacillaceae*) (=DSM 113004, =LMG 33018)

(StrainInfo: 10.60712/SI-ID414150.1, genome: GCA_040096375.1), was isolated
from human faeces.

1373

Description of *Ruminococcoides intestinale* sp. nov.

Ruminococcoides intestinale (in.tes.ti.na'le. N.L. neut. adj. *intestinalis*, pertaining to
the intestines, from where the type strain was isolated).

The genome size is 2.32 Mbp, G+C percentage is 40.88%, with 99.33%
completeness and 1.01% contamination. The isolate was determined to be similar to
Ruminococcus bromii (98.91%) and more distantly related to *Ruminococcoides bili*
(96.76%) based on 16S rRNA gene analysis. While POCP comparison of strain
CLA-JM-H38 to *R. bromii* was 59.79%, and 53.56% to *Ruminococcus bovis,*
suggesting they belong to the same genus, all other comparisons to *Ruminococcus*
species were below 50%, including to the type species, *Ruminococcus flavefaciens*
(27.58%). POCP to *R. bili*, the type species of the genus *Ruminococcoides*, was
64.44%. GTDB-Tk classified strain CLA-JM-H38 as "Ruminococcus_E bromii_B",
confirming it is not a member of the genus *Ruminococcus.* These results support
GTDB assignment that both *R. bovis* and *R. bromii* should be reclassified as
members of the genus *Ruminococcoides.* Strain CLA-JM-H38 was confirmed to
represent a novel species as all ANI comparisons to close relatives were below 95%,
and it represents a distinct novel species from *Ruminococcoides intestinihominis*
described in this work (78.33%). Functional analysis showed the strain has 81
transporters, 15 secretion genes, and predicted utilization of starch, and production
of L-glutamate. In total, 108 CAZymes were identified, with 15 different glycoside
hydrolase families and 12 glycoside transferase families represented. Ecological
analysis based on 16S rRNA gene amplicons identified this species in 55.20% of
1,000 human gut samples with a relative abundance of 1.50 ± 2.49%, suggesting it is
a prevalent and dominant bacterial species within the human gut. Major (≥5 %)
cellular fatty acids after 72 h of growth in DSMZ medium 1611 included 16:0 ISO
FAME (49.5 %) and 16:0 ISO DMA (23.8 %). The type strain, CLA-JM-H38[T] (phylum
Bacillota, family *Oscillospiraceae*) (=DSM 118486, =LMG 33604) (StrainInfo:
10.60712/SI-ID414376.1, genome: GCA_040096305.1), was isolated from human
faeces.

1403

Description of *Ruminococcoides intestinihominis* sp. nov.

Ruminococcoides intestinihominis (in.tes.ti.ni.ho'mi.nis. L. neut. n. *intestinum*,
intestine; L. masc. n. *homo*, a human being; N.L. gen. n. *intestinihominis*, of the
human intestine).

The genome size is 2.26 Mbp, G+C percentage is 34.26%, with 98.66%
completeness and 0.00% contamination. It contains one plasmid (1,825 bp). Based
on 16S rRNA gene sequence identity, the isolate was closely related to
Ruminococcus bovis (99.3%) and more distant to *Ruminococcoides bili* (94.8%), the

type species of this genus. ANI comparison confirmed the similarity of strain CLA-
AA-H171 to *R. bovis* (95.4%), however classification by GTDB-Tk as
'Ruminococcus_E sp934476515' supported the creation of a novel species.
Recently, 'Ruminococcus_E' has been validly named as *Ruminococcoides*, with the
type species *R. bili* [34]. POCP comparison between the isolate and *R. bili* provided a
value of 51.3%, suggesting that strain CLA-AA-H171 represents a novel species
within the genus *Ruminococcoides*. Functional analysis showed the strain has 75
transporters, 14 secretion genes, and predicted utilization of starch, and production
of acetate. In total, 124 CAZymes were identified, with 13 different glycoside
hydrolase families and 12 glycoside transferase families represented. Ecological
analysis based on 16S rRNA gene amplicons identified this species in 10.40% of
1,000 human gut samples with a relative abundance of 0.23 ± 0.68%. The type
strain, CLA-AA-H171^T (phylum *Bacillota*, family *Oscillospiraceae*) (=DSM 114689,
=LMG 33587) (StrainInfo: 10.60712/SI-ID414279.1, genome: GCA_040096285.1),
was isolated from human faeces.

1427

Description of *Ruthenibacterium intestinale* sp. nov.

Ruthenibacterium intestinale (in.tes.ti.na'le. N.L. neut. adj. intestinale, pertaining to
the intestines, from where the type strain was isolated).

The genome size is 3.1 Mbp, G+C percentage is 54.69%, with 98.3% completeness
and 0.00% contamination. It contains a plasmid (5,063 bp). The closest relative to
strain CLA-JM-H11 is *Ruthenibacterium lactatiformans* (94.93%), the type species of
this genus, based on 16S rRNA gene analysis. Placement of the isolate within the
genus *Ruthenibacterium* was confirmed based on POCP comparison, with a value of
56.41% to the type species. Strain CLA-JM-H11 was confirmed to be distinct to *R.*
lactatiformans based on ANI comparison (79.28%). GTDB-Tk classification
confirmed this assignment as an unknown species within *Ruthenibacterium*.
Functional analysis showed the strain has 143 transporters, 16 secretion genes, and
predicted utilization of starch, and production of acetate, propionate, folate, and
cobalamin. In total, 155 CAZymes were identified, with 27 different glycoside
hydrolase families and 13 glycoside transferase families represented. The type
strain, CLA-JM-H11^T (phylum *Bacillota*, family *Oscillospiraceae*) (=DSM 114604,
=LMG 33032) (StrainInfo: 10.60712/SI-ID414367.1, genome: GCA_040096265.1),
was isolated from human faeces.

1446

Description of *Solibaculum intestinale* sp. nov.

1448   *Solibaculum intestinale (in.tes.ti.na'le. N.L. neut. adj. intestinale,* pertaining to the
1449   intestines, from where the type strain was isolated).

1450   The genome size is 2.81 Mbp, G+C percentage is 54.62 %, with 97.99%
1451   completeness and 0.67% contamination. The isolate was determined to be a new
1452   species based on 16S rRNA gene analysis, with the closest validly named match

being *Solibaculum mannosilyticum* (94.93%). Placement within *Solibaculum* was confirmed with POCP values above 50% to *S. mannosilyticum* (55.7%), the type species, and only member of this genus. ANI comparison confirmed the isolate represents a novel species, as no ANI values were above 80%. Functional analysis showed the strain has 82 transporters, 16 secretion genes, and predicted utilization of starch, and production of acetate, propionate, and L-glutamate. In total, 169 CAZymes were identified, with 20 different glycoside hydrolase families and 15 glycoside transferase families represented. The type strain, CLA-JM-H44[T] (phylum *Bacillota*, family *Oscillospiraceae*) (=DSM 114601, =LMG 33034) (StrainInfo: 10.60712/SI-ID414377.1, genome: GCA_040096205.1), was isolated from human faeces.

Description of *Ventrimonas* gen. nov.

Ventrimonas (Ven.tri.mo'nas. L. masc. n. *venter*, the belly; L. fem. n. *monas*, a monad; N.L. fem. n. *Ventrimonas*, a microbe associated with the belly (intestines/faeces)).

Based on 16S rRNA gene sequence identity, the closest relatives are members of the genus *Hungatella* (*Hungatella effluvii*, 95.15%; *Hungatella hathewayi*, 94.95%). POCP analysis against *H. effluvii* (40.38%) and *H. hathewayi* (39.86%) indicates that strain CLA-AP-H27 represents a distinct genus to *Hungatella*. GTDB-Tk supported the creation of a novel genus, placing strain CLA-AP-H27 within the proposed genus of "*Candidatus* Ventrimonas" in the family *Lachnospiraceae*. The type species of this genus is *Ventrimonas faecis*.

Description of *Ventrimonas faecis* sp. nov.

Ventrimonas faecis (fae'cis. L. gen. n. *faecis*, of dregs, pertaining to faeces, from where the type strain was isolated).

The genome size is 3.41 Mbp, G+C percentage is 49.27%, with 98.73% completeness and 0.63% contamination. Functional analysis showed the strain has 156 transporters, 21 secretion genes, and predicted utilization of starch, and production of acetate, propionate, L-glutamate, cobalamin, and folate. In total, 139 CAZymes were identified, with 21 different glycoside hydrolase families and 12 glycoside transferase families represented. Major (≥10 %) cellular fatty acids after 24 h of growth in DSMZ medium 1203a included 16:0 FAME (35.5 %), 18:1 CIS 11 DMA (10.0 %), 18:1 CIS 11 FAME (9.6 %), 18:1 CIS 9 DMA (5.9 %), and 16:0 DMA (5.5 %). The type strain, CLA-AP-H27[T] (phylum *Bacillota*, family *Lachnospiraceae*) (=DSM 118072, =LMG 33600) (StrainInfo: 10.60712/SI-ID414341.1, genome: GCA_040096075.1), was isolated from human faeces.

Description of *Waltera hominis* sp. nov.

Waltera hominis (ho'mi.nis. L. gen. masc. n. *hominis*, of a human being, pertaining to the human gut habitat, from where the type strain was isolated).

The genome size is 3.88 Mbp, G+C percentage is 45.72%, with 99.52% completeness and 2.13% contamination. The isolate was assigned to the species *Waltera intestinalis* (100.0%) based on 16S rRNA gene analysis. However, ANI comparison identified strain CLA-AA-H183 as a novel species within the genus *Waltera*, with an ANI value of 91.97% against the type species *Waltera intestinalis*. GTDB-Tk currently lacks the inclusion of *Waltera*, causing misclassification as 'Acetatifactor sp003447295', however this confirmed the proposition of a novel species. Separation of *Waltera* from *Acetatifactor* was revalidated via both phylogenomic analysis (**Supplementary Figure 3**), which shows both genera form separate monophyletic groups, and POCP analysis, which shows clear similarity within each genus (*Waltera*: 68.86%, *Acetatifactor*: 59.25%), and separation between the genera (39.94 ± 1.63 %). Functional analysis showed the strain has 141 transporters, 36 secretion genes, and predicted utilization of starch, cellulose, and production of butyrate, propionate, acetate, and folate. In total, 228 CAZymes were identified, with 38 different glycoside hydrolase families and 14 glycoside transferase families represented. The type strain, CLA-AA-H183^{T} (phylum *Bacillota*, family *Lachnospiraceae*) (=DSM 114684, =LMG 33586) (StrainInfo: 10.60712/SI-ID414283.1, genome: GCA_040096245.1), was isolated from human faeces.

Data availability

The genomes for all strains have been deposited at NCBI under BioProject: PRJNA996881. Bulk download of HiBC resources is possible via Zenodo for the genomes (https://doi.org/10.5281/zenodo.12755497), plasmid sequences (https://doi.org/10.5281/zenodo.12187897), 16S rRNA gene sequences (https://doi.org/10.5281/zenodo.12180259) and the isolates metadata (https://doi.org/10.5281/zenodo.14592301). The PacBio genome for *P. vulgatus* CLA-AA-H253 (=DSM 118718) has been deposited at ENA (PRJEB80480) and is transiently available via Zenodo: https://doi.org/10.5281/zenodo.14674027. The code for the creation of the HiBC website has been made available at; https://git.rwth-aachen.de/clavellab/hibc.

Acknowledgments

Sequencing was performed with the support of: (i) the DFG-funded NGS Competence Center Tübingen (INST 37/1049-1) and the Institute for Medical Microbiology and Hygiene at the University Hospital (Tübingen, Germany), including help by the Quantitative Biology Center (QBiC) for raw data management and storage; (ii) Nassos Typas and Carlos Geert Pieter Voogdt (EMBL, Heidelberg), for long-read sequencing; (iii) the Genomics Facility, a core facility of the Interdisciplinary Center for Clinical Research (IZKF) Aachen within the Faculty of

Medicine at RWTH Aachen University. We are also thankful to: (iv) Patrick Buchta from the Audio-Visual department at the University Hospital of RWTH Aachen for designing the HiBC logo; (v) Marzena Wyschkon and Meina Neumann-Schaal from the Leibniz Institute DSMZ for their help with the deposition of strains and CFA analysis, respectively; (vi) Catherine Gonzalez and Maurice Heizer from the Functional Microbiome Research Group (Institute of Medical Microbiology; University Hospital of RWTH Aachen) for data management with Coscine and for the curation of HiBC, respectively; (vii) Peter Vandamme, Claudine Vereecke, and Anneleen Wieme for handling the deposition of strains at the BCCM/LMG Bacteria Collection.

Funding

TCAH received funding from the German Research Foundation (DFG) as part of SFB1382 Gut-liver axis and from the RWTH Aachen START program, project "LeakyGut". TC received funding from the German Research Foundation (DFG): project no. 513892404, no. 445552570, no. 395357507 – SFB1371 Microbiome signatures, no. 403224013 – SFB1382 Gut-liver axis, and no. 460129525 – NFDI4Microbiota. JO received funding from the DFG, project no. 6270054 NFDI 28/1 "NFDI4Microbiota" and 6270048 NFDI 5/1 "NFDI4Biodiversity", the BMBF, projects no. 8005512901 and 8005512001 of DZIF, and EU/Horizon IRA project MICROBE no. 101094353. MDC and MG received funding from the DFG, project no. 403224013. The data used in this publication was managed using the research data management platform Coscine with storage space granted by the Research Data Storage (RDS) of the DFG and Ministry of Culture and Science of the State of North Rhine-Westphalia (DFG: INST222/1261-1 and MKW: 214-4.06.05.08 - 139057). LM and TA received funding from The Leverhulme Trust: project no. RF-2023-286\2.

Conflicts of interest

The authors have no conflicts of interest

References

1.   Forster, S. C. *et al.* A human gut bacterial genome and culture collection for improved metagenomic analyses. *Nat Biotechnol* **37**, 186–192 (2019).
2.   Groussin, M. *et al.* Elevated rates of horizontal gene transfer in the industrialized human microbiome. *Cell* **184**, 2053-2067.e18 (2021).
3.   Liu, C. *et al.* Enlightening the taxonomy darkness of human gut microbiomes with a cultured biobank. *Microbiome* **9**, 1–29 (2021).
4.   Lagier, J. *et al.* Culture of previously uncultured members of the human gut microbiota by culturomics. **1**, (2016).
5.   Browne, H. P. *et al.* Culturing of 'unculturable' human microbiota reveals novel taxa and extensive sporulation. *Nature* **533**, 543–546 (2016).
6.   Hedlund, B. P. *et al.* SeqCode: a nomenclatural code for prokaryotes described from sequence data. *Nat Microbiol* **7**, 1702–1708 (2022).

7. Göker, M., Moore, E. R. B., Oren, A. & Trujillo, M. E. Status of the SeqCode in
the International Journal of Systematic and Evolutionary Microbiology. *Int J*
Syst Evol Microbiol **72**, 10–12 (2022).
8. Arahal, D. *et al.* The best of both worlds: a proposal for further integration of
Candidatus names into the International Code of Nomenclature of
Prokaryotes. *Int J Syst Evol Microbiol* **74**, 1–21 (2024).
9. Hitch, T. C. A. *et al.* Harmonious naming across nomenclature codes
exemplified by the description of bacterial isolates from the mammalian gut.
Syst Appl Microbiol **47**, (2024).
10. Blanco-Míguez, A. *et al.* Extension of the Segatella copri complex to 13
species with distinct large extrachromosomal elements and associations with
host conditions. *Cell Host Microbe* **31**, 1804-1819.e9 (2023).
11. De Filippis, F. *et al.* Distinct Genetic and Functional Traits of Human Intestinal
Prevotella copri Strains Are Associated with Different Habitual Diets. *Cell Host*
Microbe **25**, 444-453.e3 (2019).
12. Fehlner-Peach, H. *et al.* Distinct Polysaccharide Utilization Profiles of Human
Intestinal Prevotella copri Isolates. *Cell Host Microbe* **26**, 680-690.e5 (2019).
13. Conteville, L. C. & Vicente, A. C. P. A plasmid network from the gut
microbiome of semi-isolated human groups reveals unique and shared
metabolic and virulence traits. *Sci Rep* **12**, 1–9 (2022).
14. Fogarty, E. C. *et al.* Article A cryptic plasmid is among the most numerous
genetic elements in the human gut ll ll Article A cryptic plasmid is among the
most numerous genetic elements in the human gut. *Cell* **187**, 1206-1222.e16
(2024).
15. Yu, M. K., Fogarty, E. C. & Eren, A. M. Diverse plasmid systems and their
ecology across human gut metagenomes revealed by PlasX and MobMess.
Nat Microbiol **9**, 830–847 (2024).
16. Perez, M. *et al.* A synthetic consortium of 100 gut commensals modulates the
composition and function in a colon model of the microbiome of elderly
subjects. *Gut Microbes* **13**, 1–19 (2021).
17. Clark, R. L. *et al.* Design of synthetic human gut microbiome assembly and
butyrate production. *Nat Commun* **12**, 1–16 (2021).
18. Cheng, A. G. *et al.* Design, construction, and in vivo augmentation of a
complex gut microbiome. *Cell* **185**, 3617-3636.e19 (2022).
19. Derrien, M., Vaughan, E. E., Plugge, C. M. & de Vos, W. M. Akkermansia
municiphila gen. nov., sp. nov., a human intestinal mucin-degrading bacterium.
Int J Syst Evol Microbiol **54**, 1469–1476 (2004).
20. Plovier, H. *et al.* A purified membrane protein from Akkermansia muciniphila or
the pasteurized bacterium improves metabolism in obese and diabetic mice.
Nat Med **23**, 107–113 (2017).
21. Overmann, J. Significance and future role of microbial resource centers. *Syst*
Appl Microbiol **38**, 258–265 (2015).
22. Parker, C. T., Tindall, B. J. & Garrity, G. M. International code of nomenclature
of Prokaryotes. *Int J Syst Evol Microbiol* **69**, S1 (2019).
23. Segata, N., Börnigen, D., Morgan, X. C. & Huttenhower, C. PhyloPhlAn is a
new method for improved phylogenetic and taxonomic placement of microbes.
Nat Commun **4**, 1–10 (2013).
24. Hitch, T. C. A. *et al.* Recent advances in culture-based gut microbiome
research. *International Journal of Medical Microbiology* **311**, (2021).

25. Browne, H. P. *et al.* Culturing of 'unculturable' human microbiota reveals novel
taxa and extensive sporulation. *Nature* **533**, 543–546 (2016).
26. Forster, S. C. *et al.* A human gut bacterial genome and culture collection for
improved metagenomic analyses. *Nat Biotechnol* **37**, 186–192 (2019).
27. Poyet, M. *et al.* A library of human gut bacterial isolates paired with longitudinal
multiomics data enables mechanistic microbiome research. *Nat Med* **25**,
1442–1452 (2019).
28. Groussin, M. *et al.* Elevated rates of horizontal gene transfer in the
industrialized human microbiome. *Cell* **184**, 2053-2067.e18 (2021).
29. Liu, C. *et al.* Enlightening the Taxonomy Darkness of Human Gut Microbiomes
With Cultured Biobank. *Microbiome* **9**, (2021).
30. Huang, P. *et al.* Gut microbial genomes with paired isolates from China
illustrate probiotic and cardiometabolic effects. *Cell Genomics* **4**, (2024).
31. Huang, Y. *et al.* High-throughput microbial culturomics using automation and
machine learning. *Nat Biotechnol* (2023) doi:10.1038/s41587-023-01674-2.
32. Lagier, J. C. *et al.* Culture of previously uncultured members of the human gut
microbiota by culturomics. *Nat Microbiol* **1**, (2016).
33. Leviatan, S., Shoer, S., Rothschild, D., Gorodetski, M. & Segal, E. An
expanded reference map of the human gut microbiome reveals hundreds of
previously unknown species. *Nat Commun* **13**, 1–14 (2022).
34. Molinero, N. *et al.* Ruminococcoides bili gen. Nov., sp. nov., a bile-resistant
bacterium from human bile with autolytic behavior. *Int J Syst Evol Microbiol* **71**,
1–11 (2021).
35. Schmitz, M. A., Dimonaco, N. J., Clavel, T. & Hitch, T. C. A. Lineage-specific
microbial protein prediction enables large-scale exploration of protein ecology
within the human gut. *bioRxiv* (2024) doi:10.1101/2024.05.29.596415.
36. Ma, L. *et al.* Spermidine improves gut barrier integrity and gut microbiota
function in diet-induced obese mice. *Gut Microbes* **12**, 1–19 (2020).
37. Wiedmann, S., Eudy, J. D. & Zempleni, J. Biotin supplementation increases
expression of genes encoding interferon-γ, interleukin-1β, and 3-
methylcrotonyl-CoA carboxylase, and decreases expression of the gene
encoding interleukin-4 in human peripheral blood mononuclear cells. *Journal of*
Nutrition **133**, 716–719 (2003).
38. Agrawal, S., Agrawal, A. & Said, H. M. Biotin deficiency enhances the
inflammatory response of human dendritic cells. *Am J Physiol Cell Physiol*
311, C386–C391 (2016).
39. Skupsky, J. *et al.* Biotin Supplementation Ameliorates Murine Colitis by
Preventing NF-κB Activation. *Cmgh* **9**, 557–567 (2020).
40. Mao, B. *et al.* Blautia producta displays potential probiotic properties against
dextran sulfate sodium-induced colitis in mice. *Food Science and Human*
Wellness **13**, 709–720 (2024).
41. van der Lelie, D. *et al.* Rationally designed bacterial consortia to treat chronic
immune-mediated colitis and restore intestinal homeostasis. *Nat Commun* **12**,
1–17 (2021).
42. Lee, C. *et al.* P926 Blautia Obeum Aggravates Colitis in a Murine Model. *J*
Crohns Colitis **17**, i1035–i1035 (2023).
43. Huang, T. Q. *et al.* Bergenin Alleviates Ulcerative Colitis By Decreasing Gut
Commensal Bacteroides vulgatus-Mediated Elevated Branched-Chain Amino
Acids. *J Agric Food Chem* **72**, 3606–3621 (2024).

44. Sugihara, K. *et al.* Dietary phosphate exacerbates intestinal inflammation in
experimental colitis. *J Clin Biochem Nutr* **61**, 91–99 (2017).
45. Kolachala, V. L. *et al.* A2B Adenosine Receptor Gene Deletion Attenuates
Murine Colitis. *Gastroenterology* **135**, 861–870 (2008).
46. Jones, B. V, Sun, F. & Marchesi, J. R. Comparative metagenomic analysis of
plasmid encoded functions in the human gut microbiome. *BMC Genomics* **11**,
(2010).
47. Groussin, M. *et al.* Elevated rates of horizontal gene transfer in the
industrialized human microbiome. *Cell* **184**, 2053-2067.e18 (2021).
48. Sanders, J. G. *et al.* A low-cost genomics workflow enables isolate screening
and strain-level analyses within microbiomes. *Genome Biol* **23**, (2022).
49. Rozov, R. *et al.* Recycler: An algorithm for detecting plasmids from de novo
assembly graphs. *Bioinformatics* **33**, 475–482 (2017).
50. Bankevich, A. *et al.* SPAdes: A new genome assembly algorithm and its
applications to single-cell sequencing. *Journal of Computational Biology* **19**,
455–477 (2012).
51. Li, Y. *et al.* Distribution of megaplasmids in Lactobacillus salivarius and other
Lactobacilli. *J Bacteriol* **189**, 6128–6139 (2007).
52. Bottacini, F. *et al.* Discovery of a conjugative megaplasmid in Bifidobacterium
breve. *Appl Environ Microbiol* **81**, 166–176 (2015).
53. Yu, M. K., Fogarty, E. C. & Eren, A. M. Diverse plasmid systems and their
ecology across human gut metagenomes revealed by PlasX and MobMess.
Nat Microbiol **9**, 830–847 (2024).
54. Schmartz, G. P. *et al.* PLSDB: Advancing a comprehensive database of
bacterial plasmids. *Nucleic Acids Res* **50**, D273–D278 (2022).
55. Novicki, T. J. & Hecht, D. W. *Characterization and DNA Sequence of the*
Mobilization Region of PLV22a from Bacteroides Fragilis. JOURNAL OF
BACTERIOLOGY vol. 177 (1995).
56. Nieto, C. *et al.* The yefM-yoeB toxin-antitoxin systems of Escherichia coli and
Streptococcus pneumoniae: Functional and structural correlation. *J Bacteriol*
189, 1266–1278 (2007).
57. Smith, J. A. & Magnuson, R. D. Modular Organization of the Phd
Repressor/Antitoxin Protein. *J Bacteriol* **186**, 2692–2698 (2004).
58. Evans, J. C. *et al.* A proteolytically activated antimicrobial toxin encoded on a
mobile plasmid of Bacteroidales induces a protective response. *Nat Commun*
13, (2022).
59. García-Bayona, L., Coyne, M. J. & Comstock, L. E. Mobile Type VI secretion
system loci of the gut Bacteroidales display extensive intra-ecosystem transfer,
multi-species spread and geographical clustering. *PLoS Genet* **17**, 1–25
(2021).
60. García-Bayona, L. *et al.* A pervasive large conjugative plasmid mediates
multispecies biofilm formation in the intestinal microbiota increasing resilience
to perturbations. Preprint at https://doi.org/10.1101/2024.04.29.590671 (2024).
61. Sokurenko, E. V, Chesnokova, V., Doyle, R. J. & Hasty, D. L. *Diversity of the*
Escherichia Coli Type 1 Fimbrial Lectin DIFFERENTIAL BINDING TO
MANNOSIDES AND UROEPITHELIAL CELLS. and the iResearch Service*
http://www.jbc.org.
62. Jønsson, R. *et al.* Aggregative adherence fimbriae form compact structures as
seen by SAXS. *Sci Rep* **13**, (2023).

63. García-Bayona, L. *et al.* A pervasive large conjugative plasmid mediates
multispecies biofilm formation in the intestinal microbiota increasing resilience
to perturbations. *bioRxiv* 2024.04.29.590671 (2024).
64. Riepe, S. P., Goldstein, J. & Alpers, D. H. Effect of secreted Bacteroides
proteases on human intestinal brush border hydrolases. *Journal of Clinical*
Investigation **66**, 314–322 (1980).
65. Mills, R. H. *et al.* Multi-omics analyses of the ulcerative colitis gut microbiome
link Bacteroides vulgatus proteases with disease severity. *Nat Microbiol* 1–15
(2022) doi:10.1038/s41564-021-01050-3.
66. Sokol, H. *et al.* Faecalibacterium prausnitzii is an anti-inflammatory commensal
bacterium identified by gut microbiota analysis of Crohn disease patients. *Proc*
Natl Acad Sci U S A **105**, 16731–16736 (2008).
67. Ueda, A. *et al.* Identification of Faecalibacterium prausnitzii strains for gut
microbiome-based intervention in Alzheimer's-type dementia. *Cell Rep Med* **2**,
100398 (2021).
68. D'hoe, K. *et al.* Integrated culturing, modeling and transcriptomics uncovers
complex interactions and emergent behavior in a three-species synthetic gut
community. *Elife* **7**, 1–30 (2018).
69. Fitzgerald, C. B. *et al.* Comparative analysis of Faecalibacterium prausnitzii
genomes shows a high level of genome plasticity and warrants separation into
new species-level taxa. *BMC Genomics* **19**, 1–20 (2018).
70. Vital, M., Karch, A. & Pieper, D. H. Colonic Butyrate-Producing Communities in
Humans□: an Overview Using Omics Data. *mSystems* **2**, 1–18 (2017).
71. Louis, P. & Flint, H. J. Formation of propionate and butyrate by the human
colonic microbiota. *Environmental Microbiology* vol. 19 29–41 Preprint at
https://doi.org/10.1111/1462-2920.13589 (2017).
72. Panicker, I. S., Browning, G. F. & Markham, P. F. The effect of an alternate
start codon on heterologous expression of a PhoA fusion protein in
mycoplasma gallisepticum. *PLoS One* **10**, 1–10 (2015).
73. Duncan, S. H., Barcenilla, A., Stewart, C. S., Pryde, S. E. & Flint, H. J. Acetate
utilization and butyryl coenzyme A (CoA): Acetate-CoA transferase in butyrate-
producing bacteria from the human large intestine. *Appl Environ Microbiol* **68**,
5186–5190 (2002).
74. Stackebrandt, E. Diversification and focusing: Strategies of microbial culture
collections. *Trends Microbiol* **18**, 283–287 (2010).
75. Afrizal, A. *et al.* Enhanced cultured diversity of the mouse gut microbiota
enables custom-made synthetic communities. *Cell Host Microbe* 1–16 (2022)
doi:10.1016/j.chom.2022.09.011.
76. Wylensek, D. *et al.* A collection of bacterial isolates from the pig intestine
reveals functional and taxonomic diversity. *Nat Commun* **in press**, 1–26
(2020).
77. Stackebrandt, E. *et al.* Deposit of microbial strains in public service collections
as part of the publication process to underpin good practice in science.
Springerplus **3**, 1–4 (2014).
78. Burz, S. D. *et al.* A Guide for Ex Vivo Handling and Storage of Stool Samples
Intended for Fecal Microbiota Transplantation. *Sci Rep* **9**, 1–16 (2019).
79. Afrizal, A. *et al.* Anaerobic single-cell dispensing facilitates the cultivation of
human gut bacteria. *Environ Microbiol* **00**, (2022).

80. Vieira, S. *et al.* Usitatibacter rugosus gen. nov., sp. nov. and Usitatibacter
palustris sp. nov., novel members of Usitatibacteraceae fam. nov. within the
order Nitrosomonadales isolated from soil. *Int J Syst Evol Microbiol* (2021).

81. Spitzer, V. Structure analysis of fatty acids by gas chromatography - Low
resolution electron impact mass spectrometry of their 4,4-dimethyloxazoline
derivatives - A review. *Prog Lipid Res* **35**, 387–408 (1996).

82. Godon, J. J., Zumstein, E., Dabert, P., Habouzit, F. & Moletta, R. Molecular
microbial diversity of an anaerobic digestor as determined by small-subunit
rDNA sequence analysis. *Appl Environ Microbiol* **63**, 2802–2813 (1997).

83. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for
Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

84. Bushnell, B. BBMap: A Fast, Accurate, Splice-Aware Aligner. in *9th Annual
Genomics of Energy & Environment Meeting* (2014).

85. Antipov, D. *et al.* PlasmidSPAdes: Assembling plasmids from whole genome
sequencing data. *Bioinformatics* **32**, 3380–3387 (2016).

86. Yilmaz, P. *et al.* Minimum information about a marker gene sequence
(MIMARKS) and minimum information about any (x) sequence (MIxS)
specifications. *Nat Biotechnol* **29**, 415–420 (2011).

87. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W.
CheckM: assessing the quality of microbial genomes recovered from. *Cold
Spring Harbor Laboratory Press Method* **1**, 1–31 (2015).

88. Mikheenko, A., Prjibelski, A., Saveliev, V., Antipov, D. & Gurevich, A. Versatile
genome assembly evaluation with QUAST-LG. *Bioinformatics* **34**, i142–i150
(2018).

89. Bengtsson-Palme, J. *et al.* metaxa2: Improved identification and taxonomic
classification of small and large subunit rRNA in metagenomic data. *Mol Ecol
Resour* **15**, 1403–1414 (2015).

90. Schwengers, O. *et al.* Bakta: Rapid and standardized annotation of bacterial
genomes via alignment-free sequence identification. *Microb Genom* **7**, (2021).

91. Mölder, F. *et al.* Sustainable data analysis with Snakemake [version 2; peer
review: 2 approved]. *F1000Res* **10**, 1–29 (2021).

92. Hitch, T. C. A. *et al.* Automated analysis of genomic sequences facilitates high-
throughput and comprehensive description of bacteria. *ISME Communications*
1, (2021).

93. Asnicar, F. *et al.* Precise phylogenetic analysis of microbial isolates and
genomes from metagenomes using PhyloPhlAn 3.0. *Nat Commun* **11**, 1–10
(2020).

94. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation
site identification. *BMC Bioinformatics* **11**, 119 (2010).

95. Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile
HMM and adaptive score threshold. *Bioinformatics* **36**, 2251–2252 (2020).

96. Nelson, K. E. *et al.* A catalog of reference genomes from the human
microbiome. *Science (1979)* **328**, 994–999 (2010).

97. Hitch, T. C. A. *et al.* Automated analysis of genomic sequences facilitates high-
throughput and comprehensive description of bacteria. *ISME Communications*
1, (2021).

98. Lagkouvardos, I. *et al.* IMNGS: A comprehensive open resource of processed
16S rRNA microbial profiles for ecology and diversity studies. *Sci Rep* **6**, 1–9
(2016).

99. Fodor, A. A. *et al.* The 'most wanted' taxa from the human microbiome for
whole genome sequencing. *PLoS One* **7**, (2012).
100. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local
alignment search tool. *J Mol Biol* **215**, 403–410 (1990).
101. Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S.
High throughput ANI analysis of 90K prokaryotic genomes reveals clear
species boundaries. *Nat Commun* **9**, 5114 (2018).
102. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of
biomolecular interaction networks. *Genome Res* **13**, 2498–2504 (2003).
103. Sullivan, M. J., Petty, N. K. & Beatson, S. A. Easyfig: A genome comparison
visualizer. *Bioinformatics* **27**, 1009–1010 (2011).
104. Beverley, S. M. Estimation of circular DNA size using γ-irradiation and pulsed-
field gel electrophoresis. *Anal Biochem* **177**, 110–114 (1989).
105. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold.
Nature **596**, 583–589 (2021).
106. Bittrich, S., Segura, J., Duarte, J. M., Burley, S. K. & Rose, Y. RCSB protein
Data Bank: exploring protein 3D similarities via comprehensive structural
alignments. *Bioinformatics* **40**, (2024).
107. Zhang, Y. & Skolnick, J. TM-align: A protein structure alignment algorithm
based on the TM-score. *Nucleic Acids Res* **33**, 2302–2309 (2005).
108. Yarza, P. *et al.* Uniting the classification of cultured and uncultured bacteria
and archaea using 16S rRNA gene sequences. *Nat Rev Microbiol* **12**, 635–645
(2014).
109. Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S.
High throughput ANI analysis of 90K prokaryotic genomes reveals clear
species boundaries. *Nat Commun* **9**, 5114 (2018).
110. Qin, Q. L. *et al.* A proposed genus boundary for the prokaryotes based on
genomic insights. *J Bacteriol* **196**, 2210–2215 (2014).
111. Meile, L., Rohr, L. M., Geissmann, T. A., Herensperger, M. & Teuber, M.
Characterization of the D-xylulose 5-phosphate/D-fructose 6-phosphate
phosphoketolase gene (xfp) from Bifidobacterium lactis. *J Bacteriol* **183**, 2929–
2936 (2001).
112. Mishra, A. K., Hugon, P., Robert, C., Raoult, D. & Fournier, P. E. Non
contiguous-finished genome sequence and description of Peptoniphilus
grossensis sp. nov. *Stand Genomic Sci* **7**, 320–330 (2012).
1855
1856

Supplementary Tables

Supplementary Table 1: List of existing isolate collections from the human gut. For each isolate collection we detail the accessibility of the published strains. Additional sheets provide specific information on strain collections

Supplementary Table 2: HPLC measurement of the production and/or utilisation of multiple metabolites.

Supplementary Table 3: Statistical comparison of genome sizes of phyla present within the HiBC, UHGG (isolates), and complete UHGG.

Supplementary Table 4: Statistical comparison of the normalised carbohydrate active enzyme investment of phyla within the HiBC.

Tree scale: 0.1

Phyla
- Bacteroidota
- Bacillota
- Desulfobacterota
- Fusobacteriota
- Pseudomonadota
- Actinomycetota
- Verrucomicrobiota
- **Novel taxa**

a

F. gallinarum JCM 17207ᵀ
F. prausnitzii F (GCF 002549775.1)
F. prausnitzii D (GCF 002549755.1)
F. taiwanense CLA-JM-H7B
F. prausnitzii M (GCA 004558805.1)
F. intestinale CLA-AA-H281ᵀ
F. prausnitzii J (GCF 003287405.1)
F. prausnitzii ATCC 27768ᵀ
F. prausnitzii A (GCF 002550015.1)
F. tardum CLA-AA-H175ᵀ
F. prausnitzii I (GCF 003287495.1)
F. prausnitzii L (GCA 002313795.1)
F. butyricigenerans DSM 103434ᵀ
F. prausnitzii G (GCF 003293635.1)
F. longum DSM 103432ᵀ
F. prausnitzii E (GCF 002550035.1)
F. prausnitzii H (GCF 002549975.1)
F. hattorii JCM 39210ᵀ
F. hominis DSM 113716ᵀ
F. duncaniae JCM 31915ᵀ
F. prausnitzii C (GCF 010509575.1)

Tree scale: 0.1

c

Acetyl-CoA —(Thl (thiolase) K00626)→ Acetoacetyl-CoA —(Hbd (β-hydroxybutyryl-CoA dehydrogenase) K00074)→ 3-Hydroxy butanoyl-CoA —(Cro (crotonase) K01692 / K01515)→ Crotonoyl-CoA —(Bcd (butyryl-CoA dehydrogenase) K00248)→ Butyryl-CoA —(But (butyryl-CoA:acetate CoA transferase) NA)→ Butyrate

d

Butyrate production (mM) 0 — 15
OD⁶⁰⁰ 0 — 1.5

F. butyricigenerans CLA-AA-H233
F. longum CLA-AA-H243
F. longum CLA-AA-H236
F. hominis CLA-AA-H283
F. hominis CLA-AA-H16
F. hominis CLA-AA-H223
F. taiwanense CLA-JM-H7B
F. intestinale CLA-AA-H281ᵀ
F. tardum CLA-AA-H175ᵀ
F. prausnitzii CLA-AA-H222
F. prausnitzii CLA-JM-H7

Tree scale: 0.1

e

Size: 2.5kb
Identity (%) 0 — 100
Cro (1) Cro (2) Bcd Hbd Thl But

h

Pearson correlation
r = 0.71
P = 0.01

Isolation

Description

Current strategies

	Quick publication	Curated (meta)data	Long-term accessibility	Traceability

→ 1. Publication

→ 2. Publication, then deposition*

→ 3. Parallelled publication & deposition*

→ 4. Deposition*, then publication

*Single strain deposition

1, Only original strain ID in publication
2, Strain may fail to be deposited
3, Deposition is time consuming

Proposed system

Curated (meta)data

1. Unique identifier (doi)

Pure Strains

2. BULK submission

Iterative strain processing