Title

Deficiency in a special dynein DNAH12 causes male infertility by impairing DNAH1 and DNALI1 recruitment in humans and mice

Authors

Menglei Yang[1], Hafiz Muhammad Jafar Hussain[1], Manan Khan[1], Zubair Muhammad[1], Jianteng Zhou[1], Ao Ma[1], Xiongheng Huang[1], Jingwei Ye[1], Min Chen[1], Aoran Zhi[1], Tao Liu[1], Ranjha Khan[1], Asim Ali[1], Wasim Shah[1], Aurang Zeb[1], Nisar Ahmad[1], Huan Zhang[1], Bo Xu[1], Hui Ma[1]*, Qinghua Shi[1]*, Baolu Shi[1]*

Affiliations

1. Center for Reproduction and Genetics, First Affiliated Hospital of USTC, Hefei National Research Center for Physical Sciences at the Microscale, the CAS Key Laboratory of Innate Immunity and Chronic Disease, School of Basic Medical Sciences, Division of Life Sciences and Medicine, Biomedical Sciences and Health Laboratory of Anhui Province, Institute of Health and Medicine, Hefei Comprehensive National Science Center, University of Science and Technology of China, Hefei 230027, China.

*Correspondence:

bl625@ustc.edu.cn (B.S.), qshi@ustc.edu.cn (Q.S.), clsmh@ustc.edu.cn (H.M.)

Abstract

Asthenoteratozoospermia, a prevalent cause of male infertility, lacks a well-defined etiology. DNAH12 is a special dynein featured by the absence of a microtubule-binding domain, however, its functions in spermatogenesis remain largely unknown. Through comprehensive genetic analyses involving whole-exome sequencing and subsequent Sanger sequencing on infertile patients and fertile controls from six distinct families, we unveiled six biallelic mutations in *DNAH12* that co-segregate recessively with male infertility in the studied families. Transmission electron microscopy (TEM) revealed pronounced axonemal abnormalities, including inner dynein arms (IDAs) impairment and central pair (CP) loss in sperm flagella of the patients. Mouse models (*Dnah12*$^{-/-}$ and *Dnah12*$^{mut/mut}$) were generated and recapitulated the reproductive defects in the patients. Noteworthy, DNAH12 deficiency did not show effects on cilium organization and function. Mechanistically, DNAH12 was confirmed to interact with two other IDA components DNALI1 and DNAH1, while disruption of DNAH12 leads to failed recruitment of DNALI1 and DNAH1 to IDAs and compromised sperm development. Furthermore, DNAH12 also interacts with radial spoke head proteins RSPH1, RSPH9, and DNAJB13 to regulate CP stability. Moreover, the infertility of *Dnah12*$^{-/-}$ mice could be overcome by intracytoplasmic sperm injection (ICSI) treatment. Collectively, DNAH12 plays a crucial role in the proper organization of axoneme in sperm flagella, but not cilia, by recruiting DNAH1 and DNALI1 in both humans and mice. These findings expand our comprehension of dynein component assembly in flagella and cilia and provide a valuable marker for genetic counseling and diagnosis of asthenoteratozoospermia in clinical practice.

Keywords

male infertility; DNAH12; knockout mice; inner dynein arms assembly; flagella and cilia; flagellar development

Introduction

Infertility has emerged as a global health concern, impacting around 8-12% of couples during their reproductive years, approximately 50% of infertility cases are attributed to male factors (1). Among these, asthenoteratozoospermia, characterized by poor sperm motility and notable morphological abnormalities, stands out as a prevalent contributor to male infertility (2, 3). Recent investigations have unveiled a multitude of gene mutations associated with asthenoteratozoospermia and male infertility, including Dynein Axonemal Heavy Chain (DNAH) inner dynein arm (IDA) genes like *DNAH1, DNAH2, DNAH6, DNAH7,* and *DNAH10*, as well as outer dynein arm (ODA) genes such as *DNAH8* and *DNAH17*, all implicated in primary male infertility with asthenoteratozoospermia (4-11). Additionally, genes encoding cilia and flagella-associated proteins (CFAP), including *CFAP43, CFAP44, CFAP57,* and *CFAP61*, have been identified as linked to asthenoteratozoospermia (12-15). Despite these advances, the intricate and heterogeneous features of asthenoteratozoospermia leave a significant number of cases unexplained. It is anticipated that additional genetic factors play a role in sperm anomalies and dysfunction.

The axoneme, a fundamental structure of cilia and flagella, exhibits a well-organized pattern with nine peripheral doublet microtubules (DMTs) and a central pair of microtubules (CP), constituting the classical "9+2" microtubular arrangement (16). In motile cilia or flagella, the radial spoke (RS) head contacts the CP accessory structures, transmitting mechanical and chemical signals from the central microtubules to the axonemal dynein arms, thus coordinating the flagellar or ciliary movement (17).

Within mammalian species, the DNAH family plays a crucial role in the axonemal structure,
encompassing eight IDA members (DNAH1-3, DNAH6, DNAH7, DNAH10, DNAH12, DNAH14) and
five ODA members (DNAH5, DNAH8, DNAH9, DNAH11, DNAH17). Despite the collaborative effort
of IDAs and ODAs in generating the force necessary for flagellar and ciliary motion, they serve distinct
functions. Chlamydomonas ODAs, composed of α, β, and γ chains, primarily regulate beat frequency,
while IDAs, with seven major subspecies (a-g), contribute to bending formation and beating form (18).
Mutations in genes encoding ODA or IDA components have been linked to asthenoteratozoospermia,
with or without symptoms of primary ciliary dyskinesia (PCD), a genetic disorder characterized by
chronic airway infections or situs inversus (19). Notably, deficiency in certain IDAs and ODAs coding
genes, such as *DNAH7* or *DNAH11*, may lead to PCD (20-22). Patients carrying mutations in *DNAH1*
and *DNAH2* typically lack PCD symptoms (4, 23), although a study suggested the potential contribution
of the DNAH1 p.Lys1154Gln mutation to PCD (24). The extensive genetic heterogeneity and clinical
variabilities underscore the intricacy and uniqueness of dynein arms in flagella and cilia. However, the
susceptibility of uncharacterized dynein factors that related to asthenoteratozoospermia and/or PCD
remains unclear, given the absence of human genetic evidence or corresponding animal models.

Dynein axonemal heavy chain 12, DNAH12 (MIM: 603340, GenBank: NM_001366028.2), also
known as DHC3, is a unique dynein arm component (25). Notably, it is the shortest member and lacks a
typical microtubule-binding domain (MTBD) in the DNAH family. Mutations of *DNAH12* have been
predicted to be associated with male patients' infertility, however, limited pathogenic evidence was
provided, and the underlying pathogenesis remained unclear (26, 27). Besides, although DNAH12 has
been identified as a possible axonemal dynein and its mRNA is predominantly detected in two tissues
containing axonemal structures: trachea and testis (25, 28), its physiological functions remain largely
elusive.

In this study, we enrolled 796 infertile patients from unrelated Pakistani and Chinese families
diagnosed with asthenozoospermia. Utilizing whole-exome sequencing (WES) followed by Sanger
sequencing, we identified six novel bi-allelic mutations in *DNAH12* in infertile males from six unrelated
families. All these mutations resulted in the loss of DNAH12 protein in the sperm of patients. Two mouse
models, *Dnah12*$^{-/-}$ and *Dnah12*$^{mut/mut}$, were generated and exhibited asthenoteratozoospermia and male
infertility, mimicking our patients' phenotypes. Deficiency in DNAH12 led to axoneme defects
characterized by impaired IDAs and CP loss in sperm flagella, while no such defects were observed in
cilia. DNAH12 interacts with other IDA components like DNALI1 and DNAH1 in mouse testes, and
disruption of DNAH12 impaired the recruitment of dynein components into sperm flagella. Our findings
unveil, for the first time, that DNAH12 plays a pivotal role in facilitating the recruitment of IDA
components specifically in flagella, but not in cilia. Furthermore, we demonstrate that bi-allelic mutations
in *DNAH12* are identified as causative factors for asthenoteratozoospermia and male infertility in both
humans and mice. These results enhance our understanding of the development of axonemal structures
in flagella versus cilia and provide valuable clues for genetic counseling and future individualized
treatment for infertile males carrying *DNAH12* mutations in clinical practice.

Materials and methods
Human subjects
Informed consent and peripheral blood were obtained from the participants and the patients' information

103 was registered in the Human Reproductive Disease Resource Bank database. Sperm analyses were
104 performed according to the WHO guidelines (29). During their medical consultation, all patients
105 answered a questionnaire regarding clinical presentations. This study was approved by the institutional
106 ethics committee of University of Science and Technology of China (USTC) with the approval number
107 2019-KY-168.

108 **Whole exome sequencing (WES) and bioinformatics analyses**

109 Whole-exome capture and sequencing were performed by capturing exons from genomic DNAs of
110 peripheral blood cells, using the SureSelect XT 50 Mb Exon Capture Kit (Agilent, California, USA), and
111 then high-throughput sequencing was conducted on the HiSeq 2000 platform (Illumina, San Diego, CA,
112 USA) following instructions. Variants were called using the Genome Analysis Toolkit HaplotypeCaller
113 and annotated through ANNOVAR (30). Variants within exons or exon-intron boundaries were retained
114 and filtered through the criteria we previously described (31). Bcftools was applied to detect runs of
115 homozygosity (32). Runs of homozygosity over 1.5 Mb were used to calculate the inbreeding coefficients.
116 The genotypes were validated using Sanger sequencing validation. Additional primers are listed in
117 Supplemental Table 1. For phylogenetic analysis, different DNAH12 protein sequences were extracted
118 from the UniProt database, and the sequence alignment and phylogenetic tree construction were
119 generated using MEGA11 (33). Newick format tree files were extracted and processed in the Evolview
120 v2 tool (34). For gene family and conserved domain analyses, the batch sequence search tool in Pfam
121 was used to analyze features of proteins and further visualized with the assistance of Evolview v2 tool
122 and TBtools (35, 36). For protein structure analysis, the structure models were predicted and constructed
123 with the Phyre2 tool (37), interactions were analyzed by the HDOCK (http://hdock.phys.hust.edu.cn/)
124 and Alphafold3 (https://golgi.sandbox.google.com/). The predicted protein structures and possible
125 interactions were further explored and visualized by UCSF Chimera (Version: 1.15, California, USA)
126 according to the manual instruction (38).

127 **Scanning electron microscopy (SEM) and Transmission electron microscopy (TEM)**

128 Fresh semen samples were obtained and centrifuged at 900 g for 3 min. Spermatozoa were washed with
129 phosphate buffer saline (PBS) three times. Then the sediment was fixed in 0.1 M phosphate buffer (pH
130 7.4) containing 4% paraformaldehyde, 8% glutaraldehyde, and 0.2% picric acid at 4°C for at least 8
131 hours. Scanning electron microscopy (SEM) and Transmission electron microscopy (TEM) were then
132 conducted as we previously described (39).

133 **Generation and verification of the polyclonal anti-DNAH12 antibody**

134 The characteristics of the DNAH12 were analyzed and the 1-200 amino acids sequence of mouse
135 DNAH12 protein (UniProt accession no. Q69Z23) was regarded as the ideal antigen for immunization.
136 Subsequently, the DNAH12 polyclonal antibody was generated in rabbits and rats using the mentioned
137 peptide by Dia-An Biotech, Inc., in Wuhan, China. The specificity of antibodies was further validated
138 using $Dnah12^{+/+}$ and $Dnah12^{-/-}$ mice in this study.

139 **Immunoblotting**

140 Protein lysate was prepared as we previously described (40, 41). Tissues or cells were lysed using lysis
141 buffer (50 mM tris (pH 7.5), 150 mM NaCl, 5 mM EDTA, and 0.1% NP-40). For human sperm cells,
142 proteins were extracted using the TRIzol reagent. The protein sediment was washed with 95% ethanol
143 containing 0.3M guanidine hydrochloride, dissolved in 1% SDS. Then the protein lysate was boiled for
144 10 minutes, and diluted in SDS-sample buffer (120 mM Tris/HCl, 10% SDS, 20% glycerol, 20% 2-
145 mercaptoethanol, and bromophenol blue (pH 6.8). Whereafter the proteins were separated by SDS-PAGE
146 and transferred onto nitrocellulose blotting membranes with a pore size of 0.45 μm (GE Healthcare,

10600002). The membranes were blocked with TBS-Tween buffer (50 mM Tris, 150 mM NaCl, and 0.5% Tween-20, pH 7.4) containing 5% skim milk for 30 minutes to reduce unspecific binding and then incubated with primary antibodies diluted in TBST buffer containing 5% skim milk at 4°C overnight. Incubation with secondary antibodies for 1.5 hours was followed by chemiluminescence development system (GE Healthcare, Pittsburgh, USA). Antibodies details are listed in Supplemental Table 2-3.

Immunofluorescence (IF) staining

Human semen smear slides were prepared following the guidelines of WHO (29). Mouse sperm smear slide preparation and immunofluorescence staining were conducted as we described (42). For tissue IF staining, tissues were collected and fixed in 4% paraformaldehyde (PFA), the 5μm tissue sections were deparaffinized in xylene, rehydrated in a graded series of ethanol (100%, 95%, 90%, 80%, 70%, and 50% ethanol) and sterile water, and permeabilized with 0.2% Triton X-100 in PBS for 30min. Antigen retrieval was then conducted by boiling slides in a microwave for 30 minutes in 10 mM citrate buffer (pH 6.0). The other steps are consistent with sperm IF staining. Images of spermatozoa were captured using a Nikon ECLIPSE 80i microscope equipped with a charge-coupled device (Hamamatsu Photonics, Hamamatsu, Japan). Information of antibodies is available in Supplemental Table 2-3.

Animal models and fertility test

Two mouse models ($Dnah12^{-/-}$ and $Dnah12^{mut/mut}$) were generated using CRISPR/Cas9 Technology (43). Briefly, to generate the $Dnah12^{-/-}$ mouse model, guide RNAs (gRNAs) targeting exon 5 of $Dnah12$, a key exon shared by all its transcripts, were transcribed in vitro (Thermo Fisher, CA, USA). Then, the gRNAs and the Cas9 protein were electroporated into C57BL/6 mouse zygotes. The zygotes were introduced into ICR female mice, and the genotypes of the pups were checked using Polymerase Chain Reaction (PCR) with designed primers. Similar conductions were performed targeting exon 18 of $Dnah12$ to generate the $Dnah12^{mut/mut}$ mice. The genotyping primers are listed in Supplemental Table 1. For male fertility test, each 8-10 weeks male mouse was caged with wild-type female mice; for female fertility test, each 8-10 weeks female mouse was caged with one 8-10 weeks $Dnah12^{+/-}$ male mouse, the number of offspring was recorded.

Reverse-transcription PCR and real-time quantitative PCR

For reverse-transcription PCR (RT-PCR), total RNA was extracted from mouse tissues using TRIzol reagent (TAKARA, Japan). Whereafter 1μg of RNA was reverse-transcribed using the PrimeScript RT Reagent Kit (TAKARA, Japan). The cDNA was diluted to a comparable concentration and then used for RT–PCR. All amplified DNA fragments were visualized by 2% agarose gel electrophoresis containing GelRed (APExBIO, Houston, TX, USA). For real-time quantitative PCR (qPCR), the experiments were performed by using Hieff UNICON® Universal Blue qPCR SYBR Green Master Mix reagent (Yeasen Biotechnology, Shanghai, China) on CFX96® Real-Time System (Bio-Rad, CA, USA) according to the manufacturer's guidelines. Details of the primers were listed in Supplemental Table 1.

Histological analysis

As previously described (31), after sacrificing mice, the testes, epididymides, tracheas, and oviducts were harvested and fixed in Bouin's fixative solution (Sigma–Aldrich, Missouri, USA) for hematoxylin and eosin (H&E) staining or 4% PFA fixative solution for immunofluorescence or TUNEL assay. The pictures were captured using a microscope equipped with a charge-coupled device camera (Nikon, Tokyo, Japan).

Sperm counts, motility and morphology analysis

As previously described (44), H&E staining or Papanicolaou staining kit (Solabio, Beijing, China) was used for the evaluation of sperm morphology of patients. For mouse sperm parameters analysis, 8-10-week-old mice were sacrificed by cervical dislocation, the sperm number per epididymis was then

assessed by cutting the epididymis into pieces in PBS and incubating them for 30 minutes at 37°C to

release the sperm, then counting with a hemocytometer. Sperm motility were assessed by computer-aided

sperm analysis (CASA) as we described (14). Sperm morphology was assessed on sperms collected from

epididymal cauda after H&E staining.

Testicular cell smear slides and seminiferous tubule squash slides preparation

To prepare testicular cell smears, mice were sacrificed and testes were then detached with one time of

PBS washing. Then testes were peeled off the testicular capsule, torn into small pieces, and then cut into

tiny pieces in PBS. Finally, the cell suspension was filtered through a 200-mesh cell strainer, the filtered

cell suspensions were added 10% FBS before spread onto glass slides. For testicular seminiferous tubule

squash slide preparation, the testes were dissected and carefully isolated into single testicular

seminiferous tubules in PBS dishes. After 10 mins of prefixation in fix/lysis solution (2% PFA, 50%

PBS), we then applied 100μL fix/lysis solution onto poly-L-lysine coated glass slides and carefully teased

apart approximately 20 seminiferous tubules with the assistance of 2 sterile forceps on the slides. After

that, coverslips were applied using slide forceps and pressed with the heel of the palm for 10 seconds,

then the glass slides were frozen in a small container with liquid N_2 immediately till the liquid ceases to

bubble. The slides were saved in the -80°C fridge keeping the coverslip on the slides or removing

coverslips for immunofluorescence staining.

TUNEL assay

As previously described (44), In Situ Cell Death Detection Kit, Fluorescein (Sigma–Aldrich, Missouri,

USA) was applied according to the manufacturer's protocol. Images were captured using a Nikon Eclipse

80i microscope equipped with Hamamatsu C4742-80 digital camera and analyzed using the NIS-Element

Microscope imaging software (Nikon, Tokyo, Japan).

Cell culture and transfection.

As previously described (40), HEK293T (ATCC, VA, USA) cells were cultured to 70%-80% density in

12-well plates, followed by transfected with plasmids expressing mCherry-tagged mouse DNAH12 and

GFP-tagged mouse DNALI1 using Lipofectamine 3000 (Thermo Fisher, CA, USA) according to the

manufacturer's instructions. Cells were harvested for in-vitro IP and immunoblotting the next day after

transfection. The sequences of the primers for plasmid construction are listed in Supplemental Table 4.

Co-immunoprecipitation (Co-IP)

Tissue extracts were prepared using lysis buffer supplemented with PMSF, Protease Inhibitor Cocktail

(Solarbio, Beijing, China) on ice using a glass homogenizer. Lysate was then centrifuged at 12,000 g at

4°C for 10 min, and the supernatant was divided into equal aliquots. Each aliquot was incubated with 1.5

μg of target antibody or rabbit IgG nonspecific antibody. After incubation at 4°C overnight with rotation,

beads were washed five times with lysis buffer. Finally, the beads were resuspended in SDS-sample

buffer and boiled at 100°C for 10 min. The samples were either analyzed by immunoblotting immediately

or stored at -80°C. For Co-IP in cultured cells, HEK293T cells were plated in six-well plates, followed

by transfection using Lipofectamine 3000 reagents. MG132 was added into the cultured cells 6 hours

before cell harvesting. Cells were washed with PBS for 10 min, cells were then collected and lysed in

the same lysis buffer used for Co-IP from testicular samples. The cell lysate was then incubated with

prewashed Magnetic Beads (Bimake, Texas, USA) and primary antibodies at 4°C overnight, followed by

washing and boiling as described above for the Co-IP protocols for testicular samples.

Mass spectrometry proteomics

To identify potential partners that interact with DNAH12, we performed in vivo IP of mouse testes lysate.

Tissue extracts were prepared using lysis buffer supplemented with PMSF, Protease Inhibitor Cocktail

(Solarbio, Beijing, China) with the same operation of the Co-IP protocol. Beads were washed 3 times with lysis buffer and 3 times with 50 mM NH_4HCO_3, the pull-down proteins were digested with 0.2μg Trypsin overnight and stopped by 5% Formic acid, (Thermo Fisher, CA, USA). The peptide mixture was then desalted, dried with a SpeedVac, and resuspended in 0.1% Formic acid, followed by the MS analysis using timsTOF Pro 2 mass spectrometer (Bruker, Massachusetts, USA). The proteomics data are available on the ProteomeXchange Consortium via the iProX partner repository (45), with the dataset identifier PXD051681.

Intracytoplasmic sperm injection (ICSI)

3.5-4-week-old female mice were given an intraperitoneal injection of 5 IU Pregnant Mare Serum Gonadotropin (PMSG, Ningbo Sansheng, China). After 48 hours, a second intraperitoneal injection of 5 IU human Chorionic Gonadotropin (hCG, Ningbo Sansheng, China) was administered. The mice were euthanized by cervical dislocation after 15 hours post-hCG injection. The oviducts were excised and the ampullary regions were opened to release the cumulus-oocyte complexes (COCs). The cumulus cells surrounding the oocytes were digested using hyaluronidase, and the morphologically round MII stage oocytes were selected and washed three times in M2 medium (Sigma–Aldrich, Missouri, USA) before being set aside for use. Male mice are euthanized by cervical dislocation, the epididymides are excised and gently punctured with a syringe, allowing the spermatozoa to flow out. The spermatozoa were then placed in the human tubal fluid (HTF) medium (EasyCheck, Nanjing Aibei Biotechnology, China) and then frozen and thawed repeatedly to remove sperm tails, the single sperm head was microinjected into an MII oocyte and then cultured in potassium simplex optimized medium (KSOM, EasyCheck, Nanjing Aibei Biotechnology, China) at 37 °C under 5% CO_2.

Statistics analysis

Statistical analyses were performed using GraphPad Prism 8.0 (GraphPad Software, San Diego, CA, USA). Values in tables and graphs are expressed as mean ± SEM. Two groups of mice were compared with Student's 2-tailed t-test for independent data and $P > 0.05$ was considered not significant (ns); *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$ were considered significant.

Results

Identification of biallelic *DNAH12* variants in men with asthenoteratozoospermia

To identify the genetic factors contributing to male infertility, we recruited patients with primary infertility and their family members from Pakistan and China. WES was performed on these individuals, followed by comprehensive analyses of the WES data (Supplemental Figure 1, A-D). Seven mutations in *DNAH12* were identified as potential pathogenic factors for male infertility in patients P1-3 of family 1, P4-5 of family 2, P8 of family 3, and P9, P10, and P11, all of whom suffered from asthenoteratozoospermia (Fig. 1A). Initially, a homozygous frameshift mutation c.944_945 del (p. Phe315*) in family 1, and two homozygous missense mutations, c. A164G (p. Gln55Arg) in family 2 and c. A2560G (p. Ile854Val) in family 3, were identified in *DNAH12* as potential causes of infertility. Subsequently, four other mutations in *DNAH12* were discovered in three sporadic Chinese patients with asthenoteratozoospermia: a homozygous stop-gain mutation c.2680 C>T (p. Gln2077*) in subject P9, a homozygous missense mutation c.2813T>C (p. Val938Ala) in P10, and compound heterozygous mutations c.2680C>T (p. Arg894Cys) and c.5964-3A>G in P11 (Fig. 1A and B).

All seven variants were rare in public human genome databases, including the 1000 Genome Project and gnomAD (v3). Additionally, these variants were predicted to be deleterious by in-silico software

tools such as SIFT, Polyphen2, fathmm-MKL, and CADD2 (Table 1). Sanger Sequencing further
validated these variants and revealed the recessive pathogenic pattern of these mutations (Fig. 1A).
Collectively, these results suggested that the identified bi-allelic *DNAH12* variants could potentially be
the cause of asthenoteratozoospermia in our patients.

DNAH12 protein is conserved in mammals and highly expressed in the testes
DNAH12 encodes a functionally uncharacterized protein and is predicted to be abundantly expressed in
testes of humans and mice based on our bioinformatics tool FertilityOnline (46). Experimentally, our RT-
PCR, qPCR, and immunoblotting detection of DNAH12 expression in different tissues indicated that
both DNAH12 mRNA and protein showed high expression in testes and relatively lower expression in
lungs and tracheas of male mice and oviducts of female mice (Supplemental Figure 2, A-C). It's worth
noting that in epididymides, DNAH12 protein, but not mRNA, was detected, indicating the DNAH12
protein detected in epididymides is from testes, most probably from sperm. Furthermore, DNAH12
protein was initially detected in testes around 21 days postpartum (dpp) and thereafter maintained a high
level (Supplemental Figure 2D). The expression pattern of DNAH12 in testes suggests that it is possibly
involved in spermiogenesis, and its abnormal expression could lead to asthenozoospermia or
asthenoteratozoospermia, further indicating that the identified *DNAH12* mutations are candidate
pathogenic mutations for patients.

Further phylogenetic analysis revealed the conservative nature of DNAH12 from Chlamydomonas
to humans (Supplemental Figure 3). All the amino acids affected by mutations were conserved in
mammals and were located in functional regions, including the ATPase domains or the Stem domain of
DNAH12 (Fig.1C), indicating that these mutations are highly likely to hinder the structure, function, or
stability of the DNAH12 protein. Altogether, our results suggested that the identified mutations in
DNAH12 are probable pathogenic factors contributing to asthenoteratozoospermia in our patients.

Impaired sperm motility and morphology in patients with bi-allelic *DNAH12* variants
To assess the pathological effects of the identified bi-allelic *DNAH12* variants, semen analyses were
conducted following WHO guidelines (29). All affected individuals exhibited significantly reduced
sperm motility and progressive motility (Table 2). Detailed sperm morphology analyses were performed
on patients P3 from family 1, P5 from family 2, P8 from family 3, and three Chinese sporadic patients
(P9, P10, and P11). Spermatozoa from fertile controls displayed normal heads and tails, while
spermatozoa from the patients exhibited a range of abnormal morphologies, including short, coiled,
absent, bent, or irregular caliber flagella, accompanied by aberrant heads (Fig. 2A, Supplemental Figure
4, A and B). Scanning electron microscopy (SEM) analyses were further carried out on a fertile control
and P10, confirming the presence of abnormal morphologies of the sperm head and flagellum in the
patients (Fig. 2B). Thus, patients with the bi-allelic *DNAH12* variants demonstrated significantly
decreased sperm motility and increased abnormal sperm morphology.

The bi-allelic *DNAH12* variants result in absence of DNAH12 protein in patients' sperm
To investigate the effects of the identified bi-allelic *DNAH12* variants on the DNAH12 protein, we
obtained sperm samples from three affected Chinese patients, and the expression of DNAH12 was
detected using a custom-made DNAH12 rabbit antibody that recognized the N-terminal 1-200 amino
acids of the DNAH12 protein. This validated antibody (Supplemental Figure 6B) could theoretically

detect the truncated protein if it were present in P9 with a homozygous stop-gain mutation M4 (p. Gln2077*). The predicted DNAH12 protein size was detected in the sperm lysate from a fertile control but not in P9 with the homozygous stop-gain mutation M4, P10 with a homozygous missense mutation M5, and P11 with compound heterozygous mutations M6 and M7, indicating these mutations possibly lead to the disruption of the DNAH12 protein (Fig. 2C).

To further explore the effects of the identified *DNAH12* variants on the DNAH12 protein, we performed immunofluorescence assays on sperm from patients. The results showed that DNAH12 signals were localized along the flagella of the fertile control, however, no DNAH12 signals were observed in sperm flagella of patients P3 (family 1), P5 (family 2), P8 (family 3), P9, P10, and P11 (Fig. 2D). Collectively, these findings indicated that all the detected *DNAH12* variants cause the absence of DNAH12 protein in the sperm of our patients, reinforcing that these *DNAH12* mutations are pathogenic mutations of asthenoteratozoospermia in the patients.

Patients with bi-allelic *DNAH12* variants exhibit flagellar axoneme defects

Since patients with the bi-allelic *DNAH12* variants exhibited compromised sperm motility and abnormal sperm morphology, we then investigated whether sperm flagellar ultrastructure was also affected in these patients. TEM examination was performed on sperm flagella from both patients and normal fertile controls. The flagella of fertile controls displayed the typical "9+2" microtubule structure with well-organized DMTs and CP. In contrast, various defects were found in our patients, including loss of CP (54.5%), disorganization of microtubules (26%), and absence of DMTs (16.5%) in the examined cross-sections (Fig. 3A, 3C and Supplemental Figure 5A). To examine the IDAs impairment in axonemes, the sections with intact ODAs were analyzed. Compared with axonemes in controls that almost all the sections presented intact IDAs, approximately 70% of sections examined showed impaired IDA structures in the analyzed three patients (Fig. 3B, 3D and Supplemental Figure 5A).

We then performed immunofluorescence assays on sperm smears to assess the expression and localization of specific proteins of the axonemes. These included SPAG6 (a CP component), DNAH1 and DNALI1 (IDA components), DNAH17 (an ODA component), and ODF2 (an outer dense fiber, ODF component). The sperm flagella of patients showed remarkably reduced SPAG6 signals when compared with those from controls (Fig. 3E). Similarly, DNAH1 and DNALI1 signals were both obviously decreased in the patients' sperm flagella, indicating impaired IDAs in the patients (Fig. 3F and G). Besides, DNAH17 and ODF2 signals in patients' flagella were comparable to those in a normal control (Fig. 3H and Supplemental Figure 5B). In summary, these findings indicated that the bi-allelic *DNAH12* variants cause pronounced flagellar axoneme defects, including impaired IDAs and loss of CP, but do not affect ODAs and ODFs.

Deletion of *Dnah12* impairs spermiogenesis and causes male infertility in mice

Considering the bi-allelic *DNAH12* variants resulting in absence of DNAH12 protein in patients' sperm, to functionally confirm the deletion of *DNAH12*, we generated *Dnah12* knockout mice using CRISPR-Cas9 approach (Supplemental Figure 6A). Fortunately, we obtained a mouse model with a 4-bp deletion (c.386_389del) in exon 5 of *Dnah12* (Supplemental Figure 6A). In contrast to the observations in control mice, immunoblotting of testis lysate using two antibodies targeting the N-terminal of DNAH12 failed to detect DNAH12 proteins, and immunofluorescence staining of sperm flagella did not show DNAH12

signals in *Dnah12⁻ᐟ⁻* mice (Supplemental Figure 6, B and C). *Dnah12⁻ᐟ⁻* males exhibited normal development but were infertile (Fig. 4A), while *Dnah12⁻ᐟ⁻* female mice showed normal development and fertility, and no obvious defective oviductal and tracheal cilia (Supplemental Figure 7, A-C). The body weight of *Dnah12⁻ᐟ⁻* and *Dnah12⁺ᐟ⁺* males was comparable, while smaller testes were observed in *Dnah12⁻ᐟ⁻* males (Fig. 4B). Besides, the average number of sperm per epididymis of *Dnah12⁻ᐟ⁻* mice dramatically decreased to approximately 2% of that of *Dnah12⁺ᐟ⁺* mice (Fig. 4C). The epididymal caudal spermatozoa of *Dnah12⁻ᐟ⁻* male mice were 100% immotile, compared to those of *Dnah12⁺ᐟ⁺* mice (Fig. 4D, Supplemental Videos 1 and 2). H&E staining of testis and epididymis sections also showed a striking reduction in the number of sperm in *Dnah12⁻ᐟ⁻* mice (Fig. 4E). Importantly, morphological analysis of sperm from the caudal epididymis of *Dnah12⁻ᐟ⁻* mice showed similar types and frequencies of flagellum and head abnormalities to those observed in our patients (Fig. 4F and G).

To understand how *DNAH12* depletion causes abnormal sperm morphology, we conducted immunofluorescence assays on testicular cells from *Dnah12⁺ᐟ⁺* and *Dnah12⁻ᐟ⁻* mice, in *Dnah12⁻ᐟ⁻* mice, while DNAH12 signals are missing in round, elongating, and elongated spermatids, the abnormal manchette structure and flagellum were observed (Fig. 4H). We then performed immunofluorescence assays on testicular sections using α-Tubulin and lectin peanut agglutinin (PNA, a marker of the acrosome) to examine the process of spermiogenesis. The manchette structure in *Dnah12⁻ᐟ⁻* was roughly normal until steps 9-10 (Stage IX-X) but was aberrantly stretched from steps 11-12 (Stage XI-XII) when compared with those of *Dnah12⁺ᐟ⁺* mice (Supplemental Figure 8). To meticulously characterize these defects, we stained α-Tubulin and PNA on squashed seminiferous tubules and found obvious manchette defects and abnormal sperm heads in *Dnah12⁻ᐟ⁻* mice (Fig. 4I). TEM analysis showed severely abnormal head morphology in elongated spermatids of *Dnah12⁻ᐟ⁻* males when compared with those of *Dnah12⁺ᐟ⁺* males (Fig. 4J). Additionally, TUNEL assays of testis sections showed dramatically increased proportions of seminiferous tubules containing TUNEL-positive cells in *Dnah12⁻ᐟ⁻* mice, compared with those in *Dnah12⁺ᐟ⁺* mice, suggesting the aberrant spermatids may undergo apoptosis (Figure 4K-M). Altogether, these results indicated that deletion of *Dnah12* impairs spermiogenesis, results in decreased sperm counts, and increased sperm morphological abnormalities in mice.

Deletion of *Dnah12* causes axoneme defects in sperm flagella but not cilia in mice

TEM analysis was conducted on sperm from cauda epididymis of *Dnah12⁺ᐟ⁺*, *Dnah12⁺ᐟ⁻* and *Dnah12⁻ᐟ⁻* mice. Flagellar axonemes of *Dnah12⁺ᐟ⁺* and *Dnah12⁺ᐟ⁻* mice displayed a typical "9+2" microtubular arrangement with clearly visible CP, IDAs, and ODAs. In contrast, sperm from *Dnah12⁻ᐟ⁻* mice exhibited severe abnormalities in all the analyzed sections, characterized by CP loss and disorganized axoneme structure (Fig. 5A). Immunoblotting was performed to assess the expression levels of DNAH1, DNALI1, and SPAG6 in sperm lysates, *Dnah12⁻ᐟ⁻* sperm showed a significant reduction in the abundance of DNAH1 and DNALI1, while the amount of DNAI2 (a component of ODAs) remained comparable to *Dnah12⁺ᐟ⁺* and *Dnah12⁺ᐟ⁻* sperm (Fig. 5B). Additionally, immunofluorescence assays revealed the near absence of signals for SPAG6, SPEF2 (a component of CP complex), DNAH1, and DNALI1 in *Dnah12⁻ᐟ⁻* sperm flagella, whereas DNAH17 and DNAI2 (components of ODAs) remained unaffected (Fig. 5C-H). These results indicate that DNAH12 deficiency specifically impairs CP and IDAs, but not ODAs, in sperm flagella. Furthermore, immunoblotting of SPAG6 and DNALI1 in testis lysate exhibited a sharp decrease in *Dnah12⁻ᐟ⁻* mice (Supplemental Figure 9A). Additionally, TEM examination of testicular sperm axonemes showed defects like loss of CP and disarranged axonemes in *Dnah12⁻ᐟ⁻* mice

(Supplemental Figure 9B). These results indicate that impaired IDAs and loss of CP possibly occur during testicular sperm flagellar development.

Another *Dnah12* mutant mouse model (*Dnah12^{mut/mut}*) targeting exon18 was also generated (Supplemental Figure 10A), which may produce a truncated DNAH12 protein (p.Cys793Glu, fs*10). The mutation effect was checked but no truncated DNAH12 protein was detected, indicating that DNAH12 expression was disrupted in *Dnah12^{mut/mut}* mice (Supplemental Figure 10B). Moreover, *Dnah12^{mut/mut}* mice developed normally, while decreased testicular size was observed (Supplemental Figure 10, C and D). Consistent with *Dnah12^{-/-}* mice and patients carrying bi-allelic *DNAH12* mutations, *Dnah12^{mut/mut}* presented obviously decreased sperm count, increased sperm morphological abnormalities, and sperm axoneme defects similar to *Dnah12^{-/-}* mice (Supplemental Figure 10, E-H). Therefore, these results reconfirmed that DNAH12 is essential for sperm flagellar development.

Given the detection of DNAH12 protein in the trachea and lung (Supplemental Figure 2, A-C), we also examined whether the deficiency of DNAH12 would cause PCD symptoms in males. Intriguingly, both *Dnah12^{mut/mut}* and *Dnah12^{-/-}* males developed normally and showed no tracheal defects, despite the expression of DNAH12 being detected in tracheal cilia cells (Supplemental Figure 11, A-E). These findings collectively suggested that DNAH12 deficiency is crucial for sperm axonemal integrity in humans and mice without manifesting PCD symptoms.

DNAH12 interacts with IDA components DNAH1 and DNALI1

Through phylogenetic and domain analyses of DNAH family members, we found that DNAH12 is a distinctive IDA lacking the microtubule-binding domain (MTBD), which is crucial for ATP-sensitive microtubule binding and motor direction control (Fig.6A, Supplemental Figure 12). Therefore, DNAH12 likely collaborates with other members to carry out its functions. To delve into the molecular mechanism of DNAH12, we conducted co-immunoprecipitation (Co-IP) followed by mass spectrometry analysis using a DNAH12 rabbit antibody in testicular lysate from adult *Dnah12^{+/+}* mice (Fig. 6B). Gene ontology analysis showed that the potential interactors were mainly enriched in motor protein, actin binding, myosin and RNA binding proteins (Fig. 6C). The motor proteins were predominantly categorized into three groups: dynein, kinesin, and myosin protein members, interestingly, among these, dynein proteins such as DNAH1, DNALI1, DNAH1, DNAH3, and DNAH2 were annotated to be involved in IDA assembly, which is closely related to the defects found in *DNAH12* mutants (Fig. 6D). This piqued our interest to explore these proteins in detail. Besides the predominant enrichment of DNAH12 (top1), potential IDA interactors such as DNALI1 (top2), DNAH1 (top3), DNAH3 (top4), and DNAH2 (top5) were co-enriched in the candidate interacting dynein protein list (Supplemental Table 5). Immunoblotting confirmed the co-enrichment of DNAH12, DNALI1, and DNAH1 in the immunoprecipitated complex by the DNAH12 rabbit antibody in *Dnah12^{+/+}* mice but not in *Dnah12^{-/-}* mice, indicating specific interactions between DNAH12 and DNALI1, DNAH1 (Fig. 6E). Reciprocal in vivo Co-IP using an anti-DNALI1 antibody further validated strong interactions among these three IDA components (Fig. 6F).

Given the potential role of DNALI1 as an IDA assembly factor (47), we delved into the interaction details between DNAH12 and DNALI1. Employing Phyre2, we constructed the structural models of both proteins and further elucidated their interaction dynamics via DNAH12-DNALI1 docking using the HDOCK server. The results showed that the N-terminal Stem domain of DNAH12 possibly interacts

with DNALI1 in the top three best-fit conformations (Fig. 6G). Considering DNAH12 is encoded by 3092 amino acids and their full-length protein interactions have been proved in vivo, we constructed the Stem domain (1-1212 amino acids) of DNAH12 and full-length DNALI1 plasmid, further in vitro Co-IP assays validated that the Stem domain of DNAH12 indeed mediates interactions with DNALI1 (Fig. 6H-J). Together, these findings proved the specific interactions among DNAH12, DNALI1, and DNAH1. Given the significantly decreased protein levels of DNALI1 and DNAH1 observed in the sperm of *Dnah12*-/-mice (Fig.5B), we proposed that the interactions involving DNAH12 (possibly restricted to the Stem domain of DNAH12), DNALI1, and DNAH1 are crucial for flagella development. As dynein proteins typically share similar domains, our findings also suggested that the Stem domain of DNAH family members may facilitate the interaction with DNALI1.

Deletion of DNAH12 impairs the recruitment of IDA components during flagella development

To evaluate the effect of DNAH12 deletion on DNALI1 and DNAH1 during spermiogenesis, we examined the abundance of the DNAH12 interaction partners, DNALI1 and DNAH1, in *Dnah12*+/+ and *Dnah12*-/- mouse testis lysate. The protein levels of DNALI1 and DNAH1 were decreased in *Dnah12*-/- testes, while the level of ODA member DNAI1 was comparable (Fig. 7A). We then stained DNALI1 or DNAH1 on testicular cell smear of *Dnah12*+/+ and *Dnah12*-/- mice. Although the signals of DNALI1 or DNAH1 in round spermatids of *Dnah12*-/- mice were comparable to those of *Dnah12*+/+ mice (Fig. 7B and 7C), the recruitment of DNALI1 and DNAH1 to sperm flagella was severely impaired in elongated spermatids of *Dnah12*-/- mice (Fig.7D and 7E). Considering that the deficiency of DNAH12 also caused scarsely visible immunofluorescence signals of DNALI1 and DNAH1 in sperm flagella of humans and mice (Fig. 3F, 3G, 5E, and 5F), we proposed that the interaction among DNAH12, DNALI1, and DNAH1 is conserved in mammals, and is required for flagellar development during spermiogenesis. Since the deficiency of DNAH12 does not cause PCD symptoms in humans and mice, we hypothesized that flagella and cilia may exhibit distinct features in the process of dynein assembly. We thus checked the abundance of DNALI1 and DNAH1 in the oviduct and trachea, the results showed that DNALI1 and DNAH1 were comparable in the *Dnah12*+/+ and *Dnah12*-/- groups (Fig. 7F and 7G). Furthermore, no discernible disparities were observed in the intensity and localization of DNALI1 and DNAH1 signals within the oviduct and trachea (Fig. 7H-K). In summary, these results underscore the specific involvement of DNAH12 in flagellar development, rather than in cilial development. Moreover, we supposed that the assembly of IDA components differs between flagella and cilia.

DNAH12 is prerequisite for the recruitment of DNALI1 and DNAH1 to sperm flagella

In patients and mice with DNALI1 deficiency, flagellar DNAH1 but not DNAH12 signals were absent (47). To unravel the recruitment relationships among DNAH12, DNALI1, and DNAH1 during sperm flagellar formation, we conducted staining of DNALI1 and DNAH12 on sperm flagella obtained from the infertile asthenoteratozoospermic male P13 with a homozygous *DNAH1* mutation (c.11275C>T) of a family (Fig. 7L, and Supplemental Figure13 A and B), wherein DNAH1 signals were absent along the flagella (Fig. 7M). In comparison to those in the control, the DNALI1 signals were absent, while no discernible differences in DNAH12 signals were observed in the flagella of the patient with the *DNAH1* mutation (Fig. 7N and 7O). These findings suggested that the recruitment of DNAH12 to flagella is independent of DNAH1, and DNAH12 is the first protein to be recruited to flagella among the three proteins.

DNAH12 deficiency causes the decrease of RS head associated proteins

In addition to the enrichment of dynein proteins, we noticed that a clusters of RS head-related proteins such as RSPH1, RSPH9, or DNAJB13 were enriched in the testicular interactome of DNAH12. Considering that deficiency in RSPH1 (48), RSPH9 (49), or DNAJB13 (50) were associated with CP loss phenotype. We then explored their interaction with DNAH12 in the testis lysate, the results proved the interactions of these proteins with DNAH12 in the testis (Fig. 8A). Further analyses by AlphaFold3 showed that these proteins may interacts with DNAH12 through Stem domain (Fig. 8B). The expression of RSPH1, RSPH9, and DNAJB13 showed obvious decrease in testis lysate of $Dnah12^{-/-}$ mice (Fig. 8C). Additionally, immunofluorescence assays revealed the loss of signals for RSPH1, a marker of RS head, in $Dnah12^{-/-}$ sperm flagella (Fig. 8D). Together, these results indicate that deficiency of DNAH12 lead to the impairment of radial spoke head, which could contribute to CP loss in flagella of $Dnah12^{-/-}$ mice.

ICSI may overcome the infertility caused by DNAH12 deficiency

Since the DNAH12 mutated patients and the $Dnah12^{-/-}$ mice have similar deficiency, to evaluate the prospects of whether the asthenoteratozoospermia patients caused by $DNAH12$ variants could be treated with assisted reproductive technology, spermatozoa from D$nah12^{+/+}$ and $Dnah12^{-/-}$ mice were used for ICSI. Promisingly, the percentages of 2-cell and blastocyst-stage embryos in $Dnah12^{-/-}$ group showed no difference compared to those of $Dnah12^{+/+}$ group (Fig. 8E-G), indicating that ICSI could also be an effective treatment for infertile men carrying $DNAH12$ variants.

Taken together, we suggest that DNAH12 interacts with DNALI1 and DNAH1, and this interaction is a prerequisite for the proper localization of DNALI1 and DNAH1 to sperm flagella, which is essential for the intact IDAs and CP structures in both humans and mice. Additionally, DNALI1 and DNAH1 exhibit mutual dependence for their localization to sperm flagella. Deficiency of DNAH12 leads to the failure of DNALI1 and DNAH1 recruitment to sperm flagella, resulting in abnormal sperm morphology characterized by compromised axoneme organization, causing male infertility (Fig. 8H). Moreover, our results indicate that the assembly of IDA components occurs intricately, possibly not synchronously in time and space.

Discussion

In this study, we identified six novel bi-allelic mutations in $DNAH12$ that cause the loss of DNAH12 proteins in asthenoteratozoospermia patients. Spermatozoa from these patients exhibited abnormal morphology and compromised axonemal defects, including a high percentage of impaired IDAs and loss of CP. Two mouse models ($Dnah12^{-/-}$ and $Dnah12^{mut/mut}$) were generated, recapitulating the abnormal flagellum morphology and structure observed in patients, thereby confirming the pathogenic role of $DNAH12$ mutations in asthenoteratozoospermia. Through testicular Co-IP and mass spectrometry proteomics, we screened and validated that DNAH12 interacts with IDA components like DNALI1 and DNAH1 and is indispensable for recruiting DNALI1 and DNAH1 to sperm flagella, highlighting the crucial role of DNAH12 in sperm flagellar development. Notably, deficiency of DNAH12 does not manifest PCD symptoms in humans and mice, underscoring the unique role of DNAH12 in flagellar development. Collectively, our study not only unveils the functional significance of the previously unexplored gene $DNAH12$ but also elucidates the pathogenic link between $DNAH12$ mutations and asthenoteratozoospermia.

DNAH12 and other dynein arm proteins share the N-terminal Stem domain, which is crucial for
structure and accessory subunit binding (51), and the motor domain which encompasses ~6 AAA+
ATPase modules forming the ring-shaped oligomeric complexes responsible for force production (52).
Five of the identified *DNAH12* mutations, primarily in the Stem domain (critical for DNALI1 interaction),
result in loss-of-function effects, indicating a weak tolerance property of the Stem domain. Similar weak
tolerance features were observed in mutations of other dynein arm components like *DNAH1* and *DNAH2*
(53, 54). The profound impact of these variants, particularly missense mutations, underscores delicate
features of the Stem domain and DNAH family members. Notably, DNAH12 stands out as the shortest
in sequence length and the only one lacking the essential MTBD domain within the DNAH family (55).
MTBD, a 15-nanometer coiled-coil stalk structure, crucially supports ATP-sensitive microtubule binding
properties and determines dynein motility direction (56). Our investigation revealed that DNAH12
interacts with other IDA components, specifically DNALI1 and DNAH1. Despite the mutual dependence
between DNAH1 and DNALI1 for their flagellar localization, DNAH12 is prerequisite for the proper
assembly of DNALI1 and DNAH1 onto sperm flagella in both humans and mice. Notably, *Dnah12*$^{-/-}$
mice exhibit more severe phenotypes than *Dnali1* and *Dnah1* knockout mice (4, 47, 57), including a
more severe reduced sperm count (~ 2% of normal values), this observation further supports our
perspective.

Deficiency in DNAH12 presented sperm ultrastructural defects like CP loss and impaired IDAs
indicating essential roles of DNAH12 in the maintenance of axonemal integrity and arrangement. These
results indicate essential roles of DNAH12 in the maintenance of axonemal integrity and arrangement.
Noteworthy, deficiency of DNAH12 potential partner DNAH1, a well-known asthenoteratozoospermia-
related gene, presents quite similar defects with DNAH12, indicating DNAH12 and DNAH1 possibly
exhibit synergic functions in axonemal proper arrangement (58-60). In addition to DNAH12 and DNAH1,
deficiency in DNAH2 or DNAH6 also leads to high proportions of CP loss and disorganized axonemal
structures (5, 54, 61). However, DNALI1, DNAH7, or DNAH10 deficiency results in no obvious CP loss
but impaired IDA integrity (7, 8, 22, 47). These studies indicate that specific IDA components, like
DNAH12 and DNAH1, are required for the maintenance of CP integrity. Moreover, discrepancies
observed in the phenotypes resulting from deficiencies in specific IDAs suggest that multiple dynein
proteins are produced to support the intricate organization of sperm flagella. Additionally, these proteins
may exhibit distinct characteristics and perform unique functions to some extent. Noteworthy, deficiency
in RS head proteins RSPH1, RSPH9, or DNAJB13 were related to axonemal CP loss phenotype (48, 50,
62), indicating that DNAH12, and other IDAs, possibly regulate the stability of CP through the
cooperation with RS head proteins.

The localization of DNAH12 presents dynamic changes during spermiogenesis. DNAH12 is
initially expressed in the cytoplasm of the round spermatid, while it's enriched in the manchette structure
of the elongating spermatid and the tail of the elongated spermatid. The manchette and the sperm tail
axoneme are two microtubular platforms that transport proteins to the developing head and tail (63).
Manchette is a transient skirt-like structure and functions as a scaffold for morphological remodeling,
contributing to the formation of the sperm head, neck, and tail (64). Here, we found that DNAH12
deficiency results in abnormal manchette structure and compromised sperm head and tail morphology.
Additionally, in sperm of *Dnah12*$^{-/-}$ males, several CP complex proteins like SPAG6, and SPEF2 are not
properly assembled and the axoneme structure of the tail was severely disorganized, indicating dynein

components are responsible for proper axonemal components organization. Noteworthy, DNALI1, a
potential DNAH12 interacting partner showed similar localization patterns to DNAH12, a study
presented that DNALI1 recruits and stabilizes manchette component PACRG, and DNALI1 deficiency
causes impaired sperm flagellar assembly (57). These data hint that dynein proteins like DNAH12 and
DNALI, possibly work together in the manchette organization as well as sperm flagellar components
assembly during sperm development.

DNAH12 was also detected in the lungs, tracheas, and female oviducts while no cilia morphology
defects in these tissues or PCD symptoms were observed in both two mice models. Besides, none of the
patients carrying bi-allelic *DNAH12* mutations claimed symptoms of PCD via visits except for the
reproductive problems. *Dnah12*$^{-/-}$ female mice showed normal development and fertility and the mother
(III:2) of patients from family 2, carrying homozygous *DNAH12* mutation M2 had 6 offspring.
Intriguingly, we found that the localization and abundance of DNALI1 and DNAH1 in oviductal or
tracheal cilia were not affected after *Dnah12* deletion, indicating that DNAH12 works differently in
flagella and cilia. Deficiency of specific DNAH members like DNAH6 (65), or DNAH7 (22), but not
DNAH2 (54), results in PCD symptoms. Hence, DNAH family members possibly present disparate
processing of axonemal biogenesis and organization through evolution in flagella and cilia, further
explorations could contribute to a more comprehensive understanding of the differences between flagella
and cilia, and personalized clinical diagnosis of related diseases like male/female infertility and/or PCD
(66).

Previously, two genetic screening studies mentioned unverified compound heterozygous mutations
and a homozygous missense mutation in *DNAH12*, predicting their association with patients' infertility.
However, limited genetic evidence was provided, and the pathogenesis remained unclear (26, 27).
Through investigations involving patients with *DNAH12* variants and mutant mouse models, we provide
direct genetic and molecular evidence that variants in *DNAH12* are genetic causes of
asthenoteratozoospermia, and DNAH12 along with DNAH1, DNALI1 is indispensable for sperm
flagellar biogenesis. These data strongly support DNAH12 as a valuable marker in genetic counseling
and diagnosis of male infertility.

ICSI is a widely used assisted reproductive technology in treating infertile men with
asthenoteratozoospermia. Deficiency in most dynein proteins leads to asthenoteratozoospermia with
severely compromised sperm motility and morphology. Previous studies have shown that patients
carrying mutations in *DNAH17* (10) have poor ICSI outcomes while patients with *DNAH1*(59), *DNAH2*
(5, 61), *DNAH3* (67), *DNAH7* (22), or *DNAH8* (68) mutations have achieved favorable clinical pregnancy.
Thus, the comparative study of asthenoteratozoospermia-related genes on their ICSI outcomes will be
beneficial to developing individualized treatment recommendations in clinical practice. In our study, the
successful ICSI outcomes of *Dnah12*$^{-/-}$ mice suggest that ICSI could be used as a potential treatment for
infertile men carrying *Dnah12* mutations. It's meaningful to evaluate the prognosis for ICSI treatments
in patients carrying *DNAH12* variants in a large population.

In conclusion, our study functionally and mechanistically sheds light on a special and poorly
characterized protein DNAH12, during the intricate process of sperm flagellar biogenesis. These results
pave the way for an enhanced understanding of the roles of dynein proteins in sperm flagellar

development and their functional difference in flagella and cilia, offering a diagnostic target for genetic
counseling and future individualized treatment of male infertility in clinical practice.
605

606 **Data availability**
607 All data that support the findings in this study are available from the corresponding author upon
608 reasonable request. Proteomics data supporting this study have been deposited to the ProteomeXchange
609 Consortium via the iProX partner repository with the dataset identifier PXD051681.
610

Author contributions
B.S. and Q.S. conceived and supervised the study; M.Y., X.H., J.Y, M.C., A.Z., T.L., A.Z., and N.A.
performed the experiments and analyzed the data; H.M.J, M.K., Z.M., R.K., A.A., and W.S. collected the
patients' samples; J.Z. and H.Z. performed the WES sequencing and data analyses; M.Y. wrote the
manuscript and B.S., Q.S., H.M., and M.Y. modified the manuscript; B.S., Q.S, H.M., B.X., H.Z., and
A.M. were involved in discussions of the manuscript.
617

618 **Ethics statement**
619 This work was approved by the institutional ethics committee of University of Science and Technology
620 of China (USTC) with the approval number 2019-KY-168, and informed consent was obtained from
621 patients and controls. All animal experiments were approved by the Institutional Animal Care and Use
622 Committee of the USTC with the approval number USTCACUC1301021.
623

Funding
This work was supported by the National Key Research and Development Program of China
(2022YFC2702601, 2022YFA0806303, and 2019YFA0802600), the National Natural Science
Foundation of China (82171599, 32270901, and 32070850), the Global Select Project (DJK-LX-
2022010) of the Institute of Health and Medicine, Hefei Comprehensive National Science Center, the
Joint Fund for New Medicine of USTC (YD9100002034) and Scientific Research Foundation for
Scholars of the First Affiliated Hospital of USTC (RC2023054).
631

632 **Declaration of competing interest**
633 The authors declare that they have no known competing financial interests or personal relationships that
634 could have appeared to influence the work reported in this paper.
635

636 **Acknowledgments**
637 We are grateful to all the participants for their cooperation. We appreciated Li Wang and Dandan Song
638 at the Center of Cryo-Electron Microscopy, Zhejiang University, for their technical assistance with TEM
639 and SEM, respectively. We thank the Bioinformatics Center of the USTC, School of Life Sciences, for
640 providing supercomputing resources.
641

References
1.   Agarwal A, Baskaran S, Parekh N, Cho C-L, Henkel R, Vij S, et al. Male infertility. *Lancet.*
2021;397(10271):319-33.
2.   Krausz C, and Riera-Escamilla A. Genetics of male infertility. *Nat Rev Urol.* 2018;15(6):369-
84.

3.   Shang X, Shen C, Liu J, Tang L, Zhang H, Wang Y, et al. Serine protease PRSS55 is crucial for
648   male mouse fertility via affecting sperm migration and sperm-egg binding. *Cell Mol Life Sci.*
649   2018;75(23):4371-84.

650   4.   Khan R, Zaman Q, Chen J, Khan M, Ma A, Zhou J, et al. Novel Loss-of-Function Mutations in
651   DNAH1 Displayed Different Phenotypic Spectrum in Humans and Mice. *Front Endocrinol*
652   *(Lausanne).* 2021;12:765639.

653   5.   Li Y, Sha Y, Wang X, Ding L, Liu W, Ji Z, et al. DNAH2 is a novel candidate gene associated
654   with multiple morphological abnormalities of the sperm flagella. *Clin Genet.* 2019;95(5):590-
655   600.

656   6.   Tu C, Nie H, Meng L, Yuan S, He W, Luo A, et al. Identification of DNAH6 mutations in
657   infertile men with multiple morphological abnormalities of the sperm flagella. *Sci Rep.*
658   2019;9(1):15864.

659   7.   Gao Y, Liu L, Shen Q, Fu F, Xu C, Geng H, et al. Loss of function mutation in DNAH7 induces
660   male infertility associated with abnormalities of the sperm flagella and mitochondria in human.
661   *Clinical Genetics.* 2022;102(2):130-5.

662   8.   Li K, Wang G, Lv M, Wang J, Gao Y, Tang F, et al. Bi-allelic variants in DNAH10 cause
663   asthenoteratozoospermia and male infertility. *J Assist Reprod Genet.* 2022;39(1):251-9.

664   9.   Liu C, Miyata H, Gao Y, Sha Y, Tang S, Xu Z, et al. Bi-allelic DNAH8 Variants Lead to Multiple
665   Morphological Abnormalities of the Sperm Flagella and Primary Male Infertility. *Am J Hum*
666   *Genet.* 2020;107(2):330-41.

667   10.   Whitfield M, Thomas L, Bequignon E, Schmitt A, Stouvenel L, Montantin G, et al. Mutations
668   in DNAH17, Encoding a Sperm-Specific Axonemal Outer Dynein Arm Heavy Chain, Cause
669   Isolated Male Infertility Due to Asthenozoospermia. *Am J Hum Genet.* 2019;105(1):198-212.

670   11.   Zhang B, Ma H, Khan T, Ma A, Li T, Zhang H, et al. A DNAH17 missense variant causes
671   flagella destabilization and asthenozoospermia. *J Exp Med.* 2020;217(2).

672   12.   Coutton C, Vargas AS, Amiri-Yekta A, Kherraf ZE, Ben Mustapha SF, Le Tanno P, et al.
673   Mutations in CFAP43 and CFAP44 cause male infertility and flagellum defects in Trypanosoma
674   and human. *Nat Commun.* 2018;9(1):686.

675   13.   Tang S, Wang X, Li W, Yang X, Li Z, Liu W, et al. Biallelic Mutations in CFAP43 and CFAP44
676   Cause Male Infertility with Multiple Morphological Abnormalities of the Sperm Flagella. *Am J*
677   *Hum Genet.* 2017;100(6):854-64.

678   14.   Ma A, Zhou J, Ali H, Abbas T, Ali I, Muhammad Z, et al. Loss-of-function mutations in CFAP57
679   cause multiple morphological abnormalities of the flagella in humans and mice. *JCI Insight.*
680   2023;8(3).

681   15.   Hu T, Meng L, Tan C, Luo C, He WB, Tu C, et al. Biallelic CFAP61 variants cause male
682   infertility in humans and mice with severe oligoasthenoteratozoospermia. *J Med Genet.*
683   2023;60(2):144-53.

684   16.   Toure A, Martinez G, Kherraf ZE, Cazin C, Beurois J, Arnoult C, et al. The genetic architecture
685   of morphological abnormalities of the sperm tail. *Hum Genet.* 2021;140(1):21-42.

686   17.   Walton T, Gui M, Velkova S, Fassad MR, Hirst RA, Haarman E, et al. Axonemal structures
687   reveal mechanoregulatory and disease mechanisms. *Nature.* 2023;618(7965):625-33.

688   18.   Viswanadha R, Sale WS, and Porter ME. Ciliary Motility: Regulation of Axonemal Dynein
689   Motors. *Cold Spring Harb Perspect Biol.* 2017;9(8).

690   19.   Kleinboelting S, Diaz A, Moniot S, van den Heuvel J, Weyand M, Levin LR, et al. Crystal

691		structures of human soluble adenylyl cyclase reveal mechanisms of catalysis and of its
692		activation through bicarbonate. *Proc Natl Acad Sci U S A.* 2014;111(10):3727-32.
693	20.	Shoemark A, Burgoyne T, Kwan R, Dixon M, Patel MP, Rogers AV, et al. Primary ciliary
694		dyskinesia with normal ultrastructure: three-dimensional tomography detects absence of
695		DNAH11. *Eur Respir J.* 2018;51(2).
696	21.	Sironen A, Shoemark A, Patel M, Loebinger MR, and Mitchison HM. Sperm defects in primary
697		ciliary dyskinesia and related causes of male infertility. *Cell Mol Life Sci.* 2020;77(11):2029-48.
698	22.	Wei X, Sha Y, Wei Z, Zhu X, He F, Zhang X, et al. Bi-allelic mutations in DNAH7 cause
699		asthenozoospermia by impairing the integrality of axoneme structure. *Acta Biochim Biophys*
700		*Sin (Shanghai).* 2021;53(10):1300-9.
701	23.	Gao Y, Liu L, Shen Q, Fu F, Xu C, Geng H, et al. Loss of function mutation in DNAH7 induces
702		male infertility associated with abnormalities of the sperm flagella and mitochondria in human.
703		*Clin Genet.* 2022;102(2):130-5.
704	24.	Imtiaz F, Allam R, Ramzan K, and Al-Sayed M. Variation in DNAH1 may contribute to primary
705		ciliary dyskinesia. *BMC Med Genet.* 2015;16:14.
706	25.	Vaisberg EA, Grissom PM, and McIntosh JR. Mammalian cells express three distinct dynein
707		heavy chains that are localized to different cytoplasmic organelles. *J Cell Biol.*
708		1996;133(4):831-42.
709	26.	Li Y, Wang Y, Wen Y, Zhang T, Wang X, Jiang C, et al. Whole-exome sequencing of a cohort of
710		infertile men reveals novel causative genes in teratozoospermia that are chiefly related to sperm
711		head defects. *Hum Reprod.* 2021.
712	27.	Oud MS, Houston BJ, Volozonoka L, Mastrorosa FK, Holt GS, Alobaidi BKS, et al. Exome
713		sequencing reveals variants in known and novel candidate genes for severe sperm motility
714		disorders. *Hum Reprod.* 2021;36(9):2597-611.
715	28.	Chapelin C, Duriez B, Magnino F, Goossens M, Escudier E, and Amselem S. Isolation of several
716		human axonemal dynein heavy chain genes: genomic structure of the catalytic site, phylogenetic
717		analysis and chromosomal assignment. *FEBS Lett.* 1997;412(2):325-30.
718	29.	World Health Organization. *WHO Laboratory Manual for the Examination and Processing of*
719		*Human Semen.*: 6th edn. Geneva： World Health Organization; 2021.
720	30.	Wang K, Li M, and Hakonarson H. ANNOVAR: functional annotation of genetic variants from
721		high-throughput sequencing data. *Nucleic Acids Res.* 2010;38(16):e164.
722	31.	Yin H, Ma H, Hussain S, Zhang H, Xie X, Jiang L, et al. A homozygous FANCM frameshift
723		pathogenic variant causes male infertility. *Genet Med.* 2019;21(1):62-70.
724	32.	Narasimhan V, Danecek P, Scally A, Xue Y, Tyler-Smith C, and Durbin R. BCFtools/RoH: a
725		hidden Markov model approach for detecting autozygosity from next-generation sequencing
726		data. *Bioinformatics.* 2016;32(11):1749-51.
727	33.	Tamura K, Stecher G, and Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis
728		Version 11. *Mol Biol Evol.* 2021;38(7):3022-7.
729	34.	He Z, Zhang H, Gao S, Lercher MJ, Chen WH, and Hu S. Evolview v2: an online visualization
730		and management tool for customized and annotated phylogenetic trees. *Nucleic Acids Res.*
731		2016;44(W1):W236-41.
732	35.	Chen C, Chen H, Zhang Y, Thomas HR, Frank MH, He Y, et al. TBtools: An Integrative Toolkit
733		Developed for Interactive Analyses of Big Biological Data. *Mol Plant.* 2020;13(8):1194-202.
734	36.	Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al. Pfam:

735   The protein families database in 2021. *Nucleic Acids Res.* 2021;49(D1):D412-D9.

37. Kelley LA, Mezulis S, Yates CM, Wass MN, and Sternberg MJ. The Phyre2 web portal for protein modeling, prediction and analysis. *Nat Protoc.* 2015;10(6):845-58.

38. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem.* 2004;25(13):1605-12.

39. Ma H, Zhang B, Khan A, Zhao D, Ma A, Zhou J, et al. Novel frameshift mutation in STK33 is associated with asthenozoospermia and multiple morphological abnormalities of the flagella. *Hum Mol Genet.* 2021;30(21):1977-84.

40. Ma H, Li T, Xie X, Jiang L, Ye J, Gong C, et al. RAD51AP2 is required for efficient meiotic recombination between X and Y chromosomes. *Sci Adv.* 2022;8(2):eabk1789.

41. Zhang B, Khan I, Liu C, Ma A, Khan A, Zhang Y, et al. Novel loss-of-function variants in DNAH17 cause multiple morphological abnormalities of the sperm flagella in humans and mice. *Clin Genet.* 2021;99(1):176-86.

42. Dil S, Khan A, Unar A, Yang ML, Ali I, Zeb A, et al. A novel homozygous frameshift variant in DNAH8 causes multiple morphological abnormalities of the sperm flagella in a consanguineous Pakistani family. *Asian J Androl.* 2022.

43. Wang H, Yang H, Shivalila CS, Dawlaty MM, Cheng AW, Zhang F, et al. One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell.* 2013;153(4):910-8.

44. Xu J, Gao J, Liu J, Huang X, Zhang H, Ma A, et al. ZFP541 maintains the repression of pre-pachytene transcriptional programs and promotes male meiosis progression. *Cell Rep.* 2022;38(12):110540.

45. Chen T, Ma J, Liu Y, Chen Z, Xiao N, Lu Y, et al. iProX in 2021: connecting proteomics data sharing with big data. *Nucleic Acids Res.* 2022;50(D1):D1522-d7.

46. Gao J, Zhang H, Jiang X, Ali A, Zhao D, Bao J, et al. FertilityOnline: A Straightforward Pipeline for Functional Gene Annotation and Disease Mutation Discovery. *Genomics Proteomics Bioinformatics.* 2022;20(3):455-65.

47. Wu H, Liu Y, Li Y, Li K, Xu C, Gao Y, et al. DNALI1 deficiency causes male infertility with severe asthenozoospermia in humans and mice by disrupting the assembly of the flagellar inner dynein arms and fibrous sheath. *Cell Death Dis.* 2023;14(2):127.

48. Lin J, Yin W, Smith MC, Song K, Leigh MW, Zariwala MA, et al. Cryo-electron tomography reveals ciliary defects underlying human RSPH1 primary ciliary dyskinesia. *Nat Commun.* 2014;5:5727.

49. Zhu L, Liu H, Chen Y, Yan X, and Zhu X. Rsph9 is critical for ciliary radial spoke assembly and central pair microtubule stability. *Biol Cell.* 2019;111(2):29-38.

50. Liu M, Li J, Jiang C, Zhou Y, Sun Y, Yang Y, et al. A novel homozygous mutation in DNAJB13-a gene associated with the sperm axoneme-leads to teratozoospermia. *J Assist Reprod Genet.* 2022;39(3):757-64.

51. Kollmar M. Fine-Tuning Motile Cilia and Flagella: Evolution of the Dynein Motor Proteins from Plants to Humans at High Resolution. *Mol Biol Evol.* 2016;33(12):3249-67.

52. Canty JT, Tan R, Kusakci E, Fernandes J, and Yildiz A. Structure and Mechanics of Dynein Motors. *Annu Rev Biophys.* 2021;50:549-74.

53. Sha Y, Yang X, Mei L, Ji Z, Wang X, Ding L, et al. DNAH1 gene mutations and their potential

association with dysplasia of the sperm fibrous sheath and infertility in the Han Chinese
population. *Fertil Steril.* 2017;107(6):1312-8 e2.

54.   Hwang JY, Nawaz S, Choi J, Wang H, Hussain S, Nawaz M, et al. Genetic Defects in DNAH2
Underlie Male Infertility With Multiple Morphological Abnormalities of the Sperm Flagella in
Humans and Mice. *Front Cell Dev Biol.* 2021;9:662903.

55.   Lacey SE, He S, Scheres SH, and Carter AP. Cryo-EM of dynein microtubule-binding domains
shows how an axonemal dynein distorts the microtubule. *Elife.* 2019;8.

56.   Carter AP, Garbarino JE, Wilson-Kubalek EM, Shipley WE, Cho C, Milligan RA, et al.
Structure   and   functional   role   of   dynein's   microtubule-binding   domain.   *Science.*
2008;322(5908):1691-5.

57.   Yap YT, Li W, Huang Q, Zhou Q, Zhang D, Sheng Y, et al. DNALI1 interacts with the
MEIG1/PACRG complex within the manchette and is required for proper sperm flagellum
assembly in mice. *Elife.* 2023;12.

58.   Ben Khelifa M, Coutton C, Zouari R, Karaouzene T, Rendu J, Bidart M, et al. Mutations in
DNAH1, which encodes an inner arm heavy chain dynein, lead to male infertility from multiple
morphological abnormalities of the sperm flagella. *Am J Hum Genet.* 2014;94(1):95-104.

59.   Wambergue C, Zouari R, Fourati Ben Mustapha S, Martinez G, Devillard F, Hennebicq S, et al.
Patients with multiple morphological abnormalities of the sperm flagella due to DNAH1
mutations have a good prognosis following intracytoplasmic sperm injection. *Hum Reprod.*
2016;31(6):1164-72.

60.   Wang M, Yang QY, Zhou JP, Tan HP, Hu J, Jin L, et al. Novel compound heterozygous mutations
in DNAH1 cause primary infertility in Han Chinese males with multiple morphological
abnormalities of the sperm flagella. *Asian J Androl.* 2022.

61.   Gao Y, Tian S, Sha Y, Zha X, Cheng H, Wang A, et al. Novel bi-allelic variants in DNAH2 cause
severe asthenoteratozoospermia with multiple morphological abnormalities of the flagella.
Reprod Biomed Online. 2021;42(5):963-72.

62.   Zou W, Lv Y, Liu ZI, Xia P, Li H, and Jiao J. Loss of Rsph9 causes neonatal hydrocephalus with
abnormal development of motile cilia in mice. *Sci Rep.* 2020;10(1):12435.

63.   Lehti MS, and Sironen A. Formation and function of the manchette and flagellum during
spermatogenesis. *Reproduction.* 2016;151(4):R43-54.

64.   Hu W, Zhang R, Xu H, Li Y, Yang X, Zhou Z, et al. CAMSAP1 role in orchestrating structure
and   dynamics   of   manchette   microtubule   minus-ends   impacts   male   fertility   during
spermiogenesis. *Proc Natl Acad Sci U S A.* 2023;120(45):e2313787120.

65.   Li Y, Yagi H, Onuoha EO, Damerla RR, Francis R, Furutani Y, et al. DNAH6 and Its Interactions
with   PCD   Genes   in   Heterotaxy   and   Primary   Ciliary   Dyskinesia.   *PLoS   Genet.*
2016;12(2):e1005821.

66.   Zhang X, Xiao Z, Zhang J, Xu C, Liu S, Cheng L, et al. Differential requirements of IQUB for
the assembly of radial spoke 1 and the motility of mouse cilia and flagella. *Cell Rep.*
2022;41(8):111683.

67.   Meng GQ, Wang Y, Luo C, Tan YM, Li Y, Tan C, et al. Bi-allelic variants in DNAH3 cause male
infertility   with   asthenoteratozoospermia   in   humans   and   mice.   *Hum   Reprod   Open.*
2024;2024(1):hoae003.

68.   Weng M, Sha Y, Zeng YU, Huang N, Liu W, Zhang X, et al. Mutations in DNAH8 contribute
to multiple morphological abnormalities of sperm flagella and male infertility. *Acta Biochim*

Figure Legends

Figure 1. Identification of bi-allelic *DNAH12* variants in infertile men with
asthenoteratozoospermia. (A) Pedigrees of family 1 with 3 infertile males, P1 (IV:1), P2 (IV:2), and P3
(IV:4), family 2 with 3 infertile males, P4 (IV:3), P5 (IV:4) and P6 (IV:5), family 3 with 2 infertile males,
P7 (IV:1) and P8 (IV:3), and three sporadic infertile cases P9, P10 and P11. Red arrows point to the
individuals whom whole-exome sequencing was performed. Candidate mutations were validated by
Sanger sequencing and the chromatograms were shown on the right. Red arrowheads point to mutant
sites. Double horizontal lines represent consanguineous marriages. WT, wild-type allele; MT, the mutant
allele. (B) The positions of the *DNAH12* variants at the transcript (ENST00000351747.2) and protein
levels (Q6ZR08-1). MT-like, Microtubule-binding-like domain. (C) Conservation analyses of affected
amino acids across different species, the arrowheads indicate the mutation sites.

Figure 2. Abnormal sperm morphology and undetectable DNAH12 expression in men with bi-
allelic *DNAH12* variants. (A) Representative micrographs show normal sperm morphology from a
fertile control and abnormal spermatozoa from P10 with absent, short, coiled, bent, irregular flagella or
abnormal head under light microscopy. Scale bars, 5μm. (B) Representative SEM micrographs show
details about the sperm morphology of a fertile control and P10. The normal sperm morphology can be
observed in the fertile control (left) and abnormal types like absent, short, coiled, bent, irregular tails,
and head abnormality in P10 (right). Scale bars, 5μm. (C) Immunoblotting assay revealed that DNAH12
was absent in the spermatozoa from P9, P10, and P11 harboring *DNAH12* variants. β-actin was used as
a loading control. (D) Sperm cells were co-stained with α-Tubulin (green) and DNAH12 (red) antibodies
while no DNAH12 signals were observed in sperm flagella of patients. DNA was counterstained with
Hoechst 33342. Scale bars, 10μm.

Figure 3. Flagellar axoneme defects were detected in the patients with *DNAH12* variants. (A-B)
Representative TEM micrographs showing cross-sections of the midpiece, principal piece, and end piece
of sperm flagella from a fertile control and P11. The axoneme structure in control presents a "9+2"
microtubules arrangement, including mitochondrial sheath (MS, indicated with red arrows), central pair
of microtubules (CP, indicated with yellow arrows), outer dense fibers (ODFs, indicated with cerulean
arrows), peripheral doublet microtubules (DMTs, indicated with orange arrows), inner dynein arms
(IDAs, indicated with green arrows) and outer dynein arms (ODAs, indicated with purple arrows), while
in P11, axonemal defects like missing CP, disorganized axonemal structures or missing DMTs were
observed, the red asterisks mark CP loss (A). Impaired IDAs in the sperm axoneme of patient P11 (B).
The red triangle marks impaired IDAs while adjacent ODAs are identifiable. Scale bars, 200 nm. (C-D)
The proportion of different categories of observed cross-sections in the control and patients. Cross-
sections were classified into four categories: Intact "9+2", missing CP, disorganization, and missing
DMTs (C), and the proportion of intact IDAs or impaired IDAs in the cross-sections of flagellar axoneme
of which ODAs were identifiable (D). n, the total number of cross-sections for quantification. (E-G)
Representative images of spermatozoa from fertile controls and patients carrying bi-allelic *DNAH12*
variants co-stained α-Tubulin with SPAG6 (E), DNAH1 (F), DNALI1 (G), or DNAH17 (H), and Hoechst
33342 for DNA (blue). Scale bars, 10 μm.

Figure 4. Disruption of *Dnah12* results in impaired spermatogenesis and abnormal sperm morphology. (A) Fertility test of adult *Dnah12$^{+/+}$* and *Dnah12$^{-/-}$* male mice. (B) The testes to body weight ratios of *Dnah12$^{+/+}$ and Dnah12$^{-/-}$* mice. (C) The number of sperm per epididymis of *Dnah12$^{+/+}$ and Dnah12$^{-/-}$* male mice. (D) Percentages of motile sperms in *Dnah12$^{+/+}$ and Dnah12$^{-/-}$* male mice. (E) Histological sections of testis and epididymis from *Dnah12$^{+/+}$* and *Dnah12$^{-/-}$* mice after H&E staining. Scale bars, 50 μm. (F) Morphology of the sperm from caudal epididymis of *Dnah12$^{+/+}$* and *Dnah12$^{-/-}$* mice. Scale bars,10 μm. (G) Quantitative analysis of sperm morphology of *Dnah12$^{+/+}$ and Dnah12$^{-/-}$* male mice. The experiments were repeated three times with at least 200 sperms counted every time. The above data were obtained from 3 adult mice for each genotype. (H) Representative images of testicular cells from *Dnah12$^{+/+}$* and *Dnah12$^{-/-}$* mice co-stained DNAH12 and α-Tubulin antibodies. Scale bars, 10 μm. The red asterisk indicates abnormal manchette structure and the red arrow indicates abnormal sperm flagellum at the elongated spermatid stage of *Dnah12$^{-/-}$* mice. (I) Representative immunofluorescence images of seminiferous tubule squash from 8-week-old *Dnah12$^{+/+}$* and *Dnah12$^{-/-}$* mice, co-stained by PNA (red) and α-Tubulin (green) antibody. DNA was stained with Hoechst 33342. Scale bars, 10 μm. The asterisks indicate the manchette defects and the red triangles mark the abnormal sperm heads. Scale bars,10 μm. (J) Representative TEM micrographs of testicular elongated spermatids from *Dnah12$^{+/+}$* and *Dnah12$^{-/-}$* mice. Scale bars, 1μm. (K) TUNEL assay on the testicular sections from *Dnah12$^{+/+}$* and *Dnah12$^{-/-}$* mice. The green triangles mark the apoptotic elongating or elongated spermatids. Scale bars, 50 μm. (L) The proportion of tubules with apoptotic cells in *Dnah12$^{+/+}$* and *Dnah12$^{-/-}$* mice. (M) Average numbers of TUNEL-positive cells per tubule in *Dnah12$^{+/+}$* and *Dnah12$^{-/-}$* mice. Data are obtained from three mice with at least 50 tubules scored in each repeated experiment. **p < 0.01; ***p <0.001; ****p < 0.0001, Student's t test.

Figure 5. Deletion of *Dnah12* causes axoneme defects in mice. (A) Representative TEM micrographs showing cross-sections of sperm flagella from *Dnah12$^{+/+}$*, *Dnah12$^{+/-}$*, and *Dnah12$^{-/-}$* mice. In *Dnah12$^{+/+}$*, *Dnah12$^{+/-}$* groups, the normal axoneme "9+2" microtubule arrangement mainly consisted of the MS, (indicated with red arrows), CP (yellow arrows), ODFs (cerulean arrows), DMTs (orange arrows), IDAs (green arrows) and ODAs (purple arrows). However, the CP and DMTs structures were missing or disarranged in the *Dnah12$^{-/-}$* group. Red asterisks mark the loss of CP. Scale bars, 200 nm. (B) Immunoblotting of sperm lysate from *Dnah12$^{+/+}$, DNAH12$^{+/-}$* and *Dnah12$^{-/-}$* mice using DNAH1, DNALI1 or SPAG6 antibodies. DNAI2 and α-Tubulin were used as the loading controls. (C-H) Representative images of caudal epididymal sperm from *Dnah12$^{+/-}$* and *Dnah12$^{-/-}$* mice co-stained α-Tubulin and SPAG6 (C), SPEF2 (a component of CP complex) (D), DNAH1 (E), DNALI1 (F), DNAH17 (G) or DNAI2 (H) antibodies. Scale bars, 10 μm.

Figure 6. DNAH12 interacts with IDA components DNALI1 and DNAH1 (A) Phylogenetic and protein domain analyses of the DNAH family. The dashed box indicates the shortest-length member, DNAH12, while the arrow points to the missing microtubule-binding domain(MTBD) region. (B) Schematic diagram showing the procedure of the immunoprecipitation/mass spectrometry (IP/MS) proteomics experiment. (C) GO term enrichment analysis of DNAH12 potential interactors. (D) Protein-protein interaction network of DNAH12 among the motor protein term. The blue circle marks the proteins involved in IDA assembly. (E-F) Co-IP assays using DNAH12 antibody (E) or DNALI1 antibody (F) showed strong and specific interactions among DNAH12, DNALI1, and DNAH1. (G)

Predicted DNAH12-DNALI1 interaction details by the HDOCK server. The top three of DNALI1 possible binding conformations were shown in types 1-3. (H-J) Construction of mouse-mCherry-DNAH12-Stem (H) and mouse-GFP-DNALI1-full length (FL) (I) plasmids, and the validation of the interaction between the Stem domain of DNAH12 and full-length of DNALI1 (J).

Figure 7. DNAH12 facilitates the recruitment of DNALI1 and DNAH1 to flagella, but not cilia (A) Immunoblotting of testis lysate from *Dnah12+/+, DNAH12+/-* and *Dnah12-/-* mice using DNAH12, DNAH1, and DNALI1 antibodies. DNAI1 (a component of ODAs) and α-Tubulin were used as the loading controls. (B-C) Immunofluorescence assays of α-Tubulin and DNAH1 (B), or DNALI1 (C) antibodies on round spermatids from adult *Dnah12+/−* and *Dnah12−/−* mice. Scale bars, 10 μm. (D-E) Immunofluorescence assays of α-Tubulin and DNAH1 (D), or DNALI1 (E) antibodies on elongated spermatids from adult *Dnah12+/−* and *Dnah12−/−* mice. Scale bars, 10 μm. (F-G) Immunoblotting of oviduct (F) or trachea (G) lysate from *Dnah12+/+, DNAH12+/-* and *Dnah12-/-* mice using DNAH12, DNAH1, and DNALI1 antibodies. GAPDH was used as the loading control. (H-I) Immunofluorescence assays of the oviduct (H) or trachea (I) sections with α-Tubulin and DNAH1 antibodies. Scale bars, 50 μm. (J-K) Immunofluorescence assays of the oviduct (J) or trachea (K) sections with α-Tubulin and DNALI1 antibodies. Scale bars, 50 μm. (L) Pedigree of a family with 3 infertile males, P12 (III-2), P13 (III:6), and P14 (III:8); MT: the *DNAH1* mutation c.11275C>T; p.Arg3759Cys. (M-O) Representative images of spermatozoa from a fertile control and P13 co-stained by α-Tubulin antibody and DNAH1 (M), DNALI1(N), or DNAH12 (O) antibodies, respectively. Scale bars,10 μm.

Figure 8. DNAH12 may interact with RS head associated proteins to regulate CP stability (A) Co-IP assays of the potential interactions of DNAH12 and RS head proteins RSPH1, RPSH9, or DNAJB13. DNAI2 antibody was used as an negative control. (B) Interaction analyses of DNAH12 and RS head proteins RSPH1, RPSH9, or DNAJB13 by AlphaFold3. (C) Immunoblotting of testis lysate from *Dnah12+/+* and *Dnah12-/-* mice using RSPH1, RPSH9, DNAJB13 antibodies. DNAI2 (a component of ODAs), α-Tubulin and β -Tubulin were used as the loading controls. (D) Immunofluorescence assays of α-Tubulin and RSPH1 antibodies on sperm collected from *Dnah12+/−* and *Dnah12−/−* mice caudal epididymis. Scale bars, 10 μm. (E) Representative two-cell embryos and blastocysts of *Dnah12+/+,* and *Dnah12-/* male mice after intracytoplasmic sperm injection. Scale bar, 100 μm. (F) Percentages of 2-cell-stage embryos of *Dnah12+/+* and *Dnah12-/-* groups. (G) Percentages of blastocyst-stage embryos of *Dnah12+/+* and *Dnah12-/-* groups. (H) Schematic diagram showing the proposed function of DNAH12 in sperm flagellar development in humans and mice. DNAH12 is essential for recruiting DNALI1 and DNAH1 to sperm flagella and maintaining the proper axonemal arrangement, especially IDAs and CP structures. Loss of DNAH12 causes the failure of DNALI1 and DNAH1 to be recruited to sperm flagella and results in abnormal sperm morphology with compromised axoneme organization, causing male infertility. ODA, outer dynein arm; IDA, inner dynein arm; N-DRC, nexin-dynein regulatory complex; CP, central pair of microtubules; MTBD, microtubule-binding domain; RS, radial spoke.

Table 1. Detailed information of bi-allelic *DNAH12* variants identified in infertile men

	Genomic position on chr3	cDNA Change [a]	Mutation type	Genetic pattern	SIFT	Polyphen2	fathmm-MKL	CADD	1000 Genomes	gnomAD
Family 1	57,489,883	MT1: c.944_945del	frameshift substitution	AR	NA	NA	NA	NA	0	0.00002844
Family 2	57,528,434	MT2: c.164A>G	nonsynonymous SNV [b]	Male-limited AR	T	D	D	D	0.000599042	0.0004
Family 3	57,447,323	MT3: c.2560A>G	nonsynonymous SNV	AR	T	D	D	D	0.000599042	0.0002
P9	57,391,670	MT4: c.6229C>T	stop-gain	AR	NA	NA	D	NA	0	0.00002714
P10	57,445,368	MT5: c.2813T>C	nonsynonymous SNV	AR	D	D	D	D	0.000199681	0.00003383
P11	57,394,265	MT6: c. 5964-3A>G	splicing	AR	NA	NA	NA	NA	0.000199681	0.0002
	57,445,501	MT7: c.2680C>T	nonsynonymous SNV		D	D	D	D	0	0.000007178

a. The accession number of human *DNAH12* is ENST00000351747.2. b. Abbreviations are listed as follows: SNV, single nucleotide variants; NA, not assessed; T,

tolerated; D, deleterious. SIFT, Sorting Intolerant From Tolerant; PolyPhen-2, Polymorphism Phenotype v2; FATHMM-MKL, Functional Analysis Through Hidden

Markov Models-Multiple Kernel Learning; CADD, Combined Annotation-Dependent Depletion; and a mutation is predicted deleterious if the CADD Phred score > 20

in CADD. gnomAD, Genome Aggregation Database.

Table 2. Clinical characteristics of the infertile patients with bi-allelic *DNAH12* mutations

	Reference values [a]	Fertile controls [b]	IV-1(P1)	Family1 IV-2(P2)	IV-4(P3)	Family2 IV-5(P4)	IV-7(P5)	Family3 IV-4(P8)	P9	P10	P11
cDNA mutation	–	–	MT1/MT1	MT1/MT1	MT1/MT1	MT2/MT2	MT2/MT2	MT3/MT3	MT4/MT4	MT5/MT5	MT6/MT7
Age (y) [c]	–	–	58	49	42	40	38	32	33	40	31
Semen parameters											
Semen volume (ml)	>1.5	–	ND	1.5±0.5	2.8±0.3	4.0	3.0	3.0	3.2	3.1	2.4
Sperm concentration (10^6/ml)	>15	–	ND	58±28.5	50.0±15.0	6.0	8.0	2.8	24.7	83.1	57.1
Semen pH	Alkaline	–	ND	Alkaline	Alkaline	Alkaline	Alkaline	Alkaline	Alkaline	Alkaline	Alkaline
Motile sperm (%)	>40	–	ND	29.3±9.0	7.5±7.5	14.0	20.0	20	15.1	17.4	7.4
Progressively motile sperm (%)	>32	–	ND	13.3±10.3	16.7±17	5.0	8.0	5	13.4	16.8	24
Sperm morphology											
Normal (%)	–	76.5±2.0	ND	ND	19.9 [d]	ND	21.7	17.0	14.0	17.4	7.4
Abnormal head (%)	–	7.0±1.9	ND	ND	16.4	ND	5.4	6.0	3.5	16.4	20.0
Abnormal tail (%)	–	12.4±3.4	ND	ND	43.8	ND	41.4	44.0	42.4	37.4	16.3
Abnormal head &tail (%)	–	4.1±1.3	ND	ND	19.9	ND	31.5	33.0	40.2	28.8	56.3
Sperm flagella morphology											
Normal (%)	–	83.1±2.3	ND	ND	26.4	ND	26.6	22.7	17.9	31.7	27.4
Absent (%)	–	2.7±1.1	ND	ND	16.4	ND	7.9	15.8	9.6	16.6	11.6
Short (%)	–	3.6±0.6	ND	ND	16.4	ND	11.3	14.9	20.1	11.7	8.4
Coiled (%)	–	2.8±0.7	ND	ND	17.3	ND	24.6	12.9	11.8	14.6	7.0
Bent (%)	–	6.3±0.6	ND	ND	12.7	ND	15.8	10.0	13.5	7.8	33.5
Irregular caliber (%)	–	1.5±0.2	ND	ND	10.9	ND	13.8	23.8	27.1	17.6	12.1

a. The reference limits are shown according to WHO 6th edition standards. **b.** Reference values are shown by observation of sperm morphology in 3 fertile individuals.

c. "Age" represents age in 2024. ND, not determined. **d.** Values in red colors show obvious declined proportions of normal sperm morphology compared with fertile controls.

Figure 1

Figure 2

A

B

C

D

Figure 3

Figure 4

Figure 5

Figure 6

A

Stem domain   AAA+ ATPase modules

ODA
- DNAH5[Q8TE73]
- DNAH8[Q96JB1]
- DNAH10[Q8IVF4]
- DNAH11[Q96DT5]
- DNAH9[Q9NYC9]
- DNAH17[Q9UFH2]
- DNAH2[Q9P225]

IDA
- DNAH6[Q9C0G6]
- DNAH1[Q9P2D7]
- DNAH3[Q8TD57]
- DNAH12[Q6ZR08]
- DNAH7[Q8WXX0]
- DNAH14[Q0VDD8]

0.1

Legend:
- AAA_6
- DHC_N2
- DHC_N1
- AAA_8
- Dynein_C
- AAA_9
- AAA_7
- MTBD
- AAA_lid_11
- Dynein_heavy
- AAA_5
- Dynein_AAA_lid
- AAA_lid_1
- AAA

B

Testis lysate

DNAH12 antibody   IgG antibody

Intensity / m/z

IP-MS → Potential DNAH12 Interactors

C Molecular function

Motor protein
Actin-binding
Myosin
RNA-binding
Helicase
Calmodulin-binding
Muscle protein
Actin capping
Initiation factor
Kinase

GeneRatio

Count: 10, 20, 30, 40
-log10(PValue)

D

- IDA assembly
- Microtubule-based Movement
- Cytoskeleton-dependent intracellular transport
- Vesicle cytoskeletal trafficking

Kif7, Kif3a, Klc2, Kif21b, Kif5b, Myo7b, Kif5a, Kif5c, Myo6, Kif21a, Myo1d, Dnah12, Dnah3, Dnah1, Dnali1, Dnah5, Dnah2, Kif15, Kif9, Kif2c, Kif4

Myh7, Myh1, Myh3, Myh7b, Myl3, Myh8, Myl4

E

Input   IgG   IP-DNAH12

Dnah12+/+ Dnah12-/- Dnah12+/+ Dnah12-/- Dnah12+/+ Dnah12-/-

IB:
- DNAH12 — 356 kDa
- DNAH1 — 487 kDa
- DNALI1 — 30 kDa

Testis lysate

F

Input   IgG   IP-DNALI1

Dnah12+/+ Dnah12-/- Dnah12+/+ Dnah12-/- Dnah12+/+ Dnah12-/-

IB:
- DNAH12 — 356 kDa
- DNAH1 — 487 kDa
- DNALI1 — 30 kDa

Testis lysate

G

DNALI1

Stem domain of DNAH12

top three of DNALI1 possible binding conformations

- DNAH12
- Type1
- Type2
- Type3

H

1-1212aa

mCherry   m-DNAH12-Stem

I

258aa

GFP   m-DNALI1-FL

J

Input   IP-mCherry

mCherry-DNAH12-Stem - + - +
GFP-DNALI1-FL + + + +

IB:
- mCherry — 169 kDa
- GFP — 57 kDa

Figure 7

A Testis lysate

B α-Tubulin / DNAH1 / Merge — Dnah12+/+, Dnah12-/-

C α-Tubulin / DNALI1 / Merge — Dnah12+/+, Dnah12-/-

D α-Tubulin / DNAH1 / Merge — Dnah12+/+, Dnah12-/-

E α-Tubulin / DNALI1 / Merge — Dnah12+/+, Dnah12-/-

F Oviduct lysate

G Trachea lysate

H Oviduct — Dnah12+/+, Dnah12-/- — α-Tubulin / DNAH1 / Hoechst

I Trachea — Dnah12+/+, Dnah12-/- — α-Tubulin / DNAH1 / Hoechst

J Oviduct — Dnah12+/+, Dnah12-/- — α-Tubulin / DNALI1 / Hoechst

K Trachea — Dnah12+/+, Dnah12-/- — α-Tubulin / DNALI1 / Hoechst

L DNAH1 MT: c.11275C>T; p.Arg3759Cys — For WES

M α-Tubulin / DNAH1 / Merge / Bright field — Control, P13

N α-Tubulin / DNALI1 / Merge / Bright field — Control, P13

O α-Tubulin / DNAH12 / Merge / Bright field — Control, P13

Figure 8

A

IDA | DNAH12 — 356 kDa

RS head | RSPH1 — 34 kDa | RSPH9 — 31 kDa | DNAJB13 — 36 kDa

ODA | DNAI2 — 71 kDa

Input | IgG | DNAH12-IP

Testis lysate

B

Stem domain — RSPH1 — DNAH12

Stem domain — RSPH9 — DNAH12

Stem domain — DNAJB13 — DNAH12

C

Dnah12+/+ | *Dnah12*−/−

RSPH1 — 34 kDa
RSPH9 — 31 kDa
DNAJB13 — 36 kDa
DNAI2 — 71 kDa
β-Tubulin — 50 kDa
α-Tubulin — 50 kDa

Testis lysate

D

α-Tubulin | RSPH1 | Merge

Dnah12+/+

Dnah12−/−

E

Dnah12+/+ | *Dnah12*−/−

2-cell

Blastocyst

F

2-cell percentage (%)

ns

Dnah12+/+ | *Dnah12*−/−

G

Blastocys percentage (%)

ns

Dnah12+/+ | *Dnah12*−/−

H

Control :

DNALI1 — Mutual dependent

DNAH12 — Recruit

DNAH1

ODA — N-DRC — IDA

CP

Mutant :

DNALI1

Loss of DNAH12 — Fail to recruit — Mutual dependent — Fail to locate on flagella

DNAH1

Impaired IDA

ODA — N-DRC — Impaired IDA

CP loss

Asthenoteratozoospermia | **Male infertility**

Legends

DNAH12

DNAH1

MTBD domain

DNALI1

RS

CP