1

Cell type-specific immune regulation under symbiosis in a facultatively symbiotic coral

Maria Valadez-Ingersoll[1], Hanny E. Rivera[1, 2], JK Da-Anoy[1], Matthew R. Kanke[3], Kelly Gomez-Campo[4, 5], M. Isabel Martinez-Rugerio[4, 5], Julian Kwan[6], Ryan Hekman[6], Andrew Emili[1, 6, 7], Thomas D. Gilmore[1], Sarah W. Davies[1,*]

[1]Boston University, Department of Biology; Boston, MA, US
[2]Gingko Bioworks, Boston, MA, USA
[3]Amgen Research, Research Bioimics, South San Francisco, CA, USA
[4]Pennsylvania State University, Department of Biology, State College, PA, USA
[5]Helmholtz Institute for Functional Marine Biodiversity (HIFMB), Ammerländer Heerstraße 231, 26129 Oldenburg, Germany
[6]Department of Biochemistry, Boston University Chobanian & Avedisian School of Medicine, Boston, MA, USA
[7]Division of Oncological Sciences, Knight Cancer Institute, Oregon Health & Science University, Portland, OR, USA

*Corresponding author: Sarah Davies, Boston University, Department of Biology, 5 Cummington Mall, Boston, MA 02215; Phone: 617-353-8980; email: daviessw@bu.edu

Competing interests: The authors declare no competing interests.

ABSTRACT

Many cnidarians host single-celled algae within gastrodermal cells, yielding a mutually beneficial exchange of nutrients between host and symbiont, and dysbiosis can lead to host mortality. Previous research has uncovered symbiosis tradeoffs, including suppression of immune pathways in cnidarians hosting intracellular algae and correlations between symbiotic state and pathogen susceptibility. Here, we used a multiomic approach to characterize symbiotic states of the facultatively symbiotic coral *Oculina arbuscula* by generating genotype-controlled fragments of symbiotic and aposymbiotic tissue. 16S metabarcoding showed no difference in bacterial communities between symbiotic states. Whole-organism proteomics revealed differential abundance of proteins related to immunity, confirming immune suppression during symbiosis. Finally, single-cell RNAseq identified diverse cell clusters within seven cell types across symbiotic states. Specifically, the gastrodermal cell clusters containing algal-hosting cells from symbiotic tissue had higher expression of nitrogen cycling and sugar transport genes than aposymbiotic gastrodermal cells. Furthermore, differential enrichment of immune system gene pathways and lower expression of genes involved in immune regulation were observed in these gastrodermal cells from symbiotic tissue. However, no differences in immune gene expression in the immune cell cluster were observed between symbiotic states. This work reveals a compartmentalization of immune system regulation in specific gastrodermal cells in symbiosis, which may limit symbiosis tradeoffs by simultaneously dampening immunity in algal-hosting cells while maintaining general organismal immunity.

KEY WORDS: coral, symbiosis, immunity, single-cell RNA-sequencing, proteomics

INTRODUCTION

Symbioses exist across the Tree of Life along a spectrum from parasitisms – in which one

symbiotic partner benefits at the cost of the other partner – to mutualisms – in which both partners

benefit from the relationship (1). Endosymbioses, symbiotic relationships where one organism

resides within another, pose immunity challenges to hosts as they must differentiate between self,

mutualistic symbionts, and pathogens (2,3). In mutualisms, the host immune system is often

regulated in order to allow for the existence of the endosymbiont (4), whereas in parasitisms the

endosymbiont can evade detection by the host's immune system (5). Additionally, in mutualisms,

hosts must tightly regulate the density of their symbionts in order to prevent overcrowding, which

can lead to a shift from mutualism to parasitism (6).

Reef-building corals participate in a mutually beneficial symbiosis with algae that reside

inside their gastrodermal cells, wherein the algae carry out photosynthesis and translocate fixed

carbon sugars to the host (7,8). In return, the host provides CO_2, as well as nitrogen and phosphorus

species to the symbiont (9,10). Translocation of nitrogen from the host to the symbiont has been

proposed as a mechanism to promote nutrient exchange while controlling algal proliferation via

nitrogen limitation to prevent overcrowding (11). As symbiont densities increase, the greater

nutrient requirements of the algae lead to resource competition with the host, thereby destabilizing

the mutualism (12,13). In support of this model, whole-organism RNA-sequencing has revealed

differences in the expression of genes involved in sugar transport and nitrogen cycling across

cnidarians in and out of symbiosis (14), and enrichment of nitrogen deprivation genes in algae

exhibiting greater cell densities when associated with the sea anemone *Exaiptasia pallida* (11).

These whole organism studies reflect the importance of carbon and nitrogen transport between

symbiotic partners in the maintenance of this symbiosis.

In stony corals, environmental change can lead to loss of the algal symbiont in a process

termed coral bleaching (15). If prolonged, this dysbiosis can lead to host mortality. Since this

symbiotic relationship in corals is often obligate, it is challenging to disentangle the effects of

dysbiosis from nutritional losses (16). Accordingly, facultatively symbiotic cnidarian models,

which are viable and naturally occur in both symbiotic and aposymbiotic states, have been useful

for understanding mechanisms that promote symbiosis establishment, maintenance, and loss, from

both nutritional and immune system perspectives (14,16–18). In contrast to obligate symbioses,

facultatively symbiotic hosts can exist in an aposymbiotic state (with few or no algal symbionts)

and can buffer nutrient loss through heterotrophy. This facultative nature allows for studies of the

influence of symbiosis on host molecular function in the absence of starvation stress experienced

by bleached obligate corals (19). Importantly, the aposymbiotic state can be induced in

facultatively symbiotic cnidarians in the laboratory through chemical (with menthol) and heat

treatments, additionally allowing for the ability to control for genetic background. The sea

anemone *E. pallida* is a facultatively symbiotic cnidarian that has served as a useful model for

understanding how the host immune system is modulated in the presence of the algal symbiont

(20,21). Additionally, facultatively symbiotic stony corals, such as *O. arbuscula* (14) and

Astrangia poculata (22), are emerging models for studying these processes in calcifying

cnidarians.

Research using these facultatively symbiotic models has identified pathways within the

host immune system as key for the cnidarian-algal symbiosis. For example, activation of the

transcription factor NF-κB pathway, which has been implicated in immunity across metazoans,

has been shown to be downregulated in symbiotic compared to aposymbiotic hosts (14,20).

Additional evidence suggests that the attenuation of immune pathways associated with symbiosis

corresponds to reduced organismal immunity (23). However, this dampening of immunity is not

always correlated with increased susceptibility to pathogen challenge (24,25). For example, recent

work has shown that the NF-κB pathway can be independently modulated by cnidarians capable

of symbiosis under energetic limitation (26). Additionally, many antioxidant and oxidative

response pathways are constitutively higher in symbiotic compared to aposymbiotic cnidarians

(27). Taken together, a nuanced picture of the symbiotic cnidarian immune system has emerged,

with the question remaining of how the immune system challenge of endosymbiosis is mitigated

without compromising whole-organism immunity.

Herein, we have used a multiomic approach to investigate symbiosis-immunity tradeoffs

in the facultatively symbiotic coral *O. arbuscula*. Whole-organism proteomics confirms broad

immune dampening in symbiosis. Single-cell RNA-seq of genotype-controlled fragments was

used to profile *O. arbuscula* cellular diversity under different symbiotic states. We provide

evidence for how tradeoffs of endosymbiosis and immunity manifest by gene expression rewiring

across different cell states. Strikingly, we found gene expression differences in immune pathways

occur in gastrodermal cells capable of hosting symbiotic algae. These results suggest that cell type-

specific dampening of immunity explains how some symbiotic organisms balance the immune

tradeoffs of symbiosis with organismal immunity.

MATERIALS AND METHODS

Detailed information on sample preparations, bioinformatic analyses, and R packages can be found

in the **Supplemental Materials**.

Coral husbandry and manipulation of symbiotic state

Symbiotic (in symbiosis with *Brevioulum psygmophilum*) (28) colonies from seven genetic

backgrounds (genets A-G) of *Oculina arbuscula* were collected at Radio Island Jetty, North

Carolina and have been maintained at Boston University (BU) in common garden aquaria since

May, 2018. Aposymbiotic branches were generated via menthol bleaching. Aposymbiotic status

was confirmed by a lack of symbiont autofluorescence under fluorescence microscopy (Leica

M165 FC), and aposymbiotic branches were transferred back to common garden aquaria and

maintained for at least 2 months of recovery prior to physiological and multiomic profiling.

Tissue removal for symbiont cell quantification

To measure symbiont cell densities in symbiotic and aposymbiotic *O. arbuscula*, small branches

were fragmented from symbiotic (N=10) and aposymbiotic (N=10) samples from five genets.

Tissue was removed via airbrushing, homogenized, and symbiont counts were normalized to

skeletal surface area using an Einscan-SE scanner. Statistical differences between symbiotic and

aposymbiotic fragments were calculated using the Kruskal-Wallis rank sum test, as assumptions

of normality were not met.

Coral spectroscopic determinations

Light absorption capacities of symbiotic and aposymbiotic *O. arbuscula* were compared using

reflectance and absorptance (29–31). Coral reflectance (R), the fraction of light reflected, was

measured between 400-750 nm in intact fragments using a miniature spectroradiometer (Flame-T-

UV-Vis, Ocean Optics Inc.). Reflectance was expressed as the ratio of the measurement from the

tissue surface relative to the reflectance of a bleached *O. arbuscula* skeleton. Coral absorptance

(A), which describes the fraction of incident light absorbed by coral tissue, was calculated from

the reflectance spectra as $A = 1 - R$ (29,31). The absorptance peak of chlorophyll a (Chl a) at 675

nm was calculated as $A_{675} = 1 - R_{675}$, assuming that transmission through the skeleton is negligible.

Bacterial community profiling

To identify bacterial communities associated with symbiotic and aposymbiotic fragments, one

symbiotic and one aposymbiotic fragment from five genotypes were flash frozen, and a subsample

was preserved in ethanol for 16S metabarcoding (N=10). Metabarcoding libraries were generated

using a series of PCR amplifications for the V4/V5 region of the bacterial 16S rRNA gene (32,33).

Five negative controls were prepared and used to remove contamination. Samples were sequenced

on an Illumina Miseq at Tufts University Core Facility (paired-end 250 bp).

16S read processing inferred 661 amplicon sequence variants (ASVs) with assigned

taxonomy across samples. Bacterial communities across symbiotic states were compared via alpha

diversity (Shannon index, Simpson's index, ASV richness, and evenness) and beta diversity

(PCoA with Bray–Curtis dissimilarities). Alpha diversity metrics were compared using linear

mixed effects models with symbiotic state as the predictor and a random effect of genotype. The

effect of symbiotic state on beta diversity was assessed using the function *betadisper* (vegan

package; v2.6-4 (34)). DESeq2 v1.40.2 (35) explored differentially abundant ASVs to ensure that

subtle differences in specific taxa were not overlooked.

Proteomic profiling

Mass spectroscopy (MS) was used to identify differentially enriched proteins from total protein

isolated from fragments of four symbiotic genets and three aposymbiotic genets. Total protein was

isolated as described previously (36) and was stored at -80 °C before analysis by MS.

For MS, tryptic peptide mixtures were analyzed by nano-scale high-performance liquid

chromatography (Proxeon EASY-Nano system, Thermo Fisher Scientific) coupled with online

nanoelectrospray ionization tandem MS (Q-Exactive HF-X mass spectrometer; Thermo Fisher

Scientific) (37). For protein identification and analysis, data files were searched using the

workflow of MaxQuant version 2.4 (http://www.maxquant.org/) under standard settings using the

Astrangia poculata genome (22). A false discovery rate (FDR) threshold of 1% was used to filter

candidate peptides and protein identifications. Following data filtration, normalization, and

clustering with Omics Notebook (38), 2,191 proteins were retained. Differential abundance

analysis of proteomic profiles between aposymbiotic and symbiotic samples was based on a

moderated t-test (38) in which proteins were considered differentially abundant if they had a

Bonferroni adjusted p-value < 0.1. Raw intensity counts were *rlog*-normalized in DESeq2 (35) and

visualized using a Principal Component Analysis (PCA) using vegan v2.6-4 (34). Effects of

symbiotic state and genet were assessed using a PERMANOVA with the adonis2 function in vegan

v2.6-4 (34). Predicted *O. arbuscula* peptides were searched against the human proteome v.11.5

from the STRING v.11 database (39) with an e-value cut-off of $1x10^{-5}$. Protein–protein interactions

of select differentially expressed proteins (FDR < 0.1) were retrieved from the STRING v.11

database (39). Interaction networks were visualized using Cytoscape v.3.7.2 (40).

Single-cell RNA sequencing

To create single-cell libraries, live cells from one symbiotic and one aposymbiotic branch of

genotype F of *O. arbuscula* were sampled for single-cell isolation and 10X cDNA sequencing

(41). A detailed protocol for cell isolation can be found on protocols.io at DOI:

dx.doi.org/10.17504/protocols.io.rm7vzkx72vx1/v1 and is also described in the **Supplemental**

Materials. To minimize cell death, cells were not sorted prior to processing. Cell isolation samples

were analyzed by BU's Single Cell Sequencing Core Facility. The symbiotic sample had a

concentration of 2,625 cells/µl and a viability of 81.6%. The aposymbiotic sample had a

concentration of 3,313 cells/µl and a viability of 86.4%. These values are largely in line with other

single cell preparation methods reported in the literature (e.g., viability threshold of 80%,

Nematostella vectensis (42,43)).

Libraries were generated following the 10X Genomics Chromium Single Cell 3'v3

protocol, and quality was assessed via a Bioanalyzer. Samples were pooled in equimolar

concentration and sequenced on an Illumina NextSeq 2000 (P3 100 kit) with a target of 50,000

reads per cell. CellRanger (v7.2.0 (44)) processed reads, which were aligned to concatenated

genomes of *Astrangia poculata* (host) (22) and the algal symbiont *Breviolum psygmophilum* (14).

Reads aligning confidently to the host and symbiont references were used to generate CellRanger

output files. Default CellRanger cell calling algorithms were employed. The *A. poculata* genome

was used as a reference for mapping because there is currently no published genome for *O.*

arbuscula, and because alignment rates to a *de novo* assembled transcriptome from the 10X data

generated here and to a previously published *O. arbuscula* transcriptome were of lower quality

(**Supplementary Dataset 1**) (14,19,45). In addition, it is worth noting that species in the genera

Astrangia and *Oculina* are closely related phylogenetically (19), and previous work has found that

the *O. arbuscula* population from Radio Island, NC (source for this study) is more genetically

similar to the *A. poculata* population from Woods Hole, MA (coral genome source) than it is to

other, more southern *Oculina* populations (45).

Following alignment, host reads from symbiotic and aposymbiotic samples were analyzed

using Seurat (v.5.0.2) (46). Datasets were cleaned (genes expressed in fewer than three cells were

discarded, as were cells expressing fewer than 200 genes or greater than 3,000 genes), normalized,

and scaled, and a PCA reduction was performed. No duplicated barcodes were detected.

Mitochondrial reads could not be removed, as the reference genomes and transcriptome do not

have annotated mitochondrial genomes. Datasets were integrated using the Canonical Correlation

Analysis (CCA) Integration method on the PCA reduction on the top 2,000 most variable genes

(nearest neighbors parameters). Cell clusters were identified using 30 dimensions (resolution: 0.5)

and visualized using Uniform Manifold Approximation and Projection (UMAP). Marker genes for

each cluster were identified using FindAllMarkers (Wilcoxin Rank Sum Test; $\log_2$foldchange

threshold: 0.5) and cell types were informed by gene annotations and marker gene comparisons

from other cnidarian single-cell datasets (e.g., *S. pistallata* and *Xenia* sp.) (**Supplementary**

Dataset 2) (47,48). Expression patterns of top marker genes were assigned to cell clusters using

violin plots, bubble plots, and visualization of gene expression in individual cells within the

UMAP. 28 cell clusters across 7 cell types were identified in the UMAP. Broad transcriptomic

differences between cell clusters were assessed by identifying the top 10 genes enriched in each

cluster compared to all the other clusters.

To identify cells containing algal symbionts, we identified cells in which over 50% of the

total reads corresponded to *B. psygmophilum* genes, and then visualized these cells on the UMAP.

These cells were then removed, as the low percentage of host reads in these cells confounded

downstream analyses.

To determine how symbiosis alters enrichment of gene pathways at the whole-organism

level, and how this differential enrichment is reflected in specific cell types, we performed Mann–

Whitney U Gene Ontology (GO) term enrichment analysis on all genes from the full single-cell

dataset (49). Significantly enriched GO terms within the 'Biological Process' (BP) GO division

were defined with a false-discovery rate (FDR) of <0.1. Significantly over- and under-represented

GO terms relating explicitly to immunity, antigen signaling, or the NF-κB pathway were identified

(**Table S1**). Cells in which over 7% of the total reads corresponded to genes annotated with the

GO terms of interest were visualized on the UMAP of all cells from symbiotic and aposymbiotic

O. arbuscula samples.

To compare expression profiles of specific cell states between symbiotic and aposymbiotic

samples, each cell cluster was independently reclustered, and the effect of symbiotic state on gene

expression in each cell state was analyzed. First, the cell cluster of interest (*e.g.,* Immune Cell,

Gastrodermis 1) was subsetted and new variable features were identified. Data were rescaled, and

dimensionality reduction was re-run (first 30 dimensions). Subclusters were identified with

resolutions between 0.25-0.5 (**Table S2**). The distribution of cells across subclusters between

symbiotic states was visualized using UMAPs. Differentially expressed genes between symbiotic

states in all cell clusters were identified using DESeq2 within the FindMarkers function.

Additionally, marker genes for each subcluster were identified. To highlight differences in the

gene expression of the Immune Cell states, the top 20 genes enriched in each Immune Cell

subcluster were identified and plotted (unannotated genes were removed).

We used Monocle 3 (v.1.3.7) (50) to determine cell trajectories and outcomes for all

gastrodermal cells from symbiotic and aposymbiotic samples. We plotted the UMAPs of the

symbiotic and aposymbiotic subsets of Gastrodermis clusters 1-4 and overlaid trajectory graphs,

noting nodes and outcomes.

To compare expression of genes involved in sugar transport and nutrient cycling in

gastrodermal cells between symbiotic aposymbiotic states, the expression of these genes in the

reclustered Gastrodermis 1 and Gastrodermis 2 cells was analyzed. First, normalized expression

of genes involved in nitrogen cycling/symbiont density control (Glutamate Dehydrogenase DHE4

and Glutamate Synthase [NADH] GLSN) and sugar transport (Apolipophorin APOB) (14) were

compared across all cell clusters using violin plots (function VlnPlt in the Seurat package (46)).

FeaturePlot (46) was then used to visualize co-expression of these genes across the subclusters of

Gastrodermis 1 and Gastrodermis 2 cells from the symbiotic sample (scaling per-cell mean

expression to a maximum value of 10). To determine whether the coexpression of genes observed

in the symbiotic gastrodermal cells is seen in other cell types, we also visualized coexpression of

DHE4, APOB, and GLSN across Epidermis 1 and Neuron 1 from symbiotic tissue (scaling per-

cell mean expression to a maximum value of 10).

To identify genes driving separation between symbiotic states within each cell cluster, the

distribution of cells from a particular cluster was visualized along the first 10 PCs of the PCA

assay used to define the UMAP for that cluster. An analysis of variance determined whether the

cells were separated by symbiotic state (symbiotic vs. aposymbiotic) along each PC. In

Gastrodermis 1 and Gastrodermis 2 clusters, the top 20 genes driving the distribution of cells along

each significant PC (10 genes driving each PC in the positive direction and 10 genes driving each

PC in the negative direction) were selected, and unannotated genes were removed. In Gastrodermis

1, PCs 1, 3-8, and 10 were selected for a total of 96 genes. In Gastrodermis 2, PCs 1, 2, 5, 7, and

8 were selected for a total of 59 genes. Log$_2$foldchange values from DESeq2 with accompanying

p-adjusted values (<0.05) for each gene were plotted in a bar graph. Genes annotated with immune-

function related GO terms were noted, as were genes annotated with the Clusters of Orthologous

Genes (COG) annotation 'posttranslational modification, protein turnover, and chaperones.'

Finally, to compare *O. arbuscula* cells potentially hosting algal symbionts with algal-

hosting cells from an obligate coral species (*Xenia* sp. (48)), we identified orthologous genes

between *A. poculata* (the mapping reference in this study) and *Xenia* using the Broccoli algorithm

using 13 cnidarian proteomes with default parameters (51). We re-ran Seurat normalization, PCA,

and UMAP clustering on the pre-integrated *Xenia* dataset from non-regeneration samples (48) to

identify 16 clusters (resolution of 0.18 and 30 dimension), including the previously characterized

algal-hosting cell cluster and the other two gastrodermal cell clusters. We identified differentially

expressed genes between the algal-hosting cells and the gastrodermal cells (*p*-adjusted <0.05 from

the FindMarkers function, using both gastrodermal cell clusters as the comparison to the algal-

hosting cell cluster), and subsetted these genes for those that had orthologs to *A. poculata*.

Additionally, we identified differentially expressed genes between algal-hosting and non-algal

hosting *O. arbuscula* cells in two ways: 1) from symbiotic *O. arbuscula* cells from Gastrodermis

1/Gastrodermis 2 and aposymbiotic *O. arbuscula* cells from Gastrodermis 1/Gastrodermis 2 (*p*-

adjusted <0.05 from the FindMarkers function), and 2) from symbiotic *O. arbuscula* cells from

Gastrodermis 1/Gastrodermis 2 and from symbiotic *O. arbuscula* cells from Gastrodermis

3/Gastrodermis 4. We subsetted these genes to include only *Xenia* orthologs. We compared which

genes among these three sets of orthologs (*Xenia*-sym-to-gastro, *Oculina*-sym-to-apo, and

Oculina-sym-to-gastro respectively) were differentially expressed using a Venn diagram (52),

creating six groups of differentially expressed orthologs (DEOs). We performed Fisher Exact Tests

(DEO presence/absence) (49) to identify enrichment differences across these groups of DEOs in

the BP GO division.

RESULTS

For the following experiments, genetically controlled symbiotic and aposymbiotic fragments of *O.*

arbuscula were used. Symbiotic fragments had symbiont densities that were 400-fold greater

(13,678 +/- 6,204 cells/cm^2) than those of aposymbiotic fragments (30 +/- 43 cells/cm^2) (p<0.001)

(**Fig. 1a**). These fragments were then used to generate light-scattering profiles, 16S metabarcoding

libraries, tissue proteomics, and single-cell profiles.

Light absorption is higher in symbiotic *O. arbuscula*

Differences in symbiont cell densities induced significant differences in *in vivo* coral light

absorption. Reflectance spectra measurements and absorption properties of intact corals showed

statistically significant pigment content differences between symbiotic and aposymbiotic *O.*

arbuscula. Average reflectance in the PAR region (400–700 nm) of symbiotic fragments was 22%

+/- 4%, whereas aposymbiotic coral reflectance was 77% +/- 3% (p < 0.01) (**Fig. 1b**). In symbiotic

fragments, the absorptance value at the peak for Chlorophyll a (A_{675}) was 85% +/- 4%, compared

to 24% +/- 2% in aposymbiotic samples (p<0.0001) (**Fig. 1c**). Absorptance describes the relative

amount of solar energy/incident light that can potentially be used in photosynthesis for organic

carbon fixation.

Bacterial communities do not differ across symbiotic states

We profiled the bacterial communities of aposymbiotic and symbiotic fragments from 5 genets

(N=10). A total of 208,330 sequences were acquired with a mean depth of coverage of 20,833 +/-

16050 per sample (**Table S3**). Sample rarefication yielded a total of 8,298 reads per sample with

a total of 661 ASVs identified across samples. No differences in bacterial communities were

observed between symbiotic and aposymbiotic fragments (**Fig. 1d**), and both states were

dominated by Proteobacteria and Bacteroidota (**Fig. S1**). Additionally, no differences in alpha

diversity were observed between symbiotic states regardless of the metric tested, nor were

differences in beta diversity by symbiotic state detected (**Fig. S2a-d**). DESeq2 tested for

differentially abundant ASVs across symbiotic states, while modeling genetic background, but no

ASVs were identified.

The proteome is affected by symbiotic state in *O. arbuscula*

To compare proteomes of symbiotic and aposymbiotic *O. arbuscula*, we performed mass

spectrometry on whole-cell lysates from coral fragments. We identified a total of 2,343 proteins

in *O. arbuscula*, 2,191 of which were retained following filtering. A PCA showed that samples

clustered by symbiotic state (p_{state}<0.1) and genet (p_{genet}<0.05) (**Fig. 2a**). A total of 187 proteins

were differentially enriched across symbiotic states (*p*-adjusted<0.01: 105 enriched in symbiotic,

82 enriched in aposymbiotic *O. arbuscula*; **Figs. 2b-c**).

Differentially enriched proteins (DEPs) were assigned functional categories based on a

homology search of each DEP sequence in the UniProtKB database (evalue < 1×10^{-5}). DEPs

involved in Immune Response, Repair and Stress Response, and Metabolic Processes were

identified (**Fig. 2d**)

Single-cell RNA-seq identifies cell states shared between symbiotic and aposymbiotic *O.*

arbuscula

To profile gene expression in different cell types and states from symbiotic and aposymbiotic *O.*

arbuscula, we used the 10X single-cell RNA-sequencing (scRNA-seq) platform. We captured a

total of 6,992 cells (3,158 cells from the symbiotic sample; 3,834 cells from the aposymbiotic

sample), with an average of 157,534.5 reads per cell, 1,121 median UMI counts per cell, and 651.5

median genes per cell. The average number of reads was 540,436,426; the average percentage of

valid barcodes was 94.85%; the average percentage of valid UMIs was 99.90%; the average

sequencing saturation was 89.75%. Further quality control data for both samples are summarized

in **Supplementary Dataset 1**.

Clustering of the integrated 10X gene expression datasets from symbiotic and

aposymbiotic samples resulted in the identification of 28 transcriptomically distinct cell clusters

across seven broad cell types (**Fig. 3a**). Cells from symbiotic and aposymbiotic samples were

represented in all clusters (**Fig. 3a**). Additionally, each cluster defined in the UMAP displayed

distinct transcriptomic programming as exemplified by expression patterns of the ten highest

enriched genes in each cluster (**Fig. 3b**).

Identification of symbiont-hosting cells

We identified 34 cells in which over 50% of the total reads corresponded to *B. psygmophilum*

genes. 33 of these 34 cells belonged to the symbiotic Gastrodermis 2 cluster (**Fig. S3**), indicating

that this cell cluster contained the algal-hosting cells from symbiotic *O. arbuscula*.

Symbiotic gastrodermal cell populations fated for symbiosis function in nutrient exchange

To investigate the cell fates of the gastrodermal cells, we performed trajectory analysis on the four

gastrodermal cell clusters in the symbiotic and aposymbiotic *O. arbuscula* samples separately. In

the symbiotic sample, we identified two fates, one terminating in Gastrodermis 2 where symbiont-

hosting cells were identified, and one terminating in Gastrodermis 3 (**Fig. 4a**). One fate was

identified in the aposymbiotic gastrodermal cells, terminating in Gastrodermis 4 (**Fig. S4**). To

probe gene expression differences between gastrodermal cells from symbiotic compared to

aposymbiotic *O. arbuscula*, we subsetted and reclustered cells from each of the four gastrodermal

states. Four subclusters were revealed within Gastrodermis 1 (**Fig. 4b**), three subclusters were

revealed within Gastrodermis 2 (**Fig. 4c**), two were revealed in Gastrodermis 3 (**Fig. 4d**), and one

was revealed within Gastrodermis 4 (**Fig. S5a**), suggesting additional cell states. Gastrodermis 1

Subcluster 3 and Gastrodermis 2 Subcluster 3 were composed almost entirely of cells from the

symbiotic sample (**Fig. 4b** and **Fig. 4c** respectively). Independent reclustering of all other cell

types was also performed, and the resulting UMAPs displayed varying, though lesser, degrees of

structure by symbiotic state (**Fig. S5b-w**).

To help identify the localization of the gastrodermal cells participating in active symbiotic

nutrient exchange within our dataset, we selected sugar transport and nitrogen cycling genes,

which have been previously shown to have higher expression in symbiotic compared to

aposymbiotic *O. arbuscula* (14). DESeq2 identified differences in gene expression in gastrodermal

cells between symbiotic and aposymbiotic samples, which showed that Gastrodermis 1 cells from

symbiotic *O. arbuscula* have higher expression of genes related to nitrogen cycling (Glutamate

Dehydrogenase DHE4 and Glutamate Synthase [NADH] GLSN) and sugar transport

(Apolipophorin APOB) (p-adjusted <0.05) (**Fig. 4e-g**). Furthermore, DHE4, GLSN, and APOB

were identified as marker genes for Subcluster 3 (indicating that these genes were more highly

expressed inside than they were outside Subcluster 3 (p-adjusted <0.05). Additionally, there was

strong co-expression of these symbiosis-marker genes in Subcluster 3, suggesting that this cell

state is actively transporting fixed carbon sugars while mediating the growth and division of algal

symbionts (**Fig. 4h-j**). There was less co-expression of these genes in Subclusters 1, 2, and 4 of

Gastrodermis 1, suggesting that other gastrodermal cell states perform nitrogen and carbon sugar

processing independently or through distinct pathways. Similar, though weaker, patterns of

coexpression were observed in Gastrodermis 2 (**Fig. S6a-c**). Additionally, there was no or limited

coexpression of DHE4, APOB, and GLSN observed in symbiotic Epidermis 1 and Neuron 1 cells

(**Fig. S6d-i**). The presence of symbiotic-only subclusters within Gastrodermis 1 and 2, coupled

with the co-expression of highly expressed markers for symbiosis within one Gastrodermis 1

subcluster suggest that certain gastrodermal cells states in our dataset are actively hosting

symbionts.

No differences in immune cell gene expression between symbiotic state

To determine whether downregulation of genes associated with the immunity observed in whole-

organism studies of symbiotic compared to aposymbiotic coral (this study and (14)) was observed

in the Immune Cell gene expression, we identified one Immune Cell cluster with high expression

of cell markers for immunity. These immune marker genes included Interferon Regulatory Factors

(IRFs), Activating Transcription Factor 3 (ATF3), and multiple chaperone proteins in cells from

both symbiotic and aposymbiotic tissue (**Fig. 5a, b**). It is worth noting that we did not identify cell

type- or symbiotic state- specific expression of other commonly studied cnidarian immune genes,

including NF-κB, cGAS, and STING (20,53). After subsetting and independently reclustering

these Immune Cells, we identified three cell states with distinct transcriptomic profiles (**Fig. 5c**).

However, cells did not cluster by symbiotic state after reclustering, and no differential expression

between Immune Cells from symbiotic and aposymbiotic samples was observed (*p*-adjusted <0.05;

Fig. 5d).

Immune response genes are differentially regulated in symbiotic Gastrodermis 1 and 2 cells

In agreement with previous bulk studies (14,20,54), our dataset demonstrated that, at the whole-

organismal scale, immune system gene pathways identified through GO term analysis are

differentially regulated under symbiosis. Interestingly, genes belonging to these differentially

enriched immunity GO terms are most highly expressed in gastrodermal cells. In addition,

expression of these genes is higher in gastrodermal cells from aposymbiotic (**Fig. 6a**) compared to

symbiotic *O. arbuscula* (**Fig. 6b**), demonstrating that symbiont-specific immune gene regulation

occurs in gastrodermal cells rather than immune cells. Furthermore, because Gastrodermis 1 and

2 cells clustered by symbiotic state, we identified the factors (genes) driving this pattern in the

Gastrodermis 1 and 2 cell UMAPs. We analyzed the distribution of the cells along each of the first

10 Principal Components (PCs) in the PCA dimensionality reduction used to generate the UMAPs.

Gastrodermis 1 cells exhibited significant separation by symbiotic state along 8 of the first 10 PCs

(*p*-value < 0.05), and Gastrodermis 2 cells showed significant separation along 5 of the first 10

PCs. In Gastrodermis 1, of the 96 annotated genes driving the distribution of the cells along the 8

significant PCs, 33 were differentially expressed (*p*-adjusted <0.05), and 32 of the 96 genes were

annotated with GO terms involved in immunity (**Fig. 6b**). Additionally, 9 genes were annotated

with the COG term 'posttranslational modification, protein turnover, and chaperones' (**Fig. 6b**).

In Gastrodermis 2, 4 of the 59 genes were differentially expressed (*p*-adjusted <0.05), 13 were

annotated with immunity GO terms, and 5 were annotated with the COG term 'posttranslational

modification, protein turnover, and chaperones' (**Fig. 6c**).

In two coral species, gastrodermal cells capable of symbiosis share functions relating to

metabolism

To compare conserved mechanisms by which specific types of gastrodermal cells host algae across

coral species, we analyzed the overlap of differentially expressed orthologs (DEOs) between algal-

hosting cells (Gastrodermis 1 and 2 cells along the same cell fate trajectory) and non-algal-hosting

gastrodermal cells from our facultatively symbiotic coral to algal-hosting cells and other gastrodermal cells from an obligate coral species (*Xenia* sp. (48)). We identified 200 DEOs between *Xenia* algal-hosting and non-algal hosting gastrodermal cells (*Xenia*-sym-to-gastro), 233 DEOs between *O. arbuscula* Gastrodermis 1 and 2 cells from symbiotic and aposymbiotic tissue (*Oculina*-sym-to-apo), and 184 DEOs between *O. arbuscula* Gastrodermis 1 and 2 cells from symbiotic tissue and Gastrodermis 3 and 4 cells from symbiotic tissue (*Oculina*-sym-to-gastro). We performed this analysis in two ways to account for the facultatively symbiotic nature of *O. arbuscula*. First, we analyzed overlapping and unique DEOs between *Xenia*-sym-to-gastro and *O. arbuscula Oculina*-sym-to-apo: 71 DEOs were shared, 129 were unique to *Xenia*-sym-to-gastro, and 162 were unique to *Oculina*-sym-to-apo (**Fig. S7a**). Second, we analyzed overlapping and unique DEOs between *Xenia*-sym-to-gastro and *O. arbuscula Oculina*-sym-to-gastro: 51 DEOs were shared, 149 were unique to *Xenia*-sym-to-gastro, and 133 were unique to *Oculina*-sym-to-gastro (**Fig. S7b**). Using a Fisher Exact Test of BP GO terms, we probed functional enrichment of processes across each DEO group from the Venn diagrams. Across shared and *Oculina*-unique DEO groups, the most common GO terms were involved in metabolism, protein localization, and ribosomal function. Interestingly, the functions relating to cellular stress (e.g., "Cellular Response to DNA Damage Stimulus") and cell cycle regulation (e.g., "Regulation of Cell Cycle") were enriched in *Oculina*-sym-to-gastro-unique DEO set; furthermore, the "Immune Response" GO term was enriched in both *Xenia*-sym-to-gastro-unique DEO sets (**Supplementary Dataset 3**).

DISCUSSION

We characterized how host immunity is modulated to facilitate endosymbiosis by quantifying phenotypic metrics of symbiosis and profiling bacterial communities, whole-organism proteomics, and single-cell RNA sequencing (scRNA-seq) in symbiotic (with algal symbionts) and aposymbiotic (without algal symbionts) samples of the facultatively symbiotic stony coral *Oculina arbuscula.* This work reveals cell type-specific compartmentalization of the immune system that allows symbiotic hosts to simultaneously downregulate key immune pathways in algal-containing cells while maintaining constitutive organismal immunity across other cell types. Our results are consistent with emerging data indicating that the cnidarian-algal symbiosis is a balancing act between nutritional status and immunity (14,20,55). Hosts must balance beneficial nutrient acquisition from photosynthetic algae with the downregulation of the immune system to allow for intracellular symbiont establishment and maintenance.

Whole-organism transcriptomic and proteomic research has yielded many insights into how symbiosis influences cnidarian host functions. For example, transcriptomic (14) and proteomic ((55) and this study) experiments have demonstrated how symbiotic state affects gene expression and protein abundance. Our whole-organism proteomic data across different symbiotic states under baseline conditions support results from previous experiments (14,20,54) demonstrating that differential regulation of the host immune system is required to maintain symbiosis. Specifically, our whole-organism proteomic analysis and bulk analysis of our single-cell data confirmed the pattern of differential regulation of immune system pathways under symbiosis in a facultatively symbiotic cnidarian, which has been previously observed in *O. arbuscula* (14), *E. pallida* (20) and *A. poculata* (27). However, signals from whole-organism studies may be confounded by differences in the proportions of cell types sampled and opposing

expression signals within these cell types (56). Furthermore, while observed patterns between

scRNAseq and proteomic profiling are consistent between aposymbiotic and symbiotic fragments,

we cannot determine whether the signature in the bulk proteomic data is driven by gastrodermal

algal-hosting cells. Future work coupling scRNAseq with single cell proteomic profiling (57) or

immunohistochemistry would reveal more information about the localization of differentially

enriched proteins across cell types.

scRNA-seq has been gaining traction in research on the coral immune system. In one study,

Levy et al. characterized two immune cell states with divergent expression patterns in obligate

symbiotic stony coral *Stylaphora pistillata* adults (47). In additional studies, Hu et al. described

how LePin, a lectin involved in the phagocytic pathways of the cnidarian immune system, is

required for symbiont establishment in algal-hosting cells of the soft coral *Xenia* sp. (48,58). Cell

state-specific regulation of the cnidarian immune system has also been proposed to occur in the

sea anemone *E. pallida* via whole-larvae fluorescence imaging and transcriptomics (59). Notably,

MYD88, a protein that transduces extracellular signals into the host to induce the NF-κB pathway

(60,61) is repressed in *E. pallida* algal-hosting cells; this local immune modulation only occurs in

symbiosis with compatible algae, thereby allowing host cells to sort and expel other non-symbiotic

microbes (59). Our scRNAseq approach further highlights how the *O. arbuscula* immune system

is compartmentalized and reveals how cell state-specific immune modulation may be key to the

maintenance of a functionally symbiotic host. Here, we observe a downregulation of immune

pathway genes in populations of symbiotic gastrodermal cells along the cell fate trajectory that

terminates in algal-hosting cells, indicating that immune suppression is key to symbiosis

maintenance in these cell types. Although we were unable to confidently identify LePin in our

data, we noted the clear involvement of immune gene regulation in driving transcriptomic

differences in gastrodermal cells from symbiotic and aposymbiotic tissue. Notably, these genes

included Heat-Shock Proteins (HSPs) and Cathepsins, which can be involved in the NF-κB

pathway and protein chaperoning and are hypothesized to function in symbiosis regulation (62–

65). Induction of HSPs has been observed following thermal stress in symbiotic cnidarians,

implicating the unfolded protein response in ensuring the stability of symbiosis (63,66).

Interestingly, HSPs and Cathepsins have been previously shown to be markers for immune cells

in the stony coral *S. pistillata* (47), but it remains to be determined whether this immune signature

in gastrodermal cells is conserved in cnidarians engaged in obligate symbioses. It is important to

note that of the two gastrodermal cell types that we identified as displaying symbiosis-induced

immune gene suppression, Gastrodermis 1 exhibited stronger signals than Gastrodermis 2. This

observation is most likely due to the fact that Gastrodermis 1 contains more than twice the number

of cells as Gastrodermis 2 (1,682 cells versus 802 cells) and therefore had more statistical power.

Evidence for localized regulation of immune system pathways in algal-hosting

gastrodermal cells has now been shown in a soft coral (58), a sea anemone (59), and a stony coral

(present study). This compartmentalized immune system regulation allows for the maintenance of

whole-organism immunity across other cell types, most notably in immune cells. We observed no

differences in gene expression in the *O. arbuscula* Immune Cell cluster between symbiotic and

aposymbiotic tissues. The maintenance of immunity function across other cell types in *O.*

arbuscula in and out of symbiosis provides nuance to the proposed immunity-nutrient tradeoff that

symbiotic cnidarians must balance; that is, hosting algal symbionts within gastrodermal cells may

not reduce whole-organism immunity. The immune system has evolved in different ways across

metazoans. For example, in the three-spined stickleback *Gasterosteus aculeatus*, eight distinct

immune cell states were identified with distinct abundances and transcriptomic profiles across

populations, demonstrating specialized and rapid evolution of compartmentalized immune

regulation (67). Here, we observe that "microevolution" of the immune system in *O. arbuscula*

has occurred in three transcriptomically distinct immune cell states, indicating diversified immune

cell function that is in agreement with other single-cell studies in stony coral (47).

"Microevolution" may additionally occur in specialized algal-hosting gastrodermal cells, allowing

these gastrodermal cells to function as pseudo-immune cells without affecting organismal

immunity in immune cells. Notably, because canonical immune cells are able to maintain

constitutive immunity across symbiotic states, previous work in field-collected tropical corals

showing differences in pathogen susceptibility by symbiotic state (23,25) may be due to other

factors (e.g., capacity for heterotrophy, efficiency of host-algal nutrient exchange, competition for

nitrogen between host and algae) and not due to baseline immune cell tradeoffs due to symbiosis

(26). However, additional work focused on characterizing differences between cell type-specific

immune regulation in obligate versus facultative cnidarians is necessary to validate this hypothesis.

Our comparative analysis between *O. arbuscula* and *Xenia* sp. (48) algal-hosting and non-algal-

hosting gastrodermal cells suggests that obligate and facultatively symbiotic cnidarians may

maintain their algal symbionts through different regulatory pathways, such as through classical

immune response pathways in *Xenia* algal-hosting cells compared to cellular stress response

pathways in *O. arbuscula* algal-hosting cells, although further research is needed to explore this

hypothesis.

Our single-cell atlas also provides insights into carbon and nitrogen cycling within the

gastrodermal cells of symbiotic and aposymbiotic *O. arbuscula*. Notably, we find that the

gastrodermal cells that suppress immunity to allow for symbiosis are the same cells that undergo

active nutrient cycling, most likely to regulate algal cell proliferation (11,14,68). Previous work

has proposed that increased nitrogen cycling and assimilation occurring in symbiotic host cells

controls algal cell proliferation to maintain optimal nutrient exchange between host and symbiont,

preventing parasitism of the symbiont on the host (11,55). In our data, the co-expression of sugar

transport and nitrogen cycling genes within the same symbiotic cells in specific gastrodermal cell

clusters supports recent work showing that the stability of cnidarian-algal relationships relies on

glucose-dependent nitrogen competition between the host and symbiont (12). In this model, the

metabolites and energy produced from host glycolysis of symbiont-derived glucose may be used

by the host for amino acid synthesis, through which nitrogen species, such as ammonium, can be

assimilated. Hosts can thus modulate the amount of ammonium translocated to the symbiont for

proliferation (12). However, in the present study, gastrodermal cells from aposymbiotic tissue

exhibited lower expression of sugar transport and nitrogen cycling genes, which is likely driven

by symbiotic cnidarians obtaining glucose sugars from symbiotic algae, whereas aposymbiotic *O.*

arbuscula rely on heterotrophic sources to meet their metabolic needs (9,69). Thus, metabolic

processing of carbon and nitrogen are likely performed through distinct pathways in cnidarians

that rely predominantly on heterotrophy (i.e., aposymbiotic corals) compared to those that rely

predominantly on autotrophy (i.e., symbiotic corals) (55).

Finally, while we observed no differences in the composition or diversity of the bacterial

communities between aposymbiotic and symbiotic *O. arbuscula*, microbial partners may still play

an important role in nutrient cycling and possibly host immunity. Differences in bacterial

composition by symbiotic state may not have been observed in our study because corals were

housed in common garden conditions. Future experiments with increased sample sizes collected

in situ coupled with metabolomics or metatranscriptomics of the bacterial communities would

more fully capture the functional variation of bacterial communities between symbiotic states.

Our scRNAseq approach applied in a facultative coral provides a novel way to study the

tradeoffs and cellular regulation associated with the cnidarian-algal symbiosis. Previous work

using whole-organism experimental techniques suggested that immunity costs are associated with

hosting algal symbionts (14,20). Here, we showcase how differences in cell states and gene

expression can conflate patterns observed in whole-organism 'omics studies and, importantly,

facilitate cell type-specific immune regulation to allow for endosymbiosis. Our description of

highly specialized cell state-specific immune system regulation in a basal metazoan indicates that

the immune costs of symbiosis previously observed in whole organism studies might not be as

costly as previously assumed. Future work should compare how compartmentalized immunity is

modulated in different host-algal pairings in facultative versus obligate cnidarians and also

determine how these cellular processes are shaped by changing environmental conditions.

Acknowledgements

This research was supported by a National Science Foundation grant IOS-1937650 (to T.D.G. and

S.W.D.). M,V.-I. was supported by an NSF Graduate Research Fellowship, NSF NRT DGE

1735087, and a Boston University Marine Program (BUMP) Warren McLeod Marine Fellowship.

J.K.D.-A. was supported by a BUMP Warren McLeod Marine Fellowship. We thank Diana

Burkart-Waco of 10X Genomics Support for her assistance with data processing. We also thank

Boston University Microarray & Sequencing Resource Core Facility, the Boston University Single

Cell Sequencing Core Facility, Boston University Shared Computing Cluster, and the Tufts

University Core Facility.

Data availability

Raw fastq files for 16S and single-cell RNA-seq data can be found on NCBI's Short Read Archive

Bioproject PRJNA1122932. All data and code for analyses can be found at

https://github.com/mariaingersoll/Oculina_sc_manuscript.git.

Competing interests

The authors declare no competing interests.

Figure 1:   Symbiosis facilitates higher light absorptance but no changes to bacterial
communities in *O. arbuscula*. a Algal cell counts are 400x higher in symbiotic (brown, right) *O.*
arbuscula fragments than in aposymbiotic (white/gray, left) fragments (p<0.001). **b** Average
reflectance (R) in the PAR region (400–700 nm) was 22% +/- 4% in symbiotic *O. arbuscula*
(brown) and 77% +/- 3% in aposymbiotic *O. arbuscula* (gray) (p < 0.01) **c** Average absorptance
(A) (A = 1-R) demonstrating relative amount of solar energy with potential for use in
photosynthesis. Symbiotic *O. arbuscula* (brown) has an A_{675} of 85% (+/- 4%), suggesting a
functional symbiosis, and aposymbiotic (gray) has an A_{675} of 24% (+/- 2%) (p<0.001). In **b** and **c**,
shaded colored areas represent standard deviation. **d** PCoA of bacterial communities from all
ASVs in 16S data from symbiotic (brown) and aposymbiotic (gray) *O. arbuscula* genotypes. No
differences in bacterial communities were observed between symbiotic states.

612
Figure 2: Proteomic profiles of symbiotic and aposymbiotic *O. arbuscula* reveal differential
enrichment of proteins by symbiotic state. a Principal component analysis of proteomic profiles
across four genets, with symbiotic (sym) and aposymbiotic (apo) states distinguished by color, and
genets by shape. **b** Volcano plot of differentially enriched proteins (DEPs) identified through
pairwise comparison (p-value < 0.1). Upregulated proteins (105) are represented by brown dots,
downregulated proteins (82) by gray dots, while black dots indicate non-DEPs in symbiotic
relative to aposymbiotic corals (total N=2,191). **c** Heatmap of all DEPs across symbiotic states of
O. arbuscula. The color scale represents the log_2foldchange of each protein (row) for each coral
sample (column) relative to the protein's mean across all samples. **d** Interaction network of
proteins associated with Immune Response, Repair and Stress Response, and Metabolic Processes,
derived from human protein–protein interactions in the STRING database. Node colors indicate
upregulation (brown) or downregulation (gray) in symbiotic coral samples.

625
Figure 3: scRNA-seq reveals 28 transcriptomically distinct cell clusters across 7 cell types in
symbiotic and aposymbiotic *O. arbuscula*. a A UMAP Projection of the transcriptomes of 6,992
cells from symbiotic (sym) and aposymbiotic (apo) *O. arbuscula* reveals 28 clusters across 7 broad
cell types. Each cluster is color coded. **b** Each cluster is transcriptomically distinct, with expression
similarities present between clusters of the same cell type, as represented by the 10 genes most
highly expressed in each cluster. Expression values are average log$_2$foldchanges computed from
normalized gene expression data for each cell. Cell clusters are colored as in (**a**).

634
Figure 4: Gastrodermis cells along the algal-hosting trajectory are involved in sugar
transport and nitrogen cycling. a UMAP projection of Gastrodermis 1, 2, 3, and 4 cells from
symbiotic *O. arbuscula* overlayed with a graph of cell trajectories. Outcomes (fates) are denoted
by gray circles with black numbers. Branch nodes are denoted by black circles with white numbers.
b UMAP projections of the 1,682 independently reclustered Gastrodermis 1 cells revealed 4
subclusters. Each subcluster is color-coded in the left panel. Gastrodermis 1 cells separate
according to symbiotic state across the UMAP projection (right panel). **c** UMAP projections of the

802 independently reclustered Gastrodermis 2 cells revealed 3 subclusters, color coded in the left
panel. Gastrodermis 2 cells separate according to symbiotic state (right panel). **d** UMAP
projections of the 611 independently reclustered Gastrodermis 3 cells revealed 2 subclusters, color
coded in the left panel. Limited separation of Gastrodermis 3 cells by symbiotic state (right panel)
was observed. Violin plots of Glutamate Dehydrogenase (DHE4, **e**), Glutamate Synthase [NADH]
(GLSN, **f**), and Apolipophorin (APOB, **g**) across all cell clusters in both symbiotic states. DHE4
was identified as a marker gene for Gastrodermis 1; GLSN and APOB were identified as marker
genes for both Gastrodermis 1 and 2. Asterisks denote genes differentially expressed in symbiotic
versus aposymbiotic clusters (* = p-adjusted value <0.05; *** = p-adjusted value < 0.001).
Expression levels of genes in **e-f** are normalized per cell, and significance is calculated from
DESeq2. Coexpression plots of DHE4 and GLSN (**h**), APOB and GLSN (**i**), and DHE4 and APOB
(**j**) across Gastrodermis 1 cells from symbiotic *O. arbuscula* revealed high coexpression of these
sugar transport and nitrogen cycling genes within Subcluster 3, but not in the other subclusters.
The coloration of each cell represents the per-cell mean expression value of each gene scaled to a
maximum value of 10.

Figure 5: Negligible gene expression differences between Immune Cells of symbiotic and aposymbiotic samples. Immune Cells from symbiotic (**a**) and aposymbiotic (**b**) samples have high expression of previously identified immune cell marker genes. Dot size represents the percent of cells in each cell state that express each gene, and dot color represents the average (normalized) expression of each gene within each cell state. **c** Three Immune Cell sublcusters have distinct transcriptomic profiles, as represented by the annotated genes within the top 20 most highly enriched genes in each subcluster. Expression values are average $\log_2$foldchanges computed from normalized gene expression data of each cell. **d** UMAP projections of the 115 independently reclustered Immune Cells revealed 3 subclusters (left panel) that do not separate according to symbiotic state (right panel).

669
Figure 6: Gastrodermis 1 and 2 cells from symbiotic samples downregulate immune pathway
genes compared to aposymbiotic samples. a Spatial representation showing expression of genes
annotated with differentially enriched immunity GO terms across the whole dataset. Coloration
represents the percentage of all transcripts in each cell corresponding to the genes of interest (555
genes of interest out of 24,716 total genes). **b** 8 PCs, driven by 96 annotated genes, contributed to
the separation of symbiotic state in the reclustered UMAP of Gastrodermis 1 (see **Fig. 4b**). 33 of

the 96 were differentially expressed. 32 genes are annotated with GO terms relating to immunity
(bold, italicized). **c** 5 PCs, driven by 59 annotated genes, contributed to the separation of symbiotic
state in the reclustered UMAP of Gastrodermis 2 (see **Fig 4c**). 4 of the 59 genes were differentially
expressed. 13 genes are annotated with GO terms relating to immunity (bold, italicized). In both **b**
and **c**, the red bar indicates genes annotated with the COG term 'posttranslational modification,
protein turnover, and chaperones.' Significance is calculated from DESeq2 (*** = p-adjusted value
< 0.001, *** = p-adjusted value < 0.01, * = p-adjusted value < 0.05).

REFERENCES

1. Thrall PH, Hochberg ME, Burdon JJ, Bever JD. Coevolution of symbiotic mutualists and parasites in a community context. *Trends in Ecology & Evolution*. 2007 Mar 1;22(3):120–6.

2. Baker DM, Freeman CJ, Wong JCY, Fogel ML, Knowlton N. Climate change promotes parasitism in a coral symbiosis. *ISME Journal*. 2018 Mar;12(3):921–30.

3. Lesser MP, Stat M, Gates RD. The endosymbiotic dinoflagellates (Symbiodinium sp.) of corals are parasites and mutualists. *Coral Reefs*. 2013 Sep 1;32(3):603–11.

4. Mansfield KM, Gilmore TD. Innate immunity and cnidarian-Symbiodiniaceae mutualism. *Developmental & Comparative Immunology*. 2019 Jan 1;90:199–209.

5. Zambrano-Villa S, Rosales-Borjas D, Carrero JC, Ortiz-Ortiz L. How protozoan parasites evade the immune response. *Trends in Parasitology*. 2002 Jun 1;18(6):272–8.

6. Xiang T, Lehnert E, Jinkerson RE, Clowez S, Kim RG, DeNofrio JC, et al. Symbiont population control by host-symbiont metabolic interaction in Symbiodiniaceae-cnidarian associations. *Nature Communications*. 2020 Jan 8;11(1):108.

7. Whittle M, Bonsall MB, Barreaux AMG, Ponton F, English S. A theoretical model for host-controlled regulation of symbiont density. *Journal of Evolutionary Biology*. 2023 Dec 1;36(12):1731–44.

8. Muscatine L, Cernichiari E. Assimilation of photosynthetic products of zooxanthellae by a reef coral. *The Biological Bulletin*. 1969 Dec;137(3):506–23.

9. Burriesci MS, Raab TK, Pringle JR. Evidence that glucose is the major transferred metabolite in dinoflagellate–cnidarian symbiosis. *Journal of Experimental Biology*. 2012 Oct 1;215(19):3467–77.

10. Lipschultz F, Cook C. Uptake and assimilation of 15N-ammonium by the symbiotic sea anemones *Bartholomea annulata* and *Aiptasia pallida*: conservation versus recycling of nitrogen. *Marine Biology*. 2002 Mar 1;140(3):489–502.

11. Yellowlees D, Rees TAV, Leggat W. Metabolic interactions between algal symbionts and invertebrate hosts. *Plant, Cell & Environment*. 2008;31(5):679–94.

12. Cui G, Liew YJ, Li Y, Kharbatia N, Zahran NI, Emwas AH, et al. Host-dependent nitrogen recycling as a mechanism of symbiont control in Aiptasia. *PLOS Genetics*. 2019 Jun 24;15(6):e1008189.

13. Rädecker N, Raina JB, Pernice M, Perna G, Guagliardo P, Kilburn MR, et al. Using Aiptasia as a model to study metabolic interactions in cnidarian-Symbiodinium symbioses. *Frontiers in Physiology*. 2018 Mar 16.

14. Rivera HE, Davies SW. Symbiosis maintenance in the facultative coral, *Oculina arbuscula*, relies on nitrogen cycling, cell cycle modulation, and immunity. *Scientific Reports*. 2021 Oct 27;11(1):21226.

15. Hoegh-Guldberg O. Coral reef ecosystems and anthropogenic climate change. *Regional Environmental Change*. 2011 Mar 1;11(1):215–27.

16. Bove CB, Ingersoll MV, Davies SW. Help me, symbionts, you're my only hope: Approaches to accelerate our understanding of coral holobiont interactions. *Integrative and Comparative Biology*. 2022 Dec 1;62(6):1756–69.

17. Davy SK, Allemand D, Weis VM. Cell biology of cnidarian-dinoflagellate symbiosis. *Microbiology and Molecular Biology Review*s. 2012 Jun 11;76(2):229–61.

18. Cui G, Konciute MK, Ling L, Esau L, Raina JB, Han B, et al. Molecular insights into the Darwin paradox of coral reefs from the sea anemone Aiptasia. *Science Advances*. 2023 Mar 15;9(11):eadf7108.

19. Gault JA, Bentlage B, Huang D, Kerr AM. Lineage-specific variation in the evolutionary stability of coral photosymbiosis. *Science Advances*. 2021 Sep 22;7(39):eabh4243.

20. Mansfield KM, Carter NM, Nguyen L, Cleves PA, Alshanbayeva A, Williams LM, et al. Transcription factor NF-κB is modulated by symbiotic status in a sea anemone model of cnidarian bleaching. *Scientific Reports*. 2017 Nov 22;7(1):16025.

21. Matthews JL, Crowder CM, Oakley CA, Lutz A, Roessner U, Meyer E, et al. Optimal nutrient exchange and immune responses operate in partner specificity in the cnidarian-dinoflagellate symbiosis. *Proceedings of the National Academy of Sciences*. 2017 Dec 12;114(50):13194–9.

22. Stankiewicz KH, Guiglielmoni N, Kitchen SA, Flot JF, Barott KL, Davies SW, et al. Genomic comparison of the temperate coral *Astrangia poculata* with tropical corals yields insights into winter quiescence, innate immunity, and sexual reproduction. *bioRxiv*. 2023.

23. Merselis DG, Lirman D, Rodriguez-Lanetty M. Symbiotic immuno-suppression: is disease susceptibility the price of bleaching resistance? *PeerJ*. 2018 Apr 17;6:e4494.

24. Muller EM, Bartels E, Baums IB. Bleaching causes loss of disease resistance within the threatened coral species *Acropora cervicornis*. *eLife*. 2018 Sep 11;7:e35066.

25. Shore-Maggio A, Callahan SM, Aeby GS. Trade-offs in disease and bleaching susceptibility among two color morphs of the Hawaiian reef coral, *Montipora capitata*. *Coral Reefs*. 2018 Jun 1;37(2):507–17.

26. Valadez-Ingersoll M, Aguirre Carrión PJ, Bodnar CA, Desai NA, Gilmore TD, Davies SW. Starvation differentially affects gene expression, immunity and pathogen susceptibility across symbiotic states in a model cnidarian. *Proceedings of the Royal Society B: Biological Sciences*. 2024 Feb 28;291(2017):20231685.

27. Changsut I, Womack HR, Shickle A, Sharp KH, Fuess LE. Variation in symbiont density is linked to changes in constitutive immunity in the facultatively symbiotic coral, *Astrangia poculata*. *Biology Letters*. 2022 Nov 16;18(11):20220273.

28. Lajeunesse TC, Parkinson JE, Reimer JD. A genetics-based description of *Symbiodinium minutum sp. nov.* and *S. psygmophilum sp. nov.* (Dinophyceae), two dinoflagellates symbiotic with cnidaria. *Journal of Phycology*. 2012;48(6):1380–91.

29. Enríquez S, Méndez ER, Iglesias-Prieto R. Multiple scattering on coral skeletons enhances light absorption by symbiotic algae. *Limnology and Oceanography*. 2005;50(4):1025–32.

30. Scheufen T, Iglesias-Prieto R, Enríquez S. Changes in the number of symbionts and *Symbiodinium* cell pigmentation modulate differentially coral light absorption and photosynthetic performance. *Frontiers in Marine Science*. 2017 Sep 26.

31. Vásquez-Elizondo RM, Legaria-Moreno L, Pérez-Castro MÁ, Krämer WE, Scheufen T, Iglesias-Prieto R, et al. Absorptance determinations on multicellular tissues. *Photosynthesis Research*. 2017 Jun 1;132(3):311–24.

32. Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environmental Microbiology*. 2016;18(5):1403–14.

33. Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial Ecology*. 2015 Jun 4;75(2):129–37.

34. Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, et al. vegan: Community Ecology Package. 2024.

35. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*. 2014 Dec 5;15(12):550.

36. Aguirre Carrión PJ, Williams LM, Gilmore TD. Molecular and biochemical approaches to
study the evolution of NF-κB signaling in basal metazoans. *Methods in Molecular Biology*.
2021 2366:67-91.
37. Guo H, Isserlin R, Chen X, Wang W, Phanse S, Zandstra PW, et al. Integrative network
analysis of signaling in human CD34+ hematopoietic progenitor cells by global
phosphoproteomic profiling using TiO2 enrichment combined with 2D LC-MS/MS and
pathway mapping. *Proteomics*. 2013;13(8):1325–33.
38. Blum BC, Emili A. Omics Notebook: robust, reproducible and flexible automated multiomics
exploratory analysis and reporting. *Bioinformatics Advances*. 2021 Jan 1;1(1):vbab024.
39. Mering C von, Huynen M, Jaeggi D, Schmidt S, Bork P, Snel B. STRING: a database of
predicted functional associations between proteins. *Nucleic Acids Research*. 2003 Jan
1;31(1):258–61.
40. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: A software
environment for integrated models of biomolecular interaction networks. *Genome Research*.
2003 Nov 1;13(11):2498–504.
41. Rosental B, Kozhekbaeva Z, Fernhoff N, Tsai JM, Traylor-Knowles N. Coral cell separation
and isolation by fluorescence-activated cell sorting (FACS). *BMC Cell Biology*. 2017 Aug
29;18(1):30.
42. Cole AG, Steger J, Hagauer J, Denner A, Ferrer Murguia P, Knabl P, et al. Updated single cell
reference atlas for the starlet anemone *Nematostella vectensis*. *Frontiers in Zoology*. 2024 Mar
18;21(1):8.
43. Steger J, Cole AG, Denner A, Lebedeva T, Genikhovich G, Ries A, et al. Single-cell
transcriptomics identifies conserved regulators of neuroglandular lineages. *Cell Reports*. 2022
Sep 20;40(12):111370.
44. Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively parallel
digital transcriptional profiling of single cells. *Nature Communications*. 2017 Jan
16;8(1):14049.
45. Fifer JE. Coral adaptations across ecological and evolutionary scales. 2023.
46. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al. Dictionary
learning for integrative, multimodal and scalable single-cell analysis. *Nature Biotechnology*.
2024 Feb;42(2):293–304.
47. Levy S, Elek A, Grau-Bové X, Menéndez-Bravo S, Iglesias M, Tanay A, et al. A stony coral
cell atlas illuminates the molecular and cellular basis of coral symbiosis, calcification, and
immunity. *Cell*. 2021 May 27;184(11):2973-2987.e18.
48. Hu M, Zheng X, Fan CM, Zheng Y. Lineage dynamics of the endosymbiotic cell type in the
soft coral *Xenia*. *Nature*. 2020 Jun;582(7813):534–8.
49. Wright RM, Aglyamova GV, Meyer E, Matz MV. Gene expression associated with white
syndromes in a reef building coral, *Acropora hyacinthus*. *BMC Genomics*. 2015 May
9;16(1):371.
50. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics and
regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nature
Biotechnology*. 2014 Apr;32(4):381–6.
51. Derelle R, Philippe H, Colbourne JK. Broccoli: Combining phylogenetic and network analyses
for orthology assignment. *Molecular Biology and Evolution*. 2020 Nov 1;37(11):3389–96.
52. Chen H. VennDiagram: generate high-resolution Venn and Euler plots. R package v. 1.7. 3.
2022.

53. Margolis SR, Dietzen PA, Hayes BM, Wilson SC, Remick BC, Chou S, et al. The cyclic dinucleotide 2′3′-cGAMP induces a broad antibacterial and antiviral response in the sea anemone *Nematostella vectensis*. *Proceedings of the National Academy of Sciences*. 2021 Dec 21;118(51):e2109022118.

54. Lehnert EM, Mouchka ME, Burriesci MS, Gallo ND, Schwarz JA, Pringle JR. Extensive differences in gene expression between symbiotic and aposymbiotic cnidarians. *G3 Genes|Genomes|Genetics*. 2014 Feb 1;4(2):277–95.

55. Oakley CA, Ameismeier MF, Peng L, Weis VM, Grossman AR, Davy SK. Symbiosis induces widespread changes in the proteome of the model cnidarian Aiptasia. *Cellular Microbiology*. 2016;18(7):1009–23.

56. Freedman AH, Sackton TB. Rethinking eco-evo studies of gene expression for non-model organisms in the genomic era. *Molecular Ecology*. 2024 May 9;n/a(n/a):e17378.

57. Bennett HM, Stephenson W, Rose CM, Darmanis S. Single-cell proteomics enabled by next-generation sequencing or mass spectrometry. *Nature Methods*. 2023 Mar;20(3):363–74.

58. Hu M, Bai Y, Zheng X, Zheng Y. Coral–algal endosymbiosis characterized using RNAi and single-cell RNA-seq. *Nature Microbiology*. 2023 Jul;8(7):1240–51.

59. Jacobovitz MR, Rupp S, Voss PA, Maegele I, Gornik SG, Guse A. Dinoflagellate symbionts escape vomocytosis by host cell immune suppression. *Nature Microbiology*. 2021 Jun;6(6):769–82.

60. Sugiyama K ichi, Muroi M, Kinoshita M, Hamada O, Minai Y, Sugita-Konishi Y, et al. NF-κB activation via MyD88-dependent Toll-like receptor signaling is inhibited by trichothecene mycotoxin deoxynivalenol. *The Journal of Toxicological Sciences*. 2016;41(2):273–9.

61. Gohda J, Matsumura T, Inoue J ichiro. Cutting edge: TNFR-Associated Factor (TRAF) 6 ss essential for MyD88-dependent pathway but not Toll/IL-1 Receptor Domain-Containing Adaptor-Inducing IFN-β (TRIF)-dependent pathway in TLR signaling1. *The Journal of Immunology*. 2004 Sep 1;173(5):2913–7.

62. Kenkel CD, Sheridan C, Leal MC, Bhagooli R, Castillo KD, Kurata N, et al. Diagnostic gene expression biomarkers of coral thermal stress. *Molecular Ecology Resources*. 2014;14(4):667–78.

63. Cleves PA, Krediet CJ, Lehnert EM, Onishi M, Pringle JR. Insights into coral bleaching under heat stress from analysis of gene expression in a sea anemone model system. *Proceedings of the National Academy of Sciences*. 2020 Nov 17;117(46):28906–17.

64. Sproles AE, Oakley CA, Matthews JL, Peng L, Owen JG, Grossman AR, et al. Proteomics quantifies protein expression changes in a model cnidarian colonised by a thermally tolerant but suboptimal symbiont. *ISME Journal*. 2019 Sep;13(9):2334–45.

65. Dimos BA, Mahmud SA, Fuess LE, Mydlarz LD, Pellegrino MW. Uncovering a mitochondrial unfolded protein response in corals and its role in adapting to a changing world. *Proceedings of the Royal Society B: Biological Sciences*. 2019 Jun 26;286(1905):20190470.

66. Petrou K, Nunn BL, Padula MP, Miller DJ, Nielsen DA. Broad scale proteomic analysis of heat-destabilised symbiosis in the hard coral *Acropora millepora*. *Scientific Reports*. 2021 Sep 24;11(1):19061.

67. Fuess LE, Bolnick DI. Single-cell RNA sequencing reveals microevolution of the stickleback immune system. *Genome Biology and Evolution*. 2023 Apr 1;15(4):evad053.

68. Rädecker N, Pogoreutz C, Voolstra CR, Wiedenmann J, Wild C. Nitrogen cycling in corals: the key to understanding holobiont functioning? *Trends in Microbiology*. 2015 Aug 1;23(8):490–7.

867   69. Goldberg WM. Coral food, feeding, nutrition, and secretion: A review. *Results and Problems*
868   *in Cell Differentiation.* 2018.