Multi-omic Landscape of Extracellular Vesicles in Human Carotid Atherosclerotic Plaque Reveals Endothelial Communication Networks

Sneha Raju, MD, PhD,[1, 2, 3, 4] Mandy E Turner, PhD,[5] Christian Cao,[4] Majed Abdul-Samad, BSc,[6] Neil Punwasi, MSc,[7] Mark C Blaser, PhD,[5] Rachel ME Cahalane, PhD,[5,8] Steven R Botts, MSc,[1,2,4] Kamal Prajapati, BSc,[1,6] Sarvatit Patel, PhD,[1] Ruilin Wu, MSc,[1,6] Dakota Gustafson, PhD,[1] Natalie J Galant, PhD,[9] Lindsey Fiddes, PhD,[4] Melody Chemaly, PhD,[10] Ulf Hedin, MD, PhD,[10] Ljubica Matic, PhD,[10] Michael Seidman, MD, PhD,[6,7] Vallijah Subasri, PhD,[7] Sasha A Singh, PhD,[5] Elena Aikawa, MD, PhD,[5,11] Jason E Fish, PhD,[1,2,6,7] *Kathryn L Howe, MD, PhD,[1,2,3,4,6,7]

Institutions:

[1] Toronto General Hospital Research Institute, University Health Network, Toronto, Canada.

[2] Institute of Medical Science, University of Toronto, Toronto, Canada.

[3] Division of Vascular Surgery, Toronto General Hospital, Toronto, Canada.

[4] Faculty of Medicine, University of Toronto, Toronto ON, Canada.

[5] Center for Interdisciplinary Cardiovascular Sciences, Cardiovascular Division, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

[6] Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, ON, Canada

[7] Peter Munk Cardiac Centre, Toronto General Hospital, Toronto, Canada

[8] Mechanobiology and Medical Device Research group (MMDRG), Biomedical

Engineering, College of Science and Engineering, University of Galway, Galway,

Ireland.

[9] Princess Margaret Cancer Center, Toronto, ON, Canada

[10] Vascular Surgery Division, Department of Molecular Medicine and Surgery, Karolinska

University Hospital and Karolinska Institut, Stockholm, Sweden.

[11] Center for Excellence in Vascular Biology, Brigham and Women's Hospital, Harvard

Medical School, Boston, MA, USA

Short-Title: Endothelial messaging in carotid atherosclerosis

Correspondence:

Kathryn L. Howe (Kathryn.howe@uhn.ca)

Peter Munk Cardiac Centre, University Health Network

6EN 220-200 Elizabeth Street, Toronto General Hospital, Toronto, ON, M5G 2C4,

Canada

Word Count: 14,759

1   **Abstract**

2   Background: Carotid atherosclerosis is a multifaceted disease orchestrated by a myriad

3   of cell-cell communication that drives progression along a clinical continuum

4   (asymptomatic to symptomatic). Extracellular vesicles (EVs) are lipid bilayer membrane-

5   enclosed cell-derived nanoparticles that represent a new paradigm in cellular

6   communication. Little is known about their biological cargo, cellular origin/destination,

7   and functional roles in human atherosclerotic plaque.

8

9   Methods: EVs were enriched via size exclusion chromatography from human carotid

10   endarterectomy samples dissected into plaque and marginal zones (n= 29 patients,

11   paired plaque and marginal zone; symptomatic n=16, asymptomatic n=13), with further

12   density gradient ultracentrifugation for proteomic analysis. EV cargoes were assessed

13   via whole transcriptome miRNA sequencing and mass spectrometry-based proteomics.

14   EV multi-omics were integrated with publicly available bulk and single cell RNA-

15   sequencing (scRNA-seq) datasets to predict EV cellular origin and ligand-receptor

16   interactions and multi-modal biological network integration of EV-cargo was completed.

17   EV functional impact was assessed with endothelial angiogenesis assays.

18

19   Results: Human carotid plaques contained greater quantities of EVs than adjacent

20   marginal zones. EV-miRNA and protein content was different in diseased plaque versus

21   adjacent marginal zones, with differential functions in key atherogenic pathways. EV

22   cellular origin analysis suggested that tissue EV signatures originated from endothelial

23   cells (EC), smooth muscle cells (SMC), and immune cells. Furthermore, EV signatures

from SMCs and immune cells were most enriched in the marginal and plaque zones,

respectively. Integrated tissue vesiculomics and scRNA-seq indicated complex EV-

vascular cell communication strategies that changed with disease progression and

plaque vulnerability (i.e., symptomatic disease). Plaques from symptomatic patients, but

not asymptomatic patients, were characterized by increased involvement of endothelial

pathways and more complex ligand-receptor interactions, relative to their marginal

zones. Plaque-EVs were predicted to mediate communication with ECs. Pathway

enrichment analysis delineated a strong endothelial signature with potential roles in

angiogenesis and neovascularization – well-known indices of plaque instability. This

was corroborated functionally, wherein human carotid symptomatic plaque EVs induced

sprouting angiogenesis in comparison to their matched marginal zones.

Conclusion: Our findings indicate that EVs may drive dynamic changes in plaques

through EV-vascular cell communication and effector functions that typify vulnerability to

rupture, precipitating symptomatic disease. The discovery of endothelial-directed

processes mediated by EVs creates new avenues for novel therapeutics in

atherosclerosis.

Key Words: carotid atherosclerosis, extracellular vesicles, microRNAs, proteomics,

endothelial cells

Non-Standard Abbreviations

47	cryo-EM	cryogenic electron microscopy
48	EC	endothelial cell
49	EV	extracellular vesicle
50	GO	Gene Ontology
51	KEGG	Kyoto Encyclopedia of Genes and Genomes
52	NTA	nanoparticle tracking analysis
53	miRNA	microRNA
54	MISEV	Minimal information for studies of extracellular vesicles
55	MZ	marginal zone
56	PZ	plaque zone
57	scRNA-seq	single cell RNA-sequencing
58	SEC	size exclusion chromotography
59	SMC	smooth muscle cell
60	TEM	transmission electron microscopy
61		
62		

Introduction

Atherosclerosis is a multi-cellular biological process of lipid accumulation, unresolved inflammation, and plaque development in the arterial wall.[1,2] While some plaques enlarge and impede blood flow, plaques can also become unstable and rupture, thereby releasing atheroembolic material that occludes an end-artery leading to a transient ischemic attack (TIA, i.e. neurologic recovery) or stroke.[1,3] Atherosclerosis exists along a disease continuum wherein much of the disease process is silent and presents as asymptomatic disease; low-density lipoprotein accumulation and endothelial dysfunction initiates the disease process, followed by recruitment of adaptive and innate immune cells and migration of smooth muscle cells (SMCs) leading to plaque progression and development of the necrotic core, followed by eventual fibrous cap rupture and thrombosis causing clinical events such as myocardial infarction or stroke.[2] Altered cellular communication is a critical linchpin of atherogenesis.[4-7] Recent work has identified the importance of endothelial cells (ECs) in directing differential, polarized messaging with cells found in circulation and those in the vessel wall.[8] Given that several types of plaque cells exist in transitory states (ECs, SMCs, macrophages), are highly dynamic, and can alter their phenotype in the active plaque microenvironment, an exciting possibility is that cellular states dictate and result from changes in cell-cell communication[9-12]

Much of the early and late stages of atherosclerotic disease is silent or asymptomatic, creating challenges in patient stratification. Unlike management of symptomatic carotid atherosclerotic disease, which is well established and includes surgical removal of the

plaque (carotid endarterectomy) or endovascular carotid stenting with the goal of

secondary stroke prevention [13-16], management of patients with asymptomatic carotid

disease is more nuanced, as interventions have an upfront risk of stroke and we have

not yet solved how to accurately predict plaque rupture.[17,18] Current non-operative

management with lipid reduction, anti-hypertensive therapy, and lifestyle modifications

(i.e. smoking cessation) reduces stroke risk, but some patients progress despite

meeting these targets.[17,19] Prior studies have employed whole-tissue proteomics,

transcriptomics, and metabolomics to identify major shifts in plaque biology with disease

progression.[20-23] However, despite ongoing research, there are still no universally

reliable means to identify the patient at risk and novel therapeutic targets need to be

unveiled through a better understanding of plaque biology as it evolves along the

disease continuum.

Given the multi-cellular environment of the atherosclerotic plaque, cell-cell

communication is a major factor driving the biology of the disease. Extracellular vesicles

(EVs) are lipid bilayered membrane-enclosed cell-derived nanoparticles that carry

bioactive molecules such as microRNAs (miRNAs) and proteins from a donor to a

recipient cell, where they participate in EV-ligand based interactions or are endocytosed

and release contents that can modulate recipient cell phenotype.[24] Given the clinical

potential of this field, the development of Minimal Information for the Study of

Extracellular Vesicles (MISEV) guidelines has been crucial for improving clarity,

consistency in reporting, and interpretation of EV-based studies.[25,26] Previous studies in

atherosclerotic plaques have examined larger EV particles (i.e., microparticles; primarily

>200 nm) and provided early insights into their importance in atherosclerotic biology.[27,28]

However the evolution of EV isolation and technical progress in EV cargo analysis

mandates a fresh examination of their biological roles, particularly with respect to small

EVs (<200 nm) given their abundant numbers and potential for therapeutic

modulation.[29,30] Recently, a proteomics study of human carotid plaque-derived EVs

small and large identified their role in advanced atherosclerotic calcification, and found

that EV-proteomics can identify additional phenotypes than those at the whole plaque

level, warranting independent investigation.[31] Our understanding of carotid tissue

vulnerability as modulated by EV cellular source, EV-cell communication, and the

impact of EVs on downstream cellular processes is limited. There is a need to

understand the holistic biological niche of stable and vulnerable carotid plaques in these

contexts, with the goal of identifying novel targets for preventing rupture through plaque

stabilization.

The intraplaque environment is dynamic, changes over time, and there is a significant

role for cell-cell communication. EVs contain cargo that serve as important mediators of

cell-cell communication with functional effects that alter cell biology in recipient cells.[32-34]

What is not known however, are the alterations to cell-cell communication that render an

atherosclerotic plaque vulnerable to rupture. Patients with symptomatic carotid disease

have demonstrated clinical consequences of carotid plaque vulnerability where

atheroembolic events have caused transient ischemic events or stroke.[35] Here, we

performed a novel analysis to uncover the transcriptomic and proteomic EV profile in

carotid plaque and adjacent marginal zones, in both asymptomatic and symptomatic

disease states. By integrating our data with scRNA-seq datasets, we identified the

predicted EV-cell communication pathways that may be enriched in asymptomatic and

symptomatic (vulnerable) carotid plaques. Furthermore, we found a strong endothelial

signature and demonstrated a functional role for plaque EVs in regulating angiogenesis

and neovascularization – well-known precursors of plaque instability.[36-38] Importantly,

symptomatic plaques were studied within 30 days of the last neurologic event, with

pathological evaluation of a subset that clearly demonstrated intraplaque hemorrhage,

consistent with the EV angiogenesis experiments pursued. Together, our data suggest

that EVs may contribute to dynamic changes seen in plaques - through EV-cell

communication and effector functions - including neovascularization and angiogenesis

that contribute to plaque destabilization. The discovery of an endothelial-centric process

that could be mediated by EVs points towards novel therapeutics targeting this highly

accessible and strategic cellular layer.

Methods

Detailed methods are provided in the Supplemental Material. Human carotid

endarterectomy specimens were obtained from patients undergoing surgery for

asymptomatic and symptomatic disease. Samples were handled under the institutional

research ethics board protocols set by University Health Network, Toronto Canada (18-

5282). Patients gave written informed consent and procedures were followed in

accordance with institutional guidelines. Tissues were separated by plaque and

marginal (non-atheromatous) zones and EV enrichment performed according to

MISEV2023 guidelines. Transcriptomics (next-generation miRNA sequencing) and

proteomics (mass spectrometry-based) data are deposited in GEO and

ProteomeXChange Consortium via the PRIDE repository (publicly available at the time

of publication), respectively. Cell-cell communication was evaluated by integrating data

with the TISSUES database[39] and a publicly available scRNA-sequencing dataset of

carotid endarterectomy specimens containing plaque and marginal zones

(GSE159677).[20] Multi-omic network analysis was performed using BIONIC.[40] Functional

validation of predicted roles for EVs on angiogenesis was performed in Human

Umbilical Vein Endothelial Cell (HUVEC) spheroid-based sprouting assays.

Statistical Analysis

The Fisher's exact test tested the prevalence of gene ontology (GO) terms in whole

tissue proteomics. For non-omics data with sample sizes <10, a two-sided Mann-

Whitney or Wilcoxon (for paired samples) test was completed and for n≥10, data

distribution was assessed and differences between two groups were tested using a two-

tailed unpaired Student's *t*-test. For comparisons between three or more groups, a one-

way ANOVA was used with Šidák multiple comparisons test . Differences were

significant for values of $p < 0.05$ and are indicated in the figures. Proteomics (tissue and

EV) were quantified using Discover (Version 2.5) and peptides were filtered based on a

1.0% FDR. A minimum of at least 2 unique peptides for each protein was required for

the protein to be included in the analyses. Differential enrichment analysis was

performed using Qlucore Omics Explorer statistical software (Version 3.9; Qlucore,

Sweden) and differentially enriched proteins were calculated using a two-group

comparison with a Benjamini-Hochberg false discovery rate (FDR)-corrected p-value

(i.e., q-value) <0.05. Transcriptomic data (miRNA) was quantified in Partek Flow

(v10.0.23.0326) including trimming of adaptors, sequence alignment to genome (h38

assembly, Homo Sapiens, BioProject PRJNA31257, Bowtie), and quantification

(miRBase mature miRNAs, Version 22). Differential expression of miRNA in

experimental conditions was determined via DESeq2 with adjustment for multiple

comparisons using Benjamini and Hochberg's FDR step-up (adjusted p-value<0.05).

Predicted messenger RNA (mRNA) targets of differentially expressed EV-miRNA were

identified using MicroRNA Enrichment Turned Network (MIENTURNET version 2019-11-

25) using the miRTarBase database. Pathway analysis of differentially enriched EV-

miRNAs and EV-proteins were completed using Enrichr using the Gene Ontology

(Biological Process and Cellular Component) and Kyoto Encyclopedia of Genes and

Genomes(KEGG) databases with multiple testing correction using the Benjamini-

Hochberg procedure (adjusted p<0.05). Protein-protein interaction analysis was

completed in STRING v12.0. The TISSUES and Tabula Sapiens databases were

accessed through STRING v12.0 and Enrich, respectively. EV cellular identity

enrichement analysis was completed using the AddModule score function in Seurat

(Version 5.0.1) to assess the enrichment of EV protein signatures among different cell

types. The 'CellChat' R package (version 1.6.0) was used to infer cell interaction

networks by filtering the CellChat cell-cell communication ligand database for genes

associated with EV protein gene IDs. Comparison of fold enrichments between cohorts

regarding endothelial-related GO biological processes were performed using ANOVA

and Tukey's honest significant difference.

Results

Atherosclerotic tissue proteome and extracellular vesicle (EV) concentration identify a potential role for EVs in modulating the disease continuum

As recently shown by Alsaigh et al., single cell RNA sequencing (scRNA-seq) of matched human carotid marginal zones and plaque zones harbour different cellular subpopulations, revealing distinct microenvironments, and cellular communication strategies that may provide insights into disease progression.[20] To begin understanding the continuum of disease in carotid atherosclerosis, we delineated EV-based biology in plaque versus marginal zones from all patients (asymptomatic and symptomatic). Human carotid endarterectomy specimens for whole tissue proteomics (n=5) were dissected into plaque zones (PZ) and marginal zones (MZ) based on gross morphology by a single trained surgeon (Dr. S. Raju), with another 59 paired samples similarly dissected for EV enrichment and characterization (Figure 1A). Tissue histopathology with hematoxylin and eosin and Movat pentachrome stains delineated plaque and marginal zones to be different with distinguishing features of advanced atherosclerotic plaque morphology (e.g. necrotic core, intraplaque hemorrhage) visualized exclusively in plaque regions (Figure S1A-B).

Tissue proteomes (Figure S2A) from the plaque zones and marginal zones clustered separately by principal component analysis (Figure 1B donor corrected; Figure S2C donor uncorrected), with 514 differentially enriched proteins identified by zone (Figure

S2B). Pathway analysis of differentially enriched proteins in the marginal and plaque

zone (p_{adj}<0.05) using Gene Ontology (GO) and Kyoto Encyclopedia of Genes and

Genomes (KEGG) revealed atherogenic signatures, while GO terms containing 'vesicle'

were increased in plaque zones specifically (Figure 1C, S2D-F, S3A-B). We next

pursued EV enrichment in plaque and marginal zones separately. EVs were enriched

using size exclusion chromatography that included an additional density

ultracentrifugation step to remove extracellular matrix co-isolation[31] and was validated in

accordance with the MISEV2023 criteria (Figure 1D-G, S4A-F, S5A, S6). Briefly, EVs

were enriched in fractions 7-10 as evidenced by single vesicle analysis using

nanoparticle tracking analysis (NTA) and as a protein peak using the bicinchoninic acid

assay (Figure S4E). The secondary protein peak (Figure S4E) represents smaller

proteins that elute in the later fractions.[41] The presence of common EV protein markers

in per-fraction proteomics, including CD63, CD81, CD9, FLOT1, and an absence of the

cell protein marker calnexin confirmed successful EV isolation in our fractions of interest

(Figure 1D, left and right panels respectively; Figure S5A). Several vesicle-related

pathways were enriched in plaque zones compared to marginal zones (Figure S5B),

which may suggest increased EV processes taking place. We confirmed EV

morphology via direct visualization using cryogenic electron microscopy (cryo-EM),

which revealed bilayered nanoparticles with dense cores, as well as several muti-

compartmented vesicles, recently seen to be released by cells (Figure 1E).[42] We did not

observe any morphological or size differences between EV isolates from plaque and

marginal zones (Figure 1F). Paired plaque regions liberated similar EV size distribution,

but a larger quantity of EVs per milligram of tissue relative to marginal zones (Figure

1G). Notably, we found minimal lipoprotein (APOA1[43] and APOB100, enzyme-linked

immunosorbent assay, Figure S4C-D) and no whole-cell (flow cytometry, Figure S4F)

inclusion in vesicle preparations. Given EVs in marginal and plaque zones have the

potential to contribute to the local milieu, we next compared the biological cargo of EVs

between these two anatomical locations to help delineate the potential functional effects

of the EV-vesiculome in carotid atherosclerotic disease progression.

Carotid plaque EV-vesiculome (miRNA transcriptome and proteome) is distinct in

plaque versus marginal zones and drives pathways known to govern plaque

progression

Given that EV miRNA and protein cargo can mediate their functional effects on recipient

cells in local environments,[32-34] we performed next-generation miRNA sequencing and

proteomic analysis of EVs enriched from carotid plaque zones and marginal zones

(Figures S6, S7). Exploratory analysis with donor-corrected principal component

analysis (PCA) revealed that the EV-vesiculome clustered by location; specifically

plaque zone (PZ, purple) and marginal zones (MZ, pink; Figure 2A-B; donor

uncorrected Figure S7F). Differentially enriched protein pathway analysis predicted

biological interconnectability in plaque-derived EVs, as seen by greater than expected

protein-protein interactions (Figure 2C). Differential expression analysis between EV-

miRNA in plaque versus marginal zones revealed 371 EV-miRNA enriched in plaque

zones and 290 EV-miRNA enriched in marginal zones, across 20 paired patient

samples (Figure 2D). Top differentially expressed upregulated miRNA in the plaque

zones included miR-146a, miR-155, let-7a, miR-200b, miR-223, and miR181b, known to

have roles in secretion and response to inflammatory mediators and cell proliferation

and differentiation.[44-54] Among differentially enriched proteins, we found 282 proteins

enriched in plaque zones, and 379 enriched in marginal zones, across 23 paired patient

samples (Figure 2E). Like the EV-miRNAs, several known mediators of plaque

progression including metalloproteinases (MMP9), cell adhesion proteins (ICAM1),

lipoprotein metabolism proteins (LPL), complement and coagulation proteins (F2, F10),

and inflammatory mediators (CRP, SAA2-4, BLVRA, BLVRB) were more abundant in

EVs from plaque environments (Figure 2E).

To explore the potential functions of EV-miRNA and protein cargo, we queried KEGG

pathways using the Enrichr interface comparing plaque zone versus marginal zone EV

cargo.[55] Carotid plaque EV-cargo represented 155 unique KEGG pathways

(FDR<0.05), with 60 pathways specific to EV-miRNAs and 60 pathways specific to EV-

proteins (Figure 2F, Venn diagram). Top unique pathways predicted to be modulated by

plaque EV-miRNA included cell cycle, apoptosis, and inflammatory signaling pathways

(TNF, T cell receptor, TGF-β, chemokine, NFκB; Figure 2F, top right panel). Conversely,

top unique pathways of differentially expressed plaque EV-proteins include complement

and coagulation, phagosome, cholesterol metabolism, cell adhesion, and vascular

smooth muscle contraction (Figure 2F, bottom right panel). Key pathways involved in

atherosclerotic plaque progression shared between the EV-miRNA and EV-proteome

included cellular senescence, cell adhesion and migration, leukocyte transendothelial

migration, fluid shear stress and atherosclerosis, lipid and atherosclerosis, and amino

acid and pyruvate metabolism (Figure 2F, bottom left panel). These data suggest that in

the plaque the differentially enriched EV-vesiculome (i.e., protein and miRNA) may

function together, as well as independently, to modulate several pathways known to be

involved in driving plaque biology. Given EVs are a vector of cell-cell communication

between donor and recipient cells, we next attempted to determine the cellular origin of

human plaque EVs to delineate the cellular contributors to plaque EV and further, how

they might change with disease progression from marginal to plaque zones.

Total EV-proteomics suggests diverse EV cell of origin and potential targets

modulated in recipient cells

EVs carry cargo and membrane proteins that are reflective of their cell of origin.[56,57]

Cells secrete EVs with distinct biological cargo to relay specific messages to recipient

cells. To date, little is known about the cellular communication occurring *in vivo* within

the plaque niche via EVs. We used two complementary approaches to infer the EV cell

of origin using our EV proteomics. We started by exploring our total EV proteomics (i.e.

plaque and marginal regions combined) within the TISSUES database[39] which

integrates evidence on tissue expression from proteomic and transcriptomic screens.

The predicted cellular EV origin included a range of expected cells including endothelial

(neointima), several immune (B-lymphoblastoid, plasma cell, bone marrow cell, and

foam cell), and smooth muscle cells (Figure 3A; Figure S8A,C). The analysis was then

refined with EV-protein signature enrichment analysis among carotid artery cells

specifically, by integrating our complete list of total detected EV-proteins with a publicly

available scRNA-seq dataset containing paired plaque and marginal region samples

(n=6 samples, Figure S9A-G).[20] Data integration corroborated the EV cell of origin,

identifying strong EV signatures from vascular smooth muscle cells, endothelial cells,

dendritic cells, macrophages and lymphocytes (Figure 3B). Likewise, examination of

EV-miRNA normalized counts in our plaque and marginal zones identified several

miRNAs that are frequently ascribed to endothelial,[45,58-62] smooth muscle,[63-65] and

lymphocyte cells,[47,66-70] suggesting EV cell of origin signatures that align with those

predicted by the EV-proteome (Figure S10A).

Beyond the suspected cell of origin, EV surface proteins and protein cargo deliver

ligands to recipient cells within the atherosclerotic plaque environment – the biological

consequences of this interaction will depend on the protein repertoire of the recipient

cell. Using CellChat[71], we predicted EV ligand communication (incoming signaling) with

recipient cells that would be found in atherosclerosis using the same publicly available

scRNA-seq dataset as above. Global EV-cell communication analysis was performed by

specifically filtering for known receptor-ligand interactions where ligands were derived

from EV proteomics and receptors from scRNA-seq. When the total EV proteome was

applied to scRNA-seq data annotated as plaque or marginal, it identified the predicted

distribution of cellular EV source by each region (Figure 3C). Inferred cell

communication pathways used total EV-associated ligands with total scRNA-seq (i.e.

present in both plaque and marginal regions) (Figure 3D), as well as total EV-ligands

separated by marginal and plaque regions (Figure S10C). The strongest global EV-cell

interactions were seen among immune cells including macrophages and T-cells, with

the enrichment of inflammatory signaling patterns (CypA, MIF, MHC-I, CXCL,

prostaglandin, MHC-II). The predicted distribution of enriched global EV signaling

pathways place ApoE, tenascin, cholesterol, and visfatin more likely within plaque zones

(Figure S10B), while cord diagrams demonstrate predicted global EV ligand-receptor

pairs with endothelial, macrophage, and vascular SMC populations (Figure S10D left,

middle, and right panels, respectively).

Approaching EV-miRNA communication within tissue is more complex. Using the

predicted messenger RNA (mRNA) targets of EV-miRNA, the genes that could be

altered can be determined and correlated with single cell transcriptomics to understand

potential EV-miRNA communication to recipient cells. We similarly used two approaches

to annotate expected cellular targets of plaque EV-miRNAs - Tabula Sapiens database[72]

(vascular tissue) and the publicly available scRNA-seq carotid dataset as above (Figure

3E-F). All differentially expressed EV-miRNA had over 1956 (FDR<0.05) experimentally

validated (miRTarBase)[73] mRNA targets expressed in lymphocytes, endothelial cells,

SMCs and macrophages (Figure 3E). Plaque EV-miRNA to mRNA gene-set enrichment

analysis with the carotid scRNA-seq dataset corroborated these findings, wherein

targets of plaque EV-miRNAs were enriched in lymphocytes, endothelial cells, vascular

SMCs, and macrophages (Figure 3F).

Plaque and marginal zone EV-proteomes predict different EV cells of origin,

recipient cell repertoires, and expected communication with vascular cells

As the microenvironments of plaque and marginal zones are likely distinct (also

implicated by initial exploration, Figure 3C), we recapitulated our approach to infer EV

cellular origin and predicted targets in recipient cells by region using differentially

enriched EV-proteins and differentially expressed EV-miRNA between plaque versus

marginal zones. When delineated by region, plaque zones appear to have greater

representation of macrophages as sources of EVs than marginal zones, where vascular

SMCs predominated (Figure 4A-B). TISSUES database supported this finding, with our

plaque and marginal zones highlighting diverse cellular EV sources that ranked

differently by region (Figure 4C-D, S8B-C), predominantly reflecting cells that would be

found in atherosclerosis despite the broader tissue sources included in this dataset.

We then pursued EV-cell communication signals within plaque and marginal regions

separately, by filtering EV ligands as either plaque-enriched EV-proteins or marginal-

enriched EV-proteins. Known ligand-receptor interactions were then explored using

CellChat as above. In comparison to marginal zones (Figure 4E), plaque enriched EV-

protein ligands were predicted to participate in more diverse and distinct cellular

communication strategies when paired with total scRNA-seq (i.e. present in both plaque

and marginal regions) (Figure 4F). For example, several predicted inflammatory ligand-

receptor interactions emerged in the plaque zone with ApoE, complement, ICAM,

cathepsin (CTSG), and cholesterol signatures. Similar increased communication

diversity was predicted in plaque zones when analysis was restricted to single cells in

marginal and plaque regions separately (Figure S11A-B). Chord communication

diagrams were then created to explore region-specific cell-cell communication.

Differentially enriched EV-protein ligands from our marginal and plaque zones were

paired with publicly available single cell transcripts (potential receptor recipients) from

either endothelial cells (Figure 4G), vascular SMCs (Figure 4H), or macrophages

(Figure S11C) found in respective marginal regions (left panels) or plaque regions (right

panels). For each cell type examined, plaque zones exhibited increased predicted

ligand-receptor interactions relative to marginal zones. Notably, proposed plaque EV-

cell communication included endothelial integrins (needed for cell motility), vascular

SMC integrins and AXL (phagocytosis), and macrophage complement and cell adhesion

(C3AR1, CD44). Together, EVs are predicted to communicate with several

atherosclerosis-relevant cell types, with regional (plaque and marginal) differences in

communication patterns.

Carotid plaque EVs are increased in number and contain EV-cargo (miRNA and

protein) that are differentially enriched based on plaque vulnerability

Understanding the biological potential and EV-cell communication that takes place

along the continuum of asymptomatic to symptomatic carotid disease will provide new

insights into this complex clinical problem. We assessed differences in EV-cargo

between asymptomatic (n=10 patients) and symptomatic (n=10-13 patients) groups by

anatomical zones (marginal and plaque; Figure 5A). Blinded pathological grading

performed by a staff pathologist (Dr. M. Seidman) of a representative plaque samples

(n=5) demonstrated fibroatheroma and calcification in all (AHA grade V-VI),[38,74] with

evidence of intraplaque hemorrhage reported in all symptomatic patients (Figure 5B,

S12A). While both asymptomatic and symptomatic plaque zones contained more EVs

per milligram of tissue compared to their matched marginal zones, only symptomatic

patients achieved statistical significance (Figure 5C-D, S12B). Morphologically, carotid

EVs were similar in symptomatic and asymptomatic groups (size and cryo-EM, Figure

5E). Direct comparisons by symptom status within plaque and marginal zones did not

yield significant differences in EV quantification highlighting the importance of matched

marginal zone controls (Figure S12C-D).

When analysis of EV-cargos was performed, there were consistently larger numbers of

differentially enriched EV-miRNA and EV-proteins in the symptomatic group in both

marginal zones (delineated by negative fold change) and plaque zones (delineated by

positive fold change) (Figure 5F-I – volcano plots), suggesting increased biological

complexity in clinically vulnerable plaques and their corresponding marginal regions for

symptomatic patients.

We next assessed specific differences in EV-cargo between asymptomatic plaques

versus paired marginal zones and separately, EV-cargo differences between

symptomatic plaques versus paired marginal zones (Figure 6). There were 306 unique

differentially expressed EV-miRNAs in symptomatic tissue compared to only 33 unique

EV-miRNA in asymptomatic tissue (Figure 6A). Overlap was seen with 364 differentially

expressed EV-miRNAs in the two groups. Pathway analysis of unique EV-miRNA

symptomatic predicted targets revealed more GO and KEGG pathways (Figure 6B).

Unique signatures in asymptomatic tissues included steroid biosynthesis, ABC

transporters, and the AMPK signaling pathway (Figure 6C). Pathways derived using

unique symptomatic EV-miRNAs were numerous and included lipid and atherosclerosis,

inflammatory signalling (HIF-1, TGFβ, TNF) and NOTCH signaling (endothelial

angiogenesis) (Figure 6D). Similarly, 435 unique differentially enriched EV-proteins were

seen in symptomatic tissue, compared to 6 in asymptomatic (Figure 6E) and functioned

with a high degree of biological interconnectibility (Figure S12E). Similar to the EV-

miRNA analysis, there were an increased number of GO and KEGG pathways

modulated by the symptomatic plaque EV-protein contents (Figure 6F). KEGG pathway

analysis of the unique EV-proteins from asymptomatic tissue identified 6 KEGG

pathways functioning in amino acid and nutrient metabolism, antigen presentation, renin

secretion, and complement and coagulation with only a single EV-protein (annotated in

figure) contributing to each pathway protein (Figure 6G). The unique EV-proteins from

symptomatic tissue identified several biological pathways with the top differentially

expressed pathways including complement and coagulation, cell motility (regulation of

actin cytoskeleton, focal adhesion, ECM-receptor interaction, leukocyte transendothelial

migration), and cholesterol metabolism (Figure 6H, left panel). Several proteins were

enriched in the symptomatic plaque EVs that demonstrated protein-protein interactions,

forming three units (complement and coagulation cascades, cholesterol metabolism,

and regulation of actin cytoskeleton) predicted to modulate these pathways (Figure 6H,

right panel). Together, these vesiculomic data highlight the potential for EVs to govern

several well-known aspects of plaque biology, with increasing complexity as the disease

continuum progresses.

Predicted EV-based cell communication in the symptomatic plaque reveals the

emergence of an endothelial cell signature with functional effects on

angiogenesis

The differences in EV cargo between asymptomatic and symptomatic groups led us to

query differences in predicted cellular communication based on symptom status. EV

cellular identity was inferred by integrating our enriched EV-protein signatures from

either marginal or plaque zones according to symptom status with carotid marginal and

plaque scRNA-seq data (Figure 7A-D). Greater endothelial cell (EC) enrichment was

predicted in symptomatic plaque signatures relative to asymptomatic plaque signatures

(purple bars highlight each EC subpopulation, with increasing prevalence noted as a

shift to the left; Figure 7B,D), possibly suggesting a heightened role of endothelial cell-

derived EVs in governing EV-cell communication with atherosclerotic plaque

progression. No notable differences in cell type enrichment was observed for marginal

signatures (Figure 7A,C). Potential recipient cell repertoires available in asymptomatic

and symptomatic plaques was also explored through EV-miRNA-mRNA predicted

targets (Figure S13).

We then explored differences in asymptomatic and symptomatic plaque EV-cell

communication with CellChat, by similarly examining EV-proteins as the ligands in

known ligand-receptor interactions, where receptors were predicted from scRNA-seq in

marginal and plaque zones (Figure 7E-F). Although asymptomatic plaques shared all

EV-cell signaling pathways with the symptomatic group (Figure 7E-F), symptomatic

plaque EV-protein ligands also exhibited unique signaling involving ApoE, galectin, and

visfatin. Specifically, EV/visfatin to endothelial cell-based communication was unique to

symptomatic plaques – notable given angiogenesis can be modulated through visfatin-

based secretion of vascular endothelial growth factor (VEGF) and fibroblast growth

factor 2 (FGF-2),[75,76] monocyte recruitment through monocyte chemoattractant protein 1

(MCP-1),[77] and extracellular matrix remodelling via matrix metalloproteinase

expression.[78,79] We then explored specific ligand-receptor pairs between asymptomatic

and symptomatic plaque EVs and recipient cell repertoires that might be available in

ECs (Figure 8A, S14A), SMCs (Figure S14B-D), and macrophages (Figure S14E-G).

Cell-cell communication strategies predicted in the asymptomatic plaques were mostly

recapitulated in the symptomatic group, with symptomatic plaques exhibiting more

complex EV-recipient cell signaling (Figure 8A, S14D,G). As an approach to externally

validate some of our cell communication predictions, we examined the EV marker CD63

as well as CD63-normalized visfatin and CD63-normalized VEGFA in a cohort of 18

carotid endarterectomy specimens (Biobank of Karolinska Endarterectomy, BiKE) where

proteomics had been completed on plaque zones and marginal zones (*n*=9

symptomatic, *n*=9 asymptomatic).[37] While these were bulk tissue proteomics, more of

the common EV marker CD63 protein was detected in symptomatic plaques and

increased visfatin (NAMPT) and VEGFA was detected in symptomatic carotid tissue

when normalized to CD63 protein by zone (Figure S15A-C).

Given the distinct EV-based communication strategies predicted in symptomatic versus

asymptomatic tissues, we then assessed the potential functional consequences through

analysis by GO and KEGG pathways for EV-miRNA, EV-proteins, and EV-vesiculome

(integrated analysis) (Figure S16A,B). We found differences in pathways predicted to be

modulated by EVs in symptomatic compared to asymptomatic plaques. Key EV-miRNA

based pathways enriched in symptomatic plaque included inflammatory signaling (TCR

signaling, NFkB, HIF-1, TNF, TGFβ), apoptosis, cell senescence, and cell

migration/motility (adherens junctions, VEGF) (Figure S16A,B left panels). Enriched

pathways in symptomatic plaques predicted to be modulated by EV-proteins included

complement and coagulation (includes fibrinolysis, iron metabolism), cholesterol

metabolism (includes lipid transport, response to low density lipoprotein particle

stimulus), and amino acid metabolism (Figure S16 A,B middle panels). Network analysis

integrating EV-miRNA and EV-protein content identified the importance of metabolic

pathways (amino acid biosynthesis, RNA synthesis and turnover, steroid biosynthesis,

cholesterol), complement and coagulation, and inflammatory pathways (antigen

processing and presentation, macrophage proliferation, neutrophil degranulation) as

shared targets of EV-contents in symptomatic plaques (Figure S16 A-B right panels).

Likewise, overrepresentation analysis of differentially enriched EV-miRNA and EV-

protein predicted differential representation of GO and KEGG pathways that based on

symptom status (Figure S17A-D). Upon closer examination there was a clear

endothelial and angiogenic signature in EV-miRNA-mRNA overrepresentation analysis

in symptomatic versus asymptomatic EVs with overrepresented predicted pathways in

angiogenesis, blood vessel development, sprouting angiogenesis and VEGF signaling

(Figure 8B, S18A-B), while EV-protein and EV-vesiculomics overrepresentation analysis

were more varied (Figure S18C-F, S19A-D).

As significantly more endothelial angiogenesis-related pathways were identified in

symptomatic plaque zones versus their respective marginal zones (Figure 8C, p-value =

1.84×10^{-23}), we tested the functional effects of symptomatic plaque EVs (n=3) using an

endothelial spheroid-based sprouting assay (Figure 8D-F, S19E-J). Endothelial sprouts

were significantly increased in number (110.2 ± 4.34 versus. 92.3 ± 2.59; $p = 2.7 \times 10^{-2}$)

(Figure 8E) as well as total sprout length (244.9 ± 5.06 versus 179.7 ± 4.27 μm; $p < 1.0 \times 10^{-10}$

) (Figure 8F) and average sprout length (Figure S19G) after treatment with

symptomatic plaque EVs compared to matched marginal zone EVs. Controls included

eluates (negative control, Figure 8E-F, S19F-G) and VEGF (positive control, Figure

S19F,H-J). Taken together, these results highlight the biologically relevant EV-protein

and EV-miRNA cargoes that exist within the plaque and marginal zones of carotid tissue

along the continuum of asymptomatic to symptomatic disease.

Discussion

It is currently unknown what renders a carotid atherosclerotic plaque vulnerable to

rupture and leads to stroke. Cell-cell communication – including through extracellular

vesicles (EVs)[31] – may play a role. Our study evaluates carotid tissue EV-

transcriptomics and EV-proteomics to understand changes in EV-cargo in diseased

states (plaque versus marginal zones) along the clinical continuum (asymptomatic to

symptomatic) to predict EV source, potential recipient cell responses, and a relevant

biological functional effect based upon these predictions. Increased numbers of EVs

containing distinct cargos were discovered in carotid plaque versus matched adjacent

marginal zones, implicating their role in governing plaque biology. Given EV-based cell-

cell communication necessitates an EV donor and an EV recipient cell, we integrated

publicly available datasets (TISSUES[39], Tabula Sapiens[72] and carotid plaque scRNA-

seq datasets[20]) with our region-specific carotid EV-vesiculomics to gain insight into the

cellular drivers of EV-based communication within the plaque. As plaque vulnerability

increased from asymptomatic to symptomatic disease states, there was a greater

degree of EV communication suggested in the symptomatic plaques. Our *in silico*

analysis suggested an increased abundance of endothelial cell-derived EVs with plaque

progression and predicted a greater number of unique EV-endothelial cell ligand-

receptor pairs in symptomatic plaques known to function in angiogenesis, with

downstream overrepresentation analysis corroborating increased endothelial-based

pathways. Intriguingly, we discovered an EV-derived endothelial/neovascularization

'signature' in symptomatic carotid plaques and then showed for the first time, that

human symptomatic carotid plaque EVs induce angiogenesis in a model of endothelial

sprouting. Taken together, these exciting data illustrate a complex communication

network, with renewed focus on the role of EVs and the endothelium.

EVs are mediators of cardiovascular disease states and have previously been shown to

be increased in patient samples.[31,80-84] Here, we optimized and validated a protocol to

enrich tissue-entrapped atherosclerotic plaque-derived EVs using size exclusion

chromatography and validated our EVs in accordance with the MISEV2023 criteria,[26] in

alignment with our previous EV isolation approaches using similar methodology.[31] We

found increased EVs per gram of tissue from plaque regions in comparison to their

matched adjacent marginal regions. Recent work has delineated an increase in EV

signature based on pathway analysis of differential protein expression from diseased

areas (carotid plaque and aortic valve).[31] EVs mediate their effects via delivering their

cargo such as miRNAs and proteins to recipient cells.[32-34] In carotid tissue, this has

been difficult to test empirically. To date, there is evidence that carotid plaque-derived

microparticles are thrombogenic[27] and that candidate EV-cargo (from multiomics

analysis of carotid tissue) can increase calcification in cultured human carotid SMCs.[31]

Given the challenges working with EVs[31,85], there has been no study to date using

human carotid plaque derived EVs in functional assays to determine the biological

consequences of altered EV-cargo in diseased states. By examining the EV-vesiculome

within our carotid plaque versus marginal zones, plaque EV-contents were predicted to

be involved in several pro-atherogenic pathways such as complement and coagulation,

fluid shear stress, cellular senescence, cell proliferation and differentiation, and immune

pathways. As EVs carry proteins that are indicative of their cell of origin,[56,86] we

harnessed an opportunity to examine our unique dataset within publicly available single

cell RNA -sequencing datasets to explore potential EV-based cell-cell communication in

the carotid atherosclerotic plaque milieu.

To assess EV-based communication in the plaque microenvironment, we layered our EV-vesiculomics with the only published carotid plaque scRNA-seq dataset that contained site-specific data (plaque zone and marginal zone) to reveal potential EV donor cell identities and communication partners in our two zones of interest.[20,56,86] Site-specific layering is important in atherosclerotic disease given the cellular plasticity seen in plaque progression.[12,87,88] Leveraging tissue-specific scRNA-seq data to inform the probable cell of origin of EVs has recently gained acceptance as a starting point in complex human tissue.[57] The differences in EV cell of origin identity with vascular SMCs predominating in marginal zones and immune cells in plaques zones is in keeping with recent reports of different identities and quantities of various cell populations found in plaques versus the adjacent region.[20] The role of SMCs in regulating atherosclerotic plaque progression through phenotypic switching in atherogenic environments[11,87] is well characterized and our data now suggest that SMCs could also be important sources of EVs in marginal zones, beyond their roles within advanced plaques.[12,20,88,89] The role of immune cells in governing advanced plaque biology is similarly acknowledged, and likely explains the increased EV contribution predicted from these cells in plaque regions.[90,91] Intriguingly, the ligand-receptor communication strategies (using EV-protein cargo as ligands and recipient cell repertoires from scRNA-seq) that were predicted in the plaque versus marginal zones suggested increasing complexity of the biological milieu along the disease continuum.[20,31] Our analysis implicates EVs as integral mediators of atherosclerotic plaque progression and EV-based communication as a potential target to modulate plaque biology.

Assessment and treatment of vulnerable carotid atherosclerotic plaques represents a

continuing clinical challenge with opportunities to improve.[92] With symptomatic plaques

representing culprit lesions with advanced vulnerable atherosclerotic disease where

participation from resident and circulating cells that are phenotypically dynamic

contribute to the EV-pool, disease progression lends itself to multifaceted

communication patterns.[93] We defined a predicted role for endothelial cells at both

facets (donor and recipient) of the EV cell-cell communication paradigm in symptomatic

plaques, where endothelial cells featured prominently as both potential sources of EVs

and responders to EVs – specifically with communication signaling known to function in

neovascularization and angiogenesis. Neovascularization in atherosclerotic plaque

occurs as a result of hypoxemia and leads to the growth of immature and leaky vessels

causing intra-plaque hemorrhage, a well known contributor to plaque instability and

rupture.[37,94-96] While not unique to symptomatic plaques, it does predominate in this

group of patients.[97] With this insight, we corroborated our predictions functionally and

showed for the first time that symptomatic human carotid plaque zone EVs induce

angiogenesis using an optimized low-input human endothelial spheroid sprouting assay.

The importance of endothelial EV-based communication in driving atherosclerotic

plaque instability represents an opportunity to understand the disease process from the

less-studied endothelial-centric lens. Endothelial cell-EVs are intimately involved in

autocrine and paracrine communication with cells of the atherosclerotic plaque where

they orchestrate upregulation of adhesion molecules, and activation, proliferation, and

migration of vascular cells.[7,98-104] In the atherosclerotic milieu, endothelial cells are

exposed to disturbed flow and inflammatory mediators, and can undergo endothelial-to-

mesenchymal transition wherein they lose endothelial identify and adopt mesenchymal

properties.[9,10,105] In the scRNA-seq dataset from Alsaigh *et al*, there were three

endothelial cell populations contributing to plaque-derived EVs as seen in the original

scRNA-sequencing dataset that represented quiescent, activated, and

transdifferentiated endothelial cells.[20] Given the quiescent endothelium has been shown

to participate in atheroprotective EV-messaging[104], it would be beneficial to resolve the

differences in communication between these cell types and understand how to shift the

messaging to a quiescent phenotype. The important role of endothelial cell-based EV

communication in carotid atherosclerotic plaque combined with the accessibility of the

endothelial layer, emphasizes the potential of endothelial-EV targeted therapeutics in

atherosclerotic vascular disease.

Our study explored EV-based cell-cell communication in human atherosclerotic disease

and how this might dictate plaque vulnerability in part through the functional role of EVs

in endothelial angiogenesis. There are limitations, however. While we completed tissue

EV-transcriptomics and EV-proteomics, it was not possible to generate matched scRNA-

sequencing that would have allowed us to annotate the cells and communication

occurring within the same carotid tissue.[57] The external dataset selected was the only

publicly available scRNAseq on plaque and marginal zones, however it did not contain

symptomatic patients. Given the cell identity changes seen with plaque progression, cell

annotations and ligand-receptor interactions may not have captured the diversity of the

symptomatic plaque cell populations specifically. *In silico* analysis only enables

predictions for delineating EV cells of origin and potential recipient cell responses, which

could be a limitation across all samples – regardless of patient symptom status. Lastly,

EVs are one of many extracellular signals in the plaque microenvironment. While it is

encouraging that increased CD63 (common EV marker) was similarly detected in

symptomatic plaque zones in the BiKE dataset, and was significantly increased in the

most severe Cluster 3 carotid plaques described by Mokry et al,[106] the role of

lipoproteins, non-EV inflammatory or angiogenic processes, and metabolites cannot be

excluded.[107] Our work does not preclude the role of other signalling molecules, nor can

it account for the dynamic, spatial, and other physical elements taking place. As

technologies progress, it may become possible to validate our findings at a single EV-

level.

Conclusions

This work was pursued with the challenging goal of establishing an important

conversation around EVs in carotid plaque biology. Crucially, vulnerable symptomatic

carotid disease was characterized by an increased concentration of EVs in plaque

zones, with increased contribution from endothelial cell sources. Furthermore, EVs

harbored a signature for specific EV-based communication with endothelial cells, with

predicted impact on angiogenic pathways. These predictions were corroborated

functionally by demonstrating symptomatic carotid plaque zone EVs increase

endothelial sprouting angiogenesis – a signature that might imply neovascularization

and plaque rupture potential. These provocative results underscore the importance of

the endothelium in governing vulnerable biology and highlight unique opportunities to

consider the role of EVs as a therapeutic target in plaque stabilizing therapies.

Acknowledgements

The authors thank the patients, surgical teams, and stroke neurology colleagues at University Health Network and Sunnybrook Health Sciences. For assistance with pathology: Ms. Melanie Peralta (UHN). Electron microscopy imaging of EVs was performed in the Harvard Medical School Electron Microscopy Facility.

Sources of Funding

This work was supported by Canadian Institutes of Health Research (CIHR) Project Grants PJT178006 (K. Howe) and PJT148487 (J. Fish), a Tier II Canada Research Chair from CIHR (J. Fish), Heart and Stroke Foundation of Canada (New Investigator Award, K. Howe), Vascular Cures (Wylie Scholar Award, K. Howe), Blair Early Career Professorship in Vascular Surgery (K. Howe), Peter Munk Cardiac Centre (K. Howe), University Health Network (K. Howe) and Vanier Canada Graduate Scholarship (S. Raju). Infrastructure funding was obtained from the John R. Evans Leaders Fund (J. Fish). E. Aikawa lab is supported by NIH grants (R01 HL136431, R01 HL141917, R01 HL147095. M. E. Turner is supported by a CIHR Banting Fellowship. L.M. is the recipient of fellowships and awards from the Swedish Research Council [VR, 2023-02724, 2019-02027], Karolinska Institute Consolidator program, Swedish Heart-Lung Foundation [HLF, 20240094, 20230357, 20210466, 20200621, 20200520, 20180244, 20180247, 201602877], Swedish Society for Medical Research [SSMF, P13-0171]. L.M. also acknowledges funding from Mats Kleberg's, Sven and Ebba-Christina Hagberg's, Tore Nilsson's, Magnus Bergvall's and Karolinska Institute research (KI Fonder) and

713 doctoral education (KID) foundations. Project funding was also obtained by U.H. from

714 the Swedish Heart-Lung Foundation (20180036, 20170584), the Swedish Research

715 Council (2017-01070, 2019-02027), and King Gustav Vth and Queen Victoria's

716 Foundation.

Disclosures

721 Natalie Galant is co-founder and CEO of Paradox Immunotherapeutics, but the content

722 of this manuscript is unrelated to the work of this company.

Clinical Perspective

What Is New?

- First analysis of extracellular vesicle (EV) miRNA and proteomics in carotid atherosclerosis by region (plaque zone and marginal zone) across the clinical spectrum (asymptomatic and symptomatic disease).

- Integration with single-cell RNA-sequencing of carotid plaque and marginal zones revealed predicted EV cell of origins and EV-ligand-receptor interactions with an endothelial signature involved in neovascularization of vulnerable (symptomatic) plaque zones.

- Functional assays corroborated the endothelial *in silico* predictions, with EVs from symptomatic carotid plaque zones increasing endothelial sprouting (number and length) compared to matched marginal zone EVs or controls.

What Are the Clinical Implications?

- Cell-cell communication in atherosclerotic plaques reflects a dynamic milieu that can eventually render a plaque vulnerable to rupture and cause stroke, but needs further study to better reveal novel therapeutic targets for plaque stabilization.

- Tissue-based EVs are an increasingly recognized means for cell-cell communication, with EVs from donor cells (source) being taken up by neighboring cells (recipients) to govern their function – these communication patterns need to be understood along the continuum of cellular identity and local plaque environments.

748   • Discovery of an endothelial EV communication network in symptomatic carotid

749   plaques capable of neovascularization represents a new approach to consider for

750   plaque stabilization therapies.

References

1.   Libby P. The changing landscape of atherosclerosis. *Nature.* 2021;592(7855):524-533.
2.   Libby P, Buring JE, Badimon L, et al. Atherosclerosis. *Nature Reviews Disease Primers.*
2019;5(1):56.
3.   Shahjouei S, Sadighi A, Chaudhary D, et al. A 5-decade analysis of incidence trends of
ischemic stroke after transient ischemic attack: a systematic review and meta-analysis.
JAMA neurology. 2021;78(1):77-87.
4.   Howe KL, Cybulsky M, Fish JE. The Endothelium as a Hub for Cellular Communication in
Atherogenesis: Is There Directionality to the Message? *Front Cardiovasc Med.*
2022;9:888390.
5.   Yu L, Xu L, Chu H, et al. Macrophage-to-endothelial cell crosstalk by the cholesterol
metabolite 27HC promotes atherosclerosis in male mice. *Nat Commun.* 2023;14(1):4101.
6.   Yurdagul A, Jr. Crosstalk Between Macrophages and Vascular Smooth Muscle Cells in
Atherosclerotic Plaque Stability. *Arterioscler Thromb Vasc Biol.* 2022;42(4):372-380.
7.   Hergenreider E, Heydt S, Tréguer K, et al. Atheroprotective communication between
endothelial cells and smooth muscle cells through miRNAs. *Nature cell biology.*
2012;14(3):249-256.
8.   Raju S, Botts SR, Blaser MC, et al. Directional Endothelial Communication by Polarized
Extracellular Vesicle Release. *Circulation Research.* 2024;134(3):269-289.
9.   Chen P-Y, Qin L, Baeyens N, et al. Endothelial-to-mesenchymal transition drives
atherosclerosis progression. *The Journal of clinical investigation.* 2015;125(12):4514-
4528.
10.   Evrard SM, Lecce L, Michelis KC, et al. Endothelial to mesenchymal transition is common
in atherosclerotic lesions and is associated with plaque instability. *Nature*
communications. 2016;7(1):11853.
11.   Gomez D, Owens GK. Smooth muscle cell phenotypic switching in atherosclerosis.
Cardiovascular research. 2012;95(2):156-164.
12.   Alencar GF, Owsiany KM, Karnewar S, et al. Stem cell pluripotency genes Klf4 and Oct4
regulate complex SMC phenotypic changes critical in late-stage atherosclerotic lesion
pathogenesis. *Circulation.* 2020;142(21):2045-2059.
13.   Group ECSTC. Randomised trial of endarterectomy for recently symptomatic carotid
stenosis: final results of the MRC European Carotid Surgery Trial (ECST). *The Lancet.*
1998;351(9113):1379-1387.
14.   stenosis Beoceispwh-gc. North American symptomatic carotid endarterectomy trial
collaborators. *N Engl J Med.* 1991;325(7):445-453.
15.   Walker MD, Marler JR, Goldstein M, et al. Endarterectomy for asymptomatic carotid
artery stenosis. *Jama.* 1995;273(18):1421-1428.
16.   Brott TG, Hobson RW, Howard G, et al. Stenting versus endarterectomy for treatment of
carotid-artery stenosis. *New England Journal of Medicine.* 2010;363(1):11-23.
17.   Hackam DG. Optimal medical management of asymptomatic carotid stenosis. *Stroke.*
2021;52(6):2191-2198.

18. Committee W. European Society for Vascular Surgery (ESVS) 2023 Clinical Practice
Guidelines on Radiation Safety. *European Journal of Vascular and Endovascular Surgery.*
2022.
19. Goessens BM, Visseren FL, Kappelle LJ, Algra A, van der Graaf Y. Asymptomatic carotid
artery stenosis and the risk of new vascular events in patients with manifest arterial
disease: the SMART study. *Stroke.* 2007;38(5):1470-1475.
20. Alsaigh T, Evans D, Frankel D, Torkamani A. Decoding the transcriptome of calcified
atherosclerotic plaque at single-cell resolution. *Communications biology.*
2022;5(1):1084.
21. Theofilatos K, Stojkovic S, Hasman M, et al. Proteomic atlas of atherosclerosis: the
contribution of proteoglycans to sex differences, plaque phenotypes, and outcomes.
Circulation Research. 2023;133(7):542-558.
22. Tomas L, Edsfeldt A, Mollet IG, et al. Altered metabolism distinguishes high-risk from
stable carotid atherosclerotic plaques. *European heart journal.* 2018;39(24):2301-2310.
23. Vaisar T, Hu JH, Airhart N, et al. Parallel murine and human plaque proteomics reveals
pathways of plaque rupture. *Circulation research.* 2020;127(8):997-1022.
24. Colombo M, Raposo G, Théry C. Biogenesis, secretion, and intercellular interactions of
exosomes and other extracellular vesicles. *Annual review of cell and developmental*
biology. 2014;30:255-289.
25. Théry C, Witwer KW, Aikawa E, et al. Minimal information for studies of extracellular
vesicles 2018 (MISEV2018): a position statement of the International Society for
Extracellular Vesicles and update of the MISEV2014 guidelines. *Journal of extracellular*
vesicles. 2018;7(1):1535750.
26. Welsh JA, Goberdhan DC, O'Driscoll L, et al. Minimal information for studies of
extracellular vesicles (MISEV2023): From basic to advanced approaches. *Journal of*
extracellular vesicles. 2024;13(2):e12404.
27. Leroyer AS, Isobe H, Lesèche G, et al. Cellular origins and thrombogenic activity of
microparticles isolated from human atherosclerotic plaques. *Journal of the American*
College of Cardiology. 2007;49(7):772-777.
28. Rautou P-E, Leroyer AS, Ramkhelawon B, et al. Microparticles from human
atherosclerotic plaques promote endothelial ICAM-1–dependent monocyte adhesion
and transendothelial migration. *Circulation research.* 2011;108(3):335-343.
29. Haraszti RA, Didiot M-C, Sapp E, et al. High-resolution proteomic and lipidomic analysis
of exosomes and microvesicles from different cell sources. *Journal of extracellular*
vesicles. 2016;5(1):32570.
30. Stamatikos A, Knight E, Vojtech L, et al. Exosome-mediated transfer of anti-miR-33a-5p
from transduced endothelial cells enhances macrophage and vascular smooth muscle
cell cholesterol efflux. *Human gene therapy.* 2020;31(3-4):219-232.
31. Blaser MC, Buffolo F, Halu A, et al. Multiomics of tissue extracellular vesicles identifies
unique modulators of atherosclerosis and calcific aortic valve stenosis. *Circulation.*
2023;148(8):661-678.
32. Valadi H, Ekström K, Bossios A, Sjöstrand M, Lee JJ, Lötvall JO. Exosome-mediated
transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between
cells. *Nature cell biology.* 2007;9(6):654-659.

838 33. Jiang S, Li X, Li Y, et al. APOE from patient-derived astrocytic extracellular vesicles alleviates neuromyelitis optica spectrum disorder in a mouse model. *Science Translational Medicine.* 2024;16(736):eadg5116.

841 34. Antounians L, Catania VD, Montalva L, et al. Fetal lung underdevelopment is rescued by administration of amniotic fluid stem cell extracellular vesicles in rodents. *Science translational medicine.* 2021;13(590):eaax5941.

844 35. Barrett KM, Brott TG. Stroke caused by extracranial disease. *Circulation research.* 2017;120(3):496-501.

846 36. Goncalves I, Sun J, Tengryd C, et al. Plaque vulnerability index predicts cardiovascular events: a histological study of an endarterectomy cohort. *Journal of the American Heart Association.* 2021;10(15):e021038.

849 37. Matic LP, Jesus Iglesias M, Vesterlund M, et al. Novel multiomics profiling of human carotid atherosclerotic plaques and plasma reveals biliverdin reductase B as a marker of intraplaque hemorrhage. *JACC: Basic to Translational Science.* 2018;3(4):464-480.

852 38. Stary HC, Chandler AB, Dinsmore RE, et al. A definition of advanced types of atherosclerotic lesions and a histological classification of atherosclerosis: a report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Circulation.* 1995;92(5):1355-1374.

856 39. Palasca O, Santos A, Stolte C, Gorodkin J, Jensen LJ. TISSUES 2.0: an integrative web resource on mammalian tissue expression. *Database.* 2018;2018:bay003.

858 40. Forster DT, Li SC, Yashiroda Y, et al. BIONIC: biological network integration using convolutions. *Nature Methods.* 2022;19(10):1250-1261.

860 41. Hong P, Koza S, Bouvier ES. A review size-exclusion chromatography for the analysis of protein biotherapeutics and their aggregates. *Journal of liquid chromatography & related technologies.* 2012;35(20):2923-2950.

863 42. Petersen JD, Mekhedov E, Kaur S, Roberts DD, Zimmerberg J. Endothelial cells release microvesicles that harbour multivesicular bodies and secrete exosomes. *Journal of Extracellular Biology.* 2023;2(4):e79.

866 43. Organization WH. *Strategy for optimizing national routine health information systems: strengthening routine health information systems to deliver primary health care and universal health coverage.* World Health Organization; 2024.

869 44. Cheng HS, Besla R, Li A, et al. Paradoxical suppression of atherosclerosis in the absence of microRNA-146a. *Circulation research.* 2017;121(4):354-367.

871 45. Cheng HS, Sivachandran N, Lau A, et al. Micro RNA-146 represses endothelial activation by inhibiting pro-inflammatory pathways. *EMBO molecular medicine.* 2013;5(7):1017-1034.

874 46. Raitoharju E, Lyytikäinen L-P, Levula M, et al. miR-21, miR-210, miR-34a, and miR-146a/b are up-regulated in human atherosclerotic plaques in the Tampere Vascular Study. *Atherosclerosis.* 2011;219(1):211-217.

877 47. Nazari-Jahantigh M, Wei Y, Noels H, et al. MicroRNA-155 promotes atherosclerosis by repressing Bcl6 in macrophages. *The Journal of clinical investigation.* 2012;122(11):4190-4202.

880 48. Yang Y, Yang L, Liang X, Zhu G. MicroRNA-155 promotes atherosclerosis inflammation via targeting SOCS1. *Cellular Physiology and Biochemistry.* 2015;36(4):1371-1381.

882   49. Huang S, Zeng Z, Sun Y, et al. Association study of hsa_circ_0001946, hsa-miR-7-5p and
883   PARP1 in coronary atherosclerotic heart disease. *International journal of cardiology.*
884   2021;328:1-7.
885   50. Reddy MA, Jin W, Villeneuve L, et al. Pro-inflammatory role of microrna-200 in vascular
886   smooth muscle cells from diabetic mice. *Arteriosclerosis, thrombosis, and vascular*
887   *biology.* 2012;32(3):721-729.
888   51. Zhang F, Cheng N, Du J, Zhang H, Zhang C. MicroRNA-200b-3p promotes endothelial cell
889   apoptosis by targeting HDAC4 in atherosclerosis. *BMC cardiovascular disorders.*
890   2021;21:1-12.
891   52. An T-H, He Q-W, Xia Y-P, et al. MiR-181b antagonizes atherosclerotic plaque vulnerability
892   through modulating macrophage polarization by directly targeting Notch1. *Molecular*
893   *Neurobiology.* 2017;54:6329-6341.
894   53. Di Gregoli K, Mohamad Anuar NN, Bianco R, et al. MicroRNA-181b controls
895   atherosclerosis and aneurysms through regulation of TIMP-3 and elastin. *Circulation*
896   *research.* 2017;120(1):49-65.
897   54. Sun P, Li L, Liu Y-Z, et al. MiR-181b regulates atherosclerotic inflammation and vascular
898   endothelial function through Notch1 signaling pathway. *European Review for Medical &*
899   *Pharmacological Sciences.* 2019;23(7).
900   55. Kuleshov MV, Jones MR, Rouillard AD, et al. Enrichr: a comprehensive gene set
901   enrichment analysis web server 2016 update. *Nucleic acids research.* 2016;44(W1):W90-
902   W97.
903   56. Hallal S, Tűzesi Á, Grau GE, Buckland ME, Alexander KL. Understanding the extracellular
904   vesicle surface for clinical molecular biology. *Journal of extracellular vesicles.*
905   2022;11(10):e12260.
906   57. He R, Zhu J, Ji P, Zhao F. SEVtras delineates small extracellular vesicles at droplet
907   resolution from single-cell transcriptomes. *Nature Methods.* 2024;21(2):259-266.
908   58. Wang S, Aurora AB, Johnson BA, et al. The endothelial-specific microRNA miR-126
909   governs vascular integrity and angiogenesis. *Developmental cell.* 2008;15(2):261-271.
910   59. Fish JE, Santoro MM, Morton SU, et al. miR-126 regulates angiogenic signaling and
911   vascular integrity. *Developmental cell.* 2008;15(2):272-284.
912   60. Vasa-Nicotera M, Chen H, Tucci P, et al. miR-146a is modulated in human endothelial cell
913   with aging. *Atherosclerosis.* 2011;217(2):326-330.
914   61. McCann JV, Xiao L, Kim DJ, et al. Endothelial miR-30c suppresses tumor growth via
915   inhibition of TGF-β-induced Serpine1. *J Clin Invest.* 2019;129(4):1654-1670.
916   62. Yang ZJ, Liu R, Han XJ, et al. Knockdown of the long non-coding RNA MALAT1 ameliorates
917   TNF-α-mediated endothelial cell pyroptosis via the miR-30c-5p/Cx43 axis. *Mol Med Rep.*
918   2023;27(4).
919   63. Afzal TA, Luong LA, Chen D, et al. NCK Associated Protein 1 Modulated by mi RNA-214
920   Determines Vascular Smooth Muscle Cell Migration, Proliferation, and Neointima
921   Hyperplasia. *Journal of the American Heart Association.* 2016;5(12):e004629.
922   64. Cordes KR, Sheehy NT, White MP, et al. miR-145 and miR-143 regulate smooth muscle
923   cell fate and plasticity. *Nature.* 2009;460(7256):705-710.

65.   Cheng Y, Liu X, Yang J, et al. MicroRNA-145, a novel smooth muscle cell phenotypic
marker and modulator, controls vascular neointimal lesion formation. *Circulation*
research. 2009;105(2):158-166.

66.   O'Connell RM, Zhao JL, Rao DS. MicroRNA function in myeloid biology. *Blood, The*
Journal of the American Society of Hematology. 2011;118(11):2960-2969.

67.   Das A, Ganesh K, Khanna S, Sen CK, Roy S. Engulfment of apoptotic cells by
macrophages: a role of microRNA-21 in the resolution of wound inflammation. *The*
Journal of Immunology. 2014;192(3):1120-1129.

68.   Maurya M, Barthwal MK. MicroRNA-99a: a potential double-edged sword targeting
macrophage inflammation and metabolism. *Cellular & Molecular Immunology.*
2021;18(9):2290-2292.

69.   Rodriguez A, Vigorito E, Clare S, et al. Requirement of bic/microRNA-155 for normal
immune function. *Science.* 2007;316(5824):608-611.

70.   Seeley JJ, Baker RG, Mohamed G, et al. Induction of innate immune memory via
microRNA targeting of chromatin remodelling factors. *Nature.* 2018;559(7712):114-119.

71.   Jin S, Guerrero-Juarez CF, Zhang L, et al. Inference and analysis of cell-cell
communication using CellChat. *Nature communications.* 2021;12(1):1088.

72.   Consortium* TTS, Jones RC, Karkanias J, et al. The Tabula Sapiens: A multiple-organ,
single-cell transcriptomic atlas of humans. *Science.* 2022;376(6594):eabl4896.

73.   Hsu S-D, Lin F-M, Wu W-Y, et al. miRTarBase: a database curates experimentally validated
microRNA–target interactions. *Nucleic acids research.* 2011;39(suppl_1):D163-D169.

74.   Stary HC. Natural history and histological classification of atherosclerotic lesions: an
update. *Arteriosclerosis, thrombosis, and vascular biology.* 2000;20(5):1177-1178.

75.   Adya R, Tan BK, Punn A, Chen J, Randeva HS. Visfatin induces human endothelial VEGF
and MMP-2/9 production via MAPK and PI3K/Akt signalling pathways: novel insights into
visfatin-induced angiogenesis. *Cardiovascular research.* 2008;78(2):356-365.

76.   Bae Y-H, Bae M-K, Kim S-R, Lee JH, Wee H-J, Bae S-K. Upregulation of fibroblast growth
factor-2 by visfatin that promotes endothelial angiogenesis. *Biochemical and biophysical*
research communications. 2009;379(2):206-211.

77.   Adya R, Tan BK, Chen J, Randeva HS. Pre-B cell colony enhancing factor (PBEF)/visfatin
induces secretion of MCP-1 in human endothelial cells: role in visfatin-induced
angiogenesis. *Atherosclerosis.* 2009;205(1):113-119.

78.   Adya R, Tan BK, Chen J, Randeva HS. Nuclear factor-κB induction by visfatin in human
vascular endothelial cells: its role in MMP-2/9 production and activation. *Diabetes care.*
2008;31(4):758-760.

79.   Kim S-R, Bae Y-H, Bae S-K, et al. Visfatin enhances ICAM-1 and VCAM-1 expression
through ROS-dependent NF-κB activation in endothelial cells. *Biochimica et Biophysica*
Acta (BBA)-Molecular Cell Research. 2008;1783(5):886-895.

80.   Sinning J-M, Losch J, Walenta K, Böhm M, Nickenig G, Werner N. Circulating
CD31+/Annexin V+ microparticles correlate with cardiovascular outcomes. *European*
heart journal. 2011;32(16):2034-2041.

81.   Arteaga RB, Chirinos JA, Soriano AO, et al. Endothelial microparticles and platelet and
leukocyte activation in patients with the metabolic syndrome. *The American journal of*
cardiology. 2006;98(1):70-74.

82.   Chironi G, Simon A, Hugel B, et al. Circulating leukocyte-derived microparticles predict
subclinical atherosclerosis burden in asymptomatic subjects. *Arteriosclerosis,*
thrombosis, and vascular biology. 2006;26(12):2775-2780.
83.   Koga H, Sugiyama S, Kugiyama K, et al. Elevated levels of VE-cadherin-positive
endothelial microparticles in patients with type 2 diabetes mellitus and coronary artery
disease. *Journal of the American College of Cardiology.* 2005;45(10):1622-1630.
84.   Amabile N, Cheng S, Renard JM, et al. Association of circulating endothelial
microparticles with cardiometabolic risk factors in the Framingham Heart Study.
European Heart Journal. 2014;35(42):2972-2979.
85.   Crescitelli R, Lässer C, Lötvall J. Isolation and characterization of extracellular vesicle
subpopulations from tissues. *Nature Protocols.* 2021;16(3):1548-1580.
86.   Dixson AC, Dawson TR, Di Vizio D, Weaver AM. Context-specific regulation of
extracellular vesicle biogenesis and cargo selection. *Nature Reviews Molecular Cell*
Biology. 2023;24(7):454-476.
87.   Shankman LS, Gomez D, Cherepanova OA, et al. KLF4-dependent phenotypic modulation
of smooth muscle cells has a key role in atherosclerotic plaque pathogenesis. *Nature*
medicine. 2015;21(6):628-637.
88.   Pan H, Xue C, Auerbach BJ, et al. Single-cell genomics reveals a novel cell state during
smooth muscle cell phenotypic switching and potential therapeutic targets for
atherosclerosis in mouse and human. *Circulation.* 2020;142(21):2060-2075.
89.   Bennett MR, Sinha S, Owens GK. Vascular smooth muscle cells in atherosclerosis.
Circulation research. 2016;118(4):692-702.
90.   Fernandez DM, Rahman AH, Fernandez NF, et al. Single-cell immune landscape of human
atherosclerotic plaques. *Nature medicine.* 2019;25(10):1576-1588.
91.   Milei J, Parodi J, Barone A, et al. Carotid atherosclerosis. Immunocytochemical analysis
of the vascular and cellular composition in endarterectomies. *Cardiologia (Rome, Italy).*
1996;41(6):535-542.
92.   Soehnlein O, Libby P. Targeting inflammation in atherosclerosis—from experimental
insights to the clinic. *Nature reviews Drug discovery.* 2021;20(8):589-610.
93.   Libby P. The biology of atherosclerosis comes full circle: lessons for conquering
cardiovascular disease. *Nature Reviews Cardiology.* 2021;18(10):683-684.
94.   Gao P, Chen Z-q, Bao Y-h, Jiao L-q, Ling F. Correlation between carotid intraplaque
hemorrhage and clinical symptoms: systematic review of observational studies. *Stroke.*
2007;38(8):2382-2390.
95.   Parma L, Baganha F, Quax PH, de Vries MR. Plaque angiogenesis and intraplaque
hemorrhage in atherosclerosis. *European journal of pharmacology.* 2017;816:107-115.
96.   Wadén K, Karlöf E, Narayanan S, et al. Clinical risk scores for stroke correlate with
molecular signatures of vulnerability in symptomatic carotid patients. *Iscience.*
2022;25(5).
97.   Larson AS, Benson JC, Brinjikji W, et al. Variations in the presence of carotid Intraplaque
hemorrhage across age categories: what age groups are Most likely to benefit from
plaque imaging? *Frontiers in neurology.* 2020;11:603055.

98.   Arderiu G, Peña E, Badimon L. Angiogenic microvascular endothelial cells release
microparticles rich in tissue factor that promotes postischemic collateral vessel
formation. *Arteriosclerosis, thrombosis, and vascular biology.* 2015;35(2):348-357.
99.   Boyer MJ, Kimura Y, Akiyama T, et al. Endothelial cell-derived extracellular vesicles alter
vascular smooth muscle cell phenotype through high-mobility group box proteins.
Journal of extracellular vesicles. 2020;9(1):1781427.
100.   Charla E, Mercer J, Maffia P, Nicklin S. Extracellular vesicle signalling in atherosclerosis.
Cellular signalling. 2020;75:109751.
101.   He S, Wu C, Xiao J, Li D, Sun Z, Li M. Endothelial extracellular vesicles modulate the
macrophage phenotype: Potential implications in atherosclerosis. *Scandinavian journal*
of immunology. 2018;87(4):e12648.
102.   Jansen F, Yang X, Baumann K, et al. Endothelial microparticles reduce ICAM-1 expression
in a micro RNA-222-dependent mechanism. *Journal of cellular and molecular medicine.*
2015;19(9):2202-2214.
103.   Liang H, Li S, Chen H. GW28-e0938 Endothelial microparticles-mediated transfer of
microRNA-19b, a novel messenger in cell-cell communication, plays a key role in
endothelial migration and angiogenesis. *Journal of the American College of Cardiology.*
2017;70(16S):C34-C34.
104.   Njock M-S, Cheng HS, Dang LT, et al. Endothelial cells suppress monocyte activation
through secretion of extracellular vesicles containing antiinflammatory microRNAs.
Blood, The Journal of the American Society of Hematology. 2015;125(20):3202-3212.
105.   Botts SR, Fish JE, Howe KL. Dysfunctional vascular endothelium as a driver of
atherosclerosis: emerging insights into pathogenesis and treatment. *Frontiers in*
pharmacology. 2021;12:787541.
106.   Mokry M, Boltjes A, Slenders L, et al. Transcriptomic-based clustering of human
atherosclerotic plaques identifies subgroups with different underlying biology and
clinical presentation. *Nature cardiovascular research.* 2022;1(12):1140-1155.
107.   Seeley EH, Liu Z, Yuan S, et al. Spatially resolved metabolites in stable and unstable
human atherosclerotic plaques identified by mass spectrometry imaging.
Arteriosclerosis, thrombosis, and vascular biology. 2023;43(9):1626-1635.
1040

1041

1042

1043

1044

1045

1046

1047

 Figure Legends

Figure 1. Whole tissue proteomics identifies a potential role for EVs in modulating

plaque biology corroborated with increased enrichment of tissue resident EVs

from human atherosclerotic carotid plaques. A. Experimental overview. Carotid

endarterectomy specimens (*n*=64*)* were collected from the operating room and

dissected into plaque and marginal zones for downstream analysis. **B.** Principal

component analysis of carotid tissue proteomes between the PZ and MZ (donor

matched *n*=5; donor uncorrected in Figure S2C). **C.** Percent incidence of Gene

Ontology (GO; Biological Process and Cellular Component) terms (false discovery rate

<0.05) containing "vesicle" derived from differentially enriched proteins (q-value<0.05)

from plaque and marginal zones (left panel; *n*=5). Right panel lists top 26 (by adjusted

p-value) GO Biological Process terms where vesicle-associated terms from plaque

regions are highlighted (marginal zone in Figure S2D). **D.** Heatmap representing per-

fraction proteomics identifies prototypical EV markers enriched in fractions 7-10 (left

panel). Protein abundance was Z-score normalized by protein (Representative sample,

n=4 in Figure S5A). Western blot analysis depicting EV markers (CD63, CD81, CD9,

and Flot1) in EV-enriched fractions (7-10) isolated from human atherosclerotic plaque

tissue (right panel, *n*=1). **E.** Cryo-EM of EVs isolated from plaque and adjacent

marginal zones. Arrows indicate select EV structures as bilayered nanoparticles with

dense cores (Representative Sample, *n*=4). Scale bar = 200nm. **F.** Quantification of EV

mean diameter by nanoparticle tracking analysis (NTA; *n*=20). **G.** Nanoparticle

concentration (normalized to tissue weight, NTA dilution factor, and total EV-enriched

volume; *n*=11) binned by particle size from plaque (PZ, purple) and marginal zones (MZ,

pink). Quantification of nanoparticle concentration across all sizes by NTA (*n*=11, right

panel). Bar graphs show mean ± SEM. Statistical significance assessed by Fisher's

exact test (C) and a two-tailed paired t-test (11-20 pairs; F,G).

1075

Figure 2. Carotid plaque EV-vesiculome is zone-specific (plaque versus marginal)

with differentially expressed EV-miRNA and proteins associated with pro-

atherogenic functions. A-B. Donor corrected principal component analysis (PCA)

showing EV-miRNA **(A)** and EV-protein **(B)** profiles of EVs enriched from carotid plaque

zones (PZ, purple) versus marginal zones (MZ, pink) (miRNA *n*=20 pairs; protein *n*=23

pairs). **C.** Edge quantification of protein-protein interaction network with all differentially

enriched EV-proteins (n=661) in plaque versus marginal zones. Represents full STRING

network wherein edges indicate both functional and physical protein interactions with a

medium confidence interaction score and nodes represent proteins. **D-E.** Volcano plot of

carotid tissue derived EV-miRNA **(D)** and EV-protein **(E)** contents with positive and

negative fold change (FC) representing miRNAs and proteins enriched in plaque versus

marginal zones, respectively (FDR step up/q-value<0.05). Highlighted in red are

miRNAs and proteins known to function in inflammation. **F.** Kyoto Encyclopedia of

Genes and Genomes (KEGG) pathway analysis of all differentially expressed (FDR/q-

value <0.05) predicted EV-miRNA targets (mRNA, miRTarBase via MIENTURNET) and

EV-proteins. Venn diagram of unique and shared KEGG pathways between predicted

EV-miRNA targets and EV-proteins (top left panel). Bubble plots of KEGG pathways that

were significantly (FDR<0.05) modulated by all differentially enriched plaque EV-

miRNAs (top right panel) and EV-proteins (bottom right panel), with all pathways that

were shared by EV-miRNA and EV-proteins shown in bottom left panel. Labeled are

pathways of interest (FDR<0.05). Cancer-, and infection- associated pathways were

excluded from analysis.

1098

Figure 3. *In silico* analysis of human atherosclerotic tissue derive EV-cell-of-origin

and global communication strategies in plaque and marginal zones. A. Pictorial

representation of cell of origin using the TISSUES database with all EV-proteins. Top 11

cells by strength metric calculated as a ratio of observed and background gene count

after inclusion of cell types only (tissue annotations filtered out; further details in Figure

S8A). **B.** Box plot of global EV-protein signature enrichment among cells from

atherosclerotic plaques, informed by integrating all detected EV-proteins (global EV

signature) with single cell RNA sequencing (scRNA-seq) of carotid atherosclerotic

plaques. **C.** Stacked bar plot visualizing total number of predicted cell-cell

communication interactions in plaque (purple) and marginal (pink) zones with CellChat.

D. Heatmap depicting EV-cell signaling strength employed by all EV-proteins to both

plaque and marginal single cells. Strength of EV-derived signaling molecule and cell

type shown on right and top bar graph, respectively, and represents communication

probability. **E.** Cell type annotations derived using the Tabula Sapiens database

(vascular tissue) by inputting all significant (adjusted p-value <0.05) predicted mRNA

targets of all differentially expressed EV-miRNA in plaque and marginal zones combined

(FDR<0.05). **F.** Box plot of predicted EV-miRNA-mRNA target signature enrichment

among cells of the atherosclerotic plaque, informed by integrating differentially detected

EV-miRNA - mRNA targets with scRNA-seq of carotid atherosclerotic plaques.

1118

Figure 4. Site specific (plaque versus marginal zone) differences in EV cell of

origin, recipient cell communication strategies and directed communication with

vascular cells (ECs, SMCs, macrophages). A-B. Box plot of differential marginal **(A)**

and plaque **(B)** EV-protein signature enrichment among cells of the atherosclerotic

plaque, informed by integrating differentially enriched EV-proteins with scRNA-seq of

carotid atherosclerotic plaques. **C-D.** Pictorial representation of cell of origin using the

TISSUES database with differentially expressed EV-proteins in marginal **(C)** and plaque

(D) zones. Top 10 cells by strength metric calculated as a ratio of observed and

background gene count after filtering for tissues (further details in Figure S8B). **E-F.**

Heatmap depicting differential marginal and plaque EV to cell signaling strength

employed by differentially enriched EV proteins in marginal **(E)** and plaque **(F)** zones to

all atherosclerotic single cells in plaque and marginal regions. Strength of EV-derived

signaling molecule and cell type shown on right and top bar graph, respectively. **G-H.**

Chord diagrams representing differentially expressed EV ligands to EC **(G)** and SMC

(H) ligand-receptor pairs to single cells in marginal zones (left panels) and plaque zones

(right panels). Arrows represent EV-ligand interactions. Individual cell populations

annotated in legend.

1136

Figure 5. EV concentration and greater differential EV cargo (miRNA and protein)

discriminates between symptomatic and asymptomatic atherosclerotic plaques.

A. Schematic of experimental design. Carotid endarterectomy specimens from

symptomatic and asymptomatic patients were dissected into plaque and marginal

zones, followed by EV-enrichement, quantification, and miRNA transcriptomics and

proteomics. **B.** Hematoxylin and eosin (H&E) and Movat pentachrome stain of

asymptomatic (top panel) and symptomatic (bottom panel) of carotid plaque specimens

(Patient 1 and 2). Large, calcified plaque with calcified/necrotic center, but with no

significant intraplaque hemorrhage (top panel). Large, calcified plaque with

calcified/necrotic center with intraplaque hemorrhage (bottom panel). H&E staining with

intraplaque hemorrhage indicated by green arrowheads (bottom left panel) and Movat

pentachrome staining with intraplaque hemorrhage indicated by cyan arrowheads

(bottom right panel). Scale bar (500 µm) as shown. **C-D.** Nanoparticle tracking analysis

(NTA) of nanoparticle concentration in asymptomatic **(C)** and symptomatic **(D)**

atherosclerotic plaque versus marginal zones. **E.** Assessment of EV size and

morphology. NTA of mean diameter of nanoparticles enriched from asymptomatic (left

panel) and symptomatic (middle panel) carotid plaques. Cryogenic electron microscopy

(cryo-EM) of EVs from asymptomatic (right panel, top) and symptomatic (right panel,

bottom) carotid plaques (*n*=1-2 per group). Arrows indicate select EV structures as

bilayered nanoparticles with dense cores. Scale bar = 200nm. **F-G.** Volcano plot of

carotid tissue resident EV-miRNA transcriptome in asymptomatic **(F)** and symptomatic

(G) with positive and negative fold change (FC) representing miRNAs enriched in

plaque versus marginal zones (FDR step up<0.05). **H-I.** Volcano plot of carotid tissue

resident EV-proteome in asymptomatic **(H)** and symptomatic **(I)** with positive and

negative fold change (FC) representing EV-proteins enriched in plaque versus marginal

zones (q-value<0.05). Bar graphs show mean ± SEM. Statistical significance assessed

by non-parametric paired Wilcoxon test **(C-E)**.

1164

Figure 6. Differential plaque EV-secretome highlights key pathways involved in

plaque vulnerability. A. Venn diagram depicting shared and unique plaque derived EV-

miRNA in asymptomatic (green) and symptomatic (purple) conditions (FDR step-

up<0.05, filtered for miRNAs<1000). **B.** Total numbers of significant Gene Ontology

(GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways (adjusted p-

value<0.05) modulated by the predicted mRNA targets (miRTarBase via MIENTURNET,

threshold of minimum interactions 1; FDR<0.05) of differentially expressed plaque EV-

miRNAs in asymptomatic (green), symptomatic (purple), and shared (brown) conditions.

C. Dot plot of all KEGG pathways derived from unique asymptomatic plaque-derived

predicted EV-miRNA mRNA targets (p-value<0.05). Labelled are significant (adjusted p-

value<0.05) pathways graphed by odds ratio. **D.** Dot plot of KEGG pathway analysis

from unique symptomatic plaque-derived predicted EV-miRNA mRNA targets. Graphed

are top 42 significant (adjusted p-value<0.05) pathways by odds ratio (total 120

pathways with adjusted p-value<0.05). **E.** Venn diagram depicting shared and unique

plaque derived EV-proteins in asymptomatic (green) and symptomatic (purple)

conditions (q-value<0.05). **F.** Total numbers of significant GO and KEGG pathways

(adjusted p-value<0.05) modulated by differentially expressed plaque EV-proteins (q-

value<0.05) in asymptomatic (green), symptomatic (purple), and shared (brown)

conditions. **G.** Bar graph of all significant KEGG pathways (adjusted p-value<0.05) from

unique asymptomatic plaque-derived EV-proteins (adjusted p-value<0.05). **H.** Bar graph

of all significant KEGG pathways (adjusted p-value<0.05) from unique symptomatic

plaque-derived EV-proteins (adjusted p-value<0.05). STRING (db version 12.0) used to

generate protein-protein interaction maps (right panel) of pathways of interest. Cancer-,

and infection- associated pathways were excluded from analysis.

1189

Figure 7. Endothelial cells (ECs) are cellular players modulating cell-cell

communication through EVs in vulnerable plaques. A-D. Box plots of symptom and

regional EV-protein signature enrichment among cells of the atherosclerotic plaque,

informed by integrating differentially expressed EV-proteins in asymptomatic and

symptomatic, marginal versus plaque zones with scRNA-seq of carotid atherosclerotic

plaques. Asymptomatic marginal **(A)** and plaque **(B)** zone signature enrichment among

cells of the atherosclerotic plaque. Symptomatic marginal **(C)** and plaque **(D)** zone

signature enrichment among cells of the atherosclerotic plaque. **E-F.** Heatmap depicting

differential asymptomatic plaque- **(E)** and symptomatic plaque- **(F)** EV-cell signaling

strength employed by differentially expressed EV-proteins in asymptomatic and

symptomatic cohorts with plaque cells. Strength of EV-derived signaling molecule and

cell type shown on right and top bar graph, respectively.

1202

Figure 8. Overrepresentation analysis identifies angiogenesis and

neovascularization as functional effects of vulnerable (symptomatic) plaque EVs

with support from a functional EC sprouting assay. A. Chord diagrams representing

ligand-receptor pairs between differentially enriched EV-proteins (ligands) to ECs

(receptor) in asymptomatic (top panel) and symptomatic (bottom panel) plaques. Red

arrows represent EV-ligand to receptor interactions. Individual cell populations

annotated in legend. **B.** Filtered inquiry of EC, angiogenesis, and neovascularization

specific pathways from Gene Ontology (GO) overrepresentation analysis of differentially

expressed predicted EV-miRNA-mRNA targets in asymptomatic and symptomatic

groups. Fold enrichment represented in legend. **C.** Pairwise comparison of mean fold

enrichment of EC-specific pathways in symptomatic versus asymptomatic plaques. **D-F.**

Functional effect of symptomatic marginal zone (MZ) and plaque zone (PZ) EVs on

endothelial cell angiogenesis. Representative confocal microscopy images of HUVEC-

spheroid angiogenesis after addition of EVs ($2\text{-}4\text{X}10^{10}$ EVs; 24 h), or assay control **(D)**.

Scale bar 100 μm. Representative grayscale images with annotations for sprout number

(E, left panel) and quantification of sprout number per spheroid (n=7-10 from 3

independent experiments; **E, right panel**). Representative grayscale images with

tracings representing individual sprout lengths **(F, left panel)** and quantification (n=188-

331 sprouts from 7–10 spheroids from 3 independent experiments). Bar graphs shows

the mean±SEM. Violin plots show median ± interquartile range. Statistical significance

assessed by 1-way ANOVA with Sidak multiple comparisons test **(**adjusted p-values

shown; **E,F)**. P-values meeting significance (<0.05) shown in bolded text.

1225

1226

1227

1228

1229

1230

Supplemental Figure Legends

Supplemental Figure 1. Histopathology validates tissue dissection (plaque and marginal zones) and identifies characteristic features of late-stage atherosclerosis predominantly in plaque zones. A. Hematoxylin and eosin (H&E) and Movat pentachrome stain of plaque and marginal zones from patient one. Area adjacent to plaque (marginal zone; 1-2). H&E staining: intimal hyperplasia (green bracket), media with moderate degenerative changes (cyan bracket; 1). Movat pentachrome staining: intimal hyperplasia (blue bracket), media with moderate degenerative changes (cyan bracket; 2). Middle zone of large, calcified plaque (3-4). H&E staining: surface calcification (green octagon), fibrous cap (green star), and necrotic center (yellow octagon); media (yellow star; 3). Movat pentachrome staining: surface calcification (blue octagon), fibrous cap (blue star), necrotic center (cyan octagon), media (cyan star; 4).

Edge zone of large, calcified plaque (5-6). H&E staining: calcification (green star; 5). Movat pentachrome staining: calcification (cyan star; 6). **B.** H&E and Movat pentachrome stain of plaque and marginal zones from patient two. Area adjacent to plaque from (marginal zone; 1-2) H&E staining: intimal hyperplasia (green bracket), media with moderate degenerative changes (cyan bracket;1). Movat pentachrome staining: intimal hyperplasia (blue bracket), media with moderate degenerative changes (cyan bracket; 2). Middle zone of large, calcified plaque (3-4). H&E staining: surface calcification (green octagon), fibrous cap (green stars), and necrotic center (yellow octagon); media (yellow star; 3). Movat pentachrome staining: surface calcification (blue

octagon), fibrous cap (blue star), necrotic center (cyan octagon), media (cyan star, 4).

Edge zone of large, calcified plaque (5-6). H&E staining: media with marked disruption

of the subintimal layer by plaque (green bracket), plaque (artifactually displaced from

the rest of the tissue) with fibrous cap (cyan star) and intraplaque hemorrhage (green

star, 5). Movat pentachrome staining: media with marked disruption of the subintimal

layer by plaque (cyan bracket), plaque (artifactually displaced from the rest of the

tissue) with fibrous cap (cyan star) and intraplaque hemorrhage (blue star; 6). Scale bar

(500 μm) as shown.

1262

Supplemental Figure 2. Carotid tissue proteomics delineates differential protein

enrichment in plaque versus marginal zones with pathway analysis identifying

atherogenic signatures. A. Proteomic data pre-processing and per-sample violin plots

for carotid tissue proteomes of the plaque zone and marginal zone (n=10). Following

valid value filtering, data was median normalized, log2 transformed, and imputed from a

Gaussian distribution. **B.** Volcano plot of carotid plaque and marginal zone proteome

with purple and pink representing plaque and marginal enriched, respectively (q-

value<0.05). **C.** Donor uncorrected principal component analysis of label-free

proteomics of whole-tissue carotid plaque (purple) and marginal zones (pink; n=5). **D.**

Percent incidence of Gene Ontology (GO; Biological Process and Cellular Component)

terms (false discovery rate <0.05) containing "vesicle" resulting from all differentially

expressed proteins (q-value<0.05) from plaque and marginal zones (right panel; n=5,

repeated from Figure 1C). Left panel lists top 26 (by adjusted p-value) GO Biological

Process terms and highlighted are vesicle-associated terms from marginal zones. **E.**

Top 25 (by adjusted p-value) GO Cellular Component pathways derived from all

differentially enriched proteins in plaque zones. **F.** Top 25 (by adjusted p-value) Kyoto

Encyclopedia of Genes and Genomes pathways derived from all differentially enriched

proteins in plaque zones. Statistical significance assessed by Fisher's exact test (D)

Supplemental Figure 3. Marginal zone pathway analysis features vesicle related

and atherogenic pathways. A. Top 25 (by adjusted p-value) Gene Ontology Cellular

Component pathways derived from all differentially enriched proteins in marginal zones.

B. Top 25 (by adjusted p-value) Kyoto Encyclopedia of Genes and Genomes pathways

derived from all differentially enriched proteins in marginal zones.

Supplemental Figure 4. Carotid Plaque EV-enrichment protocol and validation of

purity. A. Representative images showing dissection of whole carotid plaque into

marginal and plaque zones. **B.** Schematic for EV isolation from atherosclerotic carotid

plaque tissue. Briefly, tissues were collected in Hanks balanced salt solution (HBSS)

and dissected into plaque and adjacent marginal zones. Subsequently, tissues were

prepared for size exclusion chromatography (SEC) by enzymatic digestion, and

centrifugation and filtration to remove cellular debris. EVs were enriched via SEC in

fractions 7-10 and concentrated using ultrafiltration to a desired volume. EVs destined

for proteomics underwent an additional purification step with ultracentrifugation and

density gradient ultracentrifugation. **C-D.** Lipoprotein co-isolation in EV preparations via

enzyme-linked immunosorbent assay (ELISA) for ApoA1 (HDL; **C**) and ApoB100 (LDL;

D; *n*=1 plaque). **E.** Fractional analysis of nanoparticle isolation via nanoparticle tracking

1300   analysis (NTA) and bicinchoninic acid assay (*n*=1). **F.** Flow Cytometry based

1301   assessment of whole cell co-isolation with Hoechst nuclei stain. Bone marrow cells

1302   (positive control) and plaque-derived EV fractions depicted on left and right panel,

1303   respectively. EV, extracellular vesicle (*n*=1).

1304

Supplemental Figure 5. Proteomics of EVs enriched from carotid tissue identifies

several EV markers. A. Prototypical EV markers are enriched in fractions 7-10.

Abundance Z-score normalized by protein in four samples. **B.** Bubble plot of GO

pathway analysis of all differentially expressed EV-proteins between plaque zone and

marginal zone filtered by vesicle-related terms. Data points are sized by combined

score, and size-scaled by genes altered in pathway/total number of unique genes in

analysis and color-scaled by biological processes and cellular locations.

1312

Supplemental Figure 6. Density gradient ultracentrifugation (DGUC) after size

exclusion chromatography (SEC) enriches for EVs and depletes extracellular

matrix proteins. N=3 matched sample pre- and post-DGUC. **A.** After DGUC, 85.4%

(958/1121) of proteins IDs remain, 158 protein IDs were no longer identified, and 4 were

uniquely identified. Markers of EVs were retained following DGUC. **B.** SEC and DGUC

cluster separately in principal component analysis (PCA) after using a generalized linear

model to account for matched sample variability. **C.** Extracellular matrix proteins were

differentially enriched in samples prior to DGUC (Total: N=206 differentially enriched in

SEC pre DGUC, N=58 differentially enriched post DGUC). p-value threshold of 0.1. **D.**

Transmission electron microscopy confirms the removal of debris indicative of

extracellular matrix and retention of EV structures. Representative image, *n*=3 samples.

Scale bars = 200 nm **E.** Post-DGUC nanoparticle tracking analysis identified particles

with a size distribution consistent with the TEM imaging in both marginal zone (MZ) and

plaque zone (PZ)**.** Graphs shows mean ± SEM. **F.** Proteomic data pre-processing and

per-sample violin plots for EVs derived from carotid tissue plaque zone and marginal

zone (*n*=46). Following valid value filtering, data was median normalized, log2

transformed, and imputed from a Gaussian distribution.

1330

Supplemental Figure 7. EV-miRNA transcriptomics data post-alignment quality

metrics. A. Alignment breakdown chart features each sample as a column. The

percentage of reads with different alignment outcomes (unique paired, non-unique,

unaligned) is quantified by the y-axis and visualized by colored stacked columns. **B.**

Coverage plot showing average read depth for each sample. **C.** Total reads plot

showing total reads for each sample. **D.** Average base quality per read plot displays the

proportion of reads with a given Phred quality score wherein all base quantities within

the read are averaged. Lines represent individual samples. **E.** Average base quality per

position plot displays the average Phred score for each position in the reads. Lines

represent individual samples. **F.** Donor uncorrected principal component analysis (PCA)

showing EV-miRNA (top panel) and EV-protein (bottom panel) profiles of EVs enriched

from carotid plaque zones (PZ, purple) versus marginal zones (MZ, pink) (miRNA *n*=20

pairs; protein *n*=23 pairs).

1344

Supplemental Figure 8. TISSUES Database identifies known cell and tissues

types driving atherosclerotic vascular disease. A-B. Dot plot of top (by strength

metric, FDR<0.05) cell type and tissue annotations derived from inputting all EV-

proteins **(A)** and all differentially enriched EV-proteins in marginal **(B,** top panel**)** and

plaque zones **(B,** bottom panel**)**. **C.** Proteins used to inform cell annotations (SMC, foam

cell, and B-lymphoblastoid cell) in the TISSUES database. Colored nodes represent

query proteins and edges represent protein-protein interactions (curated and

experimentally validated) with line thickness indicating the strength of data support.

1353

Supplemental Figure 9. Publicly available carotid scRNA-sequencing data

identifies over 17,000 single cells that can be annotated to distinct cell types

based on gene signatures. A-C. Uniform manifold approximation and projection

(UMAP) of 17,000 plaque- and marginal-region derived single cells labelled by sample

(A), region **(B)**, and cell-type **(C)**. **D-F.** Dot plot of marker gene expression for T cells

(D), macrophages and dendritic cells **(E)**, and vascular smooth muscle cells (VSMCs;

F). Dot size and color darkness indicate the percentage of gene-expressing cells, and

average expression, respectively. **G)** UMAP of markers *MS4A1*, *MZB1*, *TPSAB1*, *VWF*

for identifying B-cell lymphocytes, plasma cells, mast cells, and endothelial cells (ECs),

respectively.

1364

Supplemental Figure 10. Global atherosclerotic plaque EV communication

patterns. A. Normalized (DeSeq2 median normalization) counts of EV-miRNAs by

identity in plaque and marginal zones. **B-D:** Integration with publicly available carotid

plaque scRNA-seq dataset. **B.** Stacked bar plot visualizing enriched signaling pathways

by overall information flow from global EV-proteome to plaque (purple) and marginal

(pink) single cells. Pathways are colored by their dominance within the plaque or

marginal regions. **C.** Heatmap depicting EV-cell signaling strength employed by all EV-

proteins to atherosclerotic single cells in plaque (right panel) and marginal (left panel)

regions. **D.** Chord diagrams representing global EV ligand to EC (left), macrophage

(center), and SMC (right) ligand-receptor pairs. Arrows represent EV-receptor to ligands

interactions. Individual cell populations annotated in legend. Bar graphs show mean ±

SEM. Statistical significance assessed by two-tailed paired t-test (A) with Benjamini-

Hochberg false discovery rate correction.

1378

Supplemental Figure 11. Site specific EV communication strategies in plaque and

marginal zones. A-B. Heatmap depicting differential marginal and plaque EV-cell

signaling strength stratified by tissue site; derived from layering differentially enriched

EV proteins in marginal **(A)** and plaque **(B)** zones to atherosclerotic single cells in

plaque and marginal regions, respectively. Strength of EV-derived signaling molecule

and cell type shown on right and top bar graph, respectively. **C.** Chord diagrams

representing differentially expressed EV ligand to macrophage ligand-receptor pairs.

Arrows represent EV-ligand interactions. Individual cell populations annotated in legend.

Marginal and plaque zones represented in left and right panels, respectively.

Supplemental Figure 12. Further characterization of EV release, contents, and

pathway analysis in asymptomatic and symptomatic plaques. A. Hematoxylin and

eosin (H&E) and Movat pentachrome stain of asymptomatic (Patient 3 and 4) and

symptomatic (Patient 5) of carotid plaque specimens. Large plaque with significant

inflammation, but without a necrotic core (Patient 3). Large, calcified plaque with

calcified/necrotic center with intraplaque hemorrhage (Patient 4). H&E staining with

intraplaque hemorrhage indicated by green arrowhead and Movat pentachrome staining

with intraplaque hemorrhage indicated by cyan arrowhead. Large, calcified plaque with

calcified/necrotic center with intraplaque hemorrhage (Patient 5). H&E staining with

intraplaque hemorrhage indicated by green arrowhead and Movat pentachrome staining

with intraplaque hemorrhage indicated by cyan arrowhead. Scale bar (500 µm) as

shown. **B.** Differences plot from the Wilcoxon test comparing asymptomatic and

symptomatic plaque versus marginal EV concentrations (Figure 5C-D). **C-D.**

Nanoparticle tracking analysis (NTA) of nanoparticle concentration between

asymptomatic versus symptomatic cohorts in marginal **(C)** and plaque **(D)** zones. **E.**

Protein-protein interaction network with all differentially enriched EV-proteins in

asymptomatic and symptomatic plaque versus marginal zones. Represents full STRING

network wherein edges indicate both functional and physical protein interactions with a

medium confidence interaction score (0.400). **F.** Bar graph of all significant KEGG

pathways (adjusted p-value<0.05) derived from shared asymptomatic and symptomatic

plaque-derived EV-proteins (adjusted p-value<0.05). Cancer-, and infection- associated

pathways were excluded from analysis.

1410

1411 **Supplemental Figure 13. Recipient cell repertoire in asymptomatic versus**

1412 **symptomatic (vulnerable) plaques. A-B.** Box plot of predicted EV-miRNA-mRNA

1413 target signature enrichment among cells of the atherosclerotic plaque, informed by

1414 integrating differentially detected EV-miRNA - mRNA targets in asymptomatic **(A)** and

1415 symptomatic **(B)** plaques with scRNA-seq of carotid atherosclerotic plaques.

1416

Supplemental Figure 14. Symptom (asymptomatic versus symptomatic) and site-

specific (plaque versus marginal) differences in EV-ligand-receptor interactions.

A. Chord diagrams representing ligand-receptor pairs between differentially enriched

EV-proteins (ligands) to ECs (receptor) in symptomatic marginal zones. Red arrows

represent EV-ligand to receptor interactions. Individual cell populations annotated in

legend. **B-D.** Chord diagrams representing ligand-receptor pairs between differentially

enriched EV-proteins (ligands) to SMCs (receptor) in asymptomatic plaque zone **(B)**,

symptomatic marginal zone **(C)** and symptomatic plaque zone **(D)**. Purple arrows

represent EV-ligand to receptor interactions. Individual cell populations annotated in

legend. **E-G.** Chord diagrams representing ligand-receptor pairs between differentially

enriched EV-proteins (ligands) to macrophages (receptor) in asymptomatic plaque zone

(E), symptomatic marginal zone **(F)** and symptomatic plaque zone **(G)**. Yellow arrows

represent EV-ligand to receptor interactions. Individual cell populations annotated in

legend.

1431

Supplemental Figure 15. Approach to externally validate cell communication

predictions in dataset from the Biobank of Karolinska Endarterectomy (BiKE). A.

CD63 protein expression in symptomatic (S) vs asymptomatic (AS) patients in both

central (plaque zone) and distal (marginal zone) regions. B. NAMPT (visfatin) protein

normalized to CD63 level by zone in S versus AS patients in both central (plaque zone)

and distal (marginal zone) regions. C. VEGFA protein normalized to CD63 level by zone

in S versus AS patients in both central (plaque zone) and distal (marginal zone) regions.

Protein levels are expressed in arbitrary units (AU). CD63 zone values were determined

based on CD63 protein levels in each of the central AS (0.86 AU set to 1.1), distal AS

(0.79 AU set to 1), central S (1.20 AU set to 1.5) and distal S (0.96 AU set to 1.2).

NAMPT and VEGFA protein levels were then multiplied by the corresponding CD63

zone values for each group to account for CD63 expression in these zones. Statistical

comparison was performed using a two-tailed t-test when comparing AS vs S and a

paired t-test when comparing distal vs central zones as they were collected from the

same patient. P-value threshold for significance was set to 0.05 and results are

presented as mean ± standard deviation (SD). There were n=9 patients in the AS group

and n=9 patients in the S group.

1449

Supplemental Figure 16. Bioinformatic and network analysis to delineate distinct

and shared functions of EV-miRNAs and EV-proteins. A. Bubble plots depicting top

Gene Ontology pathways (by fold enrichment) for EV-miRNA (left panel), EV-proteins

(middle panel), and network analysis (BIONIC network integration algorithm, 512

embedded features, Louvain-clustered modules) to integrate whole EV-secretome

(miRNA and proteins; right panel). **B.** Bubble plots depicting top Kyoto Encyclopedia of

Genes and Genome pathways (by fold enrichment) for EV-miRNA (left panel), EV-

proteins (middle panel), and network analysis (BIONIC network integration algorithm,

512 embedded features, Louvain-clustered modules) to integrate whole EV-vesiculome

(miRNA and proteins; right panel). Bubbles colored by q-value and sized by fold

enrichment.

1461

Supplemental Figure 17. Overrepresentation analysis delineates distinct and

shared pathways being modulated by plaque EV-miRNA and EV-proteome with

symptom status-based distinctions in pathway identity and strength. A-B.

Overrepresentation analysis with Gene Ontology (GO) pathways using differentially

expressed predicted EV-miRNA-mRNA targets **(A)** and EV-proteins **(B)** in asymptomatic

and symptomatic groups. **C-D.** Overrepresentation analysis with Kyoto Encyclopedia of

Genes and Genomes pathways using differentially expressed predicted EV-miRNA-

mRNA targets **(C)** and EV-proteins **(D)** in asymptomatic and symptomatic groups.

1470

Supplemental Figure 18. Overrepresentation analysis finds endothelial and

angiogenesis-based pathways as contributors to plaque vulnerability. A-C. Subset

analysis of differentially expressed EV-miRNAs **(A,B)** and EV-proteins **(C)** querying

specific endothelial and angiogenesis based KEGG Pathways. Pairwise comparison of

mean fold enrichment of EC-specific pathways in symptomatic versus asymptomatic

plaques **(B)**. **D.** Subset analysis of differentially expressed EV-proteins querying specific

endothelial and angiogenesis-based Gene Ontology Biological Process (GO-BP)

Pathways. **E-F.** Subset analysis of differentially expressed EV-vesiculome (miRNA and

proteins; Per-module) querying specific endothelial and angiogenesis-based Gene

Ontology Biological Process (GO-BP; **E**) and Kyoto Encyclopedia of Genes and

Genomes pathways (KEGG; **F**).

1482

Supplemental Figure 19. Functional validation of increased angiogenesis with

plaque-derived vesicles employing a newly optimized low-input sprouting assay.

A-B. Pairwise comparison of mean fold enrichment of endothelial cell (EC)-specific

pathways in EV-protein based Gene Ontology (GO; **A**) and Kyoto Encyclopedia of

Genes and Genomes (KEGG**; B**) pathways in symptomatic versus asymptomatic

plaques. Individual tests labelled above graph. **C-D.** Pairwise comparison of mean fold

enrichment of EC-specific pathways in EV-vesiculome (miRNA and protein) based Gene

Ontology (GO; **C**) and Kyoto Encyclopedia of Genes and Genomes (KEGG**; D**)

pathways in symptomatic versus asymptomatic plaques. Individual tests and p-values

labelled above graph. **E.** Schematic of sprouting angiogenesis assay depicting EV-

enrichment from plaque and marginal zones of symptomatic patients, treatment of

immortalized human umbilical endothelial cell (HUVEC) derived spheroids, followed by

staining and quantification. **F.** Representative confocal microscopy images of HUVEC-

spheroid angiogenesis after addition of EVs (2-4X10^{10} EVs; 24 h), vascular endothelial

growth factor (VEGF; positive control), plaque zone (PZ) elute (PZ EV control), marginal

zone (MZ) elute (MZ EV control), assay control (media only). Scale bar represents 100

µm. **G.** Quantification of average sprout length (n=188-331 sprouts from 7–10 spheroids

from 3 independent experiments). **H-J.** Quantification of sprout number **(H)**, sprout

length **(I)**, and average sprout length **(J)** with addition of VEGF (50 ng/ml, positive

control; *n*=6-8). Bar graphs show the mean±SEM. Violin plots show median ±

interquartile range. Statistical significance assessed by 1-way ANOVA with Sidak

multiple comparisons test **(**adjusted p-values shown, G). P-values meeting significance

(<0.05) shown in bolded text **(G)**.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A

EV
EC_1
EC_2
EC_3

ITGA10_ITGB1
F2R
CTSG
COL1A1

ITGA5_ITGB1
ITGA5_ITGB1
ITGA5_ITGB1
P4HB
ITGA10_ITGB1
COL1A1
LGALS9
NAMPT

B EV-miRNA-mRNA GO BPs

Fold Enrichment

vasculogenesis involved in coronary vascular morphogenesis
vascular endothelial growth factor production
coronary artery morphogenesis
blood vessel endothelial cell differentiation
regulation of endothelial cell development
regulation of establishment of endothelial barrier
endothelial cell activation
kidney vasculature morphogenesis
renal system vasculature morphogenesis
coronary vasculature morphogenesis
vascular endothelial growth factor signaling pathway
regulation of endothelial cell differentiation
endothelial tube morphogenesis
kidney vasculature development
morphogenesis of an endothelium
renal system vasculature development
cellular response to vascular endothelial growth factor stimulus
regulation of cell migration involved in sprouting angiogenesis
regulation of blood vessel endothelial cell migration
glomerulus vasculature development
regulation of vascular endothelial growth factor production
regulation of vascular endothelial cell proliferation
cell migration involved in sprouting angiogenesis
placenta blood vessel development
vasculogenesis
blood vessel endothelial cell migration
branching involved in blood vessel morphogenesis
regulation of morphogenesis of a branching structure
regulation of endothelial cell migration
regulation of endothelial cell apoptotic process

Asymptomatic   Symptomatic

C

$p=1.84 \times 10^{-23}$

Mean Fold Enrichment

Asymptomatic   Symptomatic

D

Media Control | Marginal Zone EV Treatment | Plaque Zone EV Treatment
Phalloidin | Phalloidin | Phalloidin

E Sprout Number

Marginal Zone
Plaque Zone

$p=1.2 \times 10^{-5}$
$p=6.0 \times 10^{-2}$
$p=1.0 \times 10^{0}$
$p=1.1 \times 10^{-5}$
$p=3.1 \times 10^{-1}$
$p=2.0 \times 10^{-4}$
$p=2.7 \times 10^{-2}$

Sprouts Number/Spheroid

Untreated   MZ Eluate   PZ Eluate   MZ-EVs   PZ-EVs

F Sprout Length

Marginal Zone
Plaque Zone

$p<1.0 \times 10^{-10}$
$p=4.8 \times 10^{-8}$
$p<1.0 \times 10^{-10}$
$p=3.6 \times 10^{-1}$
$p<1.0 \times 10^{-10}$
$p=5.6 \times 10^{-1}$
$p<1.0 \times 10^{-10}$
$p<1.0 \times 10^{-10}$

Sprouts Length (µm)

Untreated   MZ Eluate   PZ Eluate   MZ EVS   PZ EVs

Supplemental Figure 1

A Patient 1

B Patient 2

Supplemental Figure 2

A

Carotid Tissue Proteome

Consensus proteome
2616 IDs
↓
≥ 2 unique peptides
2084 IDs
↓
Uniprot ID mapping
2018 IDs
↓
≥ 70% valid values
1742 IDs

B

Marginal enriched | Plaque enriched

-log$_{10}$(qvalue) vs log$_2$ Fold Change

CCN3, SMTN, IRAG1, ITIH5, SPARCL1, LAMB2, CFL2, ACAN, MECP2, FMOD

GLUL, PON1, ITGAX, LIPA, APOM, TPSB2, ME2, FABP5, HPR, TYMP

C

(Donor uncorrected)

PC2 (12%) vs PC1 (48%)

● MZ
● PZ

D

Top 26 (by adj p-value) GO Biological Process Terms

Actomyosin Structure Organization
Platelet Aggregation
Endocytosis
Cell-Matrix Adhesion
Homotypic Cell-Cell Adhesion
Supramolecular Fiber Organization
Muscle Contraction
Muscle Organ Development
Regulation Of Cell Migration
Integrin-Mediated Signaling Pathway
Elastic Fiber Assembly
Substrate Adhesion-Dependent Cell Spreading
Contractile Actin Filament Bundle Assembly
Stress Fiber Assembly
Regulation Of Focal Adhesion Assembly
Positive Regulation Of Microtubule Nucleation
Positive Regulation Of Wound Healing
Maintenance Of Blood-Brain Barrier
Nuclear Envelope Organization
Myofibril Assembly
Regulation Of Cell-Matrix Adhesion
Negative Regulation Of Cell Migration
Regulation Of Microtubule Nucleation
Positive Regulation Of Actin Filament Bundle Assembly
Cell-Cell Junction Assembly
Actin Filament Organization

-log(adj p-value)

■ Vesicle GO Biological Process Pathways
■ Other GO Biological Process Pathways

4.3X10^{-3}

3.1x

%incidence of GO terms (FDR<0.05) containing "vesicle"

MZ | PZ

E

Plaque GO CC

Top 25 (by adj p-value) GO Cellular Component Terms

Intracellular Organelle Lumen
Secretory Granule Lumen
Ficolin-1-Rich Granule
Ficolin-1-Rich Granule Lumen
Cytoplasmic Vesicle Lumen
Tertiary Granule
Tertiary Granule Lumen
Vacuolar Lumen
Endoplasmic Reticulum Lumen
Spherical High-Density Lipoprotein Particle
Platelet Alpha Granule Lumen
Mitochondrial Respiratory Chain Complex IV
Respiratory Chain Complex IV
Lysosome
Collagen-Containing Extracellular Matrix
Lysosomal Lumen
Platelet Alpha Granule
Azurophil Granule Lumen
Azurophil Granule
Secretory Granule Membrane
Endoplasmic reticulum-Golgi Intermediate Compartment Membrane
Mitochondrial Matrix
High-Density Lipoprotein Particle
Specific Granule Lumen
Phagocytic Vesicle

-log(adj p-value)

■ Vesicle GO Cellular Component Pathways
■ Other GO Cellular Component Pathways

F

Plaque KEGG

Top 25 (by adj p-value) KEGG Terms

Complement and coagulation cascades
Phagosome
Huntington disease
Lysosome
Parkinson disease
Proteasome
Citrate cycle (TCA cycle)
Prion disease
Amyotrophic lateral sclerosis
Diabetic cardiomyopathy
Alzheimer disease
Oxidative phosphorylation
Antigen processing and presentation
Neutrophil extracellular trap formation
Pathways of neurodegeneration
Protein processing in endoplasmic reticulum
Spinocerebellar ataxia
Cholesterol metabolism
Glyoxylate and dicarboxylate metabolism
Cardiac muscle contraction
Glutathione metabolism
Platelet activation
Rheumatoid arthritis
Systemic lupus erythematosus
PPAR signaling pathway

-log(adj p-value)

Supplemental Figure 3

A Marginal GO CC

B Marginal KEGG

Supplemental Figure 4

Supplemental Figure 5

Supplemental Figure 6

A

SEC + DGUC

SEC

85.4% Shared IDs
(958/1121)

EV MARKERS

ANXA1	FLOT1
ANXA2	FLOT2
ANXA3	HSP90AA1
ANXA4	HSP90AB1
ANXA5	HSPA1B
ANXA6	HSPA2
ANXA6	MFGE8
ANXA7	PECAM1
CD81	TSG101
CD9	TSPAN9
ANXA11	

B

● SEC
● SEC + DGUC

PC2 (29%)

PC1 (48%)

958 IDs

C

SEC → ← SEC + DGUC

-log₁₀(p-value)

FBN1
COL18A1
COL15A1
COL4A2
COL8A2
COL8A1
LAMB1

log₂(Fold Change)

D

SEC   SEC + DGUC

200 nm   200 nm

E

Post-DGUC

— MZ
— PZ

% of total particles

Size (nm)

F Carotid Tissue EV Proteome

Consensus proteome
2384 IDs

↓

≥ 2 unique peptides
1939 IDs

↓

Uniprot ID mapping
1880 IDs

↓

≥ 70% valid values
1546 IDs

log₂ Normalized Abundance

Sample

log₂ Normalized Abundance

Sample

Supplemental Figure 7

Supplemental Figure 8

Supplemental Figure 9

Supplemental Figure 10

Supplemental Figure 11

Supplemental Figure 12

A

DE Asymptomatic EV-miRNA-mRNA Recipient Cell Repertoire

B

DE Symptomatic EV-miRNA-mRNA Recipient Cell Repertoire

Supplemental Figure 14

A EV-EC Symptomatic Marginal Zone

B EV-VSMC Asymptomatic Plaque Zone

C EV-VSMC Symptomatic Marginal Zone

D EV-VSMC Symptomatic Plaque Zone

E EV-Macrophage Asymptomatic Plaque Zone

F EV-Macrophage Symptomatic Marginal Zone

G EV-Macrophage Symptomatic Plaque Zone

Supplemental Figure 15

EV marker CD63 in atherosclerotic plaques

Protein values normalized to CD63 levels

Gene Ontology Pathway Analysis

EV-miRNA

EV-proteins

EV-vesiculome

Kyoto Encyclopedia of Genes and Genomes Pathway Analysis

EV-miRNA

EV-proteins

EV-vesiculome

Supplemental Figure 17

Gene Ontology Overrepresentation Analysis

A Overrepresentation Analysis of de-miRNAs against GO BP

B Overrepresentation Analysis of de-proteins against GO BP

Kyoto Encyclopedia of Genes and Genomes Overrepresentation Analysis

C Overrepresentation Analysis of de-miRNAs against KEGG Pathways

D Overrepresentation Analysis of de-proteins against KEGG Pathways

Supplemental Figure 18

Supplemental Figure 19

A Tukey DEP GO EC
p=6.06X10⁻⁰²

B Tukey DEP KEGG EC
p=6.65X10⁻⁰¹

C Tukey PerMod GO EC
p=8.2X10⁻⁰¹

D Tukey PerMod KEGG EC

E Marginal Zone (MZ) — Plaque Zone (PZ) — Symptomatic Carotid Endarterectomy Sample — EV enrichment via SEC — MZ EVs — PZ EVs — HUVEC spheroid angiogenesis assay — Immunofluorescence to quantify sprout number and length

F Media Control — MZ Elute — PZ Elute — VEGF — MZ EV — PZ EV

G Average Sprout Length
p=1.3 X10⁻⁷
p=3.3 X10⁻³
p=9.2 X10⁻¹
p=9.0 X10⁻⁶
p=1.0 X10⁰
p=1.0 X10⁻²
p=6.2 X10⁻⁴
Untreated, MZ Eluate, PZ Eluate, MZ EVs, PZ EVs

H Sprout Number VEGF
p=6.6 X10⁻⁴
Untreated, VEGF

I Sprout Length VEGF
p<1.0 X10⁻¹⁵
Untreated, VEGF

J AVERAGE Sprout length VEGF
p=1.4 X10⁻⁵
Untreated, VEGF