Wnt signaling dosage controlled by a *Ctnnb1* enhancer balances homeostasis and tumorigenesis of intestinal epithelia

Xiaojiao Hua[1,2], Chen Zhao[1,2], Jianbo Tian[4,5], Junbao Wang[1,2], Xiaoping Miao[4,5], Gen Zheng[6], Min Wu[2,3,5], Mei Ye[7], Ying Liu[1,2,*], Yan Zhou[1,2,*]

[1]Department of Neurosurgery, Medical Research Institute, Zhongnan Hospital of Wuhan University, Wuhan University, Wuhan, China

[2]Frontier Science Center of Immunology and Metabolism, Wuhan University, Wuhan, China

[3]College of Life Sciences, Wuhan University, Wuhan, China

[4]Department of Epidemiology and Biostatistics, School of Public Health, Wuhan University, Wuhan, China.

[5]TaiKang Center for Life and Medical Sciences, Wuhan University, Wuhan, China

[6]Department of Gastroenterology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

[7]Department of Gastroenterology, Zhongnan Hospital of Wuhan University, Wuhan, China

*Correspondence to Dr. Ying Liu, y.liu@whu.edu.cn; Dr. Yan Zhou, yan.zhou@whu.edu.cn

Abstract

The β-catenin-dependent canonical Wnt signaling is pivotal in organ development, tissue homeostasis, and cancer. Here we identified an upstream enhancer of *Ctnnb1*, named ieCtnnb1, that is crucial for intestinal homeostasis. ieCtnnb1 is predominantly active in the base of small intestinal crypts and throughout the epithelia of large intestine. Knockout of ieCtnnb1 led to a reduction in *Ctnnb1* transcription, compromising the canonical Wnt signaling in

intestinal crypts. Single-cell sequencing revealed that ieCtnnb1 knockout altered epithelial compositions and potentially compromised functions of small intestinal crypts. While deletion of ieCtnnb1 hampered epithelial turnovers in physiologic conditions, it prevented occurrence and progression of Wnt/β-catenin-driven colorectal cancers. Human ieCTNNB1 drove reporter gene expression in a pattern highly similar to mouse ieCtnnb1. ieCTNNB1 contains a single-nucleotide polymorphism associated with *CTNNB1* expression levels in human gastrointestinal epithelia. The enhancer activity of ieCTNNB1 in colorectal cancer tissues was stronger than that in adjacent normal tissues. HNF4α and phosphorylated CREB1 were identified as key trans-factors binding to ieCTNNB1 and regulating *CTNNB1* transcription. Together, these findings unveil an enhancer-dependent mechanism controlling the dosage of Wnt signaling and homeostasis in intestinal epithelia.

Introduction

The gastrointestinal (GI) epithelium, a high turnover organ, contains multiple cell types to fulfill complex functions including food digestion, nutrient absorption, pathogen insulation and clearance, as well as endocrine roles (**Chin et al., 2017; Kurokawa et al., 2020; Noah et al., 2011**). Enterocytes (ECs) constitute ~90% of all differentiated cells and function as highly efficient cells for absorbing nutrients and water (**Snoeck et al., 2005**). Goblet cells (GCs) produce mucus to lubricate the mucosal surface (**Gustafsson & Johansson, 2022**). Enteroendocrine cells (EECs) sense luminal nutrients and secrete peptide hormones to regulate appetite, intestinal motility, and insulin release (**Gribble & Reimann, 2019**). Tuft cells (TCs) are chemosensory cells that respond to various stimuli including infections and hypoxia (**Schneider et al., 2019**). The crypts of GI epithelium harbor various types of stem cells that continually replenish lost epithelial cells (**Ramadan et al., 2022; van der Flier & Clevers, 2009**). While the prevailing belief is that *Lgr5*-expressing crypt base

columnar (CBC) cells are actively dividing stem cells and *Bmi*-expressing "+4" cells (positioned four cells above the base of the crypt) are slowly cycling stem cells, it's important to note that crypt cells have the capacity for fate reprogramming and de-differentiation to sustain regenerative capability (**Barker et al., 2007; Beumer & Clevers, 2016; Metcalfe et al., 2014; Tian et al., 2011; Yan et al., 2012**). Intestinal stem cells generate transit-amplifying cells (TACs), which undergo multiple cell divisions and differentiate into all intestinal lineages while migrating towards the crypt orifice (**Krausova & Korinek, 2014**). In addition to stem and progenitor cells, the base of small intestinal crypts contains Paneth cells (PCs) that secret antimicrobial peptides, called defensins, thereby contributing to host defense against microbes (**Barreto et al., 2022; Lueschow & McElroy, 2020**). PCs also contribute to the stem cell niche function, *e.g.* secreting Wnt ligands, to maintain epithelial homeostasis of the small intestine (**Sato et al., 2011**).

The behavior of crypt cells is precisely regulated by multiple signals, which can act sequentially, additively, synergistically, or antagonistically (**Nakamura et al., 2007**). Among them, the Wnt/β-catenin signal pathway represents the principal force governing intestinal epithelium homeostasis, particularly in the preservation of stem cell proliferation and multipotency (**Clevers et al., 2014; Mah et al., 2016; Pinto et al., 2003**). Upon Wnt ligand binding, the β-catenin degradation complex is inactivated, leading to accumulation of β-catenin and its translocation into the nucleus, where it works with the co-activator TCF-4 to turn on the transcription of a plethora of target genes, including *c-Myc*, *CCND1*, *AXIN2*, and *CD44* (**Clevers, 2006; Clevers & Nusse, 2012; Nusse & Clevers, 2017**). Inhibition of Wnt ligand secretion reduced number of functional stem cells and led to a faster crypt fixation rate (**Huels et al., 2018**). In the process of aging, reduced Wnt signaling in stem cells and the niche of intestinal crypts causes inhibition of proliferation and reduced numbers of intestinal stem cells (**Nalapareddy et al., 2017; Pentinmikko et al., 2019**). The Wnt signal is also

essential for commitment, maturation, and location of PCs and other secretory

cells (**Andreu et al., 2008; van Es et al., 2005**). On the other hand, Wnt, EGF

and Delta-like 1/4 ligands presented by PCs sustained CBCs' proliferative

states and the production (**Sato et al., 2011**). Although the Wnt signal dosage,

largely *via* post-translational regulations, controls early lineage specification,

hematopoiesis, and tumorigenesis (**Bakker et al., 2013; Luis et al., 2011**), it

remains elusive how Wnt signaling dosage, particularly through the

transcriptional control of *Ctnnb1*, regulates composition and function of

intestinal epithelia.

Aberrant activation of canonical Wnt signaling is associated with GI cancers,

most notably colorectal cancer (CRC) (**Zhao et al., 2022; Zhu & Li, 2023**).

Germinal mutations of the *APC* gene, which encodes the negative regulator of

β-catenin, underlie the Familial Adenomatous Polyposis (FAP), a hereditary

neoplastic syndrome and the pre-cancerous condition for CRC (**Cheng et al.,**

2019; J. Li et al., 2021). Moreover, *APC* is mutated in up to 80% of human

sporadic CRCs (**J. Li et al., 2021**). Inactivation of APC causes enhanced

accumulation of β-catenin in nuclei and subsequent activation of proliferative

and pro-survival signals in CRC cells (**Cheng et al., 2019**). However, largely

due to lack of specificity, no molecular therapeutic strategy targeting the

pathway has been incorporated into CRC treatment. Interestingly, the

transcriptional activity of *CTNNB1*, the coding gene for β-catenin, is also

enhanced in CRCs (**Terrin et al., 2017**); however, the underlying mechanism

is largely unknown. Dissecting it would facilitate the development of new Wnt/β-

catenin targeting means to treat CRC.

Enhancers are essential *cis*-regulatory DNA elements that control

spatiotemporal gene expressions during cell fate specification and

maintenance (**Pachano et al., 2022**). Active enhancers are often associated

with cell fate specifiers, mostly transcription factors (TFs), and are enriched with

specific histone modification such as H3K27ac (acetylated lysine 27 of histone

H3) and H3K4me1 (mono-methylated lysine 4) of histone H3 (**Jindal & Farley, 2021**). Importantly, variations of enhancer sequences often lead to aberrant gene silence and activation, thereby causing developmental anomalies and cancers (**Kvon et al., 2021; Sengupta & George, 2017; Zabidi & Stark, 2016**). Although numerous putative enhancers have been identified by high-throughput studies, only a fraction of them were functionally annotated.

The proper expression of essential fate genes in different cell types is often controlled by specific enhancers(**Kvon et al., 2021; Pachano et al., 2022**). We previously identified a neocortical enhancer of *Ctnnb1*, the coding gene of β-catenin. The enhancer – neCtnnb1, is specifically active in the subventricular zone of developing neocortex and regulates the abundance of intermediate progenitors and production of superficial-layer pyramidal neurons (**Wang et al., 2022**). Here we have identified an intestinal enhancer for *Ctnnb1* and name it as ieCtnnb1. Reporter mice showed that ieCtnnb1 is predominantly active in the base of small intestinal crypts and across the epithelium of large intestine. Knockout of ieCtnnb1 slows down turnovers of intestinal epithelia at physiology and cancer conditions in a Wnt/β-catenin dependent manner. The human ieCTNNB1 contains a SNP that is associated with *CTNNB1* expression levels in human gastrointestinal epithelia. The enhancer activity of ieCTNNB1 in CRC tissues is higher than that in adjacent normal tissues and positively correlates with *CTNNB1* expression levels. HNF4α and phosphorylated CREB1 at serine 133 (p-S133-CREB1) were found to associate with ieCtnnb1 and regulate *Ctnnb1* transcription.

Results

ieCtnnb1 is a gastrointestinal enhancer of *Ctnnb1*

We previously identified an evolutionarily conserved upstream enhancer for *Ctnnb1* – neCtnnb1, which regulates neocortical neurogenesis (**Figure 1 - figure supplement 1A**). We reasoned that there might be other potential

enhancer regions that control *Ctnnb1* expression in a cell-specific manner. Therefore, we scanned upstream regions of *Ctnnb1* for active enhancer properties – open chromatin with enriched H3K27ac and H3K4me1 – in different tissues using data retrieved from the ENCODE database. Interestingly, we noticed that a region that is 29,068 base pair (bp) upstream of the transcription starting site (TSS) for *Ctnnb1* displayed typical enhancer feature in intestinal tissues and resides in the same topologically associating domain (TAD) with *Ctnnb1* (**Figure 1A, Figure 1 - figure supplement 2A**). The region, named as ieCtnnb1 (intestinal enhancer of *Ctnnb1*), is at open chromatin status (DNase I hypersensitive site) and enriched with H3K27ac and H3K4me1, two histone marks for active enhancers, in 8-week-old mouse small intestine tissue (**Figure 1A**). Importantly, the enrichment of H3K27ac at ieCtnnb1 is also prominent in embryonic intestine and stomach tissue, as well as in adult kidney and liver (**Figure 1 - figure supplement 1B and D**). In contrast, the enrichment was not detected in non-gastrointestinal tissues such as embryonic and adult brain (**Figure 1 - figure supplement 1D**). In line with signatures of active enhancers, the enrichment of H3K4me3 and H3K36me3 at ieCtnnb1 is negligible in all examined tissues (**Figure 1A, Figure 1 - figure supplement 1B and C**).

We next performed *in vitro* and *in vivo* analyses to validate ieCtnnb1's enhancer activity. First, the 2,153 bp core region (CR), representing the H3K27ac peak region of ieCtnnb1, could drive the expression of luciferase reporter in multiple cell types (**Figure 1 - figure supplement 2B**). Second, the ieCtnnb1 reporter mice (*H11$^{i.enh}$*) were constructed by inserting the reporter cassette – 5'-*Shh*'s promoter-ieCtnnb1-CR-*LacZ-iCre*-3' – into the *H11* locus (**Figure. 1B**). Of note, the *Shh* promoter alone at the *H11* locus did not drive the expression of reporter gene in all examined tissue (**Kvon et al., 2020**). Wholemount β-galactosidase (β-gal) staining of 8-week-old tissues revealed prominent ieCtnnb1-*LacZ* signals along the gastrointestinal (GI) tract. The

signal exhibited a proximal-low to distal-high gradient, with the ileum, cecum, and colon showing the strongest signals (**Figure. 1B, Figure 1 - figure supplement 2C**). The stomach and proximal duodenum also displayed reporter activity driven by ieCtnnb1 (**Figure. 1B, Figure 1 - figure supplement 2C**). β-gal staining of intestinal sections revealed that the ieCtnnb1-*LacZ* signal could be visualized at the bottom of small intestinal crypts and in the entire large intestinal epithelia, but is absent in sections of liver, kidney, and spleen (**Figure 1 - figure supplement 2D and data not shown**). Interestingly, the proximal-low to distal-high gradient of ieCtnnb1-*LacZ* signal along the small intestine corresponds to the expression pattern of β-catenin (**Figure 1 - figure supplement 2E**).

The transcription activity of ieCtnnb1 is strongest at the bottom of crypts, where *Lgr5*-expressing CBC cells and Paneth cells (PCs) reside. To more precisely elucidate cell types in which ieCtnnb1 is active, we crossed *H11^{i.enh}* mice with *Lgr5*-EGFP-IRES-CreERT2 (*Lgr5*-EGFP) mice to obtain 8-week-old *H11^{i.enh}*;*Lgr5-EGFP* mice. Small intestinal crypts were respectively harvested from *Lgr5-EGFP* and *H11^{i.enh}*;*Lgr5-EGFP* mice followed by enrichment of both eGFP+ and eGFP- cells (**Figure 1 - figure supplement 2F**). Quantitative RT-PCR (RT-qPCR) analyses indicated that *LacZ* transcripts could be detected in both eGFP+ and eGFP- crypt cells in *H11^{i.enh}*;*Lgr5-EGFP* mice (**Figure 1 - figure supplement 2F**). Next, sections of *H11^{i.enh}* crypts were subjected with β-gal staining followed by immunohistochemistry of OLFM4 (a CBC marker) or lysozyme (a PC marker). Data showed X-gal signals are in close proximity with both OLFM4 and lysozyme (**Figure. 1C**), suggesting that the activity of ieCtnnb1 is present in both CBCs and PCs.

We further examined the ieCtnnb1 activity in GI tracts of postnatal day 7 (P7) pups, when GI epithelia are rapidly expanding. The ieCtnnb1-*LacZ* signals could be visualized in epithelia of P7 small and large intestines, aligning well with the distribution of canonical Wnt signaling illustrated by the BAT-Gal mice

(**Figure 1 - figure supplement 2G and H**). Together, ieCtnnb1 drives transcription in embryonic intestinal epithelia, the base of small intestinal crypts and large intestinal epithelia of adult mice.

ieCtnnb1 maintains the expression of *Ctnnb1* in intestinal epithelia

We next examined whether ieCtnnb1 controls the transcription of *Ctnnb1* in intestinal epithelia. For this purpose, we employed CRISPR/Cas9-mediated gene editing to delete the genomic region containing ieCtnnb1 (**Figure 1 - figure supplement 2I**). Homozygous ieCtnnb1 knockout (*Ctnnb1$^{\Delta i.enh}$*) mice were born in Mendelian ratios and thrived through adulthood. Interestingly, the body weight of 8-week-old male *Ctnnb1$^{\Delta i.enh}$* mice was slightly but significantly lighter than that of wildtype controls (**Figure. 1D and F**), but the length of small and large intestines of *Ctnnb1$^{\Delta i.enh}$* mice was comparable to that of control mice (**Figure. 1E, G and H**). We then collected crypts from small and large intestine, finding that the expression levels of *Ctnnb1* were decreased by 26.5% and 22.3% respectively in *Ctnnb1$^{\Delta i.enh}$* crypts (**Figure. 1I and J**). In contrast, the expression level of *Ctnnb1* in the *Ctnnb1$^{\Delta i.enh}$* liver remained unaltered (**Figure 1 - figure supplement 2J**). In addition, the expression of genes located in the same TAD as *Ctnnb1* and ieCtnnb1, including *Ulk4*, *Rpl14* and *Entpd3*, was not changed in *Ctnnb1$^{\Delta i.enh}$* small intestinal crypts (**Figure 1 - figure supplement 2K**). These data suggest that ieCtnnb1 specifically maintains *Ctnnb1*'s transcription in intestinal epithelia.

In epithelial tissues, most β-Catenin is localized at cell adhesion sites to maintain barrier integrity, while a smaller fraction translocates into the nucleus to active transcription upon Wnt ligand binding. We therefore carried out nuclear/cytosol fractionation assays, which revealed a more substantial decrease in the level of nuclear β-Catenin compared to cytosolic β-Catenin (49.5% versus 29.8%) in small intestinal crypts of *Ctnnb1$^{\Delta i.enh}$* mice (**Figure. 1K and L**). RNA-seq transcriptome and RT-qPCR analyses revealed a significant

reduction in the expression levels of components in the Wnt signaling pathway,

as well as Wnt target genes, in $Ctnnb1^{\Delta i.enh}$ crypts (**Figure. 1M and N**).

Together, the loss of ieCtnnb1 undermined the Wnt signaling in intestinal

epithelia by reducing *Ctnnb1* transcription.

Loss of ieCtnnb1 altered cellular composition and expression profiles of

small intestinal crypts

The canonical Wnt/β-catenin signaling is essential for the construction and

homeostasis of intestinal epithelia (**Mah et al., 2016**). Because ieCtnnb1

maintains the expression level of *Ctnnb1* of intestinal crypts, we investigated

how the deletion of ieCtnnb1 affects cellular components and functions of

intestinal crypts. We performed single-cell sequencing on EpCAM+ epithelial

cells from small intestinal crypts of 10-week-old $Ctnnb1^{\Delta i.enh}$ and control

($H11^{i.enh}$) mice (n = 2 female mice for each genotype) (**Figure. 2A, Figure 2 -**

figure supplement 1A) (**Ayyaz et al., 2019**). Uniform Manifold Approximation

and Projection (UMAP) was employed for dimensionality reduction to cluster

cells, then well-established marker genes were mapped onto the UMAP

representation to designate cell types (**Haber et al., 2017**) (**Figure. 2B, Figure**

2 - figure supplement 1B-D). A small population of crypt cells could not be

clearly defined probably due to certain technical issues during crypt extraction,

antibody staining, or cell sorting (**Figure 2 - figure supplement 1D**). First, in

accordance with bulk RNA-seq and RT-qPCR results, the expression levels of

Ctnnb1 were downregulated in almost all $Ctnnb1^{\Delta i.enh}$ crypt cells, and the

expression of *Axin2* was decreased in CBCs and TACs (**Figure. 2C and D**).

Second, the proportion of PCs was decreased by 49.2% in $Ctnnb1^{\Delta i.enh}$ crypts,

which was confirmed by the immunohistochemical (IHC) staining of *Defensin*

5a, a PC marker (**Figure. 2E and F**). The ieCtnnb1-driving *LacZ* could be

detected in the PC cluster (n = 4 10-week-old mice) (**Figure 2 - figure**

supplement 1E). Accordingly, the expression levels of examined PC markers

– *Lyz1*, *Pla2g2a*, *Wnt3* and *Math1* - were all significantly compromised in

Ctnnb1^{Δi.enh} crypts (**Figure. 2G**). The number of goblet cells (GCs) was slightly

(not statistically significant) reduced upon loss of ieCtnnb1 (**Figure 2 - figure**

supplement 1F). Third, the proliferative capability of CSCs and TACs were

compromised in *Ctnnb1^{Δi.enh}* crypts, as the proportion of G2/M cells and the

expression of *Mki67* were decreased in the two cell types (**Figure. 2H-J**).

Consistently, the numbers of Ki67-positive cells and EdU-labeled cells (4 hours

pulse) were decreased by 16.1% and 21.9%, respectively, in *Ctnnb1^{Δi.enh}* small

intestinal epithelia (**Figure. 2K and L**). Fourth, in secretory cells (PCs GCs and

EECs), as well as enterocytes/absorptive cells (mature enterocytes (mECs)

and immature enterocytes (imECs)), and stem cells of *Ctnnb1^{Δi.enh}* crypts, the

expression of genes related to secretory functions including *Lyz1*, *Ang4*, *Muc2*,

and *Mmp7* was decreased. In contrast, the expression of markers for absorptive

cells such as *Fabp6*, *Apoa1*, *Fabp2*, and *Reg3b* was elevated (**Figure. 2M and**

N, Figure 2 - figure supplement 1G). Accordingly, gene ontology (GO)

analyses showed in stem cells and secretory cells of *Ctnnb1^{Δi.enh}* crypts,

upregulated genes were enriched with terms associated with absorptive

functions including ribosome biogenesis and oxidative phosphorylation;

whereas downregulated genes in stem cells and secretory cells were

respectively enriched with terms related to cell division and secretory functions

(**Figure 2 - figure supplement 1H and I**). In contrast, in mECs, downregulated

genes were enriched with terms associated with absorptive functions (**Figure 2**

- figure supplement 1J).

In summary, the loss of ieCtnnb1 results in a reduction in the number of

PCs, compromises stem cell proliferation, and disrupts the expression of genes

related to secretory and absorptive functions. These effects could be attributed

to a combination of reduced Wnt/β-catenin signaling and dysregulated

crosstalk among crypt cells.

ieCtnnb1 deletion inhibits tumorigenesis of intestinal cancers by suppressing Wnt/β-catenin signaling

Tumorigenesis of most colorectal cancer (CRC) is associated with aberrant activation of the Wnt/β-catenin signaling (**Zhu & Li, 2023**). To investigate whether knockout of ieCtnnb1 could suppress the carcinogenesis of Wnt-driven intestinal tumors, we crossbred $Ctnnb1^{\Delta i.enh}$ mice with $Apc^{Min/+}$ mice. In $Apc^{Min/+}$ mice, a nonsense mutation of the Apc gene results in the translation of truncated and dysfunction APC protein, leading to failed degradation and nuclear translocation of β-catenin. $Apc^{Min/+}$ mice spontaneously develop intestinal tumors due to enhanced canonical Wnt/β-catenin signaling in the intestinal epithelia, particularly in stem cells (Su et al., 1992). The survival period of $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ mice (median survival - 255 days) was significantly longer than the $Apc^{Min/+}$ mice (median survival - 165 days), with the former having a heavier body weight (**Figure. 3A and C**). Strikingly, the $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ mice barely grow colorectal cancers and have much fewer intestinal tumors compared to $Apc^{Min/+}$ mice, with significantly smaller tumor area (**Figure. 3B, D-F, Figure 3 - figure supplement 1A-C**). IHC analyses showed that tumors in $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ large and small intestines had much fewer Ki67+ proliferating cells, expressed significant less β-catenin, but contained more Muc2-expressing Goblet cells, suggesting $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ tumors behave like normal epithelia (**Figure. 3G-L, Figure 3 - figure supplement 1D-I**). Consistently, RNA-seq transcriptome experiments showed that $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ colorectal cancers significantly downregulated genes associated with the Wnt signaling pathway, pluripotency of stem cells, and gastric cancer; whereas increased expression of genes related to normal epithelial functions including oxidative phosphorylation, adhesion, protein digestion and absorption, and Mucin biosynthesis (**Figure. 3M-O**). Thus, ablation of ieCtnnb1 deters tumorigenesis of intestinal cancers by suppressing the Wnt signaling.

Human ieCTNNB1 drives the expression of reporter gene in the gastrointestinal tract

We next investigated whether the human genome also harbors an intestinal enhancer for *CTNNB1*. A region with open chromatin (DNase I hypersensitivity) and enrichment of H3K27ac/H3K4me1 was identified at 38,489 bp upstream of human *CTNNB1* in adult small intestine and colon tissues (**Figure. 4A**). Moreover, the H3K27ac enrichment is also present in esophagus but not in non-GI tissues (**Figure 4 - figure supplement 1A**). Similar to ieCtnnb1, the 3 kb-long human genomic region, named as ieCTNNB1, resides in the same TAD as the promoter of *CTNNB1* (**Figure 4 - figure supplement 1B**). We generated ieCTNNB1 reporter mice ($H11^{hi.enh}$) and observed that ieCTNNB1 could drive *LacZ* expression in the bottom of small intestinal crypts and the entire epithelia of large intestine, also exhibiting a proximal-low to distal-high pattern (**Figure. 4B**). Notably, the LacZ signal driven by human ieCTNNB1 was less intense than that driven by mouse ieCtnnb1, potentially attributed to species-specific TF-enhancer interactions.

The H3K27ac-enriched region of ieCTNNB1 was divided into five equal parts for the luciferase reporter assay. Experiments demonstrated that each of the five ieCTNNB1 subregions individually increased luciferase activity in both HCT116 and HeLa cells (**Figure. 4C, Figure 4 - figure supplement 1C**). We then conducted CRISPR mediated activation and interference in HCT-15 colorectal cancer cells using three guide RNAs (gRNAs) targeting ieCTNNB1, which respectively enhanced and inhibited transcription of *CTNNB1* (**Figure. 4D**). Together, ieCTNNB1 is the intestinal enhancer of *CTNNB1* for human GI epithelia.

ieCTNNB1 is activated in colorectal cancer and its activity positively correlates with the expression of *CTNNB1*

CRC, one of the most prevalent cancers in human, is often associated with

aberrant activation of the Wnt signaling (**J. Li et al., 2021**). We then studied

whether the occurrence of CRC is associated with enhancer activation of

ieCTNNB1. We performed paired analyses of chromatin immunoprecipitation

sequencing (ChIP-seq) and RNA-seq data derived from 68 CRC samples and

their neighboring normal (native) tissue (**Figure. 5A, Figure 5 - figure**

supplement 1A). As anticipated, the expression levels of *CTNNB1* in CRC

samples are significantly higher than those in neighboring native tissues

(**Figure. 5B**). Essentially, the enrichment of H3K27ac at ieCTNNB1 in CRC

tissues were significantly higher than that in native tissues (**Figure. 5C, Figure**

5 - figure supplement 1A). In contrast, the enrichment of H3K4me3 at

CTNNB1's promoter was comparable between cancer and native tissues

(**Figure. 5C**). Moreover, the enrichment of H3K27ac at ieCTNNB1 positively

correlated with the expression of *CTNNB1*; but the enrichment of H3K4me3 at

CTNNB1's promoter was not associated with *CTNNB1*'s expression (**Figure.**

5D). Therefore, it's the activity of enhancer but not the promoter that reflects

the occurrence of CRC and expression levels of *CTNNB1*.

By examining Genotype Tissue Expression (GTEx) eQTL data browser, a

single-nucleotide polymorphism (SNP) – rs15981379 (C>T) within ieCTNNB1

was discovered (**Figure. 5A**). Importantly, the C>T variation is inversely

correlated with expression levels of *CTNNB1* in the esophagus and transverse

colon (**Figure. 5E**). Luciferase reporter assays in HCT116 and HeLa cells

revealed that the C>T variation indeed compromised luciferase activity driven

by the ieCTNNB1-region 1 (**Figure. 5F**).

HNF4α and p-S133-CREB1 bind to ieCTNNB1 and maintains the

expression of *CTNNB1*

Characterizing associated trans-factors of ieCtnnb1 and ieCTNNB1 is

essential for revealing related transcriptional mechanisms and holds clinical

relevance. By examining Cistrome Data Browser for ChIP-seq data, eight trans-factors were found to associate with ieCTNNB1 in human colorectal cancer cells or ieCtnnb1 in mouse GI tissues (**Figure. 6A, Figure 6 - figure supplement 1A**). Moreover, they are relatively highly expressed and/or have positive correlation with *CTNNB1*'s expression in colon cancer samples. We then transduced shRNAs-expressing plasmids into HCT-15 cells to individually down-regulate their expressions, revealing that decreasing the expression of *CREB1* (cAMP responsive element binding protein 1) or *HNF4A* (hepatocyte nuclear factor 4 alpha) could compromise *CTNNB1*'s transcription (**Figure. 6B, Figure 6 - figure supplement 1B**). Of note, HNF4α plays crucial roles in the normal development of the colon and the maintenance of epithelial barrier (**Cattin et al., 2009; Garrison et al., 2006**). The phosphorylation of CREB1 at serine 133 (p-S133-CREB1) is essential for its role in activation of gene transcription (**Carlezon et al., 2005**). ChIP-qPCR experiments showed that enrichment of p-S133-CREB1 was more prominent than CREB1 at ieCtnnb1 in small intestinal crypts and at ieCTNNB1 in HCT-15 cells (**data not shown**). Notably, HNF4α and p-S133-CREB1 bind to multiple subregions of ieCTNNB1 and ieCtnnb1, as well as the promoters of *CTNNB1* and *Ctnnb1* (*pCTNNB1* and *pCtnnb1*) in HCT-15 cells and small intestinal crypts (**Figure. 6C and D, Figure 6 - figure supplement 1C and D**). By analyzing the GEIPA-COAD (colon adenocarcinoma) and READ (rectum adenocarcinoma) database, expression levels of *CREB1* and *HNF4A* were found to be significantly higher in colon and rectum cancer samples than those of normal tissues (**Figure. 6E**). Furthermore, expression levels of *CREB1* and *HNF4A* positively correlated with that of *CTNNB1* (**Figure. 6F**). HNF4α and p-S133-CREB1 were significantly enriched at the promoter of *Ctnnb1* in *Apc* mutation induced colon cancer tissues (**Figure. 6G and H**). Intriguingly, knockout of ieCtnnb1 greatly diminished the association of HNF4α and p-S133-CREB1 with *Ctnnb1*'s promoter (**Figure. 6G and H**). In summary, HNF4α and p-S133-CREB1 were identified as essential

trans-factors that bind to ieCTNNB1 and maintain *CTNNB1*'s expression.

Discussion

The intestinal epithelia replenish themselves rapidly throughout lifetime,

which enables regeneration while predisposes to GI cancers (**Clevers, 2013**).

The optimal strength and dynamics of canonical Wnt/β-catenin signaling is

central to both homeostasis and tumorigenesis of GI tracts (**Beumer & Clevers,**

2016; Pinto et al., 2003; Silvia Maretto, 2003). While the stabilization and

nuclear translocation of β-catenin is the key step for Wnt signaling activation,

the transcriptional control of *Ctnnb1* could be another layer of regulation **(Rim**

et al., 2022; Zhang & Wang, 2020). Here we unveiled an enhancer-dependent

transcriptional machinery that tunes expressions of *Ctnnb1* in multiple cell types

of intestinal crypts, thereby balancing homeostasis and tumorigenesis of

intestinal epithelia (**Figure supplement 8**).

Our findings reinforce the notion that the canonical Wnt signaling is

required for the establishment of the secretory cell lineage (**Pinto & Clevers,**

2005; Pinto et al., 2003; van Es et al., 2005). Expression levels of markers for

secretory cells were greatly reduced whereas levels of markers for absorptive

cells were elevated in all small intestinal crypt cells of *Ctnnb1$^{\Delta i.enh}$* mice. Fate

determinations of CBC daughter cells occur upon leaving the stem cell niche,

exposed to changing WNT, Notch and EGF levels. Progenitors losing contact

with PC Notch ligands upregulate DLL1 and DLL4, commit to the secretory

lineage, and induce Notch activity in neighboring cells, promoting enterocyte

differentiation (**Beumer & Clevers, 2020; van Es et al., 2012**).

WNT signaling not only directs PC-fated progenitors toward the crypt

bottom but also directly induces their differentiation (**Andreu et al., 2008;**

Beumer & Clevers, 2020; van Es et al., 2005). Single-cell sequencing and

reporter gene analyses indicated that the activity of ieCtnnb1 is present in PCs.

Knockout of ieCtnnb1 decreases number of PCs by almost 50% but the pool

size of CBCs was not significantly altered. A previous study showed that genetic
removal of PCs leads to the concomitant loss of Lgr5+ stem cells (**Sato et al.,**
2011), while other studies indicated that PCs are not mandatory niche cells for
the intestinal epithelia (**Durand et al., 2012; Garabedian et al., 1997; Kim et**
al., 2012). Although Wnt ligands provided by PCs may support Lgr5+ CBCs[32],
essential Wnt ligands are primarily supplied by subepithelial stromal
populations (**Degirmenci et al., 2018; Shoshkes-Carmel et al., 2018**). The
presence of redundancy at the stem cell niche enhances the accuracy in
maintaining intestinal homeostasis.

ieCtnnb1 knockout compromises Wnt signaling dosage, leading to PC loss,
dysregulation of genes controlling absorptive and secretory functions of
intestinal epithelia, and reduced proliferation of stem cells. These combined
effects could account for the lighter body weight observed in adult $Ctnnb1^{\Delta i.enh}$
mice.

On the other hand, colon and small intestinal tumors were rarely observed
in $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ mice. Among the few $Ctnnb1^{\Delta i.enh}$ tumors analyzed, they
exhibited lower $Ctnnb1$ expression, reduced proliferation, and behaved more
like normal intestinal epithelia. Different truncations or mutations of APC lead
to specific tumor types throughout the body in a Wnt/β-catenin dosage-
dependent manner. For example, although $Apc^{1638N};Ctnnb1^{+/-}$ mice were less
prone to GI cancers, $Apc^{1638N};Ctnnb1^{+/-}$ female animals tended to grow breast
cancers (**Bakker et al., 2013**). Therefore, the intestinal specificity of ieCtnnb1
makes it an ideal target to lower the strength of Wnt signaling in treating
colorectal cancer without affecting other tissues (**Hamdan & Johnsen, 2019;**
Orouji, 2022).

Both mouse ieCtnnb1 and human ieCTNNB1 were able to drive reporter
gene expression at the base of small intestinal crypts and throughout the
epithelium of the large intestine. Very intriguingly, unlike the highly conserved
neCtnnb1 sequence among amniotes, mouse ieCtnnb1 and human ieCTNNB1

do not share significant homology in primary sequence (**Wang et al., 2022**).

Nonetheless, both ieCtnnb1 and ieCTNNB1 localize at the intergenic region 30-

40 kb upstream of *Ctnnb1*'s promoter. Moreover, both ieCtnnb1 and ieCTNNB1

contain binding motifs for HNF4α and CREB1. HNF4α and p-S133-CREB1

were found to associate with ieCtnnb1 and *Ctnnb1*'s promoter to maintain

Ctnnb1's expression. These findings support the notion that conserved TF

binding motifs stabilize and preserve enhancer functionality over evolution

(Wong et al., 2020; Zabidi & Stark, 2016). Future studies could dissect how

sequence variations of ieCtnnb1 are adapted to intestinal homeostasis of

different species.

Methods
Mice and genotyping

All animal procedures were approved by the Animal Care and Ethical

Committee of Medical Research Institute at Wuhan University. C57BL/6 back-

ground mice were housed in a temperature- and humidity- controlled

environment with 12hr light/dark cycle and food/water ad libitum. *Ctnnb1$^{\Delta i.enh}$*

mice were generated in the Beijing Biocytogen. *H11$^{i.enh}$* reporter mice were

generated in the Shanghai Model Organisms Center Inc. BAT-Gal mice were

provided by Dr. Jun-Lei Chang (Jackson Lab, stock number 005317). *Lgr5*-

EGFP-IRES-CreERT2 mice were gifts from Dr. Ye-Guang Chen (Jackson Lab,

stock number 008875). *APC$^{Min/+}$* mice were gifts from Dr. Bo Zhong (Jackson

Lab, stock number 002020). The primer set forward 5′-atcctctgcatggtcaggtc-

3′/reverse 5′-cgtggcctgattcattcc-3′ was used for genotyping of BAT-Gal mice,

H11$^{i.enh}$ mice and *H11$^{hi.enh}$* mice with a band size of 315 base pairs (bp). The

primer set forward 5′-gtcctgtccgtcactattatcctggc-3′/reverse5′-

ccactgccctgctaaagcattggt-3′/reverse 5′-tgattagttctccgggaagcccagt-3′ were

used for genotyping of *Ctnnb1$^{\Delta i.enh}$* mice and band sizes for *Ctnnb1$^{\Delta i.enh}$* mice

are 482 bp (wild-type allele) and 867 bp (ieCtnnb1 knockout allele). The primer

set forward *5′-atctcatggcaaacagacct-3′/reverse5′-tcacaaatcatctcgcaga-3′* was

used for genotyping of *APC$^{Min/+}$* mice with a band size of 340 bp. The primer set

forward   5′-ctgctctctgctcccagtct-3′/reverse5′-ataccccatccctttttgagc-3′/reverse 5′-gaacttcagggtcagcttgc-3′ were used for genotyping of *Lgr5*-EGFP-IRES-creERT2 mice and the band sizes for 386 bp (wild-type allele) and 119 bp (mutant allele).

Cell lines

HEK293T cells were gifts from Dr. Hong-Bing Shu (Wuhan University). HCT116, HCT-15 and HeLa cells were purchased from the China Center for Type Culture Collection. HEK293T, HeLa and HCT116 cells were grown in DMEM (Gibco) supplemented with 10% fetal bovine serum (FBS) (Life Technologies) and 1×penicillin/streptomycin (Gibco). HCT-15 cells were maintained in RPMI-1640 (Gibco) containing 10% FBS (Life Technologies) and 1×penicillin/streptomycin (Gibco). For routine culturing, cells were dissociated using 0.25% trypsin (Gibco).

Tissue fixation and section

Intestinal and tumor tissues were dissected and cleaned in pre-cooled phosphate buffered saline (PBS) and immersed in 4% paraformaldehyde (PFA) overnight at 4°C. For sectioning, intestines were embedded in Tissue-Tek® O.C.T. Compound (SAKURA) and cut to 8 μm thickness with a cryostat (Leica CM1950).

CRISPR/dCas9-mediated transcription activation (CRISPRa) and interference (CRISPRi) assay

CRISPRa and CRISPRi assay were performed as previously described (**Wang et al., 2022**). In brief, SgRNAs were designed using online tool (https://zlab.bio/guide-design-resources) and were listed in **Table S2**. SgRNAs were cloned into lenti sgRNA (MS2)_zeo backbone vector (Addgene, #61427). For CRISPRi assays, pHR-SFFV-dCas9-BFP-KRAB (Addgene, #46911) is used. For CRISPRa assays, lenti MS2-P65-HSF1_Hygro (Addgene, #61426) and dCAS9-VP64_GFP (Addgene, #61422) were used. To obtain lentiviral particles, HEK293T cells (5×10^6 cells per 10-cm dish) were transiently

transfected with 12 μg CRISPRa or CRISPRi constructs, 6 μg psPAX2 and 6 μg pMD2.G. The supernatant containing lentivirus particles was harvested at 72 hours after transfection and filtered through Millex-GP Filter Unit (0.45 μm pore size, Millipore). To obtain high virus titer, the supernatant was centrifuged at 10,000×g for 100 minutes and concentrate to 100 μl. On the day before infection, HCT-15 cells (2 × 10^5 cells per well) were inoculated in 12 well plates. 10 μl high titer viruses were added to each well. 72 hours after infection, cells were harvested for RT-qPCR analysis.

RNA isolation and cDNA synthesis

RNAiso Plus (TAKARA) is used for RNA isolation. 1 ml RNAiso Plus was added to DNase/RNase free EP tube to lyse tissues or cells, followed by adding one-fifth volume of chloroform to achieve phase separation. After vigorous shaking, centrifuge at 12,000×g for 15 minutes at 4°C. The upper aqueous phase was transferred to a new DNase/RNase free EP tube and equal volume of isopropanol was added to precipitate RNA. Precipitation was dissolved in DNase/RNase-free water. Complementary DNAs (cDNAs) were synthesized by HiScript® II Q RT SuperMix for qPCR kit (Vazyme; R222-01).

Real-time quantitative reverse transcription PCR (RT-qPCR)

cDNA was subjected to RT-qPCR using the SYBR Green assay with 2× SYBR Green qPCR master mix (Bimake). PCR reactions were performed on a CFX Connect Real-Time PCR Detection System (Bio-rad) under the following condition: 5 minutes at 95°C, 40 cycles at 95°C for 15 seconds, 60°C for 20 seconds. The relative transcript level of each gene was normalized to *Gapdh* (mice) or *ACTB* (human) and calculated according to the 2$^{-\Delta\Delta Ct}$ method. All primers used are shown in the **Table S2**.

Isolation of murine intestinal crypt cells

Murine small intestines were harvested and cut open longitudinally. After 3 times wash in cold PBS with Penicillin (200 U/ml) + Streptomycin (200 μg/ml) (Invitrogen), small intestines were cut into 5 cm segments and incubated in pre-

cold PBS with 8 mM EDTA at 4°C for 2 hours. Then intestinal tissues were transferred into a centrifuge tube filled with new cold PBS. Shake vigorously for 5 minutes and collect intestinal epithelial cells (villi plus crypts) from the supernatant. Centrifuge the supernatant to collect all epithelial cells or pass through 70 μm cell filter (Corning) to filter out villi. Crypts were spun down in a centrifuge at $300 \times g$ for 5 minutes.

Enriching *Lgr5*-EGFP$^+$ cells

Crypts were collected from *Lgr5*-EGFP and *H11$^{i.enh}$;Lgr5*-EGFP mice, followed by resuspension and single-cell digestion in DMEM containing 0.25% trypsin (Gibco) at 37°C for 5 minutes. Disaggregated cells were passed through 40 μm cell strainers (Corning). *Lgr5*-EGFP$^+$ and *Lgr5*-EGFP$^-$ cells were collected using a BD Arial III cell sorter.

X-Gal staining

X-Gal staining is performed as described (**Wang et al., 2022**). In short, frozen slides were placed in fixed buffer (0.2% PFA, 0.1 M PIPES buffer (pH 6.9), 2 mM MgCl2, 5 mM EGTA) for 10 minutes, followed by rinse for two times with PBS containing 2 mM $MgCl_2$. Then slides were incubated with detergent solution (0.1 M PBS (pH 7.3), 2 mM MgCl2, 0.01% sodium deoxycholate, 0.02% Nonidet P-40) for 10 minutes. Slides were incubated with freshly prepared and filtered X-Gal staining solution (0.1 M PBS (pH 7.3), 2 mM MgCl2, 0.01% sodium-deoxycholate, 0.02% Nonidet P-40, 5 mM $K_3Fe(CN)_6$, 5 mM $K_4Fe(CN)_6 \cdot 3H_2O$ and 1 mg/ml X-Gal) overnight in the dark at 37°C. For whole gastrointestinal tissues, tissues were fixed in freshly prepared 4% PFA for 10 minutes, followed by rinse with detergent solution for 5 minutes by three times. Tissues were incubated with freshly prepared and filtered X-Gal staining solution at 37°C for a few hours.

EdU pulse-labeling experiment

Mice were intraperitoneally injected with EdU (10 mg/kg of body weight). After 4 hours, harvest the intestines for frozen sectioning. Staining was performed

592 using BeyoClick™ EdU Cell Promotion Kit (C0075S, Beyotime, China)

593 according to the manufacturer's manual.

ChIP-qPCR assay

Cells were cross-linked with 1% formaldehyde for 10 minutes at room

temperature and then quenched with 0.125 M glycine for 5 minutes and washed

twice with cold PBS. Cells were resuspended with 500 µl lysis buffer (50 mM

Tris-HCl, pH 8.0, 0.5% SDS, 5 mM EDTA) and subjected to sonication on ice

with an output power of 100 W, 12 minutes, 0.5 seconds on, 0.5 seconds off.

After centrifugation, 1% of the supernatant was taken out as input. The rest of

the sonicated lysates was diluted into 0.1% SDS and divided into several equal

parts. Immunoprecipitation was performed overnight at 4°C on a rotating wheel

with sheared chromatin, protein G agarose beads and indicated antibodies: 2

µg HNF4α (Abcam, ab181604), 2 µg p-S133-CREB1 (Abcam, ab32096), 2 µg

IgG antibody (ABclonal, AC005). The next day, beads were cleaned three times

with Wash Buffer I (20 mM Tris-HCl, pH 8.0; 1% Triton X-100; 2 mM EDTA; 150

mM NaCl; 0.1% SDS), Wash Buffer II (20 mM Tris-HCl, pH 8.0; 1% Triton X-

100; 2 mM EDTA; 500 mM NaCl; 0.1% SDS), Wash Buffer III (10 mM Tris-HCl,

pH 8.0; 1 mM EDTA; 0.25 M LiCl; 1% NP-40; 1% deoxycholate) and TE buffer.

Protein-DNA complexes were de-crosslinked in 120 µl elution buffer (0.1 M

NaHCO$_3$, 1% SDS, 20 μg/ml proteinase K) and shaken at 65°C overnight. DNA

purification kit (TIANGEN) was used to extract DNA fragments for qPCR.

Luciferase reporter assay

The luciferase reporter assay was performed as described previously (**Li et al.,**

2017). Briefly, the core region of ieCtnnb1, 5 regions of ieCTNNB1 and the

region containing the mutation site were respectively cloned into pGL3-

promoter vector (Promega). HEK293T, HeLa and HCT116 cells were

transfected with pGL3-promoter vector and the plasmids mentioned above.

Renilla basic vector was co-transfected as a control for normalization of

luciferase activity. After 48 hours of transfection, Promega Dual Glo test kit

(Promega, E1910) was used to measure luciferase activity according to

manufacturer's instructions.

Immunohistochemical staining

Frozen sections were dried at 37°C, fixed with 4% PFA for 10 minutes, and then

washed three times with PBS. Antigen retrieval was performed using citrate

buffer (pH 6, epitope retrieval solution) at 95°C for 20 minutes. Slides were then

placed in 3% H_2O_2 solution for a few hours to remove endogenous catalase.

After washing with PBS three times, slides were blocked with 10% normal goat

serum for 1 hour, then incubated with the following primary antibodies overnight

at 4°C: rabbit anti-α-defensin5 (Abcam, ab80515, 1:500), rabbit anti-Ki67

(Abcam, ab15580, 1:500), rabbit anti-β-catenin (Cell Signaling Technology,

8480T, 1:100), rabbit anti-Muc2 (Thermo Fisher Scientific, PA5-21329, 1:500).

The next day, slides were incubated with biotin labeled secondary antibodies

at room temperature for 1 hour. Then, slides were incubated with the avidin-

biotin-peroxidase complex (VECTASTAIN Elite ABC system, Vector Labs).

Finally, colorization was performed in the reaction solution (Tris-HCl (pH 7.2),

5 mg/ml 3,3'-diaminobenzidine, 0.075% H_2O_2).

H&E staining

H&E staining kit (Beyotime Biotech, C0105S) was used for experiments. Frozen

slides were dried at 65°C, then dripped with hematoxylin staining solution for 5

minutes. Excess staining solution was rinsed off with water. Slides were

immersed in distilled water for 5 minutes, then rinsed with 95% ethanol, and

stained with eosin solution for 30 seconds. Subsequently, slides were put into

anhydrous ethanol for 2 minutes, and cleared with xylene for 5 minutes. After

air dry, slides were sealed with neutral resin.

Immunoblotting analysis

Crypt cells were lysed by 1× SDS loading buffer at 95°C for 10 minutes. After

short centrifugation, the supernatants were loaded onto 10% SDS-

polyacrylamide gel electrophoresis gels. Then, proteins were transferred onto

polyvinylidene fluoride membrane (Millipore). Membranes were blocked with 10% non-fat milk in tris-buffered saline containing 0.3% Tween 20 (TBST; pH 7.4) for 1 hour at room temperature. Immunoblotting was carried out used primary antibody: rabbit anti-β-catenin (Cell Signaling Technology, 8480T, 1:100); mouse anti-GAPDH (ABclonal, AC001, 1:1000); rabbit anti-H3 (ABclonal, A2348, 1:10,000). After three times of washing with TBST, the membrane was incubated with anti-rabbit immunoglobulin G (IgG)–conjugated horseradish peroxidase secondary for 1 hour at room temperature. Signals were detected using the ECL substrate (Thermo Fisher Scientific).

Alcian Blue PAS Staining

Alcian Blue PAS Stain Kit (Abcam, ab245876) was used for experiments. all materials and prepared reagents were equilibrated to room temperature and gently agitated prior to use. Tissue sections were applied with acetic acid solution (3%) for 2 minutes. Then alcian blue solution (pH 2.5) was applied for 15-20 minutes. Slides were rinsed with running tap water for 2 minutes followed by 2 changes of distilled water. Periodic acid solution was applied to tissue sections for 5 minutes. After slides were rinsed with two changes of distilled water, Schiff's solution was applied to tissue sections for 10-20 minutes. Slides were rinsed with warm running tap water for 2 minutes followed by 2 changes of distilled water. Tissue sections were applied with hematoxylin for 2 minutes. Sections were rinsed in running tap water for 2 minutes followed by 2 changes of distilled water. Slides were dehydrated through graded alcohols and sealed with synthetic resin.

Cell Sorting

Crypt single cells were prepared as described above. Cells were resuspended using 1 ml PBS and stained with 5 μl anti-EpCAM-FITC (eBioScience, 11-5791-82) at 4°C for 30 minutes. After washing with PBS three times, cells were resuspended using 1 M FACS buffer (PBS with 1%BSA) containing 0.2 μg/ml DAPI (BD Pharmingen, Cat. No. 564907). Cells were filtered with nylon membrane with a 40 μm pore size. EpCAM positive cells were sorted and

collected based on fluorescence signals for subsequent single cell sequencing.

Flow cytometry data were acquired on a BD Arial III flow cytometer and

analyzed with FlowJo (version 10.6.2).

Bulk RNA sequencing

RNA-seq and data analyses were performed at Novogene Co., Ltd (Beijing,

China). Crypt cells or tumor tissues were harvested and total RNAs were

extracted. RNA integrity was assessed using the RNA Nano 6000 Assay Kit of

the Bioanalyzer 2100 system (Agilent Technologies, CA, USA). RNA-seq

library construction was performed as described previously (**Xu et al., 2021**).

Briefly, first strand cDNA was synthesized using random hexamer primer and

M-MuLV Reverse Transcriptase (RNase H-). Second strand cDNA synthesis

was subsequently performed using DNA Polymerase I and RNase H. Then, the

library fragments were purified with AMPure XP system (Beckman Coulter,

Beverly, USA). Then PCR was performed with Phusion High-Fidelity DNA

polymerase, Universal PCR primers and Index (X) Primer. At last, PCR

products were purified (AMPure XP system). The clustering of the index-coded

samples was performed on a cBot Cluster Generation System using TruSeq

PE Cluster Kit v3-cBot-HS (Illumia) according to the manufacturer's instructions.

After cluster generation, the library preparations were sequenced on an Illumina

Novaseq platform and 150 bp paired-end reads were generated.

Bulk RNA-seq data processing

For data analysis, raw data of fastq format were firstly processed through in-

house perl scripts. In this step, clean data were obtained by removing reads

containing adapter, reads containing ploy-N and low-quality reads from raw

data. Index of the reference genome was built using Hisat2 v2.0.5 and paired-

end clean reads were aligned to the reference genome using Hisat2 v2.0.5.

featureCounts v1.5.0-p3 was used to count reads numbers mapped to each

gene. Differential expression analysis of two groups was performed using the

DESeq2 R package (1.20.0). The resulting P-values were adjusted using the

Benjamini and Hochberg's approach for controlling the false discovery rate.

Genes with an adjusted P-value <0.05 found by DESeq2 were assigned as differentially expressed. Gene Ontology (GO) enrichment analysis of differentially expressed genes was implemented by the clusterProfiler R package, in which gene length bias was corrected. clusterProfiler R package was also used to test the statistical enrichment of differential expression genes in KEGG pathways. Local version of the GSEA analysis tool (http://www.broadinstitute.org/gsea/index.jsp), GO、KEGG、Reactome、DO and DisGeNET data sets were used for GSEA independently.

Single-Cell RNA sequencing and data analysis

ScRNA-seq and data analysis were performed at Novogene Co., Ltd (Beijing, China). The protoplast suspension was loaded into Chromium microfluidic chips with 30 chemistry (v2 or v3) and barcoded with a 10× Chromium Controller (10X Genomics). RNA from the barcoded cells was subsequently reverse-transcribed and sequencing libraries constructed with reagents from a Chromium Single Cell 30 v2 reagent kit (10X Genomics) according to the manufacturer's instructions. Sequencing was performed with Illumina HiSeq 2000 according to the manufacturer's instructions. For data analysis, we use FastQC to perform basic statistics on the quality of the raw reads. Raw reads were demultiplexed and mapped to the reference genome by 10X Genomics Cell Ranger pipeline using default parameters. All downstream single-cell analyses were performed using Cell Ranger and Seurat unless mentioned specifically. In brief, for each gene and each cell barcode (filtered by CellRanger), unique molecule identifiers were counted to construct digital expression matrices. Secondary filtration by Seurat: a gene with expression in more than 3 cells was considered as expressed, and each cell was required to have at least 200 expressed genes.

Hi-C data analysis

Hi-C data analysis completed by Frasergen Medicine Co., Ltd (Wuhan, China). Hi-C data of BALB/c mice large intestinal epithelial cells were provided by Dr. Gen Zheng(Zheng et al., 2023). The Hi-C data of SW480 cells was obtained

from deposited data(Orouji, 2022). According to the reported method, the
dynamic changes of TAD structures can be explored by calculating the relative
signal values of TAD boundaries. TAD boundaries that are common among
individuals and are not within the 2 Mb range of chromosomal ends were
included, followed by filtering out boundaries with interactions less than 10 in
the 2 Mb upstream and downstream regions. The ratio of the interaction
between the boundary and its 2 Mb upstream and downstream regions were
calculated using log2 to obtain the signal value of the TAD boundary, then the
minimum value was subtracted from it to finally obtain the distribution of the
relative signal value of each sample TAD boundary.

ChIP-seq data analysis
The human colorectal cancer patient RNA-seq FPKM files and ChIP-seq data
were from GEO dataset GSE156614 (**Q. L. Li et al., 2021**). All ChIP-seq raw
fastq data were removed of adaptor sequence. Cutadapt (version 1.16) was
used with the parameters -u 3 -u -10 -U 3 -U -10 -m 30. Cleaned reads were
aligned to the human reference genome (hg19) using bowtie2 (version 2.4.4)
with default settings. The CrossMap (0.6.5) package was used as a conversion
between different versions of the human genome (hg19 and hg38). We
calculated the normalized RPM (reads per million mapped reads) of ChIP-seq
signal at ieCTNNB1 and *CTNNB1* promoter. Briefly, reads aligned to the
specified region were calculated with bedtools multicov command. These reads
were normalized with RPM. RNA-seq data FPKM (Reads per kilobase per
million mapped reads) were converted to TPM (Transcript per Kilobase per
Million mapped reads) using R version 4.2.3.

Knockdown experiments
Target sequences for shRNAs were designed on the Merck Life Sciences
website (https://www.sigmaaldrich.cn) and were listed in **Table S2**.
Oligonucleotides for shRNAs were cloned into pLKO.1 vector (Addgene,
#21297). The day before transfection, HEK293T cells (2×10^5 cells per well)
were seeded in 12-well plates. pLKO.1 constructs, psPAX2 and pMD2.G were

co-transfected into HEK293T cells. After 72 hours, supernatant containing virus

particles was harvested and passed through Millex-GP Filter Unit (0.45 μm pore

size, Millipore). Viral particles were aliquoted and stored at -80°C. On the day

before infection, HCT-15 cells (2×10^5 cells per well) were seeded in 12-well

plates. 20 μl virus suspension were added to each well. After 72 hours, cells

were harvested for RT-qPCR to evalutate knockdown efficiency.

Reference database

ChIP-seq signals for H3K27ac, H3K4me3, H3K4me1, H3K36me3, and DNase

I hypersensitivity data of stomach, intestines, liver, and other tissues at different

stages were downloaded from the ENCODE (https://www.encodeproject.org/)

and their identifiers are listed in **Table S1**. In addition, ChIP-seq data for listed

transcription factors of cells or tissues were obtained from Cistrome Data

Browser (http://cistrome.org). Expressions of *HNF4A* and *CREB1* in human

colon (COAD) and rectal (READ) cancers were obtained from GEPIA

(gepia.cancer-pku.cn).

eQTL analysis

The GTEX database (https://www.gtexportal.org/home/) was used to search

SNPs at *CTNNB1* and the 1 MB upstream and downstream region, followed by

eQTL analysis. A variant, rs115981379 (C>T), was identified inside *CTNNB1*.

The expression levels of *CTNNB1* in 428 C/C type and 37 C/T type esophagus

samples, and 343 C/C type and 25 C/T type transverse colon samples were

also obtained in the GTEX database.

Analyses of HNF4α and CREB1 binding motifs

The JASPAR database (https://jaspar.elixir.no/) was used to analyze binding

motifs of HNF4α and CREB1 at ieCtnnb1/ieCTNNB1 and *Ctnnb1/CTNNB1*

promoter. Results are listed in **Table S3**.

Nuclear-cytoplasmic separation

Nuclear and cytoplasmic protein extraction kit (Beyotime Biotech, P0027) was

used for experiments. Small intestinal crypts were collected from wild-type and

Ctnnb1$^{\Delta i.enh}$ mice as described above. According to manufacturer's instructions,

20 μl cell precipitation was resuspended in 200 μl cytoplasmic protein extraction

reagent A with 1 mM PMSF. After placing on ice for 15 minutes, 10 μl

cytoplasmic protein extraction reagent B was added. After vigorous vortex,

solutions were centrifuged at 12,000×*g* for 5 minutes at 4°C. The supernatant

was collected as the extracted cytoplasmic protein. The precipitate was

resuspended with 50 μl nuclear protein extraction reagent with 1 mM PMSF.

Solutions were placed on ice and vortex vigorously every 2 minutes for a total

of 30 minutes. After centrifuging at 12,000×*g* for 10 minutes at 4°C, the

supernatant was collected as the extracted nuclear protein.

Statistics

All experiments reported in this study were repeated at least three independent

times. *In vivo* analyses were conducted with at least three animals per condition.

The number of animals in each experiment is depicted in Figure legends. The

signal intensity of tumor stain region was generated using ImageJ software

(version 6.0.0.260) as described previously (**Wang et al., 2021**). GraphPad

Prism (version 8.0.2) was used to determine statistical significance. Unpaired

two-tailed Student's t-test were used for analysis between two groups of equal

variances by *F*-tests. When the *F*-test of equal variance failed, the Welch *t*-tests

were used. Data comparison of three or more groups with control groups were

analyzed using one-way ANOVA followed by Dunnett's multiple comparison

test. Data comparison of three or more groups among each other were

analyzed using one-way ANOVA followed by Tukey's multiple comparison test.

The exact *P* values were reported in each figure or indicated as ****, $P < 0.0001$;

***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$, ns = not significant.

Data and code availability

The GEO accession number for the bulk RNA-seq and scRNA-seq data

reported in this paper is GSE233979. RNA-seq data of mouse small intestinal

crypt and colon tumor have been deposited at GEO: GSE233977. ScRNA-seq

data of mouse small intestinal crypt cells in this study have been deposited at

GEO: GSE233978. All data are publicly available as of the date of publication.

Custom codes were described in detail at methods part. Any additional

information required to analyze the data in this paper is available from authors

upon reasonable request.

References

Andreu, P., Peignon, G., Slomianny, C., Taketo, M. M., Colnot, S., Robine, S., Lamarque,
D., Laurent-Puig, P., Perret, C., & Romagnolo, B. 2008. A genetic study of the role
of the Wnt/beta-catenin signalling in Paneth cell differentiation. *Dev Biol, 324*, 288-
296. https://doi.org/10.1016/j.ydbio.2008.09.027

Ayyaz, A., Kumar, S., Sangiorgi, B., Ghoshal, B., Gosio, J., Ouladan, S., Fink, M., Barutcu,
S., Trcka, D., Shen, J., Chan, K., Wrana, J. L., & Gregorieff, A. 2019. Single-cell
transcriptomes of the regenerating intestine reveal a revival stem cell. *Nature, 569*,
121-125. https://doi.org/10.1038/s41586-019-1154-y

Bakker, E. R., Hoekstra, E., Franken, P. F., Helvensteijn, W., van Deurzen, C. H., van
Veelen, W., Kuipers, E. J., & Smits, R. 2013. beta-Catenin signaling dosage
dictates tissue-specific tumor predisposition in Apc-driven cancer. *Oncogene, 32*,
4579-4585. https://doi.org/10.1038/onc.2012.449

Barker, N., van Es, J. H., Kuipers, J., Kujala, P., van den Born, M., Cozijnsen, M.,
Haegebarth, A., Korving, J., Begthel, H., Peters, P. J., & Clevers, H. 2007.
Identification of stem cells in small intestine and colon by marker gene Lgr5. *Nature,*
449, 1003-1007. https://doi.org/10.1038/nature06196

Barreto, E. B. L., Rattes, I. C., da Costa, A. V., & Gama, P. 2022. Paneth cells and their
multiple functions. *Cell Biol Int, 46*, 701-710. https://doi.org/10.1002/cbin.11764

Beumer, J., & Clevers, H. 2016. Regulation and plasticity of intestinal stem cells during
homeostasis and regeneration. *Development, 143*, 3639-3649.
https://doi.org/10.1242/dev.133132

Beumer, J., & Clevers, H. 2020. Cell fate specification and differentiation in the adult
mammalian intestine. *Nature Reviews Molecular Cell Biology, 22*, 39-53.
https://doi.org/10.1038/s41580-020-0278-0

Carlezon, W. A., Jr., Duman, R. S., & Nestler, E. J. 2005. The many faces of CREB. *Trends*
Neurosci, 28, 436-445. https://doi.org/10.1016/j.tins.2005.06.005

Cattin, A. L., Le Beyec, J., Barreau, F., Saint-Just, S., Houllier, A., Gonzalez, F. J., Robine,
S., Pincon-Raymond, M., Cardot, P., Lacasa, M., & Ribeiro, A. 2009. Hepatocyte
nuclear factor 4alpha, a key factor for homeostasis, cell architecture, and barrier
function of the adult intestinal epithelium. *Mol Cell Biol, 29*, 6294-6308.
https://doi.org/10.1128/MCB.00939-09

Cheng, X., Xu, X., Chen, D., Zhao, F., & Wang, W. 2019. Therapeutic potential of targeting the Wnt/beta-catenin signaling pathway in colorectal cancer. *Biomed Pharmacother, 110*, 473-481. https://doi.org/10.1016/j.biopha.2018.11.082

Chin, A. M., Hill, D. R., Aurora, M., & Spence, J. R. 2017. Morphogenesis and maturation of the embryonic and postnatal intestine. *Semin Cell Dev Biol, 66*, 81-93. https://doi.org/10.1016/j.semcdb.2017.01.011

Clevers, H. 2006. Wnt/beta-catenin signaling in development and disease. *Cell, 127*, 469-480. https://doi.org/10.1016/j.cell.2006.10.018

Clevers, H. 2013. The intestinal crypt, a prototype stem cell compartment. *Cell, 154*, 274-284. https://doi.org/10.1016/j.cell.2013.07.004

Clevers, H., Loh, K. M., & Nusse, R. 2014. Stem cell signaling. An integral program for tissue renewal and regeneration: Wnt signaling and stem cell control. *Science, 346*, 1248012. https://doi.org/10.1126/science.1248012

Clevers, H., & Nusse, R. 2012. Wnt/beta-catenin signaling and disease. *Cell, 149*, 1192-1205. https://doi.org/10.1016/j.cell.2012.05.012

Degirmenci, B., Valenta, T., Dimitrieva, S., Hausmann, G., & Basler, K. 2018. GLI1-expressing mesenchymal cells form the essential Wnt-secreting niche for colon stem cells. *Nature, 558*, 449-453. https://doi.org/10.1038/s41586-018-0190-3

Durand, A., Donahue, B., Peignon, G., Letourneur, F., Cagnard, N., Slomianny, C., Perret, C., Shroyer, N. F., & Romagnolo, B. 2012. Functional intestinal stem cells after Paneth cell ablation induced by the loss of transcription factor Math1 (Atoh1). *Proceedings of the National Academy of Sciences, 109*, 8965-8970. https://doi.org/10.1073/pnas.1201652109

Garabedian, E. M., Roberts, L. J. J., McNevin, M. S., & Gordon, J. I. 1997. Examining the Role of Paneth Cells in the Small Intestine by Lineage Ablation in Transgenic Mice. *Journal of Biological Chemistry, 272*, 23729-23740. https://doi.org/10.1074/jbc.272.38.23729

Garrison, W. D., Battle, M. A., Yang, C., Kaestner, K. H., Sladek, F. M., & Duncan, S. A. 2006. Hepatocyte nuclear factor 4alpha is essential for embryonic development of the mouse colon. *Gastroenterology, 130*, 1207-1220. https://doi.org/10.1053/j.gastro.2006.01.003

Gribble, F. M., & Reimann, F. 2019. Function and mechanisms of enteroendocrine cells and gut hormones in metabolism. *Nature Reviews Endocrinology, 15*, 226-237. https://doi.org/10.1038/s41574-019-0168-8

Gustafsson, J. K., & Johansson, M. E. V. 2022. The role of goblet cells and mucus in intestinal homeostasis. *Nature Reviews Gastroenterology & Hepatology, 19*, 785-803. https://doi.org/10.1038/s41575-022-00675-x

Haber, A. L., Biton, M., Rogel, N., Herbst, R. H., Shekhar, K., Smillie, C., Burgin, G., Delorey, T. M., Howitt, M. R., Katz, Y., Tirosh, I., Beyaz, S., Dionne, D., Zhang, M., Raychowdhury, R., Garrett, W. S., Rozenblatt-Rosen, O., Shi, H. N., Yilmaz, O., Xavier, R. J., & Regev, A. 2017. A single-cell survey of the small intestinal epithelium. *Nature, 551*, 333-339. https://doi.org/10.1038/nature24489

Hamdan, F. H., & Johnsen, S. A. 2019. Perturbing Enhancer Activity in Cancer Therapy. *Cancers (Basel), 11*. https://doi.org/10.3390/cancers11050634

Huels, D. J., Bruens, L., Hodder, M. C., Cammareri, P., Campbell, A. D., Ridgway, R. A., Gay, D. M., Solar-Abboud, M., Faller, W. J., Nixon, C., Zeiger, L. B., McLaughlin, M. E., Morrissey, E., Winton, D. J., Snippert, H. J., van Rheenen, J., & Sansom, O. J. 2018. Wnt ligands influence tumour initiation by controlling the number of intestinal stem cells. *Nature Communications, 9*. https://doi.org/10.1038/s41467-018-03426-2

Jindal, G. A., & Farley, E. K. 2021. Enhancer grammar in development, evolution, and disease: dependencies and interplay. *Dev Cell, 56*, 575-587. https://doi.org/10.1016/j.devcel.2021.02.016

Kim, T.-H., Escudero, S., & Shivdasani, R. A. 2012. Intact function of Lgr5 receptor-expressing intestinal stem cells in the absence of Paneth cells. *Proceedings of the National Academy of Sciences, 109*, 3932-3937. https://doi.org/10.1073/pnas.1113890109

Krausova, M., & Korinek, V. 2014. Wnt signaling in adult intestinal stem cells and cancer. *Cell Signal, 26*, 570-579. https://doi.org/10.1016/j.cellsig.2013.11.032

Kurokawa, K., Hayakawa, Y., & Koike, K. 2020. Plasticity of Intestinal Epithelium: Stem Cell Niches and Regulatory Signals. *Int J Mol Sci, 22*. https://doi.org/10.3390/ijms22010357

Kvon, E. Z., Waymack, R., Gad, M., & Wunderlich, Z. 2021. Enhancer redundancy in development and disease. *Nat Rev Genet, 22*, 324-336. https://doi.org/10.1038/s41576-020-00311-x

Kvon, E. Z., Zhu, Y., Kelman, G., Novak, C. S., Plajzer-Frick, I., Kato, M., Garvin, T. H., Pham, Q., Harrington, A. N., Hunter, R. D., Godoy, J., Meky, E. M., Akiyama, J. A., Afzal, V., Tran, S., Escande, F., Gilbert-Dussardier, B., Jean-Marcais, N., Hudaiberdiev, S., Ovcharenko, I., Dobbs, M. B., Gurnett, C. A., Manouvrier-Hanu, S., Petit, F., Visel, A., Dickel, D. E., & Pennacchio, L. A. 2020. Comprehensive In Vivo Interrogation Reveals Phenotypic Impact of Human Enhancer Variants. *Cell, 180*, 1262-1271 e1215. https://doi.org/10.1016/j.cell.2020.02.031

Li, J., Ma, X., Chakravarti, D., Shalapour, S., & DePinho, R. A. 2021. Genetic and biological hallmarks of colorectal cancer. *GENES & DEVELOPMENT, 35*, 787-820. https://doi.org/10.1101/gad.348226

Li, Q. L., Lin, X., Yu, Y. L., Chen, L., Hu, Q. X., Chen, M., Cao, N., Zhao, C., Wang, C. Y., Huang, C. W., Li, L. Y., Ye, M., & Wu, M. 2021. Genome-wide profiling in colorectal cancer identifies PHF19 and TBC1D16 as oncogenic super enhancers. *Nat Commun, 12*, 6407. https://doi.org/10.1038/s41467-021-26600-5

Li, Y., Wang, W., Wang, F., Wu, Q., Li, W., Zhong, X., Tian, K., Zeng, T., Gao, L., Liu, Y., Li, S., Jiang, X., Du, G., & Zhou, Y. 2017. Paired related homeobox 1 transactivates dopamine D2 receptor to maintain propagation and tumorigenicity of glioma-initiating cells. *Journal of Molecular Cell Biology, 9*, 302-314. https://doi.org/10.1093/jmcb/mjx017

Lueschow, S. R., & McElroy, S. J. 2020. The Paneth Cell: The Curator and Defender of the Immature Small Intestine. *Front Immunol, 11*, 587. https://doi.org/10.3389/fimmu.2020.00587

Luis, Tiago C., Naber, Brigitta A. E., Roozen, Paul P. C., Brugman, Martijn H., de Haas,

Edwin F. E., Ghazvini, M., Fibbe, Willem E., van Dongen, Jacques J. M., Fodde, R., & Staal, Frank J. T. 2011. Canonical Wnt Signaling Regulates Hematopoiesis in a Dosage-Dependent Fashion. *Cell Stem Cell, 9*, 345-356. https://doi.org/10.1016/j.stem.2011.07.017

Mah, A. T., Yan, K. S., & Kuo, C. J. 2016. Wnt pathway regulation of intestinal stem cells. *J Physiol, 594*, 4837-4847. https://doi.org/10.1113/JP271754

Metcalfe, C., Kljavin, N. M., Ybarra, R., & de Sauvage, F. J. 2014. Lgr5+ stem cells are indispensable for radiation-induced intestinal regeneration. *Cell Stem Cell, 14*, 149-159. https://doi.org/10.1016/j.stem.2013.11.008

Nakamura, T., Tsuchiya, K., & Watanabe, M. 2007. Crosstalk between Wnt and Notch signaling in intestinal epithelial cell fate decision. *J Gastroenterol, 42*, 705-710. https://doi.org/10.1007/s00535-007-2087-z

Nalapareddy, K., Nattamai, K. J., Kumar, R. S., Karns, R., Wikenheiser-Brokamp, K. A., Sampson, L. L., Mahe, M. M., Sundaram, N., Yacyshyn, M.-B., Yacyshyn, B., Helmrath, M. A., Zheng, Y., & Geiger, H. 2017. Canonical Wnt Signaling Ameliorates Aging of Intestinal Stem Cells. *Cell Reports, 18*, 2608-2621. https://doi.org/10.1016/j.celrep.2017.02.056

Noah, T. K., Donahue, B., & Shroyer, N. F. 2011. Intestinal development and differentiation. *Exp Cell Res, 317*, 2702-2710. https://doi.org/10.1016/j.yexcr.2011.09.006

Nusse, R., & Clevers, H. 2017. Wnt/beta-Catenin Signaling, Disease, and Emerging Therapeutic Modalities. *Cell, 169*, 985-999. https://doi.org/10.1016/j.cell.2017.05.016

Orouji, E., Raman, A. T., Singh, A. K., Sorokin, A., Arslan, E., Ghosh, A. K., Schulz, J., Terranova, C., Jiang, S., Tang, M., Maitituoheti, M., Callahan, S. C., Barrodia, P., Tomczak, K., Jiang, Y., Jiang, Z., Davis, J. S., Ghosh, S., Lee, H. M., Reyes-Uribe, L., … Rai, K. 2022. Chromatin state dynamics confers specific therapeutic strategies in enhancer subtypes of colorectal cancer. *Gut, 71(5)*, 938–949. https://doi.org/ https://doi.org/10.1136/gutjnl-2020-322835

Pachano, T., Haro, E., & Rada-Iglesias, A. 2022. Enhancer-gene specificity in development and disease. *Development, 149*. https://doi.org/10.1242/dev.186536

Pentinmikko, N., Iqbal, S., Mana, M., Andersson, S., Cognetta, A. B., Suciu, R. M., Roper, J., Luopajärvi, K., Markelin, E., Gopalakrishnan, S., Smolander, O.-P., Naranjo, S., Saarinen, T., Juuti, A., Pietiläinen, K., Auvinen, P., Ristimäki, A., Gupta, N., Tammela, T., Jacks, T., Sabatini, D. M., Cravatt, B. F., Yilmaz, Ö. H., & Katajisto, P. 2019. Notum produced by Paneth cells attenuates regeneration of aged intestinal epithelium. *Nature, 571*, 398-402. https://doi.org/10.1038/s41586-019-1383-0

Pinto, D., & Clevers, H. 2005. Wnt control of stem cells and differentiation in the intestinal epithelium. *Exp Cell Res, 306*, 357-363. https://doi.org/10.1016/j.yexcr.2005.02.022

Pinto, D., Gregorieff, A., Begthel, H., & Clevers, H. 2003. Canonical Wnt signals are essential for homeostasis of the intestinal epithelium. *Genes Dev, 17*, 1709-1713. https://doi.org/10.1101/gad.267103

Ramadan, R., van Driel, M. S., Vermeulen, L., & van Neerven, S. M. 2022. Intestinal stem

cell dynamics in homeostasis and cancer. *Trends Cancer, 8*, 416-425. https://doi.org/10.1016/j.trecan.2022.01.011

Rim, E. Y., Clevers, H., & Nusse, R. 2022. The Wnt Pathway: From Signaling Mechanisms to Synthetic Modulators. *Annu Rev Biochem, 91*, 571-598. https://doi.org/10.1146/annurev-biochem-040320-103615

Sato, T., van Es, J. H., Snippert, H. J., Stange, D. E., Vries, R. G., van den Born, M., Barker, N., Shroyer, N. F., van de Wetering, M., & Clevers, H. 2011. Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature, 469*, 415-418. https://doi.org/10.1038/nature09637

Schneider, C., O'Leary, C. E., & Locksley, R. M. 2019. Regulation of immune responses by tuft cells. *Nature Reviews Immunology, 19*, 584-593. https://doi.org/10.1038/s41577-019-0176-x

Sengupta, S., & George, R. E. 2017. Super-Enhancer-Driven Transcriptional Dependencies in Cancer. *Trends Cancer, 3*, 269-281. https://doi.org/10.1016/j.trecan.2017.03.006

Shoshkes-Carmel, M., Wang, Y. J., Wangensteen, K. J., Tóth, B., Kondo, A., Massasa, E. E., Itzkovitz, S., & Kaestner, K. H. 2018. Subepithelial telocytes are an important source of Wnts that supports intestinal crypts. *Nature, 557*, 242-246. https://doi.org/10.1038/s41586-018-0084-4

Silvia Maretto, M. C., Sirio Dupont, Paola Braghetta, Vania Broccoli, A. Bassim Hassan, Dino Volpin, Giorgio M. Bressan, and Stefano Piccolo. 2003. Mapping Wnt/β-catenin signaling during mouse development and in colorectal tumors. *PNAS*.

Snoeck, V., Goddeeris, B., & Cox, E. 2005. The role of enterocytes in the intestinal barrier function and antigen uptake. *Microbes and Infection, 7*, 997-1004. https://doi.org/10.1016/j.micinf.2005.04.003

Su, L. K., Kinzler, K. W., Vogelstein, B., Preisinger, A. C., Moser, A. R., Luongo, C., Gould, K. A., & Dove, W. F. 1992. Multiple intestinal neoplasia caused by a mutation in the murine homolog of the APC gene. *Science, 256*, 668-670. https://doi.org/10.1126/science.1350108

Terrin, L., Agostini, M., Ruvoletto, M., Martini, A., Pucciarelli, S., Bedin, C., Nitti, D., & Pontisso, P. 2017. SerpinB3 upregulates the Cyclooxygenase-2 / beta-Catenin positive loop in colorectal cancer. *Oncotarget, 8*, 15732-15743. https://doi.org/10.18632/oncotarget.14997

Tian, H., Biehs, B., Warming, S., Leong, K. G., Rangell, L., Klein, O. D., & de Sauvage, F. J. 2011. A reserve stem cell population in small intestine renders Lgr5-positive cells dispensable. *Nature, 478*, 255-259. https://doi.org/10.1038/nature10408

van der Flier, L. G., & Clevers, H. 2009. Stem cells, self-renewal, and differentiation in the intestinal epithelium. *Annu Rev Physiol, 71*, 241-260. https://doi.org/10.1146/annurev.physiol.010908.163145

van Es, J. H., Jay, P., Gregorieff, A., van Gijn, M. E., Jonkheer, S., Hatzis, P., Thiele, A., van den Born, M., Begthel, H., Brabletz, T., Taketo, M. M., & Clevers, H. 2005. Wnt signalling induces maturation of Paneth cells in intestinal crypts. *Nat Cell Biol, 7*, 381-386. https://doi.org/10.1038/ncb1240

van Es, J. H., Sato, T., van de Wetering, M., Lyubimova, A., Yee Nee, A. N., Gregorieff, A.,

Sasaki, N., Zeinstra, L., van den Born, M., Korving, J., Martens, A. C. M., Barker, N., van Oudenaarden, A., & Clevers, H. 2012. Dll1+ secretory progenitor cells revert to stem cells upon crypt damage. *Nat Cell Biol, 14*, 1099-1104. https://doi.org/10.1038/ncb2581

Wang, A., Wang, J., Tian, K., Huo, D., Ye, H., Li, S., Zhao, C., Zhang, B., Zheng, Y., Xu, L., Hua, X., Wang, K., Wu, Q. F., Wu, X., Zeng, T., Liu, Y., & Zhou, Y. 2021. An epigenetic circuit controls neurogenic programs during neocortex development. *Development, 148*. https://doi.org/10.1242/dev.199772

Wang, J., Wang, A., Tian, K., Hua, X., Zhang, B., Zheng, Y., Kong, X., Li, W., Xu, L., Wang, J., Li, Z., Liu, Y., & Zhou, Y. 2022. A Ctnnb1 enhancer regulates neocortical neurogenesis by controlling the abundance of intermediate progenitors. *Cell Discov, 8*, 74. https://doi.org/10.1038/s41421-022-00421-2

Wong, E. S., Zheng, D., Tan, S. Z., Bower, N. L., Garside, V., Vanwalleghem, G., Gaiti, F., Scott, E., Hogan, B. M., Kikuchi, K., McGlinn, E., Francois, M., & Degnan, B. M. 2020. Deep conservation of the enhancer regulatory code in animals. *Science, 370*. https://doi.org/10.1126/science.aax8137

Xu, L., Zheng, Y., Li, X., Wang, A., Huo, D., Li, Q., Wang, S., Luo, Z., Liu, Y., Xu, F., Wu, X., Wu, M., & Zhou, Y. 2021. Abnormal neocortex arealization and Sotos-like syndrome–associated behavior in Setd2 mutant mice. *Science Advances, 7*, 1180. https://doi.org/10.1126/sciadv.aba1180

Yan, K. S., Chia, L. A., Li, X., Ootani, A., Su, J., Lee, J. Y., Su, N., Luo, Y., Heilshorn, S. C., Amieva, M. R., Sangiorgi, E., Capecchi, M. R., & Kuo, C. J. 2012. The intestinal stem cell markers Bmi1 and Lgr5 identify two functionally distinct populations. *Proc Natl Acad Sci U S A, 109*, 466-471. https://doi.org/10.1073/pnas.1118857109

Zabidi, M. A., & Stark, A. 2016. Regulatory Enhancer-Core-Promoter Communication via Transcription Factors and Cofactors. *Trends Genet, 32*, 801-814. https://doi.org/10.1016/j.tig.2016.10.003

Zhang, Y., & Wang, X. 2020. Targeting the Wnt/beta-catenin signaling pathway in cancer. *J Hematol Oncol, 13*, 165. https://doi.org/10.1186/s13045-020-00990-3

Zhao, H., Ming, T., Tang, S., Ren, S., Yang, H., Liu, M., Tao, Q., & Xu, H. 2022. Wnt signaling in colorectal cancer: pathogenic role and therapeutic target. *Mol Cancer, 21*, 144. https://doi.org/10.1186/s12943-022-01616-7

Zheng, G., Pang, S., Wang, J., Wang, F., Wang, Q., Yang, L., Ji, M., Xie, D., Zhu, S., Chen, Y., Zhou, Y., Higgins, G. A., Wiley, J. W., Hou, X., & Lin, R. 2023. Glucocorticoid receptor-mediated Nr1d1 chromatin circadian misalignment in stress-induced irritable bowel syndrome. *iScience, 26*, 107137. https://doi.org/10.1016/j.isci.2023.107137

Zhu, Y., & Li, X. 2023. Advances of Wnt Signalling Pathway in Colorectal Cancer. *Cells, 12*. https://doi.org/10.3390/cells12030447

Acknowledgments

We thank Dr. Yeguang Chen for providing the *Lgr5*-EGFP-IRES-CreERT2 mice.

We thank the Core Facility and the Animal Facility of Medical Research Institute of Wuhan University for technical support. We thank all Zhou lab members for critical reading of the manuscript. Y. Zhou was supported by grants from National Key R&D Program of China (2022YFA0806603 and 2018YFA0800700), National Natural Science Foundation of China (32270876 and 31970770), and the Fundamental Research Funds for the Central Universities (2042022dx0003 and 2042023kf0234). Y.L. was supported by grants from National Key R&D Program of China (2018YFA0800700 and 2022YFA0806603), National Natural Science Foundation of China (31970676), and Hubei Natural Science Foundation (2022CFB128).

Author contributions

Y.Z. conceived and designed the study; X.H. and J.W. performed experiments and analyzed data; C.Z. assisted with processing ChIP-seq data; J.T. and X.P. retrieved and analyzed eQTL data. M.Y. and M.W. provided RNA-seq and ChIP-seq data of colorectal cancer samples; G.Z. provided Hi-C data; Y.Z. and X.H. wrote the paper. Y.Z. and Y.L. provided funding and supervised the study. All authors discussed results and commented to the manuscript.

Competing interests

The authors claimed that they have no competing interests.

Figures

Figure 1

Figure 1. ieCtnnb1 is an intestinal enhancer of *Ctnnb1*. (A) Schematic

representation of the upstream region of mouse *Ctnnb1* gene and the location

of ieCtnnb1 (6,399 bp, pink shading), which is marked by H3K27ac and

H3K4me1 peaks, and DNase I hypersensitivity in small intestine and large intestine of 8-week-old mice. Data were obtained from ENCODE. Locations of single-guide RNAs (sgRNAs) for generating ieCtnnb1 knockout mice were marked. (B) Top left: a schematic illustration showing that the knock-in reporter construct carries the *Shh* promoter, ieCtnnb1 core region sequences (2,153 bp) and the *LacZ* reporter gene. Top right: X-Gal staining (blue) of the gastrointestinal (GI) tract of an 8-week-old *H11$^{i.enh}$* mouse. Bottom: X-Gal staining (blue) of the small intestine (left) and colon (right) of an 8-week-old *H11$^{i.enh}$* mouse. Boxed areas were enlarged at top-right corners. (C) Representative images of small intestinal crypts co-labelled by X-gal with OLFM4 (left), and X-gal with Lysozyme (right). (D-E) Representative images of whole body (d) and GIs (e) of 8-week-old male wildtype (WT) and *Ctnnb1$^{Δi.enh}$* mice. (F) Comparison of the body weight of 8-week-old male WT (n = 13) and *Ctnnb1$^{Δi.enh}$* (n = 13) mice. (G-H) Measurements of small (G) and large (H) intestine length of 8-week-old male WT (n = 6) and *Ctnnb1$^{Δi.enh}$* (n = 6) mice. (I-J) Relative mRNA levels of *Ctnnb1* in small (I) and large (J) intestinal crypts of WT (n = 6) and *Ctnnb1$^{Δi.enh}$* (n = 6) mice. (K-L) Left: immunoblotting of nuclear (K) and cytoplasmic (L) β-Catenin, GAPDH and H3 of small intestinal crypts of WT (n = 3) and *Ctnnb1$^{Δi.enh}$* (n = 3) mice. Right: histograms showing protein levels of β-Catenin normalized to H3 (K) or GAPDH (L) levels. Values of WT were set as '1'. (M) Heat map of indicated Wnt target genes and GSEA analysis of Wnt signaling pathway according to transcriptome profiles of small intestinal crypts of WT (n = 3) and *Ctnnb1$^{Δi.enh}$* (n = 3) mice. (N) RT-qPCR showing relative mRNA levels of indicated Wnt target genes (*Axin2, Lgr5* and *Mmp7*) in small intestinal crypts of WT (n = 3) and *Ctnnb1$^{Δi.enh}$* (n = 3) mice. Scale bars, 1 cm (B, top; D and E), 100 μm (B, bottom), 10 μm (B, magnified views; C). Quantification data are shown as means ± SEM, statistical significance was determined using an unpaired two-tailed Student's *t*-test (F-L). Quantification data are shown means ± SD, statistical significance was determined using

1159   Multiple t-tests - one per row (N). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P$

1160 < 0.0001. ns, not significant. NES: Normalized Enrichment Score.

Figure 2

Figure 2. ieCtnnb1 knockout altered cellular composition and expression

profiles of small intestinal crypts. (A) Schematic illustration of single-cell

sequencing. Crypts were extracted from small intestines followed by

fluorescence activated cell sorting (FACS) to enrich EpCAM+DAPI- epithelial

cells. Cells of two 10-week-old female mice for each genotype were pooled

together to perform single cell transcriptome analyses. (B) UMAP were used to

visualize the clustering of 11,824 single cells from $H11^{i.enh}$ mice and 8,094

single cells from $Ctnnb1^{\Delta i.enh}$ mice. Cell types were assigned according to

expressions of marker genes. CBC, crypt base columnar cell; TAC, transit

amplifying cell; EEC, enteroendocrine cell; imEC, immature enterocytes; mEC,

mature enterocytes; GC, goblet cell; PC, Paneth cell; TC, tuft cell. (C)

Expression and distribution of $Ctnnb1$ in small intestinal crypt cells of $H11^{i.enh}$

and $Ctnnb1^{\Delta i.enh}$ mice. (D) Violin plots showing the expression of $Ctnnb1$ in CBC,

TAC, PC, GC, imEC, mEC; and the expression of $Axin2$ in CBC and TAC, of

$H11^{i.enh}$ and $Ctnnb1^{\Delta i.enh}$ mice. (E) Comparison of the proportion of indicated small

intestinal crypt cell types in $H11^{i.enh}$ and $Ctnnb1^{\Delta i.enh}$ mice. (F)

Immunohistochemistry (left and middle) and quantification (right) of PCs in

small intestines of $H11^{i.enh}$ (n = 6) and $Ctnnb1^{\Delta i.enh}$ (n = 6) mice. (G) RT-qPCR

showing relative mRNA levels of PC marker genes ($Lyz1, Pla2g2a, Wnt3,$

$Math1$) in small intestinal crypts of $H11^{i.enh}$ (n = 3) and $Ctnnb1^{\Delta i.enh}$ (n = 3) mice.

(H) Distribution of G2M cells in $H11^{i.enh}$ and $Ctnnb1^{\Delta i.enh}$ small intestinal crypts,

based on the expression of cell cycle marker gene $Mki67$. (I) Comparison of the

proportion of G2M cells in CBC and TAC of $H11^{i.enh}$ and $Ctnnb1^{\Delta i.enh}$ small

intestinal crypts. (J) Violin plots showing the expression of $Mki67$ in CBC and

TAC of $H11^{i.enh}$ and $Ctnnb1^{\Delta i.enh}$ small intestinal crypts. (K)

Immunohistochemistry (left and middle) and quantification (right) of Ki67+ cells

in small intestines of $H11^{i.enh}$ (n = 6) and $Ctnnb1^{\Delta i.enh}$ (n = 6) mice. (L)

Immunofluorescence (left and middle) and quantification (right) of EdU+ cells

(red) in small intestines of $H11^{i.enh}$ (n = 6) and $Ctnnb1^{\Delta i.enh}$ (n = 6) mice after 4

hours EdU injection. Nuclei were labeled with DAPI (blue). (M-N) Violin plots

showing expressions of marker genes for secretory cells ($Lyz1, Ang4, Muc2,$

$Mmp7$) and absorptive cells ($Fabp6, Apoa1, Fabp2, Reg3b$) in secretory(M) and

absorptive lineages (N) of $H11^{i.enh}$ and $Ctnnb1^{\Delta i.enh}$ small intestinal crypts. Scale

bars, 50 µm (F, K, and L). Quantification data are shown as means ± SEM,

statistical significance was determined using an unpaired two-tailed Student's

t-test (D, F, J, K, L, M and N). Quantification data are shown means ± SD,

statistical significance was determined using Multiple *t*-tests - one per row (G).

*P < 0.05, **P < 0.01, ***P < 0.001, and ****P < 0.0001. ns, not significant.

1200

Figure 3

Figure 3. Knocking out ieCtnnb1 inhibits tumorigenesis of colorectal cancer. (A) Survival of *Apc*^{Min/+} (n = 15) and *Ctnnb1*^{Δi.enh};*Apc*^{Min/+} (n =17) mice. (B) Colon images of 5-month-old *Apc*^{Min/+} (n =9) and *Ctnnb1*^{Δi.enh};*Apc*^{Min/+} (n =9) mice. (C) Weight statistics of 5-month-old *Apc*^{Min/+} (n = 20) and *Ctnnb1*^{Δi.enh};*Apc*^{Min/+} (n = 20) mice. (D) The statistical analysis of tumor counts in colons of 5-month-old *Apc*^{Min/+} (n = 9) and *Ctnnb1*^{Δi.enh};*Apc*^{Min/+} (n = 9) mice.

(E) Representative H&E staining images of colon sections of 5-month-old

$Apc^{Min/+}$ and $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ mice. (F) The statistical analysis of colon

tumor area in 5-month-old $Apc^{Min/+}$ (n = 6) and $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ (n = 6) mice.

(G-H) Immunohistochemistry (G) and quantification (H) of Ki67+ cells in colon

tumors of 5-month-old $Apc^{Min/+}$ (n = 6) and $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ (n = 6) mice. (I-

J) Immunohistochemistry (I) and signal intensity statistics (J, red dashed boxes

of I) of β-Catenin in colon tumors of 5-month-old $Apc^{Min/+}$ (n = 6) and

$Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ (n = 6) mice. (K-L) Immunohistochemistry (K) and

quantification (L) of MUC2+ cells in colon tumors of 5-month-old $Apc^{Min/+}$ (n = 6)

and $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ (n = 6) mice. (M) The heat map showing relative

expressions of Wnt signaling pathway genes of colon tumors from 5-month-old

$Apc^{Min/+}$ (n = 3) and $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ (n = 3) mice. (N-O) KEGG analyses of

downregulated (N) and upregulated (O) genes in colon tumors of 5-month-old

$Apc^{Min/+}$ (n = 3) and $Ctnnb1^{\Delta i.enh};Apc^{Min/+}$ (n = 3) mice. Scale bars, 1 cm (B), 4

mm (E, G, I and K), 200 μm (magnified views in G and K), 100 μm (magnified

views in I). Quantification data are shown as means ± SEM, statistical

significance was determined using an unpaired two-tailed Student's t-test (C,

D, F, H, J and L) or log-rank analysis (A). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$,

and $****P < 0.0001$. ns, not significant.

1227

Figure 4

Figure 4. ieCTNNB1 is the intestinal enhancer of human *CTNNB1*. (A) Schematic representation of human *CTNNB1* gene and the location of ieCTNNB1 (3,000 bp, pink shading), which is marked by H3K27ac and H3K4me1 peaks, and DNase I hypersensitivity in human small intestine (30-year-old female) and colon (34-year-old male). Data were obtained from ENCODE. (B) Top: a schematic illustration showing that the knock-in construct containing the *Shh* promoter, ieCTNNB1 sequences (3,000 bp), and the LacZ reporter gene. Bottom: X-Gal staining (blue) of the gastrointestinal tract, and sections of the proximal small intestine, distal small intestine, and large intestine in 8-week-old $H11^{hi.enh}$ mice. (C) Left: ieCTNNB1 is marked by enrichment of H3K27ac in human small intestine (30-year-old female) and colon (34-year-old male). Data were obtained from ENCODE. Locations of sgRNA target sites were indicated. Five subregions of ieCTNNB1 were shown. Right: luciferase reporter assay in HCT116 cells transfected with indicated plasmids for 48 hours. (D) RT-qPCR showing relative mRNA levels of *CTNNB1* in HCT-15 cells transfected with indicated CRISPR activation or CRISPR interference vectors

1245   for 48 hours. Scale bars, 1 cm (whole mount in B), 100 μm (sections in B), 10

1246   μm (magnified views in B). Quantification data are shown as means ± SEM,

1247   statistical significance was determined using one-way ANOVA analysis (C) and

1248   an unpaired two-tailed Student's *t*-test (D). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$,

1249   and ****$P < 0.0001$. ns, not significant.

1250

Figure 5

Figure 5. ieCTNNB1 is activated in colorectal cancer and its activity positively correlates with the expression of *CTNNB1*. (A) Schematic representation of ieCTNNB1 (pink shading) and *CTNNB1* promoter (yellow shading), which is respectively marked by H3K27ac and H3K4me3 peaks, and mRNA signals in native and tumor tissues of a patient with colorectal cancer. The location of risk mutation site was indicated. (B) Comparison of *CTNNB1* expression levels in native and tumor tissues of colorectal cancer patients (n = 68). (C) Left: comparison of H3K27ac signals at ieCTNNB1 in native and tumor tissues of colorectal cancer patients (n = 64). Right: comparison of H3K4me3 signals at *CTNNB1* promoter in native and tumor tissues of colorectal cancer patients (n = 42). (D) Left: correlation between H3K27ac signals at ieCTNNB1 and *CTNNB1* expression in native and tumor tissues of colorectal cancer patients (n = 55). Right: correlation between H3K4me3 signals at *CTNNB1* promoter and *CTNNB1* expression in native and tumor tissues of colorectal cancer patients (n = 38). (E) Left: comparison of *CTNNB1* expression in esophagus between subjects with common sequence (C/C, n = 428) and variant sequence (C/T, n = 37). Right: comparison of *CTNNB1* expression in transverse colon between subjects with common sequence (C/C, n = 343) and

variant sequence (C/T, n = 25). (F) Luciferase reporter assay in HCT116 and

HeLa cells transfected with indicated plasmids for 48 hours. Quantification data

are shown as means ± SEM, statistical significance was determined using a

paired (B, C and D) or unpaired (E) two-tailed Student's t-test and Two-way

ANOVA analysis (F). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$.

ns, not significant. R: Pearson correlation.

Figure 6

Figure 6. HNF4α and p-S133-CREB1 associate with ieCTNNB1 to regulate

***CTNNB1*'s transcription.** (A) ChIP-seq tracks of indicated trans-acting factors

enriched at ieCTNNB1 (pink shading) in indicated colorectal cancer cell lines.

(B) RT-qPCR showing relative mRNA levels of *CTNNB1* in HCT-15 cells transfected with indicated shRNA-expressing plasmids for 48 hours. The expression level of *CTNNB1* in cells transfected with scramble (scr) shRNA was set to '1'. (C-D) Top: schematic diagram showing the enrichment of H3K27ac and p-S133-CREB1 at ieCTNNB1 (C) and *CTNNB1*'s promoter (D). Locations of HNF4α and CREB1 binding motif sites were indicated. Middle and bottom: ChIP-qPCR showing enrichment of HNF4α (middle) and p-S133-CREB1 (bottom) at ieCTNNB1 (C) and *CTNNB1*'s promoter (D) in HCT-15 cells. Locations of ChIP regions were indicated. (E) Comparison of expression levels of *CREB1* and *HNF4A* between native and tumor tissues in colon adenocarcinoma (COAD) and rectum adenocarcinoma (READ) tumors. (F) Correlations between the expression level of *CTNNB1* and those of *CREB1* or *HNF4A* in COAD tumors. (G-H) ChIP-qPCR showing enrichment of HNF4α (G) and p-S133-CREB1 (H) at *Ctnnb1* promoter in native colon tissues of WT (n=3) mice, tumor tissues of *Apc*^Min/+ (n=3) mice and *Ctnnb1*$^{\Delta i.enh}$;*Apc*^Min/+ (n=3) mice. Quantification data are shown as means ± SEM, statistical significance was determined using one-way ANOVA analysis (B), unpaired two-tailed Student's *t*-test (E and F) and Multiple *t*-tests - one per row (C, D, G and H). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$. ns, not significant. R: Pearson correlation.