Impact of Run-of-River Damming on Increasing Phytoplankton Biomass and Species Shift in a Large Amazonian River

Alfonso Pineda[1], Beatriz Melissa Campos[2], Felipe Zanon[2], Renata Felicio[2], Luzia C. Rodrigues[2,3]

[1] Programa de Pós-graduação em Biologia Comparada – PGB, Universidade Estadual de Maringá – UEM, Maringá, PR, Brazil

[2] Programa de Pós-graduação em Ecologia de Ambientes Aquáticos Continentais – PEA, Centro de Ciências Biológicas (CCB), Universidade Estadual de Maringá – UEM, Maringá, PR, Brazil

[3] Núcleo de Pesquisas em Limnologia e Aquicultura – Nupélia, Centro de Ciências Biológicas (CCB), Universidade Estadual de Maringá – UEM, Maringá, PR, Brazil

Correspondence: Alfonso Pineda, Programa de Pós-graduação em Ecologia de Ambientes Aquáticos Continentais, Universidade Estadual de Maringá. Avenida Colombo, 5790, Bloco H-90, Maringá, Paraná 87020-900, Brazil Amanecce@gmail.com

ORCID Ids

Pineda, A amanecce@gmail.com https://orcid.org/0000-0001-6902-9993 ;
Campos, BM https://orcid.org/0000-0003-1985-8528
Zanon, FM https://orcid.org/0000-0002-8098-2803
Felicio, RS https://orcid.org/0000-0001-7716-8408
Rodrigues, LC https://orcid.org/0000-0002-0593-0931

Running Head: algal biomass after damming an Amazon tributary

Abstract

Run-of-river (ROR) dams, often perceived as having minimal environmental impact, can induce significant hydrodynamic changes that alter aquatic ecosystems. We investigated the impacts of an ROR dam on the Madeira River, the largest Amazon tributary, focusing on phytoplankton communities, their ecological implications, and related environmental factors. Our study examined changes in biomass and environmental factors (using General Linear Mixed Models - GLMM), species composition (using PERMANOVA) before and after damming, in both the main channel and tributaries (N=549 samples). We also identified indicator species associated with different damming phases and regions through an indicator value analysis. The results showed that, following dam construction, the phytoplankton community changed in both the main channel and tributaries, with a shift from lotic diatoms to lentic phytoflagellates. This transition was likely facilitated by altered hydrodynamics and possibly influenced by the decomposition of flooded vegetation in the dam's influence zone. The decomposition of this vegetation could explain both the observed increase in oxygen consumption and the subsequent rise in phytoflagellate biomass after damming. However, despite the overall increase in phytoplankton biomass, the values remained within oligotrophic to mesotrophic conditions, consistent with the low nutrient concentrations recorded. However, we caution that the dam-created hydrodynamic conditions are optimal for phytoplankton growth, potentially exacerbating nutrient-related issues in the future. We recommend proactive management strategies to prevent nutrient enrichment from activities such as agriculture and livestock in isolated Amazon areas affected by dams, thereby mitigating potential water quality degradation linked to increased phytoplankton biomass.

Keywords: Amazon; backwaters; cyanobacteria; damming; plankton; run-of-river dam

1 Introduction

Hydroelectric energy is considered the most environmentally friendly option for meeting global energy demand. Hydroelectric projects have been constructed even in important ecological regions. For instance, in the Amazon region, the largest basin in the world, 140 dams are functioning or under construction, and 288 are planned (Latrubesse et al., 2017). Most of the hydroelectric projects built nowadays are run-of-river (ROR) structures, which in theory flood smaller areas than traditional storage reservoirs (Dias et al., 2018; Winemiller et al., 2016). However, even hydroelectric projects with a ROR dam can impact environmental conditions (Almeida et al., 2019; Moran, 2020; Tundisi, 2018), and biodiversity (Lees et al., 2016; Vasconcelos et al., 2022).

Run-of-river dams can have little effect on the main channel where they are built (Almeida et al., 2019; Zanon et al., 2024). However, they relate to the apparition of lentic or semi-lentic conditions in the back-flooded tributaries (Almeida et al., 2019). In some cases, the back-flooding area can correspond to 30% of the reservoir area (Almeida et al., 2019), and the total flooding area can be 64.5% larger than initially predicted by hydroelectric project including a ROR dam (Cochrane et al., 2017). These new lake-like conditions can harbor problematic species, such as mosquitoes which are vectors of diseases (Rodrigues et al., 2017), aquatic macrophytes that cover the water surface (Thomaz et al., 1999), and invasive species that exclude native ones (Lees et al., 2016).

The change from running water to standing water can also favor algae, as a longer water residence time enhances their reproduction. Standing water conditions together with high nutrient concentrations can trigger populational explosions of algae known as blooms (O'Neil et al., 2012). In turn, high algae biomass can threaten water quality (Ferrão-Filho and Kozlowsky-Suzuki, 2011), diversity (Krztoń et al., 2019), and human health (Pouria et al., 1998) since it decreases the transparency and oxygen of the water and even increases the concentration of toxins produced by some microalgae. Moreover, high algae biomass can degrade the services that we take advantage of from aquatic ecosystems such as recreation and fishing (Huisman et al., 2018).

The formation of standing waters in the Amazonian region could not trigger algae blooms *per se*, but together with the high temperature of the region, high nutrient input, and the hydrodynamics changes promoted by the dams may favor phytoplankton growth

(Rodrigues et al., 2018; Thornton, 1990). For instance, human activities such as agriculture
and cattle ranching in the Amazon increase phosphorous and nitrogen load in aquatic
ecosystems (Tilman et al., 2002; Vale et al., 2019). This is particularly important given the
increase in this type of enterprise in the Amazon in recent years (Carvalho et al., 2020; Gusso
et al., 2017; Macedo et al., 2012; Vale et al., 2019).

Several studies have described the response of phytoplankton communities to
damming in several regions worldwide (e.g., Okuku et al., 2016; Souza et al., 2016; Xiao et
al., 2016; Znachor et al., 2020). However, it is no clear how the recently built and futures
dams affects the spatial and temporal variation of phytoplankton communities in critical
ecological regions such as the Amazon. Moreover, there exists a knowledge gap regarding
the effect of damming on microalgae biomass of Amazonian Rivers. It is now crucial to
determine whether the environmental changes induced by hydroelectric projects in the
Amazon region are sufficient to promote the development of phytoplankton biomass and
identify the favored species. This information is essential for alerting authorities to potential
negative effects and for elucidating the appropriate management measures necessary to
prevent or mitigate impacts associated with high phytoplankton biomass or unexpected
species.

In this study, we assessed the influence of the Jirau dam on upstream biomass and
community structure (species abundance) of phytoplankton in the Madeira River, one of the
largest tributaries of the Amazon River. We collected data in the main channel and tributaries
before and after the installation of the dam. Our objectives were to determine the effects of
damming in phytoplankton biomass (structure and diversity) and identify which abiotic
effects related to damming could explain eventual changes.

Since backwaters formed in tributaries after damming can exhibit lake-like
conditions, we hypothesized that *i*) post-damming conditions in tributaries would favor an
increase in phytoplankton biomass. Therefore, we expected an increase in biomass in
tributaries, while no significant changes were anticipated in the main channel. Additionally,
due to the shift in environmental conditions caused by damming, we hypothesized that *ii*)
damming would alter the community structure (species abundance). Specifically, we
anticipated differences in community structure before and after the damming, with species
adapted to standing water conditions becoming representative after damming. Finally, we

examined the significance of our findings in the context of environmental degradation
resulting from human activities in the Amazon.

2   Methods

2.1   Study area

The Madeira River basin is in the Amazon region, between Peru, Bolivia, and Brazil,
draining much of that region (Guyot et al., 1996). The Madeira River is among the ten rivers
with the highest discharge in the world (Latrubesse et al., 2017)), it is the main tributary of
the Amazon River and contributes significantly to the phosphorus load for the entire Amazon
basin (Almeida et al., 2015). The Madeira River basin has one annual cycle with four well-
defined hydrological periods, with well-established duration intervals between years: 1. The
high-water period, which usually occurs from February to May; 2. The receding period,
occurring from June to July; 3. low water, from August to November; and 4. rising period,
occurring from December to January. This region has the largest tropical forest in the world,
being responsible for much of the global biodiversity in addition to the importance of
temperature and precipitation in South and Central America (Agudelo et al., 2019; Aragón,
2018).

The Madeira River has two dams installed, Santo Antônio and Jirau, but other dams
are in the planning and approval phases (Latrubesse et al., 2017). The construction of the
Jirau dam ended in July 2012, being considered a run-of-river, located in the city of Porto
Velho, Rondônia state, Brazil. The Jirau reservoir has an annual average water residence time
of approximately 3.0 days (Almeida et al., 2020). It is located 1,206 km from the mouth of
the Madeira River and has an installed capacity of 3.750 MW, supplying more than ten
million homes (Energia Sustentável do Brasil, 2020). In the study period, 2009 to 2017, the
average flow was 19.409 m³/s, reaching a maximum average of 45.649 m³/s in the high-water
period and a minimum average of 4.133 m³/s in the low-water period (ANA, 2022). For the
study, we cover the area upstream of the Jirau hydroelectric plant, between the coordinates
9º67'73.05" S, 65º44'10.27" O, and 9º60'45.27" S, 64º91'75.55" O (Figure 1).

Run-of-river dams have been reported to have minimal influence on river flow and
creating minor flooded regions (Pracheil et al., 2016; Souza et al., 2019). Nonetheless, the
water level of the Madeira River has changed significantly since the Jirau dam was built. The
water level ranged between 1 m and 20 m before damming, whereas it varied between 21 m
and 29 m after damming. Also, the seasonal fluctuation in the Madeira River's water level has

decreased after damming (Supplementary material S1). The Madeira River's rising water

level has reduced flow velocity and induced stratification in its tributaries (Fearnside, 2013).

The reservoir's extension varies between 207 km^2 during the dry season and 361.6 km^2 during

the wet season (Hartmann et al., 2023).

2.2   Data Samplings

Fifteen sampling stations were set up along the Madeira River (M2 to M5) and its

tributaries (T4 to T14 – Figure 1). Water samples for phytoplankton and nutrient

concentration assessments were collected directly using bottles at a depth of approximately

40 cm in the middle section of the Madeira Rivers and tributaries. At the same time, we

measured chemical and physical variables that are critical to phytoplankton growth. Between

2009 and 2017, samples were taken during the high (March to May), rising (December to

February), receding (June to August), and low (September to November) seasons. In 2010,

2011, 2016, and 2017, the campaigns were carried out quarterly, while from 2013 to 2015

they were carried out bimonthly. Five samplings were conducted in 2012. In total, we had 37

sampling campaigns (N = 550). Five samples were lost (M4 in campaign sampling 2, M3 in

April 2010, T4 in April 2012, T10 in April 2013, and T5 in August 2014). The before-dam

period is considered up until July 2012 (177 samples), while the after-dam period begins in

October 2012 and continues through the remainder of the sampled periods (373 samples).

Figure 1. Map and location of sampling sites in the Jirau Hydroelectric Power Plant, in the
Madeira River.

2.3   Phytoplankton

The sampled individuals were identified to the lowest possible taxonomic level, using

specialized literature such as taxonomic articles, identification keys and books (Pineda et al.,

2020a; van den Hoek et al., 1995; Komarek and Anagnostidis 1989, 1998, 2005).

Phytoplankton samples were fixed with 1% acetic Lugol solution. Individuals (cell,

coenobium, colonies, and filaments) were counted using an inverted microscope at 400X as

described by Utermöhl (1958). Phytoplankton density was calculated according to APHA

(American Public Health Association, 2012) and expressed as individuals per milliliter

(ind.mL^{-1}). Biomass (expressed in mm^3.L-1) was obtained by multiplying the density of each

taxon by its corresponding volume. The volume of individual cells was calculated using

geometric models that were based on the shape of the cells (Hillebrand et al., 1999; Sun and

Liu, 2003). The classification systems proposed by Reviers (2003) and Komárek &

Anagnostidis (1998) were adopted for taxonomic identification at the level of class for

eukaryotic algae and Cyanobacteria. The entire number of registered taxa was divided into

five algal groups: green algae, cyanobacteria, diatoms, phytoflagellates, and xantophyceans.

These groups are the most representative in freshwater ecosystems, accounting for most of
the phytoplankton biomass. Additionally, they have shown sensitivity to environmental
changes, making them well-suited for understanding how human activities affect biodiversity
and water quality (Reynolds, 2006) (Reynolds, 2006). Although species from the
Xanthophyceae class were identified, they were not included in the data analysis due to their
low representativity, which made the analysis challenging.

2.4 Environmental data

We measured chemical and physical variables critical to the development of
phytoplankton following the standard methods outlined in APHA (American Public Health
Association, 2012).In situ measurements were conducted to assess water temperature (WT,
°C), pH, inorganic carbon (IC, mg L^{-1}), and alkalinity (Alk, mEq L^{-1}). Water transparency
was measured with a Secchi disk (m), and the euphotic zone (Zeu, m) was estimated as 2.7
times the Secchi depth (Cole, 1994). The maximum depth (Zmax, m) was also measured, and
the Zeu:Zmax ratio was utilized as an indicator of light availability within the water column
(Jensen et al., 1994).

Using collected water samples, we measured additional water characteristics at the
sampled sites. Total iron concentrations ($\mu g\ L^{-1}$) were determined following digestion using
the phenanthroline method. Soluble reactive phosphorus (SRP, $P-PO_4^{3-}$, $\mu g\ L^{-1}$), nitrate (N-
NO_3^-, $\mu g\ L^{-1}$), nitrite ($N-NO_2^-$, $\mu g\ L^{-1}$), and ammonium ($N-NH_4^+$, $\mu g\ L^{-1}$) concentrations were
measured post-sample filtration. Phosphate concentrations were determined using the
ascorbic acid method and spectrophotometric readings, while nitrate, nitrite, and ammonium
concentrations were obtained through the cadmium reduction method, sulfanilamide method,
and phenol method, respectively, for spectrophotometric measurements. Total dissolved
inorganic nitrogen (DIN, $\mu g\ L^{-1}$) was calculated as the sum of nitrite, nitrate, and ammonium
concentrations. Reactive silica ($\mu g\ L^{-1}$) concentrations were determined using the molybdate
method. Biochemical oxygen demand (BOD, mg L^{-1}) was assessed following sample
incubation at a constant temperature (20 °C) and reading the dissolved oxygen concentrations
after 5 days. Chemical oxygen demand (COD, mg L^{-1}) determination involved the oxidation
of organic matter using a boiling mixture of chromic acid and sulfuric acid (potassium
bichromate in an acidic medium)(APHA, 2005).

    2.5   Data analysis

We utilized Generalized Mixed Linear Models (GLMM) to investigate the effect of

damming on the environmental variables in both the main channel and tributaries. To avoid

issues with multiple pairwise tests, we included the sampling point as a random factor

(Bolker et al., 2009). We selected an error distribution based on the better fit of the models,

such as Gaussian, Beta, or Gamma.

To verify the effect of damming on the structure of the phytoplankton community, we

utilized a Multivariate Permutation Analysis of Variance (PERMANOVA) on a matrix of

species biomass (considering the Bray-Curtis distance). To determine whether damming

effect interacted with hydrological periods, we considered the seasonality in the

PERMANOVA. The analysis was performed separately for tributaries and the main channel.

A Principal Coordinate Analysis (PCoA) was used to graphically depict the spatial and

temporal ordinations. Additionally, to verify associations between species and damming

phases (before and after) and regions (tributaries and main channel), we used the indicator

value (IndVal) (Dufrêne and Legendre, 1997).We analyzed four groups that combined

damming phases and regions. IndVal is a quantitative metric that accounts for both the

relative abundance and frequency of a species within a specific condition and is not

influenced by the abundance of other species. The significance of all indicator values was

assessed using a Monte Carlo randomization process with 9999 permutations. A significant

IndVal suggests that the species is more characteristic or prevalent in that condition. An

IndVal value close to 1 indicates a strong association of the species with a particular

condition (e.g., damming phase or region). In our study, this approach complemented the

PERMANOVA analysis by allowing us to identify which species were more favored in each

condition. For details on IndVal formula and its applications, see Borcard et al. (2018).

We utilized GLMM to assess the influence of damming on the biomass of the entire

phytoplankton community and each of the five algal group in both the tributaries and the

main channel. When we found that damming had a considerable influence on algal biomass,

we examined into its link with seasonality. We also utilized GLMM to assess the influence of

each standardized abiotic factor on algal biomass as well as on the presence-absence of each

group. In this case, we modeled both biomass and the presence-absence of algal groups since

several samples lacked a specific algae group, resulting in biomass values of zero (absence).

We chose the Gamma family as the error distribution for modeling biomass and the zero-

inflated distribution for occurrence data. To avoid issues with multiple testing, we incorporated the sampling point as a random effect in all models, as advised by Bolker et al. (Bolker et al., 2009). Visual examination validated the linearity assumption. The models' findings were given in ANOVA style. To determine the trophic state before and after damming, the registered total biomass values were classified according to Vollenweider's trophic classification (Vollenweider, 1968): oligotrophic ($1 - 5$ mm^3L^{-1}), mesotrophic (biomass $= 3 - 5$ mm^3L^{-1}), and eutrophic (10 mm^3L^{-1}).

Finally, we included thermal profiles from the main channel and tributaries to investigate thermal stratification following damming. These thermal profiles were limited to five sites (one in the Madeira River and four in tributaries) and were collected between 2012 and 2017. The statistical analyses together with their respective hypotheses were summarized in table 1. All analyzes were conducted utilizing the R program (R Development Core Team, 2024). Vegan package (Oksanen et al., 2007) was employed to perform PERMANOVA (adonis function) and to compute distance matrices (vegdist function). PCoA was performed with hnp package (Moral et al., 2017), GLMM (glmmTMB function) with glmmTMB package (Brooks et al., 2017) and plots with ggplot2 package (Wickham, 2016).

Table 1. Summary of data analyses and respective hypotheses

Hypotheses to test	Analyses used	References
Hypothesis I: Changes in environmental characteristics and consequently changes in phytoplankton biomass	GLMM	Brooks, Mollie et al., 2017
Hypothesis II: Change in phytoplankton composition between the pre- and post-dam phases	Indval	Dufrêne and Legendre, 1997
	PERMANOVA	Anderson, 2017

3   RESULTS

3.1   Environmental conditions

Both the main channel and tributaries exhibited changes in the environment after damming, but changes were more pronounced in the tributaries (Table S1, Figures 2 and 3). An increase in chemical oxygen demand (COD) was observed in both environments. The main channel showed a reduction in alkalinity, inorganic carbon, silica, conductivity, and pH, whereas tributaries displayed a decrease in light availability (Zeu and Zeu:Zmax) and iron,

but an increase in depth, conductivity, inorganic carbon, temperature, and alkalinity. The

variation of most environmental factors seemed to increase with damming at both the main

channel and tributaries (Table 2), except for orthophosphate, BOD, and COD which

decreased.

Figure 2. The effect of damming on the variation of environmental variables measured at the
tributaries of the Madeira River. Asterisks (*) in the figure indicate a significant difference.
The vertical axis displays the log-transformed values for each variable, except for pH and the
ratio of Zeu:Zmax. *IC* – dissolved inorganic carbon; *BOD* – biochemical oxygen demand;
COD – chemical oxygen demand; *DIN* – dissolved inorganic nitrogen; Z_{eu}:Z_{max} – light
availability in the water column; *COND* – Conductivity; *Alk* – alkalinity; *Zeu* – euphotic
zone; *SRP* – soluble reactive phosphorous; *SI* - Silica.

Figure 3. The effect of damming on the variation of environmental variables measured at Madeira River. Asterisks (*) in the figure indicate a significant effect of damming. The vertical axis displays the log-transformed values for each variable, except for pH and the ratio of Zeu:Zmax. The boxes in the plot represent the first and third quartiles, the horizontal bar represents the median, and the vertical bars indicate the minimum and maximum values, excluding any outliers, which are represented as individual points. IC – dissolved inorganic carbon; BOD – biochemical oxygen demand; COD – chemical oxygen demand; DIN – dissolved inorganic nitrogen; Zeu:Zmax – light availability in the water column; COND – Conductivity; Alk – alkalinity; Zeu – euphotic zone; SRP – soluble reactive phosphorous; SI - Silica.

287   **Table 2:** Mean and coefficient of variation (CV, %) of abiotic variables in the main channel
288   and tributaries during the construction phases of the dam.

	Madeira		Tributaries	
	Before	**After**	**Before**	**After**
Temperature (°C)	27.91 (5.68)	28.04 (6)	26.61 (7.98)	27.76 (7.32)
pH	7 (7.7)	6.41 (10.88)	6.18 (11.84)	6.06 (13.06)
Conductivity (μS.cm^{-1})	88.9 (32.17)	66.35 (52.98)	10.31 (75.87)	25.54 (123.92)
$Z_{eu}:Z_{max}$ (m)	0.03 (141.07)	0.06 (184.55)	0.59 (51.73)	0.36 (85.52)
Euphotic zone - Z_{eu} (m)	0.22 (66.38)	0.57 (136.02)	2.09 (59.31)	1.63 (73.58)
Orthophosphate (μg L^{-1})	0.09 (162.19)	0.07 (85.63)	0.01 (370.74)	0.02 (159.5)
Dissolved inorganic nitrogen (μg L^{-1})	1.74 (52.95)	1.89 (62.93)	0.97 (61.4)	1.13 (73.73)
Biochemical oxygen demand (mg L^{-1})	1.18 (66.06)	1.28 (56.22)	1.45 (66.4)	1.39 (61.85)
Chemical oxygen demand (mg L^{-1})	10.11 (74.71)	12.69 (63.76)	9.47 (71.57)	14.48 (65.46)
Iron (μg L^{-1})	5.05 (80.42)	1.39 (80.54)	1.25 (71.22)	1.03 (61.62)
Alkalinity (mEqL^{-1})	28.22 (34.25)	23.19 (45.47)	6.71 (58.59)	10.63 (72.73)
Inorganic carbon (mg L^{-1})	7.31 (36.15)	6.31 (37.72)	2.76 (71.73)	4.53 (38.89)
Silica (μg L^{-1})	22.55 (107.92)	9.28 (73.69)	6.94 (43.68)	7.94 (57.16)

WT - water temperature - °C; pH; IC – dissolved inorganic carbon; BOD - biochemical
oxygen demand; COD - chemical oxygen demand; DIN – dissolved inorganic nitrogen;
$Z_{eu}:Z_{max}$ – light availability in the water column, SRP – soluble reactive phosphorus, Si -
Silica.

3.2   *Phytoplankton structure and biomass*

In this study, we identified a total of 525 taxa, with the highest number of species
belonging to green algae (236 species, mean of 0.095 ind.mL^{-1}), Phytoflagellates comprised
118 species (mean of 0.145 ind.mL^{-1}), followed by diatoms (86 species, mean of 0.057
ind.mL^{-1}). Cyanobacteria accounted for 72 species (mean of 0.017 ind.mL^{-1}), and
xantophyceans were represented by 13 species (mean of 0.0004 ind.mL^{-1}).

Before and after damming, the main channel recorded 119 and 167 species,
respectively, with diatoms (mainly bacillariophyceans) exhibiting the highest richness both
before (30) and after (37) damming. In the tributaries, the number of species increased from
307 before damming to 457 after damming, with zygnemaphyceans having the greatest
richness both before (62) and after (94) damming.

PERMANOVA revealed a significant effect of damming on the phytoplankton
diversity for the main channel (F = 1.87, p = 0.01) and tributaries (F = 12.89, p <0.01). In
addition to impoundment, we also found a significant effect of hydrological periods on
community diversity (for main channel, F = 1.22, p = 0.001; for tributaries, F = 2.09, p
<0.01). Both the PERMANOVA results and their graphical representation (Figure 4) showed
that damming altered communities and increased the spatial and temporal variation in species
biomass in both the main channel and tributaries across all hydrological periods.

311
Figure 4: Temporal variation of phytoplankton community (species abundance) related to
damming phases and hydrological periods at both the main channel and tributaries. In all
cases, the community's heterogeneity was higher in the after damming phase. The higher
variation seemed to be related to the low water period.

IndVal analysis indicated that the number of species linked with tributaries increased
after damming, while indicators of the Madeira River decreased. Diatoms, which are related
to flowing waters, were found to be exclusive indicators of pre-damming conditions for both
tributaries and the main channel. It is noteworthy that most of the indicator species associated

with tributaries in the post-damming phase were indicative of standing water, specifically

phytoflagellates (Table 3).

Table 3. Species as indicators of the conditions at each damming phase (before and after) and
system (tributaries and Madeira River). *IndVal*, indicator value based on the relative
frequency and relative average abundance in the clusters Phase – system. The table shows
only those species with a high probability of being an indicator (p < 0.05). *Freq*, specie
frequency at each cluster. The taxonomic groups were colored differently to highlight the
representativity of each group at each phase – system: blue – Cyanobacteria; pink – Diatoms;
green – Green algae; golden – Phytoflagellates.

Phase - system	Indval	p	Freq	Taxa	G
Before - Tributaries	0.05	0.013	8	*Lemmermanniella flexa* Hindák	
	0.04	0.011	7	*Synechocystis aquatilis* Sauv.	
	0.04	0.044	12	*Pseudanabaena mucicola* (Hüb.-Pest. e W. Naumann) Bourr.	
	0.03	0.03	4	*Snowella* sp.	
	0.05	0.011	11	*Cymbopleura naviculiformis* (Auersw.) Krammer	
	0.04	0.043	12	Pennales sp1	
	0.04	0.028	12	*Pinnularia microstauron* (Ehrenb.) Cleve	
	0.1	0.008	43	*Oocystis lacustris* Chodat	
	0.09	0.031	39	*Monoraphidium minutum* (Naegeli) Komárk. - Legn.	
	0.05	0.016	11	*Cosmarium punctulatum* Bréb.	
	0.04	0.028	14	*Scenedesmus acunae* Com.	
	0.33	0.001	91	*Cryptomonas* sp.	
Before - Madeira	0.03	0.024	4	*Aphanizomenon gracile* Lemmerm.	
	0.22	0.001	99	*Fragilaria* sp.	
	0.19	0.006	111	*Aulacoseira granulata* var. *granulata* (Ehrenb.) Sim.	
	0.17	0.001	43	*Cyclotella meneghiniana* Kütz.	
	0.17	0.001	42	*Aulacoseira distans* (Ehrenb.) Sim.	
	0.17	0.001	60	*Aulacoseira granulata* var. *angustissima* (O. Müller) Sim.	
	0.13	0.001	23	*Ulnaria ulna* (Nitzsch.) Comp.	
	0.12	0.003	31	*Nitzschia* sp.	
	0.08	0.001	7	*Gyrosygma* sp1	
	0.08	0.002	9	*Cocconeis* sp.	
	0.06	0.008	19	*Aulacoseira ambigua* (Grunow) Sim.	
	0.06	0.043	26	*Navicula viridula* (Kütz.) Ehrenb.	
	0.05	0.025	16	*Achnanthidium minutissimum* (Kütz.) Czarn.	
	0.04	0.015	7	*Melosira varians* Agardh	
	0.05	0.007	5	*Actinastrum aciculare* Playf.	
	0.05	0.027	14	*Desmodesmus intermedius* var. *acutispinus* (Roll) E.	

Phase - system	Indval	p	Freq	Taxa	G
				Hegew.	
	0.04	0.005	2	*Monactinus simplex* (Meyen) Corda	
	0.04	0.05	16	*Scenedesmus* sp.	
	0.04	0.03	10	*Scenedesmus acuminatus* (Lagerh.) Chodat	
	0.13	0.002	39	*Eunotia bilunaris* (Ehrenb.) Schaar.	
	0.07	0.015	29	*Urosolenia eriensis* (H. L. Sm.) Round e Craw.	
	0.2	0.002	107	*Scenedesmus ecornis* (Ehrenb.) Chodat	
	0.19	0.001	95	*Monoraphidium griffithii* (Berk.) Komárk.-Legn.	
	0.16	0.006	59	*Crucigenia tetrapedia* (Kirch.) W. e G.S. West	
	0.13	0.005	57	*Monoraphidium irregulare* (G. M. Smith) Komárk.-Legn.	
	0.11	0.016	62	*Closteriopsis* sp.	
	0.11	0.015	64	*Monoraphidium convolutum* (Corda) Komárk. - Legn.	
	0.1	0.005	29	*Crucigenia fenestrata* (Schm.) Schm.	
	0.1	0.016	46	*Desmodesmus hystrix* (Lagerh.) E. Hegew.	
	0.09	0.045	76	*Monoraphidium komarkovae* Nygaard	
	0.09	0.005	26	*Desmodesmus brasiliensis* (Bohlin) E. Hegew.	
	0.08	0.002	23	*Nephrochlamys* sp.	
After - Tributaries	0.07	0.024	37	*Ankyra judayi* (G.W. Smith) Fott	
	0.06	0.025	23	*Willea rectangularis* (Nägeli) Komárek	
	0.06	0.023	18	*Monoraphidium circinale* (Nygaard) Nygaard	
	0.06	0.017	19	*Cosmarium regnesi* Reins.	
	0.05	0.032	14	Chlorococcales colonial não identificada 1	
	0.47	0.001	168	*Cryptomonas brasiliensis* Castro, Bic. e Bic.	
	0.29	0.001	97	*Cryptomonas erosa* Ehrenb.	
	0.25	0.001	120	*Trachelomonas volvocinopsis* Svirenko	
	0.24	0.001	115	*Peridinium* sp.	
	0.22	0.002	75	*Gonyostomum* sp.	
	0.2	0.001	108	*Chlamydomonas* sp.	
	0.18	0.019	94	*Euglena* sp.	
	0.18	0.001	55	*Trachelomonas oblonga* Lemmerm.	
	0.16	0.004	65	*Lepocinclis ovum* (Ehrenb.) Lemmerm.	
	0.15	0.007	86	*Mallomonas* sp.	
	0.09	0.046	54	*Cryptomonas curvata* Ehrenb. emend. Pen.	
	0.09	0.016	43	*Euglena acus* Ehrenb.	
	0.08	0.048	47	*Trachelomonas volvocina* Ehrenb.	
	0.07	0.009	19	*Strombomonas scabra* (Playf.) Tell e Conf.	
	0.06	0.014	20	*Chroomonas* sp.	
	0.05	0.033	20	*Trachelomonas lacustris* Dezep.	
	0.05	0.026	17	*Phacus* sp.	
	0.05	0.026	18	*Phacus acutus* Pochmann	

Phase - system	Indval	p	Freq	Taxa	G
	0.04	0.049	10	*Phacus horridus* Pochm.	
	0.04	0.045	10	*Pandorina morum* (F. Muller) Bory	
	0.03	0.048	11	*Trachelomonas similis* Stokes var. *spinosa* Hub.-Pest.	
After - Madeira	0.02	0.047	2	*Pediastrum duplex* Meyen	

In the tributaries, the total biomass ranged from 0.0001 to 4.74 mm^3L^{-1} before

damming. After damming values varied between 0.0001 and 15 mm^3L^{-1}. Most samples (98%

before and 96% after damming) showed a biomass lower than 2 mm^3L^{-1}. After damming, we

observed biomass higher than 2 mm^3L^{-1} (four samples), 3 mm^3L^{-1} (four samples), 5 mm^3L^{-1}

(1 sample), and 10 mm^3L^{-1} (one sample). In the Madeira River, the total biomass ranged

between 0.001 and 3.38 mm^3L^{-1} before damming, and after damming, values varied between

0.0001 and 0.97 mm^3L^{-1}. Biomass was less than 2 mm^3L^{-1} in 98% of the samples before

damming and 100% after damming. According to Vollenweider, the registered biomass

conditions mainly ranged from oligotrophic (biomass = 1 – 5 mm^3L^{-1}) to mesotrophic

(biomass = 3 – 5 mm^3L^{-1}).

GLMM revealed differences in total biomass between damming phases in both the

main channel and tributaries (Figure 5A and Table S2). In the tributaries, total and

phytoflagellates biomass increased (Figure 5A), with variations between phases depending on

the season considered. Specifically, no differences were observed during the rising season for

total biomass and at high water for phytoflagellates, while cyanobacteria showed differences

only in the high and receding water seasons (Figure 5B and Table S3). In the main channel,

total biomass decreased, along with diatoms and green algae biomass, while phytoflagellates

increased (Figure 5A). Differences between the phases were consistent across season for

phytoflagellates. In contrast, differences between the phases at the low- and rising-water

seasons were observed for the total biomass and diatoms, at the rising- and high-water

seasons for greens, and at the rising waters for cyanobacteria (Figure 5B).

Figure 5: A. Difference in the distribution of total biomass and for each group in the tributaries and main channel during the construction phases of the dam. Asterisks represent significant differences. B. Differences in the biomass distribution of the total community and of each group during the hydrological periods and construction phases of the dam. Asterisks indicate significant differences in the biomass values obtained.

3.3 Relationship between abiotic variables and community biomass

The GLMM also revealed that the factors leading total biomass differed between the main channel and its tributaries. (Figure 6 Table S5). Moreover, the significance of these factors varied across taxonomic groups, depending on whether the occurrence or the biomass data were considered (Figure 6).

Figure 6: Influence of abiotic variables on total biomass and on the biomass and occurrence of algae groups from the main channel and tributaries. Asterisks represent a significant effect. Biomass is shown by blue bars while occurrence is shown by red bars. Zeu:Zmax – light availability in the water column; Zeu – euphotic zone; WT - water temperature; SRP – soluble reactive phosphorous; SI - Silica; PH - hydrogen potential; COD – chemical oxygen demand; BOD – biochemical oxygen demand; IC – dissolved inorganic carbon; DIN – dissolved inorganic nitrogen; COND – Conductivity; Alk – alkalinity.

4   Discussion

Several dams will occupy rivers in the Amazon region in the coming decades. Understanding their environmental implications and the subsequent effect on organisms is crucial for developing effective management and conservation strategies, as for anticipating and mitigating potential future. In this study, we demonstrated that the Jirau dam, as expected, induced changes in the biomass and structure of the phytoplankton communities in both the main channel and tributaries, corroborating our hypotheses. Specifically, damming promoted increased phytoplankton biomass in the tributaries, driven by the proliferation of species adapted to lacustrine, particularly phytoflagellates. Our findings highlight that while hydroelectric projects with Run-of-River (ROR) dams have historically been associated with

minimal environmental impact, they can still significantly alter biotic characteristics, even in

main channels where ROR dams are presumed to have less impact. This section delves into

the impacts of damming on phytoplankton community dynamics and total biomass. We also

discuss broader implications, including the potential future degradation of the Amazonian

region, and how our results contribute to preventing undesirable scenarios.

4.1   Changes in phytoplankton structure

The construction of the Jirau dam seemed to influence changes in the phytoplankton

communities in both the main channel and its tributaries, increasing the variation in

communities at all hydrological seasons. This contrasts with previous research in the same

study zone, which showed that biotic variation of planktonic communities decreased after

damming (Pineda et al., 2024; Zanon et al., 2024). Unlike that research, our analysis focused

on community variation between damming phases, considering each hydrological season

separately. Figure 4 shows both spatial (between sampled points) and temporal (samples from

different periods but within the same hydrological season and damming phase) variations

together. However, given that the temporal water level variation decreased after damming

and that spatial variation increased due to the emergence of lentic regions (which contrast

with the lotic regions), the observed increase in sample dispersion was mainly related to

spatial variation rather than temporal variation within the same hydrological season. Our

results, together with research findings of Pineda et al. (2024) and Zanon et al. (2024),

suggest that after damming, there could be an increase in spatial variation while temporal

variation decreases. However, the spatial variation does not seem to be enough to compensate

for the loss of temporal variation since the general pattern exhibit reduction in biotic

heterogeneity (Pineda et al., 2024).

The increase in spatial variation that we observed can be attributed to the mosaic of

habitats with varying features that resulted after damming. Following damming, as in our

case, a region impacted by run-of-river projects shows not just rivers but also habitats with

lake-like properties because of backwaters formed at the tributary's junction with the main

channel (Almeida et al., 2019). Each one of those habitats can then favor different species

with different adaptations (Reynolds et al., 2002; Rodrigues et al., 2018; Zanco et al., 2017)

and increase the spatial variation in communities. Our results indicated that after damming

the number of indicator species of tributaries increased by 300%. It means that the

characteristics of tributaries favored different species than those favored by the main channel.

Furthermore, most of the species identified as tributary indicators after damming are typical

of standing waters (Supplementary material), whereas most indicator species at the main

channel were associated with lotic environments. It is consistent with our evidence

(Supplementary material – Figure S1) and from predictions from other studies (e.g.,

Fearnside, 2013) about the stratification of tributaries after damming.

4.2   Changes in phytoplankton biomass

In the case of the total biomass, it decreased in the main channel (which was

unexpected) and increased in the tributaries (as expected). We believe that the reduction of

biomass in the main channel was caused by a decrease in phytoplankton export from

tributaries, as the dam in the Madeira River increased the water level and reduced the water

flow in the tributaries. However, more research is necessary to study the effect of damming

on the dispersion between tributaries and the main channel. In any case, our result contrasts

with evidence showing that run-of-river projects have little effect on large rivers (Almeida et

al., 2019), and suggests that this kind of project may affect the flux of matter and energy in

dammed rivers by decreasing the biomass of primary producers in the main channel.

We attributed the biomass increase in the tributaries to a reduction in their water flow,

which increase water residence time, promotes stratification, and consequently favors

phytoplankton development. As support for that assumption, we demonstrated an increase in

species associated with standing waters as well as evidence of water stratification at certain

tributaries after damming (Supplementary material – Figure S1). Additionally, we observed

that water temperature had a positive effect on total biomass, and that temperature increased

after damming, which may relate to stratification. Moreover, the stratification in tributaries in

Madeira River has been documented after the construction of the Santo Antônio hydropower

project (Almeida et al., 2019), which is also a large ROR located downstream of the Jirau

dam. Unfortunately, we did not have stratification data for each sampling to include in our

biomass models. Nevertheless, our observations suggest that even a run-of-river dam can lead

to changes in the hydrodynamics of affected basins, mainly at tributaries, and that these

changes can increase the phytoplankton biomass in tributaries.

Although total biomass increased at the tributaries, not all the phytoplankters reacted

to damming in the same way. While the biomass of flagellates increased, that of

cyanobacteria decreased. We believe that while lentic characteristics appeared to favor the

whole phytoplankton community, chemical factors appeared to determine which group might

thrive best. For instance, we found that an increase in nitrogen concentration following

damming had a negative effect on cyanobacteria growth but favored green algae. In another

case, high phosphate concentrations boosted cyanobacteria while harming phytoflagellates. In

both examples, while the positive effect was associated with increased biomass, the negative

effect may be linked to the rise in biomass of antagonistic groups, which dominate and limit

the growth of other phytoplankton groups (Reynolds, 2006).

Phytoflagellates dominated the biomass, which may be related to the accumulation of

organic material in the backwater regions of the tributaries. Despite the vegetation removal

practices implemented before the reservoir filling in the Jirau Project (Energia Sustentável do

Brasil, 2020), it seems that the flooded area still covered a significant amount of vegetation

biomass (Cochrane et al., 2017), affecting the plant community and the decomposition of

organic matter (Medeiros et al., 2023). The 10-meter average increase in the water level of

the Madeira River submerged more than 160 km² of natural forest (Cochrane et al., 2017).

We observed that chemical oxygen demand (COD) increased after damming in tributaries,

indicating that the process of degradation of organic matter consumed more oxygen with

damming. In this way, organisms with the ability of take advantage of a high amount of

organic matter could domain. As changes in water features influence all aquatic species, these

alterations can have an impact on the entire trophic web. For instance, a decrease in oxygen

can relate to the death of fish (Demertzioglou et al., 2022) and the favoring of invasive

species more tolerant to those new environmental conditions, such as some macrophyte

species (Harrow-Lyle and Kirkwood, 2021).

The increase in algal biomass that we have identified here may not be restricted to our

study zone. A comparable rise in phytoplankton biomass in backwater was registered in the

influence zone of the dam of the Santo Antônio dam (Appel, 2017), downstream Jirau dam.

As in our case, damming caused the formation of standing waters in that location (Almeida et

al., 2019), which likely supported algal development as in our case. Furthermore, an increase

in phytoplankton biomass may be predicted through the amazon, since the Brazilian

government plans to expand hydroelectric power generation in the Brazilian Amazon over the

next 25 years (Nobre et al., 2016) with the construction of thirty large dams (Empresa de

Pesquisa Energética, 2013) and the flooding of $\sim$12,000 km^2 (Nobre et al., 2016).

4.3   What our findings mean for the future and how we can work to prevent

Our study revealed that the creation of backwaters post-damming in tributaries

appeared to favor an increase in algal biomass, resembling lake-like conditions. Despite this

increase, the biomass values remained low, indicative of oligotrophic to mesotrophic

conditions (Vollenweider, 1968). This limited biomass was primarily limited by low

phosphorus availability (less than $0.001 \mu g.L^{-1}$). Our findings suggest that while standing

water conditions created by the Jirau dam enhanced hydrodynamic conditions for

phytoplankton growth, current nutrient levels are insufficient to support high and problematic

phytoplankton biomass, at least in the present nutrient conditions.

However, we caution that the optimal hydrodynamic conditions created by the dam,

such as increased residence time and stratification in tributaries, could potentially synergize

with future nutrient increases in Amazonian rivers. Studies (e.g., dos Santos et al., 2014)

indicate that nutrient enrichment could lead to elevated phytoplankton biomass levels, like

issues observed in other dam-affected regions worldwide (Grill et al., 2019; Winemiller et al.,

2016).

Human activities associated with nutrient enrichment in water have reached areas of

the Amazon that would have been unimaginable a few decades ago. For example,

deforestation for pastures has been linked to phosphorus rise in rivers like the Ji-Paraná River

in the Amazon basin (Ballester et al., 2003). Furthermore, projections suggest that by 2050,

pastures and agricultural expansion could cover 40% of the entire Amazon region (Soares-

Filho et al., 2006), potentially increasing nutrient loads to waterways. Urban expansion

facilitated by dams and agro-industrial operations has also contributed to nutrient increases in

Amazonian rivers due to untreated sewage (Ferreira et al., 2021).

Given these observations, we urge authorities to implement stringent protocols for

nutrient management in future agricultural and urban projects in dam-influenced zones of the

Amazon. Proper management practices can mitigate the risks associated with phytoplankton

blooms, preserving water quality and the ecosystem services derived from aquatic

environments, including water supply, recreation, and fishing. This proactive approach to

nutrient management is essential to safeguarding the ecological integrity of Amazonian

rivers.

503   5   Conclusion

Even run-of-the-river hydroelectric projects like the Jirau hydroelectric plant can alter

water chemistry and hydrodynamics, subsequently affecting the characteristics of microbial

communities such structure and biomass of phytoplankton. This impact could potentially

extend across the Amazon region, as several similar hydropower projects are currently under

construction in the area. Given that the flooding of large swathes of vegetation is a common

consequence of these projects, it is anticipated that dams in the Amazon region will lead to an

increase in mixotrophic organisms such as flagellates, which take advantage of organic

matter. Additionally, the dams are expected to elevate the chemical demand for oxygen to

degrade the accumulated organic matter, thereby reducing oxygen availability. While

damming does promote the biomass of phytoplankton due to extended residence times and

water column stratification, it may not suffice to produce problematic levels of biomass due

to limited nutrient availability. Therefore, authorities must remain vigilant regarding activities

that may exacerbate these issues and impact water quality in the Amazon. This includes

monitoring practices that could result in heightened nutrient loading in aquatic habitats, such

as agricultural practices, urban development, industrial activities, and sewage discharge.

Acknowledgments

We thank the Programa de Pós-graduação em Ecologia de Ambientes Aquáticos
Continentais and the Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura (Nupélia)
from the Universidade Estadual de Maringá (UEM) for supplying infrastructure. We thank
the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior— Brazil (CAPES) for the
scholarship granted to BMC, FMZ and RFS. AP thanks Conselho Nacional de
Desenvolvimento Científico e Tecnológico (CNPq) for postdoctoral fellowships. LCR was
supported by productivity fellowships from CNPq. The authors state that they have no
conflicts of interest relating to this study.

Declarations

Data availability The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request

Funding AP was awarded a scholarship from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). BMC, FMZ and RFS were awarded scholarship from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior— Brazil (CAPES)

Consent to participate Not applicable.

Consent to publication Not applicable

Ethical approval Not applicable. The present study did not involve research with humans or animals

References

Agudelo, J., Arias, P.A., Vieira, S.C., Martínez, J.A., 2019. Influence of longer dry seasons in the Southern Amazon on patterns of water vapor transport over northern South America and the Caribbean. Clim Dyn 52, 2647–2665. https://doi.org/10.1007/s00382-018-4285-1

Almeida, R.M., Hamilton, S.K., Rosi, E.J., Arantes, J.D., Barros, N., Boemer, G., Gripp, A., Huszar, V.L.M., Junger, P.C., Lima, M., Pacheco, F., Carvalho, D., Reisinger, A.J., Silva, L.H.S., Roland, F., 2019. Limnological effects of a large Amazonian run-of-river dam on the main river and drowned tributary valleys. Sci Rep 9. https://doi.org/10.1038/s41598-019-53060-1

Almeida, R.M., Hamilton, S.K., Rosi, E.J., Barros, N., Doria, C.R.C., Flecker, A.S., Fleischmann, A.S., Reisinger, A.J., Roland, F., 2020. Hydropeaking operations of two run-of-river mega-dams alter downstream hydrology of the largest Amazon tributary. Front Environ Sci 8, 120. https://doi.org/10.3389/fenvs.2020.00120

Almeida, R.M., Tranvik, L., Huszar, V.L.M., Sobek, S., Mendonça, R., Barros, N., Boemer, G., Durval, J., Arantes Jr, J.D., Roland, F., 2015. Phosphorus transport by the largest Amazon tributary (Madeira River, Brazil) and its sensitivity to precipitation and damming [WWW Document]. Inland Waters. https://doi.org/10.5268/IW-5.3.815

American Public Health Association, 2012. Standard Methods for the Examination of Water and Wastewater, 21st ed, Standard Methods. American Public Health Association, American Water Works Association, and Water Environment Federation, Washington, DC.

ANA, 2022. Agência Nacional de Águas [WWW Document]. URL
https://www.snirh.gov.br/hidroweb/apresentacao (accessed 11.9.22).
APHA, 2005. Standard Methods for the Examination of Water and Wastewater. American
Public Health Association, American Water Works Association, and Water Environment
Federation, Washington DC (USA).
Appel, E., 2017. Efeitos do barramento tipo fio d'água sobre as comunidades
fitoplanctônicas do rio Madeira e seus tributários na área de influência da hidrelétrica
Santo Antônio, RO, Brasil. Universidade Federal do Rio de Janeiro.
Aragón, L.E., 2018. A dimensão internacional da Amazônia: um aporte para sua
interpretação. Revista NERA 1, 14–33.
Ballester, M.V.R., Victoria, D.D.C., Krusche, A. V., Coburn, R., Victoria, R.L., Richey, J.E.,
Logsdon, M.G., Mayorga, E., Matricardi, E., 2003. A remote sensing/GIS-based
physical template to understand the biogeochemistry of the Ji-Paraná river basin
(Western Amazônia). Remote Sens Environ 87, 429–445.
https://doi.org/10.1016/j.rse.2002.10.001
Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H.,
White, J.S.S., 2009. Generalized linear mixed models: a practical guide for ecology and
evolution. Trends Ecol Evol 24, 127–135. https://doi.org/10.1016/j.tree.2008.10.008
Borcard, D., Gillet, F., Legendre, P., 2018. Numerical Ecology with R, Use R! Springer
International Publishing, Cham. https://doi.org/10.1007/978-3-319-71404-2
Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A.,
Skaug, H.J., Mächler, M., Bolker, B.M., 2017. glmmTMB balances speed and flexibility
among packages for zero-inflated generalized linear mixed modeling. R Journal 9, 378–
400. https://doi.org/10.32614/rj-2017-066
Carvalho, R., de Aguiar, A.P.D., Amaral, S., 2020. Diversity of cattle raising systems and its
effects over forest regrowth in a core region of cattle production in the Brazilian
Amazon. Reg Environ Change 20, 1–15. https://doi.org/10.1007/s10113-020-01626-5
Cochrane, S.M.V., Matricardi, E.A.T., Numata, I., Lefebvre, P.A., 2017. Landsat-based
analysis of mega dam flooding impacts in the Amazon compared to associated
environmental impact assessments: Upper Madeira River example 2006–2015. Remote
Sens Appl 7, 1–8. https://doi.org/10.1016/j.rsase.2017.04.005
Cole, A.G., 1994. Textbook of Limnology, 4th Ed. ed. Waveland Press, Inc. Prospect
Heights, Illinois.
Demertzioglou, M., Genitsaris, S., Mazaris, A.D., Kyparissis, A., Voutsa, D., Kozari, A.,
Kormas, K.A., Stefanidou, N., Katsiapi, M., Michaloudi, E., Moustaka-Gouni, M., 2022.
A catastrophic change in a european protected wetland: From harmful phytoplankton
blooms to fish and bird kill. Environmental Pollution 312, 120038.
https://doi.org/10.1016/j.envpol.2022.120038
Dias, V. de S., da Luz, M.P., Medero, G.M., Nascimento, D.T.F., 2018. An overview of
hydropower reservoirs in Brazil: Current situation, future perspectives and impacts of
climate change. Water (Switzerland) 10. https://doi.org/10.3390/w10050592
dos Santos, J.C.N., de Andrade, E.M., de Araújo Neto, J.R., Meireles, A.C.M., de Queiroz
Palácio, H.A., 2014. Land use and trophic state dynamics in a tropical semi-arid
reservoir. Revista Ciencia Agronomica 45, 35–44. https://doi.org/10.1590/s1806-
66902014000100005
Dufrêne, M., Legendre, P., 1997. Species Assemblages and Indicator Species: the Need
for a Flexible Asymmetrical Approach. Ecol Monogr 67, 345–366.

Empresa de Pesquisa Energética, 2013. Plano Decenal de Expansão de Energia 2013–
2022. Empresa Pesquisa Energética, Rio de Janeiro.
Energia Sustentável do Brasil, 2020. Dados técnicos relacionados à Usina Hidrelétrica de
Jirau. Retrieved [WWW Document].
Fearnside, P.M., 2013. Credit for climate mitigation by Amazonian dams: loopholes and
impacts illustrated by Brazil's Jirau Hydroelectric Project. Carbon Manag 4, 681–696.
https://doi.org/10.4155/cmt.13.57
Ferrão-Filho, A.D.S., Kozlowsky-Suzuki, B., 2011. Cyanotoxins: Bioaccumulation and effects
on aquatic animals. Mar Drugs. https://doi.org/10.3390/md9122729
Ferreira, S.J.F., Pinel, S., Ríos-Villamizar, E.A., Miranda, S.Á.F., Pascoaloto, D., Vital,
A.R.T., Monteiro, M.T.F., da Silva, M. do S.R., da Cunha, T.R.B., dos Santos, A.S.,
Bender, S., da Cunha, H.B., 2021. Impact of rapid urbanization on stream water quality
in the Brazilian Amazon. Environ Earth Sci 80, 316. https://doi.org/10.1007/s12665-021-
09621-7
Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., Babu, S., Borrelli, P.,
Cheng, L., Crochetiere, H., Ehalt Macedo, H., Filgueiras, R., Goichot, M., Higgins, J.,
Hogan, Z., Lip, B., McClain, M.E., Meng, J., Mulligan, M., Nilsson, C., Olden, J.D.,
Opperman, J.J., Petry, P., Reidy Liermann, C., Sáenz, L., Salinas-Rodríguez, S.,
Schelle, P., Schmitt, R.J.P., Snider, J., Tan, F., Tockner, K., Valdujo, P.H., van
Soesbergen, A., Zarfl, C., 2019. Mapping the world's free-flowing rivers. Nature 569,
215–221. https://doi.org/10.1038/s41586-019-1111-9
Gusso, A., Ducati, J.R., Bortolotto, V.C., 2017. Analysis of soybean cropland expansion in
the southern Brazilian Amazon and its relation to economic drivers. Acta Amazon 47,
281–292. https://doi.org/10.1590/1809-4392201700543
Guyot, J.L., Filizola, N., Quintanilla, J., Cortez, J., 1996. Dissolved solids and suspended
sediment yields in the Rio Madeira basin, from the Bolivian Andes to the Amazon, in:
Erosion and Sediment Yield: Global and Regional Perspectives. IAHS Publication,
Exeter, pp. 55–63.
Harrow-Lyle, T.J., Kirkwood, A.E., 2021. Low Benthic Oxygen and High Internal
Phosphorus-Loading are Strongly Associated With the Invasive Macrophyte Nitellopsis
obtusa (starry stonewort) in a Large, Polymictic Lake. Front Environ Sci 9, 357.
https://doi.org/10.3389/FENVS.2021.735509/BIBTEX
Hartmann, J., Fonseca, A., Trias, M., Mandilovich, V., 2023. Jirau Hydropower Plant:
Assessment Report. London.
Hillebrand, H., Dürselen, C.-D., Kirschtel, D., Pollingher, U., Zohary, T., 1999. Biovolume
Calculation for Pelagic and Benthic Microalgae. J Phycol 35, 403–424.
https://doi.org/10.1046/j.1529-8817.1999.3520403.x
Huisman, J., Codd, G.A., Paerl, H.W., Ibelings, B.W., Verspagen, J.M.H., Visser, P.M., 2018.
Cyanobacterial blooms. Nat Rev Microbiol 16, 471–483. https://doi.org/10.1038/s41579-
018-0040-1
Jensen, J.P., Jeppesen, E., Olrik, K., Kristensen, P., 1994. Impact of nutrients and physical
factors on the shift from cyanobacterial to chlorophyte dominance in shallow Danish
lakes. Canadian Journal of Fisheries and Aquatic Sciences 51, 1692–1699.
https://doi.org/10.1139/f94-170
Komárek, J., Anagnostidis, K., 1998. Cyanoprokaryota. 1. Teil Chroococcales,
Sübwasserflora von Mitteleuropa. Gustav Fischer Verlag.

Krztoń, W., Kosiba, J., Pociecha, A., Wilk-Woźniak, E., 2019. The effect of cyanobacterial
blooms on bio- and functional diversity of zooplankton communities. Biodivers Conserv
28, 1815–1835. https://doi.org/10.1007/s10531-019-01758-z
Latrubesse, E.M., Arima, E.Y., Dunne, T., Park, E., Baker, V.R., D'Horta, F.M., Wight, C.,
Wittmann, F., Zuanon, J., Baker, P.A., Ribas, C.C., Norgaard, R.B., Filizola, N., Ansar,
A., Flyvbjerg, B., Stevaux, J.C., 2017. Damming the rivers of the Amazon basin. Nature
546, 363–369. https://doi.org/10.1038/nature22333
Lees, A.C., Peres, C.A., Fearnside, P.M., Schneider, M., Zuanon, J.A.S., 2016. Hydropower
and the future of Amazonian biodiversity. Biodivers Conserv 25, 451–466.
https://doi.org/10.1007/s10531-016-1072-3
Macedo, M.N., DeFries, R.S., Morton, D.C., Stickler, C.M., Galford, G.L., Shimabukuro, Y.E.,
2012. Decoupling of deforestation and soy production in the southern Amazon during
the late 2000s. Proc Natl Acad Sci U S A 109, 1341–1346.
https://doi.org/10.1073/pnas.1111374109
Medeiros, M.B., Oliveira, W.L., Rodrigues, F.R.O., Silva, R.D., Ferreira, Í.J.K., Ayala, W.E.,
Silva, S.R., Souza, R.T., Simon, M.F., 2023. Monitoring the impacts of a mega-dam on
Amazonian understorey herbs. For Ecol Manage 536, 120909.
https://doi.org/10.1016/J.FORECO.2023.120909
Moral, R.A., Hinde, J., Demétrio, C.G.B., 2017. Half-Normal Plots and Overdispersed
Models in R : The hnp Package. J Stat Softw 81. https://doi.org/10.18637/jss.v081.i10
Moran, E.F., 2020. Changing how we build hydropower infrastructure for the common good:
lessons from the Brazilian Amazon. Civitas 20, 5–15. https://doi.org/10.15448/1984-
7289.2020.1.34643
Nobre, C.A., Sampaio, G., Borma, L.S., Castilla-Rubio, J.C., Silva, J.S., Cardoso, M., 2016.
Land-use and climate change risks in the amazon and the need of a novel sustainable
development paradigm. Proc Natl Acad Sci U S A 113, 10759–10768.
https://doi.org/10.1073/pnas.1605516113
Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J.,
Suggests, M., 2007. The vegan package. Community ecology package.
Okuku, E.O., Tole, M., Kiteresi, L.I., Bouillon, S., 2016. The response of phytoplankton and
zooplankton to river damming in three cascading reservoirs of the Tana River, Kenya.
Lakes Reserv 21, 114–132. https://doi.org/10.1111/lre.12127
O'Neil, J.M., Davis, T.W., Burford, M.A., Gobler, C.J., 2012. The rise of harmful
cyanobacteria blooms: The potential roles of eutrophication and climate change.
Harmful Algae 14, 313–334. https://doi.org/10.1016/j.hal.2011.10.027
Pineda, A., Velho, L.F.M., de Carvalho, P., Rodrigues, L.C., 2024. How a mega-dam in the
Amazon's largest tributary altered the drivers of planktonic communities. Hydrobiologia.
https://doi.org/10.1007/s10750-024-05529-y
Pouria, S., De Andrade, A., Barbosa, J., Cavalcanti, R.L., Barreto, V.T.S., Ward, C.J.,
Preiser, W., Poon, G.K., Neild, G.H., Codd, G.A., 1998. Fatal microcystin intoxication in
haemodialysis unit in Caruaru, Brazil. The Lancet 352, 21–26.
https://doi.org/10.1016/S0140-6736(97)12285-1
Pracheil, B.M., DeRolph, C.R., Schramm, M.P., Bevelhimer, M.S., 2016. A fish-eye view of
riverine hydropower systems: the current understanding of the biological response to
turbine passage. Rev Fish Biol Fish 26, 153–167. https://doi.org/10.1007/s11160-015-
9416-8
R Development Core Team, 2024. R: a language and environment for statistical computing.
Reviers, B. de, 2003. Biologie et phylogénie des algues. Éditions Berlin, Paris.

Reynolds, C.S., 2006. The Ecology of Phytoplankton, 1st ed. Cambridge university press,
New York.
Reynolds, C.S., Huszar, V., Kruk, C., Naselli-Flores, L., Melo, S., 2002. Towards a functional
classification of the freshwater phytoplankton. J Plankton Res 24, 417–428.
https://doi.org/10.1093/plankt/24.5.417
Rodrigues, L.C., Pivato, B.M., Vieira, L.C.G., Bovo-Scomparin, V.M., Bortolini, J.C., Pineda,
A., Train, S., 2018. Use of phytoplankton functional groups as a model of spatial and
temporal patterns in reservoirs: a case study in a reservoir of central Brazil.
Hydrobiologia 805, 147–161. https://doi.org/10.1007/s10750-017-3289-x
Rodrigues, M.S., Batista, E.P., Silva, A.A., Costa, F.M., Neto, V.A.S., Gil, L.H.S., 2017.
Change in Anopheles richness and composition in response to artificial flooding during
the creation of the Jirau hydroelectric dam in Porto Velho, Brazil. Malar J 16, 87–93.
https://doi.org/10.1186/s12936-017-1738-7
Soares-Filho, B.S., Nepstad, D.C., Curran, L.M., Cerqueira, G.C., Garcia, R.A., Ramos,
C.A., Voll, E., McDonald, A., Lefebvre, P., Schlesinger, P., 2006. Modelling
conservation in the Amazon basin. Nature 440, 520–523.
https://doi.org/10.1038/nature04389
Souza, C.A. de, Vieira, L.C.G., Legendre, P., Carvalho, P. de, Velho, L.F.M., Beisner, B.E.,
2019. Damming interacts with the flood pulse to alter zooplankton communities in an
Amazonian river. Freshw Biol 64, 1040–1053. https://doi.org/10.1111/fwb.13284
Souza, D.G., Bueno, N.C., Bortolini, J.C., Rodrigues, L.C., Bovo-Scomparin, V.M., de Souza
Franco, G.M., 2016. Phytoplankton functional groups in a subtropical Brazilian
reservoir: responses to impoundment. Hydrobiologia 779, 47–57.
https://doi.org/10.1007/s10750-016-2798-3
Sun, J., Liu, D., 2003. Geometric models for calculating cell biovolume and surface area for
phytoplankton. J Plankton Res 25, 1331–1346. https://doi.org/10.1093/plankt/fbg096
Thomaz, S.M., Bini, L.M., De Souza, M.C., Kita, K.K., Camargo, A.F.M., 1999. Aquatic
macrophytes of Itaipu Reservoir, Brazil: Survey of species and ecological
considerations. Brazilian Archives of Biology and Technology.
https://doi.org/10.1590/S1516-89131999000100003
Thornton, K.W., 1990. Perspectives on reservoir limnology, in: Reservoir Limnology:
Ecological Perspectives. pp. 1–14.
Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R., Polasky, S., 2002. Agricultural
sustainability and intensive production practices. Nature.
https://doi.org/10.1038/nature01014
Tundisi, J.G., 2018. Reservoirs: New challenges for ecosystem studies and environmental
management. Water Secur 4–5, 1–7. https://doi.org/10.1016/j.wasec.2018.09.001
Utermöhl, H., 1958. Mitteilungen der Internationalen Vereinigung für Theoretische und
Angewandte. Mitteilungen der Internationalen Vereinigung für Theoretische und
Angewandte Limnologie 9, 1–38.
Vale, P., Gibbs, H., Vale, R., Christie, M., Florence, E., Munger, J., Sabaini, D., 2019. The
expansion of intensive beef farming to the Brazilian Amazon. Global Environmental
Change 57, 101922. https://doi.org/10.1016/j.gloenvcha.2019.05.006
Vasconcelos, L.P., Alves, D.C., Agostinho, A.A., Hahn, L., da Câmara, L.F., Gomes, L.C.,
2022. Fish eggs and larvae drifting through hydropower reservoirs: a case study in the
Brazilian Amazon. Hydrobiologia 849, 357–372. https://doi.org/10.1007/s10750-021-
04631-9

Vollenweider, 1968. Water management research. Scientific fundamentals of the
eutrophication of lakes and flowing waters with particular reference to nitrogen and
phosphorus as factors in eutrophication. Technical Repport DAS/CSI/68, Paris.
Winemiller, K.O., McIntyre, P.B., Castello, L., Fluet-Chouinard, E., Giarrizzo, T., Nam, S.,
Baird, I.G., Darwall, W., Lujan, N.K., Harrison, I., Stiassny, M.L.J., Silvano, R.A.M.,
Fitzgerald, D.B., Pelicice, F.M., Agostinho, A.A., Gomes, L.C., Albert, J.S., Baran, E.,
Petrere, M., Zarfl, C., Mulligan, M., Sullivan, J.P., Arantes, C.C., Sousa, L.M., Koning,
A.A., Hoeinghaus, D.J., Sabaj, M., Lundberg, J.G., Armbruster, J., Thieme, M.L., Petry,
P., Zuanon, J., Vilara, G.T., Snoeks, J., Ou, C., Rainboth, W., Pavanelli, C.S., Akama,
A., Soesbergen, A. van, Sáenz, L., 2016. Balancing hydropower and biodiversity in the
Amazon, Congo, and Mekong. Science (1979) 351, 128–129.
https://doi.org/10.1126/science.aac7082
Xiao, Y., Li, Z., Guo, J., Fang, F., Smith, V.H., 2016. Succession of phytoplankton
assemblages in response to large-scale reservoir operation: a case study in a tributary
of the Three Gorges Reservoir, China. Environ Monit Assess 188, 1–20.
https://doi.org/10.1007/s10661-016-5132-7
Zanco, B.F., Pineda, A., Bortolini, J.C., Jati, S., Rodrigues, L.C., 2017. Phytoplankton
functional groups indicators of environmental conditions in floodplain rivers and lakes of
the Paraná Basin. Acta Limnologica Brasiliensia 29, e119.
https://doi.org/10.1590/S2179-975X8117
Zanon, F.M., Pineda, A., Lansac-Tôha, F.M., Rodrigues, L.C., 2024. Impact of a run-of-river
dam in an Amazonian large river on the spatial and temporal patterns of phytoplankton
beta diversity. Biol Conserv 295, 1–12. https://doi.org/10.1016/j.biocon.2024.110656
Znachor, P., Nedoma, J., Hejzlar, J., Seďa, J., Komárková, J., Kolář, V., Mrkvička, T.,
Boukal, D.S., 2020. Changing environmental conditions underpin long-term patterns of
phytoplankton in a freshwater reservoir. Science of the Total Environment 710.
https://doi.org/10.1016/j.scitotenv.2019.135626