1   **Impact of Run-of-River Damming on Increasing Phytoplankton Biomass and Species Shift in a**
2   **Large Amazonian River: Ecological Implications and Recommendations.**

3   Alfonso Pineda[1], Beatriz Melissa Campos[2], Felipe Zanon[2], Renata Felicio[2], Luzia C.
4   Rodrigues[2,3]

[1] Programa de Pós-graduação em Biologia Comparada – PGB, Universidade Estadual de
Maringá – UEM, Maringá, PR, Brazil

[2] Programa de Pós-graduação em Ecologia de Ambientes Aquáticos Continentais – PEA,
Centro de Ciências Biológicas (CCB), Universidade Estadual de Maringá – UEM, Maringá,
PR, Brazil

[3] Núcleo de Pesquisas em Limnologia e Aquicultura – Nupélia, Centro de Ciências Biológicas
(CCB), Universidade Estadual de Maringá – UEM, Maringá, PR, Brazil

Correspondence: Alfonso Pineda, Programa de Pós-graduação em Ecologia de Ambientes
Aquáticos Continentais, Universidade Estadual de Maringá. Avenida Colombo, 5790, Bloco
H-90, Maringá, Paraná 87020-900, Brazil Amanecce@gmail.com
15

ORCID Ids

Pineda, A amanecce@gmail.com https://orcid.org/0000-0001-6902-9993 ;
Campos, BM https://orcid.org/0000-0003-1985-8528
Zanon, FM https://orcid.org/0000-0002-8098-2803
Felicio, RS https://orcid.org/0000-0001-7716-8408
Rodrigues, LC https://orcid.org/0000-0002-0593-0931

Running Head: algal biomass after damming an Amazon tributary

Author contributions
Alfonso Pineda: Conceptualization, Data Analysis, Writing- Original draft preparation.
Beatriz M. Campos: Data analysis and editing. **Felipe M. Zanon**: Data analysis and editing.
Renata S. Felicio: Data analysis and editing. **Luzia C. Rodrigues:** Supervision, Collecting
and Identifying samples, Reviewing and Editing.
28

29   **Competing interest** All authors certify that they have no affiliations with or involvement in
30   any organization or entity with any financial interest or non-financial interest in the subject
31   matter or materials discussed in this manuscript.
32

Abstract

Run-of-river (ROR) dams, often perceived as having minimal environmental impact, can induce significant hydrodynamic changes that alter aquatic ecosystems. We investigated the impacts of an ROR dam on the Madeira River, the largest Amazon tributary, focusing on phytoplankton communities and their ecological implications. Our study examined changes in biomass and species composition before and after damming as well as their drivers in both the main channel and tributaries (N=549 samples). Following dam construction, we observed a shift in the phytoplankton community from lotic diatoms to lentic phytoflagellates. This transition was likely facilitated by altered hydrodynamics and possibly influenced by the decomposition of flooded vegetation in the dam's influence zone. The decomposition of this vegetation could explain both the observed increase in oxygen consumption and the subsequent rise in phytoflagellate biomass after damming. On the other hand, despite an overall increase in phytoplankton biomass, the values remained within oligo- and mesotrophic conditions, which related with low nutrient concentrations observed. However, we caution that the dam-created hydrodynamic conditions are optimal for phytoplankton growth, potentially exacerbating nutrient-related issues in the future. We recommend proactive management strategies to prevent nutrient enrichment from agricultural and livestock activities in isolated Amazon areas affected by dams, thereby mitigating potential water quality degradation linked to phytoplankton biomass increase.

Keywords: Amazon; backwaters; cyanobacteria; damming; plankton; run-of-river dam

1 Introduction

Hydroelectric energy is considered the most environmentally friendly option for meeting global energy demand. Hydroelectric projects have been constructed even in important ecological regions. For instance, in the Amazon region, the largest basin in the world, 140 dams are functioning or under construction, and 288 are planned (Latrubesse et al., 2017). Most of the hydroelectric projects built nowadays are run-of-river (ROR) structures, which in theory flood smaller areas than traditional storage reservoirs. However, even hydroelectric projects with a ROR dam can impact environmental conditions (Almeida et al., 2019; Tundisi, 2018), human communities (Moran, 2020), and biodiversity (Lees et al., 2016; Vasconcelos et al., 2022).

Run-of-river dams can have little effect on the main channel where they are built. However, they relate to the apparition of lentic or semi-lentic conditions in the back-flooded tributaries (Almeida et al., 2019). In some cases, the back-flooding area can correspond to 30% of the reservoir area (Almeida et al., 2019), and the total flooding area can be 64.5% larger than initially predicted by hydroelectric project including a ROR dam (Cochrane et al., 2017). These new lake-like conditions can harbor problematic species, such as mosquitoes which are vectors of diseases (Rodrigues et al., 2017), aquatic macrophytes that cover the water surface (Thomaz et al., 1999), and invasive species that exclude native ones (Lees et al., 2016).

The change from running water to standing water can also favor algae, as a longer water residence time enhances their reproduction. Standing water conditions together with high nutrient concentrations can trigger populational explosions of algae known as blooms (O'Neil et al., 2012). In turn, high algae biomass can threaten water quality (Ferrão-Filho and Kozlowsky-Suzuki, 2011), diversity (Krztoń et al., 2019), and human health (Pouria et al., 1998) since it decreases the transparency and oxygen of the water and even increases the concentration of toxins produced by some microalgae. Moreover, high algae biomass can degrade the services that we take advantage of from aquatic ecosystems such as recreation and fishing (Huisman et al., 2018).

The formation of standing waters in the Amazonian region could not trigger algae blooms *per se*, but together with the high temperature of the region, high nutrient input, and the hydrodynamics changes promoted by the dams may favor phytoplankton growth. For instance, human activities such as agriculture and cattle ranching in the Amazon increase

phosphorous and nitrogen load in aquatic ecosystems (Tilman et al., 2002; Vale et al., 2019).

This is particularly important given the increase in this type of enterprise in the Amazon in

recent years (Carvalho et al., 2020; Gusso et al., 2017; Macedo et al., 2012; Vale et al., 2019).

Several studies have described the response of phytoplankton communities to damming

in several regions worldwide (e.g., Souza et al., 2016; Zanon et al., 2019). However, it is no

clear how the recently built and futures dams affects the spatial and temporal variation of

phytoplankton communities in critical ecological regions such as the Amazon. Moreover, there

exists a knowledge gap regarding the effect of damming on microalgae biomass of Amazonian

Rivers. It is now crucial to determine whether the environmental changes induced by

hydroelectric projects in the Amazon region are sufficient to promote the development of

phytoplankton biomass and identify the favored species. This information is essential for

alerting authorities to potential negative effects and for elucidating the appropriate management

measures necessary to prevent or mitigate impacts associated with high phytoplankton biomass

or unexpected species.

In this study, we assessed the upstream influence of the Jirau dam on the biomass and

community structure (species abundance) of phytoplankton in the Madeira River, one of the

largest tributaries of the Amazon River. We collected data in the main channel and tributaries

before and after the installation of the dam. Our objectives were to determine how damming

affected phytoplankton biomass, identify the abiotic factors influencing phytoplankton

biomass, and evaluate the impact of damming on community structure and changes in

representative species.

Since backwaters formed in tributaries after damming can exhibit lake-like conditions,

we hypothesized that *i*) post-damming conditions in tributaries would favor an increase in

phytoplankton biomass. Therefore, we expected an increase in biomass in tributaries, while no

significant changes were anticipated in the main channel. Additionally, due to the shift in

environmental conditions caused by damming, we hypothesized that *ii*) damming would alter

the community structure (species abundance). Specifically, we anticipated differences in

community structure before and after the damming, with species adapted to standing water

conditions becoming representative after damming. Finally, we examined the significance of

our findings in the context of environmental degradation resulting from human activities in the

Amazon.

2   Methods

2.1   Study area

The Madeira River basin is in the Amazon region, between Peru, Bolivia, and Brazil, draining much of that region (Guyot et al., 1996). The Madeira River is among the ten rivers with the highest discharge in the world (Latrubesse et al., 2017), it is the main tributary of the Amazon River, and contributes significantly to the phosphorus load for the entire Amazon basin (Almeida et al., 2015). The Madeira River basin has one annual cycle with four well-defined hydrological periods, with well-established duration intervals between years: 1. The high-water period, which usually occurs from February to May; 2. The receding period, occurring from June to July; 3. low water, from August to November; and 4. rising period, occurring from December to January. This region has the largest tropical forest in the world, being responsible for much of the global biodiversity in addition to the importance of temperature and precipitation in South and Central America (Agudelo et al., 2019; Aragón, 2018).

The Madeira River has two dams installed, Santo Antônio and Jirau, but other dams are in the planning and approval phases (Latrubesse et al., 2017). The construction of the Jirau dam ended in July 2012, being considered a run-of-river, located in the city of Porto Velho, Rondônia state, Brazil. The Jirau reservoir has an annual average water residence time of approximately 3.0 days (Almeida et al., 2020). It is located 1,206 km from the mouth of the Madeira River and has an installed capacity of 3.750 MW, supplying more than ten million homes(Energia Sustentável do Brasil, 2020). In the study period, 2009 to 2017, the average flow was 19.409 m³/s, reaching a maximum average of 45.649 m³/s in the high-water period and a minimum average of 4.133 m³/s in the low-water period (ANA, 2022). For the study, we cover the area upstream of the Jirau hydroelectric plant, between the coordinates 9º67'73.05" S, 65º44'10.27" O, and 9º60'45.27" S, 64º91'75.55" O (Figure 1).

Run-of-river dams have been reported to have minimal influence on river flow and creating minor flooded regions (Pracheil et al., 2016; Souza et al., 2019). Nonetheless, the water level of the Madeira River has changed significantly since the Jirau dam was built. The water level ranged between 1 m and 20 m before damming, whereas it varied between 21 m and 29 m after damming. Also, the seasonal fluctuation in the Madeira River's water level has decreased after damming (Supplementary material S1). The Madeira River's rising water level has reduced flow velocity and induced stratification in its tributaries (Fearnside, 2013). The

reservoir's extension varies between 207 km^2 during the dry season and 361.6 km^2 during the
wet season (Hartmann et al., 2023).

2.2   Data Samplings

Fifteen sampling stations were set up along the Madeira River (M2 to M5) and its
tributaries (T4 to T14 – Figure 2). Water samples for phytoplankton and nutrient concentration
assessments were collected directly using bottles at a depth of approximately 40 cm in the
middle section of the Madeira Rivers and tributaries. At the same time, we measured chemical
and physical variables that are critical to phytoplankton growth. Between 2009 and 2017,
samples were taken during the high (March to May), rising (December to February), receding
(June to August), and low (September to November) seasons. In 2010, 2011, 2016, and 2017,
the campaigns were carried out quarterly, while from 2013 to 2015 they were carried out
bimonthly. Five samplings were conducted in 2012. We had 37 sampling campaigns (N = 549)
in total. The before dam period is considered until July 2012 (177 samples), and the after dam
begins in October 2012 until continues through the remainder of the sampled periods (372
samples).

Figure 1. Map and location of sampling sites in the Jirau Hydroelectric Power Plant, in the
Madeira River.

2.3   Phytoplankton

Phytoplankton samples were fixed with 1% acetic Lugol solution. Individuals (cell,
coenobium, colonies, and filaments) were counted using an inverted microscope at 400X as
described by Utermöhl (1958). Phytoplankton density was calculated according to APHA
(American Public Health Association, 2012) and expressed as individuals per milliliter
(ind.mL^{-1}). Biomass was obtained based on biovolume (mm^3.L^{-1}), by multiplying the density
of each taxon by its corresponding volume. The volume of individual cells was calculated using
geometric models that were based on the shape of the cells (Hillebrand et al., 1999; Sun and
Liu, 2003). The entire number of registered taxa was divided into five algal groups: green algae,
cyanobacteria, diatoms, and phytoflagellates. Those groups have demonstrated sensitivity to
environmental changes making them suited for understanding how human activities affect
biodiversity and water quality (Reynolds, 2006).

2.4   Environmental data

We measured chemical and physical variables critical to the development of
phytoplankton following the standard methods outlined in APHA (American Public Health
Association, 2012). In situ measurements were conducted to assess water temperature (WT,
°C), pH, inorganic carbon (IC, mg L^{-1}), and alkalinity (Alk, mEq L^{-1}). Water transparency was
measured with a Secchi disk (m), and the euphotic zone (Zeu, m) was estimated as 2.7 times
the Secchi depth (Cole, 1994). The maximum depth (Zmax, m) was also measured, and the
Zeu:Zmax ratio was utilized as an indicator of light availability within the water column
(Jensen et al., 1994).

Using collected water samples, we measured additional water characteristics at the
sampled sites. Total iron concentrations (µg L-1) were determined following digestion using
the phenanthroline method. Soluble reactive phosphorus (SRP, P-PO$_4^{3-}$, µg L^{-1}), nitrate (N-
NO$_3^-$, µg L^{-1}), nitrite (N-NO$_2^-$, µg L^{-1}), and ammonium (N-NH$_4^+$, µg L^{-1}) concentrations were
measured post-sample filtration. Phosphate concentrations were determined using the ascorbic
acid method and spectrophotometric readings, while nitrate, nitrite, and ammonium
concentrations were obtained through the cadmium reduction method, sulfanilamide method,
and phenol method, respectively, for spectrophotometric measurements. Total dissolved
inorganic nitrogen (DIN, µg L-1) was calculated as the sum of nitrite, nitrate, and ammonium
concentrations. Reactive silica (µg L^{-1}) concentrations were determined using the molybdate
method. Biochemical oxygen demand (BOD, mg L^{-1}) was assessed following sample
incubation at a constant temperature (20 °C) and reading the dissolved oxygen concentrations

after 5 days. Chemical oxygen demand (COD, mg L^{-1}) determination involved the oxidation of

organic matter using a boiling mixture of chromic acid and sulfuric acid (potassium bichromate

in an acidic medium) (APHA, 2005).

2.5   Data analysis

We utilized Generalized Mixed Linear Models (GLMM) to investigate the effect of

damming on the environmental variables in both the main channel and tributaries. To avoid

issues with multiple pairwise tests, we included the sampling point as a random factor (Bolker

et al., 2009). We selected an error distribution based on the better fit of the models, such as

Gaussian, Beta, or Gamma. We also included thermal profiles from the main channel and

tributaries to investigate thermal stratification following damming. These thermal profiles were

limited to five sites (one in the Madeira River and four in tributaries) and were collected

between 2013 and 2017.

To verify the effect of damming on the structure of the phytoplankton community, we

utilized a Multivariate Permutation Analysis of Variance (PERMANOVA) on a matrix of

species biomass (considering the Bray-Curtis distance). To determine whether damming effect

interacted with hydrological periods, we considered the seasonality in the PERMANOVA. The

analysis was performed separately for tributaries and the main channel. A Principal Coordinate

Analysis (PCoA) was used to graphically depict the spatial and temporal ordinations.

Additionally, to verify associations between species and damming phases and regions

(tributaries and main channel), we used the indicator value (IndVal, Dufrêne and Legendre,

1997). IndVal is a quantitative metric that considers both a species relative abundance and

frequency within a given habitat. A IndVal value close to 1 indicate a stronger association of

the species with a particular condition (e.g., damming phase or habitat). In our study, it

supplements the PERMANOVA analysis since Indval allowed us to discover which species

were more favored at each condition.

We utilized GLMM to assess the influence of damming on the biomass of the entire

phytoplankton community and each of the five algal group in both the tributaries and the main

channel. When we found that damming had a considerable influence on algal biomass, we

examined into its link with seasonality. We also utilized GLMM to assess the influence of each

standardized abiotic factor on algal biomass as well as on the presence-absence of each group.

In this case, we modeled both biomass and the presence-absence of algal groups since several

samples lacked a specific algae group, resulting in biomass values of zero (absence). We chose
the Gamma family as the error distribution for modeling biomass and the zero-inflated
distribution for occurrence data. To avoid issues with multiple testing, we incorporated the
sampling point as a random effect in all models, as advised by Bolker et al (Bolker et al., 2009).
Visual examination validated the linearity assumption. The models' findings were given in
ANOVA style.

All analyzes were conducted utilizing the R program (R Development Core Team,
2023). Vegan package (Oksanen et al., 2007) was employed to perform PERMANOVA (adonis
function) and to compute distance matrices (vegdist function). PCoA was performed with hnp
package (Moral et al., 2017), GLMM (glmmTMB function) with glmmTMB package (Brooks,
Mollie et al., 2017), and plots with ggplot2 package (Wickham, 2016).

3 RESULTS

3.1 Changes in environmental conditions

Both the main channel and tributaries exhibited changes in the environment after
damming, but changes were more pronounced in the tributaries (Figures 2 and 3). An increase
in chemical oxygen demand was observed in both environments. The main channel showed a
reduction in alkalinity, inorganic carbon, silica, conductivity, and pH, whereas tributaries
displayed a decrease in light availability (Zeu and Zeu:Zmax) and iron, but an increase in depth,
conductivity, inorganic carbon, temperature, and alkalinity. The variation of most
environmental factors seemed to increase with damming at both the main channel and
tributaries (Table 1), except for orthophosphate, BOD, and COD which decreased.

Figure 2. The effect of damming on the variation of environmental variables measured at the tributaries of the Madeira River. Asterisks (*) in the figure indicate a significant effect of damming. The vertical axis displays the log-transformed values for each variable, except for pH and the ratio of Zeu:Zmax. *IC* – dissolved inorganic carbon; *BOD* – biochemical oxygen demand; *COD* – chemical oxygen demand; *DIN* – dissolved inorganic nitrogen; $Z_{eu}:Z_{max}$ – light availability in the water column; *COND* – Conductivity; *Alk* – alkalinity; *Zeu* – euphotic zone; *SRP* – soluble reactive phosphorous; *SI* - Silica.

Figure 3. The effect of damming on the variation of environmental variables measured at Madeira River. Asterisks (*) in the figure indicate a significant effect of damming. The vertical axis displays the log-transformed values for each variable, except for pH and the ratio of Zeu:Zmax. The boxes in the plot represent the first and third quartiles, the horizontal bar represents the median, and the vertical bars indicate the minimum and maximum values, excluding any outliers, which are represented as individual points. IC – dissolved inorganic carbon; BOD – biochemical oxygen demand; COD – chemical oxygen demand; DIN – dissolved inorganic nitrogen; Zeu:Zmax – light availability in the water column; COND – Conductivity; Alk – alkalinity; Zeu – euphotic zone; SRP – soluble reactive phosphorous; SI - Silica.

Table 1: Mean and coefficient of variation (CV, %) of abiotic variables in the main channel and tributaries during the construction phases of the dam.

	Madeira		Tributaries	
	Before	**After**	**Before**	**After**
Temperature (°C)	27.91 (5.68)	28.04 (6)	26.61 (7.98)	27.76 (7.32)
pH	7 (7.7)	6.41 (10.88)	6.18 (11.84)	6.06 (13.06)
Conductivity (μS.cm^{-1})	88.9 (32.17)	66.35 (52.98)	10.31 (75.87)	25.54 (123.92)
$Z_{eu}:Z_{max}$ (m)	0.03 (141.07)	0.06 (184.55)	0.59 (51.73)	0.36 (85.52)
Euphotic zone - Z_{eu} (m)	0.22 (66.38)	0.57 (136.02)	2.09 (59.31)	1.63 (73.58)
Orthophosphate (μg L^{-1})	0.09 (162.19)	0.07 (85.63)	0.01 (370.74)	0.02 (159.5)
Dissolved inorganic nitrogen (μg L^{-1})	1.74 (52.95)	1.89 (62.93)	0.97 (61.4)	1.13 (73.73)
Biochemical oxygen demand (mg L^{-1})	1.18 (66.06)	1.28 (56.22)	1.45 (66.4)	1.39 (61.85)
Chemical oxygen demand (mg L^{-1})	10.11 (74.71)	12.69 (63.76)	9.47 (71.57)	14.48 (65.46)
Iron (μg L^{-1})	5.05 (80.42)	1.39 (80.54)	1.25 (71.22)	1.03 (61.62)
Alkalinity (mEqL^{-1})	28.22 (34.25)	23.19 (45.47)	6.71 (58.59)	10.63 (72.73)
Inorganic carbon (mg L^{-1})	7.31 (36.15)	6.31 (37.72)	2.76 (71.73)	4.53 (38.89)
Silica (μg L^{-1})	22.55 (107.92)	9.28 (73.69)	6.94 (43.68)	7.94 (57.16)

WT - water temperature - °C; pH; IC – dissolved inorganic carbon; BOD - biochemical oxygen demand; COD - chemical oxygen demand; DIN – dissolved inorganic nitrogen; $Z_{eu}:Z_{max}$ – light availability in the water column , SRP – soluble reactive phosphorus, Si - Silica.

3.2 *The effect of damming and seasonality on phytoplankton structure and biomass*

In this study, we identified a total of 525 taxa, with the highest number of species belonging to green algae (236 species, mean of 0.095 ind.mL^{-1}), Phytoflagellates comprised 118 species (mean of 0.145 ind.mL^{-1}), followed by diatoms (86 species, mean of 0.057 ind.mL^{-1}). Cyanobacteria accounted for 72 species (mean of 0.017 ind.mL^{-1}), and xantophyceans were represented by 13 species (mean of 0.0004 ind.mL^{-1}).

Before and after damming, the main channel recorded 119 and 167 species,
respectively, with diatoms (mainly bacillariophyceans) exhibiting the highest richness both
before (30) and after (37) damming. In the tributaries, the number of species increased from
307 before damming to 457 after damming, with zygnemaphyceans having the greatest
richness both before (62) and after (94) damming.

PERMANOVA revealed a significant effect of damming on the phytoplankton
structure for the main channel (F = 1.87, p = 0.01) and tributaries (F = 12.89, p <0.01). We also
verified a significant effect the hydrological periods (for main channel, F = 1.22, p = 0.001; for
tributaries, F = 2.09, p <0.01). Both the PERMANOVA result, and their graphic representation
(Figure 4) showed that damming change communities and increased the spatial and temporal
variation of species biomass in both the main channel and tributaries at all the hydrological
periods.

295
Figure 4: Temporal variation of phytoplankton community (species abundance) related to
damming phases and hydrological periods at both the main channel and tributaries. In all cases,
the community's heterogeneity was higher in the after damming phase. The higher variation
seemed to be related to the low water period.

IndVal analysis indicated that the number of species linked with tributaries increased
after damming, while indicators of the Madeira River decreased. Diatoms, which are related to
flowing waters, were found to be exclusive indicators of pre-damming conditions for both
tributaries and the main channel. It is noteworthy that most of the indicator species associated

with tributaries in the post-damming phase were indicative of standing water, specifically
phytoflagellates (Table 2).

Table 2. Species as indicators of the conditions at each damming phase (before and after) and
system (tributaries and Madeira River). *IndVal*, indicator value based on the relative frequency
and relative average abundance in the clusters Phase – system. The table shows only those
species with a high probability of being an indicator ($p < 0.05$). *Freq*, specie frequency at each
cluster. The taxonomic groups were colored differently to highlight the representativity of each
group at each phase – system: blue – Cyanobacteria; pink – Diatoms; green – Green algae;
golden – Phytoflagellates.

Phase - system	Indval	p	Freq	Taxa	G
Before - Tributaries	0.05	0.013	8	*Lemmermanniella flexa* Hindák	
	0.04	0.011	7	*Synechocystis aquatilis* Sauv.	
	0.04	0.044	12	*Pseudanabaena mucicola* (Hüb.-Pest. e W. Naumann) Bourr.	
	0.03	0.03	4	*Snowella* sp.	
	0.05	0.011	11	*Cymbopleura naviculiformis* (Auersw.) Krammer	
	0.04	0.043	12	Pennales sp1	
	0.04	0.028	12	*Pinnularia microstauron* (Ehrenb.) Cleve	
	0.1	0.008	43	*Oocystis lacustris* Chodat	
	0.09	0.031	39	*Monoraphidium minutum* (Naegeli) Komárk. - Legn.	
	0.05	0.016	11	*Cosmarium punctulatum* Bréb.	
	0.04	0.028	14	*Scenedesmus acunae* Com.	
	0.33	0.001	91	*Cryptomonas* sp.	
Before - Madeira	0.03	0.024	4	*Aphanizomenon gracile* Lemmerm.	
	0.22	0.001	99	*Fragilaria* sp.	
	0.19	0.006	111	*Aulacoseira granulata* var. *granulata* (Ehrenb.) Sim.	
	0.17	0.001	43	*Cyclotella meneghiniana* Kütz.	
	0.17	0.001	42	*Aulacoseira distans* (Ehrenb.) Sim.	
	0.17	0.001	60	*Aulacoseira granulata* var. *angustissima* (O. Müller) Sim.	
	0.13	0.001	23	*Ulnaria ulna* (Nitzsch.) Comp.	
	0.12	0.003	31	*Nitzschia* sp.	
	0.08	0.001	7	*Gyrosygma* sp1	
	0.08	0.002	9	*Cocconeis* sp.	
	0.06	0.008	19	*Aulacoseira ambigua* (Grunow) Sim.	
	0.06	0.043	26	*Navicula viridula* (Kütz.) Ehrenb.	

Phase - system	Indval	p	Freq	Taxa	G
	0.05	0.025	16	*Achnanthidium minutissimum* (Kütz.) Czarn.	
	0.04	0.015	7	*Melosira varians* Agardh	
	0.05	0.007	5	*Actinastrum aciculare* Playf.	
	0.05	0.027	14	*Desmodesmus intermedius* var. *acutispinus* (Roll) E. Hegew.	
	0.04	0.005	2	*Monactinus simplex* (Meyen) Corda	
	0.04	0.05	16	*Scenedesmus* sp.	
	0.04	0.03	10	*Scenedesmus acuminatus* (Lagerh.) Chodat	
After - Tributaries	0.13	0.002	39	*Eunotia bilunaris* (Ehrenb.) Schaar.	
	0.07	0.015	29	*Urosolenia eriensis* (H. L. Sm.) Round e Craw.	
	0.2	0.002	107	*Scenedesmus ecornis* (Ehrenb.) Chodat	
	0.19	0.001	95	*Monoraphidium griffithii* (Berk.) Komárk.-Legn.	
	0.16	0.006	59	*Crucigenia tetrapedia* (Kirch.) W. e G.S. West	
	0.13	0.005	57	*Monoraphidium irregulare* (G. M. Smith) Komárk.-Legn.	
	0.11	0.016	62	*Closteriopsis* sp.	
	0.11	0.015	64	*Monoraphidium convolutum* (Corda) Komárk. - Legn.	
	0.1	0.005	29	*Crucigenia fenestrata* (Schm.) Schm.	
	0.1	0.016	46	*Desmodesmus hystrix (*Lagerh.) E. Hegew.	
	0.09	0.045	76	*Monoraphidium komarkovae* Nygaard	
	0.09	0.005	26	*Desmodesmus brasiliensis* (Bohlin) E. Hegew.	
	0.08	0.002	23	*Nephrochlamys* sp.	
	0.07	0.024	37	*Ankyra judayi* (G.W. Smith) Fott	
	0.06	0.025	23	*Willea rectangularis* (Nägeli) Komárek	
	0.06	0.023	18	*Monoraphidium circinale* (Nygaard) Nygaard	
	0.06	0.017	19	*Cosmarium regnesi* Reins.	
	0.05	0.032	14	Chlorococcales colonial não identificada 1	
	0.47	0.001	168	*Cryptomonas brasiliensis* Castro, Bic. e Bic.	
	0.29	0.001	97	*Cryptomonas erosa* Ehrenb.	
	0.25	0.001	120	*Trachelomonas volvocinopsis* Svirenko	
	0.24	0.001	115	*Peridinium* sp.	
	0.22	0.002	75	*Gonyostomum* sp.	
	0.2	0.001	108	*Chlamydomonas* sp.	
	0.18	0.019	94	*Euglena* sp.	
	0.18	0.001	55	*Trachelomonas oblonga* Lemmerm.	
	0.16	0.004	65	*Lepocinclis ovum* (Ehrenb.) Lemmerm.	
	0.15	0.007	86	*Mallomonas* sp.	
	0.09	0.046	54	*Cryptomonas curvata* Ehrenb. emend. Pen.	
	0.09	0.016	43	*Euglena acus* Ehrenb.	

Phase - system	Indval	p	Freq	Taxa	G
	0.08	0.048	47	*Trachelomonas volvocina* Ehrenb.	
	0.07	0.009	19	*Strombomonas scabra* (Playf.) Tell e Conf.	
	0.06	0.014	20	*Chroomonas* sp.	
	0.05	0.033	20	*Trachelomonas lacustris* Dezep.	
	0.05	0.026	17	*Phacus* sp.	
	0.05	0.026	18	*Phacus acutus* Pochmann	
	0.04	0.049	10	*Phacus horridus* Pochm.	
	0.04	0.045	10	*Pandorina morum* (F. Muller) Bory	
	0.03	0.048	11	*Trachelomonas similis* Stokes var. *spinosa* Hub.-Pest.	
After - Madeira	0.02	0.047	2	*Pediastrum duplex* Meyen	

In the tributaries, the total biovolume ranged from 0.0001 to 4.74 mm^{3}L^{-1} before

damming. After damming values varied between 0.0001 and 15 mm^{3}L^{-1}. Most of the samples

(98% before and 96% after damming) showed a biovolume lower than 2 mm^{3}L^{-1}. After

damming, we observed biovolumes higher than 2 mm^{3}L^{-1} (four samples), 3 mm^{3}L^{-1} (four

samples), 5 mm^{3}L^{-1} (1 sample), and 10 mm^{3}L^{-1} (one sample). In the Madeira River, the total

biovolume ranged between 0.001 and 3.38 mm^{3}L^{-1} before damming, and after damming, values

varied between 0.0001 and 0.97 mm^{3}L^{-1}. Biovolume was less than 2 mm^{3}L^{-1} in 98% of the

samples before damming and 100% after damming. According to Vollenweider (1968), the

registered biomass conditions mainly ranged from oligotrophic (biovolume = 1 – 5 mm^{3}L^{-1}) to

mesotrophic (biovolume = 3 – 5 mm^{3}L^{-1}).

GLMM revealed differences in total biomass between phases in both the main channel

and tributaries (Figure 5A and Table S2). In the tributaries, total and phytoflagellates biomass

increased (Figure 5A), with variations between phases depending on the season considered.

Specifically, no differences were observed during the rising season for total biomass and at

high water for phytoflagellates, while cyanobacteria showed differences only in the high and

receding water seasons (Figure 5B and Table S3). In the main channel, total biomass decreased,

along with diatoms and green algae biomass, while phytoflagellates increased (Figure 5A).

Differences between the phases were consistent across season for phytoflagellates. In contrast,

differences between the phases at the low- and rising-water seasons were observed for the total

biomass and diatoms, at the rising- and high-water seasons for greens, and at the rising waters

for cyanobacteria (Figure 5B).

Figure 5: A. Difference in the distribution of total biomass and for each group in the tributaries and main channel during the construction phases of the dam. Asterisks represent significant differences. B. Differences in the biomass distribution of the total community and of each group during the hydrological periods and construction phases of the dam. Asterisks indicate significant differences in the biomass values obtained.

3.3   Influence of abiotic variables on community biomass

The GLMM also revealed that the factors driving total biomass differed between the main channel and its tributaries. (Figure 6 Table S5). Moreover, the significance of these factors varied across taxonomic groups, depending on whether the occurrence or the biomass data were considered (Figure 6).

Figure 6: Influence of abiotic variables on total biomass and on the biomass and occurrence of algae groups from the main channel and tributaries. Asterisks represent a significant effect. Biomass is shown by blue bars while occurrence is shown by red bars.

4   Discussion

Several dams will occupy rivers in the Amazon region in the coming decades. Understanding their environmental implications and the subsequent effect on organisms is crucial for developing effective management and conservation strategies, as for anticipating and mitigating potential future. In this study, we demonstrated that the Jirau dam, as expected, induced changes in the biomass and structure of the phytoplankton communities in both the main channel and tributaries, corroborating our hypotheses. Specifically, damming promoted increased phytoplankton biomass in the tributaries, driven by the proliferation of species adapted to lacustrine, particularly phytoflagellates. Our findings highlight that while hydroelectric projects with Run-of-River (ROR) dams have historically been associated with minimal environmental impact, they can still significantly alter biotic characteristics, even in main channels where ROR dams are presumed to have less impact. This section delves into the

impacts of damming on phytoplankton community dynamics and total biomass. We also

discuss broader implications, including the potential future degradation of the Amazonian

region, and how our results contribute to preventing undesirable scenarios.

4.1 Changes in phytoplankton after damming.

The construction of the Jirau dam influenced temporal and spatial changes in the

phytoplankton communities in the main channel and its tributaries. Despite the hydrological

season, the spatial variation in communities appeared to increase after damming. This contrasts

with previous research in the same study zone, which showed that biotic variation decreased

after damming (Pineda et al., 2024; Zanon et al., 2024). Unlike that research, we analyzed the

community variation while removing the effect of seasonality. Our results, together with the

mentioned research, suggest that after damming, there could be an increase in spatial variation

while temporal variation decreases. However, the spatial variation does not seem to be enough

to compensate for the loss of temporal variation since the general pattern exhibit reduction in

heterogeneity (Pineda et al., 2024).

The increase in spatial variation that we observed can be attributed to the mosaic of

habitats with varying features that resulted after damming. Following damming, as in our case,

a region impacted by run-of-river projects shows not just rivers but also habitats with lake-like

properties because of backwaters formed at the tributary's junction with the main channel

(Almeida et al., 2019). Each one of those habitats can then favor different species with different

adaptations (Reynolds et al., 2002; Rodrigues et al., 2018; Zanco et al., 2017) and increase the

spatial variation in communities. Our results indicated that after damming the number of

species associated with tributaries increased by 300%. It means that the characteristics of

tributaries favored different species than those favored by the main channel. Furthermore, most

of the species identified as tributary indicators after damming are typical of standing waters

(Supplementary material), whereas most indicator species at the main channel were associated

with lotic environments. It is consistent with our evidence (Supplementary material – Figure

S1) and from predictions from other studies (e.g., Fearnside, 2013) about the stratification of

tributaries after damming.

In the case of the total biomass, it decreased in the main channel (which was

unexpected) and increased in the tributaries (as expected). We believe that the reduction of

biomass in the main channel was caused by a decrease in phytoplankton export from tributaries,

as the dam in the Madeira River increased the water level and reduced the water flow in the

tributaries. However, more research is necessary to study the effect of damming on the

dispersion between tributaries and the main channel. In any case, our result contrasts with

evidence showing that run-of-river projects have little effect on large rivers (Almeida et al.,

2019), and suggests that this kind of project may affect the flux of matter and energy in dammed

rivers by decreasing the biomass of primary producers in the main channel.

We attributed the biomass increase in the tributaries to a reduction in their water flow,

which increase water residence time, promotes stratification, and consequently favors

phytoplankton development. As support for that assumption, we demonstrated an increase in

species associated with standing waters as well as evidence of water stratification at certain

tributaries after damming (Supplementary material – Figure S1). Additionally, we observed

that water temperature had a positive effect on total biomass, and that temperature increased

after damming, which may relate to stratification. Moreover, the stratification in tributaries in

Madeira River has been documented after the construction of the Santo Antônio hydropower

project (Almeida et al., 2019), which is also a large ROR located downstream of the Jirau dam.

Unfortunately, we did not have stratification data for each sampling to include in our biomass

models. Nevertheless, our observations suggest that even a run-of-river dam can lead to

changes in the hydrodynamics of affected basins, mainly at tributaries, and that these changes

can increase the phytoplankton biomass in tributaries.

Although total biomass increased at the tributaries, not all the phytoplankters reacted to

damming in the same way. While the biomass of flagellates increased, that of cyanobacteria

decreased. We believe that while lentic characteristics appeared to favor the whole

phytoplankton community, chemical factors appeared to determine which group might thrive

best. For instance, we found that an increase in nitrogen concentration following damming had

a negative effect on cyanobacteria growth but favored green algae. In another case, high

phosphate concentrations boosted cyanobacteria while harming phytoflagellates.

Phytoflagellates dominated in the biovolume values. Its dominance could be related to

the increase of organic material accumulated in the back-waters regions at the tributaries. The

increase by 10 meters in average of the water level of the Madeira River covered more than

160 Km^2 of natural forest (Cochrane et al., 2017). We observed that chemical oxygen demand

(COD) increased after damming in tributaries, indicating that the process of degradation of

organic matter consumed more oxygen with damming. In this way, organisms with the ability

of take advantage of a high amount of organic matter could domain. As changes in water

features influence all aquatic species, these alterations can have an impact on the entire trophic

web. For instance, a decrease in oxygen can relate to the death of fish (Demertzioglou et al.,

2022) and the favoring of invasive species more tolerant to those new environmental

conditions, such as some macrophyte species (Harrow-Lyle and Kirkwood, 2021).

The increase in algal biomass that we have identified here may not be restricted to our

study zone. A comparable rise in phytoplankton biomass in backwater was registered in the

influence zone of the dam of the Santo Antônio dam (Appel, 2017). Again, damming caused

the formation of standing waters in that location (Almeida et al., 2019), which likely supported

algal development as in our case. Furthermore, an increase in phytoplankton biomass may be

predicted through the amazon, since the Brazilian government plans to expand hydroelectric

power generation in the Brazilian Amazon over the next 25 years (Nobre et al., 2016) with the

construction of thirty large dams (Empresa de Pesquisa Energética, 2013) and the flooding of

~12,000 km^2 (Nobre et al., 2016).

4.2   What our findings mean for the future and how we can work to prevent

Our study revealed that the creation of backwaters post-damming in tributaries

appeared to favor an increase in algal biomass, resembling lake-like conditions. Despite this

increase, the biomass values remained low, indicative of oligotrophic to mesotrophic

conditions (Vollenweider, 1968). This limited biomass was primarily limited by low phosphorus

availability (less than 0.001µg.L^{-1}). Our findings suggest that while standing water conditions

created by the Jirau dam enhanced hydrodynamic conditions for phytoplankton growth, current

nutrient levels are insufficient to support high and problematic phytoplankton biomass, at least

in the present nutrient conditions.

However, we caution that the optimal hydrodynamic conditions created by the dam,

such as increased residence time and stratification in tributaries, could potentially synergize

with future nutrient increases in Amazonian rivers. Studies (e.g., dos Santos et al., 2014)

indicate that nutrient enrichment could lead to elevated biomass levels, like issues observed in

other dam-affected regions worldwide (Grill et al., 2019; Winemiller et al., 2016).

Human activities associated with nutrient enrichment in water have reached areas of

the Amazon that would have been unimaginable a few decades ago. For example, deforestation

for pastures has been linked to phosphorus rise in rivers like the Ji-Paraná River in the Amazon

basin (Ballester et al., 2003). Furthermore, projections suggest that by 2050, pastures and

agricultural expansion could cover 40% of the entire Amazon region (Soares-Filho et al., 2006),

potentially increasing nutrient loads to waterways. Urban expansion facilitated by dams and

agro-industrial operations has also contributed to nutrient increases in Amazonian rivers due

to untreated sewage (Ferreira et al., 2021).

Given these observations, we urge authorities to implement stringent protocols for

nutrient management in future agricultural and urban projects in dam-influenced zones of the

Amazon. Proper management practices can mitigate the risks associated with phytoplankton

blooms, preserving water quality and the ecosystem services derived from aquatic

environments, including water supply, recreation, and fishing. This proactive approach to

nutrient management is essential to safeguarding the ecological integrity of Amazonian rivers.

Conclusion

Even run-of-the-river hydroelectric projects like the Jirau hydroelectric plant can alter

water chemistry and hydrodynamics, subsequently affecting the characteristics of microbial

communities such structure and biomass of phytoplankton. This impact could potentially

extend across the Amazon region, as several similar hydropower projects are currently under

construction in the area. Given that the flooding of large swathes of vegetation is a common

consequence of these projects, it is anticipated that dams in the Amazon region will lead to an

increase in mixotrophic organisms such as flagellates, which take advantage of organic matter.

Additionally, the dams are expected to elevate the chemical demand for oxygen to degrade the

accumulated organic matter, thereby reducing oxygen availability. While damming does

promote the biomass of phytoplankton due to extended residence times and water column

stratification, it may not suffice to produce problematic levels of biomass due to limited nutrient

availability. Therefore, authorities must remain vigilant regarding activities that may

exacerbate these issues and impact water quality in the Amazon. This includes monitoring

practices that could result in heightened nutrient loading in aquatic habitats, such as agricultural

practices, urban development, industrial activities, and sewage discharge.

Acknowledgments

We thank the Programa de Pós-graduação em Ecologia de Ambientes Aquáticos Continentais and the Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura (Nupélia) from the Universidade Estadual de Maringá (UEM) for supplying infrastructure. We thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior— Brazil (CAPES) for the scholarship granted to BMC, FMZ and RFS. AP thanks Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for postdoctoral fellowships. LCR was supported by productivity fellowships from CNPq. The authors state that they have no conflicts of interest relating to this study.

Declarations

Data availability The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request

Funding AP was awarded a scholarship from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). BMC, FMZ and RFS were awarded scholarship from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior— Brazil (CAPES)

Consent to participate Not applicable.

Consent to publication Not applicable

Ethical approval Not applicable. The present study did not involve research with humans or animals

References

Agudelo, J., Arias, P.A., Vieira, S.C., Martínez, J.A., 2019. Influence of longer dry seasons in the Southern Amazon on patterns of water vapor transport over northern South America and the Caribbean. Clim Dyn 52, 2647–2665. https://doi.org/10.1007/s00382-018-4285-1

Almeida, R.M., Hamilton, S.K., Rosi, E.J., Arantes, J.D., Barros, N., Boemer, G., Gripp, A., Huszar, V.L.M., Junger, P.C., Lima, M., Pacheco, F., Carvalho, D., Reisinger, A.J., Silva, L.H.S., Roland, F., 2019. Limnological effects of a large Amazonian run-of-river dam on the main river and drowned tributary valleys. Sci Rep 9. https://doi.org/10.1038/s41598-019-53060-1

Almeida, R.M., Hamilton, S.K., Rosi, E.J., Barros, N., Doria, C.R.C., Flecker, A.S., Fleischmann, A.S., Reisinger, A.J., Roland, F., 2020. Hydropeaking operations of two run-of-river mega-dams alter downstream hydrology of the largest Amazon tributary. Front Environ Sci 8, 120. https://doi.org/10.3389/fenvs.2020.00120

Almeida, R.M., Tranvik, L., Huszar, V.L.M., Sobek, S., Mendonça, R., Barros, N., Boemer, G., Durval, J., Arantes Jr, J.D., Roland, F., 2015. Phosphorus transport by the largest Amazon tributary (Madeira River, Brazil) and its sensitivity to precipitation and damming [WWW Document]. Inland Waters. https://doi.org/10.5268/IW-5.3.815

American Public Health Association, 2012. Standard Methods for the Examination of Water and
Wastewater, 21st ed, Standard Methods. American Public Health Association, American Water
Works Association, and Water Environment Federation, Washington, DC.

ANA, 2022. Brasil - Agência Nacional de Água [WWW Document].

Appel, E., 2017. Efeitos do barramento tipo fio d'água sobre as comunidades fitoplanctônicas do rio
Madeira e seus tributários na área de influência da hidrelétrica Santo Antônio, RO, Brasil.
Universidade Federal do Rio de Janeiro.

Aragón, L.E., 2018. A dimensão internacional da Amazônia: um aporte para sua interpretação.
Revista NERA 1, 14–33.

Ballester, M.V.R., Victoria, D.D.C., Krusche, A. V., Coburn, R., Victoria, R.L., Richey, J.E.,
Logsdon, M.G., Mayorga, E., Matricardi, E., 2003. A remote sensing/GIS-based physical
template to understand the biogeochemistry of the Ji-Paraná river basin (Western Amazônia).
Remote Sens Environ 87, 429–445. https://doi.org/10.1016/j.rse.2002.10.001

Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H., White, J.S.S.,
2009. Generalized linear mixed models: a practical guide for ecology and evolution. Trends Ecol
Evol 24, 127–135. https://doi.org/10.1016/j.tree.2008.10.008

Brooks, Mollie, E., Kristensen, K., Benthem, Koen, J., V., Magnusson, A., Berg, Casper, W., Nielsen,
A., Skaug, Hans, J., Mächler, M., Bolker, Benjamin, M., 2017. glmmTMB Balances Speed and
Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. R J 9, 378.
https://doi.org/10.32614/RJ-2017-066

Carvalho, R., de Aguiar, A.P.D., Amaral, S., 2020. Diversity of cattle raising systems and its effects
over forest regrowth in a core region of cattle production in the Brazilian Amazon. Reg Environ
Change 20, 1–15. https://doi.org/10.1007/s10113-020-01626-5

Cochrane, S.M.V., Matricardi, E.A.T., Numata, I., Lefebvre, P.A., 2017. Landsat-based analysis of
mega dam flooding impacts in the Amazon compared to associated environmental impact
assessments: Upper Madeira River example 2006–2015. Remote Sens Appl 7, 1–8.
https://doi.org/10.1016/j.rsase.2017.04.005

Cole, A.G., 1994. Textbook of Limnology, 4th Ed. ed. Waveland Press, Inc. Prospect Heights,
Illinois.

Demertzioglou, M., Genitsaris, S., Mazaris, A.D., Kyparissis, A., Voutsa, D., Kozari, A., Kormas,
K.A., Stefanidou, N., Katsiapi, M., Michaloudi, E., Moustaka-Gouni, M., 2022. A catastrophic
change in a european protected wetland: From harmful phytoplankton blooms to fish and bird
kill. Environmental Pollution 312, 120038. https://doi.org/10.1016/j.envpol.2022.120038

dos Santos, J.C.N., de Andrade, E.M., de Araújo Neto, J.R., Meireles, A.C.M., de Queiroz Palácio,
H.A., 2014. Land use and trophic state dynamics in a tropical semi-arid reservoir. Revista
Ciencia Agronomica 45, 35–44. https://doi.org/10.1590/s1806-66902014000100005

Dufrêne, M., Legendre, P., 1997. Species Assemblages and Indicator Species : the Need for a Flexible
Asymmetrical Approach. Ecol Monogr 67, 345–366.

Empresa de Pesquisa Energética, 2013. Plano Decenal de Expansão de Energia 2013– 2022. Empresa
Pesquisa Energética, Rio de Janeiro.

Energia Sustentável do Brasil, 2020. Dados técnicos relacionados à Usina Hidrelétrica de Jirau.
Retrieved [WWW Document].

Fearnside, P.M., 2013. Credit for climate mitigation by Amazonian dams: loopholes and impacts
illustrated by Brazil's Jirau Hydroelectric Project. Carbon Manag 4, 681–696.
https://doi.org/10.4155/cmt.13.57

Ferrão-Filho, A.D.S., Kozlowsky-Suzuki, B., 2011. Cyanotoxins: Bioaccumulation and effects on
aquatic animals. Mar Drugs. https://doi.org/10.3390/md9122729

Ferreira, S.J.F., Pinel, S., Ríos-Villamizar, E.A., Miranda, S.Á.F., Pascoaloto, D., Vital, A.R.T.,
Monteiro, M.T.F., da Silva, M. do S.R., da Cunha, T.R.B., dos Santos, A.S., Bender, S., da
Cunha, H.B., 2021. Impact of rapid urbanization on stream water quality in the Brazilian
Amazon. Environ Earth Sci 80, 316. https://doi.org/10.1007/s12665-021-09621-7

Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., Babu, S., Borrelli, P.,
Cheng, L., Crochetiere, H., Ehalt Macedo, H., Filgueiras, R., Goichot, M., Higgins, J., Hogan,
Z., Lip, B., McClain, M.E., Meng, J., Mulligan, M., Nilsson, C., Olden, J.D., Opperman, J.J.,
Petry, P., Reidy Liermann, C., Sáenz, L., Salinas-Rodríguez, S., Schelle, P., Schmitt, R.J.P.,
Snider, J., Tan, F., Tockner, K., Valdujo, P.H., van Soesbergen, A., Zarfl, C., 2019. Mapping the
world's free-flowing rivers. Nature 569, 215–221. https://doi.org/10.1038/s41586-019-1111-9

Gusso, A., Ducati, J.R., Bortolotto, V.C., 2017. Analysis of soybean cropland expansion in the
southern Brazilian Amazon and its relation to economic drivers. Acta Amazon 47, 281–292.
https://doi.org/10.1590/1809-4392201700543

Guyot, J.L., Filizola, N., Quintanilla, J., Cortez, J., 1996. Dissolved solids and suspended sediment
yields in the Rio Madeira basin, from the Bolivian Andes to the Amazon, in: Erosion and
Sediment Yield: Global and Regional Perspectives. IAHS Publication, Exeter, pp. 55–63.

Harrow-Lyle, T.J., Kirkwood, A.E., 2021. Low Benthic Oxygen and High Internal Phosphorus-
Loading are Strongly Associated With the Invasive Macrophyte Nitellopsis obtusa (starry
stonewort) in a Large, Polymictic Lake. Front Environ Sci 9, 357.
https://doi.org/10.3389/FENVS.2021.735509/BIBTEX

Hartmann, J., Fonseca, A., Trias, M., Mandilovich, V., 2023. Jirau Hydropower Plant: Assessment
Report. London.

Hillebrand, H., Dürselen, C.-D., Kirschtel, D., Pollingher, U., Zohary, T., 1999. Biovolume
Calculation for Pelagic and Benthic Microalgae. J Phycol 35, 403–424.
https://doi.org/10.1046/j.1529-8817.1999.3520403.x

Huisman, J., Codd, G.A., Paerl, H.W., Ibelings, B.W., Verspagen, J.M.H., Visser, P.M., 2018.
Cyanobacterial blooms. Nat Rev Microbiol 16, 471–483. https://doi.org/10.1038/s41579-018-
0040-1

Jensen, J.P., Jeppesen, E., Olrik, K., Kristensen, P., 1994. Impact of nutrients and physical factors on
the shift from cyanobacterial to chlorophyte dominance in shallow Danish lakes. Canadian
Journal of Fisheries and Aquatic Sciences 51, 1692–1699.

Krztoń, W., Kosiba, J., Pociecha, A., Wilk-Woźniak, E., 2019. The effect of cyanobacterial blooms on
bio- and functional diversity of zooplankton communities. Biodivers Conserv 28, 1815–1835.
https://doi.org/10.1007/s10531-019-01758-z

Latrubesse, E.M., Arima, E.Y., Dunne, T., Park, E., Baker, V.R., D'Horta, F.M., Wight, C.,
Wittmann, F., Zuanon, J., Baker, P.A., Ribas, C.C., Norgaard, R.B., Filizola, N., Ansar, A.,
Flyvbjerg, B., Stevaux, J.C., 2017. Damming the rivers of the Amazon basin. Nature 546, 363–
369. https://doi.org/10.1038/nature22333

Lees, A.C., Peres, C.A., Fearnside, P.M., Schneider, M., Zuanon, J.A.S., 2016. Hydropower and the
future of Amazonian biodiversity. Biodivers Conserv 25, 451–466.
https://doi.org/10.1007/s10531-016-1072-3

Macedo, M.N., DeFries, R.S., Morton, D.C., Stickler, C.M., Galford, G.L., Shimabukuro, Y.E., 2012.
Decoupling of deforestation and soy production in the southern Amazon during the late 2000s.
Proc Natl Acad Sci U S A 109, 1341–1346. https://doi.org/10.1073/pnas.1111374109

Moral, R.A., Hinde, J., Demétrio, C.G.B., 2017. Half-Normal Plots and Overdispersed Models in R :
The hnp Package. J Stat Softw 81. https://doi.org/10.18637/jss.v081.i10

Moran, E.F., 2020. Changing how we build hydropower infrastructure for the common good: lessons
from the Brazilian Amazon. Civitas 20, 5–15. https://doi.org/10.15448/1984-7289.2020.1.34643

Nobre, C.A., Sampaio, G., Borma, L.S., Castilla-Rubio, J.C., Silva, J.S., Cardoso, M., 2016. Land-use
and climate change risks in the amazon and the need of a novel sustainable development
paradigm. Proc Natl Acad Sci U S A 113, 10759–10768.
https://doi.org/10.1073/pnas.1605516113

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J., Suggests, M.,
2007. The vegan package. Community ecology package.

O'Neil, J.M., Davis, T.W., Burford, M.A., Gobler, C.J., 2012. The rise of harmful cyanobacteria
blooms: The potential roles of eutrophication and climate change. Harmful Algae 14, 313–334.
https://doi.org/10.1016/j.hal.2011.10.027

Pineda, A., Velho, L.F.M., de Carvalho, P., Rodrigues, L.C., 2024. How a mega-dam in the Amazon's
largest tributary altered the drivers of planktonic communities. Hydrobiologia.
https://doi.org/10.1007/s10750-024-05529-y

Pouria, S., De Andrade, A., Barbosa, J., Cavalcanti, R.L., Barreto, V.T.S., Ward, C.J., Preiser, W.,
Poon, G.K., Neild, G.H., Codd, G.A., 1998. Fatal microcystin intoxication in haemodialysis unit
in Caruaru, Brazil. The Lancet 352, 21–26. https://doi.org/10.1016/S0140-6736(97)12285-1

Pracheil, B.M., DeRolph, C.R., Schramm, M.P., Bevelhimer, M.S., 2016. A fish-eye view of riverine
hydropower systems: the current understanding of the biological response to turbine passage.
Rev Fish Biol Fish 26, 153–167. https://doi.org/10.1007/s11160-015-9416-8

R Development Core Team, 2023. R: a language and environment for statistical computing. R
Foundation for Statistical Computing.

Reynolds, C.S., 2006. The Ecology of phytoplankton, Ecology. Cambridge University Press, New
York, USA. https://doi.org/http://dx.doi.org/10.1017/CBO9780511542145

Reynolds, C.S., Huszar, V., Kruk, C., Naselli-Flores, L., Melo, S., 2002. Towards a functional
classification of the freshwater phytoplankton. J Plankton Res 24, 417–428.
https://doi.org/10.1093/plankt/24.5.417

Rodrigues, L.C., Pivato, B.M., Vieira, L.C.G., Bovo-Scomparin, V.M., Bortolini, J.C., Pineda, A.,
Train, S., 2018. Use of phytoplankton functional groups as a model of spatial and temporal
patterns in reservoirs: a case study in a reservoir of central Brazil. Hydrobiologia 805, 147–161.
https://doi.org/10.1007/s10750-017-3289-x

Rodrigues, M.S., Batista, E.P., Silva, A.A., Costa, F.M., Neto, V.A.S., Gil, L.H.S., 2017. Change in
Anopheles richness and composition in response to artificial flooding during the creation of the
Jirau hydroelectric dam in Porto Velho, Brazil. Malar J 16, 87–93.
https://doi.org/10.1186/s12936-017-1738-7

Soares-Filho, B.S., Nepstad, D.C., Curran, L.M., Cerqueira, G.C., Garcia, R.A., Ramos, C.A., Voll,
E., McDonald, A., Lefebvre, P., Schlesinger, P., 2006. Modelling conservation in the Amazon
basin. Nature 440, 520–523. https://doi.org/10.1038/nature04389

Souza, C.A. de, Vieira, L.C.G., Legendre, P., Carvalho, P. de, Velho, L.F.M., Beisner, B.E., 2019.
Damming interacts with the flood pulse to alter zooplankton communities in an Amazonian
river. Freshw Biol 64, 1040–1053. https://doi.org/10.1111/fwb.13284

Souza, D.G., Bueno, N.C., Bortolini, J.C., Rodrigues, L.C., Bovo-Scomparin, V.M., de Souza Franco,
G.M., 2016. Phytoplankton functional groups in a subtropical Brazilian reservoir: responses to
impoundment. Hydrobiologia 779, 47–57. https://doi.org/10.1007/s10750-016-2798-3

Sun, J., Liu, D., 2003. Geometric models for calculating cell biovolume and surface area for
phytoplankton. J Plankton Res 25, 1331–1346. https://doi.org/10.1093/plankt/fbg096

Thomaz, S.M., Bini, L.M., De Souza, M.C., Kita, K.K., Camargo, A.F.M., 1999. Aquatic
macrophytes of Itaipu Reservoir, Brazil: Survey of species and ecological considerations.
Brazilian Archives of Biology and Technology. https://doi.org/10.1590/S1516-
89131999000100003

Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R., Polasky, S., 2002. Agricultural sustainability
and intensive production practices. Nature. https://doi.org/10.1038/nature01014

Tundisi, J.G., 2018. Reservoirs: New challenges for ecosystem studies and environmental
management. Water Secur 4–5, 1–7. https://doi.org/10.1016/j.wasec.2018.09.001

Utermöhl, H., 1958. Mitteilungen der Internationalen Vereinigung für Theoretische und Angewandte.
Mitteilungen der Internationalen Vereinigung für Theoretische und Angewandte Limnologie 9,
1–38.

Vale, P., Gibbs, H., Vale, R., Christie, M., Florence, E., Munger, J., Sabaini, D., 2019. The expansion
of intensive beef farming to the Brazilian Amazon. Global Environmental Change 57, 101922.
https://doi.org/10.1016/j.gloenvcha.2019.05.006

Vasconcelos, L.P., Alves, D.C., Agostinho, A.A., Hahn, L., da Câmara, L.F., Gomes, L.C., 2022. Fish
eggs and larvae drifting through hydropower reservoirs: a case study in the Brazilian Amazon.
Hydrobiologia 849, 357–372. https://doi.org/10.1007/s10750-021-04631-9

Vollenweider, 1968. Water management research. Scientific fundamentals of the eutrophication of
lakes and flowing waters with particular reference to nitrogen and phosphorus as factors in
eutrophication. Technical Repport DAS/CSI/68, Paris.

Wickham, H., 2016. ggplot2: Elegant graphics for data analysis. Springer-Verlag, New York.

Winemiller, K.O., McIntyre, P.B., Castello, L., Fluet-Chouinard, E., Giarrizzo, T., Nam, S., Baird,
I.G., Darwall, W., Lujan, N.K., Harrison, I., Stiassny, M.L.J., Silvano, R.A.M., Fitzgerald, D.B.,
Pelicice, F.M., Agostinho, A.A., Gomes, L.C., Albert, J.S., Baran, E., Petrere, M., Zarfl, C.,
Mulligan, M., Sullivan, J.P., Arantes, C.C., Sousa, L.M., Koning, A.A., Hoeinghaus, D.J., Sabaj,
M., Lundberg, J.G., Armbruster, J., Thieme, M.L., Petry, P., Zuanon, J., Vilara, G.T., Snoeks, J.,
Ou, C., Rainboth, W., Pavanelli, C.S., Akama, A., Soesbergen, A. van, Sáenz, L., 2016.
Balancing hydropower and biodiversity in the Amazon, Congo, and Mekong. Science (1979)
351, 128–129. https://doi.org/10.1126/science.aac7082

Zanco, B.F., Pineda, A., Bortolini, J.C., Jati, S., Rodrigues, L.C., 2017. Phytoplankton functional
groups indicators of environmental conditions in floodplain rivers and lakes of the Paraná Basin.
Acta Limnologica Brasiliensia 29, e119. https://doi.org/10.1590/S2179-975X8117

Zanon, F.M., Pineda, A., Lansac-Tôha, F.M., Rodrigues, L.C., 2024. Impact of a run-of-river dam in
an Amazonian large river on the spatial and temporal patterns of phytoplankton beta diversity.
Biol Conserv 295, 1–12. https://doi.org/10.1016/j.biocon.2024.110656

Zanon, J.E., de Carvalho, P., Rodrigues, L.C., Bini, L.M., 2019. Potential mechanisms related to the
spatial synchrony of phytoplankton is dependent on the type of data. Hydrobiologia 841, 95–
108. https://doi.org/10.1007/s10750-019-04009-y

695