An improved tetracycline-inducible expression system for fission yeast

Xiao-Hui Lyu[1], Yu-Sheng Yang[1], Zhao-Qian Pan[1], Shao-Kai Ning[1], Fang Suo[1], Li-Lin Du[1,2,*]

[1]National Institute of Biological Sciences, Beijing 102206, China

[2]Tsinghua Institute of Multidisciplinary Biomedical Research, Tsinghua University, Beijing 102206, China

*Corresponding author: dulilin@nibs.ac.cn (L.-L.D.)

Running head: Inducible system for *S. pombe*.

Keywords: inducible expression system, fission yeast, *Schizosaccharomyces pombe*, tetracycline, conditional loss-of-function

Abstract

The ability to manipulate gene expression is valuable for elucidating gene function. In the fission yeast *Schizosaccharomyces pombe*, the most widely used regulatable expression system is the *nmt1* promoter and its two attenuated variants. However, these promoters have limitations, including a long lag, incompatibility with rich media, and unsuitability for non-dividing cells. Here, we present a tetracycline-inducible system free of these shortcomings. Our system features the *enotetS* promoter, which achieves a similar induced level and a higher induction ratio compared to the *nmt1* promoter, without exhibiting a lag. Additionally, our system includes four weakened *enotetS* variants, offering an expression range similar to the *nmt1* series promoters but with more intermediate levels. To enhance usability, each promoter is combined with a Tet-repressor-expressing cassette in an integration plasmid. Importantly, our system can be used in non-dividing cells, enabling the development of a synchronous meiosis induction method with high spore viability. Moreover, our system allows for the shutdown of gene expression and the generation of conditional loss-of-function mutants. This system provides a versatile and powerful tool for manipulating gene expression in fission yeast.

30 **Summary statement**

31 A new inducible expression system for fission yeast enhances control, offers compatibility

32 with non-dividing cells, and enables synchronous meiosis induction and conditional loss-of-

33 function analysis.

Introduction

Understanding gene function is a central pursuit in biological studies. One important approach to dissecting the function of individual genes is through the artificial manipulation of gene expression levels. Over the past 35 years, a large number of regulatable promoter systems have been developed to enable the manipulation of gene expression in the fission yeast *Schizosaccharomyces pombe*. These systems include glucose-repressible promoters (Hoffman and Winston, 1989; Iacovoni et al., 1999), hormone-inducible promoters (Ohira et al., 2017; Picard et al., 1990), thiamine-repressible promoters and their derivatives (Basi et al., 1993; Garg, 2020; Kjaerulff and Nielsen, 2015; Kumar and Singh, 2006; Maundrell, 1993, 1990; Moreno et al., 2000), tetracycline-inducible promoters (Erler et al., 2006; Faryar and Gatz, 1992; Patterson et al., 2019; Zilio et al., 2012), a copper-repressible promoter (Bellemare et al., 2001), a heat shock-inducible promoter (Fujita et al., 2006), a uracil-inducible promoter (Watson et al., 2011; Watt et al., 2008), an ethanol-inducible promoter (Matsuzawa et al., 2013), and pheromone-inducible promoters (Hennig et al., 2018).

Among the various regulatable promoters available for *S. pombe*, the most popular ones are three thiamine-repressible promoters based on that of the *nmt1* gene (Basi et al., 1993; Maundrell, 1993, 1990). This promoter system offers several advantages. Firstly, the *nmt1* promoter (*Pnmt1*) is a highly active promoter when induced, as *nmt1* ranks as the eighth most highly transcribed protein-coding gene in vegetative cells (340 mRNA molecules per cell) (Marguerat et al., 2012). Secondly, there are two weakened promoter variants, *P41nmt1* and *P81nmt1*, which allow for lower levels of expression, approximately one and two orders of magnitude lower than *Pnmt1*, respectively (Basi et al., 1993). Thirdly, these promoters are strongly repressed in the presence of thiamine, resulting in an around 100-fold difference in expression level between the induced and repressed states (Basi et al., 1993; Vještica et al., 2020). However, this system is not without limitations. It is not compatible with yeast extract-

based rich media that contain thiamine. Additionally, induction requires a medium switch,

which incurs time and labor costs. The most significant drawback is that the intracellular

thiamine concentration needs to be diluted through several cell doublings before induced

expression can occur (Maundrell, 1990; Tommasino and Maundrell, 1991). This results in a

long lag time of around 10-12 h and makes the system unsuitable for non-dividing cells such

as cells in a quiescent state.

Tetracycline-inducible systems based on the Tet repressor-operator system of the

bacterial transposon Tn*10* have been widely used to manipulate gene expression in

eukaryotes (Berens and Hillen, 2003; Hillen et al., 1984, 1983). A basic version of the

tetracycline-inducible systems consists of three components: the *tet* operator DNA sequence

(*tetO*), which is incorporated into a promoter; the Tet repressor protein (TetR), which has a

high binding affinity to *tetO*; and a small-molecule inducer that prevents TetR from binding to

tetO. Commonly used inducers include tetracycline and its analogs anhydrotetracycline

(ahTet) and doxycycline. In the absence of the inducer, TetR binds to *tetO* and represses the

transcription of the downstream gene. When the inducer is added, TetR no longer binds to

tetO, and the transcription of the downstream gene is induced.

Three different tetracycline-inducible promoters have been used in fission yeast. In 1992,

Faryar and Gatz established the first tetracycline-inducible promoter for fission yeast by

incorporating three *tetO* sequences into the cauliflower mosaic virus 35S (CaMV35S)

promoter (Faryar and Gatz, 1992). This tetracycline-inducible promoter can be quickly

induced, with a reporter protein product becoming detectable within two hours after the

addition of an inducer (Faryar and Gatz, 1992). However, an independent study showed that

this system has an induction ratio of only 10-fold and the induced expression level is low,

similar to that of *P81nmt1* (Forsburg, 1993). In 2012, Zilio et al. adapted a tetracycline-

inducible system, originally developed for the budding yeast *Saccharomyces cerevisiae*, for

use in the fission yeast (Bellí et al., 1998; Garí et al., 1997; Zilio et al., 2012). The promoter

in this system is composed of seven *tetO* sequences and the basal promoter of the budding

yeast *CYC1* gene. The induced level of expression of this promoter is also low, similar to that

of *P81nmt1* (Zilio et al., 2012). In 2019, a much stronger tetracycline-inducible promoter,

called the *enoTet* promoter (*PenoTet*), was developed by placing three *tetO* sequences into

the promoter of the *S. pombe eno101* gene (Patterson et al., 2019). The *eno101* gene is the

fourth most highly transcribed protein-coding gene in vegetative cells (470 mRNA molecules

per cell) (Marguerat et al., 2012). *PenoTet* achieves an induced expression level about 50%

as high as that of *Pnmt1* (Patterson et al., 2019). Unlike the *nmt1* series promoters, these

tetracycline-inducible promoters are compatible with rich media, do not need a medium

switch for induction, do not exhibit an obvious lag time, and can be used in non-dividing cells.

However, these promoters have not been widely adopted, potentially due to the lack of multi-

level options, insufficient induced expression levels, non-ideal induction ratios, and the

inconvenience associated with the need to introduce both a tetracycline-inducible promoter

and a TetR-expressing cassette into a host strain.

In the present study, we have developed an improved tetracycline-inducible expression

system for *S. pombe*. This system includes a strong promoter, *PenotetS*, which, similar to

PenoTet, is derived from the *eno101* promoter but contains only a single *tetO* sequence.

PenotetS has a high induction ratio of approximately 200-fold and its induced level is similar

to that of *Pnmt1*. Additionally, our system includes four weakened variants of *PenotetS*,

allowing for five different levels of induced expression. For ease of use, a collection of

integration plasmids was constructed, each containing an *enotetS* series promoter and a

TetR-expressing cassette. Capitalizing on the compatibility of this system with non-dividing

cells, we have developed a new method to induce meiosis, resulting in favorable spore

viability and synchrony. Moreover, we have successfully employed this system to generate

conditional lethal mutants of essential genes, underscoring its versatility and applicability.

Results

The development of a strong inducible promoter *PenotetS*

It has been shown that in the case of the *GAL1* promoter of *Saccharomyces cerevisiae*, placing a single *tetO* sequence between the TATA box and the transcription start site (TSS) is sufficient to render the promoter strongly repressible by TetR (Murphy et al., 2007). We adopted this approach to modify the 276-bp promoter of the *S. pombe eno101* gene (Wang et al., 2014), which was the basis of the *PenoTet* promoter that contains three *tetO* sequences (Patterson et al., 2019). In *S. pombe* promoters, the TATA box and the TSS, each possessing distinct sequence characteristics, are typically separated by a constant distance (Li et al., 2015; Lu and Lin, 2021; Russell, 1983; Thodberg et al., 2019). To maintain this distance, we substituted a 19-bp sequence between the TATA box and the TSS of the *eno101* promoter with a 19-bp *tetO* sequence (Fig. 1A). We named the modified promoter *PenotetS*, with the letter "*S*" indicating the presence of a single copy of the *tetO* sequence.

A drawback of some of the previously published tetracycline-inducible systems for *S. pombe* is the inconvenience caused by the need to separately introduce a tetracycline-inducible promoter and a TetR-expressing cassette. To establish a user-friendly system, we combined *PenotetS* with a TetR-expressing cassette, which consists of the cytomegalovirus (CMV) promoter and the TetR coding sequence, in a stable integration vector (SIV) (Vještica et al., 2020) (Fig. 1B). The CMV promoter is a robust constitutive promoter in *S. pombe* (Amelina et al., 2016; Toyama and Okayama, 1990). Additionally, downstream of *PenotetS*, we inserted the coding sequence for the yellow fluorescent protein mECitrine, followed by the terminator sequence derived from the *ADH1* gene of *Saccharomyces cerevisiae*. The resulting plasmid was named pDB5318. The fluorescence emitted by mECitrine serves as an indicator of the expression level of *PenotetS*.

In *S. pombe*, previous studies have shown that anhydrotetracycline (ahTet), an analog

of tetracycline, is a more effective inducer compared to tetracycline and doxycycline (Erler et

al., 2006; Zilio et al., 2012). When an *S. pombe* strain containing pDB5318 was cultured in

YES medium supplemented with 2.5 µg/mL of ahTet, a concentration used in a previous

study (Zilio et al., 2012), it was observed that the expression level of *PenotetS* was higher

than that of the strong constitutive promoter *Padh1* (Fig. 1D). In the absence of ahTet, the

expression level of *PenotetS* was markedly lower than that of *Padh1*. Thus, *PenotetS* is an

inducible promoter with a high level of induced expression and a substantial induction ratio.

To determine whether the expression of TetR or treatment with ahTet affects the growth

rate, we conducted a growth curve analysis using a strain containing pDB5318 and a control

strain lacking pDB5318 (Fig. S1). The growth rates of both strains were identical in media

with or without ahTet, indicating that neither the expression of TetR nor the presence of 2.5

µg/mL of ahTet in the culture medium impacted the cell growth rate.

The development of four weakened variants of *PenotetS*

To provide a range of different expression levels, we developed four weakened variants of

PenotetS. The design was inspired by a previous study that used GC-rich stem-loop-forming

sequences inserted in the 5′ UTR to reduce gene expression (Lamping et al., 2013). It was

shown that the extent of expression attenuation increases with the number of GC base pairs

in the stem. We selected four different stem-loop-forming sequences used in that study (Fig.

1C). Each sequence was inserted immediately downstream of *PenotetS*. The resulting

promoters were named *PenotetSW1*, *PenotetSW2*, *PenotetSW3*, and *PenotetSW4*. The

letter "*W*" indicates a weakened variant, while the number denotes the expected expression

strength, with *PenotetSW1* being the highest and *PenotetSW4* being the lowest.

These weakened promoters were combined with the TetR-expressing cassette in an SIV

plasmid, following the same procedure as *PenotetS*. In YES medium supplemented with 2.5

μg/mL of ahTet, the expression strengths of these promoters varied as expected (Fig. 1D).

PenotetSW1 exhibited an expression level similar to that of *Padh1*, while the other three

promoters showed progressively lower levels. In the absence of ahTet, the expression level

of each promoter was dramatically lower compared to when ahTet was present, with the

three weaker promoters showing fluorescence signals that were indistinguishable from the

background.

Inducer dose response

We assessed the inducer dose response of *PenotetS* in both YES and EMM media. The

inducer ahTet was used at a range of six concentrations (0, 0.004, 0.02, 0.1, 0.5, and 2.5

μg/mL). After a 24-hour induction period (which was long enough to reach a steady-state

level; see below), the expression levels were measured using the fluorescence signal of the

reporter protein mECitrine (Fig. 2). In both types of media, a dramatic transition from a near-

minimal level to a near-maximal level was observed between two doses that differed by 5-

fold. In YES, this transition occurred between 0.004 μg/mL and 0.02 μg/mL of ahTet (Fig. 2B),

while in EMM, it occurred between 0.02 μg/mL and 0.1 μg/mL of ahTet (Fig. 2D). Similar

steep dose-response patterns have been observed in other published eukaryotic

tetracycline-inducible systems that utilize constitutively expressed TetR as a repressor

(Azizoglu et al., 2021; Erler et al., 2006; Nevozhay et al., 2009). We chose to use 2.5 μg/mL

of ahTet, an inducer concentration well above the saturating level in both types of media, for

subsequent experiments.

Induction kinetics in dividing cells

We next assessed the induction kinetics of *PenotetS* and *PenotetSW2* by performing

induction time course experiments in both YES and EMM media (Fig. 3). In all time courses,

the expression level of the reporter protein mECitrine increased by more than two orders of

magnitude within the first 10 hours following the addition of ahTet and ceased to increase

after 12 hours. We used the expression level at the 24-hour time point as the steady-state

level. In YES medium, for both promoters, at the one-hour time point after adding ahTet to

the cultures, the fluorescence signal of mECitrine had already increased by more than one

order of magnitude, reaching around 10% of the steady-state level. This rapid increase

suggests that both promoters were turned on quickly after the addition of ahTet. In EMM

medium, the initial phase of the induction was slightly slower, reaching around 5% of the

steady-state level at the one-hour time point. In both types of media, at the four-hour time

point, both promoters had reached or surpassed 50% of the steady-state level. By the eight-

hour time point, both promoters had reached within 15% of the steady-state level.

Upon the activation of an inducible promoter that drives the expression of a protein-

coding gene, the level of the protein product reaches a steady state when its synthesis is

balanced by its degradation and dilution (Azizoglu et al., 2021; Rosenfeld et al., 2002). The

steady-state expression level is determined by both the synthesis rate and the degradation-

and-dilution rate, while the induction kinetics are solely determined by the degradation-and-

dilution rate (Rosenfeld et al., 2002). Fluorescent proteins are highly stable and have in vivo

half-lives longer than 20 hours (Corish and Tyler-Smith, 1999; He et al., 2019; Perruca-

Foncillas et al., 2022). In our experimental setup, given that the doubling time of proliferating

S. pombe cells is around 3 hours (Petersen and Russell, 2016), the degradation-and-dilution

rate of mECitrine should mainly be determined by its dilution through cell division. In this

scenario, without considering any delay between transcription activation and the formation

of mECitrine protein capable of emitting fluorescence, the time required to reach 50% of the

steady-state expression level theoretically equals the doubling time (Rosenfeld et al., 2002).

However, nascent fluorescent proteins only become fluorescent after undergoing a

maturation process, which takes approximately 30 minutes for mECitrine (Azizoglu et al.,

2021). Taking this delay into account, the induction kinetics observed in our experiments align

closely with immediate or near-immediate transcription activation upon the addition of ahTet.

To better characterize the induction kinetics, we performed quantitative modeling by

fitting the experimental data to an induction-kinetics model. The resulting fitted curves closely

align with our quantitative data, and the half-life parameters derived from the modeling are

consistent with our proposed explanation of the expression kinetics (Fig. 3B-D). This analysis

further supports the conclusion that transcription activation occurs rapidly upon the addition

of ahTet.

Quantification of expression levels and comparison with the *nmt1* series promoters

Next, we conducted a comprehensive quantitative analysis of the induced and repressed

expression levels of all *enotetS* series promoters in YES and EMM media, and made

comparisons with *Padh1* in YES medium and the *nmt1* series promoters in EMM medium

(Fig. 4). In YES medium, the induced expression level of *PenotetS* was approximately 1.6-

fold that of *Padh1* (Fig. 4A). The fold differences between the induced level of *PenotetS* and

the induced levels of the four weakened *enotetS* series promoters were approximately 1.7-

fold, 7.1-fold, 30-fold, and 54-fold, respectively (Fig. 4A). The fold differences between the

induced and repressed levels (induction ratios) were about 207-fold for *PenotetS* and about

226-fold for *PenotetSW1* (Fig. 4A). Since the repressed levels of *PenotetSW2*, *PenotetSW3*,

and *PenotetSW4* were very close to the background, we could only determine that these

levels were below a threshold (THR) for accurate measurement (Fig. 4A).

In EMM medium, the induced and repressed expression levels of the *enotetS* series

promoters mirrored those observed in YES medium (Fig. 4B,C). When compared to the *nmt1*

series promoters, the induced expression level of *PenotetS* was almost the same as *Pnmt1*

and the induced levels of *PenotetSW2* and *PenotetSW4* were similar to those of *Pnmt41* and

P81nmt1, respectively (Fig. 4B,C). Therefore, in comparison to the *nmt1* series promoters,

the *enotetS* series promoters span the same range of induced levels but provide more

intermediate levels. Using the mRNA level of the *nmt1* gene in vegetative cells (340 mRNA

molecules per cell) as a reference (Marguerat et al., 2012; Rutherford et al., 2024), the

induced expression levels of *PenotetS*, *PenotetSW1*, *PenotetSW2*, *PenotetSW3*, and

PenotetSW4 correspond to approximately 365, 219, 57, 13, and 8 mRNA molecules per cell,

respectively (Fig. 4D). It is important to note that mRNA levels do not necessarily correlate

with protein levels, and the expression of different proteins from the same promoter can result

in markedly different protein levels. Regarding repressed expression levels, the *enotetS*

series promoters had lower repressed levels than the *nmt1* series promoters when

comparing promoters with similar induced levels. In our experiments, *Pnmt1* exhibited only

a 26-fold induction ratio, which is lower than the reported induction ratio of around 100-fold

(Basi et al., 1993; Vještica et al., 2020). The reason behind this discrepancy is unclear.

Nevertheless, even when considering the reported induction ratio for comparison, the

enotetS series promoters still outperformed the *nmt1* series promoters in this aspect.

Induction kinetics and expression levels in quiescent cells

One limitation of the *nmt1* series promoters is their unsuitability for non-dividing cells, as the

induction of these promoters requires the dilution of intracellular thiamine through multiple

rounds of cell division after shifting the cells to a thiamine-free medium. Tetracycline-inducible

systems should be free of this limitation. Therefore, we tested whether *PenotetS* can function

in two types of non-dividing cells: cells arrested in the G0 cell cycle phase through nitrogen

starvation and spores (Fig. 5).

The basal expression levels of *PenotetS* in G0 cells and spores, as reported by the

fluorescence of mECitrine, were slightly higher than the basal level in the dividing cells (Fig.

5C), presumably due to the lack of dilution of mECitrine through cell division. Upon the

addition of ahTet to G0 cells and spores, the expression of mECitrine from *PenotetS*

increased several-fold within the first hour and reached a level similar to the steady-state

level in the dividing cells at the eight-hour time point. Unlike the situation in the dividing cells,

where the expression of mECitrine from *PenotetS* reached a steady state at the 12-hour time

point, the level of mECitrine continued to increase in G0 cells and spores after the 12-hour

time point (Fig. 5C). The increase did not stop after the 24-hour time point and the level of

mECitrine reached several-fold higher than the steady-state level in the dividing cells at the

36-hour time point (Fig. 5C). The prolonged rise time and the higher induced levels in G0

cells and spores can be explained by the degradation-and-dilution rate of mECitrine no longer

being determined by its dilution through cell division but rather by the degradation of

mECitrine, which has a long half-life.

Meiosis induction in *h+/h+* diploids using *PenotetSW1*-expressed Mc-2A-Mi

In *S. pombe,* the most commonly used method to induce synchronous meiosis is through the

inactivation of a negative regulator of meiosis, Pat1, using a temperature-sensitive allele of

the *pat1* gene, *pat1-114*, in diploid cells homozygous for the mating type (*h+/h+* or *h−/h−*)

(Bähler et al., 1991; Iino and Yamamoto, 1985). However, this approach has several

drawbacks, including the use of non-physiological temperature and low spore viability (Bähler

et al., 1991; Chikashige et al., 2004; Cipak et al., 2012; Guerra-Moreno et al., 2012;

Yamamoto and Hiraoka, 2003). The development of ATP analog-sensitive alleles of *pat1* has

enabled the induction of synchronous meiosis at physiological temperature (Cipak et al.,

2012; Guerra-Moreno et al., 2012). Moreover, combining Pat1 inactivation with the

expression of the *matP*-cassette-encoded protein Pc in *h−/h−* diploids has been shown to

elevate spore viability to near-wild-type levels (Cipak et al., 2012; Yamamoto and Hiraoka,

2003). Thus, the current state-of-the-art approach for achieving synchronous meiosis with

normal spore viability in *S. pombe* requires the combination of two genetic modifications. To

showcase the practical utility of our *enotetS* series promoters, we devised a single-plasmid-

based method for inducing synchronous meiosis.

Diploid *S. pombe* cells that are heterozygous for the mating type (*h+/h−*) can undergo

meiosis, whereas those that are homozygous for the mating type (*h+/h+* or *h−/h−*) cannot.

This is because the latter lack proteins encoded by one of the two types of mating-type

cassettes. In the case of *h+/h+* diploid cells, the missing proteins are the *matM*-cassette-

encoded proteins Mc and Mi (Kelly et al., 1988). In our new method for inducing meiosis, we

used *PenotetSW1* to induce the expression of Mc and Mi linked by a self-processing 2A

peptide (Mc-2A-Mi) in *h+/h+* diploid cells (Fig. 6A). *h+/h+* cells harboring an integration

plasmid expressing Mc-2A-Mi under the control of *PenotetSW1* were initially cultured in YES

medium and subsequently transferred to EMM−N medium to arrest the cells in G0. Following

a 24-hour incubation in EMM−N medium, ahTet was added to induce meiosis. Time course

analysis demonstrated the successful induction of synchronous meiosis (Fig. 6B). Tetrad

analysis revealed a spore viability of 97% for meiosis induced using this method (Fig. 6C).

Thus, we have established a new method for inducing meiosis that is characterized by good

synchrony and normal spore viability.

Using *enotetS* series promoters to create conditional loss-of-function mutants

Studying the functions of essential genes relies on the creation of conditional loss-of-function

mutants. Previously, we developed an auxin-inducible degron (AID) system for generating

such mutants in *S. pombe* (Zhang et al., 2022). However, not all genes are compatible with

the AID approach. In an unrelated project, we applied the AID method to construct conditional

loss-of-function mutants of three essential genes: *ypp1*, *sui1*, and *nrs1*. We first performed

endogenous AID tagging in a diploid background and obtained heterozygous tagged diploid

strains. Then, we attempted to derive haploid AID-tagged progeny through tetrad dissection

(Fig. S2). AID-tagged haploid progeny were readily obtained for *ypp1*. However, no viable

AID-tagged haploid progeny were obtained for *sui1* and *nrs1*, possibly due to AID tagging

interfering with the function of the protein products of these two genes. We decided to take

this opportunity to explore the feasibility of using the *enotetS* series promoters for generating

conditional loss-of-function mutants, especially for genes that are not amenable to the AID

approach.

We first separately deleted *sui1* and *nrs1* in a diploid background to obtain heterozygous

deletion diploid strains. Then, for each gene, a plasmid expressing the gene from

PenotetSW2 and a plasmid expressing the gene from *PenotetSW3* were separately

integrated into the heterozygous deletion diploid strain. These two promoters were chosen

because their induced expression levels are close to the mRNA levels of *sui1* and *nrs1* in

vegetative cells (11 and 20 mRNA molecules per cell, respectively) (Marguerat et al., 2012;

Rutherford et al., 2024). The resulting strains were subjected to tetrad analysis on YES plates

with and without ahTet (Fig. 7). On YES plates without ahTet, for both genes, the basal level

expression from neither promoter was able to allow the deletion haploids to form colonies

(Fig. 7A). In contrast, on YES plates with ahTet, for both genes, the induced expression from

either promoter rescued the inviability of the deletion haploids (Fig. 7A). *PenotetSW2*-driven

induced expression in deletion haploids resulted in colony growth indistinguishable from wild

type, whereas *PenotetSW3*-driven induced expression resulted in colonies slightly smaller

than wild type at 3 days but not at 6 days (Fig. 7A). Thus, we successfully obtained

conditional lethal mutants for both genes.

To determine how quickly the loss-of-function phenotype arises upon the removal of

ahTet, we conducted a growth curve analysis (Fig. 7B). Haploid deletion strains harboring a

plasmid expressing the deleted gene from *PenotetSW2* or *PenotetSW3* were pre-cultured in

YES liquid medium with ahTet. After the cells were harvested and washed with water, they

were transferred to fresh YES medium with and without ahTet. At the six-hour time point, the

cultures in the ahTet-free YES medium started to exhibit a noticeably lower growth rate

compared to the cultures in the ahTet-containing YES medium. The differences in growth

rate became more pronounced as time elapsed. Given that the loss-of-function phenotype

does not manifest immediately after the promoter is turned off, but rather arises only after

the protein product has been significantly depleted through dilution and degradation, these

findings suggest that the *enotetS* series promoters can be rapidly switched off after shifting

cells to an ahTet-free medium. Therefore, these promoters provide useful tools for analyzing

loss-of-function phenotypes of essential genes in *S. pombe*.

Discussion

In this study, we have developed the *enotetS* series promoters, a new set of tetracycline-

inducible promoters for *S. pombe* (Fig. 8). These promoters offer a wide range of expression

levels and allow for rapid induction of gene expression in both dividing and non-dividing cells.

By combining each of these promoters with a TetR-expressing cassette, this system can be

easily implemented in *S. pombe* through a single plasmid transformation. To demonstrate

the usefulness of this new inducible expression system, we have successfully developed a

new method for inducing synchronous meiosis and generated conditional loss-of-function

mutants for two essential genes.

Tetracycline and its analogs have traditionally been considered biologically inert to

eukaryotic physiology (Gossen et al., 1993). However, recent research has demonstrated

that doxycycline, a commonly used inducer in mammalian tetracycline-inducible systems,

disturbs mitochondrial protein translation, presumably due to the bacterial origin of

mitochondria (Moullan et al., 2015). This raises concerns regarding the use of tetracycline-

inducible systems (Chatzispyrou et al., 2015; De Boeck and Verfaillie, 2021; Ohira et al.,

2017). Unlike tetracycline and doxycycline, which strongly inhibit bacterial protein translation

and belong to the category of "classical tetracyclines", the inducer used in our system, ahTet,

does not effectively inhibit bacterial protein translation and is considered an "atypical

tetracycline" (Rasmussen et al., 1991). Therefore, it is unlikely that ahTet negatively affects

mitochondrial protein translation. Consistent with this, even though mitochondrial protein

translation is essential for *S. pombe* proliferation, studies that have used ahTet on *S. pombe*,

including our own, have not observed any detrimental effects on proliferation (Berg and

Moseley, 2023; Erler et al., 2006; Lemay et al., 2014; Patterson et al., 2019; Shiroiwa et al.,

2011; Sunder et al., 2012; Torres-Garcia et al., 2020; Wang et al., 2018; Zilio et al., 2012).

As a precautionary measure against potential unknown side effects of ahTet on *S. pombe*,

373 especially when investigating non-growth phenotypes, we recommend including a strain

374 without an *enotetS* series promoter as a control to rule out the possibility that a phenotype is

375 caused by side effects of ahTet.

376  ahTet is a cost-effective inducer. When used at a concentration of 2.5 µg/mL, the cost of

377 ahTet for making one liter of ahTet-containing medium is about 6 US dollars (based on the

378 list price of Sigma-Aldrich catalog #37919). The ahTet-containing medium should be

379 prepared fresh from the DMSO-dissolved ahTet stock solution before use, as ahTet is not

380 very stable in aqueous solution. It has been reported that ahTet has a half-life of around 30

381 hours in culture media at 30°C (Politi et al., 2014).

382  A higher level of ahTet is required for induction in EMM than in YES (Fig. 2), and the

383 induction kinetics appear to be slightly slower at the early stage of induction in EMM than in

384 YES (Fig. 3). We speculate that these differences may be due to a reduced uptake of ahTet

385 in EMM. It has been reported that the addition of 5 mM of Mg^{2+} to the culture medium

386 diminishes the uptake of tetracycline into *E. coli* cells (Zhang et al., 2014). EMM contains a

387 notably higher concentration of Mg^{2+} (5.2 mM) than YES (less than 0.5 mM) (Li et al., 2020;

388 Petersen and Russell, 2016).

389  The five induced expression levels of the *enotetS* series promoters offer a wide range

390 of possibilities for manipulating gene expression levels. In theory, even more intermediate

391 expression levels can be obtained by titrating the inducer dose. However, the *PenotetS*

392 promoter (and likely its weakened variants) exhibits a steep dose-response curve. This

393 means that the range of inducer doses over which the promoter is responsive, known as the

394 input dynamic range, is narrow. Consequently, precise adjustments to expression levels by

395 altering the inducer dose can be challenging (Azizoglu et al., 2021). If desired, the input

396 dynamic range can be expanded by implementing negative auto-regulation of TetR (Azizoglu

397 et al., 2021; Madar et al., 2011; Nevozhay et al., 2009; Patterson et al., 2019).

398  The use of the *enotetS* series promoters to generate conditional lethal mutants of

399 essential genes is a valuable alternative to the AID method, especially for genes that cannot

400 tolerate AID tagging. This strategy, known as promoter shut-off, has previously been

401 implemented in *S. pombe* using the *nmt1* series promoters and has successfully revealed

402 the loss-of-function phenotypes of many essential genes (Bestul et al., 2021; Kelly et al.,

403 1993; Matynia et al., 2002; Ottilie et al., 1995; Petersen and Hagan, 2003; Tajadura et al.,

404 2004; Wang et al., 2002; Westwood et al., 2004; Win et al., 2004; Yue et al., 2014). Compared

405 to the *nmt1* series promoters, the *enotetS* series promoters offer more levels of expression

406 and larger ratios between induced and repressed conditions. Therefore, they are expected

407 to be better tools for promoter shut-off.

408  While we generated the promoter shut-off strains for *sui1* and *nrs1* by first establishing

409 heterozygous deletion diploid strains and then deriving haploid strains, it should be feasible

410 to directly obtain promoter shut-off haploid strains by inserting an *enotetS* series promoter,

411 along with the TetR-expressing cassette, upstream of the target essential gene using a

412 single-step CRISPR-based marker-free knock-in (Jacobs et al., 2014; Zhang et al., 2018).

413 The intended knock-in clones can be readily selected based on their ability to grow on ahTet-

414 containing plates but not on ahTet-free plates. Promoter shut-off using the *nmt1* series

415 promoters can be initiated by simply adding thiamine to the culture, whereas promoter shut-

416 off using the *enotetS* series promoters requires the more laborious process of washing cells

417 to remove ahTet. It would be worth exploring in the future the feasibility of implementing the

418 method of light-induced inactivation of ahTet for shutting off the *enotetS* series promoters

419 (Baumschlager et al., 2020).

Materials and methods

Plasmid construction

The plasmids used in this study are listed in Table S1, and the primers used in this study are listed in Table S2. We amplified the 276-bp *eno101* promoter (Wang et al., 2014) from genomic DNA using primers oYY1 and oYY2. This amplified promoter was then cloned between the SwaI and XhoI sites in the stable integration vector (SIV) pAV0714 (Addgene plasmid 133479), replacing the *act1* promoter (Vještica et al., 2020). The resulting plasmid was named pDB5530. For the construction of the *enotetS* promoter (*PenotetS*), a 19-bp sequence between the TATA box and the transcription start site (TSS) of the *eno101* promoter in pDB5530 was replaced with a 19-bp *tetO* sequence, specifically that of the O_2 operator from the bacterial transposon Tn*10* (TCCCTATCAGTGATAGAGA) (Hillen et al., 1984, 1983; Wray and Reznikoff, 1983). This alteration was accomplished by performing overlapping PCR using pDB5530 as the template and oXH5, oXH6, oXH7, and oXH8 as primers, and then cloning the PCR product between the PacI and HpaI sites in pDB5530. The resulting plasmid was named pDB5531. To use *PenotetS* to drive the expression of the yellow fluorescent protein mECitrine, which serves as a reporter protein, we replaced the *adh1* promoter in pDB4915, a modified SIV plasmid containing mECitrine, with *PenotetS*. pDB4915 was derived from the SIV plasmid pAV0585 (pAde6[Pmel]-hphMX, Addgene plasmid 133472) (Vještica et al., 2020), by altering the linearization site from PmeI to NotI and introducing the *adh1* promoter, the coding sequence of mECitrine, and the terminator of the *Saccharomyces cerevisiae ADH1* gene. *PenotetS* was amplified from pDB5531 using primers oXH17 and oXH18, and then cloned between the ApaI and NheI sites in pDB4915. The resulting plasmid was named pDB5532.

To construct an integration plasmid that contains both *PenotetS* and a TetR-expressing cassette, we amplified a TetR expression cassette containing the cytomegalovirus promoter

(*P.CMV*) and the coding sequence of TetR from pDM291-tetR-tup11Δ70 (Addgene plasmid

41027) (Zilio et al., 2012), using primers oXH19 and oXH47. The amplified cassette was then

cloned into the KpnI site in pDB5532. The resulting plasmid was named pDB5318. To

construct weakened variants of *PenotetS*, we used primers oXH34-oXH38 to amplify four

PCR products, each containing *PenotetS* and a stem-loop-forming sequence immediately

downstream of *PenotetS*. These PCR products were then cloned separately between the

ApaI and NheI sites in pDB5318, generating four different plasmids, which were named

pDB5319-pDB5322. The four weakened variants of *PenotetS* in these four plasmids were

named *PenotetSW1*, *PenotetSW2*, *PenotetSW3*, and *PenotetSW4*. The plasmids pDB5318-

pDB5322 have been deposited at Addgene (Addgene plasmids 204828-204832) and the

Yeast Genetic Resource Center of Japan (YGRC/NBRP) (NBRP IDs FYP6242-6246). The

SnapGene files for these plasmids are provided in a compressed folder as a supplementary

file.

The DNA sequence encoding Mc-2A-Mi was synthesized by BGI Tech Solutions (Beijing

Liuhe) and consists of three parts: the coding sequence of Mi, a 60-bp segment encoding

the 20-amino acid T2A peptide (RAEGRGSLLTCGDVEENPGP) (Beekwilder et al., 2014),

and the coding sequence of Mc. To construct a plasmid expressing Mc-2A-Mi, the Mc-2A-Mi

sequence was amplified using primers oXH161 and oXH162, and then cloned between the

NheI and BglII sites of pDB5319, replacing the coding sequence of mECitrine. In the resulting

plasmid (pDB5533), Mc-2A-Mi is under the control of *PenotetSW1*.

The plasmids pDB4877, pDB4878, and pDB4879 were derived from the SIV vector

pAV0585 (pAde6^{PmeI}-hphMX, Addgene plasmid 133472) (Vještica et al., 2020), by altering

the linearization site from PmeI to NotI and introducing the *nmt1* promoter, the *41nmt1*

promoter, or the *81nmt1* promoter, respectively, along with the coding sequence of mECitrine,

and the terminator of the *Saccharomyces cerevisiae ADH1* gene. The plasmids pDB5534,

pDB5535, pDB5536, pDB5537, and pDB5538 were derived from the SIV plasmid pAV0586

(pLys3[BstZ17I]-bsdMX, Addgene plasmid 133473), by altering the linearization site from

BstZ17I to NotI and introducing the TetR expression cassette, an *enotetS* series promoter,

the coding sequence of mECitrine, and the terminator of the *Saccharomyces cerevisiae*

ADH1 gene. To construct plasmids expressing *nrs1*, the coding sequence of *nrs1* was

amplified from genomic DNA and cloned into pDB5537 and pDB5536, replacing the coding

sequence of mECitrine in each plasmid. The resulting plasmids were named pDB5539 and

pDB5541, respectively. Two plasmids expressing *sui1* were similarly constructed, except

cDNA was used as the amplification template, and they were named pDB5540 and pDB5542.

Strain construction

The *S. pombe* strains used in this study are listed in Table S3. Strains containing an

integrated plasmid were constructed by linearizing an SIV plasmid with NotI and then

transforming the plasmid into a host strain. Correct integration of the plasmid into the *ade6*

or *lys3* locus was verified using colony PCR (Vještica et al., 2020). Deletion strains were

created using PCR-based gene deletion. Strains with a gene endogenously tagged at the C-

terminus with the 3×sAID tag were generated through PCR-based gene tagging (Zhang et

al., 2022). The *h+/h−* diploid strain DY51002 was constructed by crossing two haploid strains

with *ade6-M210* and *ade6-M216* mutations, respectively. The two haploid strains were pre-

grown on YES plates and then mixed in an equal ratio in 20 µL of water. The mixture was

spotted on SPA plates with necessary supplements. After approximately 12 hours of

incubation at 30°C, cells were collected and plated on YE plates containing 200 mg/L guanine

at a density of approximately 10^4 cells per plate. Guanine was used to inhibit the uptake of

adenine, thereby preventing the growth of adenine auxotrophs and allowing only the growth

of *ade6-M210/ade6-M216* diploid cells, which are adenine prototrophs due to interallelic

complementation (Cummins and Mitchison, 1967; Smith, 2009). After incubating for 3 days

at 30°C, colonies formed on the guanine-containing plates were streaked on YE2A plates. A

diploid clone that showed high spore viability in tetrad analysis was saved as DY51002. A

prototrophic *h+/h−* diploid strain, DY50999, constructed in the same way as DY51002, was

plated on YEPD plates, and colonies that were unable to form spores were selected based

on iodine staining. The mating type of the non-sporulating diploid clones was determined by

colony PCR, and an *h+/h+* diploid strain was saved as DY51000.

Media and growth conditions

The compositions of the media were as described (Petersen and Russell, 2016).

Anhydrotetracycline hydrochloride (ahTet; J&K Scientific, catalog number 541642) was

dissolved in DMSO at a concentration of 5 mg/mL and stored at −80°C.

To analyze the dose response of the inducer, the cells were first cultured in liquid YES

medium or EMM medium with the necessary supplements until they reached the logarithmic

phase. The DMSO-diluted ahTet stock was then added to achieve the desired concentrations,

with an equal volume of DMSO serving as the control. After 24 hours of culturing (with

cultures diluted when the OD600 approaching 1.0) at 30°C, logarithmic phase cells were

collected for live-cell imaging.

For the time course assay of proliferating cells, the cells were first cultured in liquid YES

medium or EMM medium with the necessary supplements until they reached the logarithmic

phase at 30°C. After adding ahTet to the medium, the cells were collected at specific time

points. During the time course, the cultures were diluted when the OD600 approached 1.0.

To quantify the expression levels of the *enotetS* series promoters, induced expression

levels were measured after transferring logarithmic phase cells to media with 2.5 μg/mL ahTet

for 24 hours at 30°C. Repressed expression levels were measured after transfer to media

with DMSO under the same conditions.

To quantify the expression levels of the *nmt1* series promoters, the cells were first

cultured in EMM medium containing thiamine until they reached the logarithmic phase. Then,

to measure the induced expression levels, the cells were washed three times with water,

transferred to EMM medium without thiamine, and grown at 30°C for 24 hours. To measure

the repressed expression levels, the cells were cultured in EMM medium containing thiamine

at 30°C for 24 hours.

For the time course assay of G0 phase cells, logarithmically growing cells grown in EMM

medium were washed three times with water and shifted to EMM medium without nitrogen

(EMM−N) for 24 hours at 30°C to arrest the cells in the G0 phase. After adding ahTet to the

medium at a concentration of 2.5 µg/mL, the cells were collected at specific time points.

For the time course assay of spores, h^{90} cells grown in YES medium were washed three

times with water and shifted to MSL−N medium at 30°C for 48 hours to induce sporulation.

Then, the cells were collected and treated with Snailase (Beijing Solarbio Science &

Technology Co.,Ltd., catalog number S8280) in water at 30°C for 24 hours to digest non-

spore cells. After digestion, spores were purified by centrifugation in Percoll solutions. Spores

were incubated in EMM−N medium overnight. Then ahTet was added to the EMM−N medium.

After adding ahTet to the medium at a concentration of 2.5 µg/mL, spores were collected at

specific time points.

For the growth curve analysis, haploid deletion strains harboring a plasmid expressing

the deleted gene from *PenotetSW2* or *PenotetSW3*, along with a wild-type strain were pre-

cultured in YES liquid medium supplemented with ahTet until they reached the logarithmic

phase at 30°C. Then, the cells were washed three times with water and transferred to either

YES medium or YES medium supplemented with ahTet. The optical density at 600 nm

(OD600) of each culture was measured at specific time points, with cultures being diluted if

the OD600 approached 1.0. The number of generations was determined based on the

OD600 and the dilution ratio.

Fluorescence microscopy and image quantification

Live-cell imaging was performed using a Dragonfly 200 spinning disk confocal microscope

system (Andor Technology) equipped with an mCherry/YFP/CFP filter set and an

mCherry/GFP filter set. Images were acquired with a 100×, 1.4-NA objective using a Sona

sCMOS camera and analyzed with Fiji (ImageJ).

To quantify the expression level of mECitrine, first, channels of images were split and

the DIC channel was used to segment cell boundaries using Cellpose (version 2.1.1)

(Stringer et al., 2021). The resulting masked images were processed with the LabelsToRois

plugin (Waisman et al., 2021) in Fiji (Schindelin et al., 2012). This plugin generated outlines

outlining the boundaries of each cell. Subsequently, the generated outlines were

superimposed onto the mECitrine channel. Fiji was employed to calculate the mean

fluorescence intensity for each individual cell.

In total, three images were acquired for each sample, with each image containing

between 50 and 200 cells. The median fluorescence intensity of cells within each image was

calculated and averaged across the triplicate images. Background fluorescence was

measured using a parental strain that lacked mECitrine and was subsequently subtracted.

The resulting fluorescence intensity provided a measurement of the expression level. For all

fluorescence quantification results, we conducted two or more biological replicates, each

yielding consistent outcomes.

Induction and monitoring of meiosis

Log-phase cells grown in YES medium were collected, washed three times with water,

resuspended in EMM medium without nitrogen (EMM−N), and incubated at 30°C for 24 hours

to arrest cells in the G0 phase. Then, ahTet was added to the culture at a final concentration

of 2.5 μg/mL to induce meiosis. At specific time points, cells were collected and fixed using

70% ethanol. The advancement of meiosis was evaluated by quantifying the number of nuclei

per cell after staining the DNA with SYTOX Green (Thermo Fisher, catalog number S7020).

For SYTOX Green staining, the fixed cells were washed once with water, resuspended in

water containing 1 μM SYTOX Green, and incubated in the dark at room temperature for 5

minutes. The number of nuclei per cell was determined through imaging. The experiment

was conducted in three biological replicates, with one replicate presented as representative

data in the figure.

Tetrad analysis

To examine spore viability of induced meiosis, cells collected 24 hours after meiosis induction

were streaked onto YES plates. Tetrad analysis was carried out using a TDM50 tetrad

dissection microscope (Micro Video Instruments, Avon, USA) on YES plates. To assess

whether the exogenous expression of *nrs1* or *sui1* can complement the deletion of the

corresponding endogenous gene, diploid strains freshly grown on YE2A plates were spotted

onto SPASK plates. The SPASK plates were incubated at 30°C for about 24 h. Subsequently,

tetrad analysis was carried out on YES plates with and without ahTet.

Curve fitting

The induction kinetics data were fitted using Equation 1, as shown below.

$t \leq lag, y = base,$

$t > lag, y = base + (max - base) * (1 - e^{\wedge}(-(t - lag) * \frac{ln(2)}{half\text{-}life})) , (1)$

In this equation, "lag" represents the time required for the newly expressed reporter

protein to become detectable, set to 0.5 hours because mECitrine, the reporter protein used,

takes approximately 30 minutes to mature and emit fluorescence. "Base" denotes the

expression level of the reporter protein at t = 0, set to the quantified value at that time. "Max"

indicates the steady-state expression level. "Half-life" refers to the half-life of the reporter

protein, determined by both protein degradation and dilution through cell doubling. Curve

fitting was performed using the lmfit library in Python, employing the leastsq method (Newville

et al., 2016).

Disclosure on the use of artificial intelligence tools

The manuscript was proofread using the artificial intelligence tool editGPT. Following the

use of this tool, the authors carefully reviewed and edited the content as necessary. The

authors assume full responsibility for the publication's content.

Acknowledgements

We thank the Sophie Martin laboratory for making the SIV plasmids they have developed

available through Addgene and the Yeast Genetic Resource Center of Japan (YGRC/NBRP).

We thank YGRC/NBRP for providing the plasmids.

Competing interests

No competing interests declared.

Funding

This work was supported by funding from the Ministry of Science and Technology of the

People's Republic of China, the Beijing municipal government, and Tsinghua University.

Data and resource availability

Data availability: All relevant data can be found within the article and its supplementary

information.

References

Amelina, H., Moiseeva, V., Collopy, L.C., Pearson, S.R., Armstrong, C.A., Tomita, K. (2016). Sequential and counter-selectable cassettes for fission yeast. BMC Biotechnol 16, 76. https://doi.org/10.1186/s12896-016-0307-4

Azizoglu, A., Brent, R., Rudolf, F. (2021). A precisely adjustable, variation-suppressed eukaryotic transcriptional controller to enable genetic discovery. Elife 10, e69549. https://doi.org/10.7554/eLife.69549

Bähler, J., Schuchert, P., Grimm, C., Kohli, J. (1991). Synchronized meiosis and recombination in fission yeast: observations with pat1-114 diploid cells. Curr Genet 19, 445–451. https://doi.org/10.1007/BF00312735

Basi, G., Schmid, E., Maundrell, K. (1993). TATA box mutations in the Schizosaccharomyces pombe nmt1 promoter affect transcription efficiency but not the transcription start point or thiamine repressibility. Gene 123, 131–136. https://doi.org/10.1016/0378-1119(93)90552-E

Baumschlager, A., Rullan, M., Khammash, M. (2020). Exploiting natural chemical photosensitivity of anhydrotetracycline and tetracycline for dynamic and setpoint chemo-optogenetic control. Nat Commun 11, 3834. https://doi.org/10.1038/s41467-020-17677-5

Beekwilder, J., van Rossum, H.M., Koopman, F., Sonntag, F., Buchhaupt, M., Schrader, J., Hall, R.D., Bosch, D., Pronk, J.T., van Maris, A.J.A. et al. (2014). Polycistronic expression of a β-carotene biosynthetic pathway in Saccharomyces cerevisiae coupled to β-ionone production. J Biotechnol 192 Pt B, 383–392. https://doi.org/10.1016/j.jbiotec.2013.12.016

Bellemare, D.R., Sanschagrin, M., Beaudoin, J., Labbé, S. (2001). A novel copper-regulated promoter system for expression of heterologous proteins in Schizosaccharomyces pombe. Gene 273, 191–198. https://doi.org/10.1016/s0378-1119(01)00591-1

Bellí, G., Garí, E., Piedrafita, L., Aldea, M., Herrero, E. (1998). An activator/repressor dual system allows tight tetracycline-regulated gene expression in budding yeast. Nucleic Acids Res 26, 942–947. https://doi.org/10.1093/nar/26.4.942

Berens, C., Hillen, W. (2003). Gene regulation by tetracyclines. Constraints of resistance regulation in bacteria shape TetR for application in eukaryotes. Eur J Biochem 270, 3109–3121. https://doi.org/10.1046/j.1432-1033.2003.03694.x

Berg, R.A., Moseley, J.B. (2023). Regulation of cell size and Wee1 kinase by elevated levels of the cell cycle regulatory protein kinase Cdr2. J Biol Chem 299, 102831. https://doi.org/10.1016/j.jbc.2022.102831

Bestul, A.J., Yu, Z., Unruh, J.R., Jaspersen, S.L. (2021). Redistribution of

centrosomal proteins by centromeres and Polo kinase controls partial nuclear envelope
breakdown in fission yeast. Mol Biol Cell 32, 1487–1500.
https://doi.org/10.1091/mbc.E21-05-0239
Chatzispyrou, I.A., Held, N.M., Mouchiroud, L., Auwerx, J., Houtkooper, R.H.
(2015). Tetracycline antibiotics impair mitochondrial function and its experimental use
confounds research. Cancer Res 75, 4446–4449. https://doi.org/10.1158/0008-
5472.CAN-15-1626
Chikashige, Y., Kurokawa, R., Haraguchi, T., Hiraoka, Y. (2004). Meiosis induced
by inactivation of Pat1 kinase proceeds with aberrant nuclear positioning of
centromeres in the fission yeast Schizosaccharomyces pombe. Genes Cells 9, 671–684.
https://doi.org/10.1111/j.1356-9597.2004.00760.x
Cipak, L., Hyppa, R.W., Smith, G.R., Gregan, J. (2012). ATP analog-sensitive Pat1
protein kinase for synchronous fission yeast meiosis at physiological temperature. Cell
Cycle 11, 1626–1633. https://doi.org/10.4161/cc.20052
Corish, P., Tyler-Smith, C. (1999). Attenuation of green fluorescent protein half-life
in mammalian cells. Protein Eng 12, 1035–1040.
https://doi.org/10.1093/protein/12.12.1035
Cummins, J.E., Mitchison, J.M. (1967). Adenine uptake and pool formation in the
fission yeast Schizosaccharomyces pombe. Biochim Biophys Acta 136, 108–120.
https://doi.org/10.1016/0304-4165(67)90326-1
De Boeck, J., Verfaillie, C. (2021). Doxycycline inducible overexpression systems:
how to induce your gene of interest without inducing misinterpretations. Mol Biol Cell
32, 1517–1522. https://doi.org/10.1091/mbc.E21-04-0177
Erler, A., Maresca, M., Fu, J., Stewart, A.F. (2006). Recombineering reagents for
improved inducible expression and selection marker re-use in Schizosaccharomyces
pombe. Yeast 23, 813–823. https://doi.org/10.1002/yea.1396
Faryar, K., Gatz, C. (1992). Construction of a Tetracycline-inducible promoter in
Schizosaccharomyces pombe. Curr Genet 21, 345–349.
https://doi.org/10.1007/BF00351693
Forsburg, S.L. (1993). Comparison of Schizosaccharomyces pombe expression
systems. Nucleic Acids Res 21, 2955–2956. https://doi.org/10.1093/nar/21.12.2955
Fujita, Y., Tohda, H., Giga-Hama, Y., Takegawa, K. (2006). Heat shock-inducible
expression vectors for use in Schizosaccharomyces pombe. FEMS Yeast Res 6, 883–
887. https://doi.org/10.1111/j.1567-1364.2006.00093.x
Garg, A. (2020). A lncRNA-regulated gene expression system with rapid induction
kinetics in the fission yeast Schizosaccharomyces pombe. RNA 26, 1743–1752.
https://doi.org/10.1261/rna.076000.120

Garí, E., Piedrafita, L., Aldea, M., Herrero, E. (1997). A set of vectors with a
tetracycline-regulatable promoter system for modulated gene expression in
Saccharomyces cerevisiae. Yeast 13, 837–848. https://doi.org/10.1002/(SICI)1097-
0061(199707)13:9<837::AID-YEA145>3.0.CO;2-T
Gossen, M., Bonin, A.L., Bujard, H. (1993). Control of gene activity in higher
eukaryotic cells by prokaryotic regulatory elements. Trends Biochem Sci 18, 471–475.
https://doi.org/10.1016/0968-0004(93)90009-c
Guerra-Moreno, A., Alves-Rodrigues, I., Hidalgo, E., Ayté, J. (2012). Chemical
genetic induction of meiosis in Schizosaccharomyces pombe. Cell Cycle 11, 1621–
1625. https://doi.org/10.4161/cc.20051
He, L., Binari, R., Huang, J., Falo-Sanjuan, J., Perrimon, N. (2019). In vivo study of
gene expression with an enhanced dual-color fluorescent transcriptional timer. Elife 8,
e46181. https://doi.org/10.7554/eLife.46181
Hennig, S., Hornauer, N., Rödel, G., Ostermann, K. (2018). Pheromone-inducible
expression vectors for fission yeast Schizosaccharomyces pombe. Plasmid 95, 1–6.
https://doi.org/10.1016/j.plasmid.2017.11.002
Hillen, W., Gatz, C., Altschmied, L., Schollmeier, K., Meier, I. (1983). Control of
expression of the Tn10-encoded tetracycline resistance genes. Equilibrium and kinetic
investigation of the regulatory reactions. J Mol Biol 169, 707–721.
https://doi.org/10.1016/s0022-2836(83)80166-1
Hillen, W., Schollmeier, K., Gatz, C. (1984). Control of expression of the Tn10-
encoded tetracycline resistance operon. II. Interaction of RNA polymerase and TET
repressor with the tet operon regulatory region. J Mol Biol 172, 185–201.
https://doi.org/10.1016/s0022-2836(84)80037-6
Hoffman, C.S., Winston, F. (1989). A transcriptionally regulated expression vector for
the fission yeast Schizosaccharomyces pombe. Gene 84, 473–479.
https://doi.org/10.1016/0378-1119(89)90523-4
Iacovoni, J.S., Russell, P., Gaits, F. (1999). A new inducible protein expression
system in fission yeast based on the glucose-repressed inv1 promoter. Gene 232, 53–58.
https://doi.org/10.1016/S0378-1119(99)00116-X
Iino, Y., Yamamoto, M. (1985). Negative control for the initiation of meiosis in
Schizosaccharomyces pombe. Proc Natl Acad Sci U S A 82, 2447–2451.
https://doi.org/10.1073/pnas.82.8.2447
Jacobs, J.Z., Ciccaglione, K.M., Tournier, V., Zaratiegui, M. (2014).
Implementation of the CRISPR-Cas9 system in fission yeast. Nat Commun 5, 5344.
https://doi.org/10.1038/ncomms6344
Kelly, M., Burke, J., Smith, M., Klar, A., Beach, D. (1988). Four mating-type genes

control sexual differentiation in the fission yeast. EMBO J 7, 1537–1547.

Kelly, T.J., Martin, G.S., Forsburg, S.L., Stephen, R.J., Russo, A., Nurse, P. (1993).

The fission yeast cdc18+ gene product couples S phase to START and mitosis. Cell 74,

371–382. https://doi.org/10.1016/0092-8674(93)90427-r

Kjaerulff, S., Nielsen, O. (2015). An IPTG-inducible derivative of the fission yeast

nmt promoter. Yeast 32, 469–478. https://doi.org/10.1002/yea.3073

Kumar, R., Singh, J. (2006). A truncated derivative of nmt 1 promoter exhibits

temperature-dependent induction of gene expression in Schizosaccharomyces pombe.

Yeast 23, 55–65. https://doi.org/10.1002/yea.1343

Lamping, E., Niimi, M., Cannon, R.D. (2013). Small, synthetic, GC-rich mRNA

stem-loop modules 5' proximal to the AUG start-codon predictably tune gene

expression in yeast. Microb Cell Fact 12, 74. https://doi.org/10.1186/1475-2859-12-74

Lemay, J.-F., Larochelle, M., Marguerat, S., Atkinson, S., Bähler, J., Bachand, F.

(2014). The RNA exosome promotes transcription termination of backtracked RNA

polymerase II. Nat Struct Mol Biol 21, 919–926. https://doi.org/10.1038/nsmb.2893

Li, H., Hou, J., Bai, L., Hu, C., Tong, P., Kang, Y., Zhao, X., Shao, Z. (2015).

Genome-wide analysis of core promoter structures in Schizosaccharomyces pombe with

DeepCAGE. RNA Biol 12, 525–537. https://doi.org/10.1080/15476286.2015.1022704

Li, R., Jin, M., Du, J., Li, M., Chen, S., Yang, S. (2020). The Magnesium

Concentration in Yeast Extracts Is a Major Determinant Affecting Ethanol Fermentation

Performance of Zymomonas mobilis. Front Bioeng Biotechnol 8, 957.

https://doi.org/10.3389/fbioe.2020.00957

Lu, Z., Lin, Z. (2021). The origin and evolution of a distinct mechanism of

transcription initiation in yeasts. Genome Res 31, 51–63.

https://doi.org/10.1101/gr.264325.120

Madar, D., Dekel, E., Bren, A., Alon, U. (2011). Negative auto-regulation increases

the input dynamic-range of the arabinose system of Escherichia coli. BMC Syst Biol 5,

111. https://doi.org/10.1186/1752-0509-5-111

Marguerat, S., Schmidt, A., Codlin, S., Chen, W., Aebersold, R., Bähler, J. (2012).

Quantitative analysis of fission yeast transcriptomes and proteomes in proliferating and

quiescent cells. Cell 151, 671–683. https://doi.org/10.1016/j.cell.2012.09.019

Matsuzawa, T., Tohda, H., Takegawa, K. (2013). Ethanol-inducible gene expression

using gld1 (+) promoter in the fission yeast Schizosaccharomyces pombe. Appl

Microbiol Biotechnol 97, 6835–6843. https://doi.org/10.1007/s00253-013-4812-2

Matynia, A., Salus, S.S., Sazer, S. (2002). Three proteins required for early steps in the

protein secretory pathway also affect nuclear envelope structure and cell cycle

progression in fission yeast. J Cell Sci 115, 421–431.

https://doi.org/10.1242/jcs.115.2.421

Maundrell, K. (1993). Thiamine-repressible expression vectors pREP and pRIP for

fission yeast. Gene 123, 127–130. https://doi.org/10.1016/0378-1119(93)90551-d

Maundrell, K. (1990). nmt1 of fission yeast. A highly transcribed gene completely

repressed by thiamine. Journal of Biological Chemistry 265, 10857–10864.

https://doi.org/10.1016/S0021-9258(19)38525-4

Moreno, M.B., Durán, A., Ribas, J.C. (2000). A family of multifunctional thiamine-

repressible expression vectors for fission yeast. Yeast 16, 861–872.

https://doi.org/10.1002/1097-0061(20000630)16:9<861::AID-YEA577>3.0.CO;2-9

Moullan, N., Mouchiroud, L., Wang, X., Ryu, D., Williams, E.G., Mottis, A.,

Jovaisaite, V., Frochaux, M.V., Quiros, P.M., Deplancke, B. et al. (2015).

Tetracyclines Disturb Mitochondrial Function across Eukaryotic Models: A Call for

Caution in Biomedical Research. Cell Rep 10, 1681–1691.

https://doi.org/10.1016/j.celrep.2015.02.034

Murphy, K.F., Balázsi, G., Collins, J.J. (2007). Combinatorial promoter design for

engineering noisy gene expression. Proc Natl Acad Sci U S A 104, 12726–12731.

https://doi.org/10.1073/pnas.0608451104

Nevozhay, D., Adams, R.M., Murphy, K.F., Josic, K., Balázsi, G. (2009). Negative

autoregulation linearizes the dose-response and suppresses the heterogeneity of gene

expression. Proc Natl Acad Sci U S A 106, 5123–5128.

https://doi.org/10.1073/pnas.0809901106

Newville, M., Stensitzki, T., Allen, D.B., Rawlik, M., Ingargiola, A., Nelson, A.

(2016). Lmfit: Non-Linear Least-Square Minimization and Curve-Fitting for Python.

Astrophysics Source Code Library ascl:1606.014.

Ohira, M.J., Hendrickson, D.G., Scott McIsaac, R., Rhind, N. (2017). An estradiol-

inducible promoter enables fast, graduated control of gene expression in fission yeast.

Yeast 34, 323–334. https://doi.org/10.1002/yea.3235

Ottilie, S., Miller, P.J., Johnson, D.I., Creasy, C.L., Sells, M.A., Bagrodia, S.,

Forsburg, S.L., Chernoff, J. (1995). Fission yeast pak1+ encodes a protein kinase that

interacts with Cdc42p and is involved in the control of cell polarity and mating. EMBO

J 14, 5908–5919. https://doi.org/10.1002/j.1460-2075.1995.tb00278.x

Patterson, J.O., Rees, P., Nurse, P. (2019). Noisy Cell-Size-Correlated Expression of

Cyclin B Drives Probabilistic Cell-Size Homeostasis in Fission Yeast. Current Biology

29, 1379-1386.e4. https://doi.org/10.1016/j.cub.2019.03.011

Perruca-Foncillas, R., Davidsson, J., Carlquist, M., Gorwa-Grauslund, M.F.

(2022). Assessment of fluorescent protein candidates for multi-color flow cytometry

analysis of Saccharomyces cerevisiae. Biotechnol Rep (Amst) 34, e00735.

https://doi.org/10.1016/j.btre.2022.e00735

Petersen, J., Hagan, I.M. (2003). S. pombe aurora kinase/survivin is required for

chromosome condensation and the spindle checkpoint attachment response. Curr Biol

13, 590–597. https://doi.org/10.1016/s0960-9822(03)00205-7

Petersen, J., Russell, P. (2016). Growth and the Environment of Schizosaccharomyces

pombe. Cold Spring Harb Protoc 2016, pdb.top079764.

https://doi.org/10.1101/pdb.top079764

Picard, D., Schena, M., Yamamoto, K.R. (1990). An inducible expression vector for

both fission and budding yeast. Gene 86, 257–261. https://doi.org/10.1016/0378-

1119(90)90287-2

Politi, N., Pasotti, L., Zucca, S., Casanova, M., Micoli, G., Cusella De Angelis,

M.G., Magni, P. (2014). Half-life measurements of chemical inducers for recombinant

gene expression. J Biol Eng 8, 5. https://doi.org/10.1186/1754-1611-8-5

Rasmussen, B., Noller, H.F., Daubresse, G., Oliva, B., Misulovin, Z., Rothstein,

D.M., Ellestad, G.A., Gluzman, Y., Tally, F.P., Chopra, I. (1991). Molecular basis of

tetracycline action: identification of analogs whose primary target is not the bacterial

ribosome. Antimicrob Agents Chemother 35, 2306–2311.

https://doi.org/10.1128/AAC.35.11.2306

Rosenfeld, N., Elowitz, M.B., Alon, U. (2002). Negative autoregulation speeds the

response times of transcription networks. J Mol Biol 323, 785–793.

https://doi.org/10.1016/s0022-2836(02)00994-4

Russell, P.R. (1983). Evolutionary divergence of the mRNA transcription initiation

mechanism in yeast. Nature 301, 167–169. https://doi.org/10.1038/301167a0

Rutherford, K.M., Lera-Ramírez, M., Wood, V. (2024). PomBase: a Global Core

Biodata Resource-growth, collaboration, and sustainability. Genetics 227, iyae007.

https://doi.org/10.1093/genetics/iyae007

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch,

T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B. et al. (2012). Fiji: an open-

source platform for biological-image analysis. Nat Methods 9, 676–682.

https://doi.org/10.1038/nmeth.2019

Shiroiwa, Y., Hayashi, T., Fujita, Y., Villar-Briones, A., Ikai, N., Takeda, K., Ebe,

M., Yanagida, M. (2011). Mis17 is a regulatory module of the Mis6-Mal2-Sim4

centromere complex that is required for the recruitment of CenH3/CENP-A in fission

yeast. PLoS One 6, e17761. https://doi.org/10.1371/journal.pone.0017761

Smith, G.R. (2009). Genetic Analysis of Meiotic Recombination in

Schizosaccharomyces pombe. Methods Mol Biol 557, 65–76.

Stringer, C., Wang, T., Michaelos, M., Pachitariu, M. (2021). Cellpose: a generalist

algorithm for cellular segmentation. Nat Methods 18, 100–106.

https://doi.org/10.1038/s41592-020-01018-x

Sunder, S., Greeson-Lott, N.T., Runge, K.W., Sanders, S.L. (2012). A new method

to efficiently induce a site-specific double-strand break in the fission yeast

Schizosaccharomyces pombe. Yeast 29, 275–291. https://doi.org/10.1002/yea.2908

Tajadura, V., García, B., García, I., García, P., Sánchez, Y. (2004).

Schizosaccharomyces pombe Rgf3p is a specific Rho1 GEF that regulates cell wall

beta-glucan biosynthesis through the GTPase Rho1p. J Cell Sci 117, 6163–6174.

https://doi.org/10.1242/jcs.01530

Thodberg, M., Thieffry, A., Bornholdt, J., Boyd, M., Holmberg, C., Azad, A.,

Workman, C.T., Chen, Y., Ekwall, K., Nielsen, O. et al. (2019). Comprehensive

profiling of the fission yeast transcription start site activity during stress and media

response. Nucleic Acids Res 47, 1671–1691. https://doi.org/10.1093/nar/gky1227

Tommasino, M., Maundrell, K. (1991). Uptake of thiamine by Schizosaccharomyces

pombe and its effect as a transcriptional regulator of thiamine-sensitive genes. Curr

Genet 20, 63–66. https://doi.org/10.1007/BF00312766

Torres-Garcia, S., Yaseen, I., Shukla, M., Audergon, P.N.C.B., White, S.A.,

Pidoux, A.L., Allshire, R.C. (2020). Epigenetic gene silencing by heterochromatin

primes fungal resistance. Nature 585, 453–458. https://doi.org/10.1038/s41586-020-

2706-x

Toyama, R., Okayama, H. (1990). Human chorionic gonadotropin alpha and human

cytomegalovirus promoters are extremely active in the fission yeast

Schizosaccharomyces pombe. FEBS Lett 268, 217–221. https://doi.org/10.1016/0014-

5793(90)81012-d

Vještica, A., Marek, M., Nkosi, P.J., Merlini, L., Liu, G., Bérard, M., Billault-

Chaumartin, I., Martin, S.G. (2020). A toolbox of stable integration vectors in the

fission yeast Schizosaccharomyces pombe. Journal of Cell Science 133, jcs240754.

https://doi.org/10.1242/jcs.240754

Waisman, A., Norris, A.M., Elías Costa, M., Kopinke, D. (2021). Automatic and

unbiased segmentation and quantification of myofibers in skeletal muscle. Sci Rep 11,

11793. https://doi.org/10.1038/s41598-021-91191-6

Wang, H., Tang, X., Liu, J., Trautmann, S., Balasundaram, D., McCollum, D.,

Balasubramanian, M.K. (2002). The multiprotein exocyst complex is essential for cell

separation in Schizosaccharomyces pombe. Mol Biol Cell 13, 515–529.

https://doi.org/10.1091/mbc.01-11-0542

Wang, Hongcheng, Wang, Haiyang, Wang, M., Zhang, L., Wang, R., Mei, Y.,

Shao, W. (2014). Identification and refinement of two strong constitutive promoters for

gene expression system of Schizosaccharomyces pombe. World J Microbiol Biotechnol

30, 1809–1817. https://doi.org/10.1007/s11274-014-1603-6

Wang, J., Eisenstatt, J.R., Audry, J., Cornelius, K., Shaughnessy, M., Berkner,

K.L., Runge, K.W. (2018). A Heterochromatin Domain Forms Gradually at a New

Telomere and Is Dynamic at Stable Telomeres. Mol Cell Biol 38, e00393-17.

https://doi.org/10.1128/MCB.00393-17

Watson, A.T., Werler, P., Carr, A.M. (2011). Regulation of gene expression at the

fission yeast Schizosaccharomyces pombe urg1 locus. Gene 484, 75–85.

https://doi.org/10.1016/j.gene.2011.05.028

Watt, S., Mata, J., López-Maury, L., Marguerat, S., Burns, G., Bähler, J. (2008).

urg1: A Uracil-Regulatable Promoter System for Fission Yeast with Short Induction

and Repression Times. PLOS ONE 3, e1428.

https://doi.org/10.1371/journal.pone.0001428

Westwood, P.K., Martin, I.V., Fantes, P.A. (2004). Fission yeast Cdc37 is required

for multiple cell cycle functions. Mol Genet Genomics 271, 82–90.

https://doi.org/10.1007/s00438-003-0958-4

Win, T.Z., Goodwin, A., Hickson, I.D., Norbury, C.J., Wang, S.-W. (2004).

Requirement for Schizosaccharomyces pombe Top3 in the maintenance of chromosome

integrity. J Cell Sci 117, 4769–4778. https://doi.org/10.1242/jcs.01351

Wray, L.V., Reznikoff, W.S. (1983). Identification of repressor binding sites

controlling expression of tetracycline resistance encoded by Tn10. J Bacteriol 156,

1188–1191. https://doi.org/10.1128/jb.156.3.1188-1191.1983

Yamamoto, A., Hiraoka, Y. (2003). Monopolar spindle attachment of sister

chromatids is ensured by two distinct mechanisms at the first meiotic division in fission

yeast. The EMBO Journal 22, 2284–2296. https://doi.org/10.1093/emboj/cdg222

Yue, M., Zeng, L., Singh, A., Xu, Y. (2014). Rad4 mainly functions in Chk1-mediated

DNA damage checkpoint pathway as a scaffold protein in the fission yeast

Schizosaccharomyces pombe. PLoS One 9, e92936.

https://doi.org/10.1371/journal.pone.0092936

Zhang, X.-R., He, J.-B., Wang, Y.-Z., Du, L.-L. (2018). A Cloning-Free Method for

CRISPR/Cas9-Mediated Genome Editing in Fission Yeast. G3 (Bethesda) 8, 2067–

2077. https://doi.org/10.1534/g3.118.200164

Zhang, X.-R., Zhao, L., Suo, F., Gao, Y., Wu, Q., Qi, X., Du, L.-L. (2022). An

improved auxin-inducible degron system for fission yeast. G3 Genes|Genomes|Genetics

12, jkab393. https://doi.org/10.1093/g3journal/jkab393

Zhang, Y., Boyd, S.A., Teppen, B.J., Tiedje, J.M., Li, H. (2014). Role of tetracycline

speciation in the bioavailability to Escherichia coli for uptake and expression of

920 antibiotic resistance. Environ Sci Technol 48, 4893–4900.

921 https://doi.org/10.1021/es5003428

922 **Zilio, N., Wehrkamp-Richter, S., Boddy, M.N.** (2012). A new versatile system for

923 rapid control of gene expression in the fission yeast Schizosaccharomyces pombe. Yeast

924 29, 425–434. https://doi.org/10.1002/yea.2920

Figure legends

Fig. 1. Overview of the new tetracycline-inducible expression system.

(A) Schematic showing how the *enotetS* promoter (*PenotetS*) was generated. *PenotetS* was

engineered from the *eno101* promoter (*Peno101*) by introducing a single 19-bp *tetO*

sequence between the TATA box and the transcription start site (TSS) of *Peno101*, while

maintaining the original distance between the TATA box and the TSS.

(B) Diagram of the integration plasmid pDB5318. This plasmid contains *PenotetS* and a

TetR-expressing cassette, which consists of the cytomegalovirus promoter (*P.CMV*) and

the TetR coding sequence. The plasmid can be integrated at the *ade6* locus following

linearization with NotI digestion. The mECitrine coding sequence can be replaced with

the sequence of any desired gene to be expressed under the control of *PenotetS*, using

the NheI and BglII restriction sites.

(C) Schematic of four weakened variants of *PenotetS*. These variants were created by

inserting a GC-rich stem-loop-forming sequence downstream of *PenotetS*. Bases

forming the stem are highlighted in red, while those forming the loop are shown in black.

(D) Fluorescence of cells expressing the yellow fluorescent protein mECitrine from each of

the *enotetS* series promoters and from the *adh1* promoter in the presence and absence

of the inducer anhydrotetracycline (ahTet). Cells were imaged after 24 hours of growth in

YES medium in the presence of ahTet or DMSO (solvent control). The images are

presented with two different display ranges to enhance the visualization of differences in

fluorescence intensities. Scale bar, 10 μm.

Fig. 2. Inducer dose response of *PenotetS*.

(A) Inducer dose response in YES medium. Cells expressing mECitrine from *PenotetS* were

grown in YES medium in the presence of different doses of the inducer ahTet and then

imaged. Scale bar, 10 µm.

(B) Quantification of mECitrine fluorescence in the imaging data from the experiment shown

in A. The fluorescence level in the presence of 2.5 µg/ml ahTet was used as a reference

for normalization (100% expression level).

(C) Inducer dose response in EMM medium. Cells expressing mECitrine from *PenotetS* were

grown in EMM medium in the presence of different doses of the inducer ahTet and then

imaged. Scale bar, 10 µm.

(D) Quantification of mECitrine fluorescence in the imaging data from the experiment shown

in C. The fluorescence level in the presence of 2.5 µg/ml ahTet was used as a reference

for normalization (100% expression level).

Fig. 3. Induction kinetics of *PenotetS* and *PenotetSW2* in dividing cells.

(A) Time-course analysis of the induction kinetics of *PenotetS* in log-phase cells grown in

YES and EMM media. Images of cells expressing mECitrine from *PenotetS* were taken

at the indicated time points after the addition of 2.5 µg/ml ahTet to the culture. Scale bar,

10 µm.

(B, C) Quantification of mECitrine fluorescence in the imaging data from the experiment

shown in A. The fluorescence level at the 24-hour time point was used as a reference for

normalization (100% expression level). The curves depict the mathematical model of the

induction kinetics, with the modeled half-life value presented beneath each curve.

(D, E) Induction kinetics of *PenotetSW2* in log-phase cells grown in YES and EMM media.

Data processing and presentation were carried out as described in (B, C).

Fig. 4. Expression levels of different promoters.

(A) Expression levels in YES medium. Cells expressing mECitrine from *Padh1* and from each of the *enotetS* series promoters were grown in YES medium under induced conditions (abbreviated as ind., in the presence of ahTet) and repressed conditions (abbreviated as rep., in the absence of ahTet), and then imaged. The quantification results from two biological replicates and their mean values are shown. A thick dashed line represents a threshold (THR) below which the mECitrine fluorescence level cannot be accurately determined. This threshold corresponds to 2% of the average background fluorescence in a parental strain lacking mECitrine.

(B) Expression levels in EMM medium. Cells expressing mECitrine from each of the *enotetS* series promoters were grown in EMM medium under induced conditions (in the presence of ahTet) and repressed conditions (in the absence of ahTet), and then imaged. Cells expressing mECitrine from each of the *nmt1* series promoters were grown in EMM medium under induced conditions (in the absence of thiamine) and repressed conditions (in the presence of thiamine), and then imaged. The images are presented with two different display ranges to enhance the visualization of differences in fluorescence intensities. Scale bar, 10 µm.

(C) Quantification of mECitrine fluorescence in the imaging data from the experiment shown in B. The quantification results from two biological replicates and their mean values are shown.

(D) The induced expression level based on the mECitrine fluorescence (normalized to that of *Pnmt1*) and the estimated number of mRNA molecules per cell based on the reported value of 340 for the number of mRNA molecules per cell for the *nmt1* gene.

Fig. 5. Induction kinetics of *PenotetS* in quiescent cells.

(A) Time-course analysis of the induction kinetics of *PenotetS* in G0 cells in EMM−N medium.

Images were taken at the indicated time points after the addition of 2.5 µg/ml ahTet. Scale

bar, 10 µm.

(B) Time-course analysis of the induction kinetics of *PenotetS* in spores in EMM−N medium.

Images were taken at the indicated time points after the addition of 2.5 µg/ml ahTet. Scale

bar, 10 µm.

(C) Quantification of mECitrine fluorescence in the imaging data from the experiments shown

in A and B, as well as a control experiment in which log-phase cells were grown in EMM

medium. Three biological replicates were conducted, and one representative dataset is

shown here. The graph on the left uses a linear scale for the y-axis, while the graph on

the right uses a logarithmic scale. The curves depict the mathematical model of the

induction kinetics, with the modeled half-life values presented in the graph on the right.

1012

Fig. 6. A new method for synchronous meiosis induction based on the expression of

***Mc-2A-Mi* from *PenotetSW1* in *h+/h+* diploids.**

(A) Schematic illustrating the new method for synchronous meiosis induction.

(B) Progression of meiosis monitored by DNA staining. Cell samples were stained with the

DNA dye SYTOX Green and imaged at the indicated time points after the addition of

ahTet. Manual counting was performed to determine the percentages of cells containing

different numbers of nuclei. Three biological replicates were conducted, and one

representative dataset is shown here.

(C) Viability of spores from induced meiosis. Tetrad analysis was performed on YES plates.

1022

Fig. 7. Generation of conditional loss-of-function mutants using the *enotetS* series

promoters.

(A) Tetrad analysis of heterozygous deletion diploids carrying an integration plasmid

expressing the deleted gene from either *PenotetSW2* or *PenotetSW3*. Diploids were

sporulated, and the resulting tetrads were dissected on YES plates with and without

ahTet. Four spores from a single ascus are labeled A, B, C, and D. The genotype of

inviable spores was inferred based on the assumption of 2:2 segregation. Induced

expression of *sui1* and *nrs1* rescued the lethality of the respective deletions, while

repressed expression did not.

(B) Growth curve analysis showing that the growth defect caused by shutting off gene

expression through the removal of ahTet arises rapidly. Cells were initially cultured for

24 hours in YES liquid medium containing ahTet. Following this, cells were washed

three times with water and transferred to both ahTet-containing and ahTet-free YES

media. The optical density at 600 nm (OD600) of each culture was measured at the

indicated time points. Cultures were diluted when the OD600 approached 1.0 to

maintain logarithmic growth. The number of generations was calculated based on the

OD600 measurements and the dilution ratio.

1040

Fig. 8. Schematics depicting the improved tetracycline-inducible expression system

developed in this study.

(A) Schematic illustrating the repressed and induced states of the *enotetS* promoter. In the

absence of the inducer ahTet, TetR binds to *tetO*, repressing the transcription of the

downstream gene. Upon the addition of ahTet, the ahTet-bound TetR undergoes a

conformational change, preventing it from binding to *tetO* and thereby allowing the

transcription of the downstream gene to be induced. The induced state can be reverted

back to the repressed state by removing ahTet.

(B) Schematic illustrating the five distinct expression levels of the *enotetS* series promoters.

Four different stem-loop-forming sequences were individually inserted immediately

downstream of *PenotetS*, resulting in four weakened variants: *PenotetSW1*, *PenotetSW2*,

PenotetSW3, and *PenotetSW4*. The degree of expression attenuation increases with the

number of GC base pairs in the stem.

1054

A

B

C

	Promoter design	Stem-loop sequence	Predicted stem-loop
PenotetS	*tet*O — TSS — mECitrine	–	–
PenotetSW1	*tet*O — TSS — stem-loop — mECitrine	GCC GCTCG GGC	
PenotetSW2	*tet*O — TSS — stem-loop — mECitrine	GGCC GCTCG GGCC	
PenotetSW3	*tet*O — TSS — stem-loop — mECitrine	GGGCC GCTCG GGCCC	
PenotetSW4	*tet*O — TSS — stem-loop — mECitrine	CGGGCC GCTCG GGCCCG	

D

A YES + ahTet (µg/ml)

B *PenotetS* (YES)

C EMM + ahTet (µg/ml)

D *PenotetS* (EMM)

A

B

PenotetS (YES)

Expression level (%)

68% 76% 94% 98% 110%
31%
9.9%
1.9%
0.51%
100%

Half-life = 2.64 ± 0.28 h

Induction time (h)

C

PenotetS (EMM)

Expression level (%)

57% 71% 91% 95% 110%
19%
4.8%
1.2%
0.44%
100%

Half-life = 3.40 ± 0.52 h

Induction time (h)

D

PenotetSW2 (YES)

Expression level (%)

50% 69% 86% 89% 99%
28%
8.4%
1.9%
0.26%
100%

Half-life = 3.37 ± 0.23 h

Induction time (h)

E

PenotetSW2 (EMM)

Expression level (%)

57% 72% 94% 95% 108%
20%
4.8%
1.2%
0.55%
100%

Half-life = 3.33 ± 0.50 h

Induction time (h)

A YES

B EMM

C EMM

D

	Pnmt1	*PenotetS*	*PenotetSW1*	*PenotetSW2*	*PenotetSW3*	*PenotetSW4*
Induced expression level	1	1.073	0.643	0.168	0.039	0.023
Estimated mRNA level	340	365	219	57	13	8

A

h+/h+ diploid

PenotetSW1-Mc-2A-Mi

YES
cycling cells
30°C

EMM−N
G0 arrest
30°C

EMM−N+ahTet
meiosis
30°C

B

PenotetSW1-Mc-2A-Mi

Cell type (%)

- 1 nucleus
- 2 nuclei
- 3-4 nuclei

Induction time (h)

C

Tetrad analysis

Spore viability = 97%
(n = 192)

A

B