Coexistence despite competition: Effects of vegetation structural complexity on

bryophyte diversity over succession.

Maximilian Hanusch[1,2]

[1]Department of Environment and Biodiversity, Paris-Lodron-University Salzburg, 5020

Salzburg, Austria

[2]Evolutionary Ecology of Plants, Department of Biology, Philipps-University Marburg, 35043

Marburg, Germany

*Correspondence: maximilian.hanusch@uni-marburg.de

9 Abstract

Vascular plants and bryophytes are important drivers underlying the development of diverse

ecosystems following glacial retreat. Understanding biotic interactions between both groups

is crucial for understanding the generation of biodiversity under natural conditions. Focusing

on a 170-year primary succession gradient in the Austrian Alps, this study examines the

mechanisms governing the interactions between vascular plants and bryophytes during the

initial phases of ecosystem establishment. Using data on the local habitat's substrate

diversity, vascular plant and bryophyte taxonomic composition, and vegetation structural

complexity derived from multispectral 3D-scans of vegetation, this study provides a

mechanistic understanding of plant-plant interactions within a natural ecosystem. Vascular

plants dominate the vertical niche through complex growth structures, which negatively affect

bryophyte diversity. A pattern that is uniquely detectable by estimates of vegetation structural

complexity. Path analysis confirms that plant growth structure drives these competitive via a

reduction of bryophyte communities' realized niche sizes, a concept introduced here as

ecological dispersion. This study demonstrates the advantages of 3D-laser-scanning

approaches in capturing the complexity of vegetation structure and its impacts on ecosystem

dynamics. The findings underscore the importance of vegetation structural complexity in the

establishment and maintenance of biodiversity and ecosystem complexity and underscore

the potential of 3D-laser-scanning methods in future studies to inform biodiversity

conservation and restoration approaches in similar environments.

Introduction

Although the importance of physical vegetation structure for ecosystem functioning and

biodiversity was recognized early in ecological research (MacArthur, MacArthur and Preer,

1962; Odum, 1969), research in this area has only recently gained momentum, driven largely

by advances in remote sensing technology (LaRue, Fahey, *et al.*, 2023). Tools such as

portable 3D laser scanners now allow to accurately the arrangement of physical vegetation

structures at various scales and resolutions and the modeling of vegetation metrics from data

that previously required labor-intensive fieldwork or destructive field-sampling (Catchpole and

Wheeler, 1992; Gámez and Harris, 2022; Fei *et al.*, 2023; Zieschank and Junker, 2023).

Metrics of vegetation structural complexity are considered valuable surrogates for quantifying

the occupancy of plants in 3D-niche space (LaRue, Fahey, *et al.*, 2023), which is closely

associated with ecosystem functioning and biotic interactions with other organisms

(Srivastava, 2006). As it increases niche diversity, promotes resource partitioning, and

expands the three-dimensional habitat space for other organisms, vegetation structural

complexity is widely considered beneficial for biodiversity. Accordingly, positive associations

between structural complexity and species diversity have been demonstrated for plant

(Coverdale and Davies, 2023), animal (Tews *et al.*, 2004; Traylor *et al.*, 2022), and

belowground diversity (Lang et al., 2023). However, most of these insights are derived from

forest ecosystems, where trees create strong vertical stratification with a distinct vertical

layering (Davies and Asner, 2014; Gough *et al.*, 2019). Grasslands, in contrast, exhibit

limited vertical stratification, as most herbaceous plants compete horizontally within the same

layer for light and other resources. For instance, when competing with smaller-growing

plants, such as bryophytes, some vascular plants are expected to possess a competitive

advantage due to their larger size and greater structural robustness. This enables them to

dominate the vertical axis, thereby restricting bryophytes' access to essential resources like

light and space. Thus, the condensed horizontal structure of grasslands challenges the

assumption that vegetation structural complexity universally promotes the diversity of

associated organismal groups within a shared habitat.

Successional gradients, such as those developing after glacial retreat, provide an ideal

opportunity to study how changes in grassland structure impact other organisms or

interactions within the ecosystem. Primary successions are characterized by a gradual

increase in vascular plant biomass and associated vegetation structural complexity which in

turn will lead to biotic interactions with bryophytes. However, estimating biotic interactions

from observational field-data is notoriously challenging, and particularly so in the context of

ecological succession (Blanchet, Cazelles and Gravel, 2020; Hanusch *et al.*, 2023, 2024).

Accordingly, field observations should be complemented by clear assumptions on the

ecological mechanisms driving these interactions to enhance interpretation and reliability:

As succession progresses, vascular plants increasingly dominate the shared habitat by

developing more complex growth structures. These structures form dense mats that occupy

much of the available space and significantly alter local habitat conditions (Breen and

Lévesque, 2006; Cutler, Belyea and Dugmore, 2008; Gavini *et al.*, 2019). Within these mats

more uniform microenvironmental conditions prevail that are marked by altered temperature,

light, and moisture levels, ultimately affecting the growth and diversity of bryophytes growing

within them (van Tooren *et al.*, 1990; Bergamini *et al.*, 2001; Werger *et al.*, 2002; Fergus,

Gerighausen and Roscher, 2017). Bryophyte communities are expected to respond

dynamically to these changes in local environmental conditions, with corresponding shifts in

their realized ecological niches. Ecological Indicator Values (EIVs), ordinal scales

representing species' growth optima along environmental gradients, provide a tool for

assessing such shifts (Diekmann, 2003). Assessing the dispersion of growth optima across

multiple environmental factors at the community scale, provides a measure that

approximates the community's multidimensional realized niche. High dispersion indicates a

diverse habitat capable of supporting species with a variety of ecological preferences,

whereas low dispersion indicates more homogenous habitats favoring species with similar

ecological preferences. Accordingly, the dispersion of growth optima of a community,

introduced here as *ecological dispersion*, serves as a framework for exploring how

vegetation structure homogenizes local habitat conditions, ultimately impacting bryophyte

diversity in the process.

Studying structural properties of vascular plant communities in grassland usually required

destructive field measurements that can significantly alter or destroy the plant communities

(Catchpole and Wheeler, 1992; Zehm, Nobis and Schwabe, 2003). As an alternative to

traditional destructive methods, the use of 3D multispectral plant scanning technology

provides a non-invasive way to capture morphological features of entire plant communities.

The scans allow to obtain various community features (Zieschank and Junker, 2023), such

as digital biomass or estimates of vegetation structure, which, similar to community-weighted

means, represent the trait frequency distribution across an entire community. Estimating

such community features offers new opportunities to study how vegetation structural

complexity affects fine-scale ecological processes, while circumventing labor-intensive

species-specific, or individual-based measurements (He, Hanusch, Saueressig, *et al.*, 2023).

Focusing on a 170-year primary succession gradient in the Ödenwinkel glacier forefield,

Austrian Alps, this study investigates the role of vascular plant structural complexity in

shaping bryophyte diversity during the development of a natural ecosystem. It is shown that

vegetation structural complexity has a negative effect on bryophyte diversity which is caused

by a reduction of the bryophyte communities' realized niche sizes during succession.

Comparing estimates of structural complexity obtained through whole-community 3D

multispectral plant scanning with traditional measures of vascular plant communities, such as

species composition, plant cover, and taxonomic or functional diversity, revealed that

structural complexity uniquely identifies these competitive interactions. By capturing the

spatial and physical arrangement of vegetation, structural complexity offers a clearer

understanding of the natural dynamics driving the development of biodiverse ecosystems.

Figure 1: Conceptual overview of the successional increase in vascular plant structural complexity. Each triplet, comprising a photograph, 3D conceptual sketch, and 3D plant scan, represents a plot with increasing successional age and vegetation structural complexity. In the conceptual sketches, bryophyte patches are highlighted in blue to illustrate changes in the habitats exploited by bryophyte species and their realized ecological niches in response to increasing plant structure. As structural complexity increases, it homogenizes local habitat conditions, while substrate weathering facilitates bryophyte colonization of habitats such as bare rock sticking out from the vegetation. This process buffers bryophyte diversity against vascular plant dominance over time. (A) Photograph, sketch and scan of a plot with low vegetation structural complexity and high proportion of open substrate or rocks, approximate time since deglaction: 6 years (B) medium structural complexity, approximate time since deglaction: 35 years (C) high structural complexity, approximate time since deglaction: 100 years. 3D-scans adapted from He *et al*., (2023) and sketches published by OpenClipArt, permission granted under CC BY 3.0 and CC0 1.0 Universal.

Methods

Study design The study was conducted in the long-term ecological research platform Ödenwinkel which was established in 2019 in the Hohe Tauern National Park, Austria (Dynamic Ecological Information Management System – site and dataset registry: https://deims.org/activity/fefd07db-2f16-46eb-8883-f10fbc9d13a3, last access: November 2024; Junker *et al.*, 2020). A total of *n* = 135 permanent plots were established within the glacier forefield of the Ödenwinkelkees, which was covered by ice at the latest glacial maximum in the Little Ice Age around 1850. The plots represent a successional gradient

spanning over 1.7km in length at an altitude ranging from 2070 to 2170 m a.s.l. with a

minimum distance of 5 m between plots. Plots were defined as squares with an area of 1x1

m and were all oriented in the same cardinal direction. A full dataset, i.e. no NA value for any

variable, was used for this study representing a subset of $n = 107$ out the total $n = 135$ plots.

Detailed descriptions of the research platform, exact plot positions and details on the

surrounding environment can be found in Junker et al. (2020) and Hanusch et. al (2022).

Biotic and abiotic sampling.

Plot age was estimated according to plots' relative position compared with historical records

of eight deglaciation periods (year 1850, 1890, 1929, 1969, 1977, 1985, 1998, 2013; Junker

et al., 2020). To assess the substrate composition of the plots, the percentage cover of bare

ground, litter, scree and solid rock were estimated.

During the 2019 field season, all bryophyte and vascular plant species on each plot were

identified and recorded ($n = 42$ in 135 plots), noting their coverage at a 0.1% resolution. This

was accomplished by using 1m² vegetation relevés subdivided into 100 squares, each

measuring 10 × 10 cm. Likewise, all vascular plant species (n = 93 across 135 plots) were

recorded (He, Hanusch, Ruiz-Hernández, *et al.*, 2023). Both communities were sampled in

the same plots, which allows a direct comparison of potential interactions between both

groups (Junker *et al.*, 2021). Furthermore, vascular plant functional traits were measures

based on estimates of leaf area and weight, specific leaf area, and growth height as

described in Junker *et al.* (2020) and He, Hanusch, Saueressig, *et al.* (2023).

Plant community phenotyping and vegetation structural complexity.

In the vegetation period 2021 (from 16 July to 23 July), an on-site plot scanning in the field

was conducted with a multispectral 3D scanner 'PlantEye' (Phenospex, F500, Heerlen, the

Netherlands). For a detailed description of the set-up and data obtained, see Zieschank and

Junker, (2023) and Supplemental Figure 1. The plant eye is equipped with an active sensor

that projects a near-infrared (940 nm) laser line vertically onto the plant canopy (Kjaer and

Ottosen, 2015; Maphosa *et al.*, 2017). Each scan has a resolution of < 1 pixel*mm^{-1} and the

data points of each scan are merged to generate a 3D point cloud. From these clouds, a set

of community level morphological features can be estimated that have been shown to

correlate well with real world manual measurements (Zieschank and Junker, 2023). In this

study digital community measurements of biomass (BM), leaf area index (LAI) and growth

height-corrected light penetration depth (LPD). Settings of Software used to calculate

community features followed He *et al.* (2023). LPD is a measure for the depth to which light

can penetrate through the vegetation layer. The plant eye calculates this value as the

distance between the on average highest 10 % and on average lowest 20 % parts of the

vegetation layer that the light can penetrate through. As a result, the uncorrected, raw

measurements of LPD are not independent of the vegetation's growth height. Thus, a growth

height-corrected LPD-index was calculated by subtracting LPD from the on average highest

10 %. This index is an estimate of the height of the lowest and most dense 20 % of the

vegetation in which most bryophyte species grow within the vegetation mats. Low growth

height-corrected LPD values indicate a shorter, scattered vegetation; high values indicate a

higher, more dense vegetation. The LAI measures the density of vascular plant leaves in a

given area, representing the ratio of one-sided leaf area to ground area (Coverdale and

Davies, 2023); i.e. high values indicate a denser vegetation with a high number of leaves,

low numbers indicate a sparse vegetation with a low number of leaves. The estimates of BM

represent a digital approximation of the standing biomass, and is calculated as the product of

height and 3D leaf area.

Vegetation structural complexity is not directly observable and there is still no consensus

over the definition of structural complexity or how to measure it (Atkins *et al.*, 2018; Loke and

Chisholm, 2022). In this study vegetation structural complexity is quantified as a univariate

variable generated through a principal component analysis (PCA; R package 'vegan' v2.6-4;

Oksanen et al., 2019) conducted on estimates of BM, LAI, and growth height-corrected LPD

derived from 3D laser scanning. The first principal component (PC1), which explained 84.0%

of the variance, was used as the structural complexity metric.

Traditional metrics of vascular plant communities

To examine whether vegetation structural complexity adds additional information or is

redundant to traditional vascular plant community metrics in explaining bryophyte diversity,

multiple metrics were assessed: vascular plant species cover, species richness, taxonomic

composition, functional diversity. Species richness was calculated as the number of vascular

plant species per plot, taxonomic composition was summarized using the first principal

component (15% explained variance) from a PCA of vascular plant species composition.

Functional diversity was quantified as functional divergence using the fd_fdiv command in

the fundiversity-package (v1.1.1, R) based on trait measurements of leaf area and weight,

specific leaf area, and height. High values of functional divergence suggest functional niche

differentiation, which may capture similar aspects of spatial niche occupancy as structural

complexity. Species cover was measured as the total horizontal extent of vascular plants on

each plot. Pearson correlations were conducted to evaluate the relationships among these

metrics and structural complexity, as well as with bryophyte diversity.

Calculation of ecological dispersion

Ecological dispersion provides a novel framework for examining biotic interactions by

highlighting shifts in community tolerances driven by changing habitat conditions. Vascular

plant structural complexity acts as an important ecological feature that alters local habitat

conditions, which can be analyzed through the niche optima of bryophyte species growing

among the vascular plant layer on a plot. High ecological dispersion values indicate diverse

environmental conditions that support species with varying ecological preferences, whereas

low values reflect more homogenous habitats favoring species with similar ecological

preferences. To calculate ecological dispersion of bryophyte communities, EIV data for

bryophytes were obtained from the 2017 updated version of BryoATT (Preston and

Bosanquet, 2007). In total, EIV assignments for L (light), R (pH), F (humidity), and N

(nutrients) were successfully made for n = 32 bryophyte species; i.e. 76% of all recorded

bryophyte taxa.

To evaluate whether the selected Ecological Indicator Values (EIVs) represent ecologically

meaningful variables, the relationship between each EIV and the structural complexity

variable was assessed individually using Spearman's rank correlation. Spearman correlation

was chosen because a linear relationship was not expected between the community-

weighted means and structural complexity. This is due to the community-weighted means

being largely influenced by the occurrence of specific species with associated EIV values,

which can cause abrupt shifts in the community means. Community-weighted means of F

and N showed a significant positive relationship with structural complexity ($rho_F = 0.19$, $p_F =$

0.05; $rho_N = 0.23$, $p_N = 0.01$), while community-weighted means of L showed a significant

negative relationship ($rho_L = -0.23$, $p_L = 0.01$), and no significant relationship with R ($rho_R = -$

0.13, $p_R = 0.17$).

In a next step, ecological dispersion was calculated as follows: First the ordinal EIV for each

species were transformed into a continuous form. A multidimensional functional space was

built using principal coordinate analysis (ape v5.7-1 package, R) based on Euclidean-

distances of the four EIV-values (Maire *et al.*, 2015; De Bello *et al.*, 2021). Second,

ecological dispersion was calculated based on the axes coordinate of each species in each

of the four dimensions by using fd_fdis command in the fundiversity-package (v1.1.1, R).

Path analysis

To infer the effects of vascular plant structural complexity on bryophyte diversity, a path

analysis was conducted using the R package lavaan (v0.6-1.7; Rosseel, 2012; Tsukahara,

2023). Path analysis was selected for its ability to disentangle direct and indirect effects

among multiple interacting variables and obtain estimates of the explained variance (R^2) by

predictors. The model included key variables hypothesized to influence bryophyte diversity:

Time since deglaciation was included as a proxy for plot age, representing the increasing

likelihood of heterogeneous dispersal events over time (Hanusch *et al.*, 2022; Ficetola *et al.*,

2024). Vegetation structural complexity served as a proxy for competitive interactions,

capturing how the physical structure of vascular plant communities may restrict bryophytes

growing within the plant layer. To account for potential shared extrinsic factors affecting both

vascular plants and bryophytes, substrate diversity, calculated as the Shannon diversity of

relative cover of substrate types (bare ground, litter, scree, and solid rock), was included as a

univariate variable to represent environmental heterogeneity. Substrate diversity showed a

significant correlation with bryophyte and vascular plant species composition (Mantel-test:

Mantel $r_{bryophytes}$ = 0.26, $p_{bryophytes}$ < 0.001; Mantel $r_{vascular\ plants}$ = 0.36, $p_{vascular\ plants}$ < 0.001),

highlighting its relevance to both groups. As the study was conducted along a linear

successional gradient in a glacier forefield, spatial distance was highly correlated with time

since deglaciation (Mantel-test: Mantel r = 0.88, p < 0.001) and was therefore excluded from

the model to avoid multicollinearity. By incorporating these variables, the analysis aimed to

disentangle the mechanisms driving bryophyte diversity over succession. Specifically, it

sought to determine whether changes in bryophyte diversity are primarily shaped by

environmental heterogeneity, dispersal processes, or interactions with vascular plants. All

variables were standardized by z-transformation by subtracting the mean and dividing

through the standard deviation.

In the model, time since deglaciation is included as an exogenous variable, exerting direct

effects on vegetation structural complexity, substrate diversity, and bryophyte ecological

dispersion. Vegetation structural complexity and substrate diversity, in turn, have direct

impacts on bryophyte ecological dispersion, representing the influence of vascular plant

competition and environmental heterogeneity, respectively. Additionally, the model examines

indirect effects of time since deglaciation, vegetation structural complexity, and substrate

diversity on bryophyte diversity, mediated through ecological dispersion. Ecological

dispersion, representing the variation in the bryophyte communities' ecological niches, is

included to provide a mechanistic understanding of how structural and environmental factors

influence bryophyte diversity by restricting the number of species that are able to grow under

certain habitat conditions.

To evaluate whether vegetation structural complexity provides additional information

compared to traditional plant community metrics, four alternative model structures were

developed. In each model vegetation structural complexity was replaced with one of the

traditional plant community metrics: vascular plant cover, vascular plant species richness,

vascular plant species composition (first axis from PCA), or vascular plant functional

divergence. The comparison of these models aimed to determine whether structural

complexity captures unique aspects of plant-plant interactions that other metrics fail to

address.

Results

Comparison between vascular plant community metrics and their relationship with bryophyte diversity.

Structural complexity showed a significant negative relationship with bryophyte diversity ($t = -2.00$, $df = 104$, $r^2 = 0.04$, $p = 0.04$), which was not observed with the other metrics (Figure 2).

Plant cover exhibited a non-linear relationship with bryophyte diversity ($t = -2.68$, $df = 103$, $r^2 = 0.07$, $p < 0.01$), while plant species richness showed a marginally significant positive linear

relationship ($t = 1.80$, $df = 104$, $r^2 = 0.03$, $p = 0.06$). Neither species composition ($p = 0.64$)

nor functional divergence ($p = 0.50$) demonstrated a significant linear relationship with

bryophyte diversity. Among the vascular plant community metrics, structural complexity

showed the highest correlation with plant cover ($r = 0.78$, $p < 0.01$), followed by species

composition ($r = 0.58$, $p < 0.01$), species richness ($r = 0.38$, $p < 0.01$), and the lowest

correlation with functional divergence ($r = 0.23$, $p = 0.02$).

Figure 2: Relationship between bryophyte diversity (BD) and five metrics of the surrounding
vascular plant community across n = 107 plots along a primary succession gradient spanning
170 years. Darker points indicate older plots. Metrics of vegetation structural complexity
derived from three-dimensional laser scanning provide distinct information compared to
traditional two-dimensional estimates based on relative percentage cover per species (used
for vascular plant cover, community composition, and functional divergence) or vascular plant
species richness counts. (A) significant negative linear relationship between vascular plant
structural composition, estimated as the first axis of a PCA on digital biomass, leaf area
index, and height-corrected light penetration depth, and BD. (B) significant non-linear
relationship (quadratic fit) between vascular plant cover and BD. (C) marginally significant
positive linear relationship between vascular plant richness and BD (D) non-significant
relationship between vascular plant community composition and BD. (E) non-significant
relationship between vascular plant functional diversity and BD. (F) Pearson correlation
heatmap showing relationships among the five plant community metrics. The heatmap
highlights the degree of association between these metrics, providing insights into potential
overlaps or unique contributions of each measure.

Path analysis

The path analysis provided good model fit (Model fit: $p_{chi\text{-}square} > 0.05$; CFI > 0.95;

TLI > 0.9; SRMR < 0.05; RMSEA $= 0.08$ with lower bound of confidence interval = 0)

and revealed significant influences of all exogenous variables on bryophyte diversity (Figure

3). Time since deglaciation ($\beta = 0.25$, $p = 0.01$) and substrate diversity

($\beta = 0.20$, $p = 0.03$) positively affected bryophyte ecological dispersion. Conversely, plant

growth structure showed a significant negative correlation with ecological dispersion ($\beta = -$

0.29, $p < 0.01$) indicating that denser vegetation structure had a negative effect on the

realized niche size of the bryophyte community. Time since deglaciation had a negative

effect on substrate diversity ($\beta = -0.26$, $p < 0.01$) but a positive effect on plant growth

structure ($\beta = 0.42$, $p < 0.01$). Bryophyte diversity was positively affected by bryophyte

ecological dispersion ($\beta = 0.83$, $p < 0.01$) and experienced positive indirect effects by

substrate diversity ($\beta = 0.17$, $p = 0.03$) while plant growth structure had a negative indirect

effect ($\beta = -0.24$, $p < 0.01$). The total indirect effects of time since deglaciation on

ecological dispersion (i.e. temporal processes including effects of substrate diversity and

plant growth structure) were non-significant ($\beta = 0.07$, $p = 0.40$) indicating that several

simultaneous processes may act over time that offset each other. Thus, the indirect effects of

time since deglaciation on bryophyte diversity were separately calculated for the effects

mediated through substrate diversity ($\beta = -0.04$, $p = 0.09$), plant growth structure ($\beta = -$

0.10, $p = 0.02$), and time since deglaciation in isolation (i.e. temporal processes without

effects of substrate diversity and plant growth structure; $\beta = 0.21$, $p = 0.02$). The four

alternative models, in which vegetation structural complexity was replaced with one of the

four other vascular plant community metrics (vascular plant species richness, functional

diversity, cover, and composition) showed no significant effects of these variables on

bryophyte diversity or associated ecological dispersion, which highlights vegetation structural

complexity as the primary factor influencing bryophyte diversity (Supplementary tables 1-5).

Figure 3: Path analysis of environmental drivers and vegetation structural complexity on
bryophyte ecological dispersion and bryophyte diversity. The path diagram presents direct
effects as solid lines. Indirect effects are represented as dotted lines and were calculated
through mediation analysis. Standardized path coefficients are shown for significant
relationships, with red coloration indication a negative and black coloration indicating a
positive effect.

Discussion

Biotic interactions, particularly competition driven by the development of dense vascular plant

mats, emerge as a key driver of bryophyte diversity over succession.

This observation is explained by a mechanism where vascular plant physical structures

homogenize local habitat conditions, thereby reducing the multidimensional realized niche

size — or ecological dispersion — of bryophytes in shared habitats. Furthermore, traditional

metrics such as vascular plant cover, species composition, species richness, and functional

divergence failed to capture these -plant interactions. These finding underscore the

importance of vegetation structural complexity as a key determinant of biodiversity.

Vascular plants shape bryophyte succession through biotic interactions

Competition with vascular plants significantly reduces bryophyte diversity in grasslands.

These interactions arise through increasingly complex and dense physical structures of

vascular plants. (Legendre *et al.*, 2005; Lowe and McPeek, 2014; Hanusch *et al.*, 2022; He

et al., 2024). Path analysis revealed that vascular plant structural complexity negatively

impacts bryophyte diversity and ecological dispersion, resulting in bryophyte communities

with more homogeneous ecological preferences in habitats dominated by more complex

vascular plant structures. This is putatively driven by shifts in moisture, as well as intensified

competition for light and space along the vertical axis of the three-dimensional niche space.

Vascular plants overgrow and overshadow bryophytes, limiting the diversity of bryophyte

species able to persist by favoring a select set adapted to these constraints.

However, bryophyte diversity is partially buffered against vascular plant dominance over

succession by the availability of diverse growth substrates and time-driven dispersal

processes. Despite ongoing competition, bryophyte diversity remains higher on plots with

more diverse substrate compositions and on older plots. Bryophytes in glacier forefields are

generally not considered dispersal-limited (Cichini *et al.*, 2011), suggesting that of effect of

time since deglaciation cannot be solely attributed to slow dispersal processes, which are

more typical of vascular plant communities (Ulrich *et al.*, 2016; Hanusch *et al*., 2022; Cantera

et al., 2024). Instead, site-specific processes, such as substrate weathering, likely play a key

role by facilitating opportunistic and rapid colonization of newly available niches (Prach and

Walker, 2020). Bryophytes grow on a wide range of terrain features, such as rocks or

boulders that stick out of the vegetative layer. Recent research revealed a surprisingly high

degree of environmental heterogeneity even at fine spatial scales within grassland

ecosystems (Conti *et al.*, 2017; Price *et al.*, 2017). Glacier forefields, in particular, are highly

dynamic ecosystems that offer a variety of substrates, fostering the establishment of diverse

sets of bryophyte species. The positive effect of substrate diversity on bryophyte diversity

aligns with previous studies, confirming that substrate availability is a key driver of bryophyte

community composition and diversity during succession (e.g. Hylander *et al.*, 2005; Caruso

and Rudolphi, 2009; Chen *et al.*, 2017; Spitale, 2017). Accordingly, bryophyte succession in

alpine grasslands is driven by a combination of interrelated processes. The successful

colonization of microhabitats within a heterogeneous environment helps buffer bryophyte

diversity against competition with vascular plants over time.

Ecological dispersion as a measure for a community's ecological tolerance

Originally developed for vascular plants in Central Europe, EIVs have since been adapted for

bryophytesIn this study (Düll, 1992; Simmel, Ahrens and Poschlod, 2021). In this study, they

were applied to approximate the bryophyhte community's multidimensional realized niche.

Although EIV-based ecological dispersion does not capture the full range of functional traits

that influence ecological interactions, it proved to be a valuable tool for gaining mechanistic

insights into bryophyte community responses to changing biotic and abiotic conditions along

the successional gradient.

Adapting a "three-dimensional perspective" for ecological research

A key advantage of 3D-laser scanning techniques is their ability to estimate phenotypic

properties of entire plant communities in three dimensions, explicitly accounting for the

spatial arrangement of physical plant structures across both horizontal and vertical

dimensions. This capability represents a significant improvement over traditional vegetation

metrics, which are typically derived from two-dimensional assessments based on plant cover

or individual-based estimates of abundance (LaRue, Knott, *et al.*, 2023). Such two-

dimensional approaches often fail to capture the inherently three-dimensional aspects of

structural complexity, and as such overlook critical vertical features that strongly influence

ecological dynamics within an ecosystem (Gámez and Harris, 2022). For instance, size-

asymmetric competition for light and space is a well-documented driver of grassland

biodiversity (Newman, 1973; Weiner, 1990; Eskelinen *et al.*, 2022). Here, larger plants gain

disproportionate access to light, by overshadowing smaller plants and limiting their growth.

This form of competition primarily operates along the vertical axis of the three-dimensional

habitat space, as it is fueled by the directional supply of light. In this study, structural

complexity included measurements of light penetration depth, which refers to the extent light

can pass through the upper vegetation layers to reach the lower layers or ground. This metric

serves as an important indicator of how the vertical structure of the plant community

influences light availability for understory species (Grace, 1999; Zieschank and Junker,

2023). Accordingly structural complexity provides a more realistic representation of light

availability and spatial constraints that bryophytes experience within the vegetation layer. As

a result, structural complexity proved to be the only metric that captured a significant

negative correlation with bryophyte diversity in this study. This finding highlights the

importance of assessing structural properties to understand the impact of vascular plant

structures on interaction outcomes and biodiversity

Although traditional vegetation metrics may not capture the specific dynamics revealed by

structural complexity, they offer complementary rather than redundant information. The

correlations between structural complexity and other traditional metrics demonstrate that

some metrics are more strongly related, while others capture distinct aspects of vascular

plant communities. Adopting a "three-dimensional perspective" certainly enables the

exploration of ecosystem dynamics at scales beyond individual or species-level analyses. At

the same time, metrics such as species richness and functional diversity capture other critical

aspects of vascular plant communities, highlighting different aspects of their ecological roles

and functions. Ecologists are thus encouraged to select metrics thoughtfully, as each will

provide distinct and complementary insights into ecosystem dynamics.

Conclusion

This research contributes to the ongoing integration of structural complexity as a key

component of biodiversity (Coverdale and Davies, 2023; LaRue, Knott, *et al.*, 2023). The

findings provide a mechanistic understanding of the interactions between vascular plants and

bryophyte communities, highlighting the complex dynamics of ecological succession. They

reveal that bryophyte diversity is buffered against vascular plant dominance by exploiting

diverse microhabitats that develop over time.

Advanced laser-scanning methods, and the modelling of vegetation structural complexity

metrics, offer valuable insights into understanding plant-plant interactions, coexistence and

ecosystem dynamics. Integrating structural metrics with traditional measures of vascular

plant communities offers a more comprehensive understanding of ecosystem dynamics,

extending beyond successional research. Such integrated approaches will deepen insights

into ecosystem processes and support the development of effective conservation and

succession-based restoration strategies.

Supplementary Information

Supplementary Figure 1: Digital whole community phenotyping (DWCP) using a customized
PlantEye setup (Phenospex, F500, Heerlen, the Netherlands) in the field Figure adapted
from He *et al.*, (2023), permission granted under CC BY 3.0.

Bibliography

Atkins, J.W. *et al.* (2018) 'Quantifying vegetation and canopy structural complexity from
terrestrial Li DAR data using the FORESTR R package', *Methods in Ecology and Evolution*.
Edited by S. Goslee, 9(10), pp. 2057–2066. Available at: https://doi.org/10.1111/2041-
210X.13061.

Bergamini, A. *et al.* (2001) 'Relationships between productivity, number of shoots and
number of species in bryophytes and vascular plants', *Journal of Ecology*, 89(6), pp. 920–
929. Available at: https://doi.org/10.1046/j.0022-0477.2001.00613.x.

Blanchet, F.G., Cazelles, K. and Gravel, D. (2020) 'Co-occurrence is not evidence of
ecological interactions', *Ecology Letters*. Edited by E. Jeffers, 23(7), pp. 1050–1063.
Available at: https://doi.org/10.1111/ele.13525.

Breen, K. and Lévesque, E. (2006) 'Proglacial succession of biological soil crusts and
vascular plants: biotic interactions in the High Arctic', *Canadian Journal of Botany*, 84(11),
pp. 1714–1731. Available at: https://doi.org/10.1139/b06-131.

Cantera, I. *et al.* (2024) 'The importance of species addition "versus" replacement varies over
succession in plant communities after glacier retreat', *Nature Plants*, 10(2), pp. 256–267.
Available at: https://doi.org/10.1038/s41477-023-01609-4.

Caruso, A. and Rudolphi, J. (2009) 'Influence of substrate age and quality on species
diversity of lichens and bryophytes on stumps', *The Bryologist*, 112(3), pp. 520–531.
Available at: https://doi.org/10.1639/0007-2745-112.3.520.

Catchpole, W.R. and Wheeler, C.J. (1992) 'Estimating plant biomass: A review of
techniques', *Australian Journal of Ecology*, 17(2), pp. 121–131. Available at:
https://doi.org/10.1111/j.1442-9993.1992.tb00790.x.

Chen, Y. *et al.* (2017) 'Stand Structure and Substrate Diversity as Two Major Drivers for
Bryophyte Distribution in a Temperate Montane Ecosystem', *Frontiers in Plant Science*, 8.
Available at: https://doi.org/10.3389/fpls.2017.00874.

Cichini, K. *et al.* (2011) 'Colonization of experimentally created gaps along an alpine
successional gradient', *Plant Ecology*, 212(10), pp. 1613–1627. Available at:
https://doi.org/10.1007/s11258-011-9934-y.

Conti, L. *et al.* (2017) 'Environmental gradients and micro-heterogeneity shape fine-scale
plant community assembly on coastal dunes', *Journal of Vegetation Science*, 28(4), pp. 762–
773. Available at: https://doi.org/10.1111/jvs.12533.

Coverdale, T.C. and Davies, A.B. (2023) 'Unravelling the relationship between plant diversity
and vegetation structural complexity: A review and theoretical framework', *Journal of
Ecology*, 111(7), pp. 1378–1395. Available at: https://doi.org/10.1111/1365-2745.14068.

Cutler, N.A., Belyea, L.R. and Dugmore, A.J. (2008) 'The spatiotemporal dynamics of a
primary succession', *Journal of Ecology*, 96(2), pp. 231–246. Available at:
https://doi.org/10.1111/j.1365-2745.2007.01344.x.

Davies, A.B. and Asner, G.P. (2014) 'Advances in animal ecology from 3D-LiDAR ecosystem
mapping', *Trends in Ecology & Evolution*, 29(12), pp. 681–691. Available at:
https://doi.org/10.1016/j.tree.2014.10.005.

De Bello, F. *et al.* (2021) *Handbook of trait-based ecology: from theory to R tools*. Cambridge
University Press.

Diekmann, M. (2003) 'Species indicator values as an important tool in applied plant ecology
– a review', *Basic and Applied Ecology*, 4(6), pp. 493–506. Available at:
https://doi.org/10.1078/1439-1791-00185.

Düll, R. (1992) 'Indicator values of mosses and liverworts', *Scrip. Geobotany*, 18, pp. 175–
214.

Eskelinen, A. *et al.* (2022) 'Light competition drives herbivore and nutrient effects on plant
diversity', *Nature*, 611(7935), pp. 301–305. Available at: https://doi.org/10.1038/s41586-022-
05383-9.

Fei, S. *et al.* (2023) 'Structural diversity: a digital revolution', *Frontiers in Ecology and the
Environment*, 21(1), pp. 3–3. Available at: https://doi.org/10.1002/fee.2600.

Fergus, A.J.F., Gerighausen, U. and Roscher, C. (2017) 'Vascular plant diversity structures
bryophyte colonization in experimental grassland', *Journal of Vegetation Science*, 28(5), pp.
903–914. Available at: https://doi.org/10.1111/jvs.12563.

Ficetola, G.F. *et al.* (2024) 'The development of terrestrial ecosystems emerging after glacier
retreat', *Nature*, pp. 1–7. Available at: https://doi.org/10.1038/s41586-024-07778-2.

Gámez, S. and Harris, N.C. (2022) 'Conceptualizing the 3D niche and vertical space use',
Trends in Ecology & Evolution, 37(11), pp. 953–962. Available at:
https://doi.org/10.1016/j.tree.2022.06.012.

Gavini, S.S. *et al.* (2019) 'Facilitation of vascular plants by cushion mosses in high-Andean
communities', *Alpine Botany*, 129(2), pp. 137–148. Available at:
https://doi.org/10.1007/s00035-019-00222-6.

Gough, C.M. *et al.* (2019) 'High rates of primary production in structurally complex forests',
Ecology, 100(10), p. e02864. Available at: https://doi.org/10.1002/ecy.2864.

Grace, J.B. (1999) 'The factors controlling species density in herbaceous plant communities:
An assessment', *Perspectives in Plant Ecology, Evolution and Systematics*, 2(1), pp. 1–28.
Available at: https://doi.org/10.1078/1433-8319-00063.

Hanusch, M. *et al.* (2022) 'Succession comprises a sequence of threshold-induced
community assembly processes towards multidiversity', *Communications Biology*, 5(1), pp.
1–9. Available at: https://doi.org/10.1038/s42003-022-03372-2.

Hanusch, M. *et al.* (2023) 'Exploring the Frequency and Distribution of Ecological Non-
monotonicity in Associations among Ecosystem Constituents', *Ecosystems*, 26(8), pp. 1819–
1840. Available at: https://doi.org/10.1007/s10021-023-00867-9.

Hanusch, M. *et al.* (2024) 'Testing the sequence of successional processes in miniature
ecosystems', *Microbiology Spectrum*. Edited by F. Reverchon, 12(10), pp. e01227-24.
Available at: https://doi.org/10.1128/spectrum.01227-24.

He, X., Hanusch, M., Ruiz-Hernández, V., *et al.* (2023) 'Accuracy of mutual predictions of
plant and microbial communities vary along a successional gradient in an alpine glacier
forefield', *Frontiers in Plant Science*, 13(January), pp. 1–12. Available at:
https://doi.org/10.3389/fpls.2022.1017847.

He, X., Hanusch, M., Saueressig, L., *et al.* (2023) 'Tracking succession by means of 3D
scans of plant communities in a glacier forefield to infer assembly processes', *Oikos*,
2023(12), pp. 1–14. Available at: https://doi.org/10.1111/oik.10095.

He, X. *et al.* (2024) 'Adding experimental precision to the realism of field observations: Plant
communities structure bacterial communities in a glacier forefield', *Environmental*
Microbiology, 26(2), p. e16590. Available at: https://doi.org/10.1111/1462-2920.16590.

Hylander, K. *et al.* (2005) 'Substrate Form Determines the Fate of Bryophytes in Riparian
Buffer Strips', *Ecological Applications*, 15(2), pp. 674–688. Available at:
https://doi.org/10.1890/04-0570.

Junker, R.R. *et al.* (2020) 'Ödenwinkel: An Alpine platform for observational and
experimental research on the emergence of multidiversity and ecosystem complexity', *Web*
Ecology, 20(2), pp. 95–106. Available at: https://doi.org/10.5194/we-20-95-2020.

Junker, R.R. *et al.* (2021) 'Divergent assembly processes? A comparison of the plant and soil
microbiome with plant communities in a glacier forefield', *FEMS Microbiology Ecology*,
97(10), pp. 1–10. Available at: https://doi.org/10.1093/femsec/fiab135.

Kjaer, K.H. and Ottosen, C.O. (2015) '3D laser triangulation for plant phenotyping in
challenging environments', *Sensors (Switzerland)*, 15(6), pp. 13533–13547. Available at:
https://doi.org/10.3390/s150613533.

LaRue, E.A., Fahey, R.T., *et al.* (2023) 'A theoretical framework for the ecological role of
three-dimensional structural diversity', *Frontiers in Ecology and the Environment*, 21(1), pp.
4–13. Available at: https://doi.org/10.1002/fee.2587.

LaRue, E.A., Knott, J.A., *et al.* (2023) 'Structural diversity as a reliable and novel predictor for
ecosystem productivity', *Frontiers in Ecology and the Environment*, 21(1), pp. 33–39.
Available at: https://doi.org/10.1002/fee.2586.

Legendre, P. *et al.* (2005) 'Analyzing beta diversity: Partitioning the spatial variation of
community composition data', *Ecological Monographs*, 75(4), pp. 435–450. Available at:
https://doi.org/10.1890/05-0549.

Loke, L.H.L. and Chisholm, R.A. (2022) 'Measuring habitat complexity and spatial
heterogeneity in ecology', *Ecology Letters*, 25(10), pp. 2269–2288. Available at:
https://doi.org/10.1111/ele.14084.

Lowe, W.H. and McPeek, M.A. (2014) 'Is dispersal neutral?', *Trends in Ecology and*
Evolution, 29(8), pp. 444–450. Available at: https://doi.org/10.1016/j.tree.2014.05.009.

MacArthur, R.H., MacArthur, J.W. and Preer, J. (1962) 'On Bird Species Diversity. II.
Prediction of Bird Census from Habitat Measurements', *The American Naturalist*, 96(888),
pp. 167–174.

Maire, E. *et al.* (2015) 'How many dimensions are needed to accurately assess functional
diversity? A pragmatic approach for assessing the quality of functional spaces', *Global*
Ecology and Biogeography, 24(6), pp. 728–740. Available at:
https://doi.org/10.1111/geb.12299.

Maphosa, L. *et al.* (2017) 'Phenotyping wheat under salt stress conditions using a 3D laser
scanner', *Israel Journal of Plant Sciences*, 64(3–4), pp. 55–62. Available at:
https://doi.org/10.1080/07929978.2016.1243405.

Newman, E.I. (1973) 'Competition and Diversity in Herbaceous Vegetation', *Nature*,
244(5414), pp. 310–310. Available at: https://doi.org/10.1038/244310a0.

Odum, E.P. (1969) 'The strategy of ecosystem development', *Science*, 164(3877), pp. 262–
270. Available at: https://doi.org/10.1126/science.164.3877.262.

Prach, K. and Walker, L.R. (2020) *Comparative plant succession among terrestrial biomes of
the World*, *Comparative Plant Succession among Terrestrial Biomes of the World*.
Cambridge University Press. Available at: https://doi.org/10.1017/9781108561167.

Preston, C.D. and Bosanquet, S.D.S. (2007) *BRYOATT: attributes of British and Irish
mosses, liverworts and hornworts*. Centre for Ecology and Hydrology.

Price, J. *et al.* (2017) 'Within-community environmental variability drives trait variability in
species-rich grasslands', *Journal of Vegetation Science*, 28(2), pp. 303–312. Available at:
https://doi.org/10.1111/jvs.12487.

Rosseel, Y. (2012) '{lavaan}: An {R} Package for Structural Equation Modeling', *Journal of
Statistical Software*, 48(2), pp. 1–36.

Simmel, J., Ahrens, M. and Poschlod, P. (2021) 'Ellenberg N values of bryophytes in Central
Europe*', *Journal of Vegetation Science*, 32(1). Available at:
https://doi.org/10.1111/jvs.12957.

Spitale, D. (2017) 'Forest and substrate type drive bryophyte distribution in the Alps', *Journal
of Bryology*, 39(2), pp. 128–140. Available at:
https://doi.org/10.1080/03736687.2016.1274090.

Srivastava, D.S. (2006) 'Habitat structure, trophic structure and ecosystem function:
interactive effects in a bromeliad–insect community', *Oecologia*, 149(3), pp. 493–504.
Available at: https://doi.org/10.1007/s00442-006-0467-3.

Tews, J. *et al.* (2004) 'Animal species diversity driven by habitat heterogeneity/diversity: the
importance of keystone structures', *Journal of Biogeography*, 31(1), pp. 79–92. Available at:
https://doi.org/10.1046/j.0305-0270.2003.00994.x.

van Tooren, B.F. *et al.* (1990) 'Regeneration of Species Richness in the Bryophyte Layer of
Dutch Chalk Grasslands', *Lindbergia*, 16(5/6), pp. 153–160.

Traylor, C.R. *et al.* (2022) 'Compositional attributes of invaded forests drive the diversity of
insect functional groups', *Global Ecology and Conservation*, 35, p. e02092. Available at:
https://doi.org/10.1016/j.gecco.2022.e02092.

Tsukahara, J. (2023) *semoutput: SEM Output*.

Ulrich, W. *et al.* (2016) 'Species interactions and random dispersal rather than habitat filtering
drive community assembly during early plant succession', *Oikos*, 125(5), pp. 698–707.
Available at: https://doi.org/10.1111/oik.02658.

Weiner, J. (1990) 'Asymmetric competition in plant populations', *Trends in Ecology &
Evolution*, 5(11), pp. 360–364. Available at: https://doi.org/10.1016/0169-5347(90)90095-U.

Werger, M.J.A. *et al.* (2002) 'Light Partitioning among Species and Species Replacement in
Early Successional Grasslands', *Journal of Vegetation Science*, 13(5), pp. 615–626.

Zehm, A., Nobis, M. and Schwabe, A. (2003) 'Multiparameter analysis of vertical vegetation
structure based on digital image processing', *Flora - Morphology, Distribution, Functional
Ecology of Plants*, 198(2), pp. 142–160. Available at: https://doi.org/10.1078/0367-2530-
00086.

Zieschank, V. and Junker, R.R. (2023) 'Digital whole-community phenotyping: tracking
morphological and physiological responses of plant communities to environmental changes

in the field', *Frontiers in Plant Science*, 14(May), pp. 1–13. Available at:
https://doi.org/10.3389/fpls.2023.1141554.

611