Pronouns reactivate conceptual representations in human hippocampal neurons

D. E. Dijksterhuis[1]*†, M. W. Self[1]*, J. K. Possel[1], J. C. Peters[1,2], E.C.W. van Straaten[3], S. Idema[3], J. C. Baaijen[3], S. M. A. van der Salm[4], E.J. Aarnoutse[4], N. C. E. van Klink[4], P. van Eijsden[4], S. Hanslmayr[5], R. Chelvarajah[6,7], F. Roux[7], L. D. Kolibius[5,8], V. Sawlani[6,7], D. T. Rollings[6], S. Dehaene[9,10], P. R. Roelfsema[1,11-13]†

Affiliations:

[1]Vision and Cognition, Netherlands Institute for Neuroscience, Amsterdam, Netherlands

[2]Department of Cognitive Neuroscience, Faculty of Psychology and Neuroscience, Maastricht University, Maastricht, Netherlands

[3]Department of Neurosurgery, Amsterdam University Medical Centre location VUmc, Amsterdam, Netherlands

[4]Department of Neurology and Neurosurgery, University Medical Centre Utrecht, Utrecht, Netherlands

[5]Centre for Cognitive Neuroimaging, School of Psychology and Neuroscience, University of Glasgow, Glasgow, United Kingdom

[6]Complex epilepsy and surgery service, Queen Elizabeth Hospital, Birmingham, United Kingdom

[7]School of Psychology, University of Birmingham, Birmingham, United Kingdom

[8]Columbia University, Department of Biomedical Engineering, USA

[9]Université Paris Saclay, INSERM, CEA, Cognitive Neuroimaging Unit, NeuroSpin center, Saclay, France

[10]Collège de France, Paris, France

[11]Department of Integrative Neurophysiology, VU University, Amsterdam, Netherlands.

[12]Department of Neurosurgery, Amsterdam University Medical Centre, Amsterdam, Netherlands.

[13]Laboratory of Visual Brain Therapy, Sorbonne Université, Institut National de la Santé et de la Recherche Médicale, Centre National de la Recherche Scientifique, Institut de la Vision, Paris, France

* These authors contributed equally to this work
† Corresponding authors: dorisdijk@gmail.com; p.roelfsema@nin.knaw.nl

One-Sentence Summary: Pronouns activate neurons in the human hippocampus if they refer to the concepts to which the cells are tuned.

Abstract

During discourse comprehension, every new word adds to an evolving representation of

meaning that accumulates over consecutive sentences and constrains the next words. To

minimize repetition and utterance length, languages use pronouns, like the word 'she', to

refer to nouns and phrases that were previously introduced. It has been suggested that

language comprehension requires that pronouns activate the same neuronal represen-

tations as the nouns themselves. Here, we test this hypothesis by recording from

individual neurons in the human hippocampus during a reading task. We found that cells

that are selective to a particular noun are later reactivated by pronouns that refer to the

cells' preferred noun. These results imply that concept cells contribute to a rapid and

dynamic semantic memory network which is recruited during language comprehension.

This study uniquely demonstrates, at the single-cell level, how memory and language are

linked.

Consider these two sentences: "John and Mary walked into a bar. He sat down at a table.".

When we read the pronoun 'he', we realize that John, the only male character in the story, must

be the person who sat down at a table. In linguistic terms, John is the 'antecedent' of the

pronoun. This example illustrates how a narrative activates successive concepts in our brain,

including their interrelations, allowing us to incrementally build up a conceptual representation

of the discourse[1–3]. Previous brain-imaging studies have gained insight into the brain regions

that activate during sentence and discourse comprehension[4–11]. However, the resolution of

these non-invasive imaging methods does not suffice to track the neuronal assemblies that

encode individual concepts in the human brain during reading.

In recent years, it has become possible to directly record the activity of single neurons in

patients who are implanted with electrodes to locate the source of their epilepsy[12,13]. These

studies demonstrated the existence of 'concept cells' in the medial temporal lobe[14–16]. Concept

cells have an invariant and multimodal selective response to a concept. They contribute to the

representation of meaning because they not only activate when the participant sees a picture of

a specific individual for example, but also when the participant hears or reads the name of this

person, or recalls this individual from memory[14,17–19]. We hypothesized that monitoring the

activity of concept cells during reading could provide insight into the dynamics of semantic

representations during language comprehension. In the present study, we illustrate the

possibilities of this approach by examining how pronouns that are encountered during reading

influence the neuronal representation of concepts that were introduced in an earlier sentence.

Specifically, we asked if pronouns influence the activity of hippocampal neurons. Our results

reveal that hippocampal cells respond robustly to preferred nouns and that they are specifically

reactivated by a pronoun that refers to them as antecedent. The results represent the first

measurements of the magnitude, latency and duration of hippocampal single cell responses to

nouns and pronouns during reading, while participants incrementally build up a semantic

representation of a narrative.

We recorded from patients with pharmacologically intractable epilepsy who were

implanted with depth electrodes in the hippocampus, to localize seizure foci for clinical

purposes. During an initial screening session, the patients viewed many pictures of celebrities

and family members, and we identified pictures that, for a given neuron, elicited a significantly

higher response than other pictures (Methods). We used the results of the screening session,

aiming to select three nouns for the main reading task: one was the preferred person for a cell

(the preferred noun) and two nouns referred to a male and a female that did not activate that

cell (although the non-preferred nouns sometimes activated other, simultaneously recorded

cells).

In the main task, on every trial, patients read two sentences which were presented as a

stream of words on a computer screen (Fig. 1A and Table 1). The subjects of the first sentence

were always two of the three selected nouns (e.g. Courtney Love and Barack Obama), and the

second sentence started with a pronoun (either she or he). We then asked a question to verify

that the patients understood the meaning of the sentences and maintained focus throughout the

experiment. The mean accuracy was $89 \pm 2\%$ (s.e.m), significantly higher that the chance level

of 33% (Wilcoxon signed rank test, $p = 1.2 \cdot 10^{-4}$; Methods and Fig. 1B).

We identified a total of 529 single and multi-units in the hippocampus of 22 participants

(see Table S3 and Methods). We selected cells for further analyses based on their response

during the reading task, in which 307 cells responded to the nouns. We focused our analysis

on 53 noun-selective cells (recorded in 14 of the participants), which responded significantly

stronger to a preferred noun than to the other two nouns ($p < 0.05$, permutation-based Poisson

ANOVA and post-hoc Poisson t-test, see Methods) (Fig. 1C). Of the noun-selective cells, we

defined 19 to be concept cells, because they preferred the same concept as a picture in the

screening task and as a word during the reading task (Methods, Fig. S1 and Table S2; we note

that a previous study included more extensive tests of multimodality[20]). The remaining 34 cells

were noun-selective, but either did not respond to pictures or preferred a different concept

during the screening task, and we will refer to them as noun-selective, non-concept cells

(NSNCs). Figure 1D illustrates the activity of an example concept cell that responded

selectively to a picture of the animated male movie character Shrek during the screening task.

The cell also preferred the written noun 'Shrek' ($Noun_{Pref}$) (Figure 1E), but it did not respond

to Courtney Love and a male family member of the participant ($Nouns_{NonPref}$). The pronoun 'he'

in the second sentence activated the cell if it referred to Shrek in the first sentence ($Pronoun_{Ref}$).

The pronoun 'she' referred to the other noun ($Pronoun_{PresNotRef}$; Table 2 specifies the naming

conventions) and did not activate the cell. Furthermore, 'he' also did not activate the cell if

Shrek was absent from the first sentence ($Pronoun_{AbsentSame}$). Hence, only pronouns referring to

Shrek activated the neuron.

To examine the generality of this effect, we carried out an analysis across the population

of 53 noun-selective cells. We first examined the activity elicited by the nouns with a cross-

validation approach. We used half of the trials to determine the preferred noun and the other

half for the statistics. Across the population, the activity elicited by $Noun_{Pref}$ was 5.4 ± 0.5 Hz

(mean $\pm$ s.e.m.), which was significantly higher than the activity of 2.9 ± 0.5 Hz elicited by the

other, non-preferred nouns (t-test, $t = 3.9$, $p = 3.0 \cdot 10^{-4}$). We determined the neurons' response

latencies as the first of 10 subsequent time-bins where the response to $Noun_{Pref}$ was higher than

the response to the other nouns ($P < 0.05$; see Methods and Fig. 2D-F). Regardless of whether

$Noun_{Pref}$ was presented at the first or second position in the first sentence, the cells activated

on average $210ms \pm 38ms$ (s.e.m.) after the presentation of the preferred noun.

We next examined the responses elicited by the pronouns. On average, pronouns referring

to $Noun_{Pref}$ ($Pronoun_{Ref}$) activated the hippocampal neurons more strongly than pronouns that

did not (paired t-test, $t_{52} = 6.8$, $p = 3.3 \cdot 10^{-7}$) (Fig. 2A). This effect was significant for 25% of

the noun-selective cells and for 42% of the concept cells (permutation-based Poisson ANOVA;

$P < 0.05$). Interestingly, the responses of concept cells to $Pronoun_{Ref}$ (5.0 ± 0.9 Hz, $N = 19$)

were stronger than those of the NSNC cells (2.3 ± 0.5 Hz, $N = 34$; t-test corrected for unequal

variance, $t = 2.6$, $p = 0.016$; Fig. 2A). The extra activity elicited by $Pronoun_{Ref}$ emerged

gradually (Fig. 2G) with a latency of $600ms \pm 110ms$, which is longer than the latency of the

response to $Noun_{Pref}$ ($p = 0.010$, bootstrapping test), although activity built up gradually so that

the inclusion of more cells might have shortened the latency estimate. There were no significant

differences in response magnitude (independent samples t-test, $t_{51} = 0.19$, $p = 0.85$) or latency

($p = 0.54$; bootstrap test, see Methods) between the left and right hemispheres.

Is the response to $Pronoun_{Ref}$ caused by a prolonged activation by the preferred noun,

extending into the second sentence? To examine this question, we selected the first sentences

with $Noun_{Pref}$ and compared the activity elicited by pronouns that referred to $Noun_{Pref}$

(Pronoun$_{Ref}$) and to the other noun (Pronoun$_{PresNotRef}$); for example 'Shrek and Courtney Love'

followed by 'He' or 'She'. The response elicited by Pronoun$_{Ref}$ was higher than that evoked by

Pronoun$_{PresNotRef}$ (paired t-test, $p = 1.4 \cdot 10^{-7}$) (Fig. 2B), indicating that it reflects the antecedent

of the pronoun and is not a lingering memory trace of Noun$_{Pref}$.

We next investigated whether the pronoun activation reflected tuning to gender rather than

to the identity of the antecedent. We compared trials in which the pronoun referred to Noun$_{Pref}$

(e.g. 'Shrek and Courtney Love' followed by 'He') to trials in which the same pronoun referred

to a different noun of the same gender (Pronoun$_{AbsentSame}$, e.g. 'Barack Obama and Courtney

Love' followed by 'He'). Pronoun$_{Ref}$ elicited a higher response than Pronoun$_{AbsentSame}$ ($N = 45$,

paired t-test, $p = 1.5 \cdot 10^{-3}$; Fig. 2C), implying that the pronoun responses reflected the identity

of the antecedent and not the gender.

We examined whether population responses evoked by pronouns resembled those evoked

by the nouns and how different cell groups contributed to the population response. To this end,

we trained a linear support-vector machine (SVM) to discriminate between responses elicited

by Noun$_{Pref}$ and Noun$_{NonPref}$. We constructed 1,000 surrogate-populations by randomly

choosing units from the noun-selective cells with replacement ($N = 50$ cells; we excluded three

cells from this analysis, Table S3). We trained the classifier on activity in the noun time-

window and tested the classifiers on held-back trials in a series of sliding windows covering

the two sentences. We split trials based on first sentences in which Noun$_{Pref}$ 1) appeared at the

first position, 2) appeared at the second position and 3) was absent. The classifier output

represents the fraction of trials in a particular time-window in which the neuronal responses

are labelled as elicited by Noun$_{Pref}$ (y-axis in Fig. 3A). As expected, the classifier output

increased shortly after the presentation of Noun$_{Pref}$ at both the first and second position,

labelling the activity as evoked by Noun$_{Pref}$ in ~90% of trials (Fig. 3A). Interestingly, when

Noun$_{Pref}$ was at the first position, the classifier output increased slightly after the presentation

of the second, non-preferred noun (black arrow in Fig. 3A), which may reflect a mental process

that relates the two nouns.

We next tested the classifier on the second sentence, on which it had not been trained. We

compared the classifier output between trials in which the pronoun referred to the preferred

noun ($Pronoun_{Ref}$), trials on which it referred to the other noun ($Pronoun_{PresNotRef}$) and trials on

which $Noun_{Pref}$ was absent from the first sentence. The activity pattern elicited by $Pronoun_{Ref}$

was like that elicited by $Noun_{Pref}$, causing a higher decoder output than on trials in which the

pronoun referred to the other, non-preferred noun and on trials without $Noun_{Pref}$ in the first

sentence (Fig. 3A). The decoder output was also higher than on trials in which the antecedent

was the non-preferred noun with the same gender ($Pronoun_{AbsentSame}$; Fig. S2A). We replicated

the results with a cross-decoding analysis, training and testing on all combinations of time-bins

in the two sentences. Cross-decoding worked in both directions, from noun to pronoun and

from pronoun to noun (Fig. S2B).

To investigate which cells contributed most to the selectivity of the pronoun response, we

built decoders by successively adding concept cells ($N = 19$), NSNC cells ($N = 19$ selected of

a total of 31 units) and neurons that responded to nouns without significant tuning in different

orders ('untuned', $N = 215$, noun response > 0.5 Hz above baseline; see Table S3 for inclusion

of cells) (Fig. 3B). Decoders built from noun-selective cells (concept and NSNC) to decode

nouns also reliably decoded the pronoun of the second sentence, whereas adding untuned cells

did not improve this form of generalization (Fig. 3C). Interestingly, adding NSNC cells to

decoders that already included all concept-cells did not improve decoding, whereas adding

concept-cells to decoders with only NSNC cells improved decoding accuracy ($p = 0.015$, z-

test) (Fig. 3B,C). These results indicate that decoding of the pronoun's antecedent relied mostly

on the concept cells, with a smaller contribution from NSNC cells and little contribution from

the untuned cells.

Two additional observations demonstrated that the neural responses were linked to the

subjects' interpretation of the pronoun on individual trials. First, we compared the response to

pronouns that referred to the preferred noun ($Pronoun_{Ref}$) between correct trials and error trials,

on which the participants reported the incorrect noun in response to the question at the end of

the trial. The hippocampal response was higher on correct trials than on error trials (t-test, two-

tailed, $t = -2.31$, $p = 0.008$, $N = 35$; Fig S3A), indicating that it predicted the participant's

interpretation of the second sentence.

Second, we analysed ambiguous trials with two nouns of the same gender in the first

sentence, so that they competed for a status as antecedent of the pronoun ('Shrek and Barack

Obama went into a bar. He sat at the table', followed by the question 'Who sat at the table?')

(Fig. 3D; Table 2). We predicted that the strength of the response to $Noun_{Pref}$ in the first

sentence might relate to the probability that the participant chooses it as antecedent and reports

it in response to the question. In accordance with this prediction, the decoder output for the

preferred noun of the first sentence was stronger if the participant chose it in response to the

question (Fig. 3D) ($p = 0.045$, z-test). Furthermore, decoder output also ramped up at the end

of the second sentence if the participants chose the preferred noun (Fig. 3D). Hence, the activity

of hippocampal neurons during reading is related to the strength of the conceptual

representations and predicts the resolution of pronoun ambiguity.

The results, taken together, demonstrate a link between the activity of neurons in the human

hippocampus and the incremental representation of concepts and their interrelations during

sentence reading. Previous studies indicated how language understanding relies on the

integration of semantic knowledge, to which the hippocampus contributes[21–26], with linguistic

representations[27–31]. We here provided the first measurements of single unit activity in the

hippocampus during a task in which participants build up a semantic representation of a

narrative[32], in which a pronoun brings one of the characters into the foreground of thought.

Interestingly, most of the information about the antecedent of the pronoun was carried by

concept cells, which responded to a concept irrespective of whether it was presented as picture

or noun, indicating that their activity disambiguates otherwise identical pronouns.

Hippocampal neurons responded to both their preferred noun and to pronouns referring to

that noun. The hippocampal pronoun response could thereby link new information of the

narrative to the appropriate concept. For example, when we read about Shrek that 'he' put on

sunglasses, we can update Shrek's representation and predict his future appearance[14,33–35]. In

accordance with this view, damage to the hippocampus can lead to impairments in pronoun

production and comprehension and patients with hippocampal lesions often fail to retrieve a

pronoun's antecedent[36,37]. Our results support a role for the hippocampus in pronoun resolution,

because the pronoun response was weaker on error trials and hippocampal activity predicted

the perceived antecedent of the pronoun on ambiguous trials.

The latency of the pronoun response was 600ms, which was longer than the latency of

210ms of the noun response, and most likely reflects processes that link the representation of

the pronoun in language areas to the appropriate concept in the hippocampus. The long latency

is remarkable because pronouns are thought to increase discourse efficiency, allowing shorter

utterances to activate the same concepts as longer noun phrases[1,38]. The present results indicate

that this pronoun advantage may only occur for noun phrases that take several hundreds of

milliseconds. We note, however, that the latency of the pronoun response in the hippocampus

might be shorter if there is only a single possible antecedent.

Language theories propose that pronouns have additional advantages, such as signalling

that the subject of the narrative remains the same across sentences[39]. Our results inform these

theories[1,39], including 'prominence' theories that propose that pronouns usually refer to the

most prominent nouns of a discourse[40]. According to those theories, nouns that fulfil a similar

role in previous sentences compete for selection based on their prominence, which reflects on

how important they are in the narrative[40]. So far, these theories could only be tested

indirectly[41,42], short of a method to measure prominence in the brain. Our results on trials with

two nouns of the same gender, which caused the pronoun to be ambiguous, provided insight

into the neuronal underpinnings of prominence. On these ambiguous trials, the participant

could report either noun as antecedent of the pronoun. When the participants reported that the

pronoun referred to the preferred noun, the response to that noun was stronger during the first

sentence, and hippocampal activity ramped up once more at the end of the second sentence.

These results suggest that prominence is related to the strength of the conceptual

representations, which depend, in part, on the activity of hippocampal neurons.

Theories about the evolving mental representation of the narrative during reading suggest

that previously read words are stored in working memory so that they can be combined with

new information[28–30,32,43,44]. Interestingly, the hippocampal activity elicited by nouns lasted

only 300-400ms and was curtailed by later words, which has implications for the internal

structure of these representations[28]. Theories of working memory propose that memorized

items are not equivalent, because only one or a few of them can be in the focus of attention[45].

Studies in the frontal cortex of monkeys revealed that attended and non-attended items in

working memory are represented as sustained activity patterns[46,47], whereas the present results

suggest that the hippocampus selectively represents the attended noun. The activity of the

neurons largely vanished during the reading of subsequent words, but the cells became active

again in response to a pronoun referring to the cell's preferred concept. Hypothetically, the

representation could have moved into a different neuronal subspace, as observed during

sequence working memory in monkey prefrontal cortex[46], but this is unlikely given that there

were epochs in which we could no longer decode it from the population of neurons. The activity

profile rather suggests that hippocampal neurons predominantly represent concepts which

come in the focus of attention as a central topic of the narrative. These attended concepts could

then be used for the retrieval of additional associations, a process to which the hippocampus

contributes[48],

In our experiment, pronoun disambiguation was based on gender alone, but in other

sentence constructions, syntax plays an essential role. For instance, in the sentence "The fact

that he is so rich pleases John", the pronoun "he" can refer to John, but in "He is so rich that

John is pleased", it cannot. How brain networks implement such syntactic computations is a

topic for future research, which can now be investigated. The present results show how single

unit recordings in the medial temporal lobe contribute to the flexible linkage of words and

concepts during language comprehension.

References and Notes

1.   Gordon, P. C., Grosz, B. J. & Gilliom, L. A. Pronouns, names, and the centering of

attention in discourse. *Cogn. Sci.* **17**, 311–347 (1993).

2.   Nieuwland, M. S., Petersson, K. M. & Van Berkum, J. J. A. On sense and reference:

Examining the functional neuroanatomy of referential processing. *Neuroimage* **37**,

993–1004 (2007).

3.   Vosse, T. & Kempen, G. Syntactic structure assembly in human parsing: A

computational model based on competitive inhibition and a lexicalist grammar.

Cognition **75**, 105–143 (2000).

4.   Hagoort, P., Hald, L., Bastiaansen, M. & Petersson, K. M. Integration of Word

Meaning and World Knowledge in Language Comprehension. *Science (80-.).* **304**,

438–441 (2004).

5.   Buchweitz, A., Mason, R. A., Tomitch, L. M. B. & Just, M. A. Brain activation for

reading and listening comprehension: An fMRI study of modality effects and

individual differences in language comprehension. *Psychol. Neurosci.* **2**, 111–123

(2009).

6.   Pallier, C., Devauchelle, A. D. & Dehaene, S. Cortical representation of the constituent

structure of sentences. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 2522–2527 (2011).

7.   Huth, A. G., de Heer, Wendy, A., Griffiths, T. L., Theunissen, F. E. & Gallant, J. L.

Natural speech reveals the semantic maps that tile human cerebral cortex. *Nature* **532**,

453–458 (2016).

8.   Baldassano, C., Chen, J., Zadbood, A., Pillow, J. W. & Norman, K. A. Discovering

event structure in continuous narrative perception and memory. *Neuron* **95**, 709–721

(2017).

9.   Fedorenko, E., Behr, M. K. & Kanwisher, N. Functional specificity for high-level

linguistic processing in the human brain. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 16428–

16433 (2011).

10.   Friederici, A. D. The brain basis of language processing: From structure to function.

Physiol. Rev. **91**, 1357–1392 (2011).

11.   Binder, J. R., Desai, R. H., Graves, W. W. & Conant, L. L. Where is the semantic

system? A critical review and meta-analysis of 120 functional neuroimaging studies.

Cereb. Cortex **19**, 2767–2796 (2009).

12.   Engel, A. K., Moll, C. K. E., Fried, I. & Ojemann, G. A. Invasive recordings from the

human brain: Clinical insights and beyond. *Nat. Rev. Neurosci.* **6**, 35–47 (2005).

13.   Babb, T. L., Carr, E. & Crandall, P. H. Analysis of Extracellular Firing Patterns of

Deep Temporal Lobe Structures in Man. *Electroencephalogr. Clin. Neurophysiol.* **34**,

247–257 (1973).

14.   Quiroga, R. Q. Concept cells: the building blocks of declarative memory functions.

Nat. Rev. Neurosci. **13**, 587–597 (2012).

15.   Quiroga, R. Q., Reddy, L., Kreiman, G., Koch, C. & Fried, I. Invariant visual

representation by single neurons in the human brain. *Nature* **435**, 1102–1107 (2005).

16. Quiroga, R. Q., Kreiman, G., Koch, C. & Fried, I. Sparse but not 'Grandmother-cell' coding in the medial temporal lobe. *Trends Cogn. Sci.* **12**, 87–91 (2008).

17. Quiroga, R. Q., Reddy, L., Kreiman, G., Koch, C. & Fried, I. Invariant visual representation by single neurons in the human brain. *Nature* **435**, 1102–1107 (2005).

18. Gelbard-Sagiv, H., Mudrik, L., Hill, M. R., Koch, C. & Fried, I. Human single neuron activity precedes emergence of conscious perception. *Nat. Commun.* **9**, (2018).

19. Viskontas, I. V, Quian, R. & Fried, I. Human medial temporal lobe neurons respond preferentially to personally relevant images. **106**, 21329–21334 (2009).

20. Quian Quiroga, R., Kraskov, A., Koch, C. & Fried, I. Explicit Encoding of Multimodal Percepts by Single Neurons in the Human Brain. *Curr. Biol.* **19**, 1308–1313 (2009).

21. Fletcher, P. C., Frith, C. D. & Rugg, M. D. The functional neuroanatomy of episodic memory. *Trends Neurosci.* **20**, 213–218 (1997).

22. Squire, L. R. & Zola-Morgan, S. The medial temporal lobe memory system. *Science (80-.).* **253**, 1380–1386 (1991).

23. Tulving, E. Episodic : From Mind to Brain. *Annu. Rev. Psychol.* **53**, 1–25 (2002).

24. Wirth, S. *et al.* Single Neurons in the Monkey Hippocampus and Learning of New Associations. *Science (80-.).* **300**, 1578–1581 (2016).

25. Davachi, L. Item, context and relational episodic encoding in humans. *Curr. Opin. Neurobiol.* **16**, 693–700 (2006).

26. Blank, I., Duff, M., Brown-Schmidt, S. & Fedorenko, E. Expanding the language network: Domain-specific hippocampal recruitment during high-level linguistic processing. *bioRxiv* 091900 (2016) doi:10.1101/091900.

27. Almor, A., Smith, D. V., Bonilha, L., Fridriksson, J. & Rorden, C. What is in a name? Spatial brain circuits are used to track discourse references. *Neuroreport* **18**, 1215–

1219 (2007).

28.   Desbordes, T. *et al.* Dimensionality and Ramping: Signatures of Sentence Integration

in the Dynamics of Brains and Deep Language Models. *J. Neurosci.* **43**, 5350–5364

(2023).

29.   Desbordes, T., King, J. R. & Dehaene, S. Tracking the neural codes for words and

phrases during semantic composition, working-memory storage, and retrieval. *Cell*

Rep. **43**, (2024).

30.   Fedorenko, E. *et al.* Neural correlate of the construction of sentence meaning. *Proc.*

Natl. Acad. Sci. U. S. A. **113**, E6256–E6262 (2016).

31.   Nelson, M. J. *et al.* Neurophysiological dynamics of phrase-structure building during

sentence processing. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E3669–E3678 (2017).

32.   Schwering, S. C. & MacDonald, M. C. Verbal Working Memory as Emergent from

Language Comprehension and Production. *Front. Hum. Neurosci.* **14**, 1–19 (2020).

33.   Quian Quiroga, R. Neuronal codes for visual perception and memory.

Neuropsychologia **83**, 227–241 (2016).

34.   Kornblith, S., Quiroga, R. Q., Koch, C., Fried, I. & Mormann, F. Persistent Single-

Neuron Activity during Working Memory in the Human Medial Temporal Lobe.

(2017) doi:10.1016/j.cub.2017.02.013.

35.   Rutishauser, U., Reddy, L., Mormann, F. & Sarnthein, J. The Architecture of Human

Memory: Insights from Human Single-Neuron Recordings. *J. Neurosci.* **00**, JN-SY-

1648-20 (2020).

36.   Kurczek, J. & Duff, M. C. Processing Deficits in Amnesia. *J. Exp. Psychol.* **142**,

1346–1354 (2014).

37.   Duff, M. C., Gupta, R., Hengst, J. A., Tranel, D. & Cohen, N. J. The Use of Definite

References Signals Declarative Memory: Evidence From Patients With Hippocampal

Amnesia. *Psychol. Sci.* **22**, 666–673 (2011).

38. Nieuwland, M. S. & Martin, A. E. Neural Oscillations and a Nascent

Corticohippocampal Theory of Reference. *J. Cogn. Neurosci.* 1–15 (2013)

doi:10.1162/jocn.

39. Grosz, B. J., Joshi, A. K. & Weinstein, S. Centering : A Framework for Modelling the

Local Coherence of Discourse. *Comput. Linguist.* **21**, 203–225 (1995).

40. von Heusinger, K. & Schumacher, P. B. Discourse prominence: Definition and

application. *J. Pragmat.* **154**, 117–127 (2019).

41. Schumacher, P. B., Backhaus, J. & Dangl, M. Backward- and forward-looking

potential of anaphors. *Front. Psychol.* **6**, 1–14 (2015).

42. Repp, M. & Schumacher, P. B. What naturalistic stimuli tell us about pronoun

resolution in real-time processing. *Front. Artif. Intell.* **6**, (2023).

43. Shain, C., Blank, I. A., Fedorenko, E., Gibson, E. & Schuler, W. Robust Effects of

Working Memory Demand during Naturalistic Language Comprehension in

Language-Selective Cortex. *J. Neurosci.* **42**, 7412–7430 (2022).

44. Nelson, M. J. *et al.* Neurophysiological dynamics of phrase-structure building during

sentence processing. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E3669–E3678 (2017).

45. Cowan, N. The magical number 4 in short-term memory: A reconsideration of mental

storage capacity. *Behav. Brain Sci.* **24**, 87–114 (2001).

46. Xie, Y. *et al.* Geometry of sequence working memory in macaque prefrontal cortex.

Science (80-.). **375**, 632–639 (2022).

47. Panichello, M. F. & Buschman, T. J. Shared mechanisms underlie the control of

working memory and attention. *Nature* **592**, 601–605 (2021).

48. Ison, M. J., Quian Quiroga, R. & Fried, I. Rapid Encoding of New Memories by

Individual Neurons in the Human Brain. *Neuron* **87**, 220–230 (2015).

Acknowledgments: We are grateful to the staff of the EMU of the VUmc and UMCU for their

help during these recordings. We thank Peterjan Ris and Ndedi Sijsma from the VUmc, and

Luigi Rizzi, Naama Friedmann, Hilda Koopman and Klaus von Heusinger for discussions at

various stages of this project. We thank Peter Hagoort for his valuable feedback on the

manuscript.

Funding:

This work was supported by KNAW 240-846401 NWA-StartImpuls 2017, NWO Crossover

grant number 17619 'INTENSE'; DBI2 (a Gravitation program of the Dutch Ministry of

Science), the European Union (ERC grant 101052963 'NUMEROUS' and 647954

'Code4Memory', H2020 Research and Innovation programme grant number 899287

'NeuraViper', the Human Brain Project, grant number 650003).

Author contributions:

Conceptualization: MWS, PRR, SD

Clinical procedures: JCB, SI, PE, RC, VS, IES, SMAS, EA, NCEK, DTR

Ethical approval acquisition: MWS, DED, SH

Investigation: DED, MWS, JKP, JCP, SH, FR, LK

Formal analysis: DED, MWS

Visualization: DED, MWS

Funding acquisition: MWS, PRR

Supervision: MWS, PRR

Writing – original draft: DED, MWS, PRR

Writing – review & editing: DED, MWS, PRR, SD

Competing interest declarations: None.

Data availability. All source data and code to create the main and supplementary data figures

will become publicly available via EBRAINS (https://ebrains.eu/service/share-data/).

Supplementary Materials

Methods

Figs. S1 to S5

Tables S1 to S3

Fig. 1. Reading task. (A) Two sentences were presented word by word. The first sentence

contained two nouns representing individuals (Table 1 and Table S1). The second sentence

started with a pronoun. The trial ended with a question (see Methods). **(B)** Accuracy in

answering the question across all patients. Red lines, accuracy $\pm$ s.e.m. (chance level was 33%,

$N = 14$). **(C)** Locations of the electrodes in the hippocampus. **(D)** The average response of an

example neuron to different pictures during the screening session, and its waveshape. **(E)** The

average response of the 'Shrek' cell during the reading task. Sentences in which Shrek appeared

as the second noun were followed by a second sentence with the pronoun 'he' (magenta) or

'she' (green). Activity on trials in which Shrek did not appear in the first sentence is shown in

black. Black line at time zero, start of first sentence. Dashed grey lines, onset of successive

words. Black dashed lines, presentation of $_{Shrek}$ (or the family member) and the pronoun. The

activity elicited by the pronoun was significantly stronger if it referred to Shrek than if it did

not (permutation-based Poisson ANOVA, $P < 0.001$).

Fig. 2. Hippocampal responses elicited by nouns and pronouns. (A) Comparison of activity

elicited by pronouns referring to the preferred noun (Pronoun$_{Ref}$, y-axis) and all pronouns

referring to another noun (Pronoun$_{NonRef}$, x-axis) for concept cells (red dots) and NSNC cells

(blue dots). The inset shows the response difference. Red arrow, median. P-value refers to a

two-tailed paired t-test ($N = 53$, $t = 5.85$, $p = 3.3 \cdot 10^{-7}$). **(B)** Response to Pronoun$_{Ref}$ (y-axis) and

pronouns referring to the other, non-preferred noun on trials in which Noun$_{Pref}$ was present in

the first sentence (Pronoun$_{PresNotRef}$, $t = 6.09$, $p = 1.4 \cdot 10^{-7}$) **(C)** Response to Pronoun$_{Ref}$ and the

same pronoun on trials in which the preferred noun was absent from the first sentence

(Pronoun$_{\text{AbsentSame}}$, $t = 3.39$, $p = 1.5 \cdot 10^{-3}$). **(D)** Average activity on trials in which Noun$_{\text{Pref}}$ was

the first noun of the first sentence (Noun1, magenta and green traces) or was absent (black

trace), followed by Pronoun$_{\text{Ref}}$ (magenta) or Pronoun$_{\text{PresNotRef}}$ (green). Shading denotes s.e.m.

The vertical dashed lines indicate Noun1 and pronoun onset. **(E)** Average activity on trials in

which Noun$_{\text{Pref}}$ was the second noun in the first sentence. **(F)** Average activity elicited by

Noun$_{\text{Pref}}$ and Noun$_{\text{NonPref}}$. The red arrows indicate the latency of the response and the red bars

the 95%-confidence interval (determined with bootstrapping). **(G)** Average responses elicited

by Pronoun$_{\text{Ref}}$ and Pronoun$_{\text{NotRef}}$.

Fig. 3. Decoders for the preferred noun. (A) SVM classifiers trained to distinguish between

the responses to preferred and non-preferred nouns. The output (y-axis) represents the fraction

of trials in which the pattern of activity in a time-window is classified as evoked by $Noun_{Pref}$.

The classifiers were trained on the activity of 50 cells in the noun time-window and then tested

on the two sentences in a series of sliding windows of 0.5s duration. The words are marked at

the centre of the sliding window, the red dot indicates the end of the second sentence. The

shading around the traces indicates the standard deviation across surrogate populations ($N =$

1,000). Filled bars indicate significant time windows (z-test, $P < 0.05$, FDR corrected; see

Methods). **(B)** Decoder output for different cell populations (0.5-1.5 seconds after pronoun

onset) for Pronoun_{Ref} (magenta) and $\text{Pronoun}_{PresNotRef}$ (green). Error bars, standard deviation

across surrogate populations. P-values, z-test on the difference between conditions. **(C)**

Classifier output as function of the number of included units. **Left,** We first added concept cells

(CC: $N = 19$), then NSNC cells ($N = 19$ cells selected from a total of 31 cells) and then untuned

units ($N = 215$). Decoder output is shown for Pronoun_{Ref} (magenta) and $\text{Pronoun}_{PresNotRef}$

(green). **Middle,** We first added 19 NSNC cells, then 19 CC cells and then the untuned units.

Right, We added units in a random order. We compared the decoder output between decoders

with only CCs, only NSNCs ($N = 19$, length of blue vertical lines) to that when cells of the

other category were added ($N = 38$, length of black lines) with a z-test to determine significance

(P-values in the panels). **(D)** Decoder output on ambiguous trials. The decoders, which had

been trained on non-ambiguous trials to discriminate between preferred and non-preferred

nouns, were tested on ambiguous trials with two nouns of the same gender. The left panel

shows the decoder output to the preferred noun in the first sentence (aligned on Noun_{Pref} onset).

The right panel shows the decoder output during the second sentence, in sliding windows with

a duration of 0.5s. We sorted the trials based on whether the participant selected the preferred

noun (Chose_{Pref}, orange) or the non-preferred noun of the same gender ($\text{Chose}_{NotPref}$, purple) in

response to the question. Q1, onset of the question.

Contexts	First sentences:	Second sentences:
English	*Noun₁* and *noun₂* walked into the bar	*Pronoun* sat at the table
	Noun₁ and *noun₂* sat in the park	*Pronoun* put some sunglasses on
	Noun₁ and *noun₂* were watching the TV	*Pronoun* suddenly changed the channel
	Noun₁ and *noun₂* were having dinner	*Pronoun* poured out some wine

Table 1. Four different contexts in which the noun and pronouns were used for English patients.

Condition description	Nouns in the first sentence	Pronoun in the second sentence	Number of repeats*
The first sentence contains the preferred noun and a noun of the opposite gender. The pronoun refers to the preferred noun.	$\text{Noun}_{Pref} + \text{Noun}_{NonPref}$	Pronoun_{Ref}	16
The first sentence contains the preferred noun and a noun of the opposite gender. The pronoun refers to the noun with the opposite gender.	$\text{Noun}_{Pref} + \text{Noun}_{NonPref}$	$\text{Pronoun}_{PresNotRef}$	16
The preferred noun is not present in the first sentence. A noun of the opposite and of the same gender as the preferred noun are present.	$\text{Noun}_{NonPref1} + \text{Noun}_{NonPref2}$	Pronoun_{Absent}	32
- The pronoun refers to the Noun with the same gender		$\text{Pronoun}_{AbsentSame}$	16
- The pronoun refers to the noun with the opposite gender		$\text{Pronoun}_{AbsentOppo}$	16
All trials where the pronoun does not refer to Noun_{Pref}. Noun_{Pref} can be absent or present.	$\text{Noun}_{Pref}/\text{Noun}_{NonPref}$	Pronoun_{NotRef}	48
The first sentence contains the preferred noun and a noun of the same gender. The pronoun is ambiguous.	$\text{Noun}_{Pref} + \text{Noun}_{NonPrefSameGender}$	Chose_{Pref} or $\text{Chose}_{NotPref}$	16

Table 2. Noun and pronoun conditions. The trials with Pronoun_{Absent} include all trials with $\text{Pronoun}_{AbsentSame}$ and $\text{Pronoun}_{AbsentOppo}$. Similarly, the trials with Pronoun_{NotRef} include all trials with Pronoun_{Absent} and $\text{Pronoun}_{PresNotRef}$. *In two sessions of one of the patients (a total of three hippocampal neurons) we recorded only half of the number of trials.