Eco-evolutionary Guided Pathomics Analysis to Predict DCIS Upstaging

Yujie Xiao[1], Manal Elmasry[2,3], Ji Dong K. Bai[2], Andrew Chen[2], Yuzhu Chen[2], Brooke Jackson[4], Joseph O. Johnson[4], Robert J. Gillies[4], Prateek Prasanna[5], Chao Chen[5*], Mehdi Damaghi[1,2,3*]

1- Department of Applied Mathematics and Statistics, Stony Brook University, NY, USA
2- Department of Pathology, Stony Brook Medicine, Stony Brook University, NY, USA
3- Department of Pathology, Faculty of Medicine, Mansoura University, Mansoura, Egypt
4- Moffitt Cancer Center, Tampa, Fl, USA
5-Department of Biomedical Informatics, Stony Brook Medicine, Stony Brook University, NY, USA

Running title: Eco-Evolutionary Designed Biomarker

[*] Corresponding authors: Chao Chen: chao.chen.1@stonybrook.edu,
Mehdi Damaghi: mehdi.damaghi@stonybrookmedicine.edu
Stony Brook Cancer Center
MART building, 9th floor, 9M0802
Lauterbur Drive, Stony Brook, NY 11794-7263
Phone: (631) 216-2920

Declaration of interests
The authors declare no competing interest.

Abstract: Cancers evolve in a dynamic ecosystem. Thus, characterizing cancer's ecological dynamics is crucial to understanding cancer evolution and can lead to discovering novel biomarkers to predict disease progression. Ductal carcinoma in situ (DCIS) is an early-stage breast cancer characterized by abnormal epithelial cell growth confined within the milk ducts. In this study, we show that ecological analysis of hypoxia and acidosis biomarkers can significantly improve prediction of DCIS upstaging. First, we developed a novel eco-evolutionary designed approach to define habitats in the tumor intra-ductal microenvironment based on oxygen diffusion distance. Then, we identified cancer cells with metabolic phenotypes attributed to their habitats, including CA9 for hypoxia responding phenotype, and LAMP2b for acid adapted phenotype. Traditionally these markers have shown limited predictive capabilities for DCIS progression, if any. However, when analyzed from an ecological perspective, their power to differentiate between non-upstaged and upstaged DCIS increased significantly. Second, we discovered distinct niches with spatial patterns of these biomarkers and used the distribution of such niches to predict patient upstaging. The niches were characterized by pattern analysis of both cellular and spatial features. With a 5-fold validation on the biopsy cohort, we trained a random forest classifier to achieve the area under curve (AUC) of 0.74. Our results affirm the importance of tumor ecological features in eco-evolutionary-designed approaches for novel biomarkers discovery.

Significance: Our results affirm the importance of spatial and ecological features in eco-evolutionary-designed biomarkers discovery studies in the era of digital pathology.

Keywords: Tumor ecology and evolution, DCIS, Eco-evolutionary biomarkers, Metabolic phenotypes, Habitat, Niche, Pathomics, Machine learning, Digital pathology

Introduction:

In recent years, the understanding that cancer is a dynamic ecological and evolutionary process has become deeply entrenched (1,2,3). To date, several evolutionary approaches have been adapted and applied in cancer biology; however, tumor ecosystem and ecological studies are still overlooked (3,4). Within the human body and much like organisms in the nature, cancer cells follow ecological principles, utilizing resources and establishing niches within tissues habitats (5,6). This ecological perspective of cancer is crucial for discovering the natural selection driving cancer evolution. Such insights may potentially lead to improved cancer prognosis, progression prediction, risk stratification, and therapeutic strategies. If tumor evolutionary and/or its ecological state could be reliably achieved using a single biopsy tissue, clinical translation would be comparatively more manageable. Nevertheless, studies have yet to determine whether measures of tumor eco-evolution derived from a single biopsy sample are adequate, or if the inclusion of multiple samples significantly enhances predictions of clinical outcomes (7).

Breast cancer incidence in the US has been increasing over the past decade at a rate of 0.5% per year (8). With increased mammographic screening, there has been a substantial increase in detecting the early non-invasive forms of breast cancer, such as ductal carcinoma in situ (DCIS)(2,9). About one-third of breast cancers detected by mammography are DCIS (10). DCIS and IDC (invasive ductal carcinoma) are indistinguishable by (epi-)genetic mutations, gene expression, or protein biomarkers. It is also not possible to predict whether DCIS will remain non-upstaged or progress to aggressive disease. Therefore, almost all early tumors are treated the same (2,12–14). To avoid such overtreatment for non-upstaged group, more research is needed to fully understand evolution from pre-cancer to non-upstaged DCIS or progress to IDC (9).

DCIS can be defined as a heterogeneous group of neoplastic lesions confined to the mammary ducts. The confinement of proliferating neoplastic cells inside the duct and growth of pre-cancer cells toward the center of the duct, which is far from vasculature, causes limitations in oxygen access. This intraductal oxygen availability is also influenced by complex ecosystems surrounding the duct, such as vascular density and activity (15), stiffness of extracellular matrix (ECM) (16), and metabolites (6,17,18,19) (**Figure 1A**). Local microinvasion is the main difference between DCIS and IDC and might also be the first evolutionary step of progressing in the case of linear evolution(11). Microinvasion consists of cohorts of cancer cells that breach the basement membrane into the surrounding ECM (6,11,18,20). We hypothesize that non-genetic ecological factors, such as intra-ductal microenvironment, may be responsible for transitioning from DCIS to IDC. To validate this hypothesis, we propose a novel method to study DCIS evolution, by capturing and characterizing tumor ecological features such as "habitats" and "niches" in breast ecosystem (21). We started by defining the habitats based on availability of oxygen into: a) oxygenated habitat and b) hypoxic habitat defined by distance from the duct boundary (18, 22). We then defined local niches using biomarkers indicative of cancer cell response to variation in oxygen availability or acid adaptation. These biomarkers are designed based on our prior cancer eco-evolutionary studies and findings. Oxygen availability determines the source of energy production as of either mitochondrial respiration or glycolysis. Hypoxic cells switch to glycolysis, causing lactic acid production that can lead to acidosis when lactic acid is locally accumulated. Peri-luminal cells will experience hypoxia if they are far (>0.125 - 0.160 mm) from a blood supply. These cancer cells inhabit a microenvironment of hypoxia, acidosis, and

severe nutrient deprivation (18,22). Our previous studies strengthen the acid-induced evolution model
of breast cancer and our proposed evolutionary designed biomarkers including CA9 and LAMP2b in
this translational research (6,17,20,23,24). Here we examined the role of these biomarkers within an
eco-evolutionary concept as a predictor of DCIS upstaging for the first time. We used these markers as
representative of the cancer cells metabolic states to define niches that can select for more aggressive
phenotypes, leading to microinvasion and DCIS upstaging to IDC. Niches were defined as clusters of
cells with similar phenotype responding to hypoxia and/or acid.
To perform our analysis, we curated a retrospective cohort of DCIS patients, with specimens collected
from Biopsy (Bx) samples before surgery and the whole tumor after Excision (Ex). All the patients had
histologically confirmed DCIS at core Bx, followed by diagnosis confirmed on surgical excision
specimens with either DCIS or IDC (**Figure 1B**). Our prediction model is trained and tested on the Bx
samples. We first stained 3 sequentially sections for hematoxylin and eosin (HE), CA9 ab, and
LAMP2b ab. We then manually annotated every single duct bigger than 400 μms in diameter, 200 μms
in radius. The 200 μms in radius annotation ensures each duct has both oxygenated and hypoxic
habitats to build a balanced cohort for analysis. Then we scored the positivity of each habitat for both
CA9 and LAMP2b to compare both non-upstaged and upstaged group. We finally developed a novel
algorithm to detect intra-ductal niches based on biomarker expression similarity. We applied multiple
spatial functions and spatial entropies to define niche and micro-niches describing the spatial patterns
of the positive cells. Thus, we studied the spatial organization of CA9- and LAMP2b-positive cells at
four different scales: whole slide, duct, habitats, and niches. After a systematic and comprehensive
analysis, we observed that the spatial features at the finest niche level possess the most predictive
power where the micro-niches were defined by the expression of both CA9 and LAMP2b. By
characterizing these niches and micro-niches with spatial and pathomics features, we then developed a
risk scoring system by integrating pathological imaging and molecular features of early-stage breast
tumors (**Figure 1C**). We show that quantitative analyses of immuno-histological images combined
with the tumor's eco-evolution dynamics and underlying molecular pathophysiology can significantly
improve predicting if the neoplasm is invasive disease. Our developed machine learning model, fine-
tuning the tumor ecosystem into habitats and niches, provides informed decision support.
In summary, we show that eco-evolutionary applications can improve existing models of DCIS
progression. Our developed machine learning techniques to calculate the niches and habitats inside the
tumor of DCIS patients with pure DCIS and upstaged disease can provide supportive information to
understand the disease progression. Finally, we suggest that deploying eco-evolutionary principles and
machine learning techniques, can propose a novel consilient approach to improve stratifying DCIS
patients.
123

Materials and Methods

Method overview

Our evolutionary analysis pipeline takes 3 sequential sections of each patient sample, defines oxygen
habitats (normoxia and hypoxia) and abundance of CA9 and LAMP2b cells positives in those habitats,
detects intra-ductal cell niches, characterizes these niches' spatial and morphological features, and
finally use these eco-evolutionary markers to predict whether the patient will be upstaged or not. In
particular, the pipeline has 4 modules. First, we annotate single ducts to their habitats and score

habitats. Then align ducts from all three whole slide images (WSIs). This ensures cells of different
slides are aligned and we can characterize their interactions in habitats and niches. In the second
module, we detect and map all eco-evo positive cells (i.e., cells positive for CA9 and LAMP2b) into the
same duct on HE slide and detect different clusters of cells as niches. In the third module, we
characterize these niches with comprehensive spatial statistical features, as well as their morphological
features as observed in HE. Finally, we categorize these niches into different subclasses through deep
learning-based dimension reduction and feature-based clustering. We also use the distribution of niche
subclasses to characterize different samples/patients. We demonstrate the discriminative power of these
niche-based characterization in predicting whether a patient will be upstaged or not. **Figure 1C**
illustrates the overview of our pipeline.

Data preparation and usage

Data used in this study comes from biopsy samples collected after mammography and before surgery.
84 samples including 68 pure DCIS and 16 progressed to IDC were collected. This study complied with
the Health Insurance Portability and Accountability Act and was approved by the institutional review
board, with a waiver of the requirement for informed consent. Women with a core biopsy diagnosis of
DCIS between 2012 and 2022 who consented to Moffitt Cancer Center Total Cancer Care protocol
were included in this analysis. Cases were excluded if surgical excision was performed more than 6
months after the core biopsy, if there was concurrent ipsilateral invasive breast cancer or metastatic
malignancy, or if neoadjuvant chemotherapy (for a concurrent contralateral breast malignancy) or
chemotherapy for any malignancy was administered between the dates of the DCIS core Bx and
surgery. We also excluded the patient if there was a personal history of IDC or DCIS within 12 months
prior the Bx or a concurrent diagnosis of Paget disease in the ipsilateral breast. After applying these
inclusion and exclusion criteria, 84 cases of biopsy-proven DCIS were identified, of which 16 were
upstaged at surgery and 68 remained non-upstaged. From each patient one block selected with at least
one lesion in it. From each block, we obtained 3 whole slide images, including 1 HE and 2 IHC slides.
To minimize confounding effects, non-upstaged and upstaged patients were matched across clinical
features including age, race, ethnicity, tumor grade, and ER/PR status. Demographic characteristics of
the cohort (e.g., age range, race/ethnicity distribution) are summarized in **Supplementary Figure S1**.
Sample selection, immunohistochemistry and HE staining. Tumor blocks were selected by
pathologists using the archived HE stained slides. The blocks were sequentially sectioned 4 μm and de-
identified for research use. 3 slides were stained with primary antibodies of 1:100 dilution of anti-
LAMP2 (Abcam Cat# ab18529, RRID:AB_2134632)), and 1 ug/ml concentration of anti-CA9 (R&D
Systems Cat# AF2188, RRID:AB_416562), and HE staining using standard hematoxylin and eosin
protocol. Normal placenta was used as a positive control for LAMP2b and clear cell renal cell
carcinoma was used as a positive control for CA9. For the negative control, an adjacent section of the
same tissue was stained without application of primary antibody and any stain pattern observed was
considered as non-specific binding of the secondary. Primary immunohistochemical analysis was
conducted using digitally scanned slides. The scoring method used by our pathologist to determine (a)
the degree of positivity scored for each sample ranged from 0 to 3 and was derived from the product of
staining intensity (0 – 3+). A zero score was considered negative, score 1 was weak positive, score 2
was moderate positive, and score 3 was strong positive. (b) The percentage of positive cells (on a scale
of 0-3).

Whole slide imaging (WSI) of IHC and HE slides were obtained by scanning at 20X magnification (of
0.5022 micrometer per pixel) using Aperio AT2 from Leica Biosystems. Images were transferred to
cloud storage and locally to be uploaded in QuPath (v0.4.3; RRID:SCR_018257) software for analysis.
QuPath software was used to detect the positive pixels for each IHC marker (CA9 and LAMP2b) and to
segment the HE images into hypoxic and normoxic tumor habitats based on their distance from the
basement membrane. The 'Positive Cell Detection' function from QuPath was used to automatically
classify the positivity of CA9 and LAMP2b markers and validated by the study pathologist.
180

MODULE 1: Duct annotation and alignment

181
Manual annotation of ducts. We annotate and align ducts within all input slides (1 HE + 2 IHCs per
sample). After annotating ducts from WSIs of all three modalities, we align and co-register the ducts
from the three modalities. This alignment enables us to map cells into the same spatial domain in each
slide and analyze their interaction. QuPath was used as the interface to annotate ducts by the
pathologist (Dr. Bai) and reviewed by Dr. Damaghi. To ensure the same number of oxygenated and
hypoxic habitats, we only select ducts of >400 μm diameter from basement membrane. Following this,
based on distance, each duct was annotated with four layers: adjacent stroma, oxidative/normoxia,
hypoxic/hypoxia, and necrosis. Within the duct, necrosis was defined as any area containing dead cells,
as identified by a lack of nuclei. Oxidative layer was defined as the area containing cells inside the duct
within 125 μm of the basement membrane. Hypoxia was defined as the area containing cells inside the
duct further than 125 μm from the basement membrane. The annotations were done for all 84 samples
in the Bx cohort, and then were exported as standard GeoJSON files. Pathologists were blinded to
patient outcomes during annotation and scoring.
Co-registration. To build a multiplex image from monoplex slides, we register both CA9 and
LAMP2b IHC slides towards the HE slides. For that reason, we applied direct co-registration at the
whole slide level with manual landmarks for alignment at each duct. We further co-register the slides in
a duct-by-duct fashion. Using initially registered whole slides, and spatial proximity, we identify the
corresponding ducts at the HE and 2 IHC slides. Next, we register both the CA9 duct and LAMP2b
duct into the corresponding HE ducts. We use Virtual Alignment of pathology Image Series (VALIS),
which provides a fully automated pipeline to register whole slide images (WSI) using rigid and/or non-
rigid transformations (34). For each sample, we chose non-rigid registration and registered the ducts
from CA9 and LAMP2b towards the reference HE ducts. The co-registration procedure and the
qualitative results are shown in **Figure S4**. The co-registration provides a mapping of any cells detected
in CA9 or LAMP2b towards a shared spatial domain, enabling the analysis of their interactions.

MODULE 2: Cell detection and niche definition

206
Cell detection. With the duct annotations in place, we automatically detect cells from the 2 IHCs and
determine if they are positive in CA9 or LAMP2b based on their intensities. For each duct, we detect
cells using QuPath watershed cell detection algorithm (25). Based on the intensity level, we categorize
the cells into 4 groups: 'Negative', '1+', '2+', and '3+'. The detection of cells within a duct is done by
StarDist (25,35) extension in QuPath.
Graph construction for niche detection. After annotating all the positive cells (i.e., CA9 or LAMP2b
positive cells), they were mapped on HE slides, enabling us to detect niches on HE slides. We construct
a graph with the positive cells by connecting cells whose distances are smaller than a certain threshold

and detect connected components of the graph as representatives of cells living in "niches". Multiple
thresholds have been experimented and an optimum value is selected based on performance. Each niche
was set to have cells with similar eco-evo phenotype and be spatially coherent. Therefore, we overlay
both CA9 and LAMP2b positive cells into the same domain as an approximation of the local eco-evo
cell distribution (**Figure S5**). This gives us the opportunity to measure their interaction via spatial
statistical functions as defined later. Based on the same principle, we use cell morphological features
extracted from HE within the region of each niche to characterize the niche.

MODULE 3: feature extraction from niches for characterization

Once niches are detected, we extracted both spatial and morphological features. Then we characterize
spatial interaction patterns, utilizing various spatial functions as features. This is extra to the cell
features consisting of morphology and texture features that adopted in HE images.
For cellular features we measured both morphological and texture features. The morphological features
include area, eccentricities, circularity, elongation, extent, major axis length, minor axis length, solidity
and curvature. The texture features include angular second moment (ASM) of co-occurrence matrix,
contrast, correlation, entropy, homogeneity and intensity. All features were calculated following the
implementations in the sc-MTOP package (36).
Although we do not have exact cell-to-cell correspondence between the cells within a niche and cells
detected in HE, we still can aggregate morphological and texture features within the proxy of the cells
part of a niche to characterize that niche. For each niche, we identify the concave hull region enclosing
its eco-evo positive cells within a duct on HE slide. Next, we aggregate cell features across all HE-
detected cells within the corresponding region. For each cell feature dimension, we calculated its mean,
standard deviation, maximum, minimum, kurtosis and skewness.
Spatial features. We extract various spatial statistical functions (37) to characterize residing cells and
their interactions to define niches. These functions are listed below:
G Function: The G function, denoted as G(r), is the cumulative distribution function of nearest-
neighbor distance. The G function provides insights into the clustering or dispersion behavior of the
point pattern.

$$G(r) \ = \ P\{d(u, X \backslash u) \ \leq \ r | u \ \in \ X\}, d(\bullet) \ is \ the \ minimum \ distance$$

243 F Function: The F function, known as the empty space function, is the cumulative distribution
244 function of the empty-space distance. The F function is commonly used to assess the regularity
245 or inhibition patterns in point patterns.

$$F(r) \ = \ P\{d(u, X) \ \leq \ r\}, d(\bullet) \ is \ the \ minimum \ distance$$

247 K Function: Ripley's K function, denoted as K(r), is a measure of second-order intensity or spatial
248 interaction. It assesses whether points tend to be more clustered or dispersed within a certain distance r
249 compared to a CSR process. It considers both the distance and intensity of points to capture the
250 clustering behavior of the point pattern.

$$K(r) \ = \ \frac{|W|}{n(n-1)} \sum_{i=1}^{n} \sum_{j=1, j \neq i}^{n} \mathbf{1}\{d_{ij} \leq r\} e_{ij}(r), e_{ij}(\bullet) \ is \ the \ edge \ correction \ weight$$

252 L Function: L function is a variance stabled version of K function.

$$L(r) = \frac{\sqrt{K(r)}}{r}$$

We calculated G, F, and L functions in both univariate and multivariate fashions. For each of the
functions, the distances between source cell and the target cells are considered. Univariate spatial
functions sample source cells and target cells from the same type of cells while multivariate
counterparts' sample from different types of cells. Univariate G, F, and L are calculated for the single-
marker cell subsets, and multivariate G_cross and L_cross for different subsets such as CA9-LAMP2b.
'Gest' function and 'Fest' function from 'spatstat' R package (v3.3-1; https://CRAN.R-
project.org/package=spatstat) were used with Kaplan-Meier estimator (38), and 'Lest' function was
used with isotropic correction (39, 40).

MODULE 4: Diagnostic risk estimation with pattern proportion
We trained a classifier using the niches defined and characterized as described above to predict whether
a patient will be "upstaged" or "non-upstaged". This establishes the diagnostic power of the identified
niches. Worth to note that a direct aggregation of niche information within each sample/patient is not
sufficient. One technical challenge is that the niche features computed in the previous module are high
dimensional and the niche features are diversely distributed. To solve that we first find a simplified
distributional description of the niches, and then use the simplified description for prediction. First, we
cluster the niches into different sub-classes based on their features. The clustering is carried out using
K-means clustering with a tunable parameter k. Once the niche sub-classes are determined, we use their
distribution on each sample to predict its upstaged/not-upstaged status. The prediction power of the
classifier sheds light on the diagnostic power of the niches and their spatial and cellular features. Five-
fold cross-validation was employed, with one-fold designated as the test set in each run. This approach
prevents data leakage and helps mitigate overfitting. To understand the contribution of each feature to
the prediction model, we employed SHAP (SHapley Additive exPlanations) analysis. Machine learning
models were implemented using scikit-learn (RRID:SCR_019053)
276

277 **Data and Code Availability**
278 The data generated in this study are available within the article and its supplementary data files. All the
279 images and their annotations at duct or habitat levels are deposited in the physical sciences in oncology
280 network (PSON) and can be accessed at DOI: https://doi.org/10.1016/j.isci.2024.109433. Custom
281 Python and R analysis code is available at https://github.com/sbu-damaghi-ceel/DCIS_archive.
282

283 **Study Limitation**
284 This study excludes unaccounted clinical variables that can play role in DCIS upstaging. Another
285 limitation is the absence of independent external validation due to our particular annotations.
286

Results:
Sample curation and cohort building
We built a retrospective cohort of 84 patients with histologically confirmed DCIS on core Bx, followed
by surgical Ex, with available FFPE from Moffitt Cancer Center Biobank. The cohort has two arms: i)
pure DCIS: the patient diagnosed with DCIS at both Bx and Ex; ii) upstaged: patient with DCIS at Bx
and IDC at Ex (**Figure 1B**). HE slides of Bx cores were retrieved from the biobank at the Moffitt

Cancer Center tissue core and reviewed by a study pathologist (43). Then the selected blocks were pulled and sequentially cut for HE, CA9, and LAMP2b staining. The HE and subsequent 2 IHC slides are digitally scanned using the Aperio XT® high-throughput slide scanner and housed on the web-based Aperio server/Spectrum database package. Upstage status was pulled from the electronic medical record and confirmed by our study pathologist from the Ex tissues (**Figure 1C**). All images were then segmented and annotated using QuPath supervised by study pathologist (25) (43).

Non-upstaged and upstaged patients were matched across clinical features, including age, race, ethnicity, grade, and ER/PR status, to minimize their influence on the analysis (**Figure S1**). To validate the comparability of these groups, we conducted a Wilcoxon rank-sum test for the continuous variable (age) and Chi-square tests for the categorical variables (race, ethnicity, grade, ER/PR status). None of these tests showed significant differences between the two groups, with all p-values larger than 0.1, indicating that the groups were well-matched (**Figure S1**).

Annotation and eco-evolutionarily mapping of habitats at the individual duct level

We have shown previously that peri-luminal cells that are far (>0.125 - 0.160 mm) from a blood supply inhabit a microenvironment of hypoxia and acidosis (18,20,26). Thus, we created two simple annotation zones on HE slides based on oxygen diffusion distance representing oxygen habitats: i) hypoxic zone or habitat that is above 125 μm from the duct boundary, basement membrane, and ii) normoxic habitat that is the outer regions adjacent to the basement membrane (**Figure 2A**). We used the basement membrane as our zero point of reference. We also annotated necrotic zones inside the hypoxic habitats that also represent the anoxic habitat falling perfectly above 0.160 mm distance from basement membrane. Since adjacent stroma is also of interest to our group and others, we annotated adjacent stroma for each duct with binary scoring of 1 for having adjacent stroma or 0 for lacking it (**Supplementary Table 1**). To ensure a balanced representation of hypoxic and normoxic habitats, we excluded small ducts by establishing a duct size threshold of minimum 400 μm in diameter (or 200 μm radius) for manual annotation (**Figure S2**). After annotating all the ducts bigger than 200 μm of radius on HE slides, we expanded our annotations to other 2 consecutive IHC slides stained with CA9 and LAMP2b antibodies (**Figure 2B**). Subsequently, our pathologist, Dr. Bai, manually scored each duct for their architecture (Solid form, Cribriform, Papillary, and Comedo), nuclear grade, necrosis, microcalcification, reactive stroma, and lymphocyte aggregate. Interestingly inside one patient we found heterogeneity of ducts as they are different tumors implying the possible multiclonality. She then scored hypoxic and normoxic habitats based on CA9 and LAMP2b positivity using a scoring scale of 0–3 (**Supplementary Table 1**). Following this, positive cells in IHC slides were counted using QuPath (25), habitats were categorized into different classes based on the count of positive cells. The distribution of these habitat categories was compared between non-upstaged and upstaged groups (**Figure 2C, 2D, and S3**). Using the Wilcoxon test, it was shown that there existed significant differences between non-upstaged and upstaged group when habitats considered at the duct level. The tests were carried out for both hypoxic and oxidative layers for both CA9 (**Figure 2C and 2D**), and LAMP2b (**Figure S3**) as well as architecture, grade, lymphocytes, microcalcifications, and necrosis (**Supplementary Table 1**). As shown in Figure 2D, CA9 scoring within hypoxic habitats provides a much clearer distinction between pure DCIS and upstaged groups compared to the normoxic zone. Interestingly, CA9 did not show significant differences between the groups when analyzed at the whole duct or whole-slide level, as is traditionally done (**Figure S2B**). However, focusing on habitats revealed

that CA9-positive cells are distributed differently between the two patient groups. This analysis underscores the value of examining fine-scale habitats within ducts. The improved performance of habitat scoring compared to whole-duct scoring highlights the necessity and significance of exploring the cellular composition and interactions within the tumor ecosystem.

Mapping Metabolic Niches Within Habitats to Enhance Spatial Machine Learning Models

Previous analyses of hypoxic and normoxic habitats in ducts were limited to scoring each biomarker individually, focusing solely on the count of positive cells within each habitat. To broaden the scope and incorporate interactions and relationships between these two eco-evolutionary markers, we performed co-registration of two slides. This step enabled the creation of a virtual multiplex IHC (mIHC) by mapping cells onto a unified 2D reference space. HE slides were selected as the reference, and all IHC slides were registered onto it. (**Figure S4**). Note that since our analysis is carried out duct-by-duct, it is not necessary to register the whole slide. Instead, for each duct, we register its IHC stainings to its HE staining. This ensures all the downstream analyses could be performed on the same HE slide coordinates, providing consistency and precision in the spatial data integration. Then we used these mIHC images to define niches of cells that are positive for CA9, LAMP2b, or both. We hypothesized that niches characterized by both markers together would provide greater biological insight than analyzing each marker individually, given the established correlation between hypoxia and acid phenotypes. Then, we focus on the cell features such as nuclear morphology and texture and cell spatial features inside these niches to explore their predictive power on DCIS upstaging. As illustrated in **Figure 3**, we first map each positive cells to the reference HE slide using the co-registration described above. Then, by treating each positive cell as a node and connecting the cells within a distance threshold, we construct a cell-proximity graph out of mIHC positive cells whereby each connected component represents a continuous region or niche that is hypoxic, acidic, or both. The threshold is a tunable parameter that is optimized by the classifying power of the downstream analysis. And depending on the selection of the eco-evo markers, there can be CA9 rich niches, LAMP2b rich niches, or both CA9 and LAMP2b rich niches.

Post Analysis reveals the top contributing patterns and features of DCIS upstaging.

We then developed a pattern differential analysis pipeline, which comprises two stages: First, the samples are clustered based on the features and classified into patterns (**Figure S6**). We found eight distinct patterns shown by different color in Figure 4A. Then for each patient, we calculate the proportion of each pattern to build a distribution profile of each pattern (**figureS7**). By using these proportion features, we train a classifier to predict the upstaging (**Table 1**). We were able to predict the clinical outcome of a patient based on his/her spatially defined pattern distributions. Then, to test the hypothesis that finer regions (niche) with biological meanings could provide better predictive power, we conduct a multi scale analysis performing a series of experiments using the same set of features and with the same pattern differential analysis pipeline at 3 different scales: duct, habitat, and niche (**Figure 1C**). At the habitat level, normoxic and hypoxic zones are analyzed independently. At the niche level, analyses are further refined to separately examine CA9-positive cells, LAMP2b-positive cells, and cells positive for both. For all the experiments, the Bx dataset underwent 5-fold stratified cross-validation, where in each round, 4 folds served as the training dataset and 1-fold as the test dataset, with the goal of predicting the patients' clinical outcome at the time of Bx. Upon comparing the mean accuracy score

and the mean AUC score of all the classifiers, the niche level classifier yielded the best predictive
results particularly under both metrics (**Table 1**). This result confirms that niche-based analysis
outperforms our primary habitat analysis. The higher accuracy of the niche measurements may be
implying that niche measurement is better than inferring habitat. Also, it is worth mentioning that
oxygen habitat analysis is a rough estimate in our analysis since we do not know the exact location of
the vasculature and their activity. After identifying the best-performing classifier based on the AUC
metric we employed SHAP (42) (Shapley Additive exPlanations) analysis to interpret the model by
calculating SHAP values for each feature, specifically on the proportions of distinct patterns. SHAP is a
unified approach to interpreting machine learning models by assigning each feature an importance
value for a particular prediction. In our study, SHAP values were computed for the features
representing the proportions of different patterns within the niches. By calculating the SHAP values, we
could determine the impact of each feature on the model's output, thereby identifying the most
influential patterns that contribute to predicting DCIS progression. This step is crucial for ensuring the
transparency and interpretability of the machine learning model. Furthermore, we select features that
are highly relevant to the sub-classes using different approaches including covariance, mutual
information scoring and maximum relevance minimum redundancy (mRMR)(41) and choose the
features identified by all three approaches. The pattern with the maximum SHAP value, identified as
the most impactful, underwent further differential analysis to uncover features that significantly
differentiated this pattern from others. This differential analysis employed methods including
correlation analysis, mutual information (MI), and maximum relevance minimum redundancy
(MRMR), which together identified $F\_0 <= r < 10$ and solidity skewness as the top distinguishing
features for Pattern 5 (**Figure 4B**). **Figure 4C** shows the gradient map of each of these features of
niches in the latent space, while top differential features of other niche patterns can be found in **Figure**
S8.
403
Niche distribution for diagnosis. Tumor microenvironment is heterogeneous, and niches demonstrate
diverse spatial and morphological behavior. To account for the diversity, we focus on how different
niches are distributed across a sample. We show that the distributions of different niches essentially
characterize the tumor ecology in a much more refined manner compared with previous distance-based
definitions of hypoxia/oxidative layers. After assigning each duct to its sub-class, we aggregate across
all niches of each sample and use its sub-class distribution to characterize this sample. Assuming k
niche sub-classes, each sample has a k dimensional histogram to describe its niche sub-class
distribution. We call this the niche distributional (Nbd-Dist) feature. We trained a classifier to predict
whether a sample is pure DCIS or upstaged. Repeating the iteration 10 times and comparing the mean
AUC on the test set. We experimented several classifier types including lightGBM, soft vector machine
(SVM), logistic regression and random forest. Random forest classifier yields the best performance on
our data.
A prototype for Pattern 5, selected based on its alignment with the mean values of these features, was
visualized to illustrate its characteristics (**Figure 4D**). Using a multi-scale analytical approach, we
integrated spatial interactions of CA9 and LAMP2b positive cells into the machine learning pipeline to
distinguish between pure DCIS and upstaged group. Niche-level analysis yielded the highest accuracy
and AUC, emphasizing the importance of fine-scale regions in predicting clinical outcomes. The use of
SHAP analysis and differential analysis provided an interpretable framework to highlight influential

patterns and features, such as cell shape related features (Solidity) and spatial clusteredness related features (F function values), offering insights into the tumor microenvironment. This approach not only advanced our understanding of key spatial and morphological features but also demonstrated significant potential for precise diagnostic tools in clinical applications.

Discussion:

Ductal carcinoma in situ is the most prevalent type of precancer that can range from non-upstaged to aggressive. DCIS lesions are highly heterogeneous in their intra- and inter- ductal microenvironments, genetics, and molecular expression patterns. Recently, genomic analysis of matched DCIS and IDC samples has revealed that in 75% of cases, the invasive recurrence was found to be clonally related to the initial DCIS. This implies that tumor cells derived from DCIS could evolve in a linear or branching fashion with 18% new transformations and/or clonogenesis (11). These new findings emphasize the extraordinary heterogeneity in genotype and phenotypic plasticity in breast cancer that must be studied in the light of evolution and ecological studies. Tumors are complete ecosystems containing habitats and niches including normal epithelial cells, pre-cancer cells, stromal cells, vasculature, structural proteins, signaling proteins and physical factors such as pH and oxygen concentration (18). These habitats and niches can contain unique mixtures of cells with physical and biochemical characteristics, with differential evolutionary potential and trajectories (27). The niches with similar mixtures of cells usually are also similar in their physiology and phenotypes mainly due to living in similar habitats. Our hypothesis is that knowledge of these niches and their habitats can potentially provide patient benefit by stratifying their tumor progress and therapeutic choices. However, tools and techniques are lacking to distinguish them. Proper tools and techniques can identify and define habitats and niches to map pre-cancer ecosystems.

In this study, we argue that the overdiagnosis and overtreatment of DCIS stem from conventional frameworks that focus primarily on genetic signatures while neglecting the ecological heterogeneity within tumor ecosystems. Thus, we interpreted complex eco-evolutionary data of cancer cells within their niche using machine learning and pathomics, all framed within an innovative ecological and evolutionary dynamic model. Oxygen habitats are identified based on varying levels of perfusion and oxygenation, which are believed to play a crucial role in driving ecological diversity by changing cancer cells metabolism, creating new habitats, and enhancing tumor heterogeneity, ultimately leading to diverse evolutionary trajectories (28, 29). Solid tumors often exhibit an impaired vascular system, leading to habitats within tumors that vary in hypoxia, nutrient deficiency, and acidity. These habitats can significantly influence the spatial selection of cellular phenotypes in distinct subregions. Inhabiting hypoxia, acidosis, and severely nutrient deprived, face (pre-)cancer cells to strong selective pressures leading to divergence to novel phenotypes in population. These new phenotypes can reciprocally influence the microenvironment resulting in a dynamically changing tumor ecosystem. Therefore, the phenotype of the cells residing in these habitats can also be leveraged to define the habitats with a certain degree of accuracy. Previous research from our group and others demonstrated that cancer cells within breast ducts, exposed to chronic hypoxia and acidosis, develop adaptive mechanisms for survival in this challenging microenvironment including expression of CA9 or LAMP2b at the cell surface (18,20,30). However, none of these findings were used in a relevant translational study for biomarker discovery. In this study, we explore these biomarkers within an eco-evolutionary framework

for the first time, using them as indicators of the metabolic state of cancer cells residing in a niche as part of oxygen habitats that may favor the selection of more aggressive phenotypes to predict the upstaging of DCIS. While a longitudinal study would indeed be a better study design for direct observation of evolutionary changes over time, our current cross-sectional approach enables us to capture a snapshot of the tumor microenvironment at two near time points, providing valuable insight into the conditions that distinguish DCIS from IDC. We recognize the assumption that the synchronous IDC microenvironment may contribute to the progression from DCIS to IDC. However, our study design allows us to test whether specific microenvironmental factors and related habitats and niche correlate with the presence of IDC, which can provide strong hypotheses for future longitudinal investigations. A future prospective or retrospective longitudinal (multiple long time points) study would indeed help distinguish whether these microenvironmental changes in tumor ecosystem locally belonged to habitats or niches can drive progression from DCIS to IDC or if IDC-induced those changes in the tumor ecosystem contribute to the synchronous DCIS phenotype.

In our curated retrospective cohort of 84 DCIS patients with histologically confirmed DCIS on core biopsy, we manually annotated 916 single ducts and more than 3000 habitats on all three slides and scored them at habitat levels. This unique detailed eco-evolutionary annotation can be used for future similar eco-evolutionary designed studies including reactive stroma habitats. Our risk scoring system integrating principles of ecological-evolutionary dynamics with pathological imaging and molecular features of early-stage breast tumors showed improvement on prediction power of biomarkers alone and in combination.

We employed a 5-fold stratified cross-validation approach to ensure robust internal validation of our model. While this method helps mitigate overfitting and provides reliable performance estimates, we acknowledge the absence of an independent validation set, which is crucial for assessing the model's generalizability. The unique design of our cohort, which integrates specific ecological and microenvironmental factors, limits the availability of comparable external datasets for validation. As such, there is no current dataset with similar characteristics for cross-validation. We recognize this as a key limitation and emphasize that future studies should aim to validate the model on independent cohorts when such datasets become available. Furthermore, although our model achieved an AUC of 0.74, this performance is not yet sufficient for clinical translation. Additional efforts to refine the model and test it in larger, independent cohorts will be essential before its use in clinical practice can be considered. Interestingly, a recent approach using multiplex IF on DCIS cohort reached the same AUC(2). While both our study and the Risom et al. paper aim to leverage spatial relationships to predict DCIS progression, we would like to emphasize that the two approaches are fundamentally different in terms of the markers used. Risom et al. focused on a broad panel of markers, including those related to the stroma, immune cells, and tumor cells, which provide a comprehensive view of the tumor microenvironment. In contrast, our approach centers on eco-evolutionary markers derived from adaptation of cancer cells to physical microenvironment, specifically CA9 and LAMP2b, which are associated with hypoxia and tumor acidity and their spatial distribution, respectively. These differences reflect divergent hypotheses about the key drivers of DCIS progression. The fact that both studies report a similar AUC of 0.74, with the distinct marker sets and biological processes, suggests that our findings offer complementary insights into DCIS progression and combination of approaches might increase the accuracy.

Our study demonstrates the utility of eco-evolutionary principles in understanding DCIS progression. In our study, we proposed that specific tumor microenvironmental conditions, such as hypoxia and acidosis, are associated with phenotypic changes that may indicate DCIS progression. However, although we have shown previously that these microenvironments can cause aggressive phenotypes, we acknowledge that our findings here do not conclusively demonstrate that these environmental factors are causative agents in the transition from DCIS to IDC. Instead, our data suggests that these conditions could serve as biomarkers for identifying lesions that are more likely to be upstaged. However, the ability to define more refined cell phenotypes within each region of interest (ROI) could further enhance our analysis. If we can identify and characterize more detailed phenotypes, it would allow us to extract additional features that describe the spatial interactions of these phenotypes. This, in turn, could potentially improve the classifier's performance and make the results more interpretable. By capturing the intricate interactions between various cell types and their microenvironments, we could gain deeper insights into the ecological dynamics driving DCIS progression and improve predictive models for patient outcomes.

In recent years, there has been a growing trend towards adopting a "watchful waiting" approach for certain cases of DCIS, rather than immediate surgical excision (31,32). This strategy aims to reduce overtreatment by closely monitoring DCIS lesions that may not progress to invasive cancer. In this context, our upstaging predictions become particularly relevant. Identifying microenvironmental and phenotypic factors that indicate a higher likelihood of progression to IDC could help clinicians make more informed decisions about when to intervene and when to adopt a more conservative, observational approach. The ability to predict which DCIS cases are at higher risk of progressing to invasive disease would provide critical information for optimizing patient management, minimizing unnecessary treatments, and reducing the psychological and physical burdens associated with overtreatment (33). Further validation of these predictive models could therefore have important implications for guiding treatment strategies in the context of DCIS.

Lead contact

Further information and any related requests should be directed to and will be fulfilled by the lead contact Mehdi Damaghi (Mehdi.Damaghi@stonybrookmedicine.edu).

Acknowledgements

We gratefully acknowledge funding from Physical Sciences Oncology Network at the National Cancer Institute (grant U01CA261841), R01 grant R01CA272601, and R01CA249016. This work has been supported also in part by the Analytic Microscopy Core Facility at the H. Lee Moffitt Cancer Center & Research Institute, an NCI designated Comprehensive Cancer Center (P30-CA076292). We thank Dr. Alexander Borowsky and Dr. Kenneth Shroyer, for reading our manuscript and constructive comments on pathological aspect as well as Dr. Liliana Davolos and Dr. David Ray for their guidelines on ecological and evolutionary principles used in the manuscript.

Author contributions

M.D. conceptualized and designed the research; Y.X., M.A., J.D.B., A.C., Y.C., M.D., performed the experiment and analysis; J.D.B. reviewed all the slides and scored them as the project pathologist; P.P.,

C.C., and M.D. contributed to results interpretation; and M.D. wrote the paper. All authors revised the
paper.

References:

1. Greaves M, Maley CC. Clonal evolution in cancer. Nature. 2012; 481:306–13.

2. Risom T, Glass DR, Averbukh I, Liu CC, Baranski A, Kagel A, et al. Transition to
invasive breast cancer is associated with progressive changes in the structure and
composition of tumor stroma. Cell. 2022;185:299–310.e18.

3. Maley CC, Aktipis A, Graham TA, Sottoriva A, Boddy AM, Janiszewska M, et al.
Classifying the evolutionary and ecological features of neoplasms. Nat Rev Cancer.
2017;17:605–19.

4. Boutry J, Tissot S, Ujvari B, Capp J-P, Giraudeau M, Nedelcu AM, et al. The evolution
and ecology of benign tumors. Biochim Biophys Acta Rev Cancer. 2022;1877:188643.

5. Amend SR, Pienta K. Abstract 2884: Tumor-driven eutrophication of the tumor ecosystem
selects for cancer cell clones that overcome evolutionary inertia leading to increased
metastatic capacity. Cancer Res. American Association for Cancer Research;
2015;75:2884–2884.

6. Damaghi M, Mori H, Byrne S, Xu L, Chen T, Johnson J, et al. Collagen production and
niche engineering: A novel strategy for cancer cells to survive acidosis in DCIS and
evolve. Evol Appl. 2020;13:2689–703.

7. Lipinski KA, Barber LJ, Davies MN, Ashenden M, Sottoriva A, Gerlinger M. Cancer
Evolution and the Limits of Predictability in Precision Cancer Medicine. Trends Cancer
Res. 2016;2:49–63.

8. Giaquinto AN, Sung H, Miller KD, Kramer JL, Newman LA, Minihan A, et al. Breast
Cancer Statistics, 2022. CA Cancer J Clin. 2022;72:524–41.

9. Strand SH, Rivero-Gutiérrez B, Houlahan KE, Seoane JA, King LM, Risom T, et al.
Molecular classification and biomarkers of clinical outcome in breast ductal carcinoma in
situ: Analysis of TBCRC 038 and RAHBT cohorts. Cancer Cell. 2022;40:1521–36.e7.

10. Lehman CD, Arao RF, Sprague BL, Lee JM, Buist DSM, Kerlikowske K, et al. National
Performance Benchmarks for Modern Screening Digital Mammography: Update from the
Breast Cancer Surveillance Consortium. Radiology. 2017;283:49–58.

11. Lips EH, Kumar T, Megalios A, Visser LL, Sheinman M, Fortunato A, et al. Genomic
analysis defines clonal relationships of ductal carcinoma in situ and recurrent invasive
breast cancer. Nat Genet. 2022;54:850–60.

12. Sarhadi S, Salehzadeh-Yazdi A, Damaghi M, Zarghami N, Wolkenhauer O, Hosseini H.
Omics Integration Analyses Reveal the Early Evolution of Malignancy in Breast Cancer.
Cancers [Internet]. 2020;12. Available from: http://dx.doi.org/10.3390/cancers12061460

13. Heselmeyer-Haddad K, Berroa Garcia LY, Bradley A, Ortiz-Melendez C, Lee W-J,
585   Christensen R, et al. Single-cell genetic analysis of ductal carcinoma in situ and invasive
586   breast cancer reveals enormous tumor heterogeneity yet conserved genomic imbalances
587   and gain of MYC during progression. Am J Pathol. 2012;181:1807–22.

588   14. Hanna WM, Parra-Herran C, Lu F-I, Slodkowska E, Rakovitch E, Nofech-Mozes S. Ductal
589   carcinoma in situ of the breast: an update for the pathologist in the era of individualized
590   risk assessment and tailored therapies. Mod Pathol. 2019;32:896–915.

591   15. Carmeliet P, Jain RK. Principles and mechanisms of vessel normalization for cancer and
592   other angiogenic diseases. Nat Rev Drug Discov. 2011;10:417–27.

593   16. Wu B, Liu D-A, Guan L, Myint PK, Chin L, Dang H, et al. Stiff matrix induces exosome
594   secretion to promote tumour growth. Nat Cell Biol. 2023;25:415–24.

595   17. Persi E, Duran-Frigola M, Damaghi M, Roush WR, Aloy P, Cleveland JL, et al. Systems
596   analysis of intracellular pH vulnerabilities for cancer therapy. Nat Commun. 2018;9:2997.

597   18. Damaghi M, West J, Robertson-Tessi M, Xu L, Ferrall-Fairbanks MC, Stewart PA, et al.
598   The harsh microenvironment in early breast cancer selects for a Warburg phenotype. Proc
599   Natl Acad Sci U S A [Internet]. 2021;118. Available from:
600   http://dx.doi.org/10.1073/pnas.2011342118

601   19. Lobo RC, Hubbard NE, Damonte P, Mori H, Pénzváltó Z, Pham C, et al. Glucose Uptake
602   and Intracellular pH in a Mouse Model of Ductal Carcinoma In situ (DCIS) Suggests
603   Metabolic Heterogeneity. Front Cell Dev Biol. 2016;4:93.

604   20. Damaghi M, Tafreshi NK, Lloyd MC, Sprung R, Estrella V, Wojtkowiak JW, et al.
605   Chronic acidosis in the tumour microenvironment selects for overexpression of LAMP2 in
606   the plasma membrane. Nat Commun. 2015;6:8752.

607   21. Ordway B, Swietach P, Gillies RJ, Damaghi M. Causes and Consequences of Variable
608   Tumor Cell Metabolism on Heritable Modifications and Tumor Evolution. Front Oncol.
609   2020;10:373.

610   22. Gillies RJ, Verduzco D, Gatenby RA. Evolutionary dynamics of carcinogenesis and why
611   targeted therapy does not work. Nat Rev Cancer. 2012;12:487–93.

612   23. Ibrahim-Hashim A, Robertson-Tessi M, Enriquez-Navas PM, Damaghi M, Balagurunathan
613   Y, Wojtkowiak JW, et al. Defining Cancer Subpopulations by Adaptive Strategies Rather
614   Than Molecular Properties Provides Novel Insights into Intratumoral Evolution. Cancer
615   Res. 2017;77:2242–54.

616   24. Damaghi M, Gillies R. Phenotypic changes of acid-adapted cancer cells push them toward
617   aggressiveness in their evolution in the tumor microenvironment. Cell Cycle.
618   2017;16:1739–43.

619   25. Bankhead P, Loughrey MB, Fernández JA, Dombrowski Y, McArt DG, Dunne PD, et al.
620   QuPath: Open source software for digital pathology image analysis. Sci Rep.
621   2017;7:16878.

26. Freischel AR, Damaghi M, Cunningham JJ, Ibrahim-Hashim A, Gillies RJ, Gatenby RA, et
al. Frequency-dependent interactions determine outcome of competition between two
breast cancer cell lines. Sci Rep. 2021;11:4908.

27. Jardim-Perassi BV, Huang S, Dominguez-Viqueira W, Poleszczuk J, Budzevich MM,
Abdalah MA, et al. Multiparametric MRI and Coregistered Histology Identify Tumor
Habitats in Breast Cancer Mouse Models. Cancer Res. 2019;79:3952–64.

28. Sobhani F, Muralidhar S, Hamidinekoo A, Hall AH, King LM, Marks JR, et al. Spatial
interplay of tissue hypoxia and T-cell regulation in ductal carcinoma in situ. NPJ Breast
Cancer. 2022;8:105.

29. Compton ZT, Mallo D, Maley CC. Stronger together: Cancer clones cooperate to alleviate
growth barriers in critical cancer progression transitions. Cancer Res. American
Association for Cancer Research (AACR); 2023;83:4013–4.

30. Chafe SC, McDonald PC, Saberi S, Nemirovsky O, Venkateswaran G, Burugu S, et al.
Targeting hypoxia-induced carbonic anhydrase IX enhances immune-checkpoint blockade
locally and systemically. Cancer Immunol Res. American Association for Cancer
Research; 2019;7:1064–78.

31. Ryser MD, Worni M, Turner EL, Marks JR, Durrett R, Hwang ES. Outcomes of active
surveillance for ductal carcinoma in situ: A computational risk analysis. J Natl Cancer Inst.
Oxford University Press (OUP); 2016;108:djv372.

32. Glencer AC, Miller PN, Greenwood H, Maldonado Rodas CK, Freimanis R, Basu A, et al.
Identifying good candidates for active surveillance of ductal carcinoma in situ: Insights
from a large neoadjuvant endocrine therapy cohort. Cancer Res Commun. American
Association for Cancer Research (AACR); 2022;2:1579–89.

33. Fortunato A, Mallo D, Cisneros L, King LM, Khan A, Curtis C, et al. Evolutionary
Measures Show that Recurrence of DCIS is Distinct from Progression to Breast Cancer.
medRxiv [Internet]. 2024; Available from: http://dx.doi.org/10.1101/2024.08.15.24311949

34. Gatenbee CD, Baker A-M, Prabhakaran S, Swinyard O, Slebos RJC, Mandal G, et al.
Virtual alignment of pathology image series for multi-gigapixel whole slide images. Nat
Commun. 2023;14:4502.

35. Schmidt U, Weigert M, Broaddus C, Myers G. Cell Detection with Star-Convex Polygons.
Medical Image Computing and Computer Assisted Intervention – MICCAI 2018. Springer
International Publishing; 2018. page 265–73.

36. Zhao S, Chen D-P, Fu T, Yang J-C, Ma D, Zhu X-Z, et al. Single-cell morphological and
topological atlas reveals the ecosystem diversity of human breast cancer. Nat Commun.
2023;14:6796.

37. Baddeley A, Rubak E, Turner R. Spatial Point Patterns: Methodology and Applications
with R. CRC Press; 2015.

38. Baddeley A, Gill R. Kaplan-Meier estimators of interpoint distance distributions for spatial
point processes. IEEE Trans Inf Theory [Internet]. 1993; Available from:

https://ir.cwi.nl/pub/5247/05247D.pdf

39. Ohser J. On estimators for the reduced second moment measure of point processes. Series
Statistics. Taylor & Francis; 1983;14:63–71.

40. Kendall WS. Stochastic Geometry: Likelihood and Computation. Routledge; 2019.

41. Peng H, Long F, Ding C. Feature selection based on mutual information: criteria of max-
dependency, max-relevance, and min-redundancy. IEEE Trans Pattern Anal Mach Intell.
2005;27:1226–38.

42. Lundberg, S. M. & Lee, S.-I. A unified approach to interpreting model predictions. Proc.
Adv. Neural Inf. Process. Syst. 2017;30.

43. Mayfield, J. D., Ataya, D., Abdalah, M., Stringfield, O., Bui, M. M., Raghunand, N., Niell,
B., & El Naqa, I. Presurgical upgrade prediction of DCIS to invasive ductal carcinoma
using time-dependent deep learning models with DCE MRI. Radiology: Artificial
Intelligence. 2024; 6(5):e230348.

Figure Captions

Figure 1. Ecological and evolutionary designed biomarkers of DCIS upstaging. A) Model of
microenvironment-driven evolution of breast cancer from normal breast tissue to DCIS and IDC: Our schematic
is overlaid on HE staining of breast cancer specimens at different stages of DCIS and IDC. Different patients
may experience various types of evolutionary trajectory following different evolutionary models, including linear
and branched progression from DCIS to IDC shown here. Note that these events are not sequential or stepwise.
B) The patient cohort was curated from retrospective DCIS samples, with two sample collections at biopsy and
excision. The main criterion was the diagnosis of DCIS at the biopsy stage. **C)** Eco-evolutionary designed-
machine learning assisted pipeline to define cancer cell niches inside oxygen habitats in DCIS. *i)* Data
preprocessing steps including duct annotation, cell detection and classification for HE and IHC slides, followed
by co-registration to map IHC-identified cells onto the HE slides. *ii)*The analysis is carried out at multiple scales,
namely duct, habitat and niche, from the largest to smallest. At each scale the nucleus morphology texture
feature and spatial features are extracted. *iii)* The pattern differential analysis approach where the patterns are
firstly identified and then the proportions of such patterns are used as features to predict the upstaging status of a
patient.

Figure 2. **Eco-evolutionarily designed biomarker discovery to predict upstaging in DCIS. A)**
Illustration of normoxic, hypoxic and necrotic habitats in a duct. **B)** Illustration of annotation and
scoring on 2 IHCs and how cells are scored in each habitat. **C)** and **D)** Dot plots of counts of CA9
expression in each habitat per duct. Cells are scored 0 for 'negative' or '1+','2+','3+' for positive cells
based on their intensity. Scoring was performed and analyzed separately for normoxic (oxidative)
habitat (C) or hypoxic habitat (D). In the dot plot, each dot is a single duct. The color of dots reflects
their score as follows: Blue = 0, yellow ='1+', orange ='2+', and red = '3+'. The number of dots
reflects how many ducts were detected in each patient's biopsy with size bigger than 400 μm in
diameter. The distribution in hypoxic habitat is significantly different between pure DCIS and upstaged
groups in hypoxic habitats and not in oxygenated habitats. Data was analyzed using the Wilcoxon
signed-rank test. The same graph is created for LAMP2b (supplementary fig. 2).

Figure 3. Niches are defined inside habitats from the hypoxia and acidosis markers expression. A) One sample duct from CA9 slide. Top: The original IHC slide. Middle: Cell detection and intensity-based classification using QuPath overlaid on the slide. Bottom: the graph constructed from the CA9 positive cells and the connected components of the graph (Niches) highlighted in different colors. **B)** The HE staining of the same duct as A. Top: The original HE slide. Middle: Duct annotation overlaid on the HE slide. Bottom: Co-registered CA9-positive niches mapped and overlaid on HE slides as mIHC to be able to extract HE features from CA9 positive niches. Note the orientation of HE and CA9 slide was opposite, and our co-registration technique successfully created a mIHC of the ducts with similar coordinates. The same approach was used for LAMP2b and the combination.

Figure 4. Post Analysis and projection on mIHC reveals the top contributing ecological patterns and features. A) UMAP (Uniform Manifold Approximation and Projection) of the features of the niches, different colors represent different patterns **B)** Top: The impact of each pattern on the classifying result, blue and red colors represent impact on pure DCIS and progressed predictions respectively, the proportion of pattern 5 has the greatest impact for both categories. Bottom: Using correlation, MI, and MRMR to obtain the most contributing features in the pattern 5 clustering phase, identifying a common feature set that includes 2 features: $F_0 < = r < 10$ and solidity skewness. **C)** UMAP showing the value of the 2 identified features for different samples, and it indicates that samples in the pattern 5 tend to have low values for $F_0 < = r < 10$ and high values in solidity_skewness. **D)** A niche belonging to pattern 5, it exhibits a relatively dispersed distribution.

	Duct	**Habitat**		**Niche**		
		Normoxia	Hypoxia	CA9	LAMP2b	CA9 & LAMP2b
Accuracy	0.78 ± 0.06	0.86 ± 0.03	0.83 ± 0.06	0.82 ± 0.06	0.90 ± 0.03	0.90 ± 0.03
AUC	0.61 ± 0.08	0.67 ± 0.03	0.66 ± 0.10	0.64 ± 0.10	0.72 ± 0.07	0.74 ± 0.13

Table 1. **Performance scores of multi scale classifiers.** While habitat-level analysis enhanced performance, the niche-level classifier produced the most accurate predictive results.

Figure 1

Figure 2

A)

B) Segmentation and scoring

Manual annotation of Hypoxic habitat

Score hypoxic and normoxic habitats

C)

Oxidative

Pure DCIS   Progressed

Wilcoxon test between indolent and upstaged
p-value: 0.06452

Number of ducts

Patient

D)

Hypoxia

Pure DCIS   Progressed

Wilcoxon test between indolent and upstaged
p-value: 0.002253

Number of ducts

Patient

Figure 3

A)

B)

Co-registration

CA9

Figure 4