Eco-evolutionary Guided Pathomic Analysis to Predict DCIS Upstaging

Yujie Xiao[1], Manal Elmasry[2,3], Ji Dong K. Bai[2], Andrew Chen[2], Yuzhu Chen[2], Brooke Jackson[4],
Joseph O. Johnson[4], Robert J. Gillies[4], Prateek Prasanna[5], Chao Chen[5*], Mehdi Damaghi[1,2,3*]

1- Department of Applied Mathematics and Statistics, Stony Brook University, NY, USA

2- Department of Pathology, Stony Brook Medicine, Stony Brook University, NY, USA

3- Department of Pathology, Faculty of Medicine, Mansoura University, Mansoura, Egypt

4- Moffitt Cancer Center, Tampa, Fl, USA

5-Department of Biomedical Informatics, Stony Brook Medicine, Stony Brook University, NY, USA

[*] Corresponding authors: Chao Chen: chao.chen.1@stonybrook.edu, Mehdi Damaghi: mehdi.damaghi@stonybrookmedicine.edu

Abstract

Cancers evolve in a dynamic ecosystem. Thus, characterizing cancer's ecological dynamics is crucial to understanding cancer evolution and can lead to discovering novel biomarkers to predict disease progression. Ductal carcinoma in situ (DCIS) is an early-stage breast cancer characterized by abnormal epithelial cell growth confined within the milk ducts. In this study, we show that ecological habitat analysis of hypoxia and acidosis biomarkers can significantly improve prediction of DCIS upstaging. First, we developed a novel eco-evolutionary designed approach to define habitats in the tumor intra-ductal microenvironment based on oxygen diffusion distance. Then, we identify cancer cells with metabolic phenotypes attributed to their habitat conditions, such as the expression of CA9 indicating hypoxia responding phenotype, and LAMP2b indicating the acid adaptation. Traditionally these markers have shown limited predictive capabilities for DCIS upstaging, if any. However, when analyzed from an ecological perspective, their power to differentiate between pure DCIS and upstaged DCIS increased significantly. Second, using eco-evolutionary guided computational and digital pathology techniques, we discovered distinct niches with spatial patterns of these biomarkers and used the distribution of such niches to predict patient upstaging. The niches patterns were characterized by pattern analysis of both cellular and spatial features. With a 5-fold validation on the biopsy cohort, we trained a random forest classifier to achieve the area under curve (AUC) of 0.74. Our results affirm the importance of using eco-evolutionary-designed approaches in biomarkers discovery studies in the era of digital pathology by demonstrating the role of tumor ecological habitats and niches.

Keywords:

Breast cancer, Tumor ecology and evolution, DCIS, Eco-evolutionary biomarkers, Metabolic phenotypes, Habitat analysis, Niche analysis, Pathomic, Machine learning, Digital pathology

Introduction:

In recent years, the understanding that cancer is a dynamic ecological and evolutionary process has become deeply entrenched (1,2)(3). To date, several evolutionary approaches have been adapted and applied in cancer biology, such as diversity measures to predict disease progression; however, tumor

ecosystem and ecological habitat and niche studies are still overlooked (3,4). Within the human body and much like organisms in the natural world, cancer cells follow evolutionary principles, utilizing resources and establishing habitats and niches within tissues (5,6). This ecological perspective of cancer is crucial for discovering the natural selection driving cancer evolution. Recognizing the parallels between organismal ecology and the tumor microenvironment opens up untapped opportunities to incorporate ecological measures, improving our understanding of both tumor dynamics and selective pressures shaping tumors' evolutionary landscapes. Such insights may potentially lead to improved cancer prognosis, progression prediction, risk stratification, and therapeutic strategies. If tumor evolutionary state and/or its evolutionary trajectories could be reliably achieved using a single biopsy tissue, clinical translation would be comparatively more manageable. Nevertheless, studies have yet to determine whether measures of tumor evolvability derived from a single biopsy sample are adequate, or if the inclusion of multiple samples significantly enhances predictions of clinical outcomes (7).

Breast cancer incidence in the US has been increasing over the past decade at a rate of 0.5% per year(8). With increased mammographic screening, there has been a substantial increase in detecting the early non-invasive forms of breast cancer, such as ductal carcinoma in situ (DCIS)(2,9). About one-third of breast cancers detected by mammography are DCIS(10). As the most common pre-cancer state, DCIS can progress to invasive disease in a linear evolution pattern, or can be part of other clonal evolutionary dynamics such as branching, punctuated, or neutral evolution (2,9,11). Since DCIS and IDC (invasive ductal carcinoma) are indistinguishable by (epi-)genetic mutations, gene expression, or protein biomarkers, and because it is not possible to predict whether DCIS will remain indolent or progress to more aggressive disease, almost all early tumors are treated with aggressive interventions(2,12–14). To avoid such over treatment, more research is needed to fully understand evolution from pre-cancer to indolent DCIS or progress to IDC(9).

DCIS is a heterogeneous group of neoplastic lesions confined to the mammary ducts. The confinement of proliferating neoplastic cells inside the duct and growth of pre-cancer cells toward the center of the duct, which is far from vasculature, causes limitations in oxygen and nutrients. This intraductal oxygen microenvironment is also influenced by complex ecosystems surrounding the duct, such as vascular activity(15), stiffness of extracellular matrix (ECM) (16), and metabolites(6,17,18) (19) (**Figure 1A**) . Local microinvasion is the main difference between DCIS and IDC and might also be the first evolutionary step of progressing in the case of linear evolution(11). Microinvasion consists of cohorts of cancer cells that breach the basement membrane into the surrounding ECM. Recently, genomic analysis of matched DCIS and IDC samples has revealed that in 75% of cases, the invasive recurrence was found to be clonally related to the initial DCIS. This implies that tumor cells derived from DCIS could evolve in a linear or branching fashion with 18% new transformations and/or clonogenesis (11). These new findings emphasize the extraordinary heterogeneity in genotype and phenotypic plasticity in breast cancer that must be studied in light of evolution and ecological studies. Thus, we designed our study to capture the phenotypic heterogeneity of cancer cells in their selective microenvironments. We hypothesize that non-genetic ecological factors, such as intra-ductal microenvironmental conditions, may be responsible for transitioning from a DCIS to IDC phenotype, in the case of linear and branching evolution, or may select clones with pre-existing IDC phenotypes in the case of the other evolutionary trajectories, including punctuated and neutral evolution(6,11,18,20).

To validate this hypothesis, we propose a novel method to study DCIS evolution, by capturing and characterizing "tumor habitats" and "cell niches" and their interactions in the tumor ecosystem. Natural selection requires phenotypic diversity within a population undergoing microenvironmental selection forces (21). Cells that adapt in response to natural selection may present similar phenotypes,

Figure 1. Ecological and evolutionary designed biomarkers of DCIS upstaging. A) Model of microenvironment-driven evolution of breast cancer from normal breast tissue to DCIS and IDC: Our schematic is overlaid on HE staining of breast cancer specimens at different stages of DCIS and IDC. Different patients may experience various types of evolutionary trajectory following different evolutionary models, including linear and branched progression from DCIS to IDC shown here. **B)** The patient cohort was curated from retrospective DCIS samples, with two sample collections at biopsy and excision. The main criterion was the diagnosis of DCIS at the biopsy stage. **C)** Eco-evolutionary designed- machine learning assisted pipeline to define cancer cell niches inside oxygen habitats in DCIS. *i)* Data preprocessing steps including duct annotation, cell detection and classification for HE and IHC slides, followed by coregistration to map IHC-identified cells onto the HE slide. *ii)* The analysis is carried out at multiple scales, namely duct, habitat and niche, from the largest to smallest. At each scale the nucleus morphology texture feature and spatial features are extracted. *iii*)The pattern differential analysis approach where the patterns are firstly identified and then the proportions of such patterns are used as features to predict the upstaging status of a patient.

corresponding to the microenvironment exerting the selection. We started by defining the habitats based on availability of oxygen into: a) oxygenated habitat and b) hypoxic habitat. Following previous theory(18,22), these habitats are defined by distance from the duct boundary. However, a uniform distance threshold hardly captures the true oxidate/hypoxic states of cells. Therefore, we further proposed to fine-tune these habitats using protein expression indicative of phenotypes resulting from cancer cell adaptation to variation in oxygen availability. Therefore, we defined *intraductal DCIS niches* inside habitats as clusters of cells with similar phenotypic behavior responding to hypoxia. Through analysis via these niches, we can identify more aggressive phenotypes leading to

microinvasion and DCIS upstaging to IDC or possible direct evolution to IDC without going through DCIS sub-stages.

Our biomarkers are designed based on prior biological knowledge. Oxygen availability determines the source of energy production as of either mitochondrial respiration or glycolysis. Hypoxic cells switch to glycolysis, causing lactic acid production that can lead to acidosis when lactic acid is not cleared from the tumor space. Peri-luminal cells will experience hypoxia if they are far (>0.125 - 0.160 mm) from a blood supply. These cancer cells inhabit a microenvironment of hypoxia, acidosis, and severe nutrient deprivation (18,22). These environmental properties exert a strong selection pressure upon the cancer cells, which in turn feeds back to the microenvironment, creating a dynamically changing tumor ecosystem containing several habitats. We have shown that cancer cells within breast ducts subjected to chronic hypoxia and acidosis evolve mechanisms of adaptations to survive in this harsh microenvironment (17,18,20). We have also shown that cells adapted to hypoxic and/or acidic niches have developed specific metabolic vulnerabilities that can be targeted to push them back to a more physiologically normal state(17). Both these studies strengthen the acid-induced evolution model of breast cancer and our proposed evolutionary designed biomarkers including CA9 and LAMP2b in this research(6,17,18,20,23,24). Here we examined the role of these biomarkers within an eco-evolutionary concept as a predictor of DCIS upstaging for the first time. We used these markers as representative of the cancer cell metabolic states to define niches inside habitats that can select for more aggressive phenotypes, leading to microinvasion and DCIS upstaging to IDC or possible direct evolution to IDC without going through DCIS sub-stages.

To perform our analysis, we curated a retrospective cohort of DCIS patients, with specimens collected from Biopsy (Bx) samples before surgery and after Excision (Ex). All the patients had histologically confirmed DCIS on core biopsy, followed by diagnosis confirmed on surgical excision specimens with either DCIS or IDC (**Figure 1B**). Our niche-based prediction model is trained and tested on the Bx samples. These best fits future clinical applications that machine learning model can be subsequently applied to predict upstaging at Bx for future patients. We then stained 3 sequentially sectioned slides for HE, CA9 and LAMP2b. We manually annotated ducts bigger than 400 μms in diameter. The 200 um in radius annotation ensures each duct has both oxygenated and hypoxic habitats to build a balanced cohort for analysis. We developed a novel algorithm to detect intra-ductal DCIS cell niches based on biomarker expression similarity. Then, we studied the spatial organization of CA9- and LAMP2b-positive cells as the eco-evolution markers of cancer cells in hypoxic and acidic habitats at three different scales: whole slide, duct, and oxygen habitats (normaxic and hypoxic). We also applied multiple spatial functions and spatial entropies were used to define niche and micro-niches describing the spatial patterns of the cell groups. After a systematic and comprehensive analysis, we observed that the spatial features at the finest habitat level possess the most predictive power where the micro-niches were defined by the expression of CA9 and LAMP2b in hypoxic habitats. By characterizing these niches and micro-niches with spatial and pathomic features, we then developed a risk scoring system by integrating principles of ecological-evolutionary dynamics with pathological imaging and molecular features of early-stage breast tumors (**Figure 1C**). We show that quantitative analyses of immuno-histological images combined with the tumor's eco-evolution dynamics and underlying molecular pathophysiology can significantly improve predicting if the neoplasm has already evolved to invasive disease and cancer. We developed a machine learning model fine-tuning the tumor habitats into micro-

niches using specific molecular signatures of resident cancer cells to provide informed decision
support.
In summary, we show that specific habitats containing micro-niches of cells with similar phenotypes
responding to hypoxia and acidosis, or adaptation to long term exposure of these conditions, are
responsible for DCIS progression, and hence would be correlated to upstaging. To test this hypothesis,
we applied machine learning techniques to calculate the niches inside the tumor to define spatial and
temporal distribution of habitats in solid tumors of DCIS patients with pure DCIS and upstaged disease.
By deploying eco-evolutionary principles and machine learning techniques, our work proposes a novel
consilient approach - as opposed to the traditional single biomarker studies - to stratify DCIS patients

Results:

Sample curation and cohort building

We built a retrospective cohort from 84 patients with histologically confirmed DCIS on core biopsy,
followed by surgical excision, with available FFPE blocks at both Bx and Ex. The cohort has two arms:
the first one is pure DCIS including the patient diagnosed with DCIS at both Bx and Ex. The second
arm includes the upstaged group with DCIS at Bx and IDC at Ex (**Figure 1B**). Hematoxylin and eosin
(HE) stained slides of DCIS biopsy cores were retrieved from both the biobank core at Stony Brook
University and the Moffitt Cancer Center tissue core and reviewed by our study pathologist. Then the
selected blocks were pulled and sequentially cut for HE staining and CA9 and LAMP2b IHC staining.
The HE and subsequent 2 IHC slides are digitally scanned using the Aperio XT® high-throughput slide
scanner and housed on the web-based Aperio server/Spectrum database package. Upstage status was
pulled from the electronic medical record and confirmed by our study pathologist from the Ex tissues
(**Figure 1C**). All images were then segmented and annotated using Qupath supervised by study
pathologist (25).

Annotation of eco-evolutionarily defined habitats at the individual duct level.

We have shown previously that peri-luminal cells that are far (>0.125 - 0.160 mm) from a blood supply
inhabit a microenvironment of hypoxia and lactic acidosis (18,20,26). Thus, we created two simple
annotation zones on HE slides based on O2 diffusion distance representing oxygen habitats: i) hypoxic
zone or habitat that is above 125 μms from the duct boundary, basement membrane, and ii) normoxic
habitat that is the outer regions adjacent to the basement membrane (**Figure 2A**). We used the basement
membrane as our zero point of reference. We also annotated necrotic zones inside the hypoxic habitats
that also represent the anoxic habitat falling perfectly above 0.160 mm distance from basement
membrane. Since adjacent stroma is also of interest to our group and others, we annotated adjacent
stroma for each duct with binary scoring of 1 for having adjacent stroma or 0 for lacking it
(**Supplementary Table 1**). To ensure a balanced representation of hypoxic and normoxic habitats, we
established a duct size threshold of minimum 400 μms in diameter (or 200 μms radius) for manual
annotation (**Figure S1**). After annotating all the ducts bigger than 200 ums of radius on HE slides, we
expanded our annotations to other 2 consecutive IHC slides stained with CA9 and LAMP2b antibodies
(**Figure 2B**). Then our pathologist manually scored each duct for both hypoxic and normoxic habitats
separately for positivity of CA9 (0-3) (**Supplementary Table 1**). Following this, positive cells in IHC
slides were counted using Qupath(25), habitats were categorized into different classes based on the
count of positive cells, and the count of each category of habitats for each patient was then compared
between the pure DCIS and progressed DCIS patient groups.

Figure 2. **Eco-evolutionarily designed biomarker discovery to predict upstaging in DCIS. A)** Illustration of normoxic, hypoxic and necrotic habitats in a duct. **B)** Illustration of annotation and scoring on 2 IHCs and how cells are scored in each habitats. **C)** and **D)** Dot plots of counts of CA9 expression in each habitat per duct. Cells are scored 0 for 'negative' or '1+','2+','3+' for positive cells based on their intensity. Scoring was performed and analyzed separately for normoxic (oxidative) habitat (C) or hypoxic habitat (D). In the dot plot, each dot is a single duct. The color of dots reflect their score as follows: Blue=0, yellow='1+', orange='2+', and red='3+'. The number of dots reflects how many ducts were detected in each patient's biopsy with size bigger than 400 um in diameter. The distribution in hypoxic habitat is significantly different between pure DCIS and upstaged groups in hypoxic habitats and not in oxygenated habitat. Data was analyzed using the Wilcoxon signed-rank test. The same graph is created for LAMP2b (supplementary fig. 2)

Using the Wilcoxon test, it was shown that there existed significant differences. The tests were carried
out for both hypoxic and oxidative layers for 2 stains; the result for CA9 (**Figure 2C and 2D)**, and for
LAMP2b markers **(Figure S2)** as well as architecture, grade, lymphocytes, microcalcifications, and
necrosis (**Supplementary Table 1**). As shown in **Figure 2D**, CA9 scoring in hypoxic habitat is much
more definitive between pure DCIS and upstaged groups compared to normoxic zone, or the whole
duct, or the whole slide scoring as done traditionally (**Figure S1B**). We showed that the CA9-positive
cells distribute differently between the 2 patient groups if we focused on hypoxic habitats or oxidative
habitats. The performance improvement comparing the count scoring in different habitats and duct
scoring in the whole duct suggests that exploring the cell composition and interaction in fine habitats
inside ducts are meaningful and necessary.
194

Defining metabolic niches inside habitats to build spatial machine learning model and improve
performance
Previous scoring of hypoxic and normoxic habitats was carried out for each biomarker individually and
was restricted to only the count of the positive cells in each habitat. In order to expand the analysis to
involve the interaction and relationship between different types of eco-evolutionary marker positive
cells, a co-registration step is crucial to create multiplex IHC (mIHC) and map cells onto a common
reference 2D space. We chose the HE slides as the reference and registered all IHC slides on to it
(**Figure S2**). Note that since our analysis is carried out duct-by-duct, it is not necessary to register the
whole slides. Instead, for each duct, we register its IHC staining with the HE staining. This ensures all
the downstream analyses could be performed on the same HE coordinates system for each duct. Then
we used these mIHC images to define niches of cells representing CA9, LAMP2b, or combination
phenotype. We hypothesized that these niches inside habitats using the two phenotypes can be more
informative than each marker alone. Thus, we focus on the cell features such as nuclear morphology
and texture and cell spatial features inside these niches to explore their effect on upstaging. As
illustrated in **Figure 3**, we first map each marker IHC positive cells to the reference HE slide using the
co-registration described above. Then, by treating each positive cell as a node and connecting the cells
within a distance threshold, we construct a cell-proximity graph out of mIHC positive cells whereby
each connected component of this graph represents a continuous region or niche that is hypoxic, acidic,
or both. The threshold is a tunable parameter that is optimized by the classifying power of the
downstream analysis. And depending on the selection of the eco-evo markers, there can be CA9

positive niches, LAMP2b positive niches, and CA9 and LAMP2b positive niches. We then develop a
pattern differential analysis pipeline, which comprises two stages. First, the samples are clustered based
on the features and classified into one of the clusters or patterns. Then for each patient, we calculate the
proportion of each pattern, forming a distribution profile of the patterns.

Figure 3. Niches are defined inside habitats from the hypoxia and acidosis markers expression. A) One sample duct from CA9 slide. Top: The original IHC slide. Middle: Cell detection and intensity-based classification using Qupath overlaid on the slide. Bottom: the graph constructed from the CA9 positive cells and the connected components of the graph (Niches) highlighted in different colors. **B)** The HE staining of the same duct as A. Top: The original HE slide. Middle: Duct annotation overlaid on the HE slide. Bottom: Co-registered CA9-positive niches mapped and overlaid on HE slide as mIHC to be able to extract HE features from CA9 positive niches. Note the orientation of HE and CA9 slide was opposite and our co-registration technique successfully created a mIHC of the ducts with similar coordinates. The same approach was used for LAMP2b and the combination.

By using these proportion features, we train a classifier aiming to predict the upstaging status. From
this pipeline, we are able to predict the clinical outcome of a patient based on his/her spatially-defined
pattern distributions.(**Figure 1C**). Then, to test the hypothesis that finer regions with biological
meanings could provide better predictive power, we conduct a multi scale analysis performing a series
of experiments using the same set of features and with the same pattern differential analysis pipeline at
3 different scales: duct, habitat and niche (**Figure 1C**). Under the habitat level, normoxic and hypoxic
zones are analyzed separately. And under the niche level, CA9-positive cells, LAMP2b-positive cells,
and CA9- and LAMP2b-positive cells are analyzed separately.
For all the experiments, the biopsy dataset underwent 5-fold stratified cross-validation, where in each
round, 4 folds served as the training dataset and 1 fold as the test dataset, with the goal of predicting the
patients' clinical outcome at the biopsy stage. Upon comparing the mean accuracy score and the mean
AUC score of all the classifiers, the niche level classifier yielded the best predictive results under both
metrics (**Table 1**). This result confirms that niche-based analysis outperforms our primary habitat
analysis. The higher accuracy of the niche measurements may be implying the phenotype-based niche
measurement is better than inferring habitat from oxygen diffusion rate measure based on the distance
of the cells from basement membrane. Also, it is worth mentioning the oxygen habitat analysis is a
rough estimate in our analysis since we do not know the exact location of the vasculature and their
activity.

	Duct	Habitat		Niche		
		Normoxia	Hypoxia	CA9	LAMP2b	CA9 & LAMP2b
Accuracy	0.78 ± 0.06	0.86 ± 0.03	0.83 ± 0.06	0.82 ± 0.06	0.90 ± 0.03	0.90 ± 0.03
AUC	0.61 ± 0.08	0.67 ± 0.03	0.66 ± 0.10	0.64 ± 0.10	0.72 ± 0.07	0.74 ± 0.13

238 **Table 1**. **Performance scores of multi scale classifiers.**

Figure 4. Post Analysis reveals the top contributing patterns and features. A) UMAP of the features of the niches, different colors represent different clusters(patterns) **B)** Top: The impact of each pattern on the classifying result, blue and red colors represent impact on pure DCIS and progressed predictions respectively, the proportion of pattern 5 has the greatest impact for both categories. Bottom: Using correlation, MI, and MRMR to obtain the most contributing features in the pattern 5 clustering phase, identifying a common feature set that includes 4 features: Area_min, Perimeter_min, AreaBbox_min, and F_0<=r<10. **C)** UMAP showing the value of the 4 identified features for different samples, and it can be seen that samples in the pattern 5 tend to have higher values in Area_min, Perimeter_min, AreaBbox_min and low values for F_0<=r<10. **D)** A niche belonging to pattern 5, it contains no small size cells and exhibits a relatively dispersed distribution.

Post analysis to reveal contributing features and prototype visualization on mIHC.

After selecting the best classifier based on the AUC metric, we ran SHAP[48] (Shapley Additive
exPlanations) analysis to obtain the SHAP values for each feature, specifically the proportions of each
pattern. The most contributing pattern was identified as the one with the maximum SHAP value.
Subsequently, a differential analysis was performed for this pattern to identify the top features that
significantly differ from other patterns. These features were determined using correlation, mutual
information (MI), and maximum relevance minimum redundancy (MRMR) methods. For pattern 5, the
common feature set identified included Area_min, Perimeter_min, AreaBbox_min, and $F\_0 <= r < 10$.
A prototype for pattern 5 was then selected, which closely matched the mean values of these top
features, and visualized. This process is illustrated in **Figure 4**. By leveraging multi-scale analysis and
integrating spatial interactions of CA9 and LAMP2b positive cells through a comprehensive machine
learning pipeline, we were able to identify key patterns and features that differentiate between pure
DCIS and progressed DCIS. The use of SHAP analysis and differential analysis allowed us to pinpoint
the most impactful patterns and their contributing features, such as Area_min, Perimeter_min,
AreaBbox_min, and $F\_0 <= r < 10$, enhancing the interpretability of the model. Overall, the niche-level
analysis yielded the highest accuracy and AUC, underscoring the importance of fine-scale, biologically
meaningful regions in predicting clinical outcomes. This approach not only advances our understanding
of tumor microenvironments, but also holds promise for more precise diagnostic tools in clinical
settings.
259

Discussion:

Ductal carcinoma in situ is the most prevalent type of precancer that can range from indolent to
aggressive. DCIS lesions are highly heterogeneous in their intra- and inter- ductal physical
microenvironments, genetics, and molecular expression patterns. They can be described as complete
ecosystems containing habitats and niches including normal epithelial cells, pre-cancer cells, stromal
cells, vasculature, structural proteins, signaling proteins and physical factors such as pH and oxygen
concentration(18). These habitats and niches of micro-domains can contain unique mixtures of cells
with physical and biochemical characteristics, with differential evolutionary potential and trajectories
(27). The niches with similar mixtures of cells usually are also similar in their physiology and
phenotypes mainly due to living in similar habitats. Our hypothesis is that knowledge of these niches
and their habitats can potentially provide patient benefit by stratifying their tumor progress and
therapeutic choices. However, tools and techniques are lacking to distinguish them. Proper tools and
techniques can identify and define habitats and niches to map (pre-)cancer ecosystems to discriminate
between the different types of DCIS in order to design the right treatment for breast cancer patients.
Here, we argue that the reason for DCIS overdiagnosis and overtreatment results from conventional
frameworks focusing on genetic signature and ignoring phenotypic heterogeneity in tumor ecosystems.
Then we interpret complex eco-evolutionary data of cancer cells in their niche using machine learning
and pathomics within an innovative ecological and evolutionary dynamic framework. Oxygen habitats
are recognized from their variable levels of perfusion and oxygenation. It has been suggested that this
variability serves as a significant factor in influencing the ecological diversity, new habitats, and
increased tumor heterogeneity leading to diverse evolutionary trajectories(28,29). Solid tumors often
exhibit an impaired vascular system, leading to habitats within tumors that vary in hypoxia, nutrient
deficiency, and acidity. These habitats can significantly influence the regional selection of cellular

phenotypes in distinct subregions. On the other hand, the phenotype of the cells inhabiting these niches can be used to define the habitats. Inhabiting hypoxia, acidosis, and severe nutrient deprivation niches, face (pre-)cancer cells to strong selective pressures leading to divergence to novel phenotypes in population. These new phenotypes can reciprocally influence the microenvironment reshaping due to their new metabolic phenotypes resulting in a dynamically changing tumor ecosystem with multiple habitats. Previous research from our group and others demonstrated that cancer cells within breast ducts, exposed to chronic hypoxia and acidosis, develop adaptive mechanisms for survival in this challenging microenvironment(18,20,30). However, none of these findings were used in an eco-evolutionary designed translational study for biomarker discovery or treatment design. In this study, we explore these biomarkers within an eco-evolutionary framework for the first time, using them as indicators of the metabolic state of cancer cells to define habitats that may favor the selection of more aggressive phenotypes, which in turn can be used to predict the upstaging of DCIS. While a longitudinal study would indeed be a better study design for direct observation of evolutionary changes over time, our current cross-sectional approach enables us to capture a snapshot of the tumor microenvironment at two time points, providing valuable insight into the conditions that distinguish DCIS from IDC. We recognize the assumption that the synchronous IDC microenvironment may contribute to the progression from DCIS to IDC. However, our study design allows us to test whether specific microenvironmental factors correlate with the presence of IDC, which can provide strong hypotheses for future longitudinal investigations. A future prospective or retrospective longitudinal (multiple long time points) study would indeed help distinguish whether microenvironmental changes drive progression from DCIS to IDC or if IDC-induced changes in the microenvironment contribute to the synchronous DCIS phenotype.

We curated a retrospective cohort of 84 DCIS patients with histologically confirmed DCIS on core biopsy, followed by surgical excision diagnosed as either DCIS or IDC. We then stained the 2 sequentially sectioned slides for HE, CA9 and LAMP2b and manually annotated 916 single ducts and more than 3000 habitats on all three slides and scored them. This unique detailed eco-evolutionary annotation can be used for future similar eco-evolutionary designed studies including stroma habitats. Our risk scoring system integrating principles of ecological-evolutionary dynamics with pathological imaging and molecular features of early-stage breast tumors showed improvement on prediction power of biomarkers alone and in combination.

In our study, we employed a 5-fold stratified cross-validation approach to ensure robust internal validation of our model. While this method helps mitigate overfitting and provides reliable performance estimates, we acknowledge the absence of an independent validation set, which is crucial for assessing the model's generalizability. The unique design of our cohort, which integrates specific ecological and microenvironmental factors, limits the availability of comparable external datasets for validation. As such, there is no current dataset with similar characteristics for cross-validation. We recognize this as a key limitation and emphasize that future studies should aim to validate the model on independent cohorts when such datasets become available. Furthermore, although our model achieved an AUC of 0.74, this performance is not yet sufficient for clinical translation. Additional efforts to refine the model and test it in larger, independent cohorts will be essential before its use in clinical practice can be considered. Interestingly, a recent approach using multiplex IF on DCIS cohort reached the same AUC(2). While both our study and the Risom et al. paper aim to leverage spatial relationships to predict DCIS progression, we would like to emphasize that the two approaches are fundamentally

different in terms of the markers used. Risom et al. focus on a broad panel of markers, including those
related to the stroma, immune cells, and tumor cells, which provide a comprehensive view of the tumor
microenvironment. In contrast, our approach centers on eco-evolutionary markers, specifically CA9
and LAMP2b, which are associated with hypoxia and tumor acidity and their spatial distribution,
respectively, to capture distinct microenvironmental pressures and adaptive cancer cell responses.
These differences reflect divergent hypotheses about the key drivers of DCIS progression. The fact that
both studies report a similar AUC of 0.74, with the distinct marker sets and biological processes,
suggests that our findings offer complementary insights into DCIS progression and combination of
approaches might actually increase the accuracy.
Our study demonstrates the utility of eco-evolutionary principles in understanding DCIS progression.
In our study, we proposed that specific tumor microenvironmental conditions, such as hypoxia and
acidosis, are associated with phenotypic changes that may indicate DCIS progression. However,
although we have shown previously that these microenvironments can cause aggressive phenotypes, we
acknowledge that our findings here do not conclusively demonstrate that these environmental factors
are causative agents in the transition from DCIS to IDC. Instead, our data suggest that these conditions
could serve as biomarkers for identifying lesions that are more likely to be upstaged. However, the
ability to define more refined cell phenotypes within each region of interest (ROI) could further
enhance our analysis. If we can identify and characterize more detailed phenotypes, it would allow us
to extract additional features that describe the spatial interactions of these phenotypes. This, in turn,
could potentially improve the classifier's performance and make the results more interpretable. By
capturing the intricate interactions between various cell types and their microenvironments, we could
gain deeper insights into the ecological dynamics driving DCIS progression and improve predictive
models for patient outcomes.
In recent years, there has been a growing trend towards adopting a "watchful waiting" approach for
certain cases of DCIS, rather than immediate surgical excision(31,32). This strategy aims to reduce
overtreatment by closely monitoring DCIS lesions that may not progress to invasive cancer. In this
context, our upstaging predictions become particularly relevant. Identifying microenvironmental and
phenotypic factors that indicate a higher likelihood of progression to IDC could help clinicians make
more informed decisions about when to intervene and when to adopt a more conservative, observational
approach. The ability to predict which DCIS cases are at higher risk of progressing to invasive disease
would provide critical information for optimizing patient management, minimizing unnecessary
treatments, and reducing the psychological and physical burdens associated with overtreatment(33).
Further validation of these predictive models could therefore have important implications for guiding
treatment strategies in the context of DCIS.

Method:

Method overview

Our evolutionary analysis pipeline takes 3 consecutive slides of each patient sample, detects intra-
ductal cell niches, characterizes these niches with their spatial and morphological features, and then
predicts whether the patient will be pure DCIS or upstaged based on the distribution of these niches. In

particular, the pipeline has 4 modules. First, we annotate and align ducts from different whole slide
images (WSIs) of the same patient sample. This ensures cells of different slides are aligned and we can
characterize their interactions. In the second module, we detect and map all eco-evo positive cells (i.e.,
cells activated with the selected stains) into the same duct and detect different clusters of cells as
niches. In the third module, we characterize these niches with comprehensive spatial statistical features,
as well as their morphological features as observed in HE. Finally, we categorize these niches into
different subclasses through deep learning-based dimension reduction and clustering based on their
features. We use the distribution of different niche subclasses to characterize different samples/patients.
We demonstrate the discriminative power of this niche-based characterization in predicting whether a
patient will be pure DCIS or upstaged in the future. **Figure 1C** illustrates the overview of our pipeline.
377

Data preparation and usage

The data used in this study is the biopsy samples collected after mammography and before surgery. 84
samples including 68 pure DCIS plus 16 upstaged were analyzed. For each sample, we obtained 3
whole slide images, including 1 HE and 2 IHC slides. We conduct 5-fold stratified cross validation,
where 4 folds are used for niche clustering and for the training of the pure DCIS/upstaged classifier and
1 fold is used for validation. This fits the clinical application we are aiming for; we would like our
model to estimate the risk based on biopsy samples, which are much less invasive and can be used for
patient stratifications before surgery and hopefully decrease over treatment. Further details on HE and
IHC acquisition are provided below.
387

Sample selection, immunohistochemistry and HE staining. Patients' tumor blocks were selected by
pathologists using the archived HE stained slides. The blocks were sequentially sectioned 4 μms and
de-identified for research use. 3 slides were used to be stained with primary antibodies of 1:100 dilution
of anti-LAMP2 (#ab18529, Abcam), and 1 ug/ml concentration of anti-CA9 (#AF2188, R&D), and HE
staining using standard hematoxylin and eosin protocol. Positive and negative controls were used.
Normal placenta was used as a positive control for LAMP2 and clear cell renal cell carcinoma was used
as a positive control for CA9. For the negative control, an adjacent section of the same tissue was
stained without application of primary antibody and any stain pattern observed was considered as non-
specific binding of the secondary. Primary immunohistochemical analysis was conducted using
digitally scanning slides. The scoring method used by the pathologist reviewer to determine (a) the
degree of positivity scored the positivity of each sample ranged from 0 to 3 and was derived from the
product of staining intensity (0–3+). A zero score was considered negative, score 1 was weak positive,
score 2 was moderate positive, and score 3 was strong positive. (b) The percentage of positive tumors
stained (on a scale of 0-3).
402

Whole slide imaging (WSI) of IHC and HE slides were obtained by scanning at 20X magnification (of
0.5022 micrometer per pixel) using Aperio AT2 from Leica Biosystems. Images were transferred to
cloud storage and also locally to be uploaded in QuPath software for analysis. QuPath software was
used to detect the positive pixels for each IHC marker (CA9 and LAMP2b) and to segment the HE
images into hypoxic and normoxic tumor habitats based on their distance from the basement
membrane.
409

MODULE 1: Duct annotation and alignment

We annotate and align ducts within all input slides (1 HE + 2 IHCs per sample). For the Bx cohort, there are a small number of ducts per slide. They were annotated by 1 pathologist and trained students. After annotating ducts, we align the ducts from the three modalities via co-registration. This alignment enables us to map cells into the same spatial domain and analyze their interaction. Details are provided below.

Manual annotation of aucts in the Bx cohort.

QuPath was used as the interface to annotate ducts by the pathologist (Dr. Bai) and the trained students. We annotate ducts from WSIs of all three modalities. To ensure best characterization, we only identify ducts of >400 μms diameter, with visible myoepithelial layer and basement membrane. Following this, based on distance, each duct was annotated with four layers: adjacent stroma, oxidative/normoxia, hypoxic/hypoxia, and necrosis. Adjacent stroma was defined as the stroma up to 125 μms outside a given duct. Within the duct, necrosis was defined as any area containing dead cells, as identified by a lack of nuclei. Oxidative layer was defined as the area containing cells inside the duct within 125 μms of the basement membrane. Hypoxia was defined as the area containing cells inside the duct further than 125 μms from the basement membrane. The annotations were done for all 84 samples in the Bx cohort, and then were exported as standard GeoJSON files.

Co-registration.

To characterize the interactions of different modalities from single-plexed slides, an alignment strategy was utilized. We register both CA9 and LAMP2b IHC slides towards the HE slides. A direct co-registering at the whole slide level with manual landmarks does not give us satisfactory alignment at each duct, due to the variable deformations across slides. We further co-register the slides in a duct-by-duct fashion. Using initially registered whole slides, and spatial proximity, we identify the corresponding ducts at the HE and 2 IHC slides. Next, we register both the CA9 duct and LAMP2b duct into the corresponding HE duct. We use Virtual Alignment of pathology Image Series (VALIS), which provides a fully automated pipeline to register whole slide images (WSI) using rigid and/or non-rigid transformations (34). For each sample, we chose non-rigid registration and registered the ducts from CA9 and LAMP2b towards the reference HE duct. The co-registration procedure and the qualitative results are shown in **Figure S3 and S4**. The co-registration provides a mapping of any cells detected in CA9 or LAMP2b towards a shared spatial domain, enabling the analysis of their interactions.

MODULE 2: Cell and niche detection

Cell detection. With the duct annotations in place, we automatically detect cells from the 2 IHCs and determine if they are positive in CA9 or LAMP2b based on their intensities. As we are only interested in intra-ductal cell niches, we only detect cells within each duct. For each IHC duct, we detect cells using Qupath watershed cell detection algorithm(25). Based on the intensity level, we categorize the cells into 4 groups: 'Negative', '1+', '2+', and '3+'. The detection of cells within an HE duct is done by starDist (25,35) extension on Qupath.

Graph construction for niche detection. After all eco-evo positive cells (i.e., CA9 or LAMP2b positive cells) were annotated and mapped on HE ducts, we were able to perform analysis on an intra-duct level. Since there are still a large amount of eco-evo positive cells within each duct, with diverse spatial context and morphological features, we construct a graph with these cells by connecting cells whose distances are smaller than a certain threshold and detect connected components of the graph as "niches". Multiple thresholds have been experimented and an optimum value is selected based on performance. Each eco-evo positive cell niche is supposed to have a similar eco-evo phenotype and be spatially coherent. We overlay both CA9 positive and LAMP2b positive cells into the same domain as an approximation of the local eco-evo cell distribution, because the two IHCs are consecutive sections from the same tissue block (**Figure S5**). This gives us the opportunity to measure their interaction via spatial statistical functions as defined later. Based on the same principle, we use cell morphological features extracted from HE within the region of each niche to characterize the niche.

MODULE 3: niche characterization and feature extraction

Once niches are detected. We extracted both spatial and morphological features to characterize them. To describe the spatial interaction patterns, we utilized various spatial functions as features. We also extract cell features consisting of morphology features and texture features that are commonly adopted in HE image analysis.

Cellular features. The morphological features include area, eccentricities, circularity, elongation, extent, major axis length, minor axis length, solidity and curvature, the texture features include angular second moment (ASM) of co-occurrence matrix, contrast, correlation, entropy, homogeneity and intensity .All of these features were calculated following the implementations in the sc-MTOP(36) package.

Although we do not have exact cell-to-cell correspondence between the cells within a niche and cells detected in HE, we still can aggregate morphological and texture features within the proxy of the eco-evo cells of a niche to characterize the niche. For each niche, we identify the concave hull region enclosing its eco-evo positive cells within the HE duct. Next, we aggregate cell features across all HE-detected cells within the corresponding region. For each cell feature dimension, we calculated its mean, standard deviation, maximum, minimum, kurtosis and skewness.

Spatial features. We extract various spatial statistical functions(37) to characterize eco-evo positive cells and their interactions. These functions are listed below.

G Function: The G function, denoted as G(r), is the cumulative distribution function of nearest-neighbor distance. The G function provides insights into the clustering or dispersion behavior of the point pattern.

$$G(r) \ = \ P\{d(u, X \backslash u) \ \leq \ r \,|\, u \ \in \ X\}, d(\bullet) \ is \ the \ minimum \ distance$$

F Function: The F function, known as the empty space function, is the cumulative distribution function of the empty-space distance. The F function is commonly used to assess the regularity or inhibition patterns in point patterns.

$$F(r) \ = \ P\{d(u, X) \ \leq \ r\}, d(\bullet) \ is \ the \ minimum \ distance$$

491 K Function: Ripley's K function, denoted as K(r), is a measure of second-order intensity or spatial
492 interaction. It assesses whether points tend to be more clustered or dispersed within a certain distance r
493 compared to a CSR process. It considers both the distance and intensity of points to capture the
494 clustering behavior of the point pattern.

$$K(r) \ = \ \frac{|W|}{n(n-1)} \sum_{i=1}^{n} \sum_{j=1, j \neq i}^{n} \mathbf{1}\{d_{ij} \leq r\} e_{ij}(r), e_{ij}(\bullet) \; is \; the \; edge \; correction \; weight$$

495 L Function: L function is a variance stabled version of K function.

$$L(r) \ = \ \frac{\sqrt{K(r)}}{r}$$

We calculated G, F, and L functions in both univariate and multivariate fashions. For each of the
functions, the distances between source cell and the target cells are considered. Univariate spatial
functions sample source cells and target cells from the same type of cells while multivariate
counterparts' sample from different types of cells. Univariate G,F,L are calculated for the single-marker
cell subsets, and multivariate G_cross, L_cross for different subsets such as CA9-LAMP2b. 'Gest'
function and 'Fest' function from 'spatstat' R package were used with Kaplan-Meier estimator(38), and
'Lest' function was used with isotropic correction(39,40).
503

MODULE 4: Diagnostic risk estimation with pattern proportion

In the last module, we train a classifier using these niches to predict whether a patient will be
"upstaged" or "pure DCIS". This establishes the diagnostic power of these niches. A direct aggregation
of niche information within each sample/patient is not sufficient. Tumor microenvironment is
heterogeneous, and niches demonstrate diverse spatial and morphological behavior. To account for the
diversity, we will focus on how different niches are distributed across a sample. We show that the
distributions of different niches essentially characterize the tumor ecology in a much more refined
manner compared with previous distance-based definitions of hypoxia/oxidative layers.
One technical challenge is that the niche features computed in the previous module are high
dimensional and the niche features are diversely distributed. We propose to first find a simplified
distributional description of the niches, and then use the simplified description for prediction. First, we
cluster the niches into different sub-classes based on their features. The clustering is carried out using
K-means clustering with a tunable parameter k. Once the niche sub-classes are determined. We use
their distribution on each sample to predict its upstage/pure DCIS status. The prediction power of the
classifier sheds light on the diagnostic power of the niches and their spatial and cellular features.
To understand the contribution of each feature to the prediction model, we employed SHAP (SHapley
Additive exPlanations) analysis. SHAP is a unified approach to interpreting machine learning models
by assigning each feature an importance value for a particular prediction. In our study, SHAP values
were computed for the features representing the proportions of different patterns within the niches. By
calculating the SHAP values, we could determine the impact of each feature on the model's output,
thereby identifying the most influential patterns that contribute to predicting DCIS upstaging. This step
is crucial for ensuring the transparency and interpretability of the machine learning model.
Furthermore, we select features that are highly relevant to the sub-classes using different approaches
including covariance, mutual information scoring and maximum relevance minimum redundancy

(mRMR)(41) and choose the features identified by both approaches. **Figure 4C** shows the gradient map
of each of these features on niches in the latent space.
530
Niche distribution for diagnosis. After assigning each duct to its sub-class, we aggregate across all
niches of each sample and use its sub-class distribution to characterize this sample. Assuming k niche
sub-classes, each sample has a k dimensional histogram to describe its niche sub-class distribution. We
call this the niche distributional (Nbd-Dist) feature. We trained a classifier to predict whether a sample
is pure DCIS or upstage. Repeating the iteration 10 times and comparing the mean area under curve
(AUC) on the test set. The classifier types experimented include lightGBM, soft vector machine
(SVM), logistic regression and random forest, and the random forest classifier yields the best
performance.
539

540 **Resource availability**
541 **Lead contact**
542 Further information and any related requests should be directed to and will be fulfilled by the lead
543 contact Mehdi Damaghi (Mehdi.Damaghi@stonybrookmedicine.edu). All the staining and annotations
544 are deposited in the physical sciences in oncology network (PSON).
545

546 **Materials availability**
547 This study did not generate new unique reagents.
548

Acknowledgements

We gratefully acknowledge funding from Physical Sciences Oncology Network at the National Cancer
Institute (grant U01CA261841), R01 grant R01CA272601, and R01CA249016. This work has been
supported also in part by the Analytic Microscopy Core Facility at the H. Lee Moffitt Cancer Center &
Research Institute, an NCI designated Comprehensive Cancer Center (P30-CA076292). We thank Dr.
Liliana Davolos and Dr. David Ray for their guidelines on ecological and evolutionary dynamics
principles used in the manuscript, as well as for critical reading and providing comments for the
manuscript.

Author contributions

M.D. conceptualized and designed the research; Y.X., M.A., J.D.B., A.C., Y.C., M.D., performed the
experiment and analysis; J.D.B. reviewed all the slides and scored them as the project pathologist; P.P.,
C.C., and M.D. contributed to results interpretation; and M.D. wrote the paper. All authors revised the
paper.
563

564 **Declaration of interests**
565 The authors declare no competing interest.
566

Figure Captions
Figure 1. Ecological and evolutionary designed biomarkers of DCIS upstaging. A) Model of
microenvironment-driven evolution of breast cancer from normal breast tissue to DCIS and IDC: Our schematic

is overlaid on HE staining of breast cancer specimens at different stages of DCIS and IDC. Different patients
may experience various types of evolutionary trajectory following different evolutionary models, including linear
and branched progression from DCIS to IDC shown here. **B)** The patient cohort was curated from retrospective
DCIS samples, with two sample collections at biopsy and excision. The main criterion was the diagnosis of
DCIS at the biopsy stage. **C)** Eco-evolutionary designed- machine learning assisted pipeline to define cancer cell
niches inside oxygen habitats in DCIS. *i)* Data preprocessing steps including duct annotation, cell detection and
classification for HE and IHC slides, followed by co-registration to map IHC-identified cells onto the HE slide.
ii) The analysis is carried out at multiple scales, namely duct, habitat and niche, from the largest to smallest. At
each scale the nucleus morphology texture feature and spatial features are extracted. *iii)* The pattern differential
analysis approach where the patterns are firstly identified and then the proportions of such patterns are used as
features to predict the upstaging status of a patient.
581
Figure 2. **Eco-evolutionarily designed biomarker discovery to predict upstaging in DCIS. A)**
Illustration of normoxic, hypoxic and necrotic habitats in a duct. **B)** Illustration of annotation and
scoring on 2 IHCs and how cells are scored in each habitat. **C)** and **D)** Dot plots of counts of CA9
expression in each habitat per duct. Cells are scored 0 for 'negative' or '1+','2+','3+' for positive cells
based on their intensity. Scoring was performed and analyzed separately for normoxic (oxidative)
habitat (C) or hypoxic habitat (D). In the dot plot, each dot is a single duct. The color of dots reflects
their score as follows: Blue = 0, yellow ='1+', orange ='2+', and red = '3+'. The number of dots
reflects how many ducts were detected in each patient's biopsy with size bigger than 400 um in
diameter. The distribution in hypoxic habitat is significantly different between pure DCIS and upstaged
groups in hypoxic habitats and not in oxygenated habitat. Data was analyzed using the Wilcoxon
signed-rank test. The same graph is created for LAMP2b (supplementary fig. 2).
593
Figure 3. Niches are defined inside habitats from the hypoxia and acidosis markers expression. A)
One sample duct from CA9 slide. Top: The original IHC slide. Middle: Cell detection and intensity-based
classification using Qupath overlaid on the slide. Bottom: the graph constructed from the CA9 positive cells and
the connected components of the graph (Niches) highlighted in different colors. **B)** The HE staining of the same
duct as A. Top: The original HE slide. Middle: Duct annotation overlaid on the HE slide. Bottom: Co-registered
CA9-positive niches mapped and overlaid on HE slides as mIHC to be able to extract HE features from CA9
positive niches. Note the orientation of HE and CA9 slide was opposite, and our co-registration technique
successfully created a mIHC of the ducts with similar coordinates. The same approach was used for LAMP2b
and the combination.
603
Figure 4. Post Analysis reveals the top contributing patterns and features. A) UMAP of the
features of the niches, different colors represent different clusters(patterns) **B)** Top: The impact of each
pattern on the classifying result, blue and red colors represent impact on pure DCIS and progressed
predictions respectively, the proportion of pattern 5 has the greatest impact for both categories. Bottom:
Using correlation, MI, and MRMR to obtain the most contributing features in the pattern 5 clustering
phase, identifying a common feature set that includes 4 features: Area_min, Perimeter_min,
AreaBbox_min, and F_0<=r<10. **C)** UMAP showing the value of the 4 identified features for different
samples, and it can be seen that samples in the pattern 5 tend to have higher values in Area_min,
Perimeter_min, AreaBbox_min and low values for F_0<=r<10. **D)** A niche belonging to pattern 5, it
contains no small size cells and exhibits a relatively dispersed distribution.

References:

1. Greaves M, Maley CC. Clonal evolution in cancer. Nature. 2012;481:306–13.

2. Risom T, Glass DR, Averbukh I, Liu CC, Baranski A, Kagel A, et al. Transition to invasive breast cancer is associated with progressive changes in the structure and composition of tumor stroma. Cell. 2022;185:299–310.e18.

3. Maley CC, Aktipis A, Graham TA, Sottoriva A, Boddy AM, Janiszewska M, et al. Classifying the evolutionary and ecological features of neoplasms. Nat Rev Cancer. 2017;17:605–19.

4. Boutry J, Tissot S, Ujvari B, Capp J-P, Giraudeau M, Nedelcu AM, et al. The evolution and ecology of benign tumors. Biochim Biophys Acta Rev Cancer. 2022;1877:188643.

5. Amend SR, Pienta K. Abstract 2884: Tumor-driven eutrophication of the tumor ecosystem selects for cancer cell clones that overcome evolutionary inertia leading to increased metastatic capacity. Cancer Res. American Association for Cancer Research; 2015;75:2884–2884.

6. Damaghi M, Mori H, Byrne S, Xu L, Chen T, Johnson J, et al. Collagen production and niche engineering: A novel strategy for cancer cells to survive acidosis in DCIS and evolve. Evol Appl. 2020;13:2689–703.

7. Lipinski KA, Barber LJ, Davies MN, Ashenden M, Sottoriva A, Gerlinger M. Cancer Evolution and the Limits of Predictability in Precision Cancer Medicine. Trends Cancer Res. 2016;2:49–63.

8. Giaquinto AN, Sung H, Miller KD, Kramer JL, Newman LA, Minihan A, et al. Breast Cancer Statistics, 2022. CA Cancer J Clin. 2022;72:524–41.

9. Strand SH, Rivero-Gutiérrez B, Houlahan KE, Seoane JA, King LM, Risom T, et al. Molecular classification and biomarkers of clinical outcome in breast ductal carcinoma in situ: Analysis of TBCRC 038 and RAHBT cohorts. Cancer Cell. 2022;40:1521–36.e7.

10. Lehman CD, Arao RF, Sprague BL, Lee JM, Buist DSM, Kerlikowske K, et al. National Performance Benchmarks for Modern Screening Digital Mammography: Update from the Breast Cancer Surveillance Consortium. Radiology. 2017;283:49–58.

11. Lips EH, Kumar T, Megalios A, Visser LL, Sheinman M, Fortunato A, et al. Genomic analysis defines clonal relationships of ductal carcinoma in situ and recurrent invasive breast cancer. Nat Genet. 2022;54:850–60.

12. Sarhadi S, Salehzadeh-Yazdi A, Damaghi M, Zarghami N, Wolkenhauer O, Hosseini H. Omics Integration Analyses Reveal the Early Evolution of Malignancy in Breast Cancer. Cancers [Internet]. 2020;12. Available from: http://dx.doi.org/10.3390/cancers12061460

13. Heselmeyer-Haddad K, Berroa Garcia LY, Bradley A, Ortiz-Melendez C, Lee W-J, Christensen R, et al. Single-cell genetic analysis of ductal carcinoma in situ and invasive

breast cancer reveals enormous tumor heterogeneity yet conserved genomic imbalances
and gain of MYC during progression. Am J Pathol. 2012;181:1807–22.

14.   Hanna WM, Parra-Herran C, Lu F-I, Slodkowska E, Rakovitch E, Nofech-Mozes S. Ductal
carcinoma in situ of the breast: an update for the pathologist in the era of individualized
risk assessment and tailored therapies. Mod Pathol. 2019;32:896–915.

15.   Carmeliet P, Jain RK. Principles and mechanisms of vessel normalization for cancer and
other angiogenic diseases. Nat Rev Drug Discov. 2011;10:417–27.

16.   Wu B, Liu D-A, Guan L, Myint PK, Chin L, Dang H, et al. Stiff matrix induces exosome
secretion to promote tumour growth. Nat Cell Biol. 2023;25:415–24.

17.   Persi E, Duran-Frigola M, Damaghi M, Roush WR, Aloy P, Cleveland JL, et al. Systems
analysis of intracellular pH vulnerabilities for cancer therapy. Nat Commun. 2018;9:2997.

18.   Damaghi M, West J, Robertson-Tessi M, Xu L, Ferrall-Fairbanks MC, Stewart PA, et al.
The harsh microenvironment in early breast cancer selects for a Warburg phenotype. Proc
Natl Acad Sci U S A [Internet]. 2021;118. Available from:
http://dx.doi.org/10.1073/pnas.2011342118

19.   Lobo RC, Hubbard NE, Damonte P, Mori H, Pénzváltó Z, Pham C, et al. Glucose Uptake
and Intracellular pH in a Mouse Model of Ductal Carcinoma In situ (DCIS) Suggests
Metabolic Heterogeneity. Front Cell Dev Biol. 2016;4:93.

20.   Damaghi M, Tafreshi NK, Lloyd MC, Sprung R, Estrella V, Wojtkowiak JW, et al.
Chronic acidosis in the tumour microenvironment selects for overexpression of LAMP2 in
the plasma membrane. Nat Commun. 2015;6:8752.

21.   Ordway B, Swietach P, Gillies RJ, Damaghi M. Causes and Consequences of Variable
Tumor Cell Metabolism on Heritable Modifications and Tumor Evolution. Front Oncol.
2020;10:373.

22.   Gillies RJ, Verduzco D, Gatenby RA. Evolutionary dynamics of carcinogenesis and why
targeted therapy does not work. Nat Rev Cancer. 2012;12:487–93.

23.   Ibrahim-Hashim A, Robertson-Tessi M, Enriquez-Navas PM, Damaghi M, Balagurunathan
Y, Wojtkowiak JW, et al. Defining Cancer Subpopulations by Adaptive Strategies Rather
Than Molecular Properties Provides Novel Insights into Intratumoral Evolution. Cancer
Res. 2017;77:2242–54.

24.   Damaghi M, Gillies R. Phenotypic changes of acid-adapted cancer cells push them toward
aggressiveness in their evolution in the tumor microenvironment. Cell Cycle.
2017;16:1739–43.

25.   Bankhead P, Loughrey MB, Fernández JA, Dombrowski Y, McArt DG, Dunne PD, et al.
QuPath: Open source software for digital pathology image analysis. Sci Rep.
2017;7:16878.

26.   Freischel AR, Damaghi M, Cunningham JJ, Ibrahim-Hashim A, Gillies RJ, Gatenby RA, et
al. Frequency-dependent interactions determine outcome of competition between two

breast cancer cell lines. Sci Rep. 2021;11:4908.

27. Jardim-Perassi BV, Huang S, Dominguez-Viqueira W, Poleszczuk J, Budzevich MM,
Abdalah MA, et al. Multiparametric MRI and Coregistered Histology Identify Tumor
Habitats in Breast Cancer Mouse Models. Cancer Res. 2019;79:3952–64.

28. Sobhani F, Muralidhar S, Hamidinekoo A, Hall AH, King LM, Marks JR, et al. Spatial
interplay of tissue hypoxia and T-cell regulation in ductal carcinoma in situ. NPJ Breast
Cancer. 2022;8:105.

29. Compton ZT, Mallo D, Maley CC. Stronger together: Cancer clones cooperate to alleviate
growth barriers in critical cancer progression transitions. Cancer Res. American
Association for Cancer Research (AACR); 2023;83:4013–4.

30. Chafe SC, McDonald PC, Saberi S, Nemirovsky O, Venkateswaran G, Burugu S, et al.
Targeting hypoxia-induced carbonic anhydrase IX enhances immune-checkpoint blockade
locally and systemically. Cancer Immunol Res. American Association for Cancer
Research; 2019;7:1064–78.

31. Ryser MD, Worni M, Turner EL, Marks JR, Durrett R, Hwang ES. Outcomes of active
surveillance for ductal carcinoma in situ: A computational risk analysis. J Natl Cancer Inst.
Oxford University Press (OUP); 2016;108:djv372.

32. Glencer AC, Miller PN, Greenwood H, Maldonado Rodas CK, Freimanis R, Basu A, et al.
Identifying good candidates for active surveillance of ductal carcinoma in situ: Insights
from a large neoadjuvant endocrine therapy cohort. Cancer Res Commun. American
Association for Cancer Research (AACR); 2022;2:1579–89.

33. Fortunato A, Mallo D, Cisneros L, King LM, Khan A, Curtis C, et al. Evolutionary
Measures Show that Recurrence of DCIS is Distinct from Progression to Breast Cancer.
medRxiv [Internet]. 2024; Available from: http://dx.doi.org/10.1101/2024.08.15.24311949

34. Gatenbee CD, Baker A-M, Prabhakaran S, Swinyard O, Slebos RJC, Mandal G, et al.
Virtual alignment of pathology image series for multi-gigapixel whole slide images. Nat
Commun. 2023;14:4502.

35. Schmidt U, Weigert M, Broaddus C, Myers G. Cell Detection with Star-Convex Polygons.
Medical Image Computing and Computer Assisted Intervention – MICCAI 2018. Springer
International Publishing; 2018. page 265–73.

36. Zhao S, Chen D-P, Fu T, Yang J-C, Ma D, Zhu X-Z, et al. Single-cell morphological and
topological atlas reveals the ecosystem diversity of human breast cancer. Nat Commun.
2023;14:6796.

37. Baddeley A, Rubak E, Turner R. Spatial Point Patterns: Methodology and Applications
with R. CRC Press; 2015.

38. Baddeley A, Gill R. Kaplan-Meier estimators of interpoint distance distributions for spatial
point processes. IEEE Trans Inf Theory [Internet]. 1993; Available from:
https://ir.cwi.nl/pub/5247/05247D.pdf

39. Ohser J. On estimators for the reduced second moment measure of point processes. Series
Statistics. Taylor & Francis; 1983;14:63–71.

40. Kendall WS. Stochastic Geometry: Likelihood and Computation. Routledge; 2019.

41. Peng H, Long F, Ding C. Feature selection based on mutual information: criteria of max-
dependency, max-relevance, and min-redundancy. IEEE Trans Pattern Anal Mach Intell.
2005;27:1226–38.

48.   Lundberg, S. M. & Lee, S.-I. A unified approach to interpreting model predictions. *Proc.*

Adv. Neural Inf. Process. Syst. 30 (2017).

Non-obligatory stages of breast cancer

Basement membrane
Luminal cells
Myoepithelial cells

Normal duct | Hyperplasia | Atypical Hyperplasia | DCIS | IDC

Human breast

Biopsy — Surgery

Ductal Carcinoma In Situ (DCIS)

<6 Month

DCIS at Bx — DCIS at Ex

Upstaged

Biopsy — Surgery

Ductal Carcinoma In Situ (DCIS) — Invasive Ductal Carcinoma (IDC)

<6 Month

DCIS at Bx — IDC at Ex

CA9
LAMP2

CA9
LAMP2b

Features

Morphology & Texture

Minor axis
Major axis

Pattern differential analysis