1

Full title: Between-day reliability of local and global muscle-tendon unit assessments in

female athletes whilst controlling for menstrual cycle phase.

Short title: Reliability of muscle-tendon unit assessments in female athletes

Scott Newbould[1*], Josh Walker[1], Alexander J Dinsdale[1], Sarah Whitehead[1,2], Gareth

Nicholson[1]

[1]Carnegie School of Sport, Leeds Beckett University, Leeds, United Kingdom

[2]Leeds Rhinos Netball, Leeds, United Kingdom

* Corresponding author

E-mail: s.newbould@leedsbeckett.ac.uk (SN)

Abstract

Measurements of muscle-tendon unit (MTU) function can be categorised into local (e.g. tendon strain) or global (e.g. jump height) assessments. Although menstrual cycle phase may be a key consideration when implementing these assessments in female athletes, the reliability of many MTU assessments is not well defined within female populations. Therefore, the purpose of this study was to report the test-retest reliability of local and global MTU function assessments during the early follicular phase of the menstrual cycle. Seventeen naturally menstruating females (age 28.5 ± 7.3 years) completed local and global assessments of MTU function during two testing sessions separated over 24-72 hours. Local tests included Achilles' tendon mechanical testing and isometric strength of ankle plantar flexors and knee extensors, whereas global tests included countermovement, squat, and drop jumps, and the isometric midthigh pull. Based on intraclass correlation coefficient (ICC) statistics, *poor* to *excellent* reliability was found for local measures (ICC: 0.096-0.936). *Good* to *excellent* reliability was found for all global measures (ICC: 0.788-0.985), excluding the eccentric utilisation ratio (ICC 0.738) and most rate of force development metrics (ICC: 0.635-0.912). Isometric midthigh pull peak force displayed *excellent* reliability (ICC: 0.966), whereas force-time metrics ranged from *moderate* to *excellent* (ICC: 0.635-0.970). Excluding rate of force development (coefficient of variation [CV]: 10.6-35.9%), global measures (CV: 1.6-12.9%) were more reproducible than local measures (CV: 3.6-64.5%). However, local metrics directly measure specific aspects of MTU function, and therefore provide valuable information despite lower reproducibility. The novel data reported here provides insight into the natural variability of MTU function within female athletes, which can be used to enhance the interpretation of other female athlete data, especially that which aims to investigate other aspects of variability, such as the menstrual cycle.

Introduction

Measures of muscle-tendon unit (MTU) function are of interest in sport science because of their relationships with athletic performance tasks [1–3]. Sport scientists frequently use local and global assessments of MTU function to gain insight into these characteristics and to inform training practices. Local assessments are those that directly measure aspects of MTU function (e.g., tendon mechanical properties [i.e., strain or stiffness]), whilst global tests focus on movement outcomes, such as jump height or reactive strength. Global measures are underpinned by the aspects of MTU function measured in the local assessments, for example, rate of force development (RFD) and jump height are influenced by tendon stiffness [4]. Thus, gaining further insight into the interrelationships between local measures of MTU function and global outcomes can help inform training priorities. Reliably assessing both local and global MTU function is therefore of great importance to sport scientists. However, much of the sport science literature does not reflect the increasing participation in female sport seen in recent years [5] and there is currently limited data on the sources of variability in MTU function in females.

The scarcity of female-specific reliability data is a concern because differences in the hormonal environment between males and females might result in different magnitudes of biological variation within and between days [6]. Additionally, it must be considered that females present with comparatively higher and more variable levels of hormones such as oestrogen, which can impact both neuromuscular output [7] and tendon function [8]. However, the true effect of changes in the hormonal environment on MTU function remains unknown [9]. Until these effects are better understood, reliability data should not be directly transferred between sexes, and when collecting female-specific data, consideration should be made for the hormonal environment at the time of testing. Understanding the natural variability of MTU function without the confounding effects of hormonal changes would allow more objective inferences to be made regarding MTU testing and

monitoring data in female athletes, especially when comparing different hormonal profiles (e.g.,

different phases of the menstrual cycle).

Regarding local assessments of MTU function, the triceps surae muscle group and the Achilles'

tendon (AT) comprise one of the most profiled muscle-tendon complexes in athletes. This is because

of the role that the AT has in overall MTU strategy [10,11], and the relationships that exist between

AT mechanical properties, such as elongation and stiffness, and performance in sprint [12], jump

[13], and long-distance running tasks [14]. However, if this field of research is to progress to

individualised monitoring and prescription of MTU function [15], it is important to first understand

the reliability and sensitivity to change of AT metrics. To this end, several studies have reported test-

retest reliability data, with mixed findings for both maximal AT elongation (intraclass correlation

coefficients [ICC] from 0.56-0.95, and coefficient of variations [CV] from 4.1-12.3% [16–20] and AT

stiffness (ICC: 0.64-0.98 and CV: 4.7-13.5% [13,16–18,20–23]). However, to the authors' knowledge,

only one study has included female participants (7 males and 4 females in Hansen et al. [16]), and

therefore the reliability of these measures within female populations is not well defined.

For global tests of MTU function, force plate analyses of the countermovement jump (CMJ), squat

jump (SJ), drop jump (DJ), and the isometric mid-thigh pull (IMTP) are frequently used to assess and

monitor athletes [e.g., 24,25]. Although recommendations have been made for data collection and

analysis procedures for these tests [27–29], the interpretation of these metrics is limited without

coinciding reliability data. The reliability of both basic (e.g., jump height, peak force) and more

complex CMJ metrics (e.g., RFD) have been investigated within female populations in studies by Moir

et al. [30] and Keogh et al. [26], with both studies reporting high reliability for nearly all metrics (ICC:

0.80-0.97). However, reliability data has scarcely been reported in adult females for the squat jump,

drop jump, or isometric midthigh pull, reducing the ability to interpret these tests in this population.

Therefore, research which provides reliability data for a variety of global MTU function assessments

in a female population will provide novel and valuable information for researchers and practitioners.

Many studies have aimed to measure the variability in performance over the course of the

menstrual cycle [9], although true understanding of menstrual cycle variability cannot be complete

without first understanding the natural variability that exists within this population. Therefore, the

assessment of variability without potentially confounding changes in the hormonal environment will

provide meaningful context in which both past and future female athlete research can be

interpreted. Thus, the aim of this study was to report the test-retest reliability of local and global

tests of MTU function in naturally menstruating female athletes, whilst controlling for menstrual

cycle phase.

Materials and methods

Study design

104   This study used a within-subject repeated measures design, whereby participants completed two

105   testing sessions separated over 24-72 hours during the early follicular phase of the menstrual cycle

106   (days 1-5). The early follicular phase was chosen because it is the phase with the most stable

107   hormonal profile [31], and the timing of this phase was predicted by calendar-based counting using

108   previous cycle length data and confirmed by the presence of menstrual bleeding [32].

Participants

Seventeen naturally menstruating females volunteered to participate in this study (age 28.5 ± 7.3

years, stature 166.8 ± 5.6 cm, body mass 62.6 ± 8.2 kg). 'Naturally menstruating' was self-reported

and was defined as having cycle lengths between 21-35 days and having had nine or more periods in

the previous year [32]. Participants had an average training experience of 8.0 ± 5.9 years and

performed 4.8 ± 2.6 training sessions per week, and therefore aligned with Tier 2 of the Participant

Classification Framework [33]. Additionally, participants were required to be free from lower-body

injuries for six months prior to participation, and to not have a previous AT or patellar tendon

rupture. Participant recruitment took place between 6/10/22 and 20/10/23, and all participants

provided their written informed consent to participate. The study was approved by the Leeds

Beckett Ethics Committee (approval reference 102193) and conformed to the Declaration of Helsinki

[34].

Procedures

All participants completed a familiarisation session approximately 7 days prior to the first testing

visit, which included submaximal and maximal attempts of each test. For the testing sessions,

participants first completed a standardised warm up before completing all tests in the following

order: CMJ, SJ, DJ, isometric plantar flexion (including measurements of AT mechanical function),

isometric knee extension, and the IMTP. At least 2 minutes rest was provided between each test.

Definitions and calculations for all metrics are provided in S1 and S2 Tables.

Mechanical properties of the muscle-tendon unit

All measurements of the triceps surae MTU were performed on the left leg to quasi-randomise for

limb dominance. First, resting AT length was measured using the techniques of Barfod et al. [35].

Participants were then seated on an isokinetic dynamometer (CSMI, Cybex Humac Norm, Stoughton,

MA, USA), sampling at a frequency of 1,000 Hz, with a hip joint angle of 120° (180° in anatomical

position), knee joint angle of 180° (fully extended), and an ankle joint angle of 90° (neutral).

Recording at 15 Hz, a 60 mm, 128-element linear array ultrasound probe (LV7.5/60/128Z-2, 5.0-8.0

MHz; EchoBlaster128 CEXT-1Z, Telemed UAB; Vilnius, Lithuania) was placed over the gastrocnemius

medialis myotendinous junction (defined as the junction between the most distal point of the

gastrocnemius medialis and the AT) to measure the displacement of this landmark throughout all

protocols below. Firstly, the AT moment arm was calculated using the tendon excursion method

[36]. Next, participants performed four submaximal and one maximal plantar flexion contraction to

precondition the AT [37]. Then, participants performed graded isometric contractions of 5 seconds

each with 20 seconds of rest between each contraction, starting at 20 N.m and increasing in steps of

10 N.m until a maximum was reached, with 3-5 maximal attempts being performed [19]. The

displacement of the myotendinous junction during each contraction was taken as a measure of AT

elongation. Throughout all dynamometry procedures, participants were securely strapped into

position to minimise joint rotation. A 2D video camera (Sony RX10 Mk3, Sony Group Corporation,

Tokyo, Japan) in video mode (50 Hz) was used to record the lower limb in the sagittal plane, and any

observed change in ankle joint angle (i.e., from heel lift and dynamometer movement) was

measured. These data were used in conjunction with the tendon excursion data to correct the

observed myotendinous junction displacement during the contractions for the displacement due to

passive rotation [38].

To investigate the reliability of the manual measurement of AT elongation, a random sample of 10

maximal contractions were selected from the whole dataset. For intra-rater reliability, these images

were analysed twice by the lead researcher with an intervening period of 48 hours, and for inter-

rater reliability, a second experienced ultrasound operator analysed the same videos. The ICCs and

CVs for intra- and inter-rater reliability were 0.994 (95% confidence interval [CI]: 0.976-0.999) and

1.47% (CI: 0.69-2.26%), and 0.936 (CI: 0.762-0.984) and 4.80% (CI: 2.50-7.10%), respectively.

Vertical jumps

All vertical jumps were performed on a Kistler force plate (Kistler 9286BA, Switzerland) sampling at

1,000 Hz, and all force data were analysed using BioWare software (version 5.3.0.7, Kistler,

Switzerland). The contact threshold for take-off and landing was defined as the time point when

force fell below or rose above, respectively, the value equal to 5 standard deviations of the residual

force during the final 200 ms of the flight phase [39,40]. Participants completed four successful trials

of each jump, with the average of the best three trials (determined by the greatest flight time for

CMJ and SJ, and the greatest reactive strength index [RSI] for DJ) used for analysis [41,42].

Throughout all jumps, participants were required to keep their hands on their hips to remove the

effect of arm swing and isolate the lower body [43]. A 30 second rest was provided between each

jump trial.

For the CMJ, participants were instructed to start standing upright, then perform a

countermovement to a self-selected depth immediately into a maximal effort vertical jump, aiming

to maximise vertical displacement of the centre of mass [27]. For the SJ, participants were instructed

to descend to a self-selected squat depth [44], hold this posture for 3 seconds, then perform a

maximum effort concentric-only vertical jump, once again aiming to maximise vertical displacement.

Any trials with a visible countermovement or that displayed a force unload of ≥10% of body mass

before the initiation of the jump were excluded [45]. The onset of both the CMJ and SJ was defined

as the time point 30 ms before the vertical force trace moves beyond ± 5 standard deviations of

body weight from the quiet weighing period [46]. Finally, for the DJ, participants were instructed to

step forwards off a 30 cm box and to perform a maximal effort vertical jump immediately once they

make ground contact, whilst keeping ground contact time to a minimum [28].

Isometric strength tests

As well as the isometric plantar flexion contractions above, participants also performed unilateral

isometric knee extensions and the IMTP. For both tests, participants performed three submaximal

trials (at approximately 25, 50, and 75% of maximum effort) and three maximal effort trials. For the

knee extension, participants were seated on the Cybex dynamometer with a knee joint angle of 90°

and a hip joint angle of 110°, and all contractions were held for 5 seconds. All participants used their

left leg to quasi-randomise for limb dominance. IMTP testing followed the setup and protocol

guidelines set out by Comfort et al. [29]. In brief, participants stood in a squat rack on a force plate

(Kistler 9253B, Switzerland) with a knee joint angle of 125-145° and hip joint angle of 140-150°, with

the bar fixed at mid-thigh level. The participants were then instructed to push their feet into the

floor and pull on the bar "as fast and hard as possible" for 3 seconds [47,48]. Onset for the IMTP was

defined in the same manner as for the CMJ and SJ [49]. Participants had 30 and 60 seconds of rest

between each trial of the KE and IMTP, respectively. Strong verbal encouragement was used

throughout all strength tests.

Statistical analysis

All data are presented as mean ± standard deviation (SD). Data were checked for normality using a

Shapiro-Wilk test, and any variables that were not normally distributed were logarithmically

transformed before the calculation of ICC. To investigate relative and absolute test-retest reliability,

the ICC (two-way mixed effects, mean of k measurements, absolute agreement [50]) and CV were

used, respectively, both with 95% CI. The ICC was interpreted as ≤ 0.50 = *poor*, 0.50 to 0.75 =

moderate, 0.75 to 0.90 = *good*, and ≥ 0.90 = *excellent* [50]. The CV was calculated for each

participant as (*within-subject SD / mean) * 100*, with the mean of values from all participants used as

the test-retest CV (e.g., [24]). Additionally, the standard error of measurement (SEM, calculated as

SD_{pooled} ** √(1-ICC)*) and the minimum detectable change (MDC, calculated as *1.96 * √2 * SEM*) were

calculated. For intra-rater reliability the ICC form used was a two-way mixed effects, single

measurement, absolute agreement, whereas for inter-rater reliability, the two-way random effects,

single rater, absolute agreement was used [50]. Finally, a two-tailed paired *t*-test (or Wilcoxon test in

the case of non-normally distributed data) was used to assess systematic bias between testing

sessions, with α = 0.05. Microsoft Excel (Microsoft Corporations, Redmond, WA, USA) and SPSS

Statistics (IBM Corporations, Version 28.0, Armonk, NY, USA) were used for calculations and

statistical analyses. Due to technical issues, data are missing for one participant for AT elongation,

strain, and stiffness measures, and for two participants for the IMTP and knee extension tests.

Results

All results, including the ICC, CV, mean change, effect size and p-value of change, SEM and MDC, as

well as the relevant ICC-based descriptor are displayed for each variable in tables 1-5. For local MTU

metrics, the plantar flexion and knee extension MVCs displayed *excellent* reliability, the AT moment

arm and elongation measures all displayed *good* to *excellent* reliability, whereas all stiffness metrics

displayed *moderate* or *poor* reliability, except ATk_{index} which showed *good* reliability (table 1). All

CMJ variables displayed *good* to *excellent* reliability (table 2). For the SJ, all metrics showed *excellent*

reliability, except RFD time bands which were all *good*, and the eccentric utilisation ratio, which was

moderate (table 3). All DJ variables displayed *excellent* reliability (table 4). Peak force from the IMTP

displayed *excellent* reliability, whereas results for force-time variables were mixed, ranging from

moderate to *excellent*. Finally, the DSI displayed *good* reliability (table 5).

Table 1. Test-retest reliability of local muscle tendon unit metrics.

Variable	Session 1 (M ± SD)	Session 2 (M ± SD)	Change (M ± SD)	p	d	ICC (95% CI)	CV (95% CI)	SEM	MDC
Plantar flexion MVC (N.m)	124.2 (15.9)	126.2 (20.4)	2.0 (8.2)	0.328	0.23	0.948[e] (0.859-0.981)	3.6 (2.2-5.0)	4.2	11.6
Knee extension MVC (N.m)	125.1 (30.7)	130.8 (34.3)	5.7 (11.5)	0.079	0.46	0.961[e] (0.875-0.987)	5.9 (4.0-7.7)	6.4	17.8
Raw elongation (mm)	16.8 (4.0)	16.9 (4.3)	0.15 (2.71)	0.834	0.05	0.885[g] (0.666-0.960)	9.0 (5.8-12.2)	1.4	3.9
Passive elongation (mm)	5.1 (1.3)	4.9 (1.3)	−0.21 (0.63)	0.194	−0.32	0.931[e] (0.805-0.976)	7.7 (4.8-10.5)	0.3	0.9
Corrected elongation (mm)	11.7 (3.2)	12.0 (3.5)	0.36 (2.71)	0.604	0.13	0.812[g] (0.458-0.934)	13.9 (9.1-18.7)	1.5	4.0
Strain (%)	6.0 (2.0)	6.2 (2.0)	0.16 (1.42)	0.653	0.11	0.864[g] (0.610-0.953)	13.9 (9.1-18.7)	0.7	2.1
AT force (N)	4332.0 (687.8)	4335.9 (861.9)	3.9 (477.1)	0.974	0.01	0.902[e] (0.727-0.965)	5.8 (3.0-8.6)	244.1	676.6
ATk_{all} (N/m)	377.9 (143.4)	324.0 (98.5)	−53.9 (150.8)	0.173	−0.34	0.497[p] (−0.308-0.818)	20.0 (12.5-27.5)	96.6	267.7
ATk_{low} (N/m)	223.7 (94.0)	194.6 (61.3)	−29.1 (81.9)	0.176	−0.34	0.623[m] (−0.012-0.865)	21.8 (12.1-31.5)	48.7	135.1
ATk_{high} (N/m)	375.5 (287.7)	344.7 (357.4)	−30.8 (348.4)	0.729	−0.08	0.608[m] (−0.165-0.865)	30.9 (−28.7-90.5)	203.1	563.0
NATk_{all} (N/strain)	744.2 (263.8)	615.5 (193.3)	−128.7 (312.0)	0.120	−0.39	0.096[p] (−1.221-0.665)	22.9 (12.0-33.7)	213.1	590.6
NATk_{low} (N/strain)	448.0 (199.1)	328.4 (309.5)	−119.6 (339.2)	0.179	−0.34	0.687[m] (0.403-0.896)	−43.9 (−166.0-78.1)	225.38	624.71
NATk_{high} (N/strain)	732.7 (568.1)	576.8 (532.7)	−155.9 (559.1)	0.282	−0.27	0.340[p] (−1.295-0.795)	64.5 (11.3-117.7)	326.25	904.33
NATk_{index} (N/strain)	780.6 (219.8)	747.3 (208.0)	−33.3 (172.8)	0.452	−0.18	0.809[g] (0.460-0.933)	12.4 (7.6-17.1)	93.5	259.2
AT moment arm (mm)	28.9 (2.9)	29.5 (3.6)	0.6 (2.41)	0.337	0.23	0.842[g] (0.574-0.942)	4.0 (2.1-5.9)	1.3	3.6
Ankle rotation (°)	11.0 (2.7)	10.8 (3.1)	−0.3 (1.3)	0.415	−0.19	0.946[e] (0.853-0.980)	7.8 (5.2-10.3)	0.7	1.9

M, mean; SD, standard deviation; ICC, intraclass correlation coefficient; CV, coefficient of variation; SEM, standard error of measurement; MDC, minimum detectable change; AT, Achilles' tendon; ATk, Achilles' tendon stiffness; NATk, normalised Achilles' tendon stiffness.
Superscript letters refer to ICC classification, [p] = poor, [m] = moderate, [g] = good, [e] = excellent.

Table 2. Test-retest reliability of countermovement jump metrics.

Variable	Session 1 (M ± SD)	Session 2 (M ± SD)	Change (M ± SD)	p	d	ICC (95% CI)	CV (95% CI)	SEM	MDC
Take-off velocity (m/s)	2.061 (0.146)	2.082 (0.149)	0.021 (0.051)	0.084	0.39	0.966^e (0.906-0.988)	1.6 (1.1-2.1)	0.027	0.075
Jump height from take-off velocity (cm)	21.75(3.00)	22.20 (3.10)	0.40 (1.07)	0.102	0.40	0.966^e (0.905-0.988)	3.1 (2.2-4.1)	0.57	1.59
Flight time (s)	0.442 (0.033)	0.448 (0.034)	0.006 (0.011)	0.049*	−0.49	0.966^e (0.895-0.988)	1.7 (1.2-2.2)	0.006	0.017
Jump height from flight time (cm)	24.06(3.50)	24.70 (3.56)	0.64 (1.21)	0.049*	0.50	0.966^e (0.895-0.988)	3.4 (2.3-4.4)	0.68	1.88
Peak force (N)	1356.9 (233.2)	1394.1 (227.6)	37.2 (126.0)	0.241	0.28	0.917^e (0.777-0.970)	4.6 (2.6-6.6)	66.4	184.0
Peak power (W)	2386.0 (391.7)	2416.3 (395.5)	30.3 (106.1)	0.256	0.27	0.981^e (0.949-0.993)	2.2 (1.2-3.1)	54.3	150.4
Concentric RPD (W/s)	12495.7 (3688.9)	12842.8 (4060.1)	347.2 (3231.4)	0.943	0.10	0.788^g (0.402-0.924)	12.9 (5.7-20.1)	1743.4	4832.4
Rate of force development (N/s)	3993.6 (1641.1)	4763.7 (2371.8)	770.2 (1902.7)	0.163	0.39	0.791^g (0.443-0.923)	17.9 (9.3-26.6)	1115.2	3091.2
Positive impulse (N.s)	468.2 (80.3)	450.7 (80.5)	−17.6 (57.2)	0.653	−0.29	0.875^g (0.665-0.954)	5.2 (1.7-8.7)	31.1	86.3
Countermovement depth (cm)	25.93 (5.15)	25.65 (5.70)	0.28 (3.19)	0.718	−0.09	0.910^e (0.750-0.967)	6.7 (3.6-9.9)	1.63	4.52
Vertical stiffness (N/m)	5526.3 (1656.6)	5744.8 (1669.9)	218.6 (1433.8)	0.795	0.15	0.840^g (0.561-0.942)	10.3 (5.2-15.4)	783.7	2172.2
RSImod	0.302 (0.064)	0.319 (0.060)	0.017 (0.0416)	0.116	−0.38	0.860^g (0.620-0.949)	7.2 (4.2-10.2)	0.023	0.064

M, mean; SD, standard deviation; ICC, intraclass correlation coefficient; CV, coefficient of variation; SEM, standard error of measurement; MDC, minimum detectable change; RFD, rate of force development; RPD, rate of power development; RSImod, reactive strength index modified. Superscript letters refer to ICC classification, ^{p} = poor, ^{m} = moderate, ^{g} = good, ^{e} = excellent. * indicates a significant (p < 0.05) difference between sessions.

Table 3. Test-retest reliability of squat jump metrics.

Variable	Session 1 (M ± SD)	Session 2 (M ± SD)	Change (M ± SD)	p	d	ICC (95% CI)	CV (95% CI)	SEM	MDC
Take-off velocity (m/s)	1.984 (0.166)	1.983 (0.156)	−0.01 (0.080)	0.949	−0.02	0.937[e] (0.826-0.977)	2.2 (1.3-3.0)	0.040	0.112
Jump height from take-off velocity (cm)	20.21 (3.23)	20.16 (3.11)	−0.05 (1.63)	0.911	−0.03	0.933[e] (0.814-0.976)	4.3 (2.6-6.1)	0.08	2.28
Flight time (s)	0.420 (0.034)	0.421 (0.033)	0.002 (0.014)	0.619	0.11	0.962[e] (0.895-0.986)	2.0 (1.5-2.5)	0.007	0.019
Jump height from flight time (cm)	21.80 (3.40)	21.90 (3.23)	0.10 (1.00)	0.723	0.07	0.960[e] (0.891-0.986)	4.1 (3.1-5.1)	0.73	2.03
Peak force (N)	1262.6 (198.4)	1297.4 (221.7)	34.8 (69.6)	0.056	0.48	0.967[e] (0.897-0.988)	3.1 (1.9-4.3)	38.2	105.9
Peak power (W)	2305.9 (419.5)	2327.0 (416.8)	21.2 (104.6)	0.416	0.19	0.984[e] (0.958-0.994)	2.3 (1.4-3.3)	52.9	146.6
RPD (W/s)	6825.8 (1983.8)	7485.7 (2464.4)	659.8 (848.0)	0.005*	0.74	0.944[e] (0.727-0.983)	8.4 (5.9-11.0)	529.4	1467.4
RFD (N/s)	2196.9 (991.4)	2692.7 (1475.3)	495.8 (642.1)	0.010*	0.74	0.912[e] (0.625-0.972)	16.2 (11.2-21.2)	403.4	1118.1
RFD_{50} (N/s)	1564.6 (984.6)	1735.2 (1039.5)	170.7 (784.1)	0.383	0.21	0.825[g] (0.526-0.936)	24.7 (14.8-34.6)	423.5	1173.9
RFD_{100} (N/s)	2367.4 (1410.3)	2873.8 (1674.2)	506.4 (1125.9)	0.082	0.43	0.829[g] (0.532-0.938)	26.3 (16.5-36.1)	640.1	1774.2
RFD_{250} (N/s)	2094.5 (808.8)	2081.6 (773.9)	−12.9 (632.9)	0.934	−0.02	0.819[g] (0.490-0.935)	16.4 (9.4-23.4)	336.7	933.4
Eccentric Utilisation Ratio	1.08 (0.10)	1.11 (0.08)	0.02 (0.08)	0.312	0.24	0.738[m] (0.294-0.904)	4.4 (2.9-5.8)	0.05	0.13

M, mean; SD, standard deviation; ICC, intraclass correlation coefficient; CV, coefficient of variation; SEM, standard error of measurement; MDC, minimum detectable change; RPD, rate of power development; RFD, rate of force development. Superscript letters refer to ICC classification, p = poor, m = moderate, g = good, e = excellent. * indicates a significant (p < 0.05) difference between sessions.

Table 4. Test-retest reliability of drop jump metrics.

Variable	Session 1 (M ± SD)	Session 2 (M ± SD)	Change (M ± SD)	p	d	ICC (95% CI)	CV (95% CI)	SEM	MDC
Flight time (s)	0.391 (0.056)	0.402 (0.051)	0.011 (0.013)	0.006*	0.81	0.946[e] (0.868-0.993)	2.6 (1.7-3.6)	0.008	0.023
Jump height from flight time (cm)	19.16 (4.93)	20.11 (4.75)	0.95 (1.15)	0.006*	0.79	0.976[e] (0.872-0.993)	5.2 (3.4-7.1)	0.730	2.040
Ground contact time (s)	0.227 (0.031)	0.226 (0.028)	−0.001 (0.013)	0.723	−0.08	0.953[e] (0.869-0.983)	3.1 (2.0-4.2)	0.006	0.018
Reactive strength index	0.876 (0.277)	0.915 (0.263)	0.038 (0.053)	0.008*	0.69	0.985[e] (0.925-0.995)	4.8 (2.6-7.0)	0.032	0.089
Peak force (N)	3204.5 (808.4)	3145.4 (765.2)	−59.1 (212.1)	0.268	−0.27	0.981[e] (0.949-0.993)	3.4 (2.2-4.6)	108.5	300.7

M, mean; SD, standard deviation; ICC, intraclass correlation coefficient; CV, coefficient of variation; SEM, standard error of measurement; MDC, minimum detectable change. Superscript letters refer to ICC classification, p = poor, m = moderate, g = good, e = excellent. * indicates a significant ($p < 0.05$) difference between sessions.

Table 5. Test-retest reliability of isometric midthigh pull metrics.

Variable	Session 1 (M ± SD)	Session 2 (M ± SD)	Change (M ± SD)	p	d	ICC (95% CI)	CV (95% CI)	SEM	MDC
Peak force (N)	1614.1 (211.9)	1587.6 (209.9)	−26.5 (73.9)	0.187	−0.34	0.966[e] (0.902-0.989)	2.8 (1.8-3.7)	38.9	107.8
$Force_{50}$ (N)	726.3 (89.0)	716.7 (104.4)	−9.6 (38.7)	0.353	−0.24	0.959[e] (0.881-0.986)	3.1 (1.8-4.4)	19.6	54.4
$Force_{100}$ (N)	909.3 (145.5)	875.4 (148.4)	−33.8 (132.8)	0.341	−0.24	0.744[m] (0.256-0.913)	7.3 (3.4-11.1)	74.4	206.1
$Force_{250}$ (N)	1337.2 (247.9)	1330.4 (218.9)	−6.9 (145.3)	0.857	−0.05	0.899[g] (0.697-0.966)	6.4 (3.7-9.2)	74.3	206.0
$Impulse_{50}$ (N.s)	33.6 (3.9)	32.9 (4.3)	−0.7 (1.3)	0.051	−0.53	0.970[e] (0.893-0.990)	2.6 (1.6-3.5)	0.7	2.0
$Impulse_{100}$ (N.s)	74.2 (8.9)	72.9 (10.2)	−1.3 (4.6)	0.288	−0.27	0.937[e] (0.819-0.979)	3.5 (1.9-5.1)	2.4	6.7
$Impulse_{250}$ (N.s)	239.4 (43.5)	239.6 (39.4)	0.2 (33.1)	0.978	0.01	0.821[g] (0.452-0.940)	7.1 (2.8-11.4)	17.6	48.7
RFD_{50} (N/s)	1629.8 (1057.3)	1811.7 (1193.2)	181.9 (987.9)	0.488	0.17	0.768[g] (0.313-0.922)	35.9 (18.2-53.7)	543.0	1505.1
RFD_{100} (N/s)	2484.1 (1234.5)	2488.9 (1248.1)	4.8 (1304.5)	0.989	0.00	0.635[m] (−0.147-0.879)	29.0 (11.4-46.6)	749.9	2078.7
RFD_{250} (N/s)	2816.7 (931.8)	2821.3 (817.8)	4.6 (449.0)	0.696	0.01	0.934[e] (0.802-0.978)	10.6 (4.4-16.7)	225.2	624.3
Dynamic strength index	0.866 (0.100)	0.904 (0.151)	0.038 (0.105)	0.183	0.34	0.788[g] (0.398-0.928)	5.9 (3.3-8.5)	0.059	0.163

M, mean; SD, standard deviation; ICC, intraclass correlation coefficient; CV, coefficient of variation; SEM, standard error of measurement; MDC, minimum detectable change; RFD, rate of force development. Superscript letters refer to ICC classification, p = poor, m = moderate, g = good, e = excellent.

Discussion

The aim of this study was to report the test-retest reliability of local and global tests of MTU function

in naturally menstruating female athletes, whilst controlling for menstrual cycle phase at the time of

testing. Local measures, which included metrics of AT function and single-joint isometric strength

tests, displayed *good* to *excellent* reliability, except for traditional AT stiffness metrics which were

poor to *moderate*. For global measures, *good* to *excellent* reliability was found for all CMJ metrics, all

SJ metrics except eccentric utilisation ratio (*moderate*), and all DJ metrics. IMTP peak force also

showed *excellent* reliability, whereas IMTP force-time metrics ranged from *moderate* to *excellent*. An

overview of the ICC for each variable is provided in Fig 1. To the authors' knowledge, this is the first

study to investigate the reliability of many of these metrics in a female sample, and the first study to

standardise menstrual cycle phase of the participants at the time of testing. Therefore, these results

provide highly novel insight into the natural variability of MTU function within female athletes, data

which can be used to enhance the interpretation of a vast array of past and future female athlete

research.

Fig 1. Intraclass correlation coefficients (ICC) for all metrics. MTU, muscle-tendon unit; ATk, Achilles'

tendon stiffness; NATk normalised Achilles' tendon stiffness; ATkindex, Achilles' tendon stiffness

index; CMJ, countermovement jump; SJ, squat jump; DJ, drop jump; IMTP, isometric midthigh pull;

ISO, isometric; RFD, rate of force development; RPD, rate of power development; RSI, reactive

strength index; RSImod, reactive strength index modified; kY, vertical stiffness; DSI, dynamic

strength index; TOV, take-off velocity; FT, flight time; MVC, maximum voluntary contraction; PF,

plantar flexion; KE, knee extension.

A wide range of reliability scores were seen across the local and global measures of MTU function in

this study (Fig 1), and local measures generally had higher CVs than global ones (except for single-

joint isometric moments and global RFD metrics). When interpreting these findings, it is important to

consider that any change between sessions is comprised of a combination of measurement error

and biological variation [51]. It is plausible that any biological variation that is present could directly

affect local measures, but only indirectly affect global ones. This is because local metrics directly

measure some aspect of MTU function (e.g., tendon strain) whereas global measures have more

degrees of freedom and may employ many different motor strategies to achieve a consistent

performance outcome. Therefore, the performance outcome in a global test can still be maintained

from session to session despite potential changes to MTU function that underpin the movement. For

example, there is the potential for jump strategies to differ between sessions, but the overall

outcome of jump height does not (e.g., [52]). As such, global measures such as jump height might be

considered a 'blunt instrument' when it comes to detecting biological changes in MTU function,

whereas local measures can provide more sensitive insights into specific aspects of MTU function.

Two local measures of MTU function, plantar flexion and knee extension MVCs, displayed *excellent*

reliability (Table 1). These single joint isometric actions largely remove the 'skill' aspect of strength

testing [53] and provide reliable insights into the force-generating capacity of a muscle group. The

consistent plantar flexion MVC provides a conducive environment for measuring the mechanical

properties of the AT, and this analysis found that all AT metrics displayed *good* to *excellent*

reliability, except for stiffness measures which all displayed either *moderate* or *poor* reliability (Table

1). In contrast, ATk_{index}, calculated as maximum strain divided by maximum AT force [19], displayed

good reliability. In congruence with the present study, Rogers et al. [23] found that a similar variable

(stiffness calculated as maximum torque divided by maximum elongation) was more reliable than

traditional stiffness variables (ICC: 0.87), and Hunter et al. [19] also reported *good* reliability for

ATk_{index} in males (ICC: 0.75). This metric could be more reliable than traditional stiffness variables

(that calculate stiffness over certain ranges of the force-elongation curve) as it only takes into

account maximal contractions, meaning that any differences in the lower and mid-portions of the

force-elongation curve between days would not influence the outcome. In traditional stiffness

variables, these small differences are combined, and possibly amplified, in the subsequent

calculations, which may partially explain their lower reliability.

As traditional stiffness metrics account for AT force-elongation properties over a greater range of

forces, these metrics might provide a greater fidelity of information, but at the cost of lower

between-day reliability. This discrepancy in reliability between different stiffness metrics might also

explain the mixed reliability findings from previous research [13,16–18,20–23], as there is

considerable between-study heterogeneity in stiffness calculation methods, as well as actual

measurement methods (e.g., chosen landmark for tracking, contraction protocol). As this is the one

of the first studies to assess the reliability of these metrics in female athletes, the menstrual cycle

phase of the participants was controlled for to account for one potentially confounding source of

error. Whilst it is difficult to distinguish between the contribution of biological variation and

measurement error to the variability in these metrics, the manual measurement of AT elongation

displayed *excellent* intra- and inter-rater reliability in this study. This eliminates one key aspect of

measurement error and increases the likelihood that biological variability is the driver of the

observed changes, which means that stiffness metrics are perhaps not unreliable *per se*, but are

highly sensitive to small, biologically real changes (e.g., [54]). Ultimately, these metrics provide direct

insight into a specific aspect of MTU function, and because previous research has highlighted that AT

properties are able to distinguish between cohorts [55] and identify changes due to training

interventions [56], these metrics may still be of interest to researchers providing that the

reproducibility of the metrics is kept in mind.

Regarding global tests, most variables from the CMJ, SJ, and DJ showed *good* to *excellent* reliability,

with very little change between sessions (Tables 2-4). Insights into these global measures of lower-

body performance (e.g., jump height, force production, or RSI) are useful for measuring adaptations

to training interventions, and because of their high reliability, any observed changes are likely

meaningful. However, because of their increased degrees of freedom allowing for maintenance of

the overall performance outcome, these types of outcomes may not be sensitive enough to detect

smaller acute changes such as those expected to occur over menstrual cycle phases [9] or because of

fatigue [57]. Furthermore, although MTU function is often inferred through global tests, it must be

acknowledged that only local tests can provide objective evidence of specific aspects of MTU

function. For example, RSI and CMJ performance are often used as markers of stretch-shortening

cycle function [e.g., 58,59], although evidence has shown that true stretch-shortening cycles do not

occur for the triceps surae musculature during the CMJ [10] or DJ [60]. Therefore, as well as

considering reliability and sensitivity, practitioners should be mindful of the information that each

metric can provide when choosing tests and consider how this might inform training and testing

practices. A final consideration for global metrics is the use of composite metrics which are

comprised of two other variables, such as the eccentric utilisation ratio being calculated using jump

heights from the CMJ and SJ [61] and the dynamic strength index calculated from CMJ and IMTP

peak force [62]. Importantly, the results of the present study show that even though the input

variables show *excellent* reliability, these composite metrics displayed *moderate* and *good* reliability,

respectively. Thus, caution should be taken when considering these types of metrics and what

information they are able to provide above their constituent parts.

In this study, although high reliability was found for nearly all vertical jump metrics, systematic bias

was present between testing sessions for CMJ and DJ flight time and its associated jump height, DJ

RSI, and SJ RFD and rate of power development (RPD). The values for these variables were greater in

session two, and despite this bias, all variables displayed *excellent* ICCs and CVs <10% (excluding SJ

RFD), potentially highlighting that the bias is reflecting a small, but real change between sessions.

This change might be caused by changes in exercise technique (e.g., landing posture influencing

flight time in the CMJ and DJ) or through participants developing the neuromuscular skill of the

movement (i.e., a learning effect). This is particularly salient for the DJ, which because of its technical

complexity, has been shown to be less reliable than the CMJ and SJ [63]. Thus, the technique and

skill requirements for these global metrics and the potential for bias must be considered when

choosing tests. Familiarisation sessions can be used to reduce the risk of bias due to a learning

effect, and in the present study a familiarisation session was completed by all participants despite

previous research showing that familiarisation is not necessary for these three jump tests [64–66].

As the results presented here suggest that more familiarisation sessions are required, future

research should explore the effects of multiple familiarisations within a trained female athlete

sample. Overall, due to their generally *excellent* reliability, once participants are sufficiently

familiarised with the movement (to negate systematic bias), these vertical jump tests can reliably

assess lower-body performance capacity.

Much has been written about the importance of muscular strength and RFD for sports performance

[1]. Muscular strength, measured either locally via plantar flexion and knee extension MVCs or

globally via IMTP peak force, showed *excellent* reliability, underscoring the high reliability of strength

tests with limited degrees of freedom (Tables 1 and 5). The RFD measures from the CMJ, SJ and IMTP

resulted in *good* to *excellent* ICCs, but also high CVs (10.6-35.9%). In the SJ and IMTP, where RFD was

measured at 50, 100, and 250 ms from onset, the 250 ms time band proved to be the most reliable.

It is expected that calculating RFD over longer time epochs may result in improved reliability, as the

absolute change in force measurement produces a smaller relative error over longer time periods

and greater force outputs, a trend which has been reported in previous studies [49,67,68].

Additionally, early RFD is driven largely by neuromuscular contributions [47] and has been shown to

be more responsive to training [2], highlighting that it may be more sensitive to change than the

later time bands, explaining its poorer reliability scores. Therefore, if RFD is to be used, whilst longer

time bands (i.e., 250 ms) provide better reliability, it should be considered which aspects of

neuromusculoskeletal function are of interest when choosing metrics. It should also be noted that

for the IMTP, poorer reliability was found for RFD than the other IMTP force-time metrics (force and

impulse over the same 50, 100, and 250 ms time bands as RFD; Table 5), which is in accordance with

previous research [67,69,70]. Similarly, RPD displayed greater reliability than RFD in the CMJ and SJ

both in the present study and previous investigations [43,71]. Therefore, these findings caution

against the use of RFD and promote the use of other force-time metrics instead, such as RPD or

impulse.

This study controlled for menstrual cycle phase, using best practice recommendations [32], to

understand the natural variability of the chosen metrics in female athletes without the potentially

confounding influence of menstrual cycle related hormone changes. Although this design provides a

suitable level of control, future research should measure serum hormone concentrations and record

menstrual cycle symptoms to identify the effects that these factors have on outcomes. In addition,

to provide a greater scope of support to female athletes, research should also investigate the

variability of measures within individuals with other hormonal profiles, such as hormonal

contraceptive users or post-menopausal athletes.

Conclusions

This study investigated the test-retest reliability of local and global measures of MTU function in

female athletes, with the reliability of many variables reported for the first time. This is the first

reliability study investigating MTU, strength, and jump data to standardise the menstrual cycle phase

of the participants at the time of testing. In summary, local metrics were generally less reproducible

than global metrics. However, these metrics allow measurement of specific MTU characteristics that

determine function, and as such, can provide valuable information. *Good* to *excellent* reliability was

found for single-joint isometric strength and measures of AT function, except for traditional stiffness

metrics, which were *poor* to *moderate*. On the other hand, global metrics generally displayed high

reliability, with *good* to *excellent* ICCs reported for almost all CMJ, SJ, DJ, and IMTP metrics. These

tests can reliably assess overall performance outcomes; however, their increased degrees of

freedom means that they could be seen as 'blunt instruments' when it comes to monitoring and

inferring specific aspects of MTU function. The data reported here provides insight into the natural

variability of MTU function within a female athlete population, and therefore this data can be used

to contextualise and enhance the interpretation of other female athlete data, particularly that which

investigates other aspects of variability, such as the menstrual cycle.

Acknowledgements

382 The authors would like to thank the participants for their time and cooperation throughout the

383 study.

References

1.   Suchomel TJ, Nimphius S, Stone MH. The importance of muscular strength in athletic

performance. Sports Medicine. 2016 Oct 1;46(10):1419–49.

2.   Luis Hernández-Davó J, Sabido R. Rate of force development: reliability,

improvements and influence on performance. A review. European Journal of Human

Movement. 2014;33:46–69.

3.   Jarvis P, Turner A, Read P, Bishop C. Reactive strength index and its associations

with measures of physical and sports performance: a systematic review with meta-

analysis. Sports Medicine. 2021;52(2):301–30.

4.   Bojsen-Møller J, Magnusson SP, Rasmussen LR, Kjaer M, Aagaard P. Muscle

performance during maximal isometric and dynamic contractions is influenced by the

stiffness of the tendinous structures. J Appl Physiol. 2005;99(3):986–94.

5.   Cowley ES, Olenick AA, McNulty KL, Ross EZ. "Invisible sportswomen": The sex data

gap in sport and exercise science research. Women Sport Phys Act J.

2021;29(2):146–51.

6.   Sims ST, Heather AK. Myths and Methodologies: Reducing scientific design

ambiguity in studies comparing sexes and/or menstrual cycle phases. Exp Physiol.

2018;103(10):1309–17.

7.   Ansdell P, Brownstein CG, Škarabot J, Hicks KM, Simoes DCM, Thomas K, et al.

Menstrual cycle-associated modulations in neuromuscular function and fatigability of

the knee extensors in eumenorrheic women. Journal of Appl Physiology (1985).

2019;126(1):1701–12.

8.   Leblanc DR, Schneider M, Angele P, Vollmer G, Docheva D. The effect of estrogen

on tendon and ligament metabolism and function. Journal of Steroid Biochemistry and

Molecular Biology. 2017;172(April):106–16.

9.   McNulty KL, Elliott-Sale KJ, Dolan E, Swinton PA, Ansdell P, Goodall S, et al. The

effects of menstrual cycle phase on exercise performance in eumenorrheic women: a

systematic review and meta-analysis. Sports Medicine. 2020 Oct 1;50(10):1813–27.

10.   Kurokawa S, Fukunaga T, Nagano A, Fukashiro S. Interaction between fascicles and

tendinous structures during counter movement jumping investigated in vivo. J Appl

Physiol. 2003;95(6):2306–14.

11.   Lichtwark GA, Bougoulias K, Wilson AM. Muscle fascicle and series elastic element

length changes along the length of the human gastrocnemius during walking and

running. J Biomech. 2007;40(1):157–64.

12.   Houghton L, Dawson B, Rubenson J. Achilles tendon mechanical properties after both

prolonged continuous running and prolonged intermittent shuttle running in cricket

batting. J Appl Biomech. 2013;29(4):453–62.

13.   Burgess K, Connick M, Graham-Smith P, Pearson S. Plyometric vs. isometric training

influences on tendon properties and muscle output. J Strength Cond Res.

2007;21(3):986–9.

14.   Kubo K, Tabata T, Ikebukuro T, Igarashi K, Tsunoda N. A longitudinal assessment of
running economy and tendon properties in long-distance runners. J Strength Cond
Res. 2010;24(7):1724–31.

15.   Arampatzis A, Mersmann F, Bohm S. Individualized muscle-tendon assessment and
training. Front Physiol. 2020 Jun 26;11.

16.   Hansen P, Aagaard P, Kjaer M, Larsson B, Magnusson SP. Effect of habitual running
on human Achilles tendon load-deformation properties and cross-sectional area.
Journal of Experimental Medicine. 2003;95(6):2375–80.

17.   Fouré A, Nordez A, Cornu C. Plyometric training effects on Achilles tendon stiffness
and dissipative properties. J Appl Physiol [Internet]. 2010;109:849–54.

18.   Wu YK, Lien YH, Lin KH, Shih TTF, Wang TG, Wang HK. Relationships between
three potentiation effects of plyometric training and performance. Scand J Med Sci
Sports. 2010 Feb;20(1).

19.   Hunter S, Werth J, James D, Lambrianides Y, Smith K, Karamanidis K, et al.
Reliability and accuracy of a time-efficient method for the assessment of Achilles
tendon mechanical properties by ultrasonography. Sensors. 2022 Mar 26;22(7):2549.

20.   Robshaw DC. Gastrocnemius medialis muscle-tendon properties and their
relationship with power and reactive strength in elite youth soccer players. 2021.

21.   Kubo K, Kanehisa H, Fukunaga T. Effects of resistance and stretching training
programmes on the viscoelastic properties of human tendon structures in vivo.
Journal of Physiology. 2002;538(1):219–26.

22.   Ackermans TMA, Epro G, McCrum C, Oberländer KD, Suhr F, Drost MR, et al. Aging
and the effects of a half marathon on Achilles tendon force–elongation relationship.
Eur J Appl Physiol. 2016 Dec 1;116(11–12):2281–92.

23.   Rogers SA, Whatman CS, Pearson SN, Kilding AE. Assessments of mechanical

450   stiffness and relationships to performance determinants in middle-distance runners.

451   Int J Sports Physiol Perform. 2017 Nov 1;12(10):1329–34.

452   24.   Beattie K, Flanagan EP. Establishing the reliability & meaningful change of the drop-

453   jump reactive-strength index. Journal of Australian Strength and Conditioning.

454   2015;23(5):12–8.

455   25.   Stone MH, O'Bryant HS, Hornsby G, Cunanan A, Mizuguchi S, Suarez DG, et al.

456   Using the isometric mid-thigh pull in the monitoring of weightlifters: 25+ years of

457   experience. Professional Strength and Conditioning. 2019;(54):19–26.

458   26.   Keogh JAJ, Bishop C, Ruder MC, Kobsar D. Tracking the reliability of force plate-

459   derived countermovement jump metrics over time in female basketball athletes: a

460   comparison of principal component analysis vs. Conventional methods. Journal of

461   Science in Sport and Exercise [Internet]. 2023 Aug 15.

462   27.   McMahon JJ, Suchomel TJ, Lake JP, Comfort P. Understanding the key phases of

463   the countermovement jump force-time curve. Strength Cond J. 2018;40(4):96–106.

464   28.   Pedley JS, Lloyd RS, Read P, Moore IS, Oliver JL. Drop jump: a technical model for

465   scientific application. National Strength and Conditioning Association. 2017;39(5):36–

466   44.

467   29.   Comfort P, Dos Santos T, Beckham GK, Stone MH, Guppy SN, Gregory Haff G.

468   Standardization and methodological considerations for the isometric midthigh pull.

469   Strength Cond J. 2019;41(2):57–79.

470   30.   Moir GL, Garcia A, Dwyer GB. Intersession reliability of kinematic and kinetic

471   variables during vertical jumps in men and women. Int J Sports Physiol Perform.

472   2009;4(3):317–30.

31.   Schmalenberger KM, Tauseef HA, Barone JC, Owens SA, Lieberman L, Jarczok MN,

et al. How to study the menstrual cycle: Practical tools and recommendations.

Psychoneuroendocrinology. 2021;123.

32.   Elliott-Sale KJ, Minahan CL, de Jonge XAKJ, Ackerman KE, Sipilä S, Constantini NW,

et al. Methodological considerations for studies in sport and exercise science with

women as participants: a working guide for standards of practice for research on

women. Sports Medicine. 2021;51(5):843–61.

33.   McKay AKA, Stellingwerff T, Smith ES, Martin DT, Mujika I, Goosey-Tolfrey VL, et al.

Defining training and performance caliber: a participant classification framework. Int J

Sports Physiol Perform. 2021;1–15.

34.   World Medical Association. World Medical Association declaration of Helsinki: Ethical

principles for medical research involving human subjects. J Am Med Assoc. 2013 Nov

27;310(20):2191–4.

35.   Barfod KW, Riecke AF, Boesen A, Hansen P, Maier JF, Døssing S, et al. Validation of

a novel ultrasound measurement of achilles tendon length. Knee Surgery, Sports

Traumatology, Arthroscopy. 2015 Nov 1;23(11):3398–406.

36.   Fath F, Blazevich AJ, Waugh CM, Miller SC, Korff T. Direct comparison of in vivo

Achilles tendon moment arms obtained from ultrasound and MR scans. J Appl

Physiol. 2010 Dec;109(6):1644–52.

37.   Maganaris CN. Tendon conditioning: Artefact or property? Proceedings of the Royal

Society B: Biological Sciences. 2003;270(SUPPL. 1).

38.   Arampatzis A, Karamanidis K, Morey-Klapsing G, De Monte G, Stafilidis S.

Mechanical properties of the triceps surae tendon and aponeurosis in relation to

intensity of sport activity. J Biomech. 2007;40(9):1946–52.

39.   McMahon JJ, Murphy S, Rej SJE, Comfort P. Countermovement-jump-phase

characteristics of senior and academy rugby league players. Int J Sports Physiol

Perform. 2017 Jul 1;12(6):803–11.

40.   Pérez-Castilla A, Fernandes JFT, Rojas FJ, García-Ramos A. Reliability and

magnitude of countermovement jump performance variables: influence of the take-off

threshold. Meas Phys Educ Exerc Sci. 2021;25(3):227–35.

41.   Venegas-Carro M, Kramer A, Moreno-Villanueva M, Gruber M. Test–retest reliability

and sensitivity of common strength and power tests over a period of 9 weeks. Sports.

2022 Nov 1;10(11).

42.   Claudino JG, Cronin J, Mezêncio B, McMaster DT, McGuigan M, Tricoli V, et al. The

countermovement jump to monitor neuromuscular status: A meta-analysis. J Sci Med

Sport. 2017 Apr 1;20(4):397–402.

43.   Heishman AD, Daub BD, Miller RM, Freitas EDS, Frantz BA, Bemben MG.

Countermovement jump reliability performed with and without an arm swing in NCAA

Division 1 intercollegiate basketball players. J Strength Cond Res. 2020;34(2):546–

58.

44.   Petronijevic MS, Ramos AG, Mirkov DM, Jaric S, Valdevit Z, Knezevic OM. Self-

preferred initial position could be a viable alternative to the standard squat jump

testing procedure. J Strength Cond Res. 2018;32(11):3267–75.

45.   Sheppard JM, Doyle TLA. Increasing compliance to instructions in the squat jump. J

Strength Cond Res. 2008;22(2):648–551.

46.   Owen NJ, Watkins J, Kilduff LP, Bevan HR, Bennett MA. Development of a criterion

method to determine peak mechanical power output in a countermovement jump. J

Strength Cond Res. 2014;28(6):1552–8.

47.   Maffiuletti NA, Aagaard P, Blazevich AJ, Folland J, Tillin N, Duchateau J. Rate of

force development: physiological and methodological considerations. Eur J Appl

Physiol. 2016 Jun 1;116(6):1091–116.

48.   Rendos NK, Harriell K, Qazi S, Regis RC, Alipio TC, Signorile JF. Variations in verbal

encouragement modify isokinetic performance. J Strength Cond Res.

2018;33(3):708–16.

49.   Dos'santos T, Jones PA, Comfort P, Thomas C. Effect of different onset thresholds on

isometric midthigh pull force-time variables. J Strength Cond Res. 2017;31(12):3463–

73.

50.   Koo TK, Li MY. A guideline of selecting and reporting intraclass correlation

coefficients for reliability research. J Chiropr Med. 2016 Jun 1;15(2):155–63.

51.   Hopkins WG. Measures of reliability in sports medicine and science. Sports Medicine.

2000;30(1):1–15.

52.   Bishop C, Jordan M, Torres-Ronda L, Loturco I, Harry J, Virgile A, et al. Selecting

metrics that matter: comparing the use of the countermovement jump for performance

profiling, neuromuscular fatigue monitoring, and injury rehabilitation testing. Strength

Cond J. 2023;45(5):545–53.

53.   Buckner SL, Jessee MB, Mattocks KT, Mouser JG, Counts BR, Dankel SJ, et al.

Determining strength: a case for multiple methods of measurement. Sports Medicine.

2017 Feb 1;47(2):193–5.

54.   Walker J, Bissas A, Wainwright B, Hanley B, Cronin NJ. Repeatability and sensitivity

of passive mechanical stiffness measurements in the triceps surae muscle-tendon

complex. Scand J Med Sci Sports. 2022 Jan 1;32(1):83–93.

55.   Wiesinger HP, Rieder F, Kösters A, Müller E, Seynnes OR. Are sport-specific profiles

of tendon stiffness and cross-sectional area determined by structural or functional

integrity? PLoS One. 2016;11(6):40158441–6.

56.   Lazarczuk SL, Maniar N, Opar DA, Duhig SJ, Shield A, Barrett RS, et al. Mechanical,

material and morphological adaptations of healthy lower limb tendons to mechanical

loading: a systematic review and meta-analysis. Sports Medicine. 2022 Jun 3;

57.   Gathercole R, Sporer B, Stellingwerff T, Sleivert G. Alternative countermovement-

jump analysis to quantify acute neuromuscular fatigue. Int J Sports Physiol Perform.

2015 Jan 1;10(1):84–92.

58.   Southey B, Willshire M, Connick MJ, Austin D, Spits D, Beckman E. Reactive strength

index as a key performance indicator in different athlete populations – a systematic

review. Science and Sports. Elsevier Masson s.r.l.; 2023.

59.   Kozinc Ž, Žitnik J, Smajla D, Šarabon N. The difference between squat jump and

countermovement jump in 770 male and female participants from different sports. Eur

J Sport Sci. 2021.

60.   Ishikawa M, Niemelä E, Komi P V. Interaction between fascicle and tendinous tissues

in short-contact stretch-shortening cycle exercise with varying eccentric intensities. J

Appl Physiol. 2005;99(1):217–23.

61.   Mcguigan MR, Doyle TLA, Newton M, Edwards DJ, Nimphius S, Newton RU.

Eccentric utilization ratio: effect of sport and phase of training. J Strength Cond Res.

2006;20(4):992–5.

62.   Comfort P, Thomas C, Dos'Santos T, Jones PA, Suchomel TJ, McMahon JJ.

Comparison of methods of calculating dynamic strength index. Int J Sports Physiol

Perform. 2018 Mar 1;13(3):320–5.

63.   Gathercole RJ, Sporer BC, Stellingwerff T, Sleivert GG. Comparison of the capacity of
569   different jump and sprint field tests to detect neuromuscular fatigue. J Strength Cond
570   Res. 2015;29(9):2522–31.

571   64.   Moir G, Sanders R, Button C, Glaister M. The influence of familiarization on the
572   reliability of force variables measured during unloaded and loaded vertical jumps. J
573   Strength Cond Res. 2005;19(1):140–5.

574   65.   Nibali ML, Tombleson T, Brady PH, Wagner P. Influence of familiarization and
575   competitive level on the reliability of countermovement vertical jump kinetic and
576   kinematic variables. J Strength Cond Res. 2015;29(10):2827–35.

577   66.   Costley L, Wallace E, Johnston M, Kennedy R. Reliability of bounce drop jump
578   parameters within elite male rugby players. Journal of Sports Medicine and Physical
579   Fitness. 2018 Oct 1;58(10):1390–7.

580   67.   Ishida A, Suarez DG, Travis SK, Slaton JA, White JB, Bazyler CD, et al. Intrasession
581   and intersession reliability of isometric squat, midthigh pull, and squat jump in
582   resistance-trained individuals. J Strength Cond Res. 2021;37(1):18–26.

583   68.   Suarez DG, Carroll KM, Slaton JA, Rochau KG, Davis MW, Stone MH. Utility of a
584   shortened isometric midthigh pull protocol for assessing rapid force production in
585   athletes. J Strength Cond Res. 2022;36(7):1819–25.

586   69.   Thomas C, Dos'santos T, Comfort P, Jones PA. Between-session reliability of
587   common strength-and power-related measures in adolescent athletes. Sports. 2017
588   Mar 1;5(15).

589   70.   Guppy SN, Brady CJ, Kotani Y, Stone MH, Medic N, Haff GG. The effect of altering
590   body posture and barbell position on the between-session reliability of force-time
591   curve characteristics in the isometric mid-thigh pull. Sports. 2018 Dec 1;6(4).

592   71.   Souza AA, Bottaro M, Rocha Junior VA, Lage V, Tufano JJ, Vieira A. Reliability and

593   test-retest agreement of mechanical variables obtained during countermovement

594   jump. Int J Exerc Sci. 2020;13(4):6–17.

596   Supporting information

597   S1 Table. Definitions and calculations of local muscle-tendon unit metrics.

598   S2 Table. Definitions and calculations of global muscle-tendon unit metrics.

Figure