Chemical mate choice copying in *Drosophila melanogaster*

Bloo Mitchell[1], Alexandria Abbott[1], Ashanti Brown[1], Lacy Skinner[2], Elmira Umarova[1] & Philip Kohlmeier[1]

[1] Department of Biological Sciences, University of Memphis, Tennessee, USA

[2] Biology Department, Christian Brothers University, Memphis, Tennessee, USA

Contact author: Philip Kohlmeier; Email: pkhlmier@memphis.edu

Keyword: Behavior, Cuticular Hydrocarbons, Social learning, Mate choice, Sexual Selection, Pheromones

Abstract

Mate choice is a critical decision, requiring time and energy to assess potential partners' genetic

quality. Consequently, in many species females have evolved the ability to utilize social

information by copying the mate choices of others, usually based on visual cues. However,

chemical cues offer advantages, such as not requiring active observation of copulations. Using

Drosophila melanogaster, we provide the first demonstration of chemical mate choice copying.

Student females selected the same male genotype that a teacher female mated at higher

frequently than expected by chance. Chemical mate choice copying requires sensing both male

and female cues, which might indicate that that male genotype has been chosen by other

females. Our work suggests that females, in the presence of mated females, increase choosiness

at the virgin stage, elevating sexual selection on male traits. This study provides novel evidence

that exploiting social information is more prevalent in flies than previously assumed.

Introduction

Selecting a mating partner is a critical decision with significant fitness consequences, particularly for females, as reproduction-associated costs are typically higher in females than in males (Bateman 1948; Trivers 1972; Milinski and Bakker 1992). As a result, females have developed numerous mechanisms to evaluate the genetic quality of potential mates to ensure the acquisition of high-quality sperm. Male traits assessed by females for instance include coloration (Hill 2002), courtship song (Champagnon and Cueva del Castillo 2008), courtship intensity (Lin et al. 2016), and pheromonal profiles (Billeter and Levine 2013).

Collecting private information about the fitness of each male can be time and energy-intense especially if group size is large and males are abundant (Witte 2007). As an alternative strategy, females across species evolved the ability to exploit publicly available social information by monitoring, memorizing, and integrating mate choice decisions of other females. Such mate choice copying behaviors, whereby a female is more likely to select a male after observing this or a similar male being selected by another female, are wide-spread and have been documented across taxa including in birds (Drullion and Dubois 2008), fish (Dugatkin and Godin 1993), mammals (Kavaliers et al. 2017), and invertebrates (Mery et al. 2009). Most of the studies that reported the existence of mate choice copying relied on experimental setups that tested for mate copying via visual cues even if the used model organism primarily communicates via other sensory modalities, such as chemical cues (Scauzillo and Ferkin 2019). This is surprising as olfactory or gustatory cues can convey individual information about the male, such as age (Scauzillo and Ferkin 2019), dominance status (Scauzillo and Ferkin 2019), or parasite load (Claudio-Piedras et al. 2021) that can be relevant for female mate choice. Moreover, females that include chemical cues into their mate copying decisions do not have to be present and actively observe copulations of other females as residuals of the males can still be present on the female after mating. Unlike chemical mate choice copying (CMCC), visual mate choice copying might also be restricted species that mate during the day due to its dependency on light conditions. Despite the obvious advantage of CMCC, to our knowledge, empirical evidence for the existence of CMCC is lacking.

We tested whether the fruit fly *Drosophila melanogaster* uses CMCC. Fruit flies are social and form mixed sex groups in which multiple males are available as potential mating partners (Billeter et al., in revision). Female flies assess a large variety of phenotypic traits of males during mate choice decisions. These traits include body size (Markow 1988), age (Lin et al. 2016; Zhang et al. 2021), pheromonal profiles (Billeter and Levine 2013), courtship (Villella

and Hall 2008), nutritional status (Fricke et al. 2008), and symmetry (Møller and Pomiankowski

1993) among others which might be costly to collect individually. *D. melanogaster* is one of

the very few invertebrate species in which mate choice copying has been reported. Female flies

that observe another female mating with a poor-condition male are more likely to mate with a

poor-condition male themselves compared to naïve females that did not observe any matings

before (Mery et al. 2009). This effect was also observable when two similar males dusted with

differently colored fluorescent powder were used (Mery et al. 2009; Dagaeff et al. 2016;

Danchin et al. 2018; Nöbel et al. 2022*a*) demonstrating that observer females can monitor and

use neutral publicly available information and not differences in male condition or

attractiveness for mate copying. Females copy mate choices even if the selected male has visible

genetic defects suggesting that social information can overwrite privately collected information

(Nöbel et al. 2018). However, *D. melanogaster*, like most other insects (Wyatt 2003), primarily

communicates via chemical cues, and Cuticular Hydrocarbon (CHC) profiles are important

modulators of attractiveness and mate choice (Billeter and Levine 2013; Kohlmeier et al. 2021).

CHC profiles display stable quantitative differences between different *D. melanogaster* strains

which are genetically encoded (Jallon 1984; Sureau and Ferveur 1999; Grillet et al. 2006;

Billeter et al. 2009). Several male pheromones involved in female mate choice have been

identified and include *cis*-Vaccenyl Acetate, Palmitoleic acid, and 7-Tricosene as well as its

oxidation product heptanal (Grillet et al. 2006; Kurtovic et al. 2007; Billeter et al. 2009;

Kohlmeier et al. 2021; Doubovetzky et al. 2023; Verschut et al. 2023). Some of these

pheromones are transferred from the male to the female during copulation and hence, can

potentially be sensed by other females in the context of CMCC (Mane et al. 1983; Zawistowski

and Richmond 1986; Everaerts et al. 2010; Laturney and Billeter 2016). Despite of this, it

remains unknown whether *D. melanogaster* or any other invertebrate species uses CMCC.

We used an experimental paradigm in which student females are first exposed to recently mated

females and then tested whether they select the same male genotype as the teacher or not. Many

studies on female mate choice selected male genotypes that do not co-occur in the same

environment and potentially overpronounce variation in CHC profiles (Kohlmeier et al. 2021;

Doubovetzky et al. 2023) which questions to what extent the studied behaviors are ecologically

relevant. We circumvented this issue by using two isofemale lines generated from flies collected

in the same environment and thus, representing naturally occurring variation on male traits.

Material and Methods

Drosophila melanogaster flies were collected using banana traps in a residential area in

Memphis, TN, USA in July 2023. Two isofemale lines, named *Memphis-4* (*M4*) and *Memphis-*

7 (*M7*), were generated by transferring two females individually into standard vials (25x 95mm)

filled with 10ml fly food following the recipe described in (Kohlmeier et al. 2021) at +25°C

and 12:12 Light:Dark photoperiod. Lines were kept in the lab for more than 20 generations

before the onset of the experiments.

To test whether *D. melanogaster* females utilize chemical cues in mate choice copying, virgin

females of the laboratory strain *Oregon-R* (*OR*) were aged three to seven days in groups of 15.

Student females were collected without CO_2 anesthesia to prevent potential effects on memory

formation (Muria et al. 2021). These student females underwent one of four treatments (Figure

1): In the "No teacher" treatment, each female was individually placed into a Greiner Bio-One

arena (35mm diameter) containing fly food and one male from each of the *M4* and *M7* lines.

To distinguish between the males, they were dusted with pink or green fluorescent powder

(Slice of Moon™) at least 30 minutes prior to the experiment following established protocols

(Verschut et al. 2022). Females can distinguish even nuanced chemical profile differences

between males treated with different dusts (Kohlmeier et al., 2021). Colors for the males were

assigned randomly in each trial. Mate choice was recorded, and females that did not mate within

60 minutes were excluded. For the "*M4*-mated teacher" and "*M7*-mated teacher" treatments,

teacher *OR* virgins were placed in a dish with either two *M4* or two *M7* males for mating. After

copulation ended, these teacher females were paired in a new dish with a single virgin *OR*

student female for 45 minutes. After this teaching phase, student females were identified by the

absence of a fluorescent mating plug (Lung and Wolfner 2001) using a 395nm Ultraviolet

Flashlight (Morplot) and transferred to a new dish containing one colored *M4* and one colored

M7 male to record mate choice. In the "Virgin teacher" treatment, two *OR* virgin females

collected without CO_2 were kept together in a dish for 45 minutes. and one female was randomly

selected and transferred to a new arena with one colored *M4* and one colored *M7* male to assess

mate choice.

We observed that students prefer the male the teacher was mated with. This observation could

result from mate choice copying or a "habituation-based preference", where exposure to a

male's odor—via pheromonal residues left on the teacher after mating—increases the female's

preference for that male. To differentiate between these possibilities, student females were

exposed to CHC extracts from *M4*-mated teachers and *M4* males. To prepare these extracts, 15

virgin *OR* females and 15 *M4* males were placed together in a vial for 60 minutes to mate. After

mating, the flies were immobilized on ice, and mated females were identified by the presence

of a mating plug. Three mated females or three virgin *M4* males that had never encountered a

female were then transferred to a 2ml Silanized glass vial (Thermo Scientific). CHCs were

extracted by adding 150μl of hexane and vortexing the vial for 3 minutes (Verschut et al. 2022).

The extracts were then transferred onto a Whatman filter paper disc (5mm diameter), and the

hexane was allowed to evaporate for 30 minutes. The prepared discs were placed in a mating

arena, and a student female was added for a 45-minute exposure period. After that, mate choice

between *M4* and *M7* males was assessed as previously described.

In the analysis, mate choice outcomes were organized into two columns: one for *M4* males and

one for *M7* males, where '1' indicates that the respective male was selected for mating whereas

'0' indicates that this male was not selected. Both columns were combined using the *cbind*()

function in R to create a matrix of responses. One Generalized Linear Model (GLM) with a

binomial distribution was used for each experiment. Each GLM included the outcomes of the

mate choice tests as a response variable and treatment (*e.g.*, Mated female extract vs. male

extract) as an explanatory factor. In addition, one binomial test per treatment has been used to

test whether the preference for *M4* males differs from 0.5 *i.e.*, random choice.

Results

The proportion of students selecting *M4* males was influenced by teacher treatment (GLM: χ^2

= 12.1, *p* = 0.007; Figure 2A). Students exposed to a *M4*-mated teacher selected the *M4* male

more frequently than females exposed to no teachers (Model summary: z = 2.3, *p* = 0.019),

virgin teachers (Model summary: z = 2.4, *p* = 0.015), and *M7* mated teachers (Model summary:

z = 3.3, *p* = 0.001). There was no difference in the preference for *M4* males among the other

three treatments (Model summary: z < 1.1, *p* > 0.284). Mate choice of students exposed to *M4*-

mated teachers was higher than by random choice (Binomial test: *p* = 0.013), whereas mate

choice of students exposed to *M7*-mated teachers was lower than by random choice (Binomial

test: *p* = 0.048). Mate choice of students exposed to no teacher or a virgin teacher was not

different from random choice (Binomial test: *p* > 0.532).

Student females exposed to the CHC extracts of *M4*-mated teachers selected the *M4* male more

frequently than females exposed to the CHC extracts of *M4* males (GLM: χ^2 = 4.2, *p* = 0.041;

Figure 2B) and selected *M4* males more frequently than predicted by chance (Binomial test: *p*

= 0.049) whereas students exposed to extracts of *M4* males displayed no preference for one of

the two males (Binomial test: *p* = 0.533).

Discussion

The use of social information during mate choice has been suggested as a fast and low-cost strategy to identify high-quality mating partners (Scauzillo and Ferkin 2019). Most studies on mate choice copying are biased towards vertebrates and towards visual cues even if the used species primarily uses other sensory modalities such as olfaction (Scauzillo and Ferkin 2019). In this study, we provide the first evidence of CMCC. Exposure to a recently mated teacher female increases the likelihood that student females will choose a male similar to the one with whom the teacher mated. Our findings reveal that while an exposure to CHC extracts of mated females is sufficient to induce mate choice copying, an exposure to male odors alone does not enhance student female preference for the respective males.

Mate choice copying via visual cues has previously been reported in *D. melanogaster* (Mery et al. 2009; Dagaeff et al. 2016; Danchin et al. 2018; Nöbel et al. 2018, 2022*b*, 2022*a*, 2023). Our study now provides a novel pathway via which social information can be collected and exploited in the context of mate choice copying. Unlike visual cues, chemical signals enable females to gather social information without directly witnessing mating events, allowing them to assess the mate choices of a broader array of females within their group and to also evaluate mate choice decisions made by other females in the past. The extent to which the neural substrates that integrate visual and chemical cues overlap, and how flies respond when these cues provide conflicting information, remains to be elucidated. For example, visual mate choice copying requires the expression of the coincidence detector gene Rutabaga in the γ-Kanyon cells of the mushroom bodies (Nöbel et al. 2023). These cells receive input not only from visual but also from olfactory sensory neurons and are involved in the regulation of social behavior and responses to social experiences (Sun et al. 2020). It will thus be interesting to study if chemical and visual mate choice copying employ similar neural mechanisms to encode social information and whether γ-Kanyon Cells are involved in this process. Besides from mate choice, *D. melanogaster* flies utilize chemical cues provided by the social environment to assess whether another female has chosen strawberries or bananas as an oviposition substrate and to copy this decision (Battesti et al. 2012). Our findings on chemical mate choice copying suggest that the exploitation of social information and copying decisions of other flies via chemical cues may be more prevalent in flies than previously recognized.

Most studies investigating visual mate choice copying have employed artificial male phenotypes by marking males with fluorescent powder (Dagaeff et al. 2016; Danchin et al. 2018; Nöbel et al. 2022*a*, 2023). This method complicates the identification of visual cues that

females naturally assess, leaving it unclear under which conditions visual mate choice copying
occurs (Belkina et al. 2021). On the other hand, many studies investigating male traits assessed
by females during mate choice decisions, such as different concentrations of pheromones, often
use male genotypes that do not co-occur naturally within the same environment (*e.g.*, Kohlmeier
et al. 2021). This raises the question whether similar degrees of variation in the investigated
male traits occur in nature and consequently, whether the observed behaviors are ecologically
relevant. Our assay involved males from two different isofemale lines, both collected on the
same day and in the same environment, ensuring that we examined naturally occurring variation
in male traits. By using male genotypes that co-occur and focusing exclusively on their
chemical profiles, our approach supports the idea that mate choice copying is an ecologically
significant behavior exhibited by females in natural settings.

Our findings reveal that exposure to male extracts alone does not increase student females'
preference for that specific genotype. This indicates that student females must detect cues from
both males and mated females simultaneously to enhance their preference for a particular type
of male. Functionally, the reliance on cues specific to mated females may increase the
likelihood of a female choosing a male already selected by another female, thereby potentially
enhancing reproductive success. It is likely that the female cues assessed by the students include
compounds only found on females *i.e.*, 7,11-Heptacosadiene and 7,11-Nonacosadiene (Billeter
and Levine 2013) and/or those compounds, characteristic of mated females, such as *cis*-
Vaccenyl acetate and 7-Tricosene which are both transferred from the male to the female during
copulation (Ejima et al. 2007; Laturney and Billeter 2016). Relying of pheromones
characteristic for mated females would additionally protect the female from developing a
preference for a specific male in presence of other virgin females indicating that this male has
not yet been selected by other females. On a mechanistic level, these data suggest that the
neuronal circuitry regulating CMCC requires inputs from neurons sensitive to both male and
female cues. Which male pheromones are evaluated by the female to identify similar males is
unclear. Potential candidates include pheromones previously found to be involved in mate
choice such as Palmitoleic Acid (Lin et al. 2016; Kohlmeier et al. 2021), 7-Tricosene (Grillet
et al. 2006), and 7-Pentacosene (Ferveur 2005).

Mate choice copying can increase sexual selection on males, particularly under high-density
conditions. Typically, *D. melanogaster* virgin females choose their first mating partner
indiscriminately to secure a quick acquisition of sperm. Following the initial mating, they
become more selective about male quality to enhance offspring fitness (Kokko and Mappes
2005; Kohlmeier et al. 2021). In scenarios where population density is high and mated females

are abundant, mate choice copying can enable virgin females to develop an increased selectiveness for males frequently chosen by others. This allows virgin females to reduce the risk of mating with low-quality males due to the initial lack of choosiness. This mechanism suggests that sexual selection on males is likely to be more intense in populations with high densities. It would be interesting to explore whether the presence of one or a few mated females can initiate a cascade effect among virgin females, leading to a group-level preference for specific male types.

In summary, we describe chemical mate choice copying as a novel behavior *in D. melanogaster* and our data suggest that female and male cues have to be sensed in parallel to induce copying behavior potentially protecting females from developing a preference for males that have not selected by other females. This new behavior might impact sexual selection dynamics particularly when population density is high.

Acknowledgements

We thank Rebecca Crenshaw, student of the University of Memphis 2023 Neurogenetics CURE class for bringing this work to the lab and Jean-Christophe Billeter for his feedback on an earlier version of this manuscript.

Figures

Teacher mating   **Teaching phase**   **Mate choice phase**

Figure 1: Experimental setup of the first experiment. Teacher females were mated with either *M4* or *M7* males. After mating ended, teacher females were transferred to student females for a 45min teaching phase. Student females were then offered one colored *M4* and one colored *M7* male for mate choice. In the "Virgin teacher" treatment, virgin teachers that had no prior contact with *M4* or *M7* males. In the "No teacher" treatment, mate choice of student females was tested without any teaching phase. Figure created using BioRender.

Figure 2: Student preferences for M4 males after teacher and CHC extract exposure. (**A**) Exposure to *M4*-mated teachers increases the preference for *M4* males in student females whereas an exposure to *M7*-mated teachers results in a preference for *M7* males. (**B**) Exposing students to the CHC extracts of *M4*-mated teachers increases their preference for *M4* males, whereas students exposed to the extracts of unmated *M4* males do not exhibit a preference for one of the two males. Bars labelled with the same letters are not statistically different. Numbers in each bar indicate sample size. * above a bar indicates a statistically significant difference from random choice *i.e.*, 0.5, based on binomial tests. No * indicates no difference from random choice.

References

Bateman, A. J. 1948. Intra-sexual selection in *Drosophila*. Heredity 2:349–368.

Battesti, M., C. Moreno, D. Joly, and F. Mery. 2012. Spread of social information and dynamics of social transmission within *Drosophila* groups. Current Biology 22:309–313.

Belkina, E. G., A. Shiglik, N. G. Sopilko, S. N. Lysenkov, and A. V. Markov. 2021. Mate choice copying in *Drosophila* is probably less robust than previously suggested. Animal Behaviour 176:175–183.

Billeter, J. C., J. Atallah, J. J. Krupp, J. G. Millar, and J. D. Levine. 2009. Specialized cells tag sexual and species identity in *Drosophila melanogaster*. Nature 461:987–991.

Billeter, J. C., and J. D. Levine. 2013. Who is he and what is he to you? Recognition in *Drosophila*
melanogaster. Current Opinion in Neurobiology 23:17–23.

Champagnon, J., and R. Cueva del Castillo. 2008. Female mate choice, calling song and genetic
variance in the cricket, *Gryllodes sigillatus*. Ethology 114:223–230.

Claudio-Piedras, F., B. Recio-Tótoro, J. Cime-Castillo, R. Condé, M. Maffei, and H. Lanz-Mendoza.
2021. Dietary and Plasmodium challenge effects on the cuticular hydrocarbon profile of *Anopheles*
albimanus. Scientific Reports 11:11258.

Dagaeff, A. C., A. Pocheville, S. Nöbel, A. Loyau, G. Isabel, and E. Danchin. 2016. *Drosophila* mate
copying correlates with atmospheric pressure in a speed learning situation. Animal Behaviour
121:163–174.

Danchin, E., S. Nöbel, A. Pocheville, A.-C. Dagaeff, L. Demay, M. Alphand, S. Ranty-Roby, et al.
2018. Cultural flies: Conformist social learning in fruitflies predicts long-lasting mate-choice
traditions. Science 362:1025–1030.

Doubovetzky, N., P. Kohlmeier, S. Bal, and J.-C. Billeter. 2023. Cryptic female choice in response to
male pheromones in *Drosophila melanogaster*. bioRxiv
12/21/2023:https://doi.org/10.1101/2023.12.20.572608.

Drullion, D., and F. Dubois. 2008. Mate-choice copying by female zebra finches, *Taeniopygia guttata*:
What happens when model females provide inconsistent information? Behavioral Ecology and
Sociobiology 63:269–276.

Dugatkin, L. A., and J.-G. J. Godin. 1993. Female mate copying in the guppy (*Poecilia reticulata*):
age-dependent effects. Behavioral Ecology 4:289–292.

Ejima, A., B. P. C. Smith, C. Lucas, W. Van Der Goes Van Naters, C. J. Miller, J. R. Carlson, J. D.
Levine, et al. 2007. Generalization of courtship learning in *Drosophila* is mediated by *cis*-vaccenyl
acetate. Current Biology 17:599–605.

Everaerts, C., J. P. Farine, M. Cobb, and J. F. Ferveur. 2010. *Drosophila* cuticular hydrocarbons
revisited: Mating status alters cuticular profiles. PLoS ONE 5:e9607.

Ferveur, J. F. 2005. Cuticular hydrocarbons: Their evolution and roles in *Drosophila* pheromonal
communication. Behavior Genetics 35:279–295.

Fricke, C., A. Bretman, and T. Chapman. 2008. Adult male nutrition and reproductive success in
Drosophila melanogaster. Evolution 62:3170–3177.

Grillet, M., L. Dartevelle, and J. F. Ferveur. 2006. A *Drosophila* male pheromone affects female
sexual receptivity. Proceedings of the Royal Society B: Biological Sciences 273:315–323.

Hill, G. E. 2002. A red bird in a brown bag. Oxford University Press, New York.

Jallon, J. M. 1984. A few chemical words exchanged by *Drosophila* during courtship and mating.
Behavior Genetics 14:441–478.

Kavaliers, M., R. Matta, and E. Choleris. 2017. Mate-choice copying, social information processing,
and the roles of oxytocin. Neuroscience and Biobehavioral Reviews 72:232–242.

Kohlmeier, P., Y. Zhang, J. Gortner, C.-Y. Su, and J.-C. Billeter. 2021. Mating increases *Drosophila*
melanogaster females' choosiness by reducing olfactory sensitivity to a male pheromone. Nature
Ecology & Evolution 5:1165–1173.

Kokko, H., and J. Mappes. 2005. Sexual selection when fertilization is not guaranteed. Evolution
59:1876–1885.

Kurtovic, A., A. Widmer, and B. J. Dickson. 2007. A single class of olfactory neurons mediates
behavioural responses to a *Drosophila* sex pheromone. Nature 446:542–546.

Laturney, M., and J.-C. Billeter. 2016. *Drosophila melanogaster* females restore their attractiveness
after mating by removing male anti-aphrodisiac pheromones. Nature Communications 7:12322.

Lin, H. H., D. S. Cao, S. Sethi, Z. Zeng, J. S. R. Chin, T. S. Chakraborty, A. K. Shepherd, et al. 2016.
Hormonal modulation of pheromone detection enhances male courtship success. Neuron 90:1272–
1285.

Lung, O., and M. F. Wolfner. 2001. Identification and characterization of the major *Drosophila*
melanogaster mating plug protein. Insect Biochemistry and Molecular Biology 31:543–551.

Mane, S. D., L. Tompkins, and R. C. Richmond. 1983. Male Esterase 6 catalyzes the synthesis of a sex
pheromone in *Drosophila melanogaster* Females. Science 222:419–421.

Markow, T. A. 1988. Reproductive behavior of *Drosophila melanogaster* and *D. nigrospiracula* in the
field and in the laboratory. Journal of Comparative Psychology 102:169–173.

Mery, F., S. A. M. Varela, É. Danchin, S. Blanchet, D. Parejo, I. Coolen, and R. H. Wagner. 2009.
Public versus personal information for mate copying in an invertebrate. Current Biology 19:730–734.

Milinski, M., and T. C. M. Bakker. 1992. Costs influences sequential mate choice in sticklebacks,
Gasterosteus aculeatus. Proceedings of the Royal Society of London. Series B: Biological Sciences
250:229–233.

Møller, A. P., and A. Pomiankowski. 1993. Fluctuating asymmetry and sexual selection. Genetica
89:267–279.

Muria, A., P. Y. Musso, M. Durrieu, F. R. Portugal, B. Ronsin, M. D. Gordon, R. Jeanson, et al. 2021.
Social facilitation of long-lasting memory is mediated by CO_2 in *Drosophila*. Current Biol 31:2065–
2074.

Nöbel, S., E. Danchin, and G. Isabel. 2018. Mate-copying for a costly variant in *Drosophila*
melanogaster females. Behavioral Ecology 29:1150–1156.

Nöbel, S., E. Danchin, and G. Isabel. 2023. Mate copying requires the coincidence detector Rutabaga
in the mushroom bodies of *Drosophila melanogaster*. iScience 26:107682.

Nöbel, S., M. Monier, L. Fargeot, G. Lespagnol, E. Danchin, and G. Isabel. 2022*a*. Female fruit flies
copy the acceptance, but not the rejection, of a mate. Behavioral Ecology 33:1018–1024.

Nöbel, S., M. Monier, D. Villa, É. Danchin, and G. Isabel. 2022*b*. 2-D sex images elicit mate copying
in fruit flies. Scientific Reports 12:22127.

Scauzillo, R. C., and M. H. Ferkin. 2019. Factors that affect non-independent mate choice. Biological
Journal of the Linnean Society 128:499–514.

Sun, Y., R. Qiu, X. Li, Y. Cheng, S. Gao, F. Kong, L. Liu, et al. 2020. Social attraction in *Drosophila*
is regulated by the mushroom body and serotonergic system. Nature Communications 11:5350.

Sureau, G., and J. F. Ferveur. 1999. Co-adaptation of pheromone production and behavioural
responses in *Drosophila melanogaster* males. Genetical Research 74:129–137.

Trivers, R. L. 1972. Parental investment and sexual selection. Pages 52–95 *in* Sexual Selection and the
Descent of Man: The Darwinian Pivot. Aldine, Chicago.

Verschut, T. A., P. Kohlmeier, and J.-C. Billeter. 2022. Bioassaying the function of pheromones in
Drosophila melanogaster's social behavior. Pages 123–157 *in* Behavioral Neurogenetics.

Verschut, T. A., R. Ng, N. P. Doubovetzky, G. Le Calvez, J. L. Sneep, A. J. Minnaard, C.-Y. Su, et al.
2023. Aggregation pheromones have a non-linear effect on oviposition behavior in *Drosophila*
melanogaster. Nature Communications 14:1544.

Villella, A., and J. C. Hall. 2008. Chapter 3 Neurogenetics of courtship and mating in *Drosophila*.
Pages 67–184 *in* Advances in Genetics.

Witte, K. 2007. Learning and Mate Choice. Pages 70–95 *in* Fish Cognition and Behavior. Wiley
Blackwell.

Wyatt, T. D. 2003. Pheromones and animal behaviour (1st ed.). Cambridge University Press,
Cambridge.

Zawistowski, S., and R. C. Richmond. 1986. Inhibition of courtship and mating of *Drosophila*
melanogaster by the male-produced lipid, cis-vaccenyl acetate. Journal of Insect Physiology 32:189–
192.

Zhang, S., E. Glantz, D. Rogulja, and M. Crickmore. 2021. Hormonal control of motivational circuitry
orchestrates the transition to sexuality in *Drosophila*. Science Advances 7:eabg6926.

423