Hybridization load introduced by Alpine ibex hybrid swarms

F. Gözde Çilingir[1,2+], Fabio Landuzzi[3+], Alice Brambilla[2,4], Debora Charrance[3], Federica Furia[3], Sara Trova[5], Alberto Peracino[4], Glauco Camenisch[2], Dominique Waldvogel[2], Jo Howard-McCombe[6], Yeraldin Chiquinquira Castillo De Spelorzi[7], Edoardo Henzen[7], Andrea Bernagozzi[8], Alessandro Coppe[3], Jean Marc Christille[8], Manuela Vecchi[5], Diego Vozzi[7], Andrea Cavalli[3,9], Bruno Bassano[4], Stefano Gustincich[5,10], Daniel Croll[11,*], Luca Pandolfini[5,10**] and Christine Grossen[1***]

[1] Swiss Federal Research Institute WSL, Biodiversity and Conservation Biology Research Unit, Birmensdorf, Switzerland

[2] University of Zurich, Department of Evolutionary Biology and Environmental Studies, Zurich, Switzerland

[3] Computational and Chemical Biology, Italian Institute of Technology (IIT), CMP^{3}VdA, Aosta, Italy

[4] Alpine Wildlife Research Center, Gran Paradiso National Park, Turin, Italy

[5] Non-coding RNA and RNA-based therapeutics, Italian Institute of Technology (IIT), CMP^{3}VdA, Aosta, Italy

[6] WildGenes Laboratory, Royal Zoological Society of Scotland, Edinburgh, United Kingdom

[7] Genomics Facility, Istituto Italiano di Tecnologia (IIT), Genoa, Italy

[8] Fondazione Clément Fillietroz ONLUS, Astronomical Observatory of Autonomous Region of Aosta Valley, Aosta, Italy

[9] CECAM-EPFL, Lausanne, Switzerland

[10] Non-coding RNA and RNA-based therapeutics, Center for Human Technology, Italian Institute of Technology (IIT), Genoa, Italy

[11] Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland

[+] These authors equally contributed to the work

*co-corresponding authors (daniel.croll@unine.ch, luca.pandolfini@iit.it, christine.grossen@wsl.ch)

lead contact (christine.grossen@wsl.ch)

Summary

Species restoration efforts can be threatened by the accumulation of deleterious mutations and inbreeding depression associated with the historic population contraction. However, even successfully restored species could face deleterious mutation swamping from hybridization with an abundant and closely related species. Here, we analyze this risk for Alpine ibex (*Capra ibex*), a flagship species of large mammal restoration in the Alps. The Alpine ibex faced near-extinction two centuries ago, resulting in exceptionally low genome-wide diversity and increased inbreeding, which facilitated the purging of severe deleterious mutation load. For this, we produced a highly contiguous chromosome-level genome assembly of the Alpine ibex capturing structural divergence from domestic goat (*Capra hircus*) and mapping immune-relevant MHC genes. Analyses of eight recent ibex-goat hybrids from two swarms in Northern Italy, combined with 29 non-hybrid Alpine ibex and 22 domestic goats, identified 215 masked loss-of-function (LOF) mutations introduced via hybridization. Yet, we found no evidence for counter-selection in early backcrosses. This exposes Alpine ibex to further backcrosses compounding the deleterious mutation load of the species by a factor of up to two. Our work provides one of the first direct estimates of hybridization load and guides conservation efforts to preserve endangered species gene pools.

Keywords: conservation genetics, deleterious mutation load, anthropogenic hybridization, hybrid swarm, hybridization load, heterosis, Alpine ibex

Results and discussion

Human activity has profoundly affected species and populations through habitat degradation and over-exploitation leading to severe population reductions and extinctions [1]. A particular concern for species conservation is increased levels of inbreeding and the expression of deleterious mutations in small populations. Recent studies provided compelling evidence for changes in deleterious mutation load in natural populations (reviewed in Dussex et al. [2]; Robinson et al. [3]). As predicted by theory, population bottlenecks were repeatedly shown to lead to the selective removal (purging) of highly deleterious (usually recessive) mutations, while less severely deleterious mutations can accumulate (*e.g.,* Xue et al.[4], Grossen et al. [5], Dussex et al. [6]; reviewed in Dussex et al. [2]). Mutation load is defined as load resulting from newly introduced deleterious mutations [7]. Empirical assessments of fitness effects confirmed the detrimental impact of mutation load and inbreeding on both individual and population fitness [8,9]. However, the origin of deleterious mutations remains often unresolved. In addition to *de novo* mutations, a potentially significant source of mutation load could be deleterious variants introduced into a population through hybridization.

Hybridization in nature contributes to genetic variation and fosters adaptation [10,11]. Hybridization in bottlenecked species may introduce new adaptive genetic variation, compensating previously lost variation [12]. In addition, high levels of heterozygosity in hybrids can also initially mask deleterious mutations, thereby causing hybrid vigor (*i.e.*, heterosis) and mitigating inbreeding. Yet, this initial masking of deleterious mutations prevents their counter-selection in early generations. Hybridization also poses the risk of disrupting local adaptation through introgression of non-native alleles, in particular if the new species contact is caused by human activities [13–15]. Such introduction of deleterious mutations is defined as hybridization load [12]. Overall, deleterious mutations played a significant role in hybridization events in animals such as fish [16,17], plants including crops [18–20], and trees [21]. These studies indicate that while hybridization can introduce beneficial alleles, linked deleterious variants can reduce fitness and create partial barriers to introgression. The interplay of adaptive and maladaptive variation introduced through hybridization is likely complex and poorly understood.

As hybridization can introduce new adaptive genetic variation and alleviate the effects of inbreeding through hybrid vigour, genetically impoverished species may be particularly susceptible to the introduction of hybridization load. To quantify load in genomes and to predict effects of introgression, high-quality genomic resources are essential [22]. The Alpine ibex (*Capra*

86 *ibex*), a mountain ungulate endemic to the European Alps (Figure 1A, B), has experienced severe
87 bottlenecks since the early 19th century and repeated hybridization events [12,23,24]. However, the
88 evolution of deleterious mutation load driven by hybridization remains poorly understood.
89
Conservation efforts have restored the species to its original distribution area, with a current
population of about 55,000 individuals [25]. However, these bottlenecks have resulted in Alpine ibex
having one of the lowest levels of genetic variation among mammals [5,26]. This reduced genetic
diversity has led to increased inbreeding and inbreeding depression, with negative effects both at
the individual [27] and population level [28]. Severe genetic load was likely reduced through purging
of highly deleterious mutations, while less severe mutations accumulated [5]. Genome-wide
analyses showed that successful hybridizations with domestic goats (*Capra hircus*) likely occurred
during the severe historic bottleneck when ibex numbers were at their lowest [29,30]. Introgression
was particularly evident at immune-relevant genes, where genetic variability in Alpine ibex is very
low. Such introgressed variants may have been under positive selection due to heterozygote
advantage at immune loci [29]. However, hybridization is also expected to have detrimental
consequences for Alpine ibex, with potential loss of adaptations to the alpine environment and
the introduction of deleterious variants. Of particular concern are the recently discovered hybrid
swarms between Alpine ibex and domestic goats in Northern Italy [24,31]. Such populations of
hybrids have survived beyond the initial hybrid (F1) generation, and there is continued
backcrossing between hybrid individuals and parental species [24]. This extensive hybridization
raises concerns of maladaptation and loss of ibex identity. Here, we generated the first reference-
quality genome for Alpine ibex to quantify the hybridization load generated by high rates of
hybridization as observed in hybrid swarms.
109

A high-quality chromosome-level assembly
We assembled a chromosome-level genome of a wild female Alpine ibex (Odissea) from the Gran
Paradiso National Park population (Lauson Valley, Cogne, Italy). Using 130 Gb of high-quality
ONT data, we generated a 2.66 Gb draft genome with 1040 contigs (N50 58.7 Mb), polished with
short read sequencing data. The contigs showed high completeness, with BUSCO and k-mer-
based scores of 96.1% and 98.6%, respectively (Figure 1C). We generated Hi-C data to anchor
contigs into 30 pseudo-chromosomes (Figure S1) with a resulting N50 of 101.2 Mb and matching
the chromosome numbers of domestic goats [32] (Table S1, Figure 1D). The Alpine ibex assembly
also includes the full mitochondrial genome (16.7 Kb, Figure S2). The nuclear genome is
constituted of 43.4% repetitive elements (1.15 Gb), dominated by retroelements (38.22%),

including Long Interspersed Nuclear Elements (LINEs: 32.56%) and Long Terminal Repeats (LTRs: 4.87%) (Table S2). We integrated gene model evidence using transcriptome-assisted predictions as well as annotations lifted over from the sheep (*Ovis aries*) genome, yielding a BUSCO completeness of 99.7% at the transcript level (Table S3). The annotated gene set consisted of 32,783 protein-coding genes and 61,420 transcripts (Table S4).

Figure 1: **Sampling and genome description.** A) Picture of a female Alpine ibex with her offspring (photo by Dario de Siena, Archivio PNGP). B) Map of species distribution and zoom in on the sampling location of Odissea (shown in yellow), the hybrids (green), and domestic goats (blue). C) Snail plot visualizing assembly statistics: scaffold statistics, sequence composition proportions, and BUSCO completeness scores. D) Riparian plot showing the results of the gene synteny analysis conducted on the cattle, sheep, domestic goat, and Alpine ibex genomes using GENESPACE. The human genome was used as an outgroup. See also Figures S1-S4.

Recent structural variants arising in Alpine ibex

We used gene order conservation analyses between Alpine ibex and three related domesticated
mammals, including cattle (*Bos taurus*), sheep (*Ovis aries*), and domestic goat, as well as the
human genome as an outgroup to detect chromosomal rearrangements (Figure 1D; Tables S5
and S6). Chromosomal synteny is largely conserved between the domestic goat and the Alpine
ibex. The sheep genome experienced three chromosome fusions, combining in each two distinct
chromosomes found in cattle and goats. The Alpine ibex genome includes a fully assembled X
chromosome compared to the domestic goat (contigs Xa and Xb; Figure 1D). In addition, the
Alpine ibex chromosomes 12 and 17 also include sequences matching to unplaced contigs in
domestic goat. Overall, domestic goat and Alpine ibex genomes share 2.4 Gb of syntenic regions
(~90.2%). Structural variant analyses with SyRI confirmed major rearrangements on
chromosomes 7, 18, and 23 (Figure S3, S4A), mainly involving indels (38 Mb, 1.4% of the
genome) and inversions (49 Mb, 1.8%), with fewer duplications and translocations (Figure S4B,
C). Highly divergent regions (HDR) covered a total of 43 Mb (1.6%; Figure S4B, C). Nearly half
(45%) of all insertions, deletions, and HDRs between domestic goat and Alpine ibex overlapped
with gene bodies, and the proportion was even higher for inversions (52%), duplications (54%),
and translocations (78%).

154   MHC conservation and high heterozygosity in recent hybrids

High levels of polymorphisms at the major histocompatibility complex (MHC) are typically favored
within species, and the locus is prone to duplications [33]. Consequently, assembling the MHC
regions can be challenging [see *e.g.,* challenges for human MHC assembly [34–36]]. Here, we report
a contiguous assembly of the Alpine ibex MHC. High-level synteny is well-conserved among
cattle, sheep, domestic goat, and Alpine ibex in both their main MHC regions (class I and II, Figure
2A). We detected a ~110 kb translocation between Alpine ibex and the domestic goat near the
MHC class II cluster that does not affect MHC coding sequences (Figure 2B). An additional
translocation of ~1.92 Mb occurred further away from the MHC class II on chromosome 23 (Figure
2B). In contrast to the human MHC, both the domestic goat and Alpine ibex MHC class II clusters
encode DR- and DQ-genes but no DP-genes [37,38]. The DR- and DQ-genes are generally
exceptionally polymorphic. Over 100 DRB1 alleles for sheep and dozens for goats have been
deposited in GenBank, reflecting their crucial role in immune diversity. Alpine ibex populations
have previously been shown to carry introgressed alleles from the domestic goat at the DRB gene
[30] and other immune-related genes [29].

To assess whether hybridization load may pose a risk to Alpine ibex, we generated whole-genome

sequencing data of eight recent hybrid individuals sampled from two hybrid swarms in Northern

Italy [24] and four domestic goats representing the local breed, and collected in the same areas as

the hybrids (Table S7). We analysed the new sequencing data together with publicly available

data representing 29 non-hybrid Alpine ibex as well as 18 domestic goats. Average individual

heterozygosity was consistently low in non-hybrid Alpine ibex individuals compared to hybrids

and domestic goats, including particularly striking contrasts in the DR- and DQ-gene regions

(25.40-25.75 Mb; Figure 2C). Diversity at the MHC is associated with disease resistance in a

number of species [39–45], including Alpine ibex [46]. The high diversity at the MHC in hybrids may

therefore confer a selective advantage under disease pressure. Hence, hybridization and MHC

introgression may be favored in the wild, posing a risk of disrupting local adaptation in wild Alpine

ibex.

182
Figure 2: Conserved synteny at the MHC and high diversity in hybrids. A) Close-up of the synteny analysis, conducted with GENESPACE, highlighting two MHC regions within Alpine ibex chromosome 23 (light blue and orange, respectively).

B) Structural variation along chromosome 23 based on SyRI, including annotations of domestic goat (top) and Alpine ibex (bottom) in the range encompassing the major MHC II region.

C) Average individual heterozygosity is shown as observed heterozygosity in Alpine ibex (blue), domestic goat (orange), and hybrids (purple). Heterozygosity excess and deficit in hybrids are shown with green (positive values) and orange (negative values), respectively. The grey box highlights a region corresponding to 25.47-25.55 Mb (encoding *DQA* and *DQB*), which showed a

local lack of coverage in domestic goats while being evenly covered in Alpine ibex (except for
Alpine ibex individuals with previously reported historic introgression in the MHC region, see
Figures S5-11 and Grossen et al. [30]).

Emerging recombinant haplotypes in an Alpine ibex hybrid swarm
The recent observation of hybrid swarms in Northern Italy renewed concerns about Alpine ibex
conservation efforts [24]. The high collinearity of Alpine ibex and domestic goat genomes should
facilitate recombination between haplotypes. At the phenotypic level, recent hybrids carry well-
recognizable domestic goat features such as horn shape and fur coloration [24] (Figure 3A, B). We
analyzed the whole-genome sequencing data to recapitulate introgression patterns in two hybrid
swarms. We furthermore sequenced four domestic goat individuals from the same two regions
(Figure 3, Table S7). Non-hybrid individuals of each species showed clear separation based on
a principal component analysis (PCA, n = 29 Alpine ibex, n = 22 domestic goats; Figure 3C)
performed on 86,650 biallelic SNPs. All eight hybrid swarm members showed clear evidence for
recent admixture (Figure 3C). Next, we performed chromosome painting using genome-wide
polymorphism of the hybrids to identify parental contributions. Individual 12 matched expectations
for a F1 hybrid with uninterrupted goat and Alpine ibex haplotypes over all chromosomes (Figure
3D). Hybrids that underwent backcrosses revealed evidence for recombination events as
interrupted haplotypes (Figure 3D). Genome-wide ancestry matched well with recombination
patterns and parental haplotype contributions. Individual 5 showed both the highest degree of
ancestry and the largest haplotypes of domestic goat. In contrast, individual 8 carried the longest
Alpine ibex haplotype blocks consistent with its strong Alpine ibex ancestry (Figure 3D). The
highly diverse set of recombinants in the Alpine ibex swarm raises significant concerns for further
introgression in the two respective regions and other populations with a high incidence of contacts
between domestic goat and Alpine ibex.

Figure 3

Figure 3: Genome-wide patterns of hybridization

A) Picture of hybrid individual 10. B) Picture of hybrid individual 8. C) Principal Component Analysis of biallelic autosomal SNPs showing the clustering of species and the distribution of hybrids along the conjunction between Alpine ibex and domestic goat/bezoar. D) Chromosome

223   paintings drawn with Mosaic, ordered according to the proportion of domestic goat ancestry
224   (decreasing proportion of goat genome shown in orange from left to right).
225
Hybridization load in two Alpine ibex hybrid swarms
We assessed the extent of hybridization load, which could be introduced into Alpine ibex through
a hybrid swarm. The cross-species polymorphism included 59,544,737 SNPs of which 0.009%,
0.311%, and 0.499% were classified as having putatively high, moderate, and low impacts on
encoded proteins, respectively (see also Tables S8 and S9). High-impact variants, such as those
causing premature stop codons or frameshifts, are predicted to severely disrupt protein function
(*e.g.*, loss-of-function, LOF variants). Moderate and low impact variants cover, respectively,
missense mutations that may alter protein structure without severe impairment and synonymous
changes with minimal expected effects on protein function. To evaluate the robustness of
predicted high-impact mutations, we tested whether these mutations were disproportionately
located in potential pseudogenes. However, we found that the presence of deleterious mutations
cannot be explained by pseudogene-like regions alone (Table S10).
238
We investigated deleterious mutations derived from either domestic goat or Alpine ibex as a proxy
for genetic load (Figure 4A). First, we analyzed candidate mutations underpinning heterosis in
hybrids. We defined these mutations as fixed derived LOF variants in Alpine ibex (fixed for the
ancestral allele in all other species). Seven LOF variants were identified as candidates for
heterosis (Table S11). Severely disruptive variants such as LOF are generally recessive; hence,
these should be masked in the heterozygous state, conferring increased fitness (heterosis) in
hybrids compared to non-hybrid Alpine ibex. We found that the analyzed hybrids each carried
between zero and all seven candidates for heterosis in the heterozygous state (in a backcrossed
individual into Alpine ibex and a F1 hybrid, respectively, Figure 4B). This suggests that the positive
effects of masking LOF variants in hybrids may quickly get lost through backcrossing. The
observed signs of hybrid vigour in Alpine ibex-domestic goat hybrids [24] may reflect a combination
of such masking effects alongside other genetic (*e.g.*, overdominance) or epigenetic factors [47].
251
252

Figure 4: Deleterious mutation load in hybrids

A) Phylogenetic tree showing species relationships and the definition of "goat-load" and "ibex-load" defined as newly derived mutations private to the respective species. B) Counts of loss-of-function alleles found in the heterozygous state for each sequenced hybrid individual (ordered according to proportion of goat). C) Total counts of loss-of-function alleles standardized by the total counts of modifier alleles. Box plots per species. Local goats refer to domestic goat breeds kept in the region where hybrid swarms were detected. See also Figure S12. D) Masked loss-of-function load (heterozygote counts) per individual. Local goats, as in C.

We also investigated the extent to which hybridization load could affect Alpine ibex. We found that the mutation load for LOF variants (standardized by neutral variation) was higher in domestic goat than in Alpine ibex (Figure 4C, $p < 10^{-9}$). As a consequence, hybrids carried a total of 215 newly acquired LOF variants (Table S11), more than doubling the total number of LOF variants

in Alpine ibex (n = 177 LOF variants detected among 29 non-hybrid Alpine ibex). With an average of 76 masked LOF variants per hybrid individual compared to 29 in non-hybrid Alpine ibex, hybrids carried on average a 2.5 times higher severe masked load (Figure 4D). Hybrids backcrossed with Alpine ibex saw their mutation load decreased with each generation as expected (Figure 4D). Yet standardized LOF load did not decrease over the first few generations of back-crossing into Alpine ibex (Figure S12). Hence, there is no indication that selection acted to reduce mutation load. Such counterselection is expected to face less interference after further back-crossings (and recombination). Since the hybrid swarm acquired hybridization load (*i.e.,* masked mutation load), our data highlights that Alpine ibex are at risk of increasing their mutation load through leakage from the hybrid swarm.

Concluding remarks

The detrimental effects of deleterious variants are well-recognized as a risk factor in endangered species [2,3]. While a large number of recent studies investigated the evolution of deleterious mutation load as a consequence of small population size[4–6,48–52], much less consideration was given to the source of deleterious mutation load. Here, we investigated the role of hybridization with domestic goat as a source of mutation load for Alpine ibex. The high heterozygosity in hybrids may confer early-generation hybrids an advantage over non-hybrid individuals, compounding the risk of further introgression and a potential loss of local adaptation. Here, we show that high rates of hybridization can introduce deleterious mutations from a closely related species without evident counterselection. Given the increased rate of human-induced hybridization [53], tracking the deleterious effects of hybridizations should be a crucial element of species conservation actions.

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to, and will be fulfilled by, the lead contact, Christine Grossen (Christine.Grossen@wsl.ch).

Materials availability

This study did not generate novel biological materials.

Data and code availability

Raw sequencing experiment data were submitted to the SRA archive under the project identifier

PRJNA1104824 (reviewer link:

https://dataview.ncbi.nlm.nih.gov/object/PRJNA1104824?reviewer=kij4fipl6g0eheslrgertb0ak9).

The final *C. ibex* genome assembly and high-confidence coding gene annotations will be publicly

available in the NCBI repository XXXXXX (currently undergoing NCBI processing). The

CapIbe1.0 genome fasta file, full transcriptome models, gene functional annotation, and GERP

scores relative to the novel genome can be downloaded from Zenodo

(https://doi.org/10.5281/zenodo.15632776; https://doi.org/10.5281/zenodo.15632854;

https://doi.org/10.5281/zenodo.15357984).

This paper does not report original code. The lead contact can provide any additional information

required to reanalyze the data reported in this paper upon request.

Acknowledgments

For domestic goat sampling, we thank Stefania Zanet, Alice Naudin, and Emilio Gugliermetti. For

hybrid sampling, we thank the Surveillance Service of Gran Paradiso National Park, the Forestry

Service of Ragione Valle d'Aosta, Claudio Scaini, and the personnel of Città Metropolitana di

Torino and of CATO4, Noel Zehnder and Luca Rossi. We thank Sara Gottardo, Francesca

Groppo, and the sequencing facility of CMP[3]VdA in Aosta for technical assistance with Illumina

whole-genome sequencing. We thank Sergio Decherchi, Alessandro Parodi, and Mattia Pini for

their support for high-performance computing and data storage, and Dario de Siena and Raffaele

Turvani for the beautiful ibex pictures.

F. Gözde Çilingir was funded by the Swiss National Science Foundation (IZCOZ0_198147 to A.B.

and C.G. and 31003A_182343 to C.G.).

This work was supported by the 5000genomi@VdA Project co-founded by "Fondo Europeo di

Sviluppo Regionale (FESR CUPB68H19005520007) to SG and AC; FL, ST, ACa, MV and LP

were supported by "Fondo Europeo di Sviluppo Regionale (FESR CUPB68H19005520007); DC

and FF were recipient of a fellowship from Programma Investimenti per la crescita e l'occupazione

2014/20" (European Social Fund, ESF CUP B65F19001200009) and continued their activity with

a fellowship from Programma Regionale Fondo Sociale Europeo Plus 2021/2027 ("European

Social Fund", ESF CUPJ51B24000170002).

This work is part of the "Technologies for Sustainability" Flagship program of IIT (L.P.).

Author contributions

AB conceived the study and managed DNA sample collection. AB, AP, GC, DW, JHM, and BB

performed and coordinated the sampling. FGÇ, FL, DC, LP, and CG designed experiments and

analyses. EH, YCCDS, and DV generated sequencing data under the supervision of FGÇ, LP,

MV, and ST. FGÇ, FL, FF, LP, and CG performed analyses with input from DC. FGÇ, FL, AB,

DC, LP, and CG wrote the manuscript with contributions from all authors. JMC, ACa, SG, AB, and

CG provided funding and resources for the study.

Declaration of interests

340   The authors declare no competing interests.

Supplemental information

Document S1 contains Figures S1–S12.

Figure S1: HiC contact map showing the data used for genome scaffolding, related to Figure 1.

Figure S2: Scheme of the Alpine ibex mitochondrial genome, related to Figure 1.

Figure S3: Structural variant calling between domestic goat and Alpine ibex (based on SyRI)

across all 29 autosomes, related to Figure 1.

Figure S4: Major translocations between domestic goat and Alpine ibex observed on

chromosomes 7 and 18 based on SyRI. B) Total counts and cumulative sizes of the structural

variants identified by SyRI between the Alpine ibex and domestic goat genomes. C) Size

distribution of translocations shown as a histogram from the same variants as in A (INDEL:

insertion/deletion, INV: inversion, DUP: duplication, TRANS: translocation, HDR: highly divergent

region),   related   to   Figure   1.

Figure S5: Read depth profiles of already available whole genome sequencing samples of Alpine

ibex (N=29), domestic goat (N=18), Iberian ibex (N=4), and 12 newly sequenced individual

samples (domestic goat, N=4; hybrids, N=8). The profiles are displayed for the genomic interval

25.47-25.55 Mb on chromosome 23, which includes portions of *DQA* and *DQB* loci, related to

Figure 2.

Figure S6: Figure S5 continued, related to Figure 2.

Figure S7: Figure S5 continued, related to Figure 2.

Figure S8: Figure S5 continued, related to Figure 2.

Figure S9: Figure S5 continued, related to Figure 2.

Figure S10: Figure S5 continued, related to Figure 2.

Figure S11: Figure S5 continued, related to Figure 2.

Figure S12: Total counts of loss-of-function alleles standardized by the total counts of modifier
alleles for each hybrid individual ordered by proportion of domestic goat (indicated in
parentheses), related to Figure 4.
368
Document S2 contains Tables S1-S13.
Table S1: Genome contiguity statistics of the draft and scaffolded Alpine ibex assemblies, related
to Figure 1.
Table S2: Summary of repeat annotations, related to Figure 1.
Table S3: BUSCO statistics for the scaffolded assembly and the protein-coding gene annotation
of Alpine ibex, related to Figure 1.
Table S4: Genome annotation stats, related to Figure 1.
Table S5: Syntenic blocks identified in the GENESPACE analysis, related to Figure 1.
Table S6: Gene orthogroups identified in the GENESPACE analysis, related to Figure 1.
Table S7: Goat and goat/ibex hybrid individuals sequenced in the present study, related to
Figures 2-4.
Table S8: Number of effects of genetic variants by type, related to Figure 4.
Table S9: Number of effects of genetic variants by region, related to Figure 4.
Table S10: Summary of deleterious mutations' PFAM annotations, related to Figure 4.
Table S11: Newly acquired loss-of-function (LOF) variants and candidates for heterosis, related
to Figure 4.
Table S12: Summary of the ONT runs used for genome assembly, related to Figure 1.
Table S13: Species and IDs of the available datasets used in this study, related to Figures 3 and
4.
388
389 **Document S3** Supplementary Text
390
Figure legends
Figure 1: **Sampling and genome description.** A) Picture of a female Alpine ibex with her
offspring (photo by Dario de Siena, Archivio PNGP). B) Map of species distribution and zoom in
on the sampling location of Odissea (shown in yellow), the hybrids (green), and domestic goats
(blue). C) Snail plot visualizing assembly statistics: scaffold statistics, sequence composition
proportions, and BUSCO completeness scores. D) Riparian plot showing the results of the gene
synteny analysis conducted on the cattle, sheep, domestic goat, and Alpine ibex genomes using
GENESPACE. The human genome was used as an outgroup. See also Figures S1-S4.

Figure 2: Conserved synteny at the MHC and high diversity in hybrids. A) Close-up of the synteny analysis, conducted with GENESPACE, highlighting two MHC regions within Alpine ibex chromosome 23 (light blue and orange, respectively).

B) Structural variation along chromosome 23 based on SyRI, including annotations of domestic goat (top) and Alpine ibex (bottom) in the range encompassing the major MHC II region.

C) Average individual heterozygosity is shown as observed heterozygosity in Alpine ibex (blue), domestic goat (orange), and hybrids (purple). Heterozygosity excess and deficit in hybrids were shown with green (positive values) and orange (negative values), respectively. The grey box highlights a region corresponding to 25.47-25.55 Mb (encoding *DQA* and *DQB*), which showed a local lack of coverage in domestic goats while being evenly covered in Alpine ibex (except for Alpine ibex individuals with previously reported historic introgression in the MHC region, see Figures S5-11 and Grossen et al. [30]).

Figure 3: Genome-wide patterns of hybridization

A) Picture of hybrid individual 10. B) Picture of hybrid individual 8. C) Principal Component Analysis of biallelic autosomal SNPs showing the clustering of species and the distribution of hybrids along the conjunction between Alpine ibex and domestic goat/bezoar. D) Chromosome paintings drawn with Mosaic, ordered according to the proportion of domestic goat ancestry (decreasing proportion of goat genome shown in orange from left to right).

Figure 4: Deleterious mutation load in hybrids

A) Phylogenetic tree showing species relationships and the definition of "goat-load" and "ibex-load" defined as newly derived mutations private to the respective species. B) Counts of loss-of-function alleles found in the heterozygous state for each sequenced hybrid individual (ordered according to proportion of goat). C) Total counts of loss-of-function alleles standardized by the total counts of modifier alleles. Box plots per species. Local domestic goat refers to breeds kept in the region where hybrid swarms were detected. See also Figure S12. D) Masked loss-of-function load (heterozygote counts) per individual. New goats, as in C.

Methods

We collected fresh samples from thirteen individuals: one female Alpine ibex (Odissea), eight

hybrids, and four domestic goats. For the female Alpine ibex, we extracted multiple organs and

performed high molecular weight DNA extractions to enable Oxford Nanopore Technologies

(ONT; Oxford, UK) long-read (including ultra-long) and Illumina NovaSeq 6000 short-read

sequencing (Table S12). These data were used to produce a draft genome assembly, which was

then integrated with Hi-C sequencing data from the same individual's liver. The samples from the

other individuals were prepared for whole-genome paired-end sequencing on an Illumina

NovaSeq 6000. Detailed protocols are provided in the Supplementary Text.

Genome   assembly   and   evaluation

We assembled the ONT data into a draft genome using Flye v2.9 [54], after which short-read data

were aligned and used for polishing with Pilon v1.23 [55]. We scanned the assembly for laboratory

or environmental contaminants using the NCBI Foreign Contamination Screen tool [56]. Following

quality filtering, we mapped Hi-C reads following the Arima Genomics pipeline

(https://github.com/ArimaGenomics/mapping_pipeline) and performed scaffolding with YaHS

v1.1 [57]. Finally, we evaluated the final assembly for contiguity, k-mer completeness, reference-

free QV, and expected gene content (further methodological details are in the Supplementary

Text). Genome assembly statistics were visualized with a snail plot in BlobToolKit v4.3.2 [58] (Figure

1C).

Gene   model   prediction,   curation,   and   functional   annotation

We softmasked the Alpine ibex genome by merging repeat coordinates identified via

RepeatModeler v2.0.3 [59] and TRF [60]. Then we incorporated RNA-seq and homology-based

evidence into the braker2 v2.1.6 pipeline [61–69] using pre-trained parameter sets for human (*Homo*

sapiens), which is the evolutionarily closest taxon for Alpine ibex available within the software.

Subsequently, an annotation lift-over from the sheep reference genome [ARS-UI_Ramb_v2.0,

NCBI Genome GCF_016772045.1 [70,71]] was conducted with Liftoff v1.6.3 [72]. In order to reduce

the redundancy of the transcript set obtained, we employed an isoform pruning pipeline as

previously described [73]. Finally, gene models were annotated using eggNOG mapper v2.1.11 [74]

in DIAMOND-search mode. The details of used external evidence for gene prediction and the

detailed methodology can be found in the Supplementary Text. Functional annotation of the gene

models showed that 54,205 out of 61,420 (88.2%) transcripts had matches in the eggNOG protein

database (corresponding to 81.1% of genes).

The mitochondrial genome encodes 13 proteins, including seven complex I proteins (ND1-3, ND4L, ND4-6), four complex IV proteins (COXI, COXII, COXIII, and CytB), and 2 ATP synthase subunits (ATPase 6 and 8; Figure S2).

Synteny analysis and structural rearrangement identification

We ran GENESPACE v1.2.3 [75] with default parameters to examine synteny among the Alpine ibex genome and the human (GRCh38.p14), cattle (ARS-UCD1.3), sheep (ARS-UI_Ramb_v2.0), and domestic goat (ARS1.2, GCF_001704415.2) references, focusing on primary chromosome scaffolds. We further explored syntenic regions and structural rearrangements between domestic goat and Alpine ibex reference genomes with SyRI v1.6.3 [76] (see Supplementary Text). Results were visualized using plotsr v1.1.1 [77].

SNP genotyping and hybridization analyses

Whole-genome sequencing data from eight newly sequenced hybrids and goats were combined with available datasets for Alpine ibex (N=29), domestic goat (N=18), bezoar (N=4), Iberian ibex (N=4), Nubian ibex (N=2), Siberian ibex (N=2), and markhor (N=1), plus four outgroup samples (chamois, cattle, sheep, bighorn sheep; see Table S13) After alignment to the new Alpine ibex genome following the *germline* pipeline of the NVIDIA Clara Parabricks v3.8 [78], variants were called, and hard filtering was performed with GATK v4.1.0 [79]. Eventually, we retained SNPs polymorphic among Capra spp. (see Supplementary Text for selection criteria), which were autosomal, bi-allelic, with a minimal genotype quality of 20 and a minimal genotyping rate of 80% using bcftools v1.2 [80]. We estimated average per-site heterozygosity with VCFtools [81] and used PLINK v2.0 [82] for principal component analysis. After data phasing using beagle v22Jul22.46e [83], the Mosaic R-toolbox [84] was employed to examine Alpine ibex and goat haplotypes in hybrids; chromosome painting was visualized via the karyogram function. Finally, putative functional changes were annotated with SnpEff v4.3t [85]. Individual (derived) genetic load was computed as masked LOF load (number of heterozygotes) or total load (number of heterozygotes + 2* number of homozygotes) and standardized by the respective numbers of MODIFIER genotypes. These standardized estimates of LOF load in domestic goat vs. Alpine ibex were then compared using a Welch two-sample t-test (see also Supplementary Text).

References

1.   Bongaarts, J. (2019). IPBES, 2019. Summary for policymakers of the global assessment

report on biodiversity and ecosystem services of the Intergovernmental Science-Policy
Platform on Biodiversity and Ecosystem Services. Preprint,
https://doi.org/10.1111/padr.12283 https://doi.org/10.1111/padr.12283.

2. Dussex, N., Morales, H.E., Grossen, C., Dalén, L., and van Oosterhout, C. (2023). Purging
and accumulation of genetic load in conservation. Trends Ecol. Evol. *38*, 961–969.

3. Robinson, J., Kyriazis, C.C., Yuan, S.C., and Lohmueller, K.E. (2023). Deleterious variation
in natural populations and implications for conservation genetics. Annu. Rev. Anim. Biosci.
11, 93–114.

4. Xue, Y., Prado-Martinez, J., Sudmant, P.H., Narasimhan, V., Ayub, Q., Szpak, M.,
Frandsen, P., Chen, Y., Yngvadottir, B., Cooper, D.N., et al. (2015). Mountain gorilla
genomes reveal the impact of long-term population decline and inbreeding. Science *348*,
242–245.

5. Grossen, C., Guillaume, F., Keller, L.F., and Croll, D. (2020). Purging of highly deleterious
mutations through severe bottlenecks in Alpine ibex. Nat. Commun. *11*, 1001.

6. Dussex, N., van der Valk, T., Morales, H.E., Wheat, C.W., Díez-Del-Molino, D., von Seth,
J., Foster, Y., Kutschera, V.E., Guschanski, K., Rhie, A., et al. (2021). Population genomics
of the critically endangered kākāpō. Cell Genom. *1*, 100002.

7. Whitlock, M.C., and Davis, B. (2011). Genetic Load. eLS.
https://doi.org/10.1002/9780470015902.a0001787.pub2.

8. Kardos, M., Zhang, Y., Parsons, K.M., A, Y., Kang, H., Xu, X., Liu, X., Matkin, C.O., Zhang,
P., Ward, E.J., et al. (2023). Inbreeding depression explains killer whale population
dynamics. Nat. Ecol. Evol. *7*, 675–686.

9. Stoffel, M.A., Johnston, S.E., Pilkington, J.G., and Pemberton, J.M. (2024). Purifying and
balancing selection on embryonic semi-lethal haplotypes in a wild mammal. Evol. Lett. *8*,
222–230.

10. Genovart, M. (2009). Natural hybridization and conservation. Biodivers. Conserv. *18*, 1435–
1439.

11. Mallet, J. (2005). Hybridization as an invasion of the genome. Trends Ecol. Evol. *20*, 229–
237.

12. Moran, B.M., Payne, C., Langdon, Q., Powell, D.L., Brandvain, Y., and Schumer, M. (2021).
The genomic consequences of hybridization. Elife *10*. https://doi.org/10.7554/eLife.69016.

13. Barton, N.H. (2001). The role of hybridization in evolution. Mol. Ecol. *10*, 551–568.

14. Burke, J.M., and Arnold, M.L. (2001). Genetics and the fitness of hybrids. Annu. Rev.
Genet. *35*, 31–52.

15. Allendorf, F.W., Leary, R.F., Spruell, P., and Wenburg, J.K. (2001). The problems with
hybrids: setting conservation guidelines. Trends Ecol. Evol. *16*, 613–622.

16. Leitwein, M., Cayuela, H., Ferchaud, A.-L., Normandeau, É., Gagnaire, P.-A., and

Bernatchez, L. (2019). The role of recombination on genome-wide patterns of local
ancestry exemplified by supplemented brook charr populations. Mol. Ecol. *28*, 4755–4769.

17. Schumer, M., Xu, C., Powell, D.L., Durvasula, A., Skov, L., Holland, C., Blazier, J.C.,
Sankararaman, S., Andolfatto, P., Rosenthal, G.G., et al. (2018). Natural selection interacts
with recombination to shape the evolution of hybrid genomes. Science *360*, 656–660.

18. Simon, M.F., Mendoza Flores, J.M., Liu, H.-L., Martins, M.L.L., Drovetski, S.V.,
Przelomska, N.A.S., Loiselle, H., Cavalcanti, T.B., Inglis, P.W., Mueller, N.G., et al. (2022).
Phylogenomic analysis points to a South American origin of Manihot and illuminates the
primary gene pool of cassava. New Phytol. *233*, 534–545.

19. Wang, L., Beissinger, T.M., Lorant, A., Ross-Ibarra, C., Ross-Ibarra, J., and Hufford, M.B.
(2017). The interplay of demography and selection during maize domestication and
expansion. Genome Biol. *18*, 215.

20. Janzen, G.M., Wang, L., and Hufford, M.B. (2019). The extent of adaptive wild introgression
in crops. New Phytol. *221*, 1279–1288.

21. Liu, S., Zhang, L., Sang, Y., Lai, Q., Zhang, X., Jia, C., Long, Z., Wu, J., Ma, T., Mao, K., et
al. (2022). Demographic history and natural selection shape patterns of deleterious
mutation load and barriers to introgression across Populus genome. Mol. Biol. Evol. *39*.
https://doi.org/10.1093/molbev/msac008.

22. Theissinger, K., Fernandes, C., Formenti, G., Bista, I., Berg, P.R., Bleidorn, C., Bombarely,
A., Crottini, A., Gallo, G.R., Godoy, J.A., et al. (2023). How genomics can help biodiversity
conservation. Trends Genet. *39*, 545–559.

23. Stüwe, M., and Nievergelt, B. (1991). Recovery of alpine ibex from near extinction: the
result of effective protection, captive breeding, and reintroductions. Appl. Anim. Behav. Sci.
29, 379–387.

24. Brambilla, A., Zehnder, N., Bassano, B., Rossi, L., and Grossen, C. (2024). Genetic
evidence of a hybrid swarm between Alpine ibex (Capra ibex) and domestic goat (C.
hircus). Evol. Appl. *17*, e13761.

25. Brambilla, A., Von Hardenberg, A., Nelli, L., and Bassano, B. (2020). Distribution, status,
and recent population dynamics of Alpine ibex Capra ibex in Europe. Mamm. Rev. *50*, 267–
277.

26. Biebach, I., and Keller, L.F. (2010). Inbreeding in reintroduced populations: the effects of
early reintroduction history and contemporary processes. Conserv. Genet. *11*, 527–538.

27. Brambilla, A., Biebach, I., Bassano, B., Bogliani, G., and von Hardenberg, A. (2015). Direct
and indirect causal effects of heterozygosity on fitness-related traits in Alpine ibex. Proc.
Biol. Sci. *282*, 20141873.

28. Bozzuto, C., Biebach, I., Muff, S., Ives, A.R., and Keller, L.F. (2019). Inbreeding reduces
long-term growth of Alpine ibex populations. Nat Ecol Evol *3*, 1359–1364.

29. Münger, X., Robin, M., Dalén, L., and Grossen, C. (2024). Facilitated introgression from
domestic goat into Alpine ibex at immune loci. Mol. Ecol., e17429.

30. Grossen, C., Keller, L., Biebach, I., International Goat Genome Consortium, and Croll, D.
(2014). Introgression from domestic goat generated variation at the major histocompatibility
complex of Alpine ibex. PLoS Genet. *10*, e1004438.

31. Moroni, B., Brambilla, A., Rossi, L., Meneguz, P.G., Bassano, B., and Tizzani, P. (2022).
Hybridization between Alpine Ibex and Domestic Goat in the Alps: A Sporadic and
Localized Phenomenon? Animals (Basel) *12*. https://doi.org/10.3390/ani12060751.

32. Bickhart, D.M., Rosen, B.D., Koren, S., Sayre, B.L., Hastie, A.R., Chan, S., Lee, J., Lam,
E.T., Liachko, I., Sullivan, S.T., et al. (2017). Single-molecule sequencing and chromatin
conformation capture enable de novo reference assembly of the domestic goat genome.
Nat. Genet. *49*, 643–650.

33. Piertney, S.B., and Oliver, M.K. (2006). The evolutionary ecology of the major
histocompatibility complex. Heredity *96*, 7–21.

34. Jain, M., Koren, S., Miga, K.H., Quick, J., Rand, A.C., Sasani, T.A., Tyson, J.R., Beggs,
A.D., Dilthey, A.T., Fiddes, I.T., et al. (2018). Nanopore sequencing and assembly of a
human genome with ultra-long reads. Nat. Biotechnol. *36*, 338–345.

35. Koren, S., Rhie, A., Walenz, B.P., Dilthey, A.T., Bickhart, D.M., Kingan, S.B., Hiendleder,
S., Williams, J.L., Smith, T.P.L., and Phillippy, A.M. (2018). De novo assembly of
haplotype-resolved genomes with trio binning. Nat. Biotechnol.
https://doi.org/10.1038/nbt.4277.

36. Chin, C.-S., Wagner, J., Zeng, Q., Garrison, E., Garg, S., Fungtammasan, A., Rautiainen,
M., Aganezov, S., Kirsche, M., Zarate, S., et al. (2020). A diploid assembly-based
benchmark for variants in the major histocompatibility complex. Nat. Commun. *11*, 4794.

37. Beck, S., and Trowsdale, J. (1999). Sequence organisation of the class II region of the
human MHC. Immunol. Rev. *167*, 201–210.

38. Kelley, J., Walter, L., and Trowsdale, J. (2005). Comparative genomics of major
histocompatibility complexes. Immunogenetics *56*, 683–695.

39. Gowane, G., Vandre, R., Nangre, M., and Sharma, A. (2014). Major histocompatibility
complex (MHC) of bovines: an insight into infectious disease resistance. Livestock
Research International, *1*, 46–57.

40. Westerdahl, H. (2007). Passerine MHC: genetic variation and disease resistance in the
wild. J. Ornithol. *148*, 469–477.

41. Hedrick, P.W., Kim, T.J., and Parker, K.M. (2001). Parasite resistance and genetic variation
in the endangered Gila topminnow. Animal Conservation forum *4*, 103–109.

42. Hedrick, P.W., Parker, K.M., Miller, E.L., and Miller, P.S. (1999). Major histocompatibility
complex variation in the endangered Przewalski's horse. Genetics *152*, 1701–1710.

43. Paterson, S., Wilson, K., and Pemberton, J.M. (1998). Major histocompatibility complex
variation associated with juvenile survival and parasite resistance in a large unmanaged
ungulate population (Ovis aries L.). Proceedings of the National Academy of Sciences *95*,
3714–3719.

44. Langefors, A., Lohm, J., Grahn, M., Andersen, O., and von Schantz, T. (2001). Association
between major histocompatibility complex class IIB alleles and resistance to Aeromonas
salmonicida in Atlantic salmon. Proc. Biol. Sci. *268*, 479–485.

45. Fu, M., Eimes, J.A., Kong, S., Lamichhaney, S., and Waldman, B. (2023). Identification of
major histocompatibility complex genotypes associated with resistance to an amphibian
emerging infectious disease. Infect. Genet. Evol. *113*, 105470.

46. Brambilla, A., Keller, L., Bassano, B., and Grossen, C. (2018). Heterozygosity-fitness
correlation at the major histocompatibility complex despite low variation in Alpine ibex (
Capra ibex). Evol. Appl. *11*, 631–644.

47. Chen, Z.J. (2013). Genomic and epigenetic insights into the molecular bases of heterosis.
Nat. Rev. Genet. *14*, 471–482.

48. Garcia-Erill, G., Liu, S., Le, M.D., Hurley, M.M., Nguyen, H.D., Nguyen, D.Q., Nguyen, D.H.,
Santander, C.G., Sánchez Barreiro, F., Gomes Martins, N.F., et al. (2025). Genomes of
critically endangered saola are shaped by population structure and purging. Cell.
https://doi.org/10.1016/j.cell.2025.03.040.

49. Henn, B.M., Botigué, L.R., Peischl, S., Dupanloup, I., Lipatov, M., Maples, B.K., Martin,
A.R., Musharoff, S., Cann, H., Snyder, M.P., et al. (2016). Distance from sub-Saharan
Africa predicts mutational load in diverse human genomes. Proc. Natl. Acad. Sci. U. S. A.
113, E440–E449.

50. Khan, A., Patel, K., Shukla, H., Viswanathan, A., van der Valk, T., Borthakur, U., Nigam, P.,
Zachariah, A., Jhala, Y., Kardos, M., et al. (2021). Genomic evidence for inbreeding
depression and purging of deleterious genetic variation in Indian tigers. bioRxiv.
https://doi.org/10.1101/2021.05.18.444660.

51. Ochoa, A., and Gibbs, H.L. (2021). Genomic signatures of inbreeding and mutation load in
a threatened rattlesnake. Mol. Ecol. *30*, 5454–5469.

52. Kleinman-Ruiz, D., Lucena-Perez, M., Villanueva, B., Fernández, J., Saveljev, A.P.,
Ratkiewicz, M., Schmidt, K., Galtier, N., García-Dorado, A., and Godoy, J.A. (2022).
Purging of deleterious burden in the endangered Iberian lynx. Proc. Natl. Acad. Sci. U. S.
A. *119*, e2110614119.

53. Ottenburghs, J. (2021). The genic view of hybridization in the Anthropocene. Evol. Appl. *14*,
2342–2360.

54. Kolmogorov, M., Yuan, J., Lin, Y., and Pevzner, P.A. (2019). Assembly of long, error-prone
reads using repeat graphs. Nat. Biotechnol. *37*, 540–546.

55. Walker, B.J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C.A.,
Zeng, Q., Wortman, J., Young, S.K., et al. (2014). Pilon: an integrated tool for
comprehensive microbial variant detection and genome assembly improvement. PLoS One
9, e112963.

56. Astashyn, A., Tvedte, E.S., Sweeney, D., Sapojnikov, V., Bouk, N., Joukov, V., Mozes, E.,
Strope, P.K., Sylla, P.M., Wagner, L., et al. (2024). Rapid and sensitive detection of
genome contamination at scale with FCS-GX. Genome Biol. *25*, 60.

57. Zhou, C., McCarthy, S.A., and Durbin, R. (2023). YaHS: yet another Hi-C scaffolding tool.
Bioinformatics *39*, btac808.

58. Challis, R., Richards, E., Rajan, J., Cochrane, G., and Blaxter, M. (2020). BlobToolKit--
interactive quality assessment of genome assemblies. G3: Genes, Genomes, Genetics *10*,
1361–1374.

59. Flynn, J.M., Hubley, R., Goubert, C., Rosen, J., Clark, A.G., Feschotte, C., and Smit, A.F.
(2020). RepeatModeler2 for automated genomic discovery of transposable element
families. Proc. Natl. Acad. Sci. U. S. A. *117*, 9451–9457.

60. Benson, G. (1999). Tandem repeats finder: a program to analyze DNA sequences. Nucleic
Acids Res. *27*, 573–580.

61. Brůna, T., Hoff, K.J., Lomsadze, A., Stanke, M., and Borodovsky, M. (2021). BRAKER2:
automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported
by a protein database. NAR Genom Bioinform *3*, lqaa108.

62. Hoff, K.J., Lomsadze, A., Borodovsky, M., and Stanke, M. (2019). Whole-genome
annotation with BRAKER. Methods Mol. Biol. *1962*, 65–95.

63. Hoff, K.J., Lange, S., Lomsadze, A., Borodovsky, M., and Stanke, M. (2016). BRAKER1:
Unsupervised RNA-seq-based genome qnnotation with GeneMark-ET and AUGUSTUS.
Bioinformatics *32*, 767–769.

64. Stanke, M., Diekhans, M., Baertsch, R., and Haussler, D. (2008). Using native and
syntenically mapped cDNA alignments to improve de novo gene finding. Bioinformatics *24*,
637–644.

65. Stanke, M., Schöffmann, O., Morgenstern, B., and Waack, S. (2006). Gene prediction in
eukaryotes with a generalized hidden Markov model that uses hints from external sources.
BMC Bioinformatics *7*, 62.

66. Buchfink, B., Xie, C., and Huson, D.H. (2015). Fast and sensitive protein alignment using
DIAMOND. Nat. Methods *12*, 59–60.

67. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,
G., Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The
Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079.

68. Barnett, D.W., Garrison, E.K., Quinlan, A.R., Strömberg, M.P., and Marth, G.T. (2011).
BamTools: a C++ API and toolkit for analyzing and managing BAM files. Bioinformatics *27*,
1691–1692.

69. Lomsadze, A., Burns, P.D., and Borodovsky, M. (2014). Integration of mapped RNA-Seq
reads into automatic training of eukaryotic gene finding algorithm. Nucleic Acids Res. *42*,
e119.

70. International Sheep Genomics Consortium, Archibald, A.L., Cockett, N.E., Dalrymple, B.P.,
Faraut, T., Kijas, J.W., Maddox, J.F., McEwan, J.C., Hutton Oddy, V., Raadsma, H.W., et
al. (2010). The sheep genome reference sequence: a work in progress. Anim. Genet. *41*,
449–453.

71. Li, R., Yang, P., Li, M., Fang, W., Yue, X., Nanaei, H.A., Gan, S., Du, D., Cai, Y., Dai, X., et
al. (2021). A Hu sheep genome with the first ovine Y chromosome reveal introgression
history after sheep domestication. Sci. China Life Sci. *64*, 1116–1130.

72. Shumate, A., and Salzberg, S.L. (2021). Liftoff: accurate mapping of gene annotations.
Bioinformatics *37*, 1639–1643.

73. Ivankovic, M., Brand, J.N., Pandolfini, L., Brown, T., Pippel, M., Rozanski, A., Schubert, T.,
Grohme, M.A., Winkler, S., Robledillo, L., et al. (2023). A comparative analysis of planarian
genomes reveals regulatory conservation in the face of rapid structural divergence. bioRxiv.
https://doi.org/10.1101/2023.12.22.572568.

74. Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., and Huerta-Cepas, J.
(2021). eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. Mol. Biol. Evol. *38*, 5825–5829.

75. Lovell, J.T., Sreedasyam, A., Schranz, M.E., Wilson, M., Carlson, J.W., Harkess, A., Emms,
D., Goodstein, D.M., and Schmutz, J. (2022). GENESPACE tracks regions of interest and
gene copy number variation across multiple genomes. Elife *11*.
https://doi.org/10.7554/eLife.78526.

76. Goel, M., Sun, H., Jiao, W.-B., and Schneeberger, K. (2019). SyRI: finding genomic
rearrangements and local sequence differences from whole-genome assemblies. Genome
Biol. *20*, 277.

77. Goel, M., and Schneeberger, K. (2022). plotsr: visualizing structural similarities and
rearrangements between multiple genomes. Bioinformatics *38*, 2922–2926.

78. Franke, K.R., and Crowgey, E.L. (2020). Accelerating next generation sequencing data
analysis: an evaluation of optimized best practices for Genome Analysis Toolkit algorithms.
Genomics Inform. *18*, e10.

79. McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A.,
Garimella, K., Altshuler, D., Gabriel, S., Daly, M., et al. (2010). The Genome Analysis
Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.
Genome Res. *20*, 1297–1303.

80. Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,
Keane, T., McCarthy, S.A., Davies, R.M., et al. (2021). Twelve years of SAMtools and
BCFtools. Gigascience *10*, giab008.

81. Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker,
R.E., Lunter, G., Marth, G.T., Sherry, S.T., et al. (2011). The variant call format and
VCFtools. Bioinformatics *27*, 2156–2158.

82. Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., and Lee, J.J. (2015).
Second-generation PLINK: rising to the challenge of larger and richer datasets.
Gigascience *4*, 7.

83. Browning, B.L., Tian, X., Zhou, Y., and Browning, S.R. (2021). Fast two-stage phasing of
large-scale sequence data. Am. J. Hum. Genet. *108*, 1880–1890.

84.   Salter-Townshend, M., and Myers, S. (2019). Fine-Scale Inference of Ancestry Segments
Without Prior Knowledge of Admixing Groups. Genetics *212*, 869–889.

85.   Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X.,
and Ruden, D.M. (2012). A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain
w1118; iso-2; iso-3. Fly *6*, 80–92.