**Magnesium modulates phospholipid metabolism to promote bacterial phenotypic resistance to antibiotics**

Hui Li[1,2,†], Jun Yang[1,2,†], Su-fang Kuang[1], Huan-zhe Fu[1], Hui-ying Lin[1], Bo Peng[1,2,*]

[1]*State Key Laboratory of Bio-Control, School of Life Sciences, Guangdong Province Key Laboratory for Pharmaceutical Functional Genes, Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Sun Yat-sen University, Higher Education Mega Center, Guangzhou 510006, People's Republic of China.*

[2]*Laboratory for Marine Biology and Biotechnology, Marine Fisheries Science and Food Production Processes, Qingdao Marine Science and Technology Center, Qingdao 266071, China.*

[†] These authors contributed equally to this work

_______________________________________

**Running title:** magnesium induces bacterial phenotypic resistance

[*]**Corresponding author:** Bo Peng (email: pengb26@mail.sysu.edu.cn) *State Key Laboratory of Bio-Control, School of Life Sciences, Sun Yat-sen University, Guangzhou 510275, People's Republic of China.*

**Abstract**

Non-inheritable antibiotic or phenotypic resistance ensures bacterial survival during antibiotic treatment. However, exogenous factors promoting phenotypic resistance are poorly defined. Here, we demonstrate that *Vibrio alginolyticus* are recalcitrant to killing by a broad spectrum of antibiotics under high magnesium. Functional metabolomics demonstrated that magnesium modulates fatty acid biosynthesis by increasing saturated fatty acid biosynthesis while decreasing unsaturated fatty acid production. Exogenous supplementation of unsaturated and saturated fatty acids increased and decreased bacterial susceptibility to antibiotics, respectively, confirming the role of fatty acids in antibiotic resistance. Functional lipidomics revealed that glycerophospholipid metabolism is the major metabolic pathway remodeled by magnesium, where phosphatidylethanolamine (PE) biosynthesis is reduced and phosphatidylglycerol (PG) production is increased. This process alters membrane composition, increasing membrane polarization, and decreasing permeability and fluidity, thereby reducing antibiotic uptake by *V. alginolyticus*. These findings suggest the presence of a previously unrecognized metabolic mechanism by which bacteria escape antibiotic killing through the use of an environmental factor.

**Keywords**: magnesium, phenotypic resistance, quinolones, phospholipid metabolism, membrane fluidity.

**Introduction**

Non-inheritable antibiotic or phenotypic resistance represents a serious challenge for

treating bacterial infections. Phenotypic resistance does not involve genetic mutations

Phenotypic resistance does not involve genetic mutations and is transient, allowing

bacteria to resume normal growth. Biofilm and bacterial persisters are two phenotypic

resistance types that have been extensively studied (Brandis *et al.*, 2023; Corona &

Martinez, 2013). Biofilms have complex structures, containing elements that impede

antibiotic diffusion, sequestering and inhibiting their activity (Ciofu *et al.*, 2022).

Biofilm-forming bacteria and persisters also have distinct metabolic states that

significantly reduce their antibiotic susceptibility (Yan & Bassler, 2019). These two types

of phenotypic resistance share the common feature in their retarded or even cease of

growth in the presence of antibiotics (Corona & Martinez, 2013). However, specific

factors that promote phenotypic resistance and allow bacteria to proliferate in the

presence of antibiotics remain poorly defined.

Metal ions have a diverse impact on the chemical, physical, and physiological processes

of antibiotic resistance    (Booth *et al*, 2011; Lu *et al*, 2020; Poole, 2017). This includes

genetic elements that confer resistance to metals and antibiotics (Poole, 2017) and metal

cations that directly hinder (or enhance) the activity of specific antibiotic drugs (Zhang

*et al.*, 2014). The metabolic environment can also impact the sensitivity of bacteria to

antibiotics (Jiang *et al.*, 2023; Lee & Collins, 2012; Peng *et al.*, 2015; Zhang *et al.*,

2020; Zhao *et al.*, 2021). Light metal ions, such as magnesium, sodium, and potassium,

can behave as cofactors for different enzymes (Du *et al.*, 2016) and influence drug

efficacy. Heavy metal ions, including $Cu^{2+}$ and $Zn^{2+}$, confer resistance to antibiotics (Yazdankhah *et al.*, 2014; Zhang *et al.*, 2018). Recent reports suggest that sodium negatively regulates redox states to promote the antibiotic resistance of *Vibrio alginolyticus* (Yang *et al.*, 2018), while actively growing *Bacillus subtilis* cope with ribosome-targeting antibiotics by modulating ion flux (Lee *et al*, 2019). In Gram-negative bacteria, by contrast, zinc enhances antibiotic efficacy by potentiating carbapenem, fluoroquinolone, and β-lactam-mediated killing (Isaei *et al.,* 2016; Zhang *et al.,* 2014). Magnesium influences bacterial structure, cell motility, enzyme function, cell signaling, and pathogenesis (Wang *et al.,* 2019). This mineral also modulates microbiota to harvest energy from the diet (Garcia-Legorreta *et al.,* 2020), allowing *Bacillus subtilis* to cope with ribosome-targeting antibiotics by modulating ion flux (Lee *et al.,* 2019). However, the role of magnesium in promoting phenotypic resistance is less well understood.

Vibrios inhabit seawater, estuaries, bays, and coastal waters, regions full of metal ions such as magnesium (Kumarage *et al.,* 2022). Magnesium is the second most dissolved element in seawater after sodium. At a salinity of 3.5% seawater, the magnesium concentration is about 54 mM (Potis, 1968), and in deep seawater, can be as high as 2,500 mM (Wang *et al.,* 2024). *Vibrio parahaemolyticus* and *V. alginilyticus* are two representative Vibrio pathogens that infect humans and aquatic animals, resulting in illness and economic loss, respectively (Grimes, 2020). (Fluoro)quinolones such as balofloxacin are used to treat Vibrio infection, however, resistance has emerged due to

overuse (Suyamud *et al.,* 2024). Indeed, (fluoro)quinolones are one of China's two

primary residual chemicals associated with aquaculture (Liu *et al.,* 2017). Vibrio can

develop quinolone resistance through mutations in the DNA gyrase gene or through

plasmid-mediated mechanisms (Dutta *et al.,* 2021). Thus, the use of *V.*

*parahaemolyticus* and *V. alginilyticus* as bacterial representatives, and balofloxacin as

a quinolone-based antibacterial representative, can help to define novel magnesium-

dependent phenotypic resistance mechanisms of pathogenic Vibrio species.

The current study evaluated whether magnesium induces phenotypic resistance in

Vibrio species and defined the molecular/genetic basis for this resistance. Genetic

approaches, GC-MS analysis of metabolite and membrane remodeling upon antibiotic

exposure, membrane physiology, and extensive antimicrobial susceptibility testing

were used for the evaluations.

**Results**

**$Mg^{2+}$ promotes phenotypic resistance to antibiotics**

Marine environments and agriculture, where antibiotics are commonly used, are rich in

magnesium. To investigate whether this mineral impacts antibiotic activity, the minimal

inhibitory concentration (MIC) of *V. alginolyticus* ATCC33787 and *V.*

*parahaemolyticus* VP01, which we referred as ATCC33787 and VP01 afterwards,

isolated from marine aquaculture, to balofloxacin (BLFX) in Luria-Bertani medium

(LB medium) plus 3% NaCl as LBS medium and "artificial seawater" (ASWT) medium

that included the major ion species in marine water (Wilson, 1975) (LB medium plus

210 mM NaCl, 35 mM $Mg_2SO_4$, 7 mM KCl, and 7 mM $CaCl_2$) were assessed (**Suppl.**

**Table 1**). The MICs of ATCC33787 to BLFX were 54 and 1 μg/mL in ASWT and LBS

medium, respectively. Similarly, the MICs to VP01 were 25 and 3 μg/mL in ASWT

and LBS medium, respectively (**Fig 1A**). The role of exogenous NaCl in antibiotic

resistance was investigated previously (Yang *et al.,* 2018). The current study found that

the MIC for BLFX was the same in LB and M9 minimal medium (M9 medium) plus 7

mM KCl or 7 mM $CaCl_2$ (**Suppl. Fig 1**). However, the MIC for BLFX was higher in

ASWT medium supplemented with $Mg_2SO_4$ or $MgCl_2$ than in LB medium (**Fig 1B**).

And $Mg_2SO_4$ or $MgCl_2$ had no difference on MIC, suggesting it is $Mg^{2+}$ not other ions

contribute to the MIC change. The MIC for BLFX increased at higher concentrations

of $MgCl_2$ in ASWT medium. Specifically, adding 50 mM or 200 mM $MgCl_2$ increased

the MIC for BLFX by 200- or 1600-fold, respectively (**Fig 1C**). At balofloxacin doses

of 1.56, 3.125, 6.25, and 12.5 μg, the zone of inhibition decreased with increasing

$MgCl_2$ (**Fig 1D**). Exogenous $MgCl_2$ also increased the MICs for other quinolone (*e.g.*

nalidixic acid, levofloxacin, ciprofloxacin, ofloxacin, and moxifloxacin) (**Fig 1E**) and

non-quinolone antibiotics including antibacterial peptides (colistin), macrolides

(roxithromycin), tetracyclines (oxytetraycline), β-lactams (ceftriaxone, ceftazidime),

and aminoglycosides (amikacin, kanamycin, and gentamicin) (**Fig 1F**). Notably,

magnesium had a reduced effect on ceftriaxone and gentamicin than other antibiotics.

Importantly, exogenous $MgCl_2$ also increased MICs of clinic isolates, carbapenem-

resistant *Escherichia coli*, carbapenem-resistant *Klebsiella pneumoniae,* carbapenem-

resistant *Pseudomonas aeruginosa* and carbapenem-resistant *Acinetobacter baumannii*

to balofloxacin (**Fig 1G**). These findings indicate that $Mg^{2+}$ promotes phenotypic

resistance.

**Figure 1. Magnesium promotes bacterial resistance to antibiotics. A.** MIC of

ATCC33787 and *V. parahaemolyticus* to BLFX in ASWT or LB medium as determined

by the microtitre-dilution method. **B.** MIC of ATCC33787 to BLFX in ASWT with

additional $MgSO_4$ or $MgCl_2$ as determined by the microtitre-dilution-method. **C.** MIC

of ATCC33787 to BLFX in ASWT at the indicated concentrations of $MgCl_2$ as

determined by the microtitre-dilution method. **D.** MIC of ATCC33787 to BLFX at the

indicated concentrations of BLFX and $MgCl_2$ as determined by the Oxford cup test. The

numbers 1, 2, 3, 4, 5, 6, and 7 represent 0, 0.39, 0.78, 1.56, 3.125, 6.25, and 12.5 µg

BLFX, respectively. **E.** MIC of ATCC33787 to other quinolones in ASWT at the

indicated concentrations of $MgCl_2$ as determined by the microtitre-dilution method. **F.**

MIC of ATCC 33787 to other classes of antibiotics in ASWT at the indicated

concentrations of $MgCl_2$ as determined by the microtitre-dilution method. **G.** MIC of

carbapenem-resistant *Escherichia coli*, carbapenem-resistant *Klebsiella pneumoniae,*

carbapenem-resistant *Pseudomonas aeruginosa,* and carbapenem-resistant

*Acinetobacter baumannii* isolates to BFLX at the indicated concentrations of $MgCl_2$.

**$MgCl_2$ affects bacterial metabolism**

To investigate how magnesium regulates phenotypic resistance, other effects of

magnesium, including its ability to quench balofloxacin activity and induce efflux

pump expression and LPS biosynthesis, were excluded (**Suppl. Text; Suppl. Fig 2**).

While $Mg^{2+}$ is a co-factor for enzymes (Garfinkel and Garfinkel, 1985), whether this

mineral can promote antibiotic resistance independent of this function remains

unknown.

To better understand how magnesium affects bacterial metabolism, *V. alginolyticus* was

cultured in M9 medium in the presence of various amounts of $MgCl_2$ (0, 0.78, 3.125,

12.5, 50, or 200 mM) and the levels of 54 metabolites were assessed by GC-MS. Five

biological and two technical replicates were evaluated for each treatment (**Suppl. Fig**

**3**). The levels of 41 metabolites were differential ($p < 0.05$) and are presented as Z-

values (**Fig 2A; Suppl. Fig 4**). Orthogonal partial least square discriminant analysis

(OPLS-DA) was conducted that separated the six treatments into three groups: Group

1 included 0 mM, 0.78 mM, and 3.125 mM $MgCl_2$; Group 2 was a singlet of 12.5 mM;

Group 3 included 50 mM and 200 mM $MgCl_2$. Component t[1] distinguishes Group 1

from Groups 2 and 3; component t[2] distinguishes Group 3 from Groups 1 and 2 (**Fig**

**2B**). Discriminating variables are shown as S-plot, where cut-off values are $\geq 0.05$

absolute value of covariance p and 0.5 for correlation p(corr). Six

biomarkers/metabolites were selected from component t[1] and p[1]. More specifically,

the abundance of cadaverine, urea, palmitic acid, aminoethanol, and fumaric acid were

increased, but pyroglutamic acid and glutamic acid were decreased (**Fig 2C; Suppl.**

**Fig 5A**). Pathway enrichment analysis suggests that twelve pathways are involved.

Notably, one of the pathways is biosynthesis of unsaturated fatty acids (terminology in

the software. Actually, it is biosynthesis of fatty acids) (**Fig 2D**), where the abundance

of palmitic acid and stearic acid were increased in an $Mg^{2+}$ dose-dependent manner (**Fig**

**2E; Suppl. Fig 6**). Moreover, palmitic acid was a crucial biomarker (**Suppl. Fig 6**).

The increase in fatty acid biosynthesis could be partially explained by an imbalanced

pyruvate cycle/TCA cycle, in which fumarate levels increased at higher $Mg^{2+}$ while

succinate levels increased at lower $Mg^{2+}$ (**Suppl. Fig 5B**). These findings indicated that

glycolysis fluxes into fatty acid biosynthesis rather than the pyruvate cycle/TCA cycle.

The relevance of fatty acids and BLFX was demonstrated by the observation that

exogenous palmitic acid increased bacterial resistance to balofloxacin (**Fig 2F**). These

results suggest that fatty acid metabolism may be critical to magnesium-based

phenotypic resistance.

**Figure 2. Mg²⁺-induced metabolomic change. A.** Differential metabolomes in the absence or presence of the indicated concentrations of MgCl₂. The yellow and blue colors indicate an increase or decrease in metabolite levels relative to the median

metabolite level, respectively (see color scale). Euclidean distance calculations were

used to generate a heatmap that shows clustering of the biological and technical

replicates of each treatment. **B.** OPLS-DA analysis of different $MgCl_2$-induced

metabolome concentrations. Each dot represents a technical replicate of samples in the

plot. **C.** S-plot generated from OPLS-DA. Predictive component p [1] and correlation

p(corr) [1] differentiate 0, 0.78, 3.125 mM $MgCl_2$ from 12.5, 50, 200 mM $MgCl_2$.

Predictive component p[2] and correlation p(corr)[2] separate 0, 0.78, 50, 200 mM

$MgCl_2$ from 3.125, 12.5 mM $MgCl_2$. The triangle represents metabolites in which

candidate biomarkers are marked. **D.** Enriched pathways by differential abundances of

metabolites. **E.** Areas of the peaks of palmitic acid and stearic acid generated by GC-

MS analysis. **F.** Synergy analysis for BFLX with palmitic acid for *V. alginolyticus*.

Synergy was performed by comparing the dose needed for 50% inhibition of the

synergistic agents (white) and non-synergistic (i.e., additive) agents (purple).

**$Mg^{2+}$ regulates fatty acid biosynthesis**

Acetyl-CoA carboxylase (ACC) catalyzes the conversion of acetyl-CoA to malonyl-

CoA, the first step of fatty acid synthesis. Fatty acid biosynthesis, which includes

saturated and unsaturated forms, shares common biosynthetic pathways with acetyl-

CoA to enoyl-ACP, and is sequentially mediated by *fabD* or *fabH*, *fabB*/*F*, *fabG,* and

*fabA*/*Z*. The enzyme, enoyl-ACP, produces unsaturated fatty acids or is metabolized to

acyl-ACP via *fabV* to produce saturated fatty acids. Unsaturated fatty acids produce

saturated fatty acids via *tesA*/*B* and *yciF* (**Fig 3A**). qRT-PCR was used to quantify the

expression of involved genes in bacteria treated with different concentrations of $MgCl_2$.

The expression of 18 genes increased following treatment with 50 and/or 200 mM

$MgCl_2$ (**Fig 3B**). The expression of *tesA*, *tesB*, and *yciA*, which convert unsaturated to

saturated fatty acids, increased at 50 or/and 200 mM $MgCl_2$ (**Fig 3C**). FabA and FabF

play key roles in unsaturated fatty acid and fatty acid biosynthesis, respectively (Feng

& Cronan, 2009; Lee *et al.,* 2013). Western blot analysis showed that FabA levels

declined and FabF levels increased at higher $MgCl_2$ concentrations (**Fig 3D**). Oxazole-

2-amine and triclosan inhibit the synthesis of fatty acids and decrease bacterial survival

in the presence of balofloxacin (**Fig 3E and 3F**). These results indicate that $Mg^{2+}$

inhibits the synthesis of unsaturated fatty acids and promotes the synthesis of saturated

fatty acids.

In addition, we also quantified gene expression during fatty acid degradation to

determine whether $Mg^{2+}$ affects this process (**Fig 3G and 3H**). Interestingly, the

expression of genes involved in unsaturated fatty acid degradation decreased in an

$MgCl_2$ dose-dependent manner (**Fig 3H**). Western blot analysis confirmed that FadL

levels decreased with increasing $MgCl_2$ (**Fig 3I**). FadH degrades unsaturated fatty acids

and influences the ratio of unsaturated to saturated fatty acids. Together, these results

suggest that $MgCl_2$ inhibits fatty acid degradation.

FabR, FadR, ArcA, and cAMP/CRP are transcriptional factors involved in regulating

fatty acid biosynthesis. FabR inhibits unsaturated fatty acid biosynthesis, FadR

promotes fatty acid biosynthesis and inhibits degradation, and ArcA and cAMP/CRP

inhibit and promote fatty acid degradation, respectively (Feng and Cronan, 2010; Fujita

et al., 2007). The expression of *fabR* and *arcA* increased, while the expression of *fadR*

decreased. The expression of N646_1885, which encodes CRP, increased only in the

presence of 200 mM MgCl$_2$ (**Fig 3J**). Western blot analysis confirmed that FadR levels

declined with increasing MgCl$_2$ concentration (**Fig 3K**). Viability of Δ*fadR* and Δ*arcA*

was reduced in the presence of MgCl$_2$, balofloxacin, or both in a dose-dependent

manner (**Fig 3L**). Thus, FadR and ArcA may play roles in bacterial resistance to

balofloxacin, presumably by a mechanism dependent on the abundance of

saturated/unsaturated fatty acids.

To investigate whether Mg$^{2+}$ functions through FadR and ArcA, expression of eight

fatty acid biosynthetic genes (*accA, accC, fabD, fabH, fabR, fabG, fabV,* and *yciA*) and

five fatty acid degradation genes (*fadL, fadD, fadB, fadA,* and *fadH*) were quantified in

wildtype, Δ*fadR* or/and Δ*arcA* bacterial strains. As expected, fatty acid biosynthetic

gene expression was reduced in Δ*fadR* cells (**Fig 3M**). While higher expression of *fadD,*

*fadB,* and *fadA* (elevated *fadA* only for Δ*fadR* without 200 mM MgCl$_2$) was detected in

Δ*fadR* or Δ*acrA* strains, the expression of these genes was negatively regulated by 200

mM MgCl$_2$ (**Fig 3N and 3O**). These results indicate that Mg$^{2+}$ promotes fatty acid

biosynthesis by positively regulating *fadR*, and inhibits the degradation of fatty acids

through *fadR* and *arcA*.

**Figure 3. Mg²⁺ promotes fatty acid biosynthesis. A.** Diagram of fatty acid biosynthesis. **B.** qRT-PCR for expression of fatty acid biosynthesis genes in the absence or presence of indicated concentrations of MgCl₂. **C.** qRT-PCR for expression of genes involved in converting unsaturated fatty acids to saturated fatty acids in the absence or presence of indicated MgCl₂ levels. **D.** Western blot for the abundance of proteins responsible for converting unsaturated fatty acids to saturated fatty acids in the absence or presence of indicated concentrations of MgCl₂. **E.** Diagram of saturated and unsaturated acid biosynthesis. **F.** Synergy analysis of balofloxacin with triclosan + oxazole-2-amine for ATCC33787. The percentage of bacterial survival was quantified

in the presence of indicated MgCl₂ levels and/or balofloxacin plus triclosan and

oxazole-2-amine and used to construct the isobologram. Synergy is represented using

a color scale or an isobologram, which compares the dose needed for 50% inhibition of

synergistic agents (blue) and non-synergistic (i.e., additive) agents (red). **G.** Diagram

of fatty acid degradation. **H.** qRT-PCR for the expression of genes encoding fatty acid

degradation in the absence or presence of the indicated concentrations of MgCl₂. **I.**

Western blot for the abundance of FadL in the absence or presence of the indicated

concentrations of MgCl₂. **J.** qRT-PCR for the expression of fatty acid biosynthesis

regulatory genes in the absence or presence of the indicated concentrations of MgCl₂.

**K.** Western blot for the abundance of FadR in the absence or presence of the indicated

concentrations of MgCl₂. **L.** Synergy analysis for MgCl₂ with BLFX for ATCC33787

(WT), Δ*fadR*, and Δ*arcA*. Synergy is represented using a color scale or an isobologram,

which compares the dose needed for 50% inhibition of the synergistic agents (blue) and

non-synergistic (i.e., additive) agents (red). **M, N,** and **O** qRT-PCR for expression of

genes involved in fatty acid biosynthesis (**M**) and degradation (**N** and **O**) in

ATCC33787, Δ*fadR* or/and Δ*arcA* (**O**) in the presence or absence of 200 mM MgCl₂.

Whole cell lysates resolved by SDS-PAGE gel was stained with Coomassie brilliant

blue as loading control **(D)**, **(I) and (K)**.

**MgCl₂ affects 16-carbon and 18-carbon fatty acid metabolism**

To investigate the effect of Mg²⁺ on the biosynthesis of saturated and unsaturated fatty

acids, LC-MS was used to quantify the levels of 16-carbon and 18-carbon fatty acids,

the main precursors of lipid biosynthesis (Zhang and Rock, 2008b). The abundance of

the saturated fatty acid, palmitic acid (C16:0), was higher while the abundance of five

unsaturated fatty acids, palmitoleic acid (C16:1), linoelaidic aid (C18:2), linoleic acid

(C18:2), alpha-linoleic acid (C18:3), and stearidonic acid (C18:4) was reduced with

increasing MgCl₂ concentration (**Fig 4A**). The levels of stearic acid (C16:0) and

vaccenic acid (C18:1) remained unaffected. Total saturated fatty acid levels increased

and total unsaturated fatty acid levels decreased with increasing $Mg^{2+}$ (**Fig 4B**). These

results suggest that $Mg^{2+}$ upregulates saturated fatty acid biosynthesis but

downregulates unsaturated fatty acid biosynthesis.

Direct exposure to palmitic acid also inhibited balofloxacin-mediated killing (**Fig 2F**)

while linolenic acid promoted balofloxacin-mediated killing in a dose-dependent

manner (**Fig 4C**). Increasing exogenous palmitic acid and linolenic acid increased

intracellular palmitic acid and linolenic acid levels, respectively (**Fig 4D**). When cells

were co-treated with palmitic acid, linolenic acid, and balofloxacin, the two fatty acids

appeared to antagonize each other, as demonstrated by the Bliss model (**Fig 4E** and **4F**).

Furthermore, magnesium had a minimal effect on the antagonistic effect of palmitic

acid, linolenic acid, and balofloxacin (**Fig 4G**), suggesting that this mineral functions

through lipid metabolism. These results indicate that saturated and unsaturated fatty

acids have an antagonistic effect on balofloxacin resistance.

**Figure 4. LC-MS targeted 16-carbon and 18-carbon fatty acids and the role of**

**palmitic acids and linolenic acids in BLFX resistance. A.** Scatter plots of 16-carbon

and 18-carbon fatty acids, detected by LC-MS. The area indicates the area of the peak

of the metabolite in total ion chromatography using GC-MS. **B.** Scatter plots of total

saturated fatty acids and unsaturated fatty acids with 16-carbon and 18-carbon (**A**). **C.** Synergy analysis for BLFX with linolenic acid for ATCC 33787. Synergy is represented using a color scale or an isobologram, which compares the dose needed for 50% inhibition of synergistic agents (while) and non-synergistic (i.e., additive) agents (purple). **D.** LC-MS for the abundance of intracellular linolenic acid and palmitic acid of ATCC33787 in synergy with the indicated exogenous linolenic acid and palmitic acid. **E.** Synergy analysis of linolenic acid and palmitic acid in BLFX-mediated killing to ATCC33787. Synergy is represented using a color scale or an isobologram, which compares the dose needed for 50% inhibition of synergistic agents (blue) and non-synergistic (i.e., additive) agents (red). **F.** Bliss analysis (**E**). **G**. Percent survival of ATCC33787 in the presence of linolenic acid, palmitic acid, and BLFX with or without 50 mM MgCl$_2$.

**Mg$^{2+}$ promotes phospholipid biosynthesis**

Fatty acids are biosynthetic precursors of lysophosphatidic acid (LPA) and phosphatidic acid (PA), two key cell membrane components (Zhang and Rock, 2008a). Thus, LC-MS was used to assess the effect of Mg$^{2+}$ on membrane lipid composition (**Suppl. Fig 7A**). Higher Mg$^{2+}$ levels increased the percentage of lipids from 61 to 67%, and saturated fatty acids from 24 to 26%, but decreased the percentage of unsaturated fatty acids from 15 to 8% (**Suppl. Fig 7B**). The abundance of 11, 32, and 53 lipids was increased in 3.125, 50, and 200 mM MgCl$_2$-treated bacteria, respectively, while the abundance of 26, 52, and 107 lipids was decreased in 3.125, 50, and 200 mM MgCl$_2$-treated bacteria, respectively (**Suppl. Fig 7C**). Saturated fatty acids and lipids increased and unsaturated fatty acids decreased in an Mg$^{2+}$ dose-dependent manner (**Suppl. Fig 7D**). A total of 52 lipids were quantified, including 17 high-abundance lipids (**Suppl.**

**Fig 7E**). Phosphatidylethanolamine (PE) and phosphatidylglycerol (PG) had the first

and second highest abundance (**Suppl. Fig 7E**) and PE levels declined while PG levels

increased with increasing $Mg^{2+}$ (**Fig 5A** and **5B**). A similar effect was observed for the

unsaturated derivatives of PE and PG (**Fig 5C**). Principal component analysis (PCA)

showed that component t [1] differentiated 0 and 3.125 mM $Mg^{2+}$ from 50 and 200 mM

$Mg^{2+}$, while component t [2] separated 50 mM $Mg^{2+}$ from 0 and 200 mM $Mg^{2+}$ and

variants in 3.125 mM of $Mg^{2+}$ treatment (**Fig 5D**). S-plot analysis showed that behenic

acid, PG[16:1(9Z)/16:1(9Z)], PG[18:2 (9Z, 12Z)/16:0], PA [14:0/22:6 (4Z, 7Z, 10Z,

13Z, 16Z, 19Z)], and lysoPE(16:0/0:0) were upregulated while linoelaidic acid,

palmitoleic acid, PE-NMe(24:0/24:0), PE[16:0/14:1(9Z)], and lysoPA(i-12:0/0:0) were

downregulated (**Fig 5E**). Of these, the abundance of PG[16:1(9Z)/16:1(9Z)],

PG[18:2(9Z,12Z)/16:0], and behenic acid increased, while the levels of linoelaidic acid,

palmitoleic acid, PE[16:0/14:1(9Z)], and lysoPA decreased with increasing $Mg^{2+}$ levels

(**Fig 5F**). Thus, altered phospholipid abundance may play a role in the effect of $Mg^{2+}$

on antibiotic resistance.

PE and PG are the two end products of phospholipid metabolism (**Fig 5G**). While

expression of most genes in the phospholipid biosynthetic pathway, including *psd, pldB,*

and *glpQ*, increased in the presence of 200 mM $MgCl_2$, the expression of *pssA* and *etuB,*

*etuC*, encoding the first enzyme and the last enzymes in the pathway, respectively,

remained the same. In addition, the expression of *gpsA,* which encodes GpsA and

transforms sn-glycerol-3P to glycerone-P, increased independent of $Mg^{2+}$ concentration

(**Fig 5H**). Phosphatidylglycerol phosphate synthase (PGS), encoded by *pgsA,* and

phosphatidylserine synthase (PSS), encoded by *pssA,* are critical enzymes for the

biosynthesis of PG and PE, respectively. PGS and PSS levels were elevated and reduced,

respectively, with increasing $Mg^{2+}$ (**Fig 5I**). The effect of $Mg^{2+}$ on the activity of PSS

was confirmed using recombinant PSS (**Fig 5J**). Deletion of *plsB* and *pgpA*, the first

and last genes involved in PG biosynthesis, respectively, lowered cell viability in the

presence of balofloxacin (note: Δ*psd* could not be obtained) (**Fig 5K** and **5L**). These

results indicate that phospholipids may play a role in the effect of $Mg^{2+}$ on balofloxacin

resistance; more specifically, PE may inhibit and PG may promote resistance to

balofloxacin.

**Figure 5. Effect of Mg2+ on phospholipid biosynthesis. A.** Heatmap showing changes in differential lipid levels at the indicated concentration of MgCl₂. **B.** Abundance of ATCC33787 phosphatidylglycerol (PG) and phosphatidylethanolamine (PE) at the indicated concentrations of MgCl₂. **C.** Scatter plots of PG and PE at different saturation levels in the presence of the indicated MgCl₂ levels. **D.** PCA analysis of different concentrations of MgCl₂-induced phospholipids metabolomes. Each dot represents a technical replicate of samples in the plot. **E.** S-plot generated from OPLS-DA. Predictive component p [1] and correlation *p* (corr) [1] differentiated 0 and 3.125

mM MgCl$_2$ from 50 and 200 mM MgCl$_2$. **F.** Scatter plot of biomarkers in data (**E**). **G.**

Diagram showing glycerophospholipid metabolism. **H.** qRT-PCR of the expression of

genes encoding glycerophospholipid metabolism in the absence of or at the indicated

concentrations of MgCl$_2$. **I.** PGS and PSS levels in the absence or presence of the

indicated concentrations of MgCl$_2$. **J.** Activity of recombinant PSS in the absence or

presence of the indicated concentrations of MgCl$_2$. **K** and **L** Synergy analysis for MgCl$_2$

with BLFX for Δ*plsB* (K) and Δ*pgpA* (L). Synergy is represented using a color scale or

an isobologram, which compares the dose needed for 50% inhibition for synergistic

agents (blue) and non-synergistic (i.e., additive) agents (red).

**Mg$^{2+}$ regulates membrane polarization, permeability and fluidity to confer**

**balofloxacin resistance**

Mg$^{2+}$ remodels membrane composition, which may impair membrane potential and

polarization, critical to membrane permeability and uptake of antibiotics (Lee *et al.,*

2019; Peng *et al.,* 2015). The voltage-sensitive dye, DiBAC4(3) showed that 12.5–200

mM MgCl$_2$ promoted membrane depolarization in a dose-dependent manner (**Fig 6A**).

Meanwhile, MgCl$_2$ had a dose-dependent (**Fig 6B**) and time-dependent (**Fig 6C**) effect

on proton motive force (PMF). These results suggest that Mg$^{2+}$-dependent changes in

membrane depolarization may influence antibiotic resistance.

The efficacy of antibiotics is strongly influenced by bacterial membrane permeability

and fluidity (Saeloh *et al.,* 2018; Zhao *et al.,* 2021). Thus, fluorescence microscopy was

used to visualize these functions after incubating *V. alginolyticus* cells with FM5-95

and various concentrations of MgCl$_2$. FM5-95 staining decreased with increasing

concentrations of Mg$^{2+}$, and no staining was observed in the presence of 200 mM Mg$^{2+}$

(**Fig 6D**). SYTO9, a green fluorescent dye that binds to nucleic acid, enters and stains

bacteria cells when there is an increase in membrane permeability (Lehtinen *et al.,* 2004;

McGoverin *et al.,* 2020). Staining decreased with increasing MgCl$_2$, indicating that

bacterial membrane permeability declined in an Mg$^{2+}$ dose-dependent manner (**Fig 6E**).

These results indicate that exogenous Mg$^{2+}$ reduces bacterial membrane permeability.

Exogenous palmitic acid also shifted the fluorescence signal peaks to the left in an

MgCl$_2$-dependent manner while palmitic acid only slightly shifted the peaks (**Fig 7F**).

In contrast, exogenous linolenic acid shifted the peak to the right in a dose-dependent

manner at 50 mM MgCl$_2$ (**Fig 6G**). These results indicate that exogenous palmitic acid

and linolenic acid decrease or increase membrane permeability, respectively.

Membrane permeability was increased in Δ*fadR* cells than in WT cells and was reduced

by Mg$^{2+}$, remaining higher than the control (**Fig 6H**). However, Δ*arcA* cells had lower

membrane permeability at lower MgCl$_2$ concentrations (**Fig 6I**). Δ*pgpA* cells exhibited

higher membrane permeability than control cells at all MgCl$_2$ concentrations (**Fig 6J**).

These data are consistent with the viability of the mutants described above (**Figs 3L**

and **5L**).

Relative membrane permeability appeared to correlate with relative intracellular

balofloxacin and antibiotic efficacy. Exogenous palmitic acid and linolenic acid

reduced and promoted the uptake of balofloxacin at 159- and 18-fold, respectively (**Fig**

**6K**). Loss of *fadR* and *pgpA* increased intracellular balofloxacin (**Fig 6K**). These data

426    suggest that bacterial membrane permeability is a critical factor required for the

427    efficacy and uptake of balofloxacin, in the presence or absence of exogenous MgCl₂.

**Figure 6. Mg$^{2+}$ regulates membrane polarization, permeability, and fluidity to confer balofloxacin resistance. A** and **B**. Depolarization (**A**) and PMF (**B**) of ATCC33787 in the absence of or at indicated concentrations of MgCl$_2$. **C** and **D**. Dynamic depolarization (**D**) Membrane fluidity of ATCC33787 in the absence or presence of indicated concentrations of MgCl$_2$, as shown by fluorescence microscopy. **E**. Membrane permeability of ATCC33787 in the absence or presence of the indicated concentrations of MgCl$_2$. **F** and **G**. Membrane permeability of ATCC33787 cultured in palmitic acid (**F**) or linolenic acid (**G**) at the indicated concentrations of MgCl$_2$. **H** and **I**. Membrane permeability of Δ*fadR* (**H**) and Δ*arcA* (**I**) in the absence or presence of the indicated concentrations of MgCl$_2$. **J**. Membrane permeability of Δ*pgpA* in the absence or presence of the indicated concentrations of MgCl$_2$. **K.** Intracellular BLFX of ATCC33787 in the presence of palmitic acid, linolenic acid, PG, and PE (left panel) or Δ*fadR* and Δ*pgpA* mutants (right panel). **L.** Diagram of mechanisms of Mg$^{2+}$-mediated resistance to BLFX.

**Discussion**

This study explored the effect of Mg$^{2+}$ on the phenotypic antibiotic resistance of *V. alginolyticus*. Exogenous Mg$^{2+}$ was found to promote saturated fatty acid biosynthesis and inhibit unsaturated fatty acid biosynthesis. Mg$^{2+}$ was also shown to upregulate PGS activity and PG abundance while downregulating PSS activity and PE abundance, decreasing membrane permeability and antibiotic uptake (**Fig 6L**) and enabling phenotypic resistance. These findings are consistent with the known impact of exogenous Mg$^{2+}$ on bacterial survival in the presence of balofloxacin and other antibiotics, and the association between membrane permeability and drug efficacy. This study further showed that exogenous fatty acids influence membrane permeability and bacterial survival in the presence of balofloxacin.

Mg$^{2+}$ is the most abundant divalent cation in cells (Pohland and Schneider, 2019; Pontes

et al., 2015) playing many essential roles, including stabilizing macromolecular

complexes and membranes, binding cytoplasmic nucleic acids and nucleotides,

interacting with phospholipid head groups and cell surface molecules, and acting as an

essential cofactor in many enzymatic reactions (Groisman *et al.,* 2013). The present

study showed for the first time that exogenous Mg$^{2+}$ influences palmitic acid and

linolenic acid levels and upregulates PGS while downregulating PSS. These findings

extend our current understanding of the biological functions of Mg$^{2+}$.

Recent findings suggest a link between fatty acid biosynthesis and the antibiotic

resistance of *E. tarda* and *V. alginolyticus* (Su *et al.,* 2021). The current study found

that Mg$^{2+}$ had opposing effects on the abundance of saturated and unsaturated fatty

acids, stimulating saturated fatty acid biosynthesis at moderately high levels, while

inhibiting unsaturated fatty acids biosynthesis at ≥3 mM. Thus, the ratio of saturated

fatty acids to unsaturated fatty acids should help in predicting antibiotic resistance.

Prior studies of Gram-positive and Gram-negative bacteria (Kumariya *et al.,* 2015; Said

*et al.,* 1987) found that 10–20 mM Mg$^{2+}$ disrupts *Staphylococcus aureus* membranes

and kills stationary-phase *S. aureus* cells but does not influence the survival of

*Escherichia coli* and *Bacillus subtilis* (Xie and Yang, 2016). Low concentrations of

Mg$^{2+}$ (≤10 mM) induce PmrAB-dependent modification of lipid A in wild-type *E. coli*

(Herrera *et al.,* 2010). In two mundticin KS-resistant *Enterococcus faecium* mutants, a

putative zwitterionic amino-containing phospholipid increased significantly, while

phosphatidylglycerol and cardiolipin levels declined (Sakayori *et al.,* 2003). However,

the impact of $Mg^{2+}$ on antibiotic resistance by regulating PGS and PSS activity was not

reported. More importantly, exogenous $MgCl_2$ was shown to target these enzymes in

opposing ways to increase the PE/PG ratio, a previously unknown mechanism of $Mg^{2+}$-

mediated antibiotic resistance.

Prior studies show that in *Stenotrophomonas maltophilia,* PhoP/Q is activated by low

magnesium levels and PhoP/Q inhibition or inactivation stimulates β-lactam antibiotic

uptake (Huang *et al.,* 2021). The effects of the outer membrane permeabilizers,

polymyxin B nonapeptide and EDTA, are completely abolished by 3 mM $Mg^{2+}$ (Kwon

and Lu, 2006). In response to $Mg^{2+}$-limited growth, enteric Gram-negative bacteria

show higher lipid A acylation, which alters membrane permeability and reduces the

uptake of cationic antimicrobial peptides (Guo *et al.,* 1998). $Mg^{2+}$ reduces the high

sensitivity of rough *Salmonella typhimurium, S. minnesota*, and *Escherichia coli* 08

(which includes defects in the lipopolysaccharide carbohydrate core) mutants to several

antibiotics (Stan-Lotter *et al.,* 1979). $Mg^{2+}$ (1 mM) inhibited aminoglycoside-mediated

outer membrane permeabilization in *Pseudomonas aeruginosa* (Hancock *et al.,* 1981).

However, mechanisms underlying the impact of $Mg^{2+}$ on antibiotic resistance have

remained largely unknown. The current study revealed that $Mg^{2+}$ plays a critical role in

regulating the membrane permeability required for antibiotic resistance by modulating

PE and PG during phospholipid metabolism.

Aquaculture is an environmental gateway to the development and globalization of

antimicrobial resistance due to the excessive use of these drugs to prevent and treat

bacterial contaminants (Cabello *et al.,* 2020; Millanao *et al.,* 2018). The current study

found that $Mg^{2+}$ increased antibiotic resistance, providing a reasonable explanation of

why antibiotics are being excessively used in aquaculture and a possible solution to

$Mg^{2+}$-induced antibiotic resistance by balancing phospholipid metabolism. This finding

should promote an overall reduction in antibacterial use that will improve environment

and food safety.

In summary, $Mg^{2+}$ modulated the phenotypic resistance of *V. alginolyticus* to

balofloxacin and potentially other quinolones and other classes of antibiotics by altering

membrane permeability and thus reducing antibiotic uptake. These findings inform the

development of methods to improve the therapeutic effect of antibiotics on magnesium-

induced phenotypic resistance.

**Materials and methods**

**Bacterial strains and culture**

*V. alginolyticus* ATCC33787 and *V. parahaemolyticus* 01 are from our laboratory

collection (Jiang et al., 2022; Kou et al., 2022). They were cultured in 0.5% yeast broth

(HuanKai Microbial, Guangdong, China) (pH 7.0) with 3% NaCl at 30 °C overnight.

The cultures were diluted 1:50 (v/v) and grown in fresh 0.5% yeast broth supplemented

with desired concentrations of $MgCl_2$ at 30 °C until OD600 = 0.6. Bacteria were

harvested by centrifugation at 8,000 rpm for 3 min and resuspended in corresponding

concentrations of $MgCl_2$ to 0.6 of OD600.

**Measurement of the minimum inhibitory concentration (MIC) by microtitre-**

**dilution-method**

MIC using microtitre-dilution-method was performed as previously described (Zhang

et al., 2019). In brief, the overnight bacterial cultures in 3% NaCl were diluted at 1:100

(v/v) in fresh 0.5% yeast broth supplemented with desired concentrations of $MgCl_2$,

cultured at 30 °C and collected when the cells arrived at an OD600 of 0.6 in medium

without $MgCl_2$. Then, $1 \times 10^5$ CFU cells were dispensed into each well of a 96-well

microtiter polystyrene tray after which a series of 2-fold dilutions of antibiotic was

added. Following incubation at 30 °C for 16 h, the MIC was defined as the lowest

antibiotic concentration that inhibited visible growth. Three biological repeats were

carried out.

**Measurement of MIC by Oxford method**

Determination of MIC by Oxford cup was performed as previously described (Liu et

al., 2015). Bacterial cells cultured overnight in yeast medium were diluted at 1:100,

shaken at 30 °C and 200 rpm until the $OD_{600}$ nm reached at 0.6. Aliquot of 100 μL cells

were spread on 0.5% yeast solid medium containing 3% NaCl and 0, 0.625, 1.25, 2.5,

5, 10, 20, or 40 mM of $MgCl_2$. Oxford cups were placed on the solid medium, and

various amounts of balofloxacin (0, 0.39, 0.78, 1.56, 3.125, 6.25, and 12.5 μg) were

added. After culturing at 30 °C for 12 h, diameter of the inhibition zone was measured

and the inhibition area was calculated.

**Measurement of intracellular balofloxacin**

Measurement of intracellular balofloxacin was performed by LC-MS analysis and

plate-counting assay. For LC-MS analysis, ATCC33787 cultured in the desired

concentrations of $MgCl_2$ were collected and adjusted to OD 0.6. Aliquot of 50 mL

bacterial cells were added into a 250 mL Erlenmeyer flask and then 300 μL of 10

mg/mL balofloxacin (final concentration is 60 μg/mL). After being cultured for 6 h at

30 °C, these bacterial cells were collected and adjusted to OD 1.0 with 3% NaCl

containing corresponding concentrations of $MgCl_2$. Aliquot of 30 mL bacteria were

collected and washed three times using mobile phase (acetonitrile: double distilled

water containing 0.1mol/l formic acid = 35:65) for LC-MS detection. The bacterial cells

were added 1 mL mobile phase, crushed in ice water bath for 10 min (crush for 2S,

pause for 3S at 35% power). Following by centrifugation, supernatants were collected,

filtered, and then measured by using liquid chromatography for balofloxacin. Different

concentrations of balofloxacin were used for a standard curve. For plate-counting assay,

ATCC33787 cells cultured overnight were diluted at 1:100, shaken at 30 $^0$C and 200

rpm until the $OD_{600}$ nm reached at 0.6. The cells were diluted at 1:100 and spread on

LB agar to preparing ATCC33787 plates. Meanwhile, ATCC33787 cells were cultured

in medium with desired $MgCl_2$ and at 30 °C and 200 rpm for 6 h. These cells were

collected, washed and sonicated. The sonicated solution was added to Oxford cups with

the ATCC33787 plates. After culturing at 30 °C for 12 h, diameter of the inhibition

zone was measured. A gradient of balofloxacin was used as a standard curve for drug

quantification.

**Measurement of intracellular magnesium**

Measurements of intracellular $Mg^{2+}$ concentration were carried out as previously

described with a modification (Yang et al., 2018). In brief, *V. alginolyticus* cultured in

0%, 0.78%, 3.125%, 12.5%, 50%, and 200% $MgCl_2$ were collected, washed, and

resuspended in the same concentrations of $MgCl_2$ to an OD600 of 1.0. Aliquots of 20

mL of bacterial suspensions were centrifuged and washed by $ddH_2O$ once. The resulting

cells were weighted, which was designated wet weight. These samples were freeze-

dried overnight, which was designated dry weight. Intracellular water volume was

calculated using the following formula: $W \times (1\text{-}W/D\text{-}0.23)$ as described previously

(Unemoto et al., 1973). Two hundred microliters of concentrated nitric acid was added

to the dried cells and then heated in 75 °C for 20 min. After diluting the samples 20-

fold, they were analyzed for $Mg^{2+}$ concentration by inductively coupled plasma mass

spectrometry (ICP-MS) (iCAP 6500, Thermo Fisher). The intracellular $Mg^{2+}$

concentration was calculated according to the volume of intracellular water.

**Western blotting**

Western blotting was carried out as described previously (Yao et al., 2019). Bacterial protein samples were prepared by ultrasound treatment, resolved on a 12% SDS-PAGE and transferred to nitrocellulose membranes (GE Healthcare Life Sciences). The membranes were incubated with 1:100 of the primary mouse antibodies, followed by goat anti-mouse secondary antibodies conjugated with horseradish peroxidase. Band intensities were detected by using a chemiluminescence imaging analysis system, Tanon-5200.

**Metabolomics analysis**

Metabolomics analysis was performed by GC-MS as described previously (Yang et al., 2018). Briefly, ATCC33787 were cultured in 0.5% yeast broth with desired $MgCl_2$. Equivalent numbers of cells were quenched with 60% (v/v) cold methanol (Sigma) and then centrifuged at 8,000 rpm at 4 °C for 5 min. One milliliter of cold methanol was used to extract metabolites. To do this, the samples were sonicated for 5 min at a 10-Wpower setting using the Ultrasonic Processor (JY92-IIDN, Scientz, China), followed by centrifugation at 12,000 rpm in 4 °C for 10 min. Supernatants were collected and 10 μL ribitol (0.1 mg per mL, Sigma-Aldrich, USA) was added into each sample as an internal quantitative standard. The supernatants were concentrated for metabolite derivatization and then used for GC-MS analysis. Every experiment was repeated by five biological replicates. GC-MS detection and spectral processing for GC-MS were carried out using the Agilent 7890A GC equipped with an Agilent 5975C VL MSD

detector (Agilent Technologies, USA) as described previously (Unemoto et al., 1973;

Yang et al., 2018). Statistical difference was obtained by Kruskal Wallis test and Manne

Whitney test using SPSS 13.0 and a $p$ value $<$ 0.01 was considered significant.

Hierarchical clustering was completed in the R platform (https://cran.r-project.org/)

with the function "heatmap. 2" of "gplots library". Z score analysis was used to scale

each metabolite. Multivariate statistical analysis included orthogonal partial least

square discriminant analysis (OPLS-DA) implemented with SIMCA 12.0 (Umetrics,

Umeå, Sweden). Control scaling was selected prior to fitting. All variables were mean

centered and scaled to pareto variance of each variable. Rank-sum test and a

permutation test were used to identify differentially expressed metabolites. OPLS-DA

was used to reduce the high dimension of the data set. Differential metabolites to their

respective biochemical pathways were outlined in the MetaboAnalyst 3.0

(http://www.metaboanalyst.ca/). Pathways were enriched by raw $p$ value $< 0.05$.

**qRT-PCR**

Quantitative real time polymerase chain reaction (qRT-PCR) was carried out as

described previously (Yang et al., 2020). Total RNA was extracted from *V.*

*alginolyticus* using TRIZOL regent (Invitrogen Life Technologies) according to the

protocol. RNA Electrophoresis was carried out in 1% (w/v) agarose gels to identify

quality of the extracted RNA. By using a PrimeScriptTM RT reagent Kit with gDNA

eraser (Takara, Japan), reverse transcription-PCR was carried out on 1 μg of total RNA

and primers are listed in **Table EV2.** qRT-PCR was performed in 384-well plates with

a total volume of 10 μL and the reaction mixtures were run on a LightCycler 480 system

(Roche, Germany). Data are shown as the relative mRNA expression compared to 0%

$MgCl_2$ test with the endogenous reference 16S rRNA gene.

**Antibiotic bactericidal assay**

Antibiotic bactericidal assay was performed as described previously with a

modification (Li et al., 2016; Zhao et al., 2021). The cultured bacteria of ATCC33787

and its mutant strains were transferred to fresh yeast medium at a dilution of 1:1000

and dispensed into test tubes and then the indicated concentrations of $MgCl_2$ and

balofloxacin were added. If desired, 2 mM 2-aminooxazole and 1 μg/mL triclosan were

or a metabolite complemented. These mixtures were incubated at 30 °C and 200 rpm

for 6 h. Cells were collected. To determine CFU per mL, 100-μL samples were 10-fold

serially diluted with 900 μL M9 buffer and an aliquot of 5 μL of each dilution was

spotted onto the LB agar plates and cultured at 30 °C for 8 h.

**Construction of gene-deleted mutants**

Construction of gene-deleted mutants was carried out as described previously (Kuang

et al., 2021). Primers were designed as shown in **Table EV3** using CE Design V1.03

software. To construct gene-deleted mutants, upstream and downstream 500-bp

fragments were amplified from the genome using two pairs of primers (primers P1 and

P2, primers P3 and P4), and then merged into a 1,000-bp fragment by overlap PCR

using a pair of primers (primers P1 and P4). After the fragments were digested by *Xba*I,

they were ligated into the pDS132 vector digested by the same enzymes, transformed

into MC1061 competent cells. Conversion products were coated on LB plate containing

25 μg/mL chloramphenicol and cultured at 37 °C overnight. The plasmids from colony

growing on the plate were identified by PCR using a pair of primers (primers P1 and

P4) and sequenced. The sequenced plasmids were transformed into MFD λ pir

competent cells as donor. MFD λpir and recipient bacterium ATCC 33787 were

cultured to an optical density (OD) of 1.0 and then mixed at a ratio of 4:1. After

centrifugation, the pellets were resuspended with 50 μL LB medium including DAP

(100 μg/mL), dropped onto sterilized filter paper on LB medium including DAP (100

μg/mL), and cultured for 16-18 h at 37 °C.

All bacteria rinsed from the filter paper with LB medium were smeared onto the LB

plate with chloramphenicol (25 μg/mL) and Ampicillin (100 μg/mL). After cultured for

16-18 h at 37 °C, bacteria were screened by LB plate with above two antibiotics. The

bacteria were identified by plasmid PCR using a pair of primers P1 and P4 and

sequencing and then were cultured and smeared onto the LB plates with 20% sucrose.

The clones were cultured and smeared onto the LB plates with 20% sucrose again. The

clones, which did not grow on the LB plates with chloramphenicol but grew on the LB

plates with 20% sucrose, were identified by PCR using primer P7P8 (primer P7 is set

at about 250 bp upstream of P1, and primer P8 is set at about 250 bp downstream of

P4), P4P7, and P5P6 (for amplification of the target gene).

**LC-MS analysis for lipidomics**

ATCC33787 were cultured in medium with desired concentrations of $MgCl_2$, washed by 3% NaCl with the desired concentrations of $MgCl_2$, and adjusted to OD 1.0. Aliquot of 30 mL the cultures was centrifuged each sample and bacterial cells were collected. Aliquot of 800 μL distilled water was added and treated for 5 min at 100 °C to inactivate phospholipase C. Protein concentration was determined with BCA kit. Sample solution with 3 mg protein was transferred into a 10 mL centrifuge tube, and supplemented with distilled water to 800 μL. Then all samples were processed as follows: 1 mL chloroform and 2 mL methanol (to make the volume ratio of chloroform: methanol: water = 1:2 : 0.8) were added and cholic acid was as an internal standard; These mixtures were vortexed for 2 min and then 1 mL chloroform was add and vortex for 30s; 1 mL 10% NaCl solution was add, vortexed for 30s, and placed at room temperature overnight. After the solution was layered, a 2.5 mL syringe was used to suck the lowest layer (chloroform layer) to a new EP tube. Solution of the chloroform layer was evaporated in the rotary evaporator for 2 h and then 1 mL mobile phase (50% and 50%B. A, 25 mM ammonium acetate / methanol (30 : 70); B, methanol) was added for analysis of lipids by LC-MS. Database (https://hmdb.ca) was used for lipid identification.

**Measurement of enzyme activity**

Activity of pyruvate dehydrogenase (PDH), α-ketoglutarate dehydrogenase (KGDH) and succinate dehydrogenase (SDH) was carried out as previously described (Jiang et al., 2020). Cells cultured in medium were collected and washed three times with saline.

The bacterial cells were suspended in Tris–HCl (pH 7.4) and disrupted by sonic

oscillation for 6 min (200 W total power with 35% output, 2 s pulse, 3 s pause over ice).

After centrifugation, supernatants were collected. The protein concentration of the

supernatant was determined using Bradford assay (Beyotime, P0009). Then, 200 μg

proteins were used for determination of pyruvate dehydrogenase (PDH), α-

ketoglutarate dehydrogenase (KGDH) and succinate dehydrogenase (SDH) activity.

Levels of PGS and PSS were quantified by ELISA kits according to manufacture's

instruction (Shanghai Fusheng Industrial Co., Ltd., China)

**Measurement of proton motive force**

Measurement of the transmembrane voltage delta PSI ($\delta$PSI), which is the electrical

component of the PMF, was performed as previously described (Cheng et al., 2018).

Bacteria cultured in medium with desired concentrations of $MgCl_2$ were collected at

centrifugation and labeled by DiO2(3). Approximately $1 \times 10^7$ CFU were added to a

flow cytometry tube containing 1 mL buffer with 10 μM DiOC2(3) (Sigma) and

incubated in the dark for 30 min at 30 $^0$C. Samples were assayed with BD FACSCalibur

flow cytometer with a 488 nm excitation wavelength. Gates for bacterial populations

were based on the control population by using forward versus side scatter and red versus

green emission. Size and membrane potential determined the intensity of Red (488 nm

excitation, 610 nm emission) fluorescence. The diverse ratios of red and green indicated

fluorescence intensity values of the gated populations. Computational formula of

membrane potential: $Log(10^{3/2} \times (\dfrac{\text{red fluorescence}}{\text{green fluorescence}}))$ .

**Measurement of depolarization**

The membrane potential was estimated by measuring the fluorescence of the potential-dependent probe DiBAC4(3) (Saint-Ruf et al., 2016). DiBAC4(3) is a s voltage-sensitive probe that penetrates depolarized cells, binding intracellular proteins or membranes exhibiting enhanced fluorescence and red spectral shift. ATCC33787 were cultured in medium with desired concentrations of $MgCl_2$, collected and then adjusted to OD 0.6. Aliquot of 100 μL bacterial cells were diluted to 1 mL, and 2 μL of 5 mM DiBAC4(3) was added. After incubated for 15 min at 30 ℃ without light and vibration, these samples were filtered and detected by BD FACSCalibur.

**Measurement of fluidity by fluorescence microscopy**

Measurement of membrane fluidity is performed as previously described (Wen et al., 2022). Briefly, ATCC33787 were cultured in medium with indicated concentrations of $MgCl_2$, collected and then adjusted to OD 0.6. Aliquot of 100 μL bacteria cells of each sample were diluted to 1 mL and 10 μL (10 mg/mL) FM5-95 (Thermo Fisher Scientific, USA) was added. FM5-95 is a lipophilic styryl dye that insert into the outer leaflet of bacterial membrane and become fluorescence. This dye preferentially bind to the microdomains with high membrane fluidity(Wen et al., 2022). After incubated for 20 min at 30 ℃ at vibration without light, the sample was centrifuged for 10 min at 12,000

rpm. The pellets were resuspended with 20 μL of 3% NaCI. Aliquot of 2 μL sample

was dropped on the agarose slide, and take photos under the inverted fluorescence

microscope.

**Measurement of membrane permeability**

Measurement of membrane permeability was carried out as described previously (Su et

al., 2021).   ATCC33787 were cultured in medium with desired concentrations of

$MgCl_2$, collected and then adjusted to OD 0.6. Aliquot of 100 μL bacteria cells of each

sample were diluted to 1 mL and 2 μL 10 mg/mL SYT09 was added. After incubated

for 15 min at 30 ℃ at vibration without light, the mixtures were filtered and measured

by flow cytometry (BD FACSCalibur, USA).

**Acknowledgments**

This work was sponsored by the International Cooperation and Exchange program of

National Natural Science Foundation of China (32273177) (to P.B.), Innovation Group

Project of Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai)

(311020006) (to P.B.), Natural Science Foundation of Guangdong Province

(2022A1515012079) (to L.H.), the Science and Technology Planning Project of

Guangdong Province (2023B1212060028), and Fundamental Research Funds for the

Central Universities, Sun Yat-sen University (24lgzy004).

**Author contributions**

B. Peng conceived the idea, designed the experiments, and supervise the projects. H.

Li, J. Yang, and S.F. Kuang conducted the experiments and interpreted the results.

**Declaration of interest statement**

Authors declare that they have no competing interests.

**References**

Cheng, Z.X., Yang, M.J., Peng, B., Peng, X.X., Lin, X.M., and Li, H. (2018). The
depressed central carbon and energy metabolisms is associated to the acquisition of
levofloxacin resistance in Vibrio alginolyticus. J Proteomics *181*, 83-91.
Deitchman, A.N., Singh, R.S.P., Rand, K.H., and Derendorf, H. (2018). Enhanced in
vitro activity of tigecycline in the presence of chelating agents. Int J Antimicrob Ag
*51*, 799-802.
Dutta, D., Kaushik, A., Kumar, D., and Bag, S. (2021). Foodborne Pathogenic
Vibrios: Antimicrobial Resistance. Front Microbiol *12*, 638331.
Feng, Y.J., and Cronan, J.E. (2010). Overlapping Repressor Binding Sites Result in
Additive Regulation of Escherichia coli FadH by FadR and ArcA. J Bacteriol *192*,
4289-4299.
Fujita, Y., Matsuoka, H., and Hirooka, K. (2007). Regulation of fatty acid metabolism
in bacteria. Mol Microbiol *66*, 829-839.
Garcia-Legorreta, A., Soriano-Perez, L.A., Flores-Buendia, A.M., Medina-Campos,
O.N., Noriega, L.G., Granados-Portillo, O., Nambo-Venegas, R., Tovar, A.R.,
Mendoza-Vargas, A., Barrera-Oviedo, D., et al. (2020). Effect of Dietary Magnesium
Content on Intestinal Microbiota of Rats. Nutrients *12*.
Garfinkel, L., and Garfinkel, D. (1985). Magnesium regulation of the glycolytic
pathway and the enzymes involved. Magnesium *4*, 60-72.
Grimes, D.J. (2020). The Vibrios: Scavengers, Symbionts, and Pathogens from the
Sea. Microb Ecol *80*, 501-506.
Groisman, E.A., Hollands, K., Kriner, M.A., Lee, E.J., Park, S.Y., and Pontes, M.H.
(2013). Bacterial Mg2+ Homeostasis, Transport, and Virulence. Annu Rev Genet *47*,
625-646.
Guo, L., Lim, K.B., Poduje, C.M., Daniel, M., Gunn, J.S., Hackett, M., and Miller,
S.I. (1998). Lipid A acylation and bacterial resistance against vertebrate antimicrobial
peptides. Cell *95*, 189-198.
Hancock, R.E., Raffle, V.J., and Nicas, T.I. (1981). Involvement of the outer
membrane in gentamicin and streptomycin uptake and killing in Pseudomonas
aeruginosa. Antimicrob Agents Chemother *19*, 777-785.
Herrera, C.M., Hankins, J.V., and Trent, M.S. (2010). Activation of PmrA inhibits
LpxT-dependent phosphorylation of lipid A promoting resistance to antimicrobial
peptides. Mol Microbiol *76*, 1444-1460.

Huang, H.H., Wu, B.K., Li, L.H., Lin, Y.T., and Yang, T.C. (2021). Role of the PhoPQ two-component regulatory system in the beta-lactam resistance of Stenotrophomonas maltophilia. J Antimicrob Chemoth *76*, 1480-1486.

Isaei, E., Mansouri, S., Mohammadi, F., Taheritarigh, S., and Mohammadi, Z. (2016). Novel Combinations of Synthesized ZnO NPs and Ceftazidime: Evaluation of their Activity against Standards and New Clinically Isolated Pseudomonas aeruginosa. Avicenna J Med Biotechnol *8*, 169-174.

Jiang, M., Chen, Z.G., Li, H., Zhang, T.T., Yang, M.J., Peng, X.X., and Peng, B. (2022). Succinate and inosine coordinate innate immune response to bacterial infection. Plos Pathog *18*.

Jiang, M., Kuang, S.F., Lai, S.S., Zhang, S., Yang, J., Peng, B., Peng, X.X., Chen, Z.G., and Li, H. (2020). Na+-NQR Confers Aminoglycoside Resistance via the Regulation of L-Alanine Metabolism. Mbio *11*.

Kobylka, J., Kuth, M.S., Mueller, R.T., Geertsma, E.R., and Pos, K.M. (2020). AcrB: a mean, keen, drug efflux machine. Ann Ny Acad Sci *1459*, 38-68.

Kou, T.S., Wu, J.H., Chen, X.W., Chen, Z.G., Zheng, J., and Peng, B. (2022). Exogenous glycine promotes oxidation of glutathione and restores sensitivity of bacterial pathogens to serum-induced cell death. Redox Biol *58*.

Kuang, S.F., Feng, D.Y., Chen, Z.G., Liang, Z.Z., Xiang, J.J., Li, H., Peng, X.X., and Zhang, T.T. (2021). Inactivation of Nitrite-Dependent Nitric Oxide Biosynthesis Is Responsible for Overlapped Antibiotic Resistance between Naturally and Artificially Evolved Pseudomonas aeruginosa. Msystems *6*.

Kumarage, P.M., De Silva, L., and Heo, G.J. (2022). Aquatic environments: A potential source of antimicrobial-resistant Vibrio spp. J Appl Microbiol *133*, 2267-2279.

Kumariya, R., Sood, S.K., Rajput, Y.S., Saini, N., and Garsa, A.K. (2015). Increased membrane surface positive charge and altered membrane fluidity leads to cationic antimicrobial peptide resistance in Enterococcus faecalis. Bba-Biomembranes *1848*, 1367-1375.

Kwon, D.H., and Lu, C.D. (2006). Polyamines increase antibiotic susceptibility in Pseudomonas aeruginosa. Antimicrob Agents Ch *50*, 1623-1627.

Lee, D.D., Galera-Laporta, L., Bialecka-Fornal, M., Moon, E.C., Shen, Z., Briggs, S.P., Garcia-Ojalvo, J., and Suel, G.M. (2019). Magnesium Flux Modulates Ribosomes to Increase Bacterial Survival. Cell *177*, 352-360 e313.

Lehtinen, J., Nuutila, J., and Lilius, E.M. (2004). Green fluorescent protein-propidium iodide (GFP-PI) based assay for flow cytometric measurement of bacterial viability. Cytometry A *60*, 165-172.

Li, W.X., Yao, Z.J., Sun, L.N., Hu, W.J., Cao, J.J., Lin, W.X., and Lin, X.M. (2016). Proteomics Analysis Reveals a Potential Antibiotic Cocktail Therapy Strategy for Aeromonas hydrophila Infection in Biofilm. J Proteome Res *15*, 1810-1820.

Liu, X.J., Zhu, W.C., Su, Y.B., Guo, C., Zeng, Z.H., Zhu, H., Li, H., and Peng, X.X. (2015). Characterization of ampicillin-stressed proteomics and development of a direct method for detecting drug-binding proteins in Edwardsiella tarda. J Proteomics *116*, 97-105.

Lunestad, B.T., and Goksøyr, J. (1990). Reduction in the antibacterial effect of oxy-
tetracycline in sea water by complex formation with magnesium and calcium. Dis.
aquat. Org. *9*, 67-72.
McGoverin, C., Robertson, J., Jonmohamadi, Y., Swift, S., and Vanholsbeeck, F.
(2020). Species Dependence of SYTO 9 Staining of Bacteria. Front Microbiol *11*,
545419.
Peng, B., Su, Y.B., Li, H., Han, Y., Guo, C., Tian, Y.M., and Peng, X.X. (2015).
Exogenous Alanine and/or Glucose plus Kanamycin Kills Antibiotic-Resistant
Bacteria. Cell Metab *21*, 249-261.
Pohland, A.C., and Schneider, D. (2019). Mg2+ homeostasis and transport in
cyanobacteria - at the crossroads of bacterial and chloroplast Mg2+ import. Biol
Chem *400*, 1289-1301.
Pontes, M.H., Sevostyanova, A., and Groisman, E.A. (2015). When Too Much ATP
Is Bad for Protein Synthesis. J Mol Biol *427*, 2586-2594.
Saeloh, D., Tipmanee, V., Jim, K.K., Dekker, M.P., Bitter, W., Voravuthikunchai,
S.P., Wenzel, M., and Hamoen, L.W. (2018). The novel antibiotic rhodomyrtone traps
membrane proteins in vesicles with increased fluidity. Plos Pathog *14*.
Said, A.A., Livermore, D.M., and Williams, R.J. (1987). Expression of H1 outer-
membrane protein of Pseudomonas aeruginosa in relation to sensitivity to EDTA and
polymyxin B. J Med Microbiol *24*, 267-274.
Saint-Ruf, C., Crussard, S., Franceschi, C., Orenga, S., Ouattara, J., Ramjeet, M.,
Surre, J., and Matic, I. (2016). Antibiotic Susceptibility Testing of the Gram-Negative
Bacteria Based on Flow Cytometry. Front Microbiol *7*.
Sakayori, Y., Muramatsu, M., Hanada, S., Kamagata, Y., Kawamoto, S., and Shima,
J. (2003). Characterization of Enterococcus faecium mutants resistant to mundticin
KS, a class IIa bacteriocin. Microbiol-Sgm *149*, 2901-2908.
Song, M.R., Liu, Y., Huang, X.Y., Ding, S.Y., Wang, Y., Shen, J.Z., and Zhu, K.
(2020). A broad-spectrum antibiotic adjuvant reverses multidrug-resistant Gram-
negative pathogens. Nat Microbiol *5*, 1040-+.
Stan-Lotter, H., Gupta, M., and Sanderson, K.E. (1979). The influence of cations on
the permeability of the outer membrane of Salmonella typhimurium and other gram-
negative bacteria. Can J Microbiol *25*, 475-485.
Su, Y.B., Kuang, S.F., Ye, J.Z., Tao, J.J., Li, H., Peng, X.X., and Peng, B. (2021).
Enhanced Biosynthesis of Fatty Acids Is Associated with the Acquisition of
Ciprofloxacin Resistance in Edwardsiella tarda. Msystems *6*.
Unemoto, T., Tsuruoka, T., and Hayashi, M. (1973). Role of Na + and K + in
preventing lysis of a slightly halophilic Vibrio alginolyticus. Can J Microbiol *19*, 563-
571.
Wang, T., Flint, S., and Palmer, J. (2019). Magnesium and calcium ions: roles in
bacterial cell attachment and biofilm structure maturation. Biofouling *35*, 959-974.
Wen, J., Vischer, N.O.E., de Vos, A.L., Manders, E.M.M., Setlow, P., and Brul, S.
(2022). Organization and dynamics of the SpoVAEa protein and its surrounding inner
membrane lipids, upon germination of Bacillus subtilis spores. Sci Rep *12*, 4944.

Xie, Y.T., and Yang, L.H. (2016). Calcium and Magnesium Ions Are Membrane-Active against Stationary-Phase Staphylococcus aureus with High Specificity. Sci Rep-Uk *6*.

Yang, J., Zeng, Z.H., Yang, M.J., Cheng, Z.X., Peng, X.X., and Li, H. (2018). NaCl promotes antibiotic resistance by reducing redox states in Vibrio alginolyticus. Environ Microbiol *20*, 4022-4036.

Yang, M.J., Xu, D., Yang, D.X., Li, L., Peng, X.X., Chen, Z.G., and Li, H. (2020). Malate enhances survival of zebrafish against Vibrio alginolyticus infection in the same manner as taurine. Virulence *11*, 349-364.

Yao, Z., Guo, Z., Wang, Y., Li, W., Fu, Y., Lin, Y., Lin, W., and Lin, X. (2019). Integrated Succinylome and Metabolome Profiling Reveals Crucial Role of S-Ribosylhomocysteine Lyase in Quorum Sensing and Metabolism of Aeromonas hydrophila. Mol Cell Proteomics *18*, 200-215.

Zhang, S., Wang, J., Jiang, M., Xu, D., Peng, B., Peng, X.X., and Li, H. (2019). Reduced redox-dependent mechanism and glucose-mediated reversal in gentamicin-resistant Vibrio alginolyticus. Environ Microbiol *21*, 4724-4739.

Zhang, Y.J., Boyd, S.A., Teppen, B.J., Tiedje, J.M., and Li, H. (2014). Organic acids enhance bioavailability of tetracycline in water to Escherichia coli for uptake and expression of antibiotic resistance. Water Res *65*, 98-106.

Zhang, Y.M., and Rock, C.O. (2008a). Membrane lipid homeostasis in bacteria. Nat Rev Microbiol *6*, 222-233.

Zhang, Y.M., and Rock, C.O. (2008b). Thematic review series: Glycerolipids - Acyltransferases in bacterial glycerophospholipid synthesis. J Lipid Res *49*, 1867-1874.

Zhao, X.L., Chen, Z.G., Yang, T.C., Jiang, M., Wang, J., Cheng, Z.X., Yang, M.J., Zhu, J.X., Zhang, T.T., Li, H., et al. (2021). Glutamine promotes antibiotic uptake to kill multidrug-resistant uropathogenic bacteria. Sci Transl Med *13*.

# Supplementary Materials for

**Magnesium modulates phospholipid metabolism to promote bacterial phenotypic**

**resistance to antibiotics**

Hui Li[1,2,†], Jun Yang[1,2,†], Su-fang Kuang[1], Huang-zhe Fu[1], Hui-ying Lin, Bo Peng[1,2, *]

*Corresponding author. Email: pengb26@mail.sysu.edu.cn

**This PDF file includes:**

Supplementary Text
Supplementary Tables 1 to 3
Supplementary Figures 1 to 7

**Supplementary text**

**Magnesium and balofloxacin binding only partially contributes to phenotypic resistance**

Prior studies suggest that the chelation of antibiotics by magnesium ions inhibits antibiotic uptake (Deitchman et al., 2018; Lunestad and Goksøyr, 1990). To investigate whether magnesium binds to balofloxacin, balofloxacin was pre-incubated with magnesium, and zone of inhibition (ZOI) analysis was conducted. Six different concentrations of balofloxacin were separately incubated with six different concentrations of $MgCl_2$, and then spotted on filter paper so that a defined amount of balofloxacin could be used for ZOI. While lower concentrations of $MgCl_2$, (0.78, 3.125, or 12.5 mM) did not alter the ZOI, higher concentrations, including 50 and 200 mM $MgCl_2$, decreased the ZOI (**Suppl. Fig 2A**), suggesting that even high doses of magnesium had only a partial effect on balofloxacin through direct binding. For example, at 200 mM $MgCl_2$ and 5 or 10 μg/mL balofloxacin, the balofloxacin ZOI was 53.2 and 70.3% of the ZOI at 0 mM MgCl2, suggesting that ≥50% of the antibiotics were still functional. Intracellular BLFX also decreased with increasing $MgCl_2$ (**Suppl. Fig 2B**), while exogenous $Mg^{2+}$ increased intracellular $Mg^{2+}$ levels in a dose-dependent manner. For example, exogenous 50 and 200 mM $MgCl_2$ increased intracellular $Mg^{2+}$ levels to 1.21 and 1.31 mM, respectively (**Suppl. Fig 2C**). The relationship between TolC, an efflux pump that transports quinolones from bacterial cells, and $Mg^{2+}$ was also assessed (Kobylka *et al.*, 2020; Song *et al.*, 2020). The expression of TolC/*tolC* was unaffected by $Mg^{2+}$ (**Suppl. Fig 2D**). Magnesium is critical for LPS stability. LPS

969     levels increased at 200 mM $Mg^{2+}$ (**Suppl. Fig 2E**), however, the loss of *waaF, lpxA,*

970     and *lpxC*, three key genes involved in LPS biosynthesis, did not influence balofloxacin

971     sensitivity/resistance in the presence of $Mg^{2+}$ (**Suppl. Fig 2F**). These findings suggest

972     that magnesium-induced LPS biosynthesis does not contribute directly to BLFX

973     resistance and demonstrate that $Mg^{2+}$ influx is involved in balofloxacin resistance.

990     **Supplementary Tables**

991     **Table 1 Comparison in components in LBS and ASWT**

|  | LBS | ASWT |
|---|---|---|
| Tryptone(%) | 1 | 1 |
| Yeast extract(%) | 0.5 | 0.5 |
| NaCl(mM) | 171 | 210 |
| $MgSO_4$(mM) | 0 | 35 |
| KCl(mM) | 0 | 7 |
| $CaCl_2$(mM) | 0 | 7 |

**Table 2 Primes used in the present study**

| gene | | KEGG entry | Sequences (5'-3') | gene | | KEGG entry | Sequences (5'-3') |
|------|------|------|------|------|------|------|------|
| accA | Forward | N646_1411 | AAAGGTCGTGAAACCAAAGA | fadD | Forward | N646_4025 | CATTATTTACTGCGATCCTCG |
| | Reverse | | AGCCGTGTCGATAAATGTG | | Reverse | | CCATATTGAGATATTTGTCTGGG |
| accB | Forward | N646_1967 | TCAAGCACCAATTCACTACG | fadE | Forward | N646_1395 | TATCCCAACTGATGTCAAAGG |
| | Reverse | | ACTTTCTGACCCACTTCTACG | | Reverse | | CAACCGACAGACACTCAACC |
| accC | Forward | N646_1968 | AGAAGTGACTGGTGCTGTGG | fadB | Forward | N646_2210 | ATCACTAAAACGCCCAGAAAA |
| | Reverse | | GGATAGTGTCTGCCTTTGGAC | | Reverse | | GCTGAAGCCGCCAAAGTA |
| accD | Forward | N646_1292 | CGAACTTGAGCCACAGGATA | fadA | Forward | N646_1311 | CAGATTAGCGGCCAGTTTCT |
| | Reverse | | ACGACTGAACCCATTGAACC | | Reverse | | TCAAGCGGCGGTTAATGT |
| accD | Forward | N646_4704 | AAGTTTGTCCGAGCAGTTGA | fadA | Forward | N646_2209 | ATTGCGGGCTGCGATC |
| | Reverse | | GAGACGCCACCATAAGTTTG | | Reverse | | GCACACGGCAGAGACTGC |
| fabD | Forward | N646_1131 | CTCGCCTGGTCAAGTCGT | fadH | Forward | N646_1258 | AAGTGCTGCCCTGCTCG |
| | Reverse | | TGCTGGCTTCATAAGTGCA | | Reverse | | ATGATGGCGTCGAGTTAATG |
| fabH | Forward | N646_1132 | GACCGTTCAACCATCATCCT | fabR | Forward | N646_2038 | TTTGCTACTACGCCAACTGATGCGT |
| | Reverse | | GCCATTTATCCGCATCTTG | | Reverse | | CGCTCACGCAACAACAAACGAA |
| fabH | Forward | N646_1439 | GACGCTCCAACATGGCTAG | fadR | Forward | N646_1147 | AATGCTAGGCGCTGGAA |
| | Reverse | | AGTTCGCTGGTGACAATAAGA | | Reverse | | TGAATGCGGCTTCCTG |
| fabH | Forward | N646_4350 | GGAAGACAACGGTTACTGGG | arcA | Forward | N646_1744 | TGGGTTGAATGGTTTAGTGATG |
| | Reverse | | TGCGGAATCACAACATCG | | Reverse | | TGAAGCCAGTGACGGTGAA |
| fabB | Forward | N646_1292 | TTGACAGTGCGCCCATC | crp | Forward | N646_1004 | AAGGTTGCTGCTACTTGCTGA |
| | Reverse | | TGTCTTGCTACGCCATTTAAGA | | Reverse | | GCTTCGGATTGGTAATGGTT |
| fabB | Forward | N646_3036 | CAACATTAATGCTGCGGCT | crp | Forward | N646_1885 | CAAAACAACCAGACGCAATG |
| | Reverse | | TGAGCGGCCCACATTAC | | Reverse | | CTCACGAGAACAGCCAACAAT |

| Gene | Direction | ID | Sequence | Gene | Direction | ID | Sequence |
|---|---|---|---|---|---|---|---|
| *fabF* | Forward | N646_1128 | GGTGTTGCTATCGGCTCA | *gpsA* | Forward | N646_1922 | GTGCGGGTATGTCTGATGG |
| | Reverse | | ATGTTTACAATGGTCGAAGGA | | Reverse | | ATAAAGGTTTCTGGCTGTGC |
| *fabF* | Forward | N646_3033 | AACTGCCAAAGCACTACAAGC | *plsB* | Forward | N646_2043 | GCTTAGTTCGCTTTAGTCCC |
| | Reverse | | TGATGAGCCGTATGCGATAC | | Reverse | | CACTTGGCGTTGTGGC |
| *fabG* | Forward | N646_1130 | CAACCTAACGCCAATCTACC | *plsY* | Forward | N646_2566 | CGCCAAGAATGATAGGGTC |
| | Reverse | | CCACGAGACGCAACTTCA | | Reverse | | TTATTAGGTTCGGTTTCCAGT |
| *fabG* | Forward | N646_3034 | GACCGAAGAAGAATGGGATG | *plsC* | Forward | N646_1749 | CGATTCGCTCGGTCAA |
| | Reverse | | TGCGCTGTAGTTAGTTTGACC | | Reverse | | TACTGTTTCTGCGGCTGT |
| *fabG* | Forward | N646_3826 | AGTAAAGGCGGCGTCATC | *cdsA* | Forward | N646_1423 | TTGCTGCCCTTGTGCC |
| | Reverse | | TCTGGCAGGTTCTTGGTCA | | Reverse | | GCCAAGATTTCGTCGTTTT |
| *fabG* | Forward | N646_4505 | CCCAAGGCGTTATCATCA | *pgsA* | Forward | N646_1026 | GTTTTGGCGCGTTTATT |
| | Reverse | | TTCGGGTTTCATTTGCTCT | | Reverse | | TTTCTCGCGCAATCATC |
| *fabZ* | Forward | N646_1416 | GAGAAGTACCTTCACGCGAT | *pgpA* | Forward | N646_2838 | ATGAGTTTGCGGGCTTT |
| | Reverse | | CTGTTGCCTGAGCCATTG | | Reverse | | TCCCGCCACGATATCAT |
| *fabV* | Forward | N646_4267 | AGTAATGGGCGGCGAAG | *pgpB* | Forward | N646_3562 | TTTCTTTGGCGGGTTGT |
| | Reverse | | TGGGTGCGTAATGTCAGG | | Reverse | | CTTCAGGGCGGTGCAT |
| *fabV* | Forward | N646_0267 | GGTGGTTTACTCACTGGCATC | *pssA* | Forward | N646_2008 | GATAAAACCGCCAACGA |
| | Reverse | | TCTTCACCGCCCATTACC | | Reverse | | TTACGTGCGGCGATATT |
| *tesA* | Forward | N646_4268 | AGCATTTCTGGCGATACAAC | *psd* | Forward | N646_1916 | CACTCTGCGCCAAATG |
| | Reverse | | GGAGGGAAACCTCTTAGGC | | Reverse | | ACGAGGCGGAGTGATC |
| *tesB* | Forward | N646_0021 | CTTTGGATTCGTGCTAATGG | *pldA* | Forward | N646_4183 | CTTTTTAGGTCGCCAGAAC |
| | Reverse | | ATAGAGTGGTCAATGGTGGC | | Reverse | | CATAACCCGCTGAGTATTGT |
| *yciA* | Forward | N646_1052 | TGTCTCAGCTCGACTTAGCG | *pldB* | Forward | N646_2069 | GATCTCTACCGCCAAGGT |
| | Reverse | | GGGTTTGACCCATACTTCCA | | Reverse | | AGTGCGCGATGATGAAG |
| *fadL* | Forward | N646_1315 | CTAAGTATTGGCGCAACGTACG | *glpQ* | Forward | N646_1500 | CTATCCAACCCGATTCCC |
| | Reverse | | GCCGCCGACTTGTTCTG | | Reverse | | ACCAAGGCGCTTTAATTT |

| Gene | Direction | Locus | Sequence | Gene | Direction | Locus | Sequence |
|------|-----------|-------|----------|------|-----------|-------|----------|
| *fadL* | Forward | N646_1316 | CTGCCTTCGCCGATGAT | *glpQ* | Forward | N646_3581 | AAAAGTCCAATGCCGCA |
|  | Reverse |  | GAAATTCGCGATTGGTTTG |  | Reverse |  | GTAACCGGCTTGTTGGAG |
| *fadL* | Forward | N646_3657 | TTATCTGCGGGTGCCTCT | *glpQ* | Forward | N646_3723 | CACAAATGGCGCTATCTG |
|  | Reverse |  | GTCACGCCCGTCACCAT |  | Reverse |  | CCATCGGCTTCTTCACTC |
| *fadD* | Forward | N646_1045 | TGTTCTTGAGCTGGTGGAAA | *glpQ* | Forward | N646_3724 | GACTTGTGAAGCCCGC |
|  | Reverse |  | AACACGCAATGACGAGGTAT |  | Reverse |  | AGATACTCAGGGCCAATAAAC |
| *fadD* | Forward | N646_2540 | GCATCGGTCGATTCTTCC | *etuC* | Forward | N646_3725 | CTTTTAAGCGCTGATGATGA |
|  | Reverse |  | CGCCAATCTTAACCTGTGC |  | Reverse |  | GTTAGACCGGGTCTCTCTC |
| *fadD* | Forward | N646_3023 | CTCTGATGATGCCAAACCTAC | *adhE* | Forward | N646_1240 | AGCGATGCTGCACGAA |
|  | Reverse |  | ACGAGGAGTGTAAAGTGGGTT |  | Reverse |  | CCTGTAACTAATATGGCTGAAC |

**Table 3 Primers used in the present study for construction of gene-deleted mutants**

| gene | | primer (5'-3') |
|------|------|----------------|
| *fadR* | P1 | aaaaaggatcgatcct TACTTCCCCTGAACTCAAGCAC |
| | P2 | TGTCAGCGCTTATCTACTTTTTATCCAATATTGATTTCTG |
| | P3 | ATAAAAAGTAGATAAGCGCTGACAAGTTGGATAAAC |
| | P4 | atcgcatgcggtacc GCAAAGAGATGACTTGCATAATGAGTA |
| | P5 | ATGGTCATAAAGGCGAAAAGCC |
| | P6 | TTAGCTGTCGTCTTCCGTGAAGTT |
| | P7 | AATAGTGCTTAAGTATCTGTTTGT |
| | P8 | ATGACCATAGTGTTGCAAGA |
| *arcA* | P1 | aaaaaggatcgatcct GACCTCAAGTTACTGTATGGGTTCC |
| | P2 | TAGCGGTTAAATTGGCGGTACCTAAATTTGTGACA |
| | P3 | TAGGTACCGCCATAGATTTAACCGCTAATATGCAAAGGG |
| | P4 | atcgcatgcggtacc ATTGATGGCAGGCTGTAGTTCC |
| | P5 | ATGCAAACCCCGCAGATTCT |
| | P6 | TGGCGATGATTTCTGGCGT |
| | P7 | CGACGGCAAAGCTTTCCTC |
| | P8 | AAAAGTTGGTTGGAACAAAGGAA |
| *plsB* | P1 | aaaaaggatcgatcct TTAGGTGTTCCTCTGCGGCAT |
| | P2 | AAGGTAAAACGGAAAGTTCACAGCCTCAATAGGTATTC |
| | P3 | GCTGTGAACTTTTAACCGTTTTACCTTCTTCCAATAAGA |
| | P4 | atcgcatgcggtacc AGCTATTTTTAGGCTTGGCTTTTA |
| | P5 | ACAACGATAATGATGTCCCAACC |
| | P6 | GAACGACCACCCTCACTGAAGTA |
| | P7 | GCTCATCATTGGCTGCGTCT |
| | P8 | TGCTGACCATGAACCCGC |
| *waaf* | P1 | aaaaaggatcgatcct AGTTGTGGGTTCGCTGTGTTGA |
| | P2 | GAACTAACATGAGCTTTGTTTTTCCAGTATAGGAATT |
| | P3 | GAAAAACAAAGCTCATGTTAGTTCGTATTGTTTATACCTT |
| | P4 | atcgcatgcggtacc TTGGCGTAGTTCTGACACGATT |
| | P5 | AACGCCATCCAGATAGCCAAAT |
| | P6 | AGGTTTCAAGCCCGCCAAGA |
| | P7 | ATATGTGGGTACTACAGTTGCTGCA |
| | P8 | CATCTAATACCTGTTCGTCCTC |
| *lpxA* | P1 | aaaaaggatcgatcct TACGACCTGTTGCCACGTTTG |
| | P2 | ATTCGCCAATTAATTAAAACTCTCGACGAGCACAC |
| | P3 | GAGAGTTTTAATTAATTGGCGAATGAACCAATAAATAA |
| | P4 | atcgcatgcggtacc ATTTTAAAGATGCGCTTCTGACG |
| | P5 | GGCAGCGGTGGTAGAAGAGG |
| | P6 | AACGGGTGAATCGCAGACAG |
| | P7 | ATTGATGAGATGAATAAGCGCCT |
| | P8 | TAACTCGCTACCACGACTGCCT |
| *lpxC* | P1 | aaaaaggatcgatcct AGACGGTTGGTAATACAGTGAAG |

| | P2 | GCCAAGTATTTACTGCTTGCCTCTCAAAAATTTGC |
|---|---|---|
| | P3 | AGAGGCAAGCAGTAAATACTTGGCATTAGAATTAATAA |
| | P4 | atcgcatgcggtacc CTGGAGAAATCTTACAGCTTAAC |
| | P5 | ATTGAATCAGACGAGCAACG |
| | P6 | CGAATGTAGTCCACTCCCAC |
| | P7 | AGAATGCCGTTCAAGGTATTGC |
| | P8 | TTTACTTAAAGTGGTGACTGATT |
| *pgpA* | P1 | aaaaaggatcgatcct AAGCACTAGAGCGAGCTCATTAC |
| | P2 | GTTCATGGTTTAATCCTTAAACCTATACTGCTAGTTT |
| | P3 | AGGTTTAAGGATTAAACCATGAACTAAACACTGGACGT |
| | P4 | atcgcatgcggtacc ATACCCTCGTGGTAGCGGCAT |
| | P5 | GGCACAATTACCATTTACGG |
| | P6 | TAGAAACAGGGCAACACCAG |
| | P7 | CATTTTTCGAACTGGCTGATT |
| | P8 | GCAACGTGGTTATGACCAATT |

3    Small letters are homologous recombination sequences with pDS132 plasmid.
4
5

**Supplementary Figures**

**Figure 1 MIC of ATCC33787 to BLFX in the absence or presence of the indicated concentrations of KCl or CaCl$_2$ in M9 media.**

**Figure 2. Mg²⁺ decreases balofloxacin uptake.**

**A.** Oxford cup test for effect of different concentrations of $Mg^{2+}$ on antibacterial action of different dose of BLFX to ATCC33787. To do this, 0 mM, 0.78 mM, 3.125 mM, 12.5 mM, 50 mM, and 200 mM $MgCl_2$ were individually mixed with 0 μg/mL, 12.5 μg/mL, 25 μg/mL, 50 μg/mL, 100 μg/mL or 200 μg/mL BLFX for 5 h. Then, 50 μL were added into Oxford cup for antibacterial efficiency, which contained 0 μg, 0.625 μg, 1.25 μg, 2.5 μg, 5 μg, or 10 μg BLFX, respectively.

**B.** Intracellular BLFX of ATCC 33787 in ASWT with the indicated concentrations of $MgCl_2$ and 60 μg/mL BLFX.

**C.** Intracellular $Mg^{2+}$ of ATCC 33787 in ASWT with the indicated concentrations of $MgCl_2$.

**D.** Western blot for abundance of TolC in the presence of $MgCl_2$. Whole cell lysates resolved by SDS-PAGE gel was stained with Coomassie brilliant blue as loading control.

60     **E.** LPS quantification at indicated concentrations of MgCl$_2$.

61     **F.** MIC of ATCC 33787 and its mutants Δ*waaF*、Δ*lpxA*、Δ*lpxC* in ASWT with the indicated

62     concentrations of MgCl$_2$, which is measured by microtitre-dilution-method.

**Figure 3 Metabolic profiles of *V. alginolyticus* in different concentrations of MgCl₂.**

**A.** Reproducibility

**B**. Percentage of metabolites in every category

**C.** Heatmap of metabolites

**Figure 4 Heatmap and Z score plots of differential metabolites.**

**A**. Heatmap of differential metabolites

**B-F**. Z score plots of differential metabolites.

**Figure 5 Pathway enrichment of differential metabolites.**

**A**. Pathway enrichment of differential metabolites

**B**. Differential metabolites in enriched pathways

88
89
90

Figure 6 Scatter plots of differential metabolites identified by S-plot.

91
92
93

**Figure 7 Lipidomes in the different concentrations of MgCl$_2$.**

A. Area of fatty acids in the presence of indicated concentrations of MgCl$_2$.

B. Percentage of lipids, saturated fatty acid and unsaturated fatty acid in the presence of indicated concentration of MgCl$_2$.

C. Volcano plots of lipidomics of indicated concentration of MgCl$_2$ as compared to non-treated control.

D. Relative abundance of saturated fatty acids, unsaturated fatty acids and lipids in the presence of indicated concentrations of MgCl$_2$.

E. Relative percentage of indicated lipids.