Aromatase in adipose tissue exerts an osteoprotective function in male mice via

phosphate regulation

Aoi Ikedo[1], Michiko Yamashita[2], Maiko Hoshino[3], Yosuke Okuno[4], Minori Uga[5],

Kazuya Tanifuji[5], Hiroko Segawa[5], Seiji Fukumoto[6], Yuuki Imai[1, 7*]

[1]Division of Integrative Pathophysiology, Proteo-Science Center, Ehime University

[2]Department of Hepato-Biliary-Pancreatic Surgery and Breast Surgery, Ehime

University Graduate School of Medicine

[3]Graduate School of Agriculture and Life Sciences, The University of Tokyo

[4]Department of Metabolic Medicine, Osaka University Graduate School of Medicine

[5]Department of Applied Nutrition, Tokushima University Graduate School of

Biomedical Sciences

[6]Department of Molecular Endocrinology, Fujii Memorial Institute of Medical Sciences,

Institute of Advanced Medical Sciences, Tokushima University

[7]Department of Pathophysiology, Ehime University Graduate School of Medicine.

*Yuuki Imai

Email: y-imai@m.ehime-u.ac.jp

Author Contributions: A.I., M.Y. and M.H. performed research, analyzed data, and

wrote the paper. M.Y., M.H., Y.O., K.T., and M.U. performed research and analyzed

data. H.S. designed research. S.F. designed research and wrote the paper. Y.I. conceived

the study, designed research, and wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Keywords: Estrogen; aromatase; adipose tissue; osteoid; phosphate metabolism

Funding: This work was supported in part by Medical Research Grant, Takeda Science
Foundation to A.I. and JSPS KAKENHI (JP23689066, JP15H04961, JP19H03786,
JP22H03203, to Y.I.; JP19K18502, JP21K16688, to A.I.; JP22K09358 to Y.I., A.I.)

Abstract

Aromatase contributes to maintenance of bone mass because male patients with loss-of-function mutations of *CYP19A1* exhibit bone loss. Treatment with aromatase inhibitor also causes bone loss in men and post-menopausal women, suggesting that part of the anabolic effect of testosterone in men is dependent on estradiol (E2) biosynthesized by aromatase in non-gonadal tissues. It remains unclear how locally biosynthesized E2 contributes to maintenance of bone mass. We examined the function of aromatase in local tissues rather than gonads using cell-type specific aromatase knockout (KO) mice. Because osteoblast-specific aromatase KO mice exhibited no bone phenotype, we focused on adipose tissue, known as a reservoir of steroid hormones and analyzed the bone phenotypes of adipose tissue-specific aromatase KO mice ($Aro^{\Delta aP2}$). Sixteen-week-old male $Aro^{\Delta aP2}$ mice exhibited significantly lower bone mineral density in tibia and femur, especially in trabecular bone, than controls. Bone histomorphometry showed that $Aro^{\Delta aP2}$ mice exhibited an insufficient calcification bone phenotype with increased osteoid volume and width, and decreased osteoclast area and numbers. Moreover, serum phosphate, renal phosphate reabsorption and FGF23 were significantly lower in $Aro^{\Delta aP2}$, suggesting that the insufficient calcification phenotype in $Aro^{\Delta aP2}$ was not caused by excessive FGF23 activities. Finally, we analyzed NaPi2a and NaPi2c, phosphate transporters localized in the kidney, and found that protein levels in renal brush border membrane vesicles were lower in $Aro^{\Delta aP2}$. These results indicate that estrogens locally biosynthesized by aromatase in adipocytes can play a significant role in bone mass maintenance via regulation of phosphate reabsorption in the kidney by NaPi2.

59 **Impact Statement**

60 Male adipocyte-specific aromatase KO mice ($Aro^{\Delta aP2}$) exhibited bone loss and increased

61 osteoid due to decreased serum phosphate and urinary phosphate reabsorption as a

62 result of reduced NaPi2 proteins expression in the kidney.

Introduction

Postmenopausal osteoporosis is a major medical problem that reduces the quality of life in elderly women (Rosen, 2005). Estrogen deficiencies can induce postmenopausal osteoporosis due to the lack of various physiological functions of estrogen/estrogen receptor signaling, such as maintenance of osteoclastic life span (Martin-Millan et al., 2010; Nakamura et al., 2007), maintenance of osteoblast/osteocyte functions by Wnt signaling (Almeida et al., 2013; Doolittle et al., 2022; Kondoh et al., 2014), inhibition of inflammatory cytokine production (Pacifici, 2008), and negative feedback of FSH secretion (Sun et al., 2006). Even in older men, low serum levels of estradiol (E2) (Mellström et al., 2008) as well as testosterone (Shahinian, Kuo, Freeman, & Goodwin, 2005) can lead to significant risks of fractures. Estrogens are synthesized from cholesterols by the activity of converting enzymes, including aromatase (*Cyp19a1*), mainly in the gonads. The *Cyp19a1* gene is expressed in various tissues with tissue-specific alternative splicing (Honda, Harada, & Takagi, 1994; Mahendroo, Mendelson, & Simpson, 1993; Means, Kilgore, Mahendroo, Mendelson, & Simpson, 1991; Simpson et al., 1993). Patients harboring a loss-of-function mutation in the *CYP19A1* gene show severe osteoporosis and greater stature due to delayed or unsuccessful growth plate closure (Morishima, Grumbach, Simpson, Fisher, & Qin, 1995), both of which are sensitive to estrogen replacement therapy (Bilezikian, Morishima, Bell, & Grumbach, 1998; Carani et al., 1997; Rochira & Carani, 2009). Also, conventional aromatase knockout mice (ArKO) exhibited bone loss caused by decreased bone formation (Oz et al., 2001) or increased bone resorption (Miyaura et al., 2001). These reports indicate that the functions of aromatase are essential for the preservation of bone health. However, it is still uncertain whether a deficiency of

aromatase alters bone metabolism directly or indirectly via systemic influences because

both patients and ArKO mice show systemic endocrine disturbances.

Aromatase inhibitors (AIs) are applied clinically to treat certain populations of breast

cancers in postmenopausal women (Goss et al., 2003). Although the serum levels of

estrogen should be remarkably decreased in postmenopausal women, AIs still have

adverse effects by increasing the risk of osteoporosis, called cancer treatment-induced

bone loss (Becker et al., 2012). Furthermore, Finkelstein et al. reported that the

administration of AIs to healthy men decreased bone mass in a serum testosterone

concentration-independent manner (Finkelstein et al., 2016). These results suggest that

part of the anabolic effect of testosterone (T) in men is dependent on E2 biosynthesis by

aromatase. This evidence may indicate that the activities of aromatase in peripheral

tissues and organs as well as the gonads are significant. Moreover, local synthesis of

estrogens by aromatase may exert a variety of biological functions. However, it remains

unclear how estrogens, which is biosynthesized locally by the action of aromatase in

non-gonadal tissues, contributes to the maintenance of bone mass.

In this study, we generated functional *Cyp19a1* floxed mice and osteoblast- or

adipocyte-specific aromatase KO mice. Male adipocyte-specific aromatase KO mice

($Aro^{\Delta aP2}$) exhibited bone loss due to decreased serum phosphate and urinary phosphate

reabsorption as a result of reduced NaPi2 expression in kidney. These results indicate

that locally synthesized estrogens in adipocytes can play a significant role in bone mass

maintenance. It also suggests the possible existence of a new phosphate regulatory

mechanism through aromatase in adipose tissue.

Results

Generation of *Cyp19a1* floxed mice.

To assess the tissue-specific function of aromatase, which is coded by the *Cyp19a1* gene,

we generated *Cyp19a1* floxed mice. Figure S1 A shows the strategy used to generate the

mutant *Cyp19a1* allele. Two loxPs were designed to be inserted at sites flanking exon 8

and exon 9 of the *Cyp19a1* allele as those exons correspond to the catalytic domain of

aromatase (Ghosh, Griswold, Erman, & Pangborn, 2009). Successful insertion of NeoR

cassette and loxPs was confirmed by Southern blotting using two different probes

(Figure S1 B). Cre-mediated recombination excised exon 8, exon 9 and the NeoR gene

from the mutant allele, resulting in the null allele. As expected, the *Cyp19a1* transcript

was not detected in systemic *Cyp19a1* knockout ($Aro^{-/-}$) mice by either qualitative or

quantitative RT-PCR (Figure S1 C).

Phenotypes of systemic and osteoblast-specific *Cyp19a1* knockout mice.

To further confirm the abnormal phenotypes of the systemically expressed Cre-mediated

deletion in *Cyp19a1* floxed mice, we determined whether $Aro^{-/-}$ mice showed the same

phenotypic abnormalities reported in conventional *Cyp19a1* knockout mice (Fisher,

Graves, Parlow, & Simpson, 1998; Jones et al., 2000). As expected, female $Aro^{-/-}$ mice

exhibited smaller atrophic uteri and male $Aro^{-/-}$ mice showed significantly larger seminal

vesicles (Figure S1 D, E). Also, $Aro^{-/-}$ mice showed a significantly increased volume of

fat tissue, regardless of gender (Figure S1 E). These results suggested that aromatase

was functionally depleted in *Cyp19a1* floxed mice when excised by Cre-mediated

recombination of the *Cyp19a1* gene locus, as these phenotypes were similar to those in

previous reports.

Next, we tried to reveal the function of aromatase in osteoblast lineage cells, because

some previous studies reported that aromatase is expressed in osteoblasts (Sasano et al., 1997; Shozu & Simpson, 1998; Yanase et al., 2003). We generated three lines of osteoblast/osteocyte-specific *Cyp19a1* knockout mice by crossing *Cyp19a1* floxed (*Aro$^{flox/flox}$*) mice with *2.3kb-Col1a1-Cre* (*Aro$^{\Delta Col1a1}$*), *OCN-Cre* (*Aro$^{\Delta OCN}$*), or *Dmp1-Cre* (*Aro$^{\Delta Dmp1}$*) mice. Bone mineral density (BMD) in *Aro$^{\Delta Col1a1}$* female mice and *Aro$^{\Delta OCN}$* male mice was comparable with controls (Figure S2 A, B). In contrast, BMD of *Aro$^{\Delta Dmp1}$* mice was significantly greater in female but not male mice (Figure S2 C). However, we did not analyze this phenotype in detail because our goal was to determine why aromatase deficiency causes bone loss. Taken together, these results suggest that local aromatase expression in osteoblast/osteocyte lineage cells either does not contribute to bone mass maintenance or has a negative effect on bone mass.

Deletion of *Cyp19a1* from adipose tissue resulted in decreased bone mass in male mice.

Adipose tissue is known as the major site of estrogen biosynthesis or restoration in postmenopausal women and old men. Previous studies reported that aromatase expression in adipose tissue was increased by age, obesity, and menopause (Agarwal, Ashanullah, Simpson, & Bulun, 1997; Brown et al., 2017; Bulun & Simpson, 1994; Mair, Gaw, & MacLean, 2020). To investigate the effects of expressed aromatase in adipose tissue on bone, we generated adipocyte-specific *Cyp19a1* knockout mice (*Aro$^{\Delta aP2}$*) by crossing *Aro$^{flox/flox}$* mice with *aP2-Cre* mice. *Cyp19a1* expression levels in epididymal white adipose tissue (eWAT) were significantly lower in *Aro$^{\Delta aP2}$* compared to controls (Figure 1 A). Although there were no differences in body weight (Figure 1 B), tibial and femoral BMD of *Aro$^{\Delta aP2}$* male mice at the age of 16-20 weeks were

significantly lower compared to control but not female mice (Figure 1 C and Figure S2

D). Especially, proximal site BMD in the tibia was significantly less in $Aro^{\Delta aP2}$ mice

compared to controls. Also, $Aro^{\Delta aP2}$ mice had low gastrocnemius muscle weight, and

high eWAT weight compared to controls (Figure 1 D, E). These phenotypes of low bone

and high fat mass were also observed in $Aro^{\Delta aP2}$ mice at the age of 24 weeks (Figure S3

A-E). It has been reported that some Cre mouse lines develop skeletal dysplasia or

abnormal regeneration in tissues (Rodda & McMahon, 2006; Sakai et al., 2022).

Therefore, we tested whether these phenotypes of $Aro^{\Delta aP2}$ mice were due to the

influence of the Cre mouse line. The results showed that *aP2-Cre* mice were

comparable in tibial and femoral BMD, fat mass, and muscle mass to littermate WT

control mice (Figure S4 A-D). We measured adipocyte size in eWAT and performed

intraperitoneal glucose tolerance tests (IPGTT) because $Aro^{\Delta aP2}$ mice were obese.

Adipocyte size was no different between groups (Figure 1 F). Fasting blood glucose was

significantly higher overall in $Aro^{\Delta aP2}$ compared to controls, but there was no difference

at any specific time point between groups (Figure 1 G). Since $Aro^{\Delta aP2}$ mice have

suppressed E2 biosynthesis from T by aromatase in adipose tissue, serum T and E2

levels were measured by LC-MS/MS. There were no differences in serum E2 and T

levels between groups (Figure 1 H), suggesting that E2 biosynthesized by aromatase in

adipocytes works only locally, not systemically.

As E2 biosynthesis in adipocytes is reduced due to aromatase ablation in $Aro^{\Delta aP2}$, it is

presumed that E2/estrogen receptor (ER) signaling is also reduced. In fact, we examined

steroid hormone receptor expression in eWAT and found that *Esr1* expression was

significantly less in $Aro^{\Delta aP2}$ compared to control mice (Figure S5 A). Therefore, to

investigate whether the $Aro^{\Delta aP2}$ phenotype is due to local ERα action in adipocytes, we

generated adipose tissue-specific *Esr1* knockout mice (*ERα^{ΔaP2}*) by crossing *ERα^{flox/flox}*

mice with *aP2-Cre* mice. *Esr1* expression levels in eWAT were significantly lower in

ERα^{ΔaP2} compared to control mice (Figure S5 B). The results showed no group

differences in body weight, bone or muscle mass (Figure S5 C, D, F). In contrast,

subcutaneous WAT (sWAT) and eWAT weight were significantly greater in *ERα^{ΔaP2}*

compared to control mice (Figure S5 E). This phenotype was similar to that of *Aro^{ΔaP2}*

mice. These results suggest that ERα in adipocytes is involved in lipid metabolism but

not bone metabolism. Therefore, the bone phenotype in *Aro^{ΔaP2}* was not an

ERα-mediated effect of adipocytes.

Aro^{ΔaP2} mice had an abnormal phosphate metabolism.

We analyzed bone structure and bone marrow fat volume of the proximal tibia of

Aro^{ΔaP2} and *Aro^{flox/flox}* mice using μCT. Trabecular bone parameters such as trabecular

bone volume (BV/TV) and trabecular thickness (Tb.Th) were significantly lower in

Aro^{ΔaP2} compared to control mice (Figure 2 A). However, cortical bone parameters, and

regulated bone marrow tissue (rBMAT) volume were no different between groups

(Figure 2 B, C). Further, serum P1NP, a bone formation marker, and serum CTX, a bone

resorption marker, were no different between groups (Figure 2 D).

To investigate the dynamics of osteoclasts and osteoblasts, we performed bone

histomorphometry in the proximal tibia of *Aro^{ΔaP2}* and *Aro^{flox/flox}* mice. Von Kossa

staining revealed significantly greater osteoid volume (OV/BV) and osteoid thickness

(O.Th) in *Aro^{ΔaP2}* compared to control mice (Figure 3 A, B). In addition, TRAP staining

revealed significantly lower osteoclast surface/bone surface (Oc.S/BS) and osteoclast

number/bone perimeter (N.Oc/B.Pm) in *Aro^{ΔaP2}* compared to control mice (Figure 3 G,

H). However, bone formation rate and osteoblast number were no different between

groups (Figure 3 C-F). These results suggest that $Aro^{\varDelta aP2}$ mice exhibit an abnormal bone

calcification phenotype.

Dysregulated phosphate metabolism is an important cause for insufficient calcification.

To assess this, we analyzed serum and urine parameters related to phosphate

metabolism of $Aro^{\varDelta aP2}$ and $Aro^{flox/flox}$ mice. Serum inorganic phosphate (IP) was

significantly lower in $Aro^{\varDelta aP2}$ mice compared to control mice but not in urine (Figure 4

A, B). Other parameters in serum and urine such as calcium (Ca), alkaline phosphatase

(ALP), and creatinine (CRE) were no different between groups (Figure 4 A, B). Then

we calculated urinary phosphate reabsorption using serum and urine parameters.

Tmp/GFR, an indicator of phosphate reabsorption in the renal proximal tubules was

significantly lower in $Aro^{\varDelta aP2}$ compared to control mice (Figure 4 C). In addition, serum

IP and OV/BV were significantly and negatively correlated (Figure 4 D). These results

suggest that $Aro^{\varDelta aP2}$ mice have high osteoid volume and low bone mass due to

decreased serum IP caused by decreased urinary phosphate reabsorption.

To identify the responsible genes for the phenotypes due to lack of $Cyp19a1$ in adipose

tissue that regulate phosphate metabolism, we performed RNA-seq analysis using total

RNA from eWAT in $Aro^{\varDelta aP2}$ and $Aro^{flox/flox}$ mice. The PCA plot showed separation

between the $Aro^{\varDelta aP2}$ and $Aro^{flox/flox}$ (Figure S6 A). The results yielded 125 genes with a

significant increase and 100 genes with a significant decrease (Figure S6 B). We

focused on the genes that showed significant changes and performed Gene Ontology

(GO) analysis. Genes involved in inflammatory responses were increased and genes

involved in nucleotide metabolic processes were decreased in $Aro^{\varDelta aP2}$ compared to

controls (Figure S6 C, D), suggesting that the adipose tissue of $Aro^{\varDelta aP2}$ mice was in an

inflammatory state and metabolically compromised. However, there were no genes

encoding proteins that act on other organs as a humoral factor among differentially

expressed genes.

Therefore, we decided to focus our analyses on existing and known factors that can

regulate phosphate metabolism. Phosphate reabsorption occurs in the proximal tubule

cells of the kidney, mainly via Na^+-Pi cotransporter 2a (NaPi2a) and Na^+-Pi

cotransporter 2c (NaPi2c). Phosphate homeostasis is under endocrine control that is

mediated by FGF23, PTH and 1,25☐dihydroxyvitamin D ($1\alpha,25(OH)_2D_3$) (Fukumoto &

Martin, 2009; Minisola et al., 2017). Of these, FGF23, PTH, and corticosterone

suppress NaPi2 expression or function in the renal tubule (Walton & Bijvoet, 1975). To

investigate why $Aro^{\Delta aP2}$ mice have low serum phosphate levels, we analyzed several

parameters that can be influenced by proximal tubular function and NaPi2 expression

levels in the kidney. Urine beta-2-microglobulin (B2M) and serum UA, indicators for

renal tubular damage, and Na, K, and CL, indicators for Fanconi syndrome, were no

different between groups (Figure 5 A-C). In addition, serum PTH, $1\alpha,25(OH)_2D_3$,

$25(OH)D_3$, and corticosterone were not different between groups (Figure 5 D, E, G). In

contrast, serum FGF23 levels were significantly lower in $Aro^{\Delta aP2}$ compared to control

mice (Figure 5 F). Initially we hypothesized that high FGF23, as seen in patients with

FGF23-related hypophosphatemic rickets/osteomalacia, was the cause of low phosphate

in $Aro^{\Delta aP2}$ mice. However, serum FGF23 probably decreased in response to lower serum

phosphate concentrations, suggesting that the insufficient calcification phenotype in

$Aro^{\Delta aP2}$ mice was not due to increased FGF23. Next, we analyzed NaPi2 expression in

the kidney. Transcript levels of *Npt2a* and *Npt2c* in the kidney were no different

between groups (Figure 5 H). As NaPi2 activity can be regulated by endocytosis and

membrane recycling of NaPi2 protein in addition to transcriptional and translational

regulations, protein levels of NaPi2a and NaPi2c in brush border membrane vesicles

(BBMVs) were confirmed by western blotting. Protein levels of NaPi2a and NaPi2c in

BBMVs were significantly decreased in $Aro^{\Delta aP2}$ compared to control mice (Figure 5 I).

In $Aro^{\Delta aP2}$ mice, phosphate-regulating hormones such as PTH, $1\alpha,25(OH)_2D_3$, and

corticosterone, previously reported to regulate NaPi2 in the kidney, were normal and

serum FGF23 was lower. Taken together, these results indicate that renal NaPi2 is

decreased in $Aro^{\Delta aP2}$ mice, independent of known NaPi2 regulation mechanisms.

Discussion

In this study, we investigated the osteoprotective function of aromatase in the

non-gonadal tissues, bone and adipose. Although three different types of

osteoblast/osteocyte-specific aromatase KO mice exhibited no low bone mass

phenotype, $Aro^{\Delta aP2}$ male mice but not female mice exhibited abnormal bone

calcification with low serum phosphate levels due to lower urinary phosphate

reabsorption caused by decreased NaPi2a and NaPi2c in the kidney (Figure 6).

Previous studies have demonstrated that obesity is associated with lower fracture risk.

In a reciprocal way, low body mass index (BMI) is associated with increased fracture

risk, mainly in the hip, but also in the spine, forearm and other sites, in post-menopausal

women (Mpalaris, Anagnostis, Goulis, & Iakovou, 2015). In addition, it has been

reported that increasing BMI was associated with a progressive increase in hip BMD

and structural measures in non-obese men, but no such improvements were seen in

obese men (BMI $\geq$ 30 kg/m^2) (Shen et al., 2015). These results suggest that moderate fat

mass is important to maintain bone mass and reduce fracture risk. These studies support

our data that E2 biosynthesized by aromatase in fat contributes to bone mass

maintenance. $Aro^{\varDelta aP2}$ mice exhibited not only bone loss but also low muscle mass, high

fat mass, and high fasting blood glucose, partially similar to symptoms of late-onset

hypogonadism (LOH), an age-related androgen deficiency (Lunenfeld, Saad, & Hoesl,

2005). Furthermore, Finkelstein et al. examined changes in body composition

(Finkelstein et al., 2013) and bone mass (Finkelstein et al., 2016) in healthy men aged

20-50 years when the aromatization of T to E2 was inhibited using goserelin acetate

which suppresses endogenous gonadal steroid production, testosterone gel, and AIs such

as anastrozole. The results showed that inhibition of aromatization of T to E2 by

anastrozole increased the percentage of body fat (Finkelstein et al., 2013) and decreased

trabecular BMD as measured by quantitative computed tomography (Finkelstein et al.,

2016). These findings suggest that the effects of T are dependent on a predominantly

estrogenic effect for fat and bone in men. As the phenotype of $Aro^{\varDelta aP2}$ mice was similar

to the results of these studies, estrogen, which is synthesized by aromatase in fat, could

play an important role in maintaining body composition as well as bone in men.

The present study suggested that $Aro^{\varDelta aP2}$ mice have lower serum phosphate levels due to

lower urinary phosphate reabsorption caused by decreased NaPi2a and NaPi2c in the

kidney. Phosphate homeostasis is strictly regulated by hormones such as FGF23, PTH

and $1,25(OH)_2D_3$. Among them, FGF23 was established as the principal hormone in

regulating blood phosphate level (Shimada et al., 2004). As a mechanism, FGF receptor

type 1 (FGFR1) works as a phosphate-sensing receptor, and its downstream intracellular

signaling pathway regulates the expression of *GALNT3*. And then, the post-translational

modifications of FGF23 protein via a gene product of *GALNT3* enhance FGF23

production. As results, FGF23 can reduce blood phosphate level by inhibiting both

proximal tubular phosphate reabsorption and intestinal phosphate absorption via downregulating the activity of NaPi2a and NaPi2c in the kidney and blood $1\alpha,25(OH)_2D_3$ level (Takashi, 2024; Takashi et al., 2019; Takashi et al., 2021). However, $Aro^{\Delta aP2}$ mice has normal phosphate-regulating hormones' level and low serum FGF23. Therefore, the existence of novel control mechanisms other than those mentioned factors above was suggested.

There are some limitations in this study. First, we tried to measure E2 level in fat tissue by LC-MS/MS, but it could not be detected. Since ER signaling in eWAT was significantly lower in $Aro^{\Delta aP2}$ compared to control mice, E2 was expected to be lower in $Aro^{\Delta aP2}$. Second, we did not analyze patient data. There are no reports of low serum phosphate levels in AI-treated patients or patients with loss-of-function mutations of the *Cyp19a1* gene. However, as the phenotype in the present study was mild, it is expected that the decrease in phosphate in patients might be a mild decrease within the reference range. This requires further investigation. Third, we did not analyze $Aro^{\Delta aP2}$ female mice in detail. It is hypothesized that the intact $Aro^{\Delta aP2}$ female mice had sufficient estrogen production from the ovaries and thus had no phenotype (Figure S2 D). Therefore, it will be necessary to analyze $Aro^{\Delta aP2}$ mice in which estrogen deficiency is induced by ovariectomy or other means. Finally, in this study, we were unable to identify a novel factor that regulates NaPi2 protein levels. All known predicted regulators were examined, suggesting the existence of a novel phosphoregulatory mechanism that has not been previously identified.

In conclusion, we have shown that E2 biosynthesis by aromatase in fat plays an important role in the maintenance of bone mass in male via phosphate metabolism.

Materials and Methods

Generation of *Cyp19a1* floxed mice by gene targeting.

The PL451 vector, which contains a neo cassette flanked by two frt sites and one loxP

site, was obtained from NCI-Frederick. A cassette vector was generated by inserting the

second loxP site and the DT-A gene from the pMC1DTpA vector into PL451. Three

fragments containing the mouse *Cyp19a1* gene were amplified by PCR and inserted into

the cassette vector to form the final *Cyp19a1* knockout construct. The *Cyp19a1*

conditional knockout targeting vector was linearized by PmeI and was introduced into

M1 mouse embryonic cells (RIKEN) by electroporation and screened by genomic

Southern blotting. Probes for Southern blotting was amplified by PCR using the

following primers: Probe A, 5'-GGCCAGAGTCTGTCTGGAGT-3', and

5'-AAGCTCTGGCAGGTGTCAAT-3'; Probe B,

5'-ACCCAAAAGCCTCAACACAG-3', and 5'-GGCTTTCAGGACAAACCTTG-3'.

Chimeric mice were generated by aggregation of ES with eight cell embryos of BDF1

mice. Chimeric mice were crossed with wild-type C57BL/6J mice (CLEA Japan) and

the resulting pups with *Cyp19a1$^{flox/+}$* were genotyped with the following primers:

5'-TAGGTGGCGAGAGAAAGGAA-3', and 5'-CCCCAGAGGCTTACACACAT-3'

(WT allele; 411 bps, floxed allele; 482 bps).

Generation of systemic and osteoblast- or adipocyte- specific knockout mice.

Cyp19a1$^{-/+}$ mice were generated by crossing *Cyp19a1$^{flox/+}$* mice with *CMV-Cre* mice

(Dupé et al., 1997), which were kindly provided by Prof. Pierre Chambon. The null

allele was genotyped with the following primers: 5'-ATTCCCGCCGAATAATCTCT-3',

and 5'-ACCTTCCTTGACCAATGTCG-3' (Floxed [L2] allele; 6118 bps, KO [L-]

allele; 315 bps). *2.3kb-Col1a1-Cre* transgenic mice, which were kindly provided by Dr.

Gerard Karsenty (Dacquin, Starbuck, Schinke, & Karsenty, 2002), were crossed with

Cyp19a1$^{flox/flox}$ to generate *Aro$^{\Delta Col1a1}$* mice. *Osteocalcin-Cre* mice (Zhang et al., 2002),

which were obtained from the Jackson Laboratory, were crossed with *Cyp19a1$^{floxflox+}$* to

generate *Aro$^{\Delta OCN}$* mice. *Dmp1-Cre* transgenic mice, kindly provided by Prof. Lynda

Bonewald (Lu et al., 2007), were crossed with *Cyp19a1$^{floxflox+}$* to generate *Aro$^{\Delta Dmp1}$* mice.

aP2-Cre mice (He et al., 2003), which were obtained from the Jackson Laboratory (JAX

stock #005069), were crossed with *Cyp19a1$^{flox/flox}$* or *Esr1$^{flox/flox}$* to generate *Aro$^{\Delta aP2}$* or

ERα$^{\Delta aP2}$ mice. *Esr1$^{flox/flox}$* mice were kindly provided by Prof. Chambon (Dupont et al.,

2000). Mice were housed in a specific pathogen-free facility under climate-controlled

conditions at room temperature (22 ± 2°C) with 25% humidity and a 12-hr light/dark

cycle. They were provided with ad libitum access to food and water. Animal

experiments were approved by the Animal Experiment Committee of Ehime University

(approval no. 37A1-1/16) and were performed in accordance with the Ehime University

Guidelines for Animal Experiments.

RT-qPCR

RNA was extracted from eWAT and kidney using a RNeasy Mini Kit (70146, QIAGEN)

with QIAzol Lysis Reagent (79306, QIAGEN) or ISOGEN (319-90211, NIPPON

GENE) and synthesized into cDNA using PrimeScript RT Master Mix (RR036, Takara

Bio Inc.). RT-qPCR was performed using cDNA to evaluate gene expression with

Thermal Cycler Dice (Takara Bio, Inc.) and SYBR Premix Ex Taq II (RR820S, Takara

Bio Inc.). *Rpl13a* or *Gapdh* was used as the housekeeping gene. Primers are shown in

Table S1.

Radiological examination

Samples were fixed in 4% paraformaldehyde phosphate buffer solution (PFA) overnight at 4°C. The areal bone mineral density (aBMD) of the isolated tibiae and femora was measured by Dual-energy X-ray absorptiometry (DXA) using a bone mineral analyzer (DCS-600EX, ALOKA). Micro-computed tomography (μCT) scanning of the femora was performed according to the manufacturer's instructions using a Scanco Medical μCT35 System (SCANCO Medical) with an isotropic voxel size of 6 μm. We defined the regions of interest (ROI) as 200 slices starting 1500 μm proximal to the proximal growth plate in the tibia. These images were used for 3D reconstruction and analysis. Structural parameters (3D) included cortical structure [bone area per total area (BA/TA), BMD, and cortical thickness (Ct.Th)], and trabecular structure [bone volume per total volume (BV/TV), trabecular number (Tb.N), trabecular thickness (Tb.Th), and BMD] according to established guidelines (Bouxsein et al., 2010).

Osmium staining

Tibiae were harvested, fixed in 4% PFA overnight, and then immersed in 0.1 M EDTA (pH 7.3) for 2 weeks for decalcification. The decalcified tibiae were immersed in a 1:1 mixture of 5% potassium dichromate and 2% osmium tetroxide (23310-10, Polysciences, Inc., USA) for 48 h and stained. The stained tibiae were scanned using μCT to evaluate marrow fat volume (Scheller et al., 2014).

Histology and histomorphometry

For bone histomorphometry, the mice were double labeled with subcutaneous injections

of 15 mg/kg of calcein (Sigma) at 4 and 2 days before sacrifice. Tibiae were removed

from each mouse and fixed with 4% PFA. After μCT analysis, tibiae were embedded

with Methylmethacrylate (MMA) after dehydration and the plastic sections were cut by

a standard microtome (LEICA) into 6 μm for Von Kossa, TRAP, and Toluidine blue

staining. The region of interest (ROI) was 750×1200 mm of secondary spongiosa in the

tibia. Each section was photographed using a cellSense Ver. 1.3 (OLYMPUS) and

analyzed using OsteoMeasure V3.3.0.1 (OsteoMetrics, Inc., GA, USA).

Serum and urinary analysis

Blood samples were collected from heart when mice were sacrificed. Blood samples

were centrifuged at 3,000 g at 4°C for 10 min, and separated serum samples were frozen

and stored at -80°C until analysis. Urine samples were collected for 24 hours on the day

before mice were sacrificed and stored at -80°C until analysis. Serum estradiol (E2),

testosterone (T), $1,25(OH)_2D_3$, and $25(OH)D_3$ were analyzed by LC-MS/MS. These

analyses were outsourced to ASKA Pharma Medical Co., Ltd. Serum P1NP

(MBS2702748, MyBioSource, Inc.), CTX (AC-06F1, Immunodiagnostic Systems),

PTH (60-2305, QuidelOrtho Corporation.), FGF23 (TCY4000 (11A735), KAINOS

Laboratories, Inc.), corticosterone (K014-H1/H5, Arbor Assays Inc.), and urine

beta-2-microglobulin (ab223590, abcam) were analyzed by ELISA respectively.

Analysis of serum IP, Ca, ALP, CRE, urine IP, Ca, and CRE were outsourced to Oriental

Yeast Co., ltd.

Protein sample purification and immunoblotting

BBMVs were prepared using the Ca^{2+} precipitation method. Cortical membrane and

whole homogenate were obtained from mouse kidneys and used for immunoblotting

analyses as described previously (Hanazaki et al., 2020; Ikuta et al., 2019; Ikuta et al.,

2018; Kaneko et al., 2018). Protein samples were heated at 95°C for 5 min in sample

buffer in the presence of 2-mercaptoethanol and subjected to sodium dodecyl

sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). The separated proteins were

transferred by electrophoresis to Immobilon-P polyvinylidene difluoride membrane

(Millipore, Billerica, MA, USA) and treated with diluted antibodies. Signals were

detected using Immobilon Western (Millipore) (Hanazaki et al., 2020; Ikuta et al., 2019;

Ikuta et al., 2018; Kaneko et al., 2018). Rabbit anti-NaPi2a and NaPi2c polyclonal

antibodies were generated as described previously (Hanazaki et al., 2020; Segawa et al.,

2009; Segawa et al., 2005; Segawa et al., 2007). Mouse anti-actin monoclonal antibody

(MAB1501, Millipore, RRID: AB_2223041) was used as an internal control.

Horseradish peroxidase-conjugated anti-rabbit IgG was utilized as the secondary

antibody (115-035-003, Jackson ImmunoResearch Laboratories, Inc, West Grove, PA,

USA, RRID: AB_10015289). The diluted antibodies for immunoblotting were:

anti-NaPi2a (1:15,000), anti-NaPi2c (1:1500), and anti-actin (1:10,000).

RNA-seq analysis

Total RNA was extracted from eWAT using RNeasy Mini Kit (70146, QIAGEN) with

QIAzol Lysis Reagent (79306, QIAGEN). BRB-seq (Alpern et al., 2019) was performed

for preparing libraries with some following modifications. Oligo-dT based primer was

used for single-stranded synthesis and Second Strand Synthesis Module (NEB, #E6111)

was used for double-stranded cDNA synthesis. In-house MEDS-B Tn5 transposase

(Picelli et al., 2014; Sato et al., 2019) was used for tagmentation and amplified by 10

cycles of PCR using Phusion High-Fidelity DNA Polymerase (Thermo Scientific,

#M0530). 51bp of insert read (Read2) were obtained on Illumina NovaSeq6000.

Obtained FASTQ files were analyzed by RaNA-seq (Prieto & Barrios, 2020). Data were

registered in GEO with the accession number GSE263849. Gene Ontology Analysis

(GO) was conducted by DAVID (Huang da, Sherman, & Lempicki, 2009; Sherman et

al., 2022) for significantly ($p < 0.01$) changed genes.

Statistical analysis

The data are presented as means ± SD using unpaired t tests with Welch's correction.

The Brown-Forsythe and Welch ANOVA test was used for comparison among the four

groups, and Dunnett T3 was used for post-hoc analysis. Outliers were excluded by

Smirnov☐Grubbs test. All statistical analyses were performed using Graph Pad Prism

Version 10.0.2 (GraphPad Software Inc., La Jolla, CA) and Microsoft Excel. P values

less than 0.05 were considered significant.

Acknowledgments

We are grateful to Dr. Erina Inoue for her technical support. We thank the staff of the

Division of Analytical Bio-Medicine and the Division of Laboratory Animal Research,

the Advanced Research Support Center (ADRES), the members of the Division of

Integrative Pathophysiology, Proteo-Science Center (PROS), and Ehime University for

their technical assistance and helpful support. We also thank the Medical Research

Center for High Depth Omics, Medical Institute of Bioregulation (MIB), Kyushu

University for the RNA-sequencing.

Data Availability

RNA-seq data have been deposited in the Gene Expression Omnibus as accession

number GSE263849.

References

Agarwal, V. R., Ashanullah, C. I., Simpson, E. R., & Bulun, S. E. (1997). Alternatively spliced transcripts of the aromatase cytochrome P450 (CYP19) gene in adipose tissue of women. *J Clin Endocrinol Metab, 82*(1), 70-74. doi:10.1210/jcem.82.1.3655

Almeida, M., Iyer, S., Martin-Millan, M., Bartell, S. M., Han, L., Ambrogini, E., . . . Manolagas, S. C. (2013). Estrogen receptor-α signaling in osteoblast progenitors stimulates cortical bone accrual. *J Clin Invest, 123*(1), 394-404. doi:10.1172/jci65910

Alpern, D., Gardeux, V., Russeil, J., Mangeat, B., Meireles-Filho, A. C. A., Breysse, R., . . . Deplancke, B. (2019). BRB-seq: ultra-affordable high-throughput transcriptomics enabled by bulk RNA barcoding and sequencing. *Genome Biol, 20*(1), 71. doi:10.1186/s13059-019-1671-x

Becker, T., Lipscombe, L., Narod, S., Simmons, C., Anderson, G. M., & Rochon, P. A. (2012). Systematic review of bone health in older women treated with aromatase inhibitors for early-stage breast cancer. *J Am Geriatr Soc, 60*(9), 1761-1767. doi:10.1111/j.1532-5415.2012.04107.x

Bilezikian, J. P., Morishima, A., Bell, J., & Grumbach, M. M. (1998). Increased bone mass as a result of estrogen therapy in a man with aromatase deficiency. *N Engl J Med, 339*(9), 599-603. doi:10.1056/nejm199808273390905

Bouxsein, M. L., Boyd, S. K., Christiansen, B. A., Guldberg, R. E., Jepsen, K. J., & Müller, R. (2010). Guidelines for assessment of bone microstructure in rodents using micro-computed tomography. *J Bone Miner Res, 25*(7), 1468-1486. doi:10.1002/jbmr.141

Brown, K. A., Iyengar, N. M., Zhou, X. K., Gucalp, A., Subbaramaiah, K., Wang, H., . . .

Dannenberg, A. J. (2017). Menopause Is a Determinant of Breast Aromatase

Expression and Its Associations With BMI, Inflammation, and Systemic Markers.

J Clin Endocrinol Metab, 102(5), 1692-1701. doi:10.1210/jc.2016-3606

Bulun, S. E., & Simpson, E. R. (1994). Competitive reverse transcription-polymerase

chain reaction analysis indicates that levels of aromatase cytochrome P450

transcripts in adipose tissue of buttocks, thighs, and abdomen of women increase

with advancing age. *J Clin Endocrinol Metab, 78*(2), 428-432.

doi:10.1210/jcem.78.2.8106632

Carani, C., Qin, K., Simoni, M., Faustini-Fustini, M., Serpente, S., Boyd, J., . . .

Simpson, E. R. (1997). Effect of testosterone and estradiol in a man with

aromatase deficiency. *N Engl J Med, 337*(2), 91-95.

doi:10.1056/nejm199707103370204

Dacquin, R., Starbuck, M., Schinke, T., & Karsenty, G. (2002). Mouse

alpha1(I)-collagen promoter is the best known promoter to drive efficient Cre

recombinase expression in osteoblast. *Dev Dyn, 224*(2), 245-251.

doi:10.1002/dvdy.10100

Doolittle, M. L., Saul, D., Kaur, J., Rowsey, J. L., Eckhardt, B., Vos, S., . . . Khosla, S.

(2022). Skeletal Effects of Inducible ERalpha Deletion in Osteocytes in Adult

Mice. *J Bone Miner Res, 37*(9), 1750-1760. doi:10.1002/jbmr.4644

Dupé, V., Davenne, M., Brocard, J., Dollé, P., Mark, M., Dierich, A., . . . Rijli, F. M.

(1997). In vivo functional analysis of the Hoxa-1 3' retinoic acid response

element (3'RARE). *Development, 124*(2), 399-410. doi:10.1242/dev.124.2.399

Dupont, S., Krust, A., Gansmuller, A., Dierich, A., Chambon, P., & Mark, M. (2000).

Effect of single and compound knockouts of estrogen receptors alpha (ERalpha) and beta (ERbeta) on mouse reproductive phenotypes. *Development, 127*(19), 4277-4291.

Finkelstein, J. S., Lee, H., Burnett-Bowie, S. A., Pallais, J. C., Yu, E. W., Borges, L. F., . . . Leder, B. Z. (2013). Gonadal steroids and body composition, strength, and sexual function in men. *N Engl J Med, 369*(11), 1011-1022. doi:10.1056/NEJMoa1206168

Finkelstein, J. S., Lee, H., Leder, B. Z., Burnett-Bowie, S. A., Goldstein, D. W., Hahn, C. W., . . . Yu, E. W. (2016). Gonadal steroid-dependent effects on bone turnover and bone mineral density in men. *J Clin Invest, 126*(3), 1114-1125. doi:10.1172/JCI84137

Fisher, C. R., Graves, K. H., Parlow, A. F., & Simpson, E. R. (1998). Characterization of mice deficient in aromatase (ArKO) because of targeted disruption of the cyp19 gene. *Proc Natl Acad Sci U S A, 95*(12), 6965-6970. doi:10.1073/pnas.95.12.6965

Fukumoto, S., & Martin, T. J. (2009). Bone as an endocrine organ. *Trends Endocrinol Metab, 20*(5), 230-236. doi:10.1016/j.tem.2009.02.001

Ghosh, D., Griswold, J., Erman, M., & Pangborn, W. (2009). Structural basis for androgen specificity and oestrogen synthesis in human aromatase. *Nature, 457*(7226), 219-223. doi:10.1038/nature07614

Goss, P. E., Ingle, J. N., Martino, S., Robert, N. J., Muss, H. B., Piccart, M. J., . . . Pater, J. L. (2003). A randomized trial of letrozole in postmenopausal women after five years of tamoxifen therapy for early-stage breast cancer. *N Engl J Med, 349*(19), 1793-1802. doi:10.1056/NEJMoa032312

Hanazaki, A., Ikuta, K., Sasaki, S., Sasaki, S., Koike, M., Tanifuji, K., . . . Segawa, H.

(2020). Role of sodium-dependent Pi transporter/Npt2c on Pi homeostasis in

klotho knockout mice different properties between juvenile and adult stages.

Physiol Rep, 8(3), e14324. doi:10.14814/phy2.14324

He, W., Barak, Y., Hevener, A., Olson, P., Liao, D., Le, J., . . . Evans, R. M. (2003).

Adipose-specific peroxisome proliferator-activated receptor gamma knockout

causes insulin resistance in fat and liver but not in muscle. *Proc Natl Acad Sci U*

S A, 100(26), 15712-15717. doi:10.1073/pnas.2536828100

Honda, S., Harada, N., & Takagi, Y. (1994). Novel exon 1 of the aromatase gene

specific for aromatase transcripts in human brain. *Biochem Biophys Res*

Commun, 198(3), 1153-1160. doi:10.1006/bbrc.1994.1163

Huang da, W., Sherman, B. T., & Lempicki, R. A. (2009). Systematic and integrative

analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc,*

4(1), 44-57. doi:10.1038/nprot.2008.211

Ikuta, K., Segawa, H., Hanazaki, A., Fujii, T., Kaneko, I., Shiozaki, Y., . . . Miyamoto, K.

I. (2019). Systemic network for dietary inorganic phosphate adaptation among

three organs. *Pflugers Arch, 471*(1), 123-136. doi:10.1007/s00424-018-2242-9

Ikuta, K., Segawa, H., Sasaki, S., Hanazaki, A., Fujii, T., Kushi, A., . . . Miyamoto, K. I.

(2018). Effect of Npt2b deletion on intestinal and renal inorganic phosphate (Pi)

handling. *Clin Exp Nephrol, 22*(3), 517-528. doi:10.1007/s10157-017-1497-3

Jones, M. E., Thorburn, A. W., Britt, K. L., Hewitt, K. N., Wreford, N. G., Proietto,

J., . . . Simpson, E. R. (2000). Aromatase-deficient (ArKO) mice have a

phenotype of increased adiposity. *Proc Natl Acad Sci U S A, 97*(23),

12735-12740. doi:10.1073/pnas.97.23.12735

Kaneko, I., Segawa, H., Ikuta, K., Hanazaki, A., Fujii, T., Tatsumi, S., . . . Miyamoto, K.

I. (2018). Eldecalcitol Causes FGF23 Resistance for Pi Reabsorption and

Improves Rachitic Bone Phenotypes in the Male Hyp Mouse. *Endocrinology,*

159(7), 2741-2758. doi:10.1210/en.2018-00109

Kondoh, S., Inoue, K., Igarashi, K., Sugizaki, H., Shirode-Fukuda, Y., Inoue, E., . . .

Imai, Y. (2014). Estrogen receptor α in osteocytes regulates trabecular bone

formation in female mice. *Bone, 60*, 68-77. doi:10.1016/j.bone.2013.12.005

Lu, Y., Xie, Y., Zhang, S., Dusevich, V., Bonewald, L. F., & Feng, J. Q. (2007).

DMP1-targeted Cre expression in odontoblasts and osteocytes. *J Dent Res, 86*(4),

320-325. doi:10.1177/154405910708600404

Lunenfeld, B., Saad, F., & Hoesl, C. E. (2005). ISA, ISSAM and EAU

recommendations for the investigation, treatment and monitoring of late-onset

hypogonadism in males: scientific background and rationale. *Aging Male, 8*(2),

59-74. doi:10.1080/13685530500163416

Mahendroo, M. S., Mendelson, C. R., & Simpson, E. R. (1993). Tissue-specific and

hormonally controlled alternative promoters regulate aromatase cytochrome

P450 gene expression in human adipose tissue. *J Biol Chem, 268*(26),

19463-19470.

Mair, K. M., Gaw, R., & MacLean, M. R. (2020). Obesity, estrogens and adipose tissue

dysfunction - implications for pulmonary arterial hypertension. *Pulm Circ, 10*(3),

2045894020952019. doi:10.1177/2045894020952023

Martin-Millan, M., Almeida, M., Ambrogini, E., Han, L., Zhao, H., Weinstein, R. S., . . .

Manolagas, S. C. (2010). The estrogen receptor-alpha in osteoclasts mediates the

protective effects of estrogens on cancellous but not cortical bone. *Mol*

Endocrinol, 24(2), 323-334. doi:10.1210/me.2009-0354

Means, G. D., Kilgore, M. W., Mahendroo, M. S., Mendelson, C. R., & Simpson, E. R. (1991). Tissue-specific promoters regulate aromatase cytochrome P450 gene expression in human ovary and fetal tissues. *Mol Endocrinol, 5*(12), 2005-2013. doi:10.1210/mend-5-12-2005

Mellström, D., Vandenput, L., Mallmin, H., Holmberg, A. H., Lorentzon, M., Odén, A., . . . Ohlsson, C. (2008). Older men with low serum estradiol and high serum SHBG have an increased risk of fractures. *J Bone Miner Res, 23*(10), 1552-1560. doi:10.1359/jbmr.080518

Minisola, S., Peacock, M., Fukumoto, S., Cipriani, C., Pepe, J., Tella, S. H., & Collins, M. T. (2017). Tumour-induced osteomalacia. *Nat Rev Dis Primers, 3*, 17044. doi:10.1038/nrdp.2017.44

Miyaura, C., Toda, K., Inada, M., Ohshiba, T., Matsumoto, C., Okada, T., . . . Ito, A. (2001). Sex- and age-related response to aromatase deficiency in bone. *Biochem Biophys Res Commun, 280*(4), 1062-1068. doi:10.1006/bbrc.2001.4246

Morishima, A., Grumbach, M. M., Simpson, E. R., Fisher, C., & Qin, K. (1995). Aromatase deficiency in male and female siblings caused by a novel mutation and the physiological role of estrogens. *J Clin Endocrinol Metab, 80*(12), 3689-3698. doi:10.1210/jcem.80.12.8530621

Mpalaris, V., Anagnostis, P., Goulis, D. G., & Iakovou, I. (2015). Complex association between body weight and fracture risk in postmenopausal women. *Obes Rev, 16*(3), 225-233. doi:10.1111/obr.12244

Nakamura, T., Imai, Y., Matsumoto, T., Sato, S., Takeuchi, K., Igarashi, K., . . . Kato, S. (2007). Estrogen prevents bone loss via estrogen receptor alpha and induction of

Fas ligand in osteoclasts. *Cell, 130*(5), 811-823. doi:10.1016/j.cell.2007.07.025

Oz, O. K., Hirasawa, G., Lawson, J., Nanu, L., Constantinescu, A., Antich, P. P., . . .

Simpson, E. R. (2001). Bone phenotype of the aromatase deficient mouse. *J*

Steroid Biochem Mol Biol, 79(1-5), 49-59. doi:10.1016/s0960-0760(01)00130-3

Pacifici, R. (2008). Estrogen deficiency, T cells and bone loss. *Cell Immunol, 252*(1-2),

68-80. doi:10.1016/j.cellimm.2007.06.008

Picelli, S., Björklund, A. K., Reinius, B., Sagasser, S., Winberg, G., & Sandberg, R.

(2014). Tn5 transposase and tagmentation procedures for massively scaled

sequencing projects. *Genome Res, 24*(12), 2033-2040.

doi:10.1101/gr.177881.114

Prieto, C., & Barrios, D. (2020). RaNA-Seq: Interactive RNA-Seq analysis from

FASTQ files to functional analysis. *Bioinformatics, 36*(6), 1955-1956.

doi:10.1093/bioinformatics/btz854

Rochira, V., & Carani, C. (2009). Aromatase deficiency in men: a clinical perspective.

Nat Rev Endocrinol, 5(10), 559-568. doi:10.1038/nrendo.2009.176

Rodda, S. J., & McMahon, A. P. (2006). Distinct roles for Hedgehog and canonical Wnt

signaling in specification, differentiation and maintenance of osteoblast

progenitors. *Development, 133*(16), 3231-3244. doi:10.1242/dev.02480

Rosen, C. J. (2005). Clinical practice. Postmenopausal osteoporosis. *N Engl J Med,*

353(6), 595-603. doi:10.1056/NEJMcp043801

Sakai, H., Sawada, Y., Tokunaga, N., Tanaka, K., Nakagawa, S., Sakakibara, I., . . . Imai,

Y. (2022). Uhrf1 governs the proliferation and differentiation of muscle satellite

cells. *iScience, 25*(3), 103928. doi:10.1016/j.isci.2022.103928

Sasano, H., Uzuki, M., Sawai, T., Nagura, H., Matsunaga, G., Kashimoto, O., & Harada,

N. (1997). Aromatase in human bone tissue. *J Bone Miner Res, 12*(9),

1416-1423. doi:10.1359/jbmr.1997.12.9.1416

Sato, S., Arimura, Y., Kujirai, T., Harada, A., Maehara, K., Nogami, J., . . . Kurumizaka,

H. (2019). Biochemical analysis of nucleosome targeting by Tn5 transposase.

Open Biol, 9(8), 190116. doi:10.1098/rsob.190116

Scheller, E. L., Troiano, N., Vanhoutan, J. N., Bouxsein, M. A., Fretz, J. A., Xi, Y., . . .

Horowitz, M. C. (2014). Use of osmium tetroxide staining with

microcomputerized tomography to visualize and quantify bone marrow adipose

tissue in vivo. *Methods Enzymol, 537*, 123-139.

doi:10.1016/B978-0-12-411619-1.00007-0

Segawa, H., Onitsuka, A., Furutani, J., Kaneko, I., Aranami, F., Matsumoto, N., . . .

Miyamoto, K. (2009). Npt2a and Npt2c in mice play distinct and synergistic

roles in inorganic phosphate metabolism and skeletal development. *Am J Physiol

Renal Physiol, 297*(3), F671-678. doi:10.1152/ajprenal.00156.2009

Segawa, H., Yamanaka, S., Ito, M., Kuwahata, M., Shono, M., Yamamoto, T., &

Miyamoto, K. (2005). Internalization of renal type IIc Na-Pi cotransporter in

response to a high-phosphate diet. *Am J Physiol Renal Physiol, 288*(3),

F587-596. doi:10.1152/ajprenal.00097.2004

Segawa, H., Yamanaka, S., Onitsuka, A., Tomoe, Y., Kuwahata, M., Ito, M., . . .

Miyamoto, K. (2007). Parathyroid hormone-dependent endocytosis of renal type

IIc Na-Pi cotransporter. *Am J Physiol Renal Physiol, 292*(1), F395-403.

doi:10.1152/ajprenal.00100.2006

Shahinian, V. B., Kuo, Y. F., Freeman, J. L., & Goodwin, J. S. (2005). Risk of fracture

after androgen deprivation for prostate cancer. *N Engl J Med, 352*(2), 154-164.

doi:10.1056/NEJMoa041943

Shen, J., Nielson, C. M., Marshall, L. M., Lee, D. C., Keaveny, T. M., & Orwoll, E. S. (2015). The Association Between BMI and QCT-Derived Proximal Hip Structure and Strength in Older Men: A Cross-Sectional Study. *J Bone Miner Res, 30*(7), 1301-1308. doi:10.1002/jbmr.2450

Sherman, B. T., Hao, M., Qiu, J., Jiao, X., Baseler, M. W., Lane, H. C., . . . Chang, W. (2022). DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update). *Nucleic Acids Res, 50*(W1), W216-w221. doi:10.1093/nar/gkac194

Shimada, T., Hasegawa, H., Yamazaki, Y., Muto, T., Hino, R., Takeuchi, Y., . . . Yamashita, T. (2004). FGF-23 is a potent regulator of vitamin D metabolism and phosphate homeostasis. *J Bone Miner Res, 19*(3), 429-435. doi:10.1359/jbmr.0301264

Shozu, M., & Simpson, E. R. (1998). Aromatase expression of human osteoblast-like cells. *Mol Cell Endocrinol, 139*(1-2), 117-129. doi:10.1016/s0303-7207(98)00069-0

Simpson, E. R., Mahendroo, M. S., Means, G. D., Kilgore, M. W., Corbin, C. J., & Mendelson, C. R. (1993). Tissue-specific promoters regulate aromatase cytochrome P450 expression. *J Steroid Biochem Mol Biol, 44*(4-6), 321-330. doi:10.1016/0960-0760(93)90235-o

Sun, L., Peng, Y., Sharrow, A. C., Iqbal, J., Zhang, Z., Papachristou, D. J., . . . Zaidi, M. (2006). FSH directly regulates bone mass. *Cell, 125*(2), 247-260. doi:10.1016/j.cell.2006.01.051

Takashi, Y. (2024). Phosphate-sensing mechanisms and functions of phosphate as a first

690   messenger. *Endocr J*. doi:10.1507/endocrj.EJ24-0082

691 Takashi, Y., Kosako, H., Sawatsubashi, S., Kinoshita, Y., Ito, N., Tsoumpra, M. K., . . .

692   Fukumoto, S. (2019). Activation of unliganded FGF receptor by extracellular

693   phosphate potentiates proteolytic protection of FGF23 by its O-glycosylation.

694   *Proc Natl Acad Sci U S A, 116*(23), 11418-11427. doi:10.1073/pnas.1815166116

695 Takashi, Y., Sawatsubashi, S., Endo, I., Ohnishi, Y., Abe, M., Matsuhisa, M., . . .

696   Fukumoto, S. (2021). Skeletal FGFR1 signaling is necessary for regulation of

697   serum phosphate level by FGF23 and normal life span. *Biochem Biophys Rep,*

698   *27*, 101107. doi:10.1016/j.bbrep.2021.101107

699 Walton, R. J., & Bijvoet, O. L. (1975). Nomogram for derivation of renal threshold

700   phosphate concentration. *Lancet, 2*(7929), 309-310.

701   doi:10.1016/s0140-6736(75)92736-1

702 Yanase, T., Suzuki, S., Goto, K., Nomura, M., Okabe, T., Takayanagi, R., & Nawata, H.

703   (2003). Aromatase in bone: roles of Vitamin D3 and androgens. *The Journal of*

704   *Steroid Biochemistry and Molecular Biology, 86*(3-5), 393-397.

705   doi:10.1016/s0960-0760(03)00349-2

706 Zhang, M., Xuan, S., Bouxsein, M. L., von Stechow, D., Akeno, N., Faugere, M. C., . . .

707   Clemens, T. L. (2002). Osteoblast-specific knockout of the insulin-like growth

708   factor (IGF) receptor gene reveals an essential role of IGF signaling in bone

709   matrix mineralization. *J Biol Chem, 277*(46), 44005-44012.

710   doi:10.1074/jbc.M208265200

Figures

Figure 1. Male *Aro^{ΔaP2}* mice exhibited low bone mass, low muscle mass, high fat mass,

and high fasting blood glucose. (A) *Cyp19a1* gene expression in eWAT. (B) Body

weight. (C) Total tibial BMD (left) and BMD at each site when the bone is divided into

20 sections (right). (D) Gastrocnemius muscle weight per body weight. (E)

Subcutaneous and epidydimal white adipose tissue weight per body weight. (F)

Representative image and quantitative data of adipocyte cross sectional area (CSA) in

eWAT. Scale bar indicates 100 μm. (G) Intraperitoneal glucose tolerance test (IPGTT)

of 12 weeks old *Aro^{ΔaP2}* and *Aro^{flox/flox}* mice. (H) Serum estradiol (E2) and testosterone

(T) by LC-MS/MS. Data are presented as means ± SD. *, $p < 0.05$; **, $p < 0.01$; ***, p

< 0.001

Figure 2. $Aro^{\Delta aP2}$ mice exhibited low trabecular bone volume at the proximal tibia.

Representative image and quantitative data of (A) bone volume per tissue volume

(BV/TV), trabecular number (Tb.N), trabecular thickness (Tb.Th), and trabecular bone

mineral density (Tb.BMD) as trabecular bone parameters, (B) bone area per tissue area

(BA/TA), cortical bone mineral density (Ct.BMD), and cortical thickness (Ct.Th) as

cortical bone parameters, and (C) regulated bone marrow tissue (rBMAT) volume at the

proximal tibia. (D) Serum P1NP and CTX concentration. Data are presented as means ±

SD. *, $p < 0.05$; ***, $p < 0.001$

Figure 3. $Aro^{\Delta aP2}$ mice exhibited high osteoid volume and low osteoclast number. (A)

Representative image of Von Kossa staining in the tibia. Scale bar of low magnification

image indicates 500 µm. Scale bar of high magnification image indicates 20 µm. (B)

Osteoid volume/bone volume (OV/BV), osteoid surface/bone surface (OS/BS), osteoid

thickness (O.Th). (C) Representative image of calcein labeling in tibia. Scale bar

indicates 100 µm. (D) Mineralizing surface (MS/BS), Mineral apposition rate (MAR),

Bone formation rate (BFR/BS). (E) Representative image of Toluidine blue staining in

tibia. Scale bar indicates 50 µm. (F) Osteoblast surface (Ob.S/BS), osteoblast

number/bone perimeter (N.Ob/B.Pm), osteoblast number/osteoblast perimeter

745 (N.Ob/Ob.Pm). (G) Representative image of TRAP staining in tibia. Scale bar indicate

746 50 µm. (H) Osteoclast surface/bone surface (Oc.S/BS), osteoclast number/bone

747 perimeter (N.Oc/B.Pm), osteoclast number/osteoclast perimeter (N.Oc/Oc.Pm). Data are

748 presented as means ± SD. *, $p < 0.05$

Figure 4. *Aro$^{\Delta aP2}$* mice exhibited low serum phosphate and decreased urinary phosphate

reabsorption. (A) Serum IP, Ca, ALP and CRE. (B) Urine IP, Ca, and CRE. (C)

Tmp/GFR calculated by serum and urine parameters. (D) Correlation between serum IP

and OV/BV. Data are presented as means ± SD. **, $p < 0.01$; ****, $p < 0.0001$.

Figure 5. $Aro^{\Delta aP2}$ mice exhibited low NaPi2a and NaPi2c protein levels in the kidney.

(A) Urine B2M and (B) Serum UA as indicators for renal tubular damage. (C) Serum

Na, K, and Cl as indicators for Fanconi syndrome. (D) Serum PTH, (E) $1\alpha,25(OH)_2D_3$,

$25(OH)D_3$, (F) FGF23, and (G) corticosterone as a phosphate metabolism regulating

hormone. (H) Gene expression of *Npt2a* and *Npt2c* in the kidney. (I) Protein expression

of NaPi2a and NaPi2c in renal BBMVs. Data are presented as means $\pm$ SD. *, $p < 0.05$;

**, $p < 0.01$.

Figure 6. In $Aro^{\Delta aP2}$ mice, E2 biosynthesis by aromatase in adipose tissue was suppressed, resulting in suppression of fat-derived novel factor (s) that regulates renal NaPi2 protein levels, suppressing renal phosphate reabsorption and reducing serum phosphate levels. As a result, $Aro^{\Delta aP2}$ mice exhibited an abnormal bone calcification phenotype.

771