Independent origins of spicules reconcile the evolutionary history of sponges

Maria Eleonora Rossi[1,2], Nathan James Kenny[2,3], Mattia Giacomelli[1], Joseph N. Keating[1], Sandra

Álvarez-Carretero[1], Astrid Schuster[4], Paco Cárdenas[5,6], Sergi Taboada[2,7,8,9], Vasiliki Koutsouveli[2,10],

Philip Donoghue[1*], Ana Riesgo[2,7,9*] and Davide Pisani[1*]

1 - Bristol Palaeobiology Group, School of Earth Sciences, University of Bristol, Life Sciences Building,

Tyndall Avenue, Bristol BS8 1TH, UK

2 - Life Sciences Department, The Natural History Museum, London SW7 5BD, UK

3 - Biochemistry Department, School of Biomedical Sciences, University of Otago, Dunedin, Aotearoa

New Zealand

4 - Department of Biology, University of Southern Denmark, Campusvej 55, Odense M 5230, Denmark

5 - Pharmacognosy, Department of Medicinal Chemistry, Uppsala University, Husargatan 3, Uppsala

751 23, Sweden

6 - Museum of Evolution, Uppsala University, Norbyvägen 16, Uppsala 752 36, Sweden

7 - Department of Biodiversity and Evolutionary Biology, National Museum of Natural Sciences (CSIC),

c/José Gutiérrez Abascal 2 28006 Madrid, Spain

8- Departamento de Biodiversidad, Ecología y Evolución, Facultad de Ciencias, Universidad

Complutense de Madrid, Madrid, Spain

9 - Departamento de Ciencias de la Vida, EU-US Marine Biodiversity Group, Universidad de Alcalá,

Alcalá de Henares, Spain

10 - Division of Marine Ecology, Marine Evolutionary Ecology, GEOMAR Helmholtz Centre for Ocean

Research Kiel, Kiel, Germany

*Corresponding authors: Philip C.J. Donoghue (Phil.Donoghue@bristol.ac.uk), Ana Riesgo

(anariesgogil@mncn.csic.es), and Davide Pisani (davide.pisani@bristol.ac.uk).

Sponges (Porifera) are highly effective ecosystem engineers, playing a critical role in global
biogeochemical processes, including the nitrogen, carbon, and silica cycles. Because of that, they have
been closely linked to the evolution of Earth's environments. However, determining the evolutionary
history of sponges has posed challenges. Molecular divergence times have pointed to a Tonian or
Cryogenian origin for sponges. This contrasts with the oldest unequivocal fossil evidence for sponges,
consisting of isolated siliceous spicules from the latest Ediacaran, resulting in a 150-million-mismatch
with molecular clock estimates. This mismatch may arise because molecular clock estimates are
inaccurate, the fossil record is incomplete, and/or the fossil record has been misinterpreted in the
calibration of molecular clock analyses or some combination of the above. To resolve the deep
evolutionary history of sponges we compiled an extensive phylogenomic dataset of sponges, revised
fossil calibration evidence to derive a new phylogeny and timescale for sponge evolution, and
investigated the evolutionary origin of the sponge spicules. Our results confirm that sponges include
two main lineages: Silicea (Hexactinellida plus Demospongiae) and Calcarea plus Homoscleromorpha.
Our molecular clock analyses identify an early Ediacaran estimate for the crown ancestor of sponges.
Our study of spicule evolution shows that neither the last common sponge ancestor, nor the last
common ancestors of Silicea and (Calcarea plus Homoscleromorpha) possessed siliceous spicules,
which independently evolved four times in sponges: once in Hexactinellida, twice in Demospongiae,
and once in Homoscleromorpha. Furthermore, diversification rate analyses demonstrate that the
origin of spicules did not align with species radiations within the phylum. Our dating analyses
substantially reduce the mismatch between molecular clock estimates for the antiquity of sponges and
their fossil record. Our study of the evolutionary history of siliceous spicules reconciles the remaining
difference as a consequence of crown-sponges primitively lacking siliceous spicules. As such we
demonstrate the power of combining molecular and palaeontological approaches to establishing
evolutionary timescales which together allow us to calibrate genome evolution to geological time and
read evolutionary history within the gaps in the fossil record.

Molecular clock estimates and fossil biomarkers suggest that sponges diverged in the Tonian or
Cryogenian[1–5] anticipating a deep Neoproterozoic appearance of sponge fossils. However, while fossils
from the Tonian, Cryogenian and Ediacaran have been claimed to represent early sponges, this
evidence is difficult to interpret[6]. For example, C_{30} steranes interpreted as evidence for Cryogenian
demosponges – that is 24-ipc and 26-mes[3], can also be interpreted as diagenetically altered C_{29} sterols
from chlorophyte algae[7]. Arsenopyrite crystals aggregates, and other fragmentary structures have
been misinterpreted as siliceous sponge spicules[8] while scanning of Tonian fossils that resemble
sponge-like ostia[9], and of the ~600 Ma old *Eocyathispongia qiania* (that was originally interpreted as

an adult asconoid sponge), demonstrated that they lack the internal structures expected in sponges[10].

Accordingly, there is a minimal mismatch of 150 Ma between molecular estimates of divergence times

and fossil evidence for the origin of the sponges, with the earliest, unambiguous, sponge fossils being

represented by loose, late Ediacaran ~543 Ma, siliceous spicules [6,11,12].

Three sponge lineages make spicules out of silicic acid (Hexactinellida, Homoscleromorpha

and Demospongiae). The fourth lineage, Calcarea, makes spicules with calcite. Calcite preservation

potential decreases with age, and it is possible that Neoproterozoic sponges were calcified. However,

even assuming that the earliest sponges might have been calcareous, the oldest fossil sponges are

represented by siliceous spicules, and the evolution of calcareous spicules is not relevant to

understand the evolution of siliceous sponges. For this reason, we will not consider calcite spicule

evolution any further in this study. Silica is chemically stable under low-temperature and high-

pressure conditions[6]. If siliceous spicules existed in the Neoproterozoic we should expect to find them.

Four hypotheses have been proposed to explain the absence of Neoproterozoic siliceous spicules: (1)

Molecular clock estimates of sponge divergence times are not reliable, putative sponge biomarkers

are not diagnostic of sponges, and sponges evolved in the early Cambrian[13]; (2) Sponges emerged in

the Neoproterozoic but early sponge spicules were unmineralized and had low preservation

potential[12]; (3) Neoproterozoic ocean geochemistry and/or depositional regimes were unfavourable

for the preservation of mineralized spicules[14,15]; (4) Spicules were not present in the Last Common

Ancestor (LCA) of sponges and in early sponges more broadly. If this was the case, then siliceous

spicules must have independently evolved multiple times during sponge history[16].

Here we test alternative hypotheses of the age of sponges and the origin of siliceous spicules

using phylogenetic analyses, divergence time estimation, ancestral state estimation (ASE), and

diversification rate shift analyses. We then interpret these results to generate an integrative history

of sponge evolution. We first constructed a time-calibrated phylogenomic tree based on 70

transcriptomes, 12 of which were newly sequenced. We tested the effect of competing calibration

strategies on the timescale of sponge evolution and used ASE to elucidate siliceous spicule evolution.

The results of our analysis indicate that sponges evolved in the early Ediacaran (~608 Ma) and that the

crown ancestors of all extant sponges and of the silicean (i.e. Hexactinellida plus Demosponges)

sponges did not possess silicified spicules. Spicules evolved independently in Homoscleromorpha,

Hexactinellida and Demospongiae. In Demospongiae, the most biodiverse sponge lineage, siliceous

spicules emerged in the stem of Heteroscleromorpha plus Haplocleromorpha (the spiculated

demosponge lineages) after they diverged from the clade composed of Keratosa and Verongimorpha

(the aspiculated demosponge lineages). One further independent (convergent) origin of siliceous

spicules is observed within the aspiculated sponges, where the Verongimorph genus *Chondrilla*

possess siliceous spicules. We performed a focal study, using diversification rate shift analyses, on the

demosponges (the most diverse sponge lineage), and found only two potentially significant

diversification shifts. These happened in Poecilosclerida and the clade comprised of Axinellida plus

Bubarida, both of which are spiculated lineages deeply nested within the Heteroscleromorpha, that

significantly postdate the emergence of siliceous spicules. Siliceous spicules emerged late in sponge

evolution, and they do not seem to have played a significant role in the diversification of the

Demospongiae. While our results imply that the emergence of siliceous spicules in the fossil record

cannot be used as a proxy for the age of Porifera, our new revised estimate of divergence time, in

combination with our study of the evolution of siliceous spicules, *de facto* reconcile the sponge fossil

record with molecular estimates of divergence times.

Results

Phylogenomic and molecular divergence time analyses

Phylogenomic analyses (see Fig S1—S3) recover Silicea (Hexactinellida plus Demospongiae) as the

sister of Calcarea plus Homoscleromorpha. In Demospongiae, we retrieve the aspiculated Keratosa

and Verongimorpha to form the monophyletic sister group of the spicule-bearing

Heteroscleromorpha. Relationships in our phylogeny mostly show full support. The only exception is

the relationships between Suberitida, Poecilosclerida, Clionaida, and Tethyida in Heteroscleromorpha,

where Suberitida clusters either with Poecilosclerida (in the Bayesian Analyses – BA) or with a

Clionaida plus Tethyida (Maximum Likelihood – M; and Astral-based Coalescent analyses – CA; see Fig

S1—S3).

Molecular divergence times (Fig. 1) were inferred for the Bayesian phylogeny (Fig. S3), using

12 fossil constraints to calibrate the ages of major sponge clades (see nodes highlighted with red circle

in Fig. 1 and supplementary material for detailed descriptions of all 12 calibrations). We calibrated the

root to have a hard minimum of 574 Ma and a soft maximum of 609 Ma. The fossil minimum is based

on the earliest occurrence of *Charnia masoni*, in the Drook Formation of Mistaken Point,

Newfoundland[17], while the maximum is based on the Lantian biota of South China which contains a

diversity of macrofossils but no credible metazoans[18]. A uniform prior was used for this calibration,

implying that we maintain an agnostic view of where the root of our phylogeny might lie between

these constraints. The maximum on the root node is soft, with 2.5% of the probability density falling

outside of the 574–609 Ma interval, on the deep side of the distribution. Accordingly, our analyses did

not preclude the inference of a deeper age for the origin of sponges. Eumetazoa was assigned a hard

maximum of 573 Ma. This value was empirically identified to avoid truncation problems in MCMCTree

(see Methods for details) and a soft minimum of 561 Ma based on age of Bed B in the Bradgate

Formation from which the earliest crown-eumetazoan, *Auroralumina attenboroughii*, was

recovered[19,20]. While it was possible to establish minimum constraints for the remaining sponge

calibrations, it was not possible to justify maximum constraints based on palaeontological,

biogeographic or geological evidence. The sponge fossil record, at least in terms of records that can

inform calibrations for molecular clock analyses, is based on sites of exceptional fossil preservation.

As such, we have no prior expectation of whether the earliest records of various sponge clades (Details

in Supplementary Information) represent close approximations of true clade ages. Cauchy

distributions with a heavy-tail (see methods) were used for these calibrations, to reflect the lack of

knowledge for maxima.

Two calibration strategies were used. First, we performed analyses where the oldest known

siliceous spicule was used to constrain the crown Heteroscleromorpha clade. We refer to this

calibration strategy as the Spiculated Demosponges Node (SDN) strategy. We then performed an

analysis where the same calibration was used to calibrate the root of Porifera – Spiculated Porifera

Node (SPN) strategy. These calibration strategies were implemented to reflect competing

interpretations of the fossil record of siliceous spicules. Under the SDN strategy, early siliceous fossil

spicules are assumed to represent remains of spicule-bearing demosponges, as suggested by the

morphological similarity between the oldest classifiable siliceous spicules[7,21] and those found in living

demosponges. Under the SPN strategy, siliceous spicules are assumed to have been present in the last

common ancestor of Porifera, with the spicule fossil record providing a bound on the origin of sponges.

We visualised the joint time priors (Fig. S4), to make sure that there was no major disagreements

between our specified and effective priors (see Fig. S5).

We then performed molecular clock analyses, under both calibration strategies (SDN and

SPN), using both the Autocorrelated Rates (AR)[21,22] relaxed molecular clock model, and Independent

rates (IR)[23,24] relaxed molecular clock model, and the Independent Rates (IR)[22,23] relaxed molecular

clock model. For each analysis (a total of four SDN-AR; SDN-IR; SPN-AR and SPN-IR), we run six

independent chains. We checked for convergence and dismissed results from chains that did not

converge, Fig. S6 and Methods for details. Under the SDN strategy, three chains converged (and were

therefore retained) under both the AR and IR models (Table S1).

Results inferred using the two clock models and different calibration strategies did not differ

significantly, and here we only discuss results from the SDN-IR analysis. Results from all other analyses

can be seen in Supplementary Results (Fig. S7 to Fig. S9; Table S3-S4). Our SDN-IR analysis inferred

the last common metazoan ancestor to have emerged between 616.9-605.9 Ma (early Ediacaran)

(Table 1). Similarly, the sponge LCA was dated to 615.3–601.3 Ma (early Ediacaran). Crown-group

Silicea (Hexactinellida + Demospongiae) was estimated at 591.3–570.3 Ma (middle Ediacaran),

Lyssacinosida (within Hexactinellida) at 457.3–445 Ma (late Ordovician), and the last common
ancestor of Calcarea and Homoscleromorpha at 566.5–487.6 Ma (late Ediacaran to late Cambrian).
Homoscleromorpha was estimated to have originated at 429.1–228.6 Ma (middle Silurian – late
Triassic), and Calcarea at 351.7–213.89 Ma (early Carboniferous – late Triassic). The last common
demosponge ancestor is estimated at 559.17–536.53 Ma (late Ediacaran – early Cambrian), and the
last common ancestor of the non-spiculated Verongimorpha and Keratosa at 521.85–430.44 Ma (early
Cambrian – middle Silurian). The last common ancestor of the spicule-bearing demosponges
(Heteroscleromorpha) is estimated to have existed between 521.97–515.05 Ma (middle Cambrian),
and the last common ancestor of the Haplosclerida was inferred at 403.84–246.54 Ma (early Devonian
– middle Triassic). The last common ancestor of the rest of the Heteroscleromorpha was inferred to
have existed 503.17–412.61 Ma (middle Cambrian – early Devonian). Within Heteroscleromorpha, the
freshwater Spongillida separated from the marine Scopalinida at 424.69–325.85 Ma (late Silurian –
late Carboniferous), while the three most speciose heteroscleromorph lineages (Tetractinellida,
Poecilosclerida and Spongillida) diverged, respectively, 299.37–231.07 (early Permian – late Triassic),
236.53–199.74 Ma (late Triassic – early Jurassic), and 123.21–71.52 Ma (Cretaceous).
181

Ancestral state estimation of siliceous spicules

Ancestral state estimation (ASE) results support the absence of spicules at the root of Porifera.
Absence is the most likely state irrespective of the calibration strategy used, ASE model used, or
whether characters were treated as independent or amalgamated. Absence of spicules at the root of
Porifera is strongly supported by the four-state model and the amalgamated Structured Markov
Model (SMM) when averaged using BIC (Fig. 2). Results are more ambiguous when using the two-state
model, or if any model is averaged using uncertainty (see Fig. S10-S11). However, in all these cases,
absence of spicules at the root of Porifera remains the most probable state (see Fig. S10-S11). The
dating method had little effect on the ASE results. Using the SMM we were able to accommodate the
hierarchical nature of the data, the results amalgamated with the BIC (Fig. 2) show that the LCA of
Silicea most likely did not have mineralised siliceous spicules and that siliceous spicules independently
evolved three times in Silicea and once in Homoscleromorpha. In Silicea, siliceous spicules evolved
once in the stem hexactinellid lineage, once in the lineage leading to Heteroscleromorpha, and once
in the verongimorph genus *Chondrilla*. Based on our sponge timetree, we can conclude that
demosponge spicules emerged in a ~31 Ma interval between 548 and 517 Ma (Fig. 1), in accordance
with the fossil record. In Homoscleromorpha, silicified spicules were only acquired by the Plakinidae,
after the divergence from Oscarellidae in the Carboniferous. When the results are amalgamated using
raw uncertainties, the sponge and Silicea LCAs are still reconstructed to have been, most likely,

without spicules. Similarly, an independent origin of siliceous spicules in Homoscleromorpha

continues to be supported (see Fig. S11).

Diversification events in Silicea

Because three independent origins of silicified spicules are inferred in Silicea, we tested the effect of

the origin of silicified spicules in sponge evolution inferring diversification rates shifts in this lineage.

To achieve this goal, following DeBiasse et al.[25], we first generated a megaphylogeny of Silicea

performing a constrained ML analysis of 807 *Cytochrome c Oxidase 1* (COI) barcodes, using our

phylogenomic tree (Fig. 1 and S3) as a backbone constraint (see Methods for details). The

megaphylogeny was then dated (see Methods) and used to perform diversification analyses using

BAMM[26] and Medusa[27] (Fig. S12 and Table S5; see Methods for details). BAMM identified two

diversification shifts in both the AR and IR timetrees (Table S6). In the AR timetree, rate shifts are

identified on the branches leading to the Axinellida plus Bubarida, and at the origin of the

Poecilosclerida (Fig. 3A). In the IR topology, rate shifts were inferred on the branch leading to

Hexactinellida, and again on the branch leading to the LCA of Axinellida and Bubarida (Fig. 3C). Using

Medusa diversification rate shifts were identified in the branches leading to Axinellida plus Bubarida

and Poecilosclerida for both models (Fig. 3B-D), with the IR tree also identifying a shift in

Dictyoceratida (Fig. 3D).

Discussion

Mineralized sponge spicules are not homologous

Our phylogenomic analyses confirmed that sponges are split into two fundamental lineages, Calcarea

plus Homoscleromorpha, and Demospongiae plus Hexactinellida (Silicea). As suggested by previous

studies[28], within Demospongiae, we recovered the monophyly of Keratosa, Verongimorpha, and

Heteroscleromorpha. Consistent with previous phylogenomic and mitochondrial genomic studies[28,29],

we found the spicule-less Keratosa and Verongimorpha to be each other's sister lineage, and the

spicule-bearing Heteroscleromorpha are found to represent the sister of Keratosa plus

Verongimorpha. A long-standing problem in Demospongiae phylogeny is that of the relationships of

Bubarida[10].   Our   tree   recovers   Bubarida   as   the   sister   of

(Agelasida+Plymastiida+Tethyida+Clionaida+Suberitida+Poecilosclerida).   However, we could not

resolve the relationships of Suberitida with reference to Clionaida, Tethyida, and Poecilosclerida, see

Fig. S1-S2.

All our molecular clock analyses recovered an Ediacaran origin of sponges and a Cambrian

origin for the spicule bearing demosponges (Heteroscleromorpha). Taxon sampling does not allow us

to date the origin of Hexactinellida. However, the oldest known Cambrian and latest Ediacaran sponge

spicules do not bear resemblance to the hexactinellid spicules, due to the different shape of the axial

filament, which always has a square cross-section in Hexactinellida, a shape never found in

Demospongiae or in the oldest fossil spicules. Our ASEs inferred that siliceous spicules were absent

from the root of sponges, Silicea, and Homoscleromorpha, suggesting instead independent origins for

this trait in the stem Hexactinellida, within Homoscleromorpha (in the stem Plakinidae), and in

Demospongiae (in the stem Heteroscleromorpha) and in the verongimorph genus *Chondrilla*.

Cambrian siliceous spicules are comparable to those of extant spicule-bearing demosponges [15,30], and

given also the results of our ASE analyses, we can only conclude that latest Ediacaran and Cambrian

siliceous spicules were from species belonging to total group Heteroscleromorpha plus

Haploslceromorpha, postdating the deepest divergences in Porifera.

245   To better understand whether siliceous biomineralisation could have evolved multiple times

independently, we turned to biochemical evidence. It has been shown that silica biomineralisation is

the result of different processes in different sponge lineages[16,31–33]. Demosponges use silicateins

enzymes[34–36] and hexactinellids use glassin, perisilin, and hexaxillin with the organic part of the spicule

(the axial filament) having different composition[33]. *In vitro* experiments in demosponges, particularly

with *Suberites domuncula*, have demonstrated that the self-assembly of silicateins is a fundamental

step in the process of spicule biogenesis. This process creates scaffolds where silica is immediately

deposited[36]. The implication is that, at the least in demosponges, spicules seem to only develop as

biomineralised structures, precluding the possibility that siliceous spicules could have evolved from

organic (non biomineralised) ones[12,33].

The appearance of silicified spicules did not drive diversification events

Sponges are globally distributed and comprise ~9,600 species[37]. In terms of biodiversity,

Demospongiae has undoubtedly been the most successful sponge lineage[37]. Furthermore, they have

been ecologically more successful being able to transition to freshwater and undergo revolutionary

body plan rearrangements, leading to the only known case of carnivory in the phylum[38,39]. Many

explanations have been proposed to explain their success. One is that the high diversity of spicule

shapes would have been advantageous for structural functions and defence, driving adaptive

radiations. Our diversification analyses however highlight a different scenario. While spicules certainly

play a role in defence and for structural support, their emergence did not seem to have driven adaptive

radiations. If that was the case, we would have expected a diversification event on the branch that

leads to the radiation of Heteroscleromorpha. Instead, we retrieve diversification events, consistently

across BAMM and Medusa, on the branch that leads to Bubarida and Axinellida (Fig. 3). Both these

groups are well nested within Heteroscleromorpha, and spicules were not an innovation specific to
them. Furthermore, their spicule types are shared across many other orders in demosponges (Table
S5) where we did not identify diversification events. The increase in diversification rates highlighted
in these two lineages might be linked to their ecology, however, it would be too speculative to attempt
understanding what aspects of their ecology might have driven their radiation.

Medusa highlighted a diversification event on the stem Poecilosclerida branch for both the AR
and IR timetrees. While Poecilosclerida represents the most speciose demosponge order, with ca.
2,488 species described to date (World Porifera Database[37]), have a worldwide distribution, and
achieved substantial ecological diversity (evolving also carnivory – family Cladorhizidae), we prefer to
maintain a conservative view and consider this result uncertain because it is software-specific. In any
case, Poecilosclerida are also members of the spiculated lineage and did not evolve spicules
independently. We therefore conclude that while the poecilosclerid evolutionary success might have
been linked to the massive expansion of microscleres observed in this clade[40,41], the origin of spicules
itself cannot have driven their diversification.

An Ediacaran origin of sponges
Irrespective of the fossil calibration strategy used, our divergence time estimates identified the origin
of sponges at a time that is significantly younger than previous studies (617.91-581.39 Ma early-
middle Ediacaran)[2,4,42]. Our new, younger, divergence times are driven by both a revised
geochronology and reinterpretation of the fossil evidence used in establishing calibrations[43]. The
earliest fossils that can be assigned to sponges are loosely mineralised siliceous spicules that first occur
at the Ediacaran-Cambrian boundary[11–13,44]. Our results indicate that these spicules are uninformative
of early sponge evolution. The absence of fossilised spicules before this time does not represent a
taphonomic bias due to Neoproterozoic Ocean geochemistry making siliceous spicules difficult to be
buried or modified[12], as confirmed by the presence of abundant cherts in the latest Ediacaran[45].
Rather, our results indicate that stem- and early crown Porifera lacked spicules (like extant
Verongimorpha and Keratosa), and Ediacaran sponges were therefore entirely soft-bodied. The
hypothesis that biomineralised sponges were predated by sponges with an organic skeleton has
previously been proposed[10] to explain the lack of precambrian sponge fossils. This hypothesis is not
supported by the current understanding of siliceous spicule development, as biomineralisation seems
to be a fundamental step in spicule biosynthesis (see above), and we therefore suggest that this
hypothesis should be considered rejected.

Since sponges branch early within the animal phylogeny (irrespective of the current
controversy on the relative relationships between sponges and ctenophores in the context of early

animal phylogeny[29,46–53]) they are informative regarding early metazoan evolution, and the origin of

animal multicellularity. Advocates of a literal reading of the fossil record would suggest that the origin

of sponges (and of Metazoa) corresponds closely with the first appearance of siliceous spicules in the

fossil record (~543 Ma; latest Ediacaran)[13]. However, molecular timescales derived using relaxed

molecular clock methods have dated the origin of sponges (and that of animals) in the Cryogenian (85

Ma before the Ediacaran)[2]. Some possible sponge body fossils have previously been suggested to

expand the fossil record of sponges into the Cryogenian[9]. However, these records are more

speculative than informative, based as they are based on fragmentary remains. Fossilised steranes

(24-ipc and 26-mes) from the late Cryogenian (660-635 Ma) have enjoyed a long history of alternative

interpretations[54–57]. While originally suggested to indicate the presence of demosponges in the

Crypogenian, they have recently been interpreted also to represent diagenetic derivatives of C_{29}

sterols from chlorophyte algae[7] (Bobrovskiy et al. 2021). Interpreting these steranes as demosponge

biomarkers would have material implications on our evolutionary timescale. However, current

ambiguity over their nature implies that, at the least at the time of writing, they cannot be used as

calibrations for molecular clock analyses.

The absence of siliceous sponge spicules from the fossil record prior to the latest Ediacaran is

especially informative since they require no special mechanisms for fossilization[2]. Rather, the absence

of Tonian, Cryogenian or deep Ediacaran sponge spicules requires either that sponges had not yet

evolved or else that sponge spicules evolved late within the sponge lineage. Our results provide a

reconciliation of the sponge fossil record with the molecular timescales of sponge evolution. First, our

ASEs show that siliceous spicules emerged relatively late in sponge evolution, in the stem of the

spiculated demosponges (heteroscleromorphs). The root of Silicea and of the sponges predate the

oldest fossil spicules and our results suggest that early crown sponges existed for ~75–38 million years

before the origin of spicules, this interval might diminish with the discovery of increasingly older

spicule records. This result finally implies that current estimates of divergence time for Metazoa[4,42]

are likely to be overestimated.

Material and methods

Sample collection, RNA extraction, library preparation, Illumina sequencing

Twelve novel sponge specimens were collected for RNA extraction (Supplementary material Table S).

Samples were preserved in RNAlater (Thermo Scientific, Waltham, USA) using an overnight incubation

at 4°C before long-term storage at -80 °C. Total RNA was extracted with TRIzol (Ambion, Austin, USA)

and mRNA purified with a Dynabeads mRNA DIRECT kit (Thermo Scientific, Waltham, USA). The cDNA

libraries were constructed with either the kits TruSeq v2 or Scriptseq v2 RNA Library Prep kit (Illumina),

according to the manufacturer's instructions, using the maximum quantity of mRNA allowed, which

was 50ng – 100ng of starting material. The quality of the libraries was checked with an Agilent

TapeStation 2200 system (Agilent Technologies) and the quantity with Qubit™ dsDNA HS Assay kit

(ThermoFisher Scientific). The sequencing was conducted with an Illumina NextSeq 500 platform at

the Sequencing Facilities of the Natural History Museum of London (Research Core Labs), using a

paired-end read strategy (bp length: 2x150 bp).

Transcriptome assembly and annotation

Over the 12 novel transcriptomes, 58 published RNA-seq data sets with Sequence Read Archive (SRA)

IDs were include, with the final total dataset was composed of 70 proteomes (Table S7). For all

transcriptomes assembled, read quality assessment was performed with FastQC[58] with trimming

performed with Trimmomatic[59] with the following settings: ILLUMINACLIP:./Adapters.fa:2:30:10

LEADING:3 TRAILING:3 SLIDINGWINDOW:4:28 MINLEN:36, with the Adapters.fa file adjusted to

include the adapter sequences specific to each read pair. Assembly of the clean paired reads, obtained

from Trimmomatic, was performed using Trinity version[60] with the following settings: --

normalize_reads –CPU 8 –inchwormcpu 8. Prior to orthology inference, nucleotide sequences were

translated using TransDecoder (https://hpc.nih.gov/apps/TransDecoder.html) for a total of 70 species

and isoforms were collapsed using CD-HIT[61]. The final total dataset was composed of 70 proteomes

(Table S7).

Orthology inference

Orthologs were inferred using Orthofinder v. 2.7.1[62], which identified 335,892 orthogroups. Putative

paralogous were removed using PhyloTreePruner[63], only orthogroups with at least 90% of the species

in the total dataset were retained for further analyses to minimise missing data for a total of 1,468

orthogroups retained. Hypervariable regions of the sequences were removed using PREQUAL[64].

Orthogroups were aligned using MUSCLE[65]. Final trimming was performed with TrimAl[66] with -auto

option for each gene family. After trimming, sequences shorter than 25 AA and with less then 50% of

species were removed with Al2Phylo[29]. Single-gene trees were built using IQTree v 1.6.12[67], where a

model test was run for each gene tree with the parameters -m MFP -mset LG+F+G, WAG+F+G,

JTT+F+G, GTR20 -madd LG+C20+F+G, LG+C10+F+G, LG+C30+F+G, LG+C40+F+G, LG+C50+F+G,

LG+C60+F+G, C10, C20, C30, C40, C50, C60, LG4M, LG4X , to allow selecting across compositionally

site homogeneous and site heterogeneous models. We further used an in-house developed script

(https://github.com/mgiacom/ParaFilter) by Mattia Giacomelli to remove putative hidden paralogs

and extremely long branched taxa. Finally, removed gene trees that could not identify the monophyly

of sponges or of the outgroup lineages using monophylyl.pl from

https://github.com/MaxTelford/Xenacoelomorpha2019. The final dataset was composed of 133

genes that were concatenated in a super alignment using the catfasta2phyml.pl script from

https://github.com/nylander/catfasta2phyml, for a total length of 103,269 AA.

Phylogeny inference

The superalignment was analysed using ML in IQTree2 v2.1.3[68]. Model testing was performed using

Model Finder plus (MFP option in IQTree but considering only across-site compositionally

homogeneous models)[69]. Across-site compositionally heterogeneous models were not considered

because analyses using such models were performed in Phylobayes instead (see below). Model

selection identified Q.INSECT+I+G4 model as the best fit across-site compositionally homogeneous

models, we used the ultrafast bootstrap (1,000 replicates)[70,71] to estimate support for the nodes in the

across-site compositionally homogeneous ML trees. In addition, we ran Phylobayes mpi.1.8[72] under

the CAT Poisson model[73] with support values estimated using Posterior Probabilities (PP). We ran two

independent chains for over 7,000 generations (Burnin = 1500). Convergence was tested using

Bpcomp (maxdiff = 0.0018315) and *Tracecomp* (minimal ESS 61 and maximum real_diff=224488)

which are part of Phylobayes (See details in table S8). Finally, we used ASTRAL[74] as a coalescent tree

reconstruction method of the final 133 gene trees, with default parameters.

Timetree inference

Based on the results of ModelFinder analyses performed on the single gene alignments during the

gene trees reconstruction described in the orthology inference section, a partitioned dataset was

generated (with five partitions – see Supplementary methods for justifications).   Prior to timetree

inference, we calculated the gradient (i.e., vector of first derivatives of the likelihood function) and

the Hessian (i.e., matrix of second derivatives of the likelihood function) for each partition with

CODEML, which is part of the PAML package v4.9i[75]. The vector and matrix are necessary to implement

approximate maximum likelihood estimation[75](see supplementary information for details).

We fixed the tree topology to our inferred Bayesian tree (Fig. S3, S12 and Fig. 1), and

constrained the age of 12 nodes representing major sponge clades in our phylogeny (red dots and

circles in Fig. 1) based on the fossil record. The oldest fossil spicule was initially used to calibrate the

origin of the Spicule-bearing Demospongiae (SDN calibration strategy). We used a uniform prior on

the root age with a soft maximum of 609 Ma (pU = 0.025) and a hard minimum of 574 Ma (pL = 1e-

300; this small value enables the usage of hard bounds in MCMCtree, pL=0 is not allowed as it would

cause numerical underflow). We constrained the age of Eumetazoa with a hard maximum of 573 Ma

(pU = 1e-300) and a soft minimum of 561 Ma (pU = 0.025) to avoid truncation issues during the MCMC

runs[42]. The age of the other ten calibrated nodes was constrained using heavy-tail Cauchy distributions

(i.e., lower-bound calibrations with an offset of p=0.1 or p=0.5, a scale parameter of c=0.5 or c=0.1,

and a left tail probability of pL=1e-300). The use of a heavy tailed distribution was preferred as it allows

to capture our interpretation of the reliability of the sponge fossil record, which is unlikely to be

reliable in deep time because of difficulties for a taxonomic classification given the alteration of

diagnostic characters by taphonomic processes. We performed a sensitivity test (SPN strategy) where

the oldest siliceous spicule was reassigned to calibrate the node representing the last common sponge

ancestor (see also Fig. 1) to investigate what effect, uncertainty in the spicule fossil record, could have

on molecular estimates of poriferan divergence times. In brief SDN strategy constrained the spicule

bearing Demospongiae (Haploscleromoprha + Heteroscleromoprha) to have a hard minimum at 515

Ma, while the SPN strategy constrained the root of Porifera to a minimal age of 515 Ma. Both

experiments used a Cauchy distribution specified in the same way (c= 0.5, p = 0.5, and pL=1e-300).

The age of 515 used in both the SDN and SPN strategies is based on a loose siliceous spicule from the

lower Cambrian (Series 2, Stage 3;) Sirius Passet Biota of North Greenland[30]. A list of all the calibrations

used can be found in Supplementary Information. We ran all analyses under both the IR[71,72] and the

AR[73,74] relaxed-clock models implemented in the MCMCtree (PAML package v4.9i[75]). Six independent

chains for every analysis (SDN+AR, SDN+IR, SPN+AR, SPN+IR) were run with 4,020,000 iterations each.

We discarded the first 20,000 as burn-in and subsequently subsampled every 20 iterations. We saved

the values sampled for each parameter until reaching a total of 200,000 samples for each chain. As

per our MCMC settings, we discarded the first 100,000 collected samples as part of the burn-in phase

and, every 100 iterations, we saved the values sampled for each parameter until reaching a total of

20,000 samples for each chain. The MCMC settings for the evolutionary model can be found in the

supplementary material. We used our in-house pipeline (see

https://github.com/MEleonoraRossi/spicules-dating) to (i) filter the six MCMC chains we ran for every

data analysis and (ii) generate convergence plots and summary statistics with the chains that passed

our quality control checks (see supplementary material and GitHub repository for details).

Ancestral state estimation (ASE)

We conducted marginal likelihood ASE in R using the fitmk() and ancr() functions from the phytools

package[76]. Character states for the full dataset were retrieved from the literature (Table S9). We coded

two discrete characters: (i) presence of mineralised spicules, absent = 0, present = 1; (ii) mineralogy

of the spicules, absent = 0, calcite = 1, aragonite = 2, silica = 3. For each character, different transition

models were considered: equal rates (ER), symmetrical (SYM) and all rates different (ARD). Note that

the symmetrical model is equivalent to the "all rates are different" model for binary characters and
was thus not used in these analyses. Observing spicule mineralogy is dependent upon spicule
presence. To accommodate the hierarchical nature of this data, we conducted a third ASE analysis in
which the characters were amalgamated into a single Structured Markov Model (SMM) equipped with
hidden states[77]. We tested 14 different SMMs representing seven patterns of transition rate
heterogeneity (Table S10) and two patterns of character correlation: general correlation (SMM_ind)
and switch-on dependency (SMM_sw). All ASE analyses were repeated using four trees resulting from
four different molecular clock analyses (SDN+AR, SDN+IR, SPN+AR, SPN+IR). Results were summarised
using model averaging. Models were weighted using two metrics: (i) Bayesian Information Criterion
(BIC); (ii) raw uncertainty of ancestral estimates[78].

Diversification Rates through Time and Across Clades
We assembled a dataset of 804 COI sequences from GenBank. The sequences were visualised using
Seaview and aligned with MUSCLE v. 3.8.31[65]. The final alignment was 609 bp. We inferred the best-
scoring maximum-likelihood (ML) tree with IQTree2 v2.1.3[68]. However, we constrained the analyses
using our phylogenomic tree (Fig. 1) as a backbone constraint[25]. Model testing was performed using
Model Finder plus (MFP option in IQTree but considering only across-site compositionally
homogeneous models)[69], because of the size of the dataset which limited our ability to use complex
models. Following the procedure detailed in Álvarez-Carretero et al., 2022[79], we fitted skew-t (ST)
distributions to the posterior densities of the divergence times inferred using PAML v4.9i[75] (see
Supplementary Method). We selected 13 ST distributions that matched the nodes that had been
calibrated prior to timetree inference so that we could use them as prior distributions in the
subsequent analyses (Table S11) on the ML topology inferred using the 804 COI sequences. We run 64
independent chains with 4,020,000 iterations each. We discarded the first 20,000 as burn-in and
subsequently subsampled every 20 iterations. We saved the values sampled for each parameter until
reaching a total of 200,000 samples for each chain. Convergence was calculated as explained in the
Time tree inference section above (Table S5).
Macroevolutionary dynamics of diversification were modelled with the software Bayesian
Analysis of Macroevolutionary Mixtures (BAMM) v.2.5.0[26] on the timetree inferred following the
procedure described above. We accounted for incomplete taxon sampling by defining a sampling
fraction of 5%, which corresponds to the number of species included in the COI dataset compared to
the total number of Silicea species accepted. For each order we also calculated the sampling
frequences (species sampled/species accepted for the order; data obtained by the World Porifera
Database; Table S12). Ten million generations of reversible jump Markov Chain Monte Carlo sampling

were run, drawing samples from the posterior every 10,000 generations. For the prior probability of

rate shift, we tested values ranging from 0.1 to 50 and chose the value leading to the highest ESS

values for LogLikelihood and NumberOfShifts (Table S4). We processed the output data with

BAMMtools[80] and, after removing the samples collected during the burn-in process (10% of the

samples), we obtained the summary statistics, and plotted the diversification rate over time. To

overcome the possible bias introduced by incomplete sampling in diversification analyses and to

address the debate around the application of BAMM[26], we confirmed our results using the taxonomic

method implemented in MEDUSA[27] where clades are collapsed to terminal lineages of equal ranks.

Only diversification events inferred by both methods were deemed reliable.

Data availability

The IDs of genomes and transcriptomes used for the dataset are in table S7 in the supplementary

material. The raw reads of the 12 RNA-Seq data generated in this study will be uploaded on the SRA

upon acceptance.

Code availability

The in-house pipeline used for the time tree inference can be found on GitHub at

https://github.com/MEleonoraRossi/spicules-dating

All code used for analysis and figure generation will be released on GitHub at

https://github.com/MEleonoraRossi/spicule-evolution upon acceptance.

Ethics statement

The authors declare no competing interests.

Authors' contributions

MER, DP, AR and PJD conceived the study. MER performed the phylogenomic, timetree inference, and

diversification event analyses, and wrote the first draft of the paper. DP, PJD and AR edited the paper.

MG, NJK, and MER performed orthology inference. JNK provided support for the ASE analyses, SAC

provided support and scripts for timetree inference analyses. PC and AS provided some of the fossil

calibrations and helped assembling the morphological matrix used in the ASE. ST, VK, AR, and PC took

part in the sampling. ST, VK, NJK and MER extracted the RNA and performed library preparations of

the samples. All authors have read and wrote the manuscript.

Acknowledgments

The authors want to thank Dr. Bruna Plese for her help during the RNA extraction and library

preparation. MER is supported by a NERC GW4+ PhD studentship. DP and PJD are supported by the

John Templeton Foundation (62220). Please note that the opinions expressed in this publication are

those of the author(s) and do not necessarily reflect the views of the John Templeton Foundation, and

the Gordon and Betty Moore Foundation (GBMF9741). DP is supported by a University of Bristol URF

fellowship. PJD was supported by a Leverhulme Trust fellowship (RF-2022-167), and a BBSRC grant

(BB/T012773/1). DP and PJD were also supported by a NERC-BETR Grant (NE/P013643/1). NJK

supported by Rutherford Discovery Fellowship RDF-UOO2001 from the Royal Society of New Zealand.

JNK is supported by ERC grant no. 788203 (INNOVATION). AS is supported by Villum Fonden grant no.

679849 and Enbie grant no. 54433.

References

1. Gold, D. A. *et al.* Sterol and genomic analyses validate the sponge biomarker hypothesis. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 2684–2689 (2016).

2. Sperling, E. A., Robinson, J. M., Pisani, D. & Peterson, K. J. Where's the glass? Biomarkers, molecular clocks, and microRNAs suggest a 200-Myr missing Precambrian fossil record of siliceous sponge spicules: Sponge biomarkers, molecular clocks and microRNAs. *Geobiology* **8**, 24–36 (2010).

3. Zumberge, J. A. *et al.* Demosponge steroid biomarker 26-methylstigmastane provides evidence for Neoproterozoic animals. *Nat Ecol Evol* **2**, 1709–1714 (2018).

4. Erwin, D. H. *et al.* The Cambrian Conundrum: Early Divergence and Later Ecological Success in the Early History of Animals. *Science* **334**, 1091–1097 (2011).

5. Dohrmann, M. & Wörheide, G. Dating early animal evolution using phylogenomic data. *Sci Rep* **7**, 3599 (2017).

6. Botting, J. P. & Muir, L. A. Early sponge evolution: A review and phylogenetic framework. *Palaeoworld* **27**, 1–29 (2018).

7. Bobrovskiy, I. *et al.* Algal origin of sponge sterane biomarkers negates the oldest evidence for animals in the rock record. *Nature Ecology & Evolution* **5**, 165–168 (2021).

8. Zhang, Y., Yuan, X. & Yin, L. Interpreting Late Precambrian Microfossils. *Science* **282**, 1783–1783 (1998).

9. Maloof, A. C. *et al.* Possible animal-body fossils in pre-Marinoan limestones from South Australia. *Nature Geosci* **3**, 653–659 (2010).

10. Yin, Z. *et al.* Sponge grade body fossil with cellular resolution dating 60 Myr before the Cambrian. *Proc. Natl. Acad. Sci. U.S.A.* **112**, (2015).

11. Chang, S., Feng, Q., Clausen, S. & Zhang, L. Sponge spicules from the lower Cambrian in

the Yanjiahe Formation, South China: The earliest biomineralizing sponge record.

Palaeogeography, Palaeoclimatology, Palaeoecology **474**, 36–44 (2017).

12. Tang, Q., Wan, B., Yuan, X., Muscente, A. D. & Xiao, S. Spiculogenesis and

biomineralization in early sponge animals. *Nat Commun* **10**, 3348 (2019).

13. Antcliffe, J. B., Callow, R. H. T. & Brasier, M. D. Giving the early fossil record of sponges a

squeeze: The early fossil record of sponges. *Biol Rev* **89**, 972–1004 (2014).

14. Borchiellini, C. *et al.* Sponge paraphyly and the origin of Metazoa. *J of Evolutionary*

Biology **14**, 171–179 (2001).

15. Chang, S., Zhang, L., Clausen, S., Bottjer, D. J. & Feng, Q. The Ediacaran-Cambrian rise of

siliceous sponges and development of modern oceanic ecosystems. *Precambrian*

Research **333**, 105438 (2019).

16. Francis, W. R. *et al.* The genome of the reef-building glass sponge *Aphrocallistes vastus*

provides insights into silica biomineralization. *R. Soc. open sci.* **10**, 230423 (2023).

17. Dunn, F. S. *et al.* The developmental biology of Charnia and the eumetazoan affinity of

the Ediacaran rangeomorphs. *Science advances* **7**, eabe0291 (2021).

18. Yang, C. *et al.* Implications for Ediacaran biological evolution from the ca. 602 Ma

Lantian biota in China. *Geology* **50**, 562–566 (2022).

19. Wilby, P. R., Carney, J. N. & Howe, M. P. A rich Ediacaran assemblage from eastern

Avalonia: evidence of early widespread diversity in the deep ocean. *Geology* **39**, 655–

658 (2011).

20. Noble, S. R. *et al.* U-Pb geochronology and global context of the Charnian Supergroup,

UK: constraints on the age of key Ediacaran fossil assemblages. *Bulletin* **127**, 250–265

(2015).

21. Thorne, J. L., Kishino, H. & Painter, I. S. Estimating the rate of evolution of the rate of

molecular evolution. *Mol Biol Evol* **15**, 1647–1657 (1998).

22. Yang, Z. & Rannala, B. Bayesian estimation of species divergence times under a

molecular clock using multiple fossil calibrations with soft bounds. *Mol Biol Evol* **23**,

212–226 (2006).

23. Rannala, B. & Yang, Z. Inferring speciation times under an episodic molecular clock. *Syst*

Biol **56**, 453–466 (2007).

24. Lemey, P., Rambaut, A., Welch, J. J. & Suchard, M. A. Phylogeography takes a relaxed

random walk in continuous space and time. *Mol Biol Evol* **27**, 1877–1885 (2010).

25. DeBiasse, M. B. *et al.* A cnidarian phylogenomic tree fitted with hundreds of 18S leaves.

bioRxiv 2022–10 (2022).

26. Rabosky, D. L. Automatic Detection of Key Innovations, Rate Shifts, and Diversity-

Dependence on Phylogenetic Trees. *PLOS ONE* **9**, e89543 (2014).

27. Alfaro, M. E. *et al.* Nine exceptional radiations plus high turnover explain species

diversity in jawed vertebrates. *Proceedings of the National Academy of Sciences* **106**,

13410–13414 (2009).

28. Lavrov, D. V. *et al. Phylomitogenomics Bolsters the High-Level Classification of*

Demospongiae (Phylum Porifera).

http://biorxiv.org/lookup/doi/10.1101/2023.06.04.543631 (2023)

doi:10.1101/2023.06.04.543631.

29. Simion, P. *et al.* A Large and Consistent Phylogenomic Dataset Supports Sponges as the

Sister Group to All Other Animals. *Current Biology* **27**, 958–967 (2017).

30. Botting, J. P., Cárdenas, P. & Peel, J. S. A crown-group demosponge from the early

Cambrian Sirius Passet Biota, North Greenland. *Palaeontology* **58**, 35–43 (2015).

31. Voigt, O. *et al.* Spicule formation in calcareous sponges: Coordinated expression of

biomineralization genes and spicule-type specific genes. *Sci Rep* **7**, 45658 (2017).

32. Riesgo, A., Maldonado, M., López-Legentil, S. & Giribet, G. A Proposal for the Evolution

of Cathepsin and Silicatein in Sponges. *J Mol Evol* **80**, 278–291 (2015).

33. Shimizu, K. *et al.* Silica-associated proteins from hexactinellid sponges support an

alternative evolutionary scenario for biomineralization in Porifera. *Nature*

Communications **15**, 181 (2024).

34. Cha, J. N. *et al.* Silicatein filaments and subunits from a marine sponge direct the

polymerization of silica and silicones in vitro. *Proceedings of the National Academy of*

Sciences **96**, 361–365 (1999).

35. Shimizu, K., Cha, J., Stucky, G. D. & Morse, D. E. Silicatein α: Cathepsin L-like protein in

sponge biosilica. *Proceedings of the National Academy of Sciences* **95**, 6234–6238

(1998).

36. Müller, W. E. G. *et al.* Analysis of the axial filament in spicules of the demosponge

Geodia cydonium: Different silicatein composition in microscleres (asters) and

megascleres (oxeas and triaenes). *European Journal of Cell Biology* **86**, 473–487 (2007).

37. de Voogd, N. *et al.* World Porifera Database. Accessed at

https://www.marinespecies.org/porifera on yyyy-mm-dd. VLIZ

https://doi.org/10.14284/359 (2023).

38. Hestetun, J. T. *et al.* The systematics of carnivorous sponges. *Mol Phylogenet Evol* **94**,

327–345 (2016).

39. Vacelet, J. & Boury-Esnault, N. Carnivorous sponges. *Nature* **373**, 333–335 (1995).

40. Łukowiak, M. *et al.* The terminology of sponge spicules. *Journal of Morphology* **283**,

1517–1545 (2022).

41. Morrow, C. & Cárdenas, P. Proposal for a revised classification of the Demospongiae

(Porifera). *Front Zool* **12**, 7 (2015).

42. dos Reis, M. *et al.* Uncertainty in the Timing of Origin of Animals and the Limits of

Precision in Molecular Timescales. *Curr Biol* **25**, 2939–2950 (2015).

43. Carlisle E., Yin Z., Pisani D. and Donoghue PCJ. Ediacaran origin and Ediacaran-Cambrian

diversification of Metazoa.

44. Chen, C. *et al.* New sponge spicules from the Ediacaran-Cambrian transition in deep-

water facies of South China. *Palaeogeography, Palaeoclimatology, Palaeoecology* **627**,

111714 (2023).

45. Muscente, A., Michel, F. M., Dale, J. G. & Xiao, S. Assessing the veracity of Precambrian

'sponge'fossils using in situ nanoscale analytical techniques. *Precambrian Research* **263**,

142–156 (2015).

46. Dunn, C. W. *et al.* Broad phylogenomic sampling improves resolution of the animal tree

of life. *Nature* **452**, 745–749 (2008).

47. Feuda, R. *et al.* Improved modeling of compositional heterogeneity supports sponges as

sister to all other animals. *Current Biology* **27**, 3864–3870 (2017).

48. Kapli, P. & Telford, M. J. Topology-dependent asymmetry in systematic errors affects

phylogenetic placement of Ctenophora and Xenacoelomorpha. *Science Advances* **6**,

eabc5162 (2020).

49. Li, Y., Shen, X.-X., Evans, B., Dunn, C. W. & Rokas, A. Rooting the Animal Tree of Life.

Molecular Biology and Evolution **38**, 4322–4333 (2021).

50. Philippe, H. *et al.* Phylogenomics Revives Traditional Views on Deep Animal

Relationships. *Current Biology* **19**, 706–712 (2009).

51. Pisani, D. *et al.* Genomic data do not support comb jellies as the sister group to all other

animals. *Proceedings of the National Academy of Sciences* **112**, 15402–15407 (2015).

52. Whelan, N. V. *et al.* Ctenophore relationships and their placement as the sister group to

all other animals. *Nat Ecol Evol* **1**, 1737–1746 (2017).

53. Whelan, N. V., Kocot, K. M. & Halanych, K. M. Employing Phylogenomics to Resolve the

Relationships among Cnidarians, Ctenophores, Sponges, Placozoans, and Bilaterians.

Integrative and Comparative Biology **55**, 1084–1095 (2015).

54. Antcliffe, J. B. Questioning the evidence of organic compounds called sponge

biomarkers. *Palaeontology* n/a-n/a (2013) doi:10.1111/pala.12030.

55. Nettersheim, B. J. *et al.* Putative sponge biomarkers in unicellular Rhizaria question an

early rise of animals. *Nat Ecol Evol* **3**, 577–581 (2019).

56. van Maldegem, L. M. *et al.* Geological alteration of Precambrian steroids mimics early

animal signatures. *Nature Ecology & Evolution* **5**, 169–173 (2021).

57. Love, G. D. *et al.* Sources of C30 steroid biomarkers in Neoproterozoic–Cambrian rocks

and oils. *Nature ecology & evolution* **4**, 34–36 (2020).

58. Andrews, S. FastQC: a quality control tool for high throughput sequence data. (2010).

59. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

60. Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a

reference genome. *Nature biotechnology* **29**, 644–652 (2011).

61. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-

generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

62. Emms, D. M. & Kelly, S. OrthoFinder2: fast and accurate phylogenomic orthology

analysis from gene sequences. *BioRxiv* **466201**, (2018).

63. Kocot, K. M., Citarella, M. R., Moroz, L. L. & Halanych, K. M. PhyloTreePruner: A

Phylogenetic Tree-Based Approach for selection of Orthologous sequences for

phylogenomics. *Evol Bioinform Online* **9**, EBO.S12813 (2013).

64. Whelan, S., Irisarri, I. & Burki, F. PREQUAL: detecting non-homologous characters in sets

of unaligned homologous sequences. *Bioinformatics* (2018)

doi:10.1093/bioinformatics/bty448.

65. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Research* **32**, 1792–1797 (2004).

66. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973

(2009).

67. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and

Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies.

Molecular Biology and Evolution **32**, 268–274 (2015).

68. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. *Molecular Biology and Evolution* **37**, 1530–1534 (2020).

69. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods*

14, 587–589 (2017).

70. Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2:

improving the ultrafast bootstrap approximation. *Molecular biology and evolution* **35**,

518–522 (2018).

71. Minh, B. Q., Nguyen, M. A. T. & Von Haeseler, A. Ultrafast approximation for

phylogenetic bootstrap. *Molecular biology and evolution* **30**, 1188–1195 (2013).

72. Lartillot, N., Rodrigue, N., Stubbs, D. & Richer, J. PhyloBayes MPI: Phylogenetic

Reconstruction with Infinite Mixtures of Profiles in a Parallel Environment. *Systematic*

Biology **62**, 611–615 (2013).

73. Lartillot, N. & Philippe, H. A Bayesian Mixture Model for Across-Site Heterogeneities in

the Amino-Acid Replacement Process. *Molecular Biology and Evolution* **21**, 1095–1109

(2004).

74. Mirarab, S. *et al.* ASTRAL: genome-scale coalescent-based species tree estimation.

Bioinformatics **30**, i541–i548 (2014).

75. Yang, Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Molecular Biology and*

Evolution **24**, 1586–1591 (2007).

76. Revell, L. J. phytools: an R package for phylogenetic comparative biology (and other

things). *Methods in Ecology and Evolution* **3**, 217–223 (2012).

77. Tarasov, S. Integration of Anatomy Ontologies and Evo-Devo Using Structured Markov

Models Suggests a New Framework for Modeling Discrete Phenotypic Traits. *Systematic*

Biology **68**, 698–716 (2019).

78. Keating, J. N. What is the best method for estimating ancestral states from discrete

characters? 2023.08.31.555762 Preprint at https://doi.org/10.1101/2023.08.31.555762

(2023).

79. Álvarez-Carretero, S. *et al.* A species-level timeline of mammal evolution integrating

phylogenomic data. *Nature* **602**, 263–267 (2022).

80. Rabosky, D. L. *et al.* BAMMtools: an R package for the analysis of evolutionary dynamics

on phylogenetic trees. *Methods in Ecology and Evolution* **5**, 701–707 (2014).

      MAIN FIGURES

Figure 1 Porifera time-tree with posterior estimates using the Spicule-bearing Demospongiae (SDN)

calibration strategy (red circle) and the ILN model in MCMCTree. Blue bars on nodes represent the

95% HPD. Red circles represent the two mutually exclusive calibrations strategies Spiculated Porifera

Node (SPN) and SDN. Full red dots represent calibrations in both strategies. Species in bold were

sequences in the framework of this study. Time scale on x-axis in hundreds of millions of years.

Abbreviation on the scale bar stands for Ne, Neoproterozoic; C, Cambrian; O, Ordovician; S, Silurian;

D, Devonian; C, Carboniferous; P, Permian; Tr, Triassic; J, Jurassic; K, Cretaceous; Pe, Paleogene.

Figure 2 Ancestral state estimations of spicule evolution obtained using the time-tree with the SDN

calibration strategy and the SMM combining the presence/absence of spicules and their mineralogy.

Cream dots indicate absence of spicules, blue stands for siliceous spicules, red for calcareous spicules,

and light orange stands for aragonite.

Figure 3 Rate diversification tests. Grey dots with numbers correspond to the shift highlighted by both

Medusa and BAMM analyses, the same number is reported when the same shift corresponds on both

topologies. A) BAMM results for the COI timetree inferred with the AR model. The two shifts

highlighted are on the branches of Bubarida + Axinellida (3), and Poecilosclerida (2). B) Medusa results

for the subsampled topology obtained with the AR molecular clock model. C) BAMM results for the

COI timetree inferred with the IR molecular clock model. The two shifts highlighted are on the

branches of Bubarida + Axinellida (2) and Hexacintellida (black full dot). D) Medusa results for the

subsampled topology obtained with the IR molecular clock model which highlights diversification rates

in Axinellida+ Bubarida(2), Poecilosclerida (3), and Dyctioceratida (4).

TABLES

Table 1 Summary of mean time estimates for all calibration strategies obtained with MCMCTree for

the classes of Porifera, the Silicea clade, and Heteroscleromorpha. Full details can be found in Table

S3-S4.

Nodes	SDN AR time estimates	SPN AR time estimates	SDN ILN time estimates	SDN ILN time estimates

Porifera	603.79-582.68	594.59-578.79	615.32-601.3	610.68-590.34
Silicea	577.48-561.72	567.69-551.55	591.32-570.3	575.87-542.68
Demospongiae	549.57-538.18	535.02-503.38	559.17-536.53	519.67-462.67
Hexactinellida	454.92-445.07	443.89-422.5	457.3-445.09	454.99-445.07
Calcarea	473.24-163.27	460.38-157.52	351.7-213.89	337.36-207.53
Homoscleromorpha	540.28-435.36	535.63-420.18	429.15-228.57	421.58-224.87
Heteroscleromorpha	521.97-515.05	500.69-455.96	520.81-515.04	454.95-396.53

737