Adiponectin pathway activation dampens inflammation and enhances alveolar macrophage

fungal killing via LC3-associated phagocytosis

Sri Harshini Goli,[1,2]* Joo-Yeon Lim,[1]* Nese Basaran-Akgul,[1] and Steven P. Templeton[1,†]

[1] Department of Microbiology and Immunology, Indiana University School of Medicine-Terre Haute,

Terre Haute, IN 47809, USA

[2] Department of Biology, Indiana State University, Terre Haute, IN 47809, USA

* These authors contributed equally to this work.

Short title: Adiponectin promotes LC3-mediated fungal killing by alveolar macrophages

†Correspondence: sptemple@iu.edu; Tel.:+1-821-237-8193

Abstract

Although innate immunity is critical for antifungal host defense against the human opportunistic fungal

pathogen *Aspergillus fumigatus*, potentially damaging inflammation must be controlled. Adiponectin

(APN) is an adipokine produced mainly in adipose tissue that exerts anti-inflammatory effects in

adipose-distal tissues such as the lung. We observed 100% mortality and increased fungal burden and

inflammation in neutropenic mice with invasive aspergillosis (IA) that lack APN or the APN receptors

AdipoR1 or AdipoR2. Alveolar macrophages (AMs), early immune sentinels that detect and respond to

lung infection, express both receptors, and APN-/- AMs exhibited an inflammatory/M1 phenotype that

was associated with decreased fungal killing. Pharmacological stimulation of AMs with AdipoR

agonist AdipoRon partially rescued deficient killing in APN-/- AMs that was dependent on both

receptors. Finally, APN-enhanced fungal killing was associated with increased activation of the non-

canonical LC3 pathway of autophagy. Thus, our study identifies a novel role for APN in LC3-mediated

killing of *A.fumigatus*.

Author Summary

Aspergillus fumigatus is a human fungal pathogen that causes an often-fatal invasive infection of the

lung in immune compromised individuals, a population that is increasing. Since current antifungal

drugs have limited efficacy, it is important to identify pathways that may be pharmaceutically targeted

to complement existing therapies. Adiponectin (APN) is an anti-inflammatory intercellular cytokine

messenger produced mainly in adipose tissue that protects against invasive aspergillosis. Alveolar

macrophages are early immune sentinel cells in the lung, and we report that AMs from mice lacking

APN exhibit an inflammatory phenotype and reduced killing of *A. fumigatus* spores that is improved

when AMs are treated with a drug (AdipoRon) that simulates APN binding to its receptors AdipoR1

and AdipoR2. Furthermore, APN was associated with activation of LC3-associated phagocytosis, a

mechanism of fungal killing important for host defense against *A. fumigatus* infection. Thus, we

identify therapeutic potential of the APN pathway in stimulation of immune-mediated fungal killing

and treatment of fungal infection.

Keywords: invasive aspergillosis, *Aspergillus fumigatus*, adiponectin, adiponectin receptor, adipoRon,

lung immune responses, alveolar macrophages, LC3-associated phagocytosis

Introduction

The human opportunistic fungal pathogen *Aspergillus fumigatus* is the primary etiologic agent of invasive pulmonary aspergillosis (IA), a frequently severe infection with a high mortality rate [1]. Effective treatment options for IA remain limited, despite increases in the at-risk population, and fungal resistance to existing antifungal drugs is increasing. Furthermore, although an underlying immune deficiency renders individuals susceptible to IA, poor disease outcomes are associated with detrimental inflammatory pathology [2,3]. It is thus critical to identify anti-inflammatory pathways that could be exploited to target detrimental immune pathology, enhance antifungal immunity, and complement existing antifungal therapies in IA patients.

Adiponectin (APN) is an anti-inflammatory adipokine produced mainly in adipose tissue and expressed at high levels in lean and healthy individuals, whereas in obese individuals, adipose tissue is inflamed and APN production and secretion are inhibited [4–6]. Numerous studies have demonstrated a protective effect for APN in autoimmune and inflammatory diseases, and immune-mediated clearance of *Listeria monocytogenes* was hampered in obese and APN-deficient mice due to detrimental inflammation and hematopoietic dysfunction [7]. More recently, we demonstrated increased mortality, fungal burden, lung inflammatory cytokine production and eosinophil recruitment in APN-deficient mice with IA [8]. However, eosinophils did not play a significant role in the pathology of APN-deficiency, as neutropenic mice deficient in both APN and eosinophils exhibited only delayed, but not decreased mortality, suggesting other unidentified mechanisms of APN-mediated protection in IA.

Alveolar macrophages (AMs) are the primary immune sentinel of lung airways, and their early responses to inhaled fungal conidia are critical for disease outcomes, particularly in immunocompromised hosts [9,10]. In our previous study, we observed increased production of the pro-inflammatory cytokine TNF in APN-/- AMs [8], suggesting that these cells contribute to the inflammatory phenotype of APN-/- mice with IA. In this study, we aimed to determine how the adiponectin pathway regulates responses of AMs to *A. fumigatus* infection. We report increased production of inflammatory cytokines, an elevated M1 phenotype, and reduced uptake and killing of conidia in APN-/- AMs that is increased by stimulation with the APN receptor agonist AdipoRon. Furthermore, APN stimulation was associated with increased AM activation of the non-canonical autophagy LC3 pathway in conidia-containing phagosomes, thus identifying a novel role for APN in

promoting LC3-associated phagocytosis (LAP) in antifungal immunity.

Results

Increased mortality, fungal burden, and inflammatory pathology in APN pathway-deficient mice
with IA.

Previously, we reported that APN inhibits inflammatory lung pathology during IA [8]. Since APN-/-
strains vary in their metabolic phenotypes[11–13], we confirmed the inflammatory phenotype of APN-
/- mice from a second strain (obtained from Dr. Philipp Scherer) (**Fig. S1**), and observed similar
increases in fungal burden (**Figs. S1B-S1D**), inflammatory cytokines and inflammatory cell recruitment
(**Figs. S1E-S1F** and **S1H**) while only survival and AdipoR1 transcription were not significantly
different from wild-type (WT) C57BL/6 controls (**Figs. S1A** and **S1G**, respectively). We also aimed to
determine the contribution of the canonical adiponectin receptors AdipoR1 and AdipoR2 to IA
pathology, and thus infected neutrophil-depleted WT, APN-/-, AdipoR1-/- and AdipoR2-/- mice to
compare mortality, pathology, fungal burden, inflammatory cell recruitment and cytokine production
(**Fig. 1A**). In contrast to WT mice, mortality was 100% in APN-/-, AdipoR1-/-, and AdipoR2-/- mice,
with mortality beginning on day 2 post-infection and 100% by day 6 p.i. (**Fig. 1B**), with extensive lung
hyphal growth and peri-bronchoalveolar inflammation (**Fig. 1C**) and increased fungal burden as
measured by qPCR of fungal DNA (**Fig. 1D**) and image quantification of fungal GMS staining (**Fig.**
1E). Inflammatory cells were increased in the bronchoalveolar lavage fluid (BALF), most significantly
in APN-/- mice (**Fig. 1F**), while lung inflammatory cytokines IL-1a, IL-6, TNF, IL-12b, and IL-17A
and the anti-inflammatory cytokine IL-10 were increased, most significantly in AdipoR1-/- mice (**Figs.**
1G and 1H). In summary, adiponectin-pathway deficiency is associated with increased mortality, lung
fungal burden, and inflammatory pathology, cell recruitment, and cytokine production. Furthermore,
disrupted signaling through either AdipoR1 or AdipoR2 results in increased IA severity.

Alveolar macrophages (AMs) from adiponectin pathway-deficient mice exhibit an inflammatory
phenotype.

Previously, we observed increased intracellular TNF in AMs from APN-/- mice with IA when
compared to WT AMs [8], and we observed mortality as early as day 2 post-infection in APN-
pathway-deficient mice with IA, suggesting a critical involvement of early immune effectors in
protection from infection. As AMs contribute to early responses to inhalation of *A. fumigatus* [9],

especially in immunocompromised hosts [10], we aimed to further examine the phenotype of AMs

from APN pathway-deficient hosts using primary ex vivo cultures. We observed expression of both

AdipoR1 and AdipoR2 in WT and APN-deficient mice that was decreased in lung tissue upon *A.*

fumigatus infection, but increased in the BALF (**Fig. S2A**), with the highest surface expression of

AdipoR1 in AMs (**Fig. S2B**). APN-pathway deficient AMs exhibited differential expression of

AdipoRs at the transcription level **Fig. S2C**), and infection of APN-/- AMs decreased AdipoR1

expression (**Fig. S2D**). Although WT, APN-/-, AdipoR1-/-, and AdipoR2-/- AMs were similar

morphologically (data not shown), transcription of IL-1α, IL-1β, IL-6, IL-17A, IL-23, and TNF was

increased in APN pathway-deficient AMs when compared WT-derived AMs, while IL-10 was similar

(**Fig. 2A**), and secreted TNF in the supernatant increased equally in response to infection in all groups

(**Fig. 2B**). To investigate the inflammatory phenotype more specifically in ex vivo cultured APN-/-

AMs, we compared expression of the macrophage polarization markers CD38 (M1/classical) and Egr2

(M2/alternative) with wild type WT AMs [14]. In a resting state, a higher percentage of APN-/- AMs

were CD38+ and lower percentage were Egr2+ when compared to WT (**Figs. 2C and 2D**), while upon

infection with *A. fumigatus*, percentages of both CD38+Egr2- and CD38-Egr2+ cells increased (**Figs.**

2C and 2E). These results suggest that APN pathway-deficient AMs exhibit an increased inflammatory

phenotype compared to WT AMs.

***A. fumigatus* uptake and killing is inhibited in APN-/- AMs.**

Although APN pathway-deficient mice displayed increased IA severity, fungal burden, and an

increased AM inflammatory phenotype, the mechanism of decreased fungal clearance remains

unknown. To determine if APN-deficient AMs were deficient in fungal uptake and killing, we used

dual viability/tracking fluorescent *Aspergillus* reporter FLARE conidia [15]. to compare changes in

conidia uptake and viability in wild-type and APN-/- AMs ex vivo and in vivo. In APN-/- AMs

cultured ex vivo, uptake and killing of dual-labeled FLARE conidia was decreased after 6 hours when

observed by microscopy (**Fig. 3A** and **Supplemental Video SV1**). When loss of *A. fumigatus* viability

was compared by image analysis, killing of FLARE conidia by APN-/- AMs was markedly decreased

compared to wild-type AMs (**Fig. 3B**). Uptake and killing were also reduced in APN-/- AMs when

analyzed by flow cytometry, even after filtration to remove hyphae (**Figs 3C and 3D**). Reduced killing

by APN-/- AMs was confirmed in vivo in FLARE-infected neutropenic mice, with a significant

difference in conidial viability by day 3 post-infection (**Figs. 3E and 3F**), while uptake of 1 μm inert

particles was not significantly different between wild-type and APN-/- AMs (**Fig. 3G**). Collectively,

these results demonstrate that AM fungal uptake and killing are decreased in the absence of APN.

The AdipoR agonist AdipoRon decreases inflammation and fungal burden in APN-deficient mice

with IA.

A small molecule agonist (AdipoRon) that binds and activates signaling through both AdipoR1 and

AdipoR2 improved insulin resistance and glucose intolerance and increased the lifespan of obese mice

[16]. A role for AdipoRon in dampening inflammatory responses in human lung macrophages via

inhibition of production of TNF, IL-6, and inflammatory chemokine production has also been reported

[17]. However, the ability of AdipoRon therapy to improve IA outcomes remained unknown. We

treated infected neutropenic wild-type or APN-/- mice with AdipoRon by involuntary aspiration daily

for up to eight days (**Fig. 4A**). Although AdipoRon treatment did not significantly improve survival

(**Fig. 4B**), lung hyphal growth and fungal burden were reduced in APN-/- mice (**Figs. 4C-4E**). Airway

eosinophil recruitment was reduced in AdipoRon treated APN-/- mice, while AMs were increased in

both AdipoRon-treated WT and APN-/- mice (**Fig. 4F**). Transcription of IL-1a, IL-6, IL-10, IL-17A,

and TNF were also decreased with AdipoRon treatment in APN-/- lungs and decreases in IL-1a and IL-

10 were also evident in wild-type mice (**Fig. 4G**). In contrast, IL-12b transcription appeared to be

increased by AdipoRon, but only in wild-type lungs. An AdipoRon-mediated decrease in TNF was also

confirmed in APN-/- BALF at the protein level (**Fig. 4H**). These results indicate that APN pathway

stimulation by AdipoRon decreases lung fungal burden and dampens lung inflammation in APN-/-

mice with IA.

AdipoRon increases fungal killing and inhibits the inflammatory phenotype of APN-/- AMs.

Although AdipoRon treatment influenced lung fungal growth and dampened inflammation in APN-/-

mice with IA, the specific effect of AMs remained unknown. To determine the effect of AdipoRon on

AM phenotype and fungal uptake and killing, we infected WT or APN pathway-deficient AMs with

Af293 or FLARE conidia after 24 hours treatment with AdipoRon or control vehicle (containing

DMSO), and harvested AMs ten hours later for microscopic, flow cytometric, qRT-PCR, and ELISA

analyses (**Fig. 5A**). When quantified by microscopy, conidia viability was decreased by AdipoRon

treatment of APN-/- but not wild-type AMs (**Fig. 5B**). Conidia viability was also reduced in APN

pathway-deficient AMs by AdipoRon when measured by flow cytometry after cell straining to remove

hyphae (**Fig. 5C**). Inflammatory cytokine transcription of IL-1α, IL-1β, and IL-6 were all decreased by

AdipoRon treatment in APN-/- AMs, while the anti-inflammatory cytokine IL-10 was not affected by

AdipoRon regardless of APN sufficiency (**Fig. 5D**). Secretion of TNF protein was also significantly

decreased in AdipoRon-treated APN-/- AMs (**Fig. 5E**). These results indicate that AdipoRon treatment

acts on APN-/- AMs to improve fungal killing and inhibit an inflammatory phenotype.

AdipoRon-enhanced killing of *A. fumigatus* requires both AdipoR1 and AdipoR2.

Since AdipoRon enhances AM-mediated fungal killing, we aimed to confirm the requirement of

AdipoR1 and AdipoR2 by using siRNA to target expression of AdipoR1 in wild-type, APN-/-, and

AdipoR2-/- AMs (**Fig. 6A**). Decreased AM transcription of *Adipor1* and *Adipor2* by infection was

partially restored by AdipoRon treatment (**Fig. S2E**). Knockdown of AdipoR1 was confirmed by flow

cytometry to reduce expression to identical levels observed in AMs from AdipoR1-/- mice (**Figs. 6B**

and 6C). Moreover, fungal killing was improved by AdipoRon treatment in wild-type, APN-/-,

AdipoR1-/- (KO mice and siRNA), and AdipoR2-/- AMs, but was not improved with siRNA-targeted

depletion of AdipoR1 in AdipoR2-/- AMs (**Fig. 6D**). These results show that both canonical APN

receptors are critical for AdipoRon-mediated enhancement of AM fungal killing.

Adiponectin promotes LC3-associated phagocytosis of *A. fumigatus* conidia.

To confirm the AdipoRon-mediated phenotypic shift in AMs, we compared AdipoRon treatment with

vehicle in infected and uninfected APN-/- AMs by global RNAseq analysis. In uninfected AMs, more

genes were upregulated in response to AdipoRon treatment, while the opposite was observed in AMs

after 12 hours post-infection, with AdipoRon treatment associated with gene silencing (**Fig S3A**).

Inflammatory/M1-associated gene expression was suppressed by AdipoRon treatment in infected AMs,

including *Il1a*, *Il1b*, *Ccl2*, *Cd38*, *Tlr4*, and *Hif1a*, while the anti-inflammatory *Tgfb* was increased, as

were autophagy genes, including *Map1lc3a* and *Map1lc3b* genes that are critical for LAP (**Fig. S3B**).

In contrast, inflammatory gene expression appeared to be increased by AdipoRon in uninfected AMs.

GSEA pathway analysis confirmed inhibition of cytokine signaling by AdipoRon in infected AMs and

also identified activation of gene pathways associated with oxidative phosphorylation (**Fig. S3C**), a

metabolic pathway favored by M2 macrophages [18]. Our RNAseq results thus confirm an AdipoRon-

mediated inhibition of the inflammatory/M1 phenotype of APN-/- AMs in response to fungal infection,

and identify a potential role for autophagy induction.

Since LAP is an important mechanism of killing of *A. fumigatus* by macrophages [19], we aimed to

determine if phagosomal LC3 activation is increased in APN-/- AMs by AdipoRon treatment. Using

immunofluorescence microscopy staining of LC3, we observed clear colocalization of LC3 staining

and DS-RED+(viable) conidia in AdipoRon-treated APN-/- AMs (**Fig. 7A**), with visibly less

colocalization in untreated AMs that was confirmed by image quantification (**Fig. 7B**). The median

fluorescence intensity (MFI) of LC3+ macrophage staining was also increased by AdipoRon in both

wild-type and APN-/- AMs as measured by flow cytometry (**Fig. 7C**). Collectively, these results

demonstrate a role for adiponectin in LAP in AMs in response to *A. fumigatus* conidia.

Discussion

To our knowledge, our study is the first to identify a role for adiponectin in LC3-associated

phagocytosis (LAP) in response to infection. Previous studies reported roles for adiponectin in LC3-

mediated autophagy in skeletal muscle cells [20,21], cardiomyocytes [22], or macrophages in a mouse

model of atherosclerosis [23]. In addition, AM autophagy is critical for prevention of spontaneous lung

inflammation driven by airway microflora [24]. Although others have identified a role for the LC3

pathway in efferocytosis by macrophages [25], and although adiponectin promotes macrophage

efferocytosis [26,27], a direct connection between adiponectin, LC3/autophagy and macrophage

efferocytosis is lacking. Since macrophage autophagy and removal of apoptotic cells are important

mechanisms to limit lung inflammatory pathology [24,27,28], it will be of interest in future studies to

determine their role in adiponectin-mediated dampening of inflammation in IA.

A number of studies have defined a critical role for LAP in protection from *A. fumigatus* infection

[19,29–31]. LAP dampens inflammation in response to *A. fumigatus* by inhibition of the NLRP3

inflammasome and IL-1b via activation of DAPK1 [29]. In our study, we observed that AdipoRon

decreased transcription of IL-1α and IL-1β genes in APN-/- AMs. Interestingly, blockade of IL-1

receptor signaling (by IL-1ra) restored autophagy/LC3 recruitment in NADPH oxidase-deficient

macrophages, a hallmark myeloid cell deficiency of Chronic Granulomatous Disease [32]. Since both

AdipoRon and IL-1ra promote LC3 activation, it will be important in future studies to determine if

AdipoRon could restore autophagy in NADPH oxidase-deficient macrophages, and to identify shared

regulatory pathways.

A previous study reported that LAP-associated fungal clearance was inhibited by fungal melanin, a

virulence factor of *A. fumigatus* [30]. How APN stimulates LAP in response to *A. fumigatus* and the

ability to overcome LAP interference by fungal melanin remain unknown. Furthermore, we observed

that chitin inhalation resulted in decreased production of lung adiponectin and AdipoR1 expression in

murine airway cells [33]. It will be of interest to determine the effect of AdipoRon on recognition of

multiple fungal pathogen-associated molecular patterns by WT and APN pathway-deficient AMs,

specifically in the context of LC3 activation.

Adiponectin has two canonical signaling receptors with broad, yet different cell/tissue distribution [34].

We observed that both AdipoR1 and AdipoR2 contribute to adiponectin-mediated protection from IA

and dampening of inflammatory pathology. However, some differences were evident, as AdipoR1-/-

mice with IA exhibited the highest levels of lung inflammatory cytokine transcription that was not as

clear in AdipoR1-/- AMs, suggesting that immune pathology in AdipoR1-/- mice was not driven solely

by the lack of AdipoR1 signaling in AMs, but also by other immune and resident lung cells, with a

contribution of potentially enhanced signaling through AdipoR2. AMs express both AdipoRs, and their

expression was modulated by *A fumigatus* infection, most significantly for AdipoR1, which was

increased in response to infection in WT AMs but decreased in APN-/- AMs. Furthermore, both

AdipoR1 and AdipoR2 were required for AdipoRon-mediated improvement in *A. fumigatus* killing, as

AdipoRon was not effective in AdipoR1/R2-/- AMs, confirming a role for both receptors in APN-

mediated enhancement of AM function. However, the relative contribution of AdipoR-expressing

AMs in adiponectin-mediated protection from infection remains unclear, and thus remains a subject of

future investigation.

AMs are dynamic cells, and their activation and polarization falls on a spectrum from M1 to M2, as

these are designations and not indicative of a static phenotype [35,36]. In our study, AM treatment with

AdipoRon decreased expression of M1 genes and activated M2 genes in *A. fumigatus*-infected APN-/-

AMs, although these phenotypes are not identical to previously published studies using classical

activation stimuli, such as LPS/IFNg for M1 and IL-4 for M2 [37,38]. It is thus more appropriate to

refer to APN-/- AMs as 'M1-like' and AdipoRon-treated AMs as 'M2-like' cells [39,40]. Perhaps more

importantly, AdipoRon treatment of APN-/- AMs resulted in activation of genes associated with

oxidative phosphorylation that is associated with M2-like macrophages, in contrast to glycolysis-

favoring M1 macrophages [18]. In support of increased oxidative metabolism for AdipoRon-treated

APN-/- AMs, arginase-2 (*Arg2*) expression was also increased, and mitochondrial *Arg2* was recently

identified as critical for induction of oxidative phosphorylation in inflammatory macrophages [41].

Collectively, the results of our study confirm a role for APN in skewing AMs toward an M2-like

phenotype.

Obesity is widely accepted as an inflammatory disease characterized by adipose tissue inflammation

and decreased APN and APN receptor expression [6,42–45], yet the effect of obesity-related APN

pathway-inhibition in response to *A. fumigatus* infection is not known. In response to infection with

Listeria monocytogenes, both obese and APN-/- mice exhibited attenuated clearance that was

characterized by an inflammatory phenotype in bone marrow macrophages that was partially restored

by AdipoRon [7]. Furthermore, obesity is associated with an increased inflammatory phenotype in

alveolar macrophages [46]. It is therefore possible that stimulation of the APN pathway with AdipoRon

will improve AM killing of *A. fumigatus* in obese mice, an outcome that would support the APN

pathway as a potential therapeutic route in IA patients with low APN levels.

Acknowledgements

We would like to thank Azriel Manalaysay, Alsace Key, and Gwen Schmiedel for assistance with

animal care and genetic screening, Brielle Batch for technical assistance, Takuya Akiyama for

assistance with confocal microscopy, Americo Lopez-Yglesias for assistance with flow cytometry, and

Tobias Hohl for review of this manuscript. The study was supported by National Institutes of Health

NIH R21AI163574 (ST) and T32 AI060519 (NBA; IUSM Department of Microbiology and

Immunology, American Lung Association Innovation Award #IA-690880 (ST), and the Indiana

Clinical and Translational Sciences Institute, funded in part by grant # UM1TR004402 from the

National Institutes of Health, National Center for Advancing Translational Sciences, Clinical and

Translational Science Award (ST).

Author contributions

Conceptualization, S.P.T., H.G., J.L.; Methodology, J.L. and H.G.; Investigation. H.G., J.L., and

N.B.A.; Resource; J.L., N.B.A., and H.G.; Writing - S.PT, J.L and H.G.; Visualization, H.G, J.L.;

Funding Acquisition. S.P.T.

Declaration of Interests

The authors declare no competing interests.

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
CD45-PerCP	BD Biosciences	557235
SiglecF-Superbright600	Thermo Fisher Scientific	63-1702-82
SiglecF-PE	BD Biosciences	552126
Cd11c-SparekBlue550	BioLegend	117366
Cd11c-PECy7	Thermo Fisher Scientific	25-0114-81
Ly6G-BV785	BioLegend	127645
Ly6G-FITC	BD Biosciences	551460
Fc Block	BD Biosciences	553142

Rabbit IgG - Isotype control	Abcam	ab171870
Anti-AdipoR1 antibody	Abcam	ab70362
Goat anti-Rabbit IgG H&L (Alexa Fluor 405)	Abcam	ab175652
Anti-mouse EGR2	Invitrogen	247632
Anti-mouse CD38	Invitrogen	2504972
Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 488	Thermo Fisher Scientific	A-11008
Adiponectin receptor 1 Recombinant Rabbit Monoclonal Antibody	Invitrogen	MA5-32249
Chemicals		
Streptavidin, Alexa Fluor 633 conjugate	Invitrogen	S21375
Biotin-XX, SSE	Invitrogen	B6352, B1606
ACK lysing buffer	Thermo Fisher Scientific	A1049201
Hygromycin B, *Streptomyces* sp.	Calbiochem	31282-04-9
Xylene	Millipore Sigma	XX0060
TRIzol reagent	Thermo Fisher Scientific	15596026
AdipoRon	Cayman chemical	924416-43-3
Albumin, Bovine Serum	Millipore Sigma	126579
Normal Donkey Serum	abcam	ab7475
Sodium Bicarbonate 7.5% solution	Thermo Fisher Scientific	25080094
Carboxymethylcellulose sodium salt	Thermo Fisher Scientific	A18105.36
0.5 M EDTA	Thermo Fisher Scientific	15575020
Dimethyl Sulfoxide	Millipore Sigma	D4540
10% Neutral Buffered Formalin	Thermo Fisher Scientific	5701
RPMI 1640, 1X with L-glutamine	CORNING	28121001
2-Mercaptoehanol	A ALDRICH	SHBF0971V
50ug Murine GM-CSF	PeproTech	032255 K1622
Lipofectamine 2000 Transfection Reagent	Thermo Fisher Scientific	11668019
Vectashield mounting medium for fluorescence with DAPI	VECTOR laboratories	H-1200-10

FluoSpheres Polystyrene Microspheres, 1.0 µm, red fluorescent (580/605), for tracer studies	Thermo Fisher Scientific	F13083
Critical commercial assays		
Silver stain (Modified GMS)	Millipore Sigma	HT100A
RNeasy Mini kit	Qiagen	74106
RNase-Free DNase set	Qiagen	79256
PowerUp SYBR Green Master Mix	Thermo Fisher Scientific	A25742
ABsolute Blue qPCR low ROX mix	Thermo Fisher Scientific	AB-4318
High Capacity cDNA Reverse Transcription Kit	Thermo Fisher Scientific	4368814
Mouse Adiponectin/Acrp30 ELISA kit	R&D Systems	DY1119
Murine TNF-α Standard TMB ELISA kit	Peprotech	900-T54
LC3B Antibody Kit for Autophagy	Invitrogen	L10382
TriFECTa RNAi Kit	Integrated DNA Technologies	N/A
Experimental models: Organism/strains		
A. fumigatus strain AF293	Fungal Genetic Stock Center	Cat#A1100
A. fumigatus strain AF293 ds-Red (FLARE strain)	Jhingran et al. [24w]	N/A
Oligonucleotides		
18S rRNA-encoding DNA primers and probe	50-/56-FAM/AGC CAG CGG/ZEN/CCC GCAAAT G/3IABkFQ/-30	
Software and algorithms		
FlowJo	FlowJo 10	N/A
Prism 9	Prism 9	N/A
Gen5	BioTek	N/A
Image J	Image J	N/A

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Steven P. Templeton. (sptemple@iu.edu)

Material Availability

All materials within the paper are available from the corresponding author upon reasonable request.

This study did not generate new reagents.

Data and Code Availity

The RNAseq data are available at Gene Expression Omnibus (Accession #GSE268782).

Materials and Methods

Mice

C57BL/6 and adiponectin-deficient (*Adipoq−/−*) and *AdipoR2+/−* mice were obtained from The Jackson Laboratory, while AdipoR1+/- mice were obtained from the Mutant Mouse Resource and Research Center (MMRRC). To obtain homozygous deletion mice for *AdipoR1−/−*, and *AdipoR2−/−* mice, heterozygous mice were breed with F1 animals heterozygous and offspring were screened by PCR, with loss of protein confirmed in a subset of animals by flow cytometry. All animal handling and experimental procedures were performed in accordance with the recommendations found in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. The work in this study was approved by the Institutional Animal Care and Use Committee of the host campus of Indiana University School of Medicine-Terre Haute, Indiana State University.

Fungal strains and cultivation

The clinical isolates AF293 was previously obtained from the Fungal Genetics Stock Center and grown on Malt Extract Agar plates at 22˚C. To generate FLARE conidia, AF293-dsRed conidia (provided by Dr. Tobias Hohl, Memorial Sloan-Kettering) were rotated in 0.5 mg/ml Biotin XX, SSE in 1 ml of 50 mM $NaHCO_3$ for 2 h at 4˚C, washed with 0.1 M Tris-HCl (pH 8.0), incubated with 0.02 mg/ml Af633-streptavidin for 30 min at RT, and resuspended in PBS for use [47].

Fungal aspiration and infection

Neutrophils were depleted by i.p. injection of 0.5 mg of α-Ly6G (1A8 clone) 24 h pre- and post-infection for invasive pulmonary aspergillosis mice model, as described [8]. Isoflurane-anesthetized mice involuntarily aspirated 50 μl of suspension including $1 - 2 \times 10^7$ of conidia. For survival test, infected mice were monitored for 8 days post infection. For further analyses, a subset of infected mice were euthanized with sodium pentobarbital and mouse BAL cells and lungs were harvested 24 h or 72 h later for further analyses.

Histology

At 3 days post infection, lungs were harvested after perfusion with 5 ml of PBS followed with 5 ml of 10 % phosphate buffered formalin and inflated with formalin. Lung tissue was embedded in parafilm after dehydration and sectioned in 4 μm slices using xylene and ethanol and stained with modified Gomori modified methanamine silver (GMS) and hematoxylin and eosin (H&E) staining [8]. Quantification of GMS staining to determine fungal burden was performed as previously described [48].

Quantitative PCR fungal burden assay

At 72 h post infection, lungs were harvested, frozen in liquid nitrogen and lyophilized. Genomic DNA was extracted from homogenized lung tissue with DNA extraction buffer for *Aspergillus* nucleic acids and subsequent phenol/chloroform extraction. Quantitative PCR (qPCR) fungal burden assay was performed using 0.2 μg of genomic DNA with 18S rRNA-encoding DNA primers and probe sets with a modified probe quencher (50-/56-FAM/AGC CAG CGG/ZEN/CCC GCAAAT G/3IABkFQ/-30) [49]. qPCR was performed using the CFX Connect Real-Time System. Threshold values were used to calculate the corresponding fungal DNA content in the lung tissues.

Tissue RNA extraction and gene expression analysis

Total RNA was extracted from homogenized lungs in TRIzol reagent. Following the upper phase, further RNA purification was performed using a Qiagen RNeasy column and DNase treatment per the manufacturer's recommendations [50]. Mouse β-actin was used as the housekeeping gene for cytokine analysis in the murine model. Most primers were designed using the Primer3 software (version 0.4.0) from the whole sequence available in GenBank.

BALF Flow cytometric analysis

BALF cell composition was determined by flow cytometric analysis with staining of surface markers.

In brief, BALF was centrifuged and the cell pellet was washed with 1 ml of FACS buffer (PBS, 5%

FBS, 50 mM EDTA). The washed pellet was lysed of red blood cells with ACK lysis buffer and stained

with 10% rat serum, Fc receptor blocking Ab, and antibodies: rat anti-mouse Ly6G-FITC or -BV785,

rat anti-mouse SiglecF-PE or -Superbright600, rat anti-mouse CD45-PerCP, and rate anti-mouse

CD11c-PE-Cy7 or -SparkBlue550 [50,51]. For staining of AdipoR1, the washed pellet was

resuspended in blocking solution containing 10% donkey serum, Fc receptor blocking Ab, and 1%

Bovine serum albumin in PBS for 30 min, incubated with the primary AdipoR1 antibody for 1 h and

incubated with the IgG secondary antibodies [33]. Analyses were performed by flow cytometric

analysis on a Guava EasyCyte HT system and Cytek. Flow cytometric data were analyzed using

FlowJo software.

BALF ELISA analysis

Serum and BALF supernatants were collected after centrifugation. ELISA analyses were performed for

TNF-α cytokine (PeproTech) and Adiponectin (R&D Systems) as described by the manufacturer's

protocol.

Alveolar macrophage extraction and culture

Seven to 8 weeks old mice were euthanized, and an 18-gauge catheter was inserted into the trachea and

alveolar macrophages were harvested from mouse lungs with sterile filtered PBS containing 2 mM

EDTA (diluted 1:250 from 0.5 M EDTA stock solution) and 0.5% Fetal bovine serum (FBS) in 5ml

BAL fluid. Cells were separated from the BALF by centrifugation at $400 \times g$ for 6 min at 4°C and the

cells were suspended in RPMI 1640 supplemented with penicillin (100 U/ml), streptomycin (100

U/ml), 5% heat-inactivated FBS, 20ng/ml GM-CSF and 143mM beta-mercaptoethanol. BALF cells

were plated in Corning Costar Flat Bottom 12 well Cell Culture Plates at a concentration of $5 \times$

10^5 cells per well. The cells were allowed to adhere for 24 hours at 37°C under a humidified

atmosphere with 5% CO2 and were washed with PBS thrice and cell culture media was added,

allowing only the adherent AMs to culture in the wells. The cell culture media was changed on day 3,

day 6 and day 9 of the cultures. AMs were fully confluent and ready for treatment and infection on day

10. The wells for divided into DMSO control, AdipoRon treatment with AF293/FLARE infection and

AF293/FLARE infection groups.

Quantitative RT-PCR for ex-vivo gene expression analysis

373   AMs were treated with 20 uM AdipoRon or vehicle-treated for 24 hours, followed by infection with

374   AF293 fixed and swollen conidia using RPMI 1640 medium at a concentration of 9 conidia per cell for

375   RT-PCR. After 10 hours of infection, cells were ready for RNA extraction and cDNA synthesis. RNA

376   extraction from AMs was done using Qiagen RNeasy Mini kit, following the manufacturer's protocol.

AM supernatant ELISA

378   After culturing for 10 days, AMs were grouped into uninfected, vehicle (DMSO), infected and

379   AdipoRon + infected groups. After the infection at a concentration of 9 conidia per cell with fixed and

380   swollen AF293 conidia for 10 hours, the supernatants were collected and analyzed for TNF-alpha

381   expression.

AM flow cytometric analysis

FLARE uptake and killing

Conidia was harvested from 7-10 days old Minimal Media (GMM) plates. AF293-dsRed conidia were

rotated in 0.5 mg/ml Biotin XX, SSE in 1 ml of 50 mM NaHCO3 for 2 h at 4˚C, washed with 0.1 M

Tris-HCl (pH 8.0), incubated with 0.02 mg/ml Af633-streptavidin for 30 min at RT, and resuspended in

PBS for use. For in-vitro conidial uptake and killing assays, $5x10^5$ FLARE conidia were added to $5x10^5$

alveolar macrophages (1:1). After 10 hours alveolar macrophages were harvested using cell scraper

with slight force from the wells in 500 ul ice cold FACS buffer per well and centrifuged at $400 \times g$ for

6 min at 4°C. The cell pellet was washed with 1 ml of FACS buffer (PBS, 5% FBS, 50 mM EDTA)

and the samples were passed through 70μm filter to clear hyphae and have clear sample. The cells were

analyzed by the flow cytometry for dsRed (Ex/Em: 558nm/583nm) and AF633 (Ex/Em:

631nm/650nm) fluorescence followed by gating using FlowJo.

M1/M2 phenotype study

395   For M1/M2 phenotype analysis cells were stained with 10% rat serum, Fc receptor blocking Ab and

396   antibody: *CD38 Monoclonal Antibody (90), APC-eFluor™ 780, eBioscience™* for 15 min at room

397   temperature. For staining with the antibody: *EGR2 Monoclonal Antibody (erongr2), PE-Cyanine7,*

398   *eBioscience™* cells were permeabilized using Cytoperm and processed for staining.

AM Microscopy

FLARE uptake and killing

401   Alveolar macrophages were plated in Cellvis 35 mm Glass bottom dish with 14 mm micro-well and
402   were infected with FLARE/DsRed at a 1:1 ratio. Cells were live imaged after 6 hours on a ZEISS
403   LSM780 confocal microscopy.

LC3 phagocytosis assay

On day 10, cells were treated with 20 uM AdipoRon for 24 hours followed by treatment with 90μm
chloroquine for 13 hours. Cells were then infected with DsRed+ conidia for 120 min and then fixed
with 3.7% formaldehyde in PBS for 15 min at room temperature. LC3B antibody kit for autophagy was
used and the manufacturer's protocol is followed for LC3 staining. Invitrogen™ Goat anti-Rabbit IgG
(H+L) Cross-Adsorbed Secondary, Antibody, Alexa Fluor™ 488 has been used for secondary staining.
Cells were washed and counter stained with DAPI using VECTASHIELD® Antifade Mounting
Medium has been done. Microscopy was performed on OLYMPUS IX83 and analyzed as follows:
nucleus: DAPI, LC3b: FITC, conidia: Cy5 channel at 20X.

siRNA knockdown of *Adipor1*

414   TriFECTa RNAi Kit was used with Lipofectamine 2000 reagent for transfection. Adherent 10 days old
415   alveolar macrophages were treated with diluted AdipoR1 SiRNA in Lipofectamine (1:1 ratio) at 1nM
416   per 5 x 10^5 cells concentration for 72 hours. The cells were then quantified for AdipoR1 receptor
417   expression using AdipoR1 RT-PCR and compared with provided positive and negative controls.

STATISTICAL ANALYSIS

419   Prism software was used for generation of figures and for statistical analyses (GraphPad). Unpaired *t*
420   tests or ANOVA tests were used to measure statistical significance. Differences between experimental
421   groups that resulted in a *p* value < 0.05 were considered significant.

References

1.   Veerdonk FL Van De, Gresnigt MS, Romani L, Netea MG, Latgé JP. Aspergillus fumigatus morphology
and dynamic host interactions. Nature Reviews Microbiology 2017 15:11. 2017;15: 661–674.
doi:10.1038/nrmicro.2017.90

2.   Carvalho A, Cunha C, Iannitti RG, Luca A De, Giovannini G, Bistoni F, et al. Inflammation in aspergillosis:
the good, the bad, and the therapeutic. Ann N Y Acad Sci. 2012;1273: 52–59. doi:10.1111/J.1749-
6632.2012.06754.X

3.   Sprenkeler EGG, Gresnigt MS, van de Veerdonk FL. LC3-associated phagocytosis: a crucial mechanism
for antifungal host defence against Aspergillus fumigatus. Cell Microbiol. 2016;18: 1208–1216.
doi:10.1111/CMI.12616

4.   Hoffstedt J, Arvidsson E, Sjölin E, Wåhlén K, Arner P. Adipose tissue adiponectin production and
adiponectin serum concentration in human obesity and insulin resistance. J Clin Endocrinol Metab.
2004;89: 1391–1396. doi:10.1210/JC.2003-031458

5.   Ren Y, Zhao H, Yin C, Lan X, Wu L, Du X, et al. Adipokines, Hepatokines and Myokines: Focus on Their
Role and Molecular Mechanisms in Adipose Tissue Inflammation. Front Endocrinol (Lausanne). 2022;13.
doi:10.3389/FENDO.2022.873699

6.   Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa JI, et al. Paradoxical decrease of an
adipose-specific protein, adiponectin, in obesity. Biochem Biophys Res Commun. 1999;257: 79–83.
doi:10.1006/BBRC.1999.0255

7.   Masamoto Y, Arai S, Sato T, Yoshimi A, Kubota N, Takamoto I, et al. Adiponectin enhances antibacterial
activity of hematopoietic cells by suppressing bone marrow inflammation. Immunity. 2016;44: 1422–
1433. doi:10.1016/j.immuni.2016.05.010

8.   Amarsaikhan N, Tsoggerel A, Hug C, Templeton SP. The metabolic cytokine adiponectin inhibits
inflammatory lung pathology in invasive aspergillosis. J Immunol. 2019;203: 956–963.
doi:10.4049/JIMMUNOL.1900174

9.   Bhatia S, Fei M, Yarlagadda M, Qi Z, Akira S, Saijo S, et al. Rapid host defense against Aspergillus
fumigatus involves alveolar macrophages with a predominance of alternatively activated phenotype.
PLoS One. 2011;6. doi:10.1371/journal.pone.0015943

10.   Chignard M, Balloy V, Sallenave JM, Si-Tahar M. Role of Toll-like receptors in lung innate defense

against invasive aspergillosis. Distinct impact in immunocompetent and immunocompromized hosts.
Clin Immunol. 2007;124: 238–243. doi:10.1016/J.CLIM.2007.05.004

11. Kubota N, Terauchi Y, Yamauchi T, Kubota T, Moroi M, Matsui J, et al. Disruption of adiponectin causes
insulin resistance and neointimal formation. J Biol Chem. 2002;277: 25863–25866.
doi:10.1074/JBC.C200251200

12. Ke M, Cabrero A, Saha PK, Kojima H, Li L, Chang BHJ, et al. Increased beta -oxidation but no insulin
resistance or glucose intolerance in mice lacking adiponectin. J Biol Chem. 2002;277: 34658–34661.
doi:10.1074/JBC.C200362200

13. Maeda N, Shimomura I, Kishida K, Nishizawa H, Matsuda M, Nagaretani H, et al. Diet-induced insulin
resistance in mice lacking adiponectin/ACRP30. Nat Med. 2002;8: 731–737. doi:10.1038/NM724

14. Jablonski KA, Amici SA, Webb LM, Ruiz-Rosado JDD, Popovich PG, Partida-Sanchez S, et al. Novel
Markers to Delineate Murine M1 and M2 Macrophages. PLoS One. 2015;10.
doi:10.1371/JOURNAL.PONE.0145342

15. Jhingran A, Mar KB, Kumasaka DK, Knoblaugh SE, Ngo LY, Segal BH, et al. Tracing conidial fate and
measuring host cell antifungal activity using a reporter of microbial viability in the lung. Cell Rep.
2012;2: 1762. doi:10.1016/J.CELREP.2012.10.026

16. Okada-Iwabu M, Yamauchi T, Iwabu M, Honma T, Hamagami K-I, Matsuda K, et al. A small-molecule
AdipoR agonist for type 2 diabetes and short life in obesity. Nature. 2013;503. doi:10.1038/nature12656

17. Salvator H, Grassin-Delyle S, Brollo M, Couderc LJ, Abrial C, Victoni T, et al. Adiponectin inhibits the
production of TNF-α, IL-6 and chemokines by human lung macrophages. Front Pharmacol. 2021;12:
718929. doi:10.3389/FPHAR.2021.718929

18. Viola A, Munari F, Sánchez-Rodríguez R, Scolaro T, Castegna A. The Metabolic Signature of Macrophage
Responses. Front Immunol. 2019;10. doi:10.3389/FIMMU.2019.01462

19. Sprenkeler EGG, Gresnigt MS, van de Veerdonk FL. LC3-associated phagocytosis: a crucial mechanism
for antifungal host defence against Aspergillus fumigatus. Cell Microbiol. 2016;18: 1208–1216.
doi:10.1111/CMI.12616

20. Gamberi T, Modesti A, Magherini F, D'Souza DM, Hawke T, Fiaschi T. Activation of autophagy by
globular adiponectin is required for muscle differentiation. Biochim Biophys Acta. 2016;1863: 694–702.
doi:10.1016/J.BBAMCR.2016.01.016

21.   Liu Y, Palanivel R, Rai E, Park M, Gabor T V., Scheid MP, et al. Adiponectin stimulates autophagy and
reduces oxidative stress to enhance insulin sensitivity during high-fat diet feeding in mice. Diabetes.
2015;64: 36–48. doi:10.2337/DB14-0267

22.   Essick EE, Wilson RM, Pimentel DR, Shimano M, Baid S, Ouchi N, et al. Adiponectin modulates oxidative
stress-induced autophagy in cardiomyocytes. PLoS One. 2013;8. doi:10.1371/JOURNAL.PONE.0068697

23.   Li C, Wang Z, Wang C, Ma Q, Zhao Y. Perivascular adipose tissue-derived adiponectin inhibits collar-
induced carotid atherosclerosis by promoting macrophage autophagy. PLoS One. 2015;10.
doi:10.1371/JOURNAL.PONE.0124031

24.   Kanayama M, He Y-W, Shinohara ML. The lung is protected from spontaneous inflammation by
autophagy in myeloid cells. J Immunol. 2015;194: 5465–5471. doi:10.4049/JIMMUNOL.1403249

25.   Asare PF, Roscioli E, Hurtado PR, Tran HB, Mah CY, Hodge S. LC3-Associated Phagocytosis (LAP): A
Potentially Influential Mediator of Efferocytosis-Related Tumor Progression and Aggressiveness. Front
Oncol. 2020;10. doi:10.3389/FONC.2020.01298

26.   Galvan MD, Hulsebus H, Heitker T, Zeng E, Bohlson SS. Complement protein C1q and adiponectin
stimulate Mer tyrosine kinase-dependent engulfment of apoptotic cells through a shared pathway. J
Innate Immun. 2014;6: 780–792. doi:10.1159/000363295

27.   Tee JH, Vijayakumar U, Shanmugasundaram M, Lam TYW, Liao W, Yang Y, et al. Isthmin-1 attenuates
allergic Asthma by stimulating adiponectin expression and alveolar macrophage efferocytosis in mice.
Respir Res. 2023;24. doi:10.1186/S12931-023-02569-1

28.   Madenspacher JH, Morrell ED, Gowdy KM, McDonald JG, Thompson BM, Muse G, et al. Cholesterol 25-
hydroxylase promotes efferocytosis and resolution of lung inflammation. JCI Insight. 2020;5.
doi:10.1172/JCI.INSIGHT.137189

29.   Oikonomou V, Moretti S, Renga G, Galosi C, Borghi M, Pariano M, et al. Noncanonical Fungal Autophagy
Inhibits Inflammation in Response to IFN-γ via DAPK1. Cell Host Microbe. 2016;20: 744–757.
doi:10.1016/J.CHOM.2016.10.012

30.   Akoumianaki T, Kyrmizi I, Valsecchi I, Beauvais A, Latge J-P, Chamilos G. Aspergillus cell wall melanin
blocks LC3-associated phagocytosis to promote pathogenicity. Cell Host Microbe. 2016;19: 79–90.
doi:10.1016/j.chom.2015.12.002

31.   Kyrmizi I, Ferreira H, Carvalho A, Figueroa JAL, Zarmpas P, Cunha C, et al. Calcium sequestration by

fungal melanin inhibits calcium-calmodulin signalling to prevent LC3-associated phagocytosis. Nat Microbiol. 2018;3: 791–803. doi:10.1038/S41564-018-0167-X

32. De Luca A, Smeekens SP, Casagrande A, Iannitti R, Conway KL, Gresnigt MS, et al. IL-1 receptor blockade restores autophagy and reduces inflammation in chronic granulomatous disease in mice and in humans. Proc Natl Acad Sci U S A. 2014;111: 3526–3531. doi:10.1073/PNAS.1322831111

33. Amarsaikhan N, Stolz DJ, Wilcox A, Sands EM, Tsoggerel A, Gravely H, et al. Reciprocal inhibition of adiponectin and innate lung immune responses to chitin and *Aspergillus fumigatus*. Front Immunol. 2019;10: 1057. doi:10.3389/FIMMU.2019.01057

34. Yamauchi T, Kadowaki T. Adiponectin receptor as a key player in healthy longevity and obesity-related diseases. Cell Metab. 2013;17: 185–196. doi:10.1016/J.CMET.2013.01.001

35. Hu G, Christman JW. Editorial: Alveolar Macrophages in Lung Inflammation and Resolution. Front Immunol. 2019;10. doi:10.3389/FIMMU.2019.02275

36. Byrne AJ, Mathie SA, Gregory LG, Lloyd CM. Pulmonary macrophages: key players in the innate defence of the airways. Thorax. 2015;70: 1189–1196. doi:10.1136/THORAXJNL-2015-207020

37. Orecchioni M, Ghosheh Y, Pramod AB, Ley K. Macrophage Polarization: Different Gene Signatures in M1(LPS+) vs. Classically and M2(LPS-) vs. Alternatively Activated Macrophages. Front Immunol. 2019;10. doi:10.3389/FIMMU.2019.01084

38. Colombo G, Pessolano E, Talmon M, Genazzani AA, Kunderfranco P. Getting everyone to agree on gene signatures for murine macrophage polarization in vitro. PLoS One. 2024;19. doi:10.1371/JOURNAL.PONE.0297872

39. Yunna C, Mengru H, Lei W, Weidong C. Macrophage M1/M2 polarization. Eur J Pharmacol. 2020;877. doi:10.1016/J.EJPHAR.2020.173090

40. Parisi L, Gini E, Baci D, Tremolati M, Fanuli M, Bassani B, et al. Macrophage Polarization in Chronic Inflammatory Diseases: Killers or Builders? J Immunol Res. 2018;2018. doi:10.1155/2018/8917804

41. Dowling JK, Afzal R, Gearing LJ, Cervantes-Silva MP, Annett S, Davis GM, et al. Mitochondrial arginase-2 is essential for IL-10 metabolic reprogramming of inflammatory macrophages. Nat Commun. 2021;12. doi:10.1038/S41467-021-21617-2

42. Steffes MW, Gross MD, Schreiner PJ, Yu X, Hilner JE, Gingerich R, et al. Serum adiponectin in young adults - Interactions with central adiposity, circulating levels of glucose, and insulin resistance: The

CARDIA study. Ann Epidemiol. 2004;14: 492–498. doi:10.1016/j.annepidem.2003.10.006

43.   Sood A, Shore SA. Adiponectin, Leptin, and Resistin in Asthma: Basic Mechanisms through Population
Studies. J Allergy (Cairo). 2013;2013: 1–15. doi:10.1155/2013/785835

44.   Ghigliotti G, Barisione C, Garibaldi S, Fabbi P, Brunelli C, Spallarossa P, et al. Adipose tissue immune
response: Novel triggers and consequences for chronic inflammatory conditions. Inflammation. Springer
New York LLC; 2014. pp. 1337–1353. doi:10.1007/s10753-014-9914-1

45.   Kadowaki T, Yamauchi T. Adiponectin and adiponectin receptors. Endocr Rev. 2005;26: 439–451.
doi:10.1210/ER.2005-0005

46.   Sharma S, Malur A, Marshall I, Huizar I, Barna BP, Pories W, et al. Alveolar macrophage activation in
obese   patients   with   obstructive   sleep   apnea.   Surgery.   2012;151:   107–112.
doi:10.1016/J.SURG.2011.06.035

47.   Jhingran A, Kasahara S, Hohl T. Flow Cytometry of Lung and Bronchoalveolar Lavage Fluid Cells from
Mice Challenged with Fluorescent Aspergillus Reporter (FLARE) Conidia. Bio Protoc. 2016;6.
doi:10.21769/bioprotoc.1927

48.   Stolz DJ, Sands EM, Amarsaikhan N, Tsoggerel A, Templeton SP. Histological quantification to determine
lung fungal burden in experimental aspergillosis. JoVE (Journal of Visualized Experiments). 2018;2018:
e57155–e57155. doi:10.3791/57155

49.   Bhattacharya S, Amarsaikhan N, Maupin AJ, Schlosser A, Füchtbauer E-M, Holmskov U, et al. FIBCD1
deficiency decreases disease severity in a murine model of invasive pulmonary aspergillosis.
Immunohorizons. 2021;5: 983–993. doi:10.4049/IMMUNOHORIZONS.2100092

50.   Amarsaikhan N, Sands EM, Shah A, Abdolrasouli A, Reed A, Slaven JE, et al. Caspofungin increases
fungal chitin and eosinophil and $\gamma\delta$ T Cell-dependent pathology in invasive aspergillosis. J Immunol.
2017;199: 624–632. doi:10.4049/JIMMUNOL.1700078

51.   Lim JY, Jung YE, Hwang HE, Kim CH, Basaran-Akgul N, Goli SH, et al. Survival Factor A (SvfA) Contributes
to Aspergillus nidulans Pathogenicity. J Fungi (Basel). 2023;9. doi:10.3390/JOF9020143

52.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq
aligner. Bioinformatics. 2013;29: 15–21. doi:10.1093/BIOINFORMATICS/BTS635

53.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence
reads to genomic features. Bioinformatics. 2014;30: 923–930. doi:10.1093/BIOINFORMATICS/BTT656

54.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression
570   analysis of digital gene expression data. Bioinformatics. 2010;26: 139–140.
571   doi:10.1093/BIOINFORMATICS/BTP616

572   55.   McCarthy DJ, Chen Y, Smyth GK. Differential expression analysis of multifactor RNA-Seq experiments
573   with respect to biological variation. Nucleic Acids Res. 2012;40: 4288–4297. doi:10.1093/NAR/GKS042

574   56.   Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological themes among
575   gene clusters. OMICS. 2012;16: 284–287. doi:10.1089/OMI.2011.0118

576   57.   Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment tool for
577   interpreting omics data. Innovation (Cambridge (Mass)). 2021;2. doi:10.1016/J.XINN.2021.100141

Figure 1.

Figure 1. Increased mortality, fungal burden, and inflammation in APN pathway-deficient mice
with invasive aspergillosis.

Wild-type (C57BL/6), APN-/- (*Adipoq*$^{-/-}$), *AdipoR1*$^{-/-}$, and *AdipoR2*$^{-/-}$ mice were neutrophil-depleted
and involuntarily aspirated a suspension of *A. fumigatus* conidia and followed for survival or sacrificed
on day 3 post-infection as described in Materials and Methods.

A.   Experimental time course for neutropenic mouse model of infection.
B.   Survival. N = 8-16 mice per group.
C.   Representative GMS and H&E lung sections.
D.   Fungal burden determined by quantitative PCR of fungal DNA from lung homogenates.
E.   Fungal burden determined by quantification of GMS staining.
F.   Total number of CD45$^+$ cells, eosinophils (CD45$^+$Ly6G$^-$CD11c$^-$SiglecF$^+$), AMs
(CD45$^+$Ly6G$^-$CD11c$^-$SiglecF$^+$), CD11c$^+$SiglecF$^-$ (CD45$^+$Ly6G$^-$ CD11c$^+$SiglecF$^-$), and CD11c$^-$SiglecF$^-$
(CD45$^+$Ly6G$^-$CD11c$^-$SiglecF$^-$) cells isolated from the mice with IA as determined by flow cytometry. N
= 4-6 mice per group.
G.   qRT-PCR analysis for mRNA expression of the indicated cytokines.
H.   TNFα secretion in BALF quantified by ELISA.

A-D, data are a summary of two to three independently performed experiments. F-H, data are

representative of two independent experiments with similar results. *p < 0.05, **p < 0.01, ***p <

0.001.

Figure 2.

A

B

C

D

Ctr

E

Infected

Figure 2. Alveolar macrophages from APN pathway-deficient mice exhibit an inflammatory
phenotype.

BALF cells from wild-type (C57BL/6), APN-/- (*Adipoq$^{-/-}$*), *AdipoR1$^{-/-}$*, and *AdipoR2$^{-/-}$* mice were
collected in PBS/EDTA. For alveolar macrophage differentiation, total BALF cells were plated and
cultured with GM-CSF. Alveolar macrophages were challenged with conidia in the ratio 1:9 (AM :
conidia) for 10 hours duration as per the details mentioned in Materials and Methods. The data
represented is AMs from n=3 mice/group.

A.   qRT-PCR analysis for mRNA expression of the indicated cytokines.
B.   TNFα secretion quantified at the protein level by ELISA using *ex vivo* cultured alveolar
macrophages.
C.   Flow cytometry staining of surface CD38 and intracellular Egr2

(D-E) Quantification of (C), showing the proportion of M1, M2 and CD38$^-$Egr2$^+$ macrophages with
CD38$^+$Egr2$^-$ (putative M1) or CD38$^-$Egr2$^+$ (putative M2) phenotype in control group (D) and infection
group (E).

Data are a summary of three independently performed experiments. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 3.

Fig. 3. *A. fumigatus* conidial uptake and killing are inhibited in adiponectin-deficient AMs.

A.   Microscopic analysis of uptake and killing of FLARE conidia with magnification 40x is depicted in Brightfield, Cy5, DsRed channels followed by the merged image. Scale bar is equal to 12μm.

B.   Quantification of *Aspergillus* viability using microscopy images via calculation of co-localization in imageJ.

C.   Representative plots that display RFP and AF633 fluorescence intensity of *ex vivo* cultured alveolar macrophages where R1 represents live conidia inside the alveolar macrophages and R2 represents dead/killed conidia .

D.   Quantification of (C), showing *Aspergillus* viability (R1/(R1 + R2)) and uptake (R1 + R2) in *ex vivo* cultured alveolar macrophages.

E.   Representative plots that display RFP and AF633 fluorescence intensity of AMs (CD45$^+$Ly6G$^-$CD11c$^+$SiglecF$^+$) *in vivo.* Mice of both groups were made immunocompromised and infected with 1-1.5 × 10^7 conidia. BAL cells were harvested 1 day and 3 days later.

F.   Quantification of (E), showing *Aspergillus* viability (R1/(R1 + R2)) and uptake (R1 + R2) in alveolar macrophages in neutropenic mice at d1 and d3 p.i.

G.   Fluorescent particle uptake by alveolar macrophages in vivo. For flow cytometric analysis, neutrophil-depleted mice were challenged with 1.5 × 10^7 of flow particles. Percentage of flow particle positive (FP+) cells was assessed in AMs (CD45$^+$Ly6G$^-$CD11c$^+$SiglecF$^+$) from the wild-type and *Adipoq* $^{-/-}$ mice by flow cytometry.

A-D, data are representative of three independent experiments. E-G, data are a summary of 2-3 independent experiments. *p < 0.05, and **p < 0.01.

Figure 4.

Figure 4. The AdipoR agonist AdipoRon improves decreases inflammation and fungal burden in APN-deficient mice with IA

Wild-type (C57BL/6), and *Adipoq⁻ᐟ⁻* mice were neutrophil depleted and involuntarily aspirated *A. fumigatus* conidia as described in Materials and Methods.

A.　　Experimental time course for neutropenic mouse model of infection and AdipoRon treatment.

B.　　Effect of AdipoRon on the survival rate of WT and *Adipoq⁻ᐟ⁻* mice. N = 9-11 mice per group.

C.　　C. Representative GMS and H&E lung sections.

D.　　Fungal burden determined by quantitative PCR of fungal DNA from lung homogenates.

E.　　Fungal burden determined by quantification of GMS staining.

F.　　Total number of CD45⁺ cells, eosinophils (CD45⁺Ly6G⁻CD11c⁻SiglecF⁺), AMs (CD45⁺Ly6G⁻CD11c⁻SiglecF⁺), CD11c⁺SiglecF⁻ (CD45⁺Ly6G⁻ CD11c⁺SiglecF⁻), and CD11c⁻SiglecF⁻ (CD45⁺Ly6G⁻CD11c⁻SiglecF⁻) cells isolated from the mice with IA as determined by flow cytometry. N = 4-6 mice per group.

G.　　qRT-PCR analysis for mRNA expression of the indicated cytokines.

H.　　TNFα secretion in BALF quantified at the protein level by ELISA.

Data are a summary of 2-3 independently performed experiments. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 5.

Fig. 5. AdipoRon increases AdipoR-dependent AM fungal killing and reverses inflammatory phenotype.

A.   Experimental time course for *ex vivo* cultured AMs challenged with conidia with or without AdipoRon treatment.

B.   Microscopic image quantification of *Aspergillus* viability (R1/(R1 + R2)) in *ex vivo* cultured AMs by using FLARE conidia. The infection was done ex-vivo in 1:1 ratio (AMs: FLARE) for 15 hours.

C.   Flow cytometric quantification of *Aspergillus* viability.

D.   qRT-PCR analysis for mRNA expression of the indicated cytokines.

E.   TNF secretion quantified at the protein level using *ex vivo* cultured AMs by ELISA.

Data are representative of three independently performed experiments. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 6.

A

C

B

AdipoR1

D

Fig. 6. AdipoRon-enhanced killing of *A. fumigatus* requires both AdipoR1 and AdipoR2.

A. Schematic timeline course for primary AM cultures treated with *AdipoR1* siRNA, details in
Methods section.
B. Histogram graphs obtained from flow cytometry analysis. The positive control represents the
ideal *Adipor1* knockdown control, while the negative control is in the absence of functional SiRNA.
C. qRT-PCR analysis for mRNA expression of the *adipor1* in AMs from the APN-deficient mice
with siRNA treatment.
D. Conidial viability (R1/(R1 + R2)) in *ex vivo* cultured AMs with siRNA and/or AdipoRon by
flow cytometry

Data are representative of two independently performed experiments. **$p < 0.01$ and ***$p < 0.001$.

Figure 7.

A

C

D

B

Fig. 7. Adiponectin promotes LC3-associated phagocytosis of *A. fumigatus* conidia.

A.   Schematic timeline course for LC3-associated phagocytosis analysis.

B.   Microscopy of LAP-dsRed conidia and LC3$_+$ AMs at a magnification of 20X in *ex vivo* cultured
AMs from infected WT and *Adipoq$^{-/-}$* mice with or without AdipoRon. The images from DsRed, FITC,
DAPI channels are presented followed by the merged image. Scale bar: 35µm.

C.   Microscopic quantification of LC3$_+$ phagosome (%) in *ex vivo* cultured AMs from infected WT
and *Adipoq$^{-/-}$* mice with or without AdipoRon.

D.   The mean fluorescent intensity (MFI) obtained by flow cytometry indicates the expression level
of LC3-AF488 in *ex vivo* cultured AMs from infected WT and *Adipoq$^{-/-}$* mice with or without
AdipoRon.

Data are representative of two independently performed experiments. *$p < 0.05$ and ***$p < 0.001$.

Supplemental Information

Supplemental Methods

Mice

An additional strain of Adipoq-/- mice was obtained from Dr. Philipp Scherer (University of Texas-

Southwestern) and used for survival, fungal burden, histology, flow cytometric analysis of BALF cells,

qRT-PCR of selected inflammatory and *Adipor* genes, and ELISA for TNF as described in the

Materials and Methods section.

Quantification of AM AdipoR1 expression using flow cytometry

After ex-vivo AM extraction and culture as mentioned in the methods, AMs were fixed using IC

fixation buffer. Fc Block was used to eliminate non-specific Fc-mediated interactions. The cells were

then stained at a 1:100 dilution with primary antibody with Adiponectin Receptor 1 Recombinant

Rabbit Monoclonal Antibody (SC69-04) followed by staining with Alexa Fluor 488-conjugated goat

anti rabbit IgG as the secondary antibody. All the unstained and only primary antibody staining

controls were included.

Quantification of AM *Adipor1* and *Adipor2* receptor gene expression

RNA extraction from AMs was done using Qiagen RNeasy Mini kit, following the manufacturer's

protocol. Quantitative RT-PCR was performed with 20ng of cDNA using AdipoR1 and AdipoR2

forward and reverse primers ordered from IDT Integrated Technologies. Gene Expression Master Mix

(ThermoFisher Scientific) was used, with β-actin used for signal normalization.

RNA sequencing and analysis

Sample Preparation

AMs were isolated from APN-/- mice, infected/uninfected for 10 hours (1:9 cells/conidia) and

AdipoRon/vehicle treated 24 hours before infection, with cells lysed and RNA isolated as described in

Materials and Methods.

Library preparation and sequencing

Total RNA samples were first evaluated for their quantity and quality using Agilent TapeStation. All

the samples were good quality with RIN (RNA Integrity Number) of 9.6-10. One hundred nanograms

of total RNA was used for library preparation with the Illumina Stranded mRNA Prep, Ligation kit

(lllumina), following the manufacturer's instruction. Each resulting uniquely dual-indexed library was

quantified and quality accessed by Qubit and Agilent TapeStation, and multiple libraries were pooled

in equal molarity. The pooled libraries were sequenced with 2×150bp paired-end configuration on an

Illumina NovaSeq X PLUS sequencer.

RNA-seq data analysis

The sequencing reads were first quality checked using FastQC (v.0.11.5, Babraham Bioinformatics,

Cambridge, UK) for quality control. The sequence data were then mapped to the mouse reference

genome mm10 using the RNA-seq aligner STAR (v.2.7.10a)[52] with the following parameter: "--

outSAMmapqUnique 60". To evaluate quality of the RNA-seq data, the number of reads that fell into

different annotated regions (exonic, intronic, splicing junction, intergenic, promoter, UTR, etc.) of the

reference genome was assessed using bamutils (from ngsutils v.0.4.17).[53] Uniquely mapped reads

were used to quantify the gene level expression employing featureCounts (subread v.2.0.3)[54] with the

following parameters: "-s 2 -Q 10". The data was normalized using TMM (trimmed mean of M values)

method. Differential expression analysis was performed using edgeR (v.4.0.1).[54,55] False discovery

rate (FDR) was computed from p-values using the Benjamini-Hochberg procedure. Gene Ontology

(GO) pathway enrichment analyses were performed with the R package clusterProfiler.[56,57]

Supplemental Figure S1.

A

B

C

D

E

F

G

H

Fig. S1. Inflammatory phenotype of APN-/- mice from a second strain.

Wild-type (C57BL/6) and APKO (Second *Adipoq*$^{-/-}$ mice from Scherer's laboratory) were neutrophil
depleted and involuntarily aspirated $1 - 1.5 \times 10^7$ of conidia as described in Materials and Methods.

A. Survival rate. $N = 9$ and 15 mice per group, respectively.
B. Representative GMS and H&E lung sections.
C. Fungal burden determined by quantitative PCR of fungal DNA from lung homogenates.
D. Fungal burden determined by quantification of GMS staining.
E. Total number of CD45$^+$ cells, eosinophils (CD45$^+$Ly6G$^-$CD11c$^-$SiglecF$^+$), AMs
(CD45$^+$Ly6G$^-$CD11c$^+$SiglecF$^+$), CD11c$^+$SiglecF$^-$ (CD45$^+$Ly6G$^-$ CD11c$^+$SiglecF$^-$), and CD11c$^-$SiglecF$^-$
(CD45$^+$Ly6G$^-$CD11c$^-$SiglecF$^-$) cells isolated from the mice with IA as determined by flow cytometry. N
= 4-6 mice per group.
F. qRT-PCR analysis for mRNA expression of the indicated cytokines.
G. qRT-PCR analysis for mRNA expression of *Adipor1* and *Adipor2*.
H. TNFα secretion in BALF quantified at the protein level by ELISA.

Data are a summary of two to three independently performed experiments. *p < 0.05, **p < 0.01, ***p
< 0.001.

Supplemental Figure S2.

A

Lung

BALF

B

C

D

E

Fig. S2. AdipoRs expression in the Lung and BALF in APN pathway-deficient mice with invasive
aspergillosis.

A.　　Expression of *Adipoq* and *Adipor* genes in lung homogenates and BALF cells from non-
infected (NI) and conidia-infected WT mice. Lung and BALF were collected from the mice at 3 dpi.
B.　　Summary of median fluorescence intensities of AdipoR1 staining on AMs
(CD45+Ly6G-CD11c+SiglecF+), eosinophils (CD45+Ly6G-CD11c-SiglecF+), CD11c+SiglecF-
(CD45+Ly6G- CD11c+SiglecF-), and CD11c-SiglecF- (CD45+Ly6G-CD11c-SiglecF-) cells from non-
infected (NI) and conidia-infected WT mice.
C.　　qRT-PCR analysis for mRNA expression of *Adipor1* and *Adipor2* in lung homogenates. Wild-
type (C57BL/6), *Adipoq*-/-, *AdipoR1*-/-, and *AdipoR2*-/-mice were neutrophil depleted and involuntarily
aspirated *A. fumigatus* conidia.
D.　　Flow cytometry staining of AdipoR1. Frequency of AdipoR1+ in ex-vivo cultured AMs in WT
and *Adipoq*-/- mice. The histogram represents the AdipoR1 peak relative to IgG control in infected vs
uninfected.
E.　　qRT-PCR analysis for mRNA expression of *Adipor1* and *Adipor2* from the ex-vivo cultured
AMs. The AMs were challenged with AF-293 conidia with or without AdipoRon treatment.

Data are a summary of two to three independently performed experiments. $*p < 0.05$, $**p < 0.01$, $***p$
< 0.001.

Supplemental Figure S3.

Fig. S3 Gene expression in AdipoRon treated/untreated and infected/ uninfected *Adipoq⁻ᐟ⁻* AMs
by RNAseq analysis.

AMs were infected with swollen AF293 conidia with 1:9 cells/conidia for 10 hours, or left uninfected,
with or without AdipoRon treatment, followed by RNA extraction for RNAseq analysis (N=3/group).

A. Volcano plot depicting relative changes in gene expression of AdipoRon-treatment in APN-/-
AMS, infected (top) or uninfected (bottom).
B. Heat map representation of the genes with highest differential expression in infected (left) and
uninfected (right) APN-/- AMs.
C. GSEA-GO analysis of gene pathways that are both differentially expressed by AdipoRon treatment
in infected (left) and uninfected (right) APN-/- AMs.
790

791   **Supplemental Video SV1. Time course of uptake and killing in WT and APN-/- AMs.**

792   Alveolar macrophages were plated and infected as mentioned in the methods. Cells were live imaged
793   by confocal microscopy for T= 8min under 40X water immersion lens. The black arrows indicate the
794   live conidia at T=0 min and the white arrows indicate the conidia at T=8min.

796