BioMapAI: Artificial Intelligence Multi-Omics Modeling of Myalgic Encephalomyelitis / Chronic Fatigue Syndrome

Author list

Ruoyun Xiong[1,2], Elizabeth Fleming[1], Ryan Caldwell[1], Suzanne D. Vernon[3], Lina Kozhaya[1], Courtney Gunter[1,2], Lucinda Bateman[3], Derya Unutmaz[1], Julia Oh[1,*]

1 The Jackson Laboratory, Farmington, Connecticut, USA. 06032

2 The University of Connecticut Health Center, Farmington, Connecticut, USA. 06030

3 Bateman Horne Center, Salt Lake City, Utah, USA. 84102

* Corresponding author and lead contact:

Julia Oh, Ph.D.

The Jackson Laboratory

10 Discovery Drive

Farmington, CT

860-837-2014

julia.oh@jax.org

Abstract

Chronic diseases like ME/CFS and long COVID exhibit high heterogeneity with multifactorial etiology and progression, complicating diagnosis and treatment. To address this, we developed BioMapAI, an explainable Deep Learning framework using the richest longitudinal multi-'omics dataset for ME/CFS to date. This dataset includes gut metagenomics, plasma metabolome, immune profiling, blood labs, and clinical symptoms. By connecting multi-'omics to asymptom matrix, BioMapAI identified both disease- and symptom-specific biomarkers, reconstructed symptoms, and achieved state-of-the-art precision in disease classification. We also created the first connectivity map of these 'omics in both healthy and disease states and revealed how microbiome-immune-metabolome crosstalk shifted from healthy to ME/CFS. Thus, we proposed several innovative mechanistic hypotheses for ME/CFS: Disrupted microbial functions – SCFA (butyrate), BCAA (amino acid), tryptophan, benzoate - lost connection with plasma lipids and bile acids, and activated inflammatory and mucosal immune cells (MAIT, γδT cells) with INFγ and GzA secretion. These abnormal dynamics are linked to key disease symptoms, including gastrointestinal issues, fatigue, and sleep problems.

Introduction

Chronic diseases, such as cancer[1], diabetes[2], rheumatoid arthritis (RA)[3], myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS)[4], and possibly long COVID[5,6], the sequela of SARS-CoV-2 infection, can evolve over decades and exhibit diverse phenotypic and physiological manifestations across individuals. This heterogeneity is reflected in disease progression and treatment responses, complicating the establishment of standardized clinical protocols, and demanding personalized therapeutic strategies[7].

However, this heterogeneity has not been well studied, leaving substantial knowledge and
technical gaps[8]. Current cohort studies often focus on identifying one or two key disease
indicators, such as HbA1C levels for diabetes[9,10] or survival rates for cancer[11], even with the
advent of multi-'omics. This approach has difficulty accommodating the highly multifactorial
etiology and progression of most chronic diseases, with different patients exhibiting varying
symptoms and disease markers[12]. To address this challenge, methods must link a more complex
matrix of disease-associated outcomes with a range of 'omics data types to enable precise
targeting of biomarkers tailored to each patient's specific symptoms.
52
Here, we introduce BioMapAI, an explainable AI framework that we developed to integrate
multi-'omics data to decode complex host symptomatology, specifically applied to ME/CFS.
Affecting at least 10 million people globally, ME/CFS is a chronic, complex, multi-system illness
characterized by impaired function and persistent fatigue, post-exertional malaise, multi-site
pain, sleep disturbances, orthostatic intolerance, cognitive impairment, gastrointestinal issues,
and other symptoms [13,14,15]. The pathogenesis of ME/CFS is not well understood, with triggers
believed to include viral infections such as Epstein-Barr Virus (EBV)[16], enteroviruses[17] and SARS
coronavirus[18]. As a chronic disease, ME/CFS can persist for years or even a lifetime, with each
patient developing distinct illness patterns[13]. Therefore, a universal approach to clinical care and
symptom management is insufficient, and a personalized approach is crucial for effectively
addressing the complex nature of ME/CFS. Additionally, given similarities in causality and
symptomatology to long COVID[19,20], studying ME/CFS specifically can provide broader insights
into post-viral syndromes, and more generally, our AI-driven approach can be applied to a range
of diseases with complex symptomatology not readily explained by a single data type.
67
We generated a rich longitudinal, multi-'omics dataset of 153 ME/CFS patients and 96 age-
gender-matched healthy controls, comprised of gut metagenomics, plasma metabolome,
immune cell profiling, activation, and cytokines, together with blood labs, detailed clinical
symptoms, and lifestyle survey data. We aimed to: 1) identify new disease biomarkers - not only
for ME/CFS but also to specify biomarkers that could explain the complex symptomatology, and
2) define interactions between microbiome, immune system, and metabolome – rather than
studying single data types in isolation, we created the first connectivity map of these 'omics.
This map critically accounts for covariates such as age and gender, providing an important
baseline in healthy individuals contrasted with aberrant connections identified in disease.
77
BioMapAI is a strategically designed Deep Neural Network (DNN) that connected the multi-
'omics profiles to a matrix of clinical symptoms. Here, applying to ME/CFS, it identifies both
disease- and symptom-specific biomarkers, accurately reconstructing key clinical symptoms,
achieves state-of-the-art precision in disease classification, and generates several innovative
mechanistic hypotheses for disease. By revealing microbiome-immune-metabolome crosstalk
shifts from healthy to diseased states, we found depletion of microbial butyrate (SCFA) and
amino acids (BCAA) in ME/CFS, linked with abnormal activation of inflammatory and mucosal
immune cells – MAIT and γδT cells with INFγ and GzA. This altered dynamic correlated with
clinical symptom scores, indicating deteriorated health perception and impaired social activity.
Microbial metabolites, like tryptophan and benzoate, lost connections with plasma lipids in

patients, in turn associated with fatigue, emotional and sleeping problems. This dataset is the
richest multi-'omics dataset for ME/CFS, as well as for numerous other chronic diseases to date.
It introduces a novel, generalizable, and explainable AI approach that captures the complexity of
chronic disease and provides new hypotheses for host-microbiome interactions in both health
and ME/CFS.
93
94   **Results**
95
Cohort Overview
We tracked 249 participants over 3-4 years, including 153 ME/CFS patients (75 'short-term' with
disease symptoms < 4 years and 78 'long-term' with disease symptoms > 10 years) and 96
healthy controls (Fig 1A; Supplemental Table 1). The cohort is 68% female and 32% male,
aligning with the epidemiological data showing that women are 3-4 times more likely to develop
ME/CFS[21],[22]. Participants ranged in age from 19 to 68 years with body mass indexes (BMI) from
16 to 43 kg/m². Throughout the study, we collected detailed clinical metadata, blood samples,
and fecal samples. In total, 1471 biological samples were collected across all participants at 515
timepoints (Methods, Supplemental Figure 1A, Supplemental Table 1).
105
Blood samples were 1) sent for clinical testing at Quest Laboratory (48 features measured,
N=503 samples), 2) fractionated into peripheral blood mononuclear cells (PBMCs), which were
examined via flow cytometry, yielding data on 443 immune cells and cytokines (N=489), 3)
plasma and serum, for untargeted liquid chromatography with tandem mass spectrometry (LC-
MS/MS), identifying 958 metabolites (N=414). Detailed demographic documentation and
questionnaires covering medication use, medical history, and key ME/CFS symptoms were
collected (Methods). Finally, whole-genome shotgun metagenomic sequencing of stool samples
(N=479) produced an average of 12,302,079 high-quality, classifiable reads per sample, detailing
gut microbiome composition (1293 species detected) and KEGG gene function (9993 genes
reconstructed).
116
Heterogeneity and Non-linear Progression of ME/CFS
First, we demonstrated the phenotypic complexity and heterogeneity of ME/CFS. Collaborating
with clinical experts, we consolidated detailed questionnaires and clinical metadata,
foundational to diagnosing ME/CFS, into twelve essential clinical scores (Methods). These scores
covered core symptoms including physical and mental health, fatigue, pain levels, cognitive
efficiency, sleep disturbances, orthostatic intolerance, and gastrointestinal issues (Supplemental
Table 1).
124
While healthy individuals consistently presented low symptom scores (Supplemental Figure 1D),
ME/CFS patients exhibited significant variability in symptom severity, with each individual
showing different predominant symptoms (Figure 1B). Principal coordinates analysis (PCoA) of
the 'omics matrices highlighted the difficulty in distinguishing patients from controls,
emphasizing the complex symptomatology of ME/CFS and the challenges in developing
predictive models (Supplemental Figure 1E). Additionally, over time, in contrast to the stable
patterns typical of healthy individuals (Supplemental Figure 1B), ME/CFS patients demonstrated

distinctly varied patterns each year, as evidenced by the diversity in symptom severity and
noticeable separation on the 'omics PCoA (Figure 1B, Supplemental Figure 1C). Despite
employing multiple longitudinal models (Methods), we found no consistent temporal signals,
confirming the non-linear progression of ME/CFS.
136
This individualized, multifaceted, and dynamic nature of ME/CFS that intensifies with disease
progression necessitates new approaches that extend beyond simple disease versus control
comparisons. Here, we created and implemented an AI-driven model that integrates the
multi-'omics profiles to learn host phenotypes. This allowed us not only to develop a state-of-
the-art classifier for disease, but for the first time, to identify biomarker sets for each clinical
symptom as well as unique interaction networks that differed between patients and controls.
143

BioMapAI, an Explainable Neural Network Connecting 'Omics to Multi-Type Outcomes

145   To connect multi-'omics data to clinical symptoms, a model must accommodate the learning of
146   multiple different outcomes within a single framework. However, traditional machine learning
147   models are generally designed to predict a single categorical outcome or continuous
148   variable[23],[24],[25]. This simplified disease classification and conventional biomarker identification
149   typically fails to encapsulate the heterogeneity of complex diseases[26],[27].
150
We developed an AI-powered multi-'omics framework, BioMapAI, a fully connected deep neural
network that inputs 'omics matrices (X), and outputs a mixed-type outcome matrix (Y), thereby
mapping multiple 'omics features to multiple clinical indicators (Figure 2A). By assigning specific
loss functions for each output, BioMapAI aims to comprehensively learn every y (i.e., each of
the 12 continuous or categorical clinical scores in this study), using the 'omics data inputs.
Between the input layer X and the output layer $Y = [y_1, y_2, …, y_n]$, the model consists of two
shared hidden layers (Z^1 with 64 nodes, and Z^2 with 32 nodes) for general pattern learning,
followed by a parallel hidden layer ($Z^3 = [z_1^3, z_2^3, …, z_n^3]$), with sub-layers ($z_n^3$, each with 8
nodes) tailored for each outcome (y_n), to capture outcome-specific patterns (Figure 2A). This
unique architecture – two shared and one specific hidden layer – allows the model to capture
both general and output-specific patterns. This model is made 1) explainable by incorporating a
SHAP (SHapley Additive exPlanations) explainer, which quantifies the feature importance of
each predictions, providing both local (symptom-level) and global (disease-level)
interpretability, and 2) flexible by automatically finding appropriate learning goals and loss
functions for each type of outcomes (without need of format refinement), facilitating
BioMapAI's adaptability to broader research applications.
167

BioMapAI Reconstructed Clinical Symptoms and Achieved State-of-the-Art Performance in Discriminating ME/CFS from Healthy Controls

BioMapAI is a versatile AI framework connecting a biological 'omics matrix to multiple
phenotypic outputs. It does not have a specific disease focus and is designed to be applicable to
a range of applications. Here, we trained and validated its usage with our ME/CFS datasets,
employing a five-fold cross-validation. This trained model, nicknamed DeepMECFS for the
ME/CFS community, accurately represented the structure of diverse clinical symptom score
types and discriminated between healthy individuals and patients (Figure 2, Supplemental

Figure 2, Supplemental Table 2-3). For example, it effectively differentiated the physical health
scores, where patients exhibited more severe conditions compared to healthy controls
(category datatype 4 vs. 0, respectively, Figure 2B, Supplemental Table 2) and pain scores
(continuous datatype ranging from 1(highest)- 0(lowest), mean 0.52±0.24 vs. 0.11±0.12 for
patients vs. controls). Though compressing some inherent variance, BioMapAI accurately
reconstructed key statistical measures such as the mean and interquartile range (25%-75%), and
highlighted the distinctions between healthy and disease. (Figure 2B, Supplemental Figure 2A-B,
Supplemental Table 2).

To determine the accuracy of reconstructed clinical scores by BioMapAI's integration of 'omics
data, we compared their ability to discriminate ME/CFS patients from controls with the original
clinical scores. We used one additional fully connected layer to regress the 12 predicted clinical
scores $\widehat{Y}(12,)$ into a binary outcome of patient vs. control $\hat{y}(1,)$. Because the diagnosis of
ME/CFS relies on clinical interpretation of key symptoms (i.e., the original clinical scores), the
original clinical scores have near-perfect accuracy in classification as expected (AUC, Area Under
the Curve >99%, Supplemental Figure 2C). Notably, BioMapAI's predicted scores based on the
'omics data achieved a 91% AUC, highlighting its leading-edge accuracy in disease vs. healthy
classification (Figure 2D, Supplemental Figure 2D), which was also superior to the performance
of three ML models - linear regression (LR), support vector machine (SVM), and gradient
boosting (GDBT) - and one deep learning model (DNN) without the hidden 3, 'spread out' layer
(Supplemental Table 3). BioMapAI particularly excelled utilizing immune features (AUC = 80%),
KEGG genes (78%), blood measure models (71%) and combined 'omics (91%). GDBT, however,
led in the microbial species (75%) and metabolome (74%) models, likely due to its emphasis on
specific features.

Finally, to assess the robustness of our BioMapAI model, we validated it with independent,
published ME/CFS cohorts (Figure 2E, Supplemental Table 4). Using data from two microbiome
cohorts, Guo, Cheng et al., 2023 (US)[28] and Raijmakers, Ruud et al., 2020 (Netherlands)[29],
BioMapAI achieved 72% and 63% accuracy in species relative abundance and 58% and 60%
accuracy in microbial KEGG gene abundance. When applied to two metabolome cohorts,
Germain, Arnaud et al., 2022 (US)[30] and Che, Xiaoyu et al., 2022 (US)[31], BioMapAI attained 68%
and 59% accuracy. These results were strong given that the metabolomic features only overlap
by 79% and 19%, respectively, due to methodological variations.

Importantly, BioMapAI significantly surpassed GDBT and DNN in external cohort validation,
supporting our theory that while commonly used models, such as tree-based GDBT, may be
effective within a single study, their overemphasis on specific key features can limit its
generalizability across different studies, which may not share the same biomarkers. BioMapAI's
effectiveness also highlighted the value of incorporating clinical symptoms into a predictive
model, proving that connecting 'omics features to clinical symptoms improves disease
classification. Given the limitations of using external cohorts – which often have significant
methodological differences and cohort characteristics – to validate traditional microbiome and
metabolite ML models[32,33,34], BioMapAI represents a breakthrough as a far more adaptable and
broadly applicable model.

220
'Omics' Strengths Varied in Symptom Prediction; Immune is the Most Predictive
A major innovation of BioMapAI is its ability to leverage different 'omics data to predict
individual clinical scores in addition to disease vs. healthy classification. We evaluated the
predictive accuracy by calculating the mean squared error between actual (y) and predicted ($\hat{y}$)
scores and observed that the different 'omics showed varying strengths in predicting clinical
scores (Figure 2C). Immune profiling consistently excelled in forecasting a wide range of
symptoms, including pain, fatigue, orthostatic intolerance, and general health perception,
underscoring the immune system's crucial role in health regulation. In contrast, blood
measurements demonstrated limited predictive ability, except for cognitive efficiency, likely
owing to their limited focus on 48 specific blood bioactives. Plasma metabolomics, which
encompasses nearly a thousand measurements, performed significantly better with notable
correlations with facets of physical health and social activity. These findings corroborate
published metabolites and mortality[35],[36], longevity[37],[38], cognitive function[39], and social
interactions[40],[41],[42]. Microbiome profiles surpassed other 'omics in predicting gastrointestinal
abnormalities (as expected[43],[44]), emotional well-being, and sleep problems, supporting recently
established links in gut-brain health[45],[46],[47].
237
BioMapAI is Explainable, Identifying Disease- and Symptom-Specific Biomarkers
Deep learning (DL) models are often referred to as 'black box', with limited ability to identify
and evaluate specific features that influence the model's predictions. BioMapAI is made
explainable by incorporating SHAP values, which quantify how each feature influenced the
model's predictions. BioMapAI's architecture – two shared layers (Z^1 and Z^2) for general
disease pattern learning and one parallel layer for each clinical score ($Z^3 = [z_1^3, z_2^3, ..., z_{12}^3]$) –
allowed us to identify both disease-specific biomarkers, which are shared across symptoms and
models (Supplemental Figure 3, Supplemental Table 5), and symptom-specific biomarkers,
which are tailored to each clinical symptom (Figure 3, Supplemental Figure 4-5, Supplemental
Table 6).
248
Disease-specific biomarkers are important features across symptoms and models (Methods,
Supplemental Figure 3). Increased B cells (CD19+CD3-), CCR6+ CD8 memory T cells
(mCD8+CCR6+CXCR3-), and CD4 naïve T cells (nCD4+FOXP3+) in patients were pivotal for most
symptoms, indicating a systemic dysregulation of the adaptive immune response. The species
model highlighted the importance of *Dysosmobacteria welbionis*, a gut microbe previously
reported in obesity and diabetes, with a critical role in bile acid and butyrate metabolism[48],[49].
The metabolome model categorized increased levels of glycodeoxycholate 3-sulfate, a bile acid,
and decreased vanillylmandelate (VMA), a catecholamine breakdown product[50]. These critical
features for all symptoms were consistently validated across ML and DL models, demonstrating
the efficacy of BioMapAI (Supplemental Table 5).
259
More uniquely, BioMapAI linked 'omics profiles to clinical symptoms and thus enabled the
identification of symptom-specific biomarkers (Figure 3A). Certain 'omics data, like species-
gastrointestinal and immune-pain associations, were especially effective in predicting specific
clinical phenotypes (Figure 2C). Utilizing SHAP, BioMapAI identified distinct sets of biomarkers

for each symptom (Supplemental Table 6, Supplemental Figure 5). We found that while disease-
specific biomarkers accounted for a substantial portion of the variance, symptom-specific
biomarkers crucially refined the predictions, aligned predicted scores – consistently across age
and gender – more closely with actual values (Figure 3A-B, Supplemental Figure 4B-D). For
example, in the case of pain, CD4 memory and CD1c+ dendritic cells (DC) were particularly
important features, and *Faecalibacterium prausnitzii* was uniquely linked as well with varying
impact across individual (Figure 3B). Similar to pain, each clinical score in ME/CFS was
characterized by its unique 'omics features, distinct from those common across other
symptoms (Supplemental Table 6).
273
In addition, we observed a spectrum of interaction types (linear, biphasic, and dispersed)
extending beyond conventional linear interactions, underscoring the heterogeneity inherent in
ME/CFS (Figure 3C). High-abundance species and immune cells often had a biphasic relationship
with symptoms, showing dual effects, while low-abundance species and metabolites displayed
a linear relationship with positive or negative associations with clinical scores (Supplemental
Figure 5).
280
An example of a relatively straightforward monotonic (linear) relationship was observed
between CD4 memory (CD4 M) cells, CD1c+ DCs and pain, with positive contributions of CD4 M
cells to pain intensity severity. Conversely, CD1c+ DCs contributed negatively to pain severity in
both patients and control (Figure 3C, E). These variations suggest alterations in inflammatory
responses and specific pathogenic processes in ME/CFS, which may be virally triggered and is
marked by prolonged infection symptoms. Many microbial biomarkers demonstrated linear
contributions to symptoms, evidenced by numerous negative peaks indicating their beneficial
role in symptom reduction (Figure 3A). For example, *Dysosmobacteria welbionis*, a disease-
specific biomarker, exacerbated sleeping and gastrointestinal issues (Supplemental Figure 3),
whereas *Clostridium sp.* and *Alistipes communis* alleviated these issues (Figure 3A,
Supplemental Figure 5B).
292
A more complex, biphasic relationship was observed in the interaction of *Faecalibacterium*
prausnitzii with pain, whose saddle curve (Figure 3C) and mixture of positive and negative
contribution peaks (Figure 3B) revealed how abnormal low and high abundances could be
associated with amplified pain. In disease, *F. prausnitzii* was associated with exacerbated pain,
while in healthy individuals, it appeared to mitigate pain (Figure 3D). *F. prausnitzii* was
identified as a biomarker in several ME/CFS cohorts[28],[29],[51], but also has been implicated in
numerous anti-inflammatory effects[52],[53],[54],[55]. Here notably, BioMapAI elaborated its role at
ME/CFS by recognizing its potential dual contribution to symptom severity. Similar biphasic
relationships were observed for plasma metabolomics biomarkers, glucuronide and glutamine,
in relation to pain (Figure 3C).
303
Distinct from other 'omics features, KEGG genes exhibited sparse and dispersed contributions
(Figure 3C, Supplemental Figure 4C). The vast feature matrix of KEGG models complicated the
identification of a universal biomarker for any single symptom, as individuals possessed distinct
symptom-specific KEGG biomarkers. For example, the gene FNR, an anaerobic regulatory

protein transcription factor, negatively impacted pain but was active in only a small portion of
patients, with the majority showing no significant impact (Figure 3C). This pattern was
consistent for other KEGG biomarkers, which contributed sparsely to symptom severity
(Supplemental Figures 4C).
312
313 Taken together, BioMapAI achieved a comprehensive mapping of the intricate nature of
314 symptom-specific biomarkers to clinical phenotypes that has been inaccessible to single models
315 to date. Our models unveil a nuanced and precise correlation between 'omics features and
316 disease symptomology, emphasizing ME/CFS' complex etiology and consequent disease
317 management approaches.
318
Healthy Microbiome-Immune-Metabolome Networks are Dysbiotic in ME/CFS
BioMapAI elucidated that each 'omics layer provided distinct insights into the disease symptoms
and influenced host phenotypes in a dynamic and complex manner. To examine crosstalk
between 'omics layers, we modeled co-expression modules for each 'omics using weighted gene
co-expression network analysis (WGCNA), identifying seven microbial species, six microbial gene
set, nine metabolome, and nine immune clusters (Methods, Supplemental Table 7). Observing
significant associations of these modules with disease classification (microbial modules), age
and gender (immune and metabolome modules) (Supplemental Figure 6A), we first established
baseline networks of inter-'omics interactions in healthy individuals as a function of these and
other clinical covariates such as age, weight, and gender (Figure 4A), and then examined how
these interactions were altered in patient populations (Figure 4B, Supplemental Figure 6B-C).
330
Healthy control-derived host-microbiome interactions, such as the microbial pyruvate module
interacting with multiple immune modules, and connections between commensal gut microbes
(*Prevotella*, *Clostridia* sp., *Ruminococcaceae*) with Th17 memory cells, plasma steroids,
phospholipids, and tocopherol (vitamin E) (Figure 4A), were disrupted in ME/CFS patients.
Increased interactions between gut microbiome and mucosal/inflammatory immune modules,
including CD8+ MAIT, and INFg+ CD4 memory cells, suggested a microbiome-mediated
intensified inflammatory in ME/CFS (Supplemental Figure 6D). Young, female, and normal-
weight patients shared those changes, while male patients showed more distinct alterations in
the interplay between microbial and plasma metabolites. Elderly and overweight patients had
more interaction abnormalities than other subgroups, with specific increases between *Blautia*,
Flavonifractor, *Firmicutes* sp. linked with TNFα cytotoxic T cells and plasma plasmalogen, and
decreased interactions between *Lachnospiraceae* sp. with Th17 cells (Figure 4B).
343
Further examining the pyruvate hub as well as several other key microbial modules whose
networks were dysbiotic in patients, we mapped the interactions of their metabolic
subpathways to plasma metabolites and immune cells and detailed the collective contributions
to host phenotypes (Figure 4C, Supplemental Table 8). We further validated these findings with
two independent cohorts (Guo 2023[28] and Raijmakers 2020[29]). For example, increased
tryptophan metabolism, linked to gastrointestinal issues, lost its inhibitory effect on Th22 cells,
and gained interactions with γδ T cells and the secretion of INFg and GzA from CD8 and CD8+
MAIT cells. Several networks linked with emotional dysregulation and fatigue – again

underscoring the gut-brain axis[47] – differed significantly in patients vs. controls, including
decreased butyrate production - especially from the pyruvate[56] and glutarate[57] sub-pathways-
and branched-chain amino acid (BCAA) biosynthesis, which lost or reversed their interactions
with Th17, Treg cells, and plasma lipids while gaining interactions with inflammatory immune
cells including γδ T and CD8+ MAIT cells in patients; and increased microbial benzoate,
synthesized by *Clostridia* sp.[58,59] then converted to hippurate in the liver[60,61], showed a strong
positive correlation with plasma hippurate in long-term ME/CFS patients, supporting enhanced
pathway activity in later stages of the disease. This change altered its interactions with
numerous plasma metabolites, including steroids, phenols, BCAAs, fatty acids, and vitamins B5
and B6. Finally, we noted that connections of short-term patients often resembled a transitional
phase, with dysbiotic health-associated networks and emergent pathological connections that
solidified in long-term ME/CFS patients.

Based on BioMapAI's outputs and network analyses, we propose that the shift in disease
pathology in ME/CFS is linked to the topological interaction of the gut microbiome, immune
function, and metabolome. (Figure 5). A decrease in key microbes, including *Faecalibacterium*
prausnitzii, and resultant dysfunction of microbial metabolic pathways such as butyrate,
tryptophan, and BCAA, contributed to critical ME/CFS phenotypes, particularly pain and
gastrointestinal abnormalities. In healthy individuals, these microbial metabolites regulate
mucosal immune cells, including Th17, Th22, and Treg cells, an interaction that is dysfunctional
in ME/CFS resulting in elevated pro-inflammatory interactions via elevated activation of γδ T
cells and CD8 MAIT cells with the secretion of INFg and GzA, particularly impacting health
perception and social activities. Additional health-associated networks between gut microbial
metabolites, particularly benzoate, with plasma metabolites such as lipids, GPE, fatty acids, and
bile acids, were weakened or reversed in ME/CFS. This breakdown in the host metabolic-
microbiome balance were collectively associated with fatigue, emotional and sleeping
problems, supporting recent findings underscoring microbial mechanisms in the gut-brain axis
that occur via modulation of plasma metabolites[62,63,64].

Discussion
Democratization of AI technologies and large-scale multi-'omics has the promise of
revolutionizing precision medicine[65,66,67,68]. This study generated among the richest, most
extensive paired multi-'omics dataset to date[4,28,29,30,31,69,70,71], with new insights not only into
ME/CFS, but potential other applications to heterogeneous and complicated diseases like
fibromyalgia[72] and long COVID[73]. BioMapAI marks the first AI trained to systematically decode
these complex, multi-system symptoms. Traditionally, diagnosing ME/CFS has been challenging,
often relying heavily on self-reported questionnaires[74,75]. However, the crux for long-term post-
viral infection syndromes like ME/CFS is not necessarily pinpointing an exact diagnosis or tracing
disease origins[76,14] (typically infections[77]), but rather addressing the chronic, multifaceted
symptoms that significantly impacts patients' quality of life[78,79]. Our study introduces a highly
nuanced approach to link physiological changes in gut microbiome, plasma metabolome, and
immune status, with host symptoms, moving beyond the initial causes of the disease[80,81].
Importantly, we validated key biomarkers in external cohorts[28,29,30,31], despite significant
demographic and methodological differences between the studies.

In addition, by integrating these datatypes, we constructed complex new host-microbiome
networks contrasted in health vs. ME/CFS. Networks constructed in healthy individuals revealed
unique microbe-immune-metabolome connections and set a baseline for comparing numerous
disease conditions while, critically, accounting for cohort covariates, including age, gender, and
weight, as these factors reshape these networks by differing degrees, just as comorbid
conditions like aging or obesity can complicate and individualize disease profiles. This approach
enhanced the reliability of our findings in ME/CFS by rigorously accounting for potential
confounders and solidified our proposed mechanisms exclusively to the disease itself[82],[83]. For
example, gut microbiome abnormalities were most relevant to ME/CFS, while changes in
immune profiles and plasma metabolome were significant but influenced by factors like age and
gender. Symptomatologically, the gut microbiome was expectedly linked to gastrointestinal
issues and unexpectedly, to pain, fatigue, and mental health problems, possibly due to
disruptions in the gut-brain axis from abnormal microbial metabolic functions, such as lost
network connections with key plasma metabolites, particularly lipids. We previously noted
immune abnormalities in ME/CFS[84]; in this study, we further analyzed activation of mucosal and
inflammatory immunity, namely MAIT and γδ T cells, which linked to dysbiosis in gut microbial
functions. These nuanced insights, while still premature for actual treatment applications, lay
the groundwork for more precise controlled experiments and interventional studies. For
instance, personalized treatment options could include supplementation of butyrate and amino
acids for patients suffering from severe gastrointestinal and emotional symptoms, or targeted
treatments for chronic inflammation for those experiencing significant pain and fatigue.

Taken together, our results underscore BioMapAI's particular suitability to complex datatypes
that collectively, better explains the phenotypic heterogeneity of diseases such as ME/CFS than
any one alone. BioMapAI's specialized deep neuron network structure with two shared general
layers and one outcome-focused parallel layer is moreover generalizable and scalable to other
cohort studies that aim to utilize 'omics data for a range of outputs (e.g., not just limited to
clinical symptoms). For instance, researchers could employ our model to link whole genome
sequencing data with blood or protein measurements. Constructed to automatically adapt to
any input matrix X and any output matrix $Y = [y_1, y_2, \ldots, y_m]$, BioMapAI defaults to parallelly
align specific layers for each output, y. Currently, the model treated all 12 studied symptoms,
$[y_1, y_2, \ldots, y_{12}]$, with equal importance due to the unclear symptom prioritization in ME/CFS[85].
We computed modules to assign different weights to symptoms to enhance diagnostic accuracy.
While this approach was not particularly effective for ME/CFS, it may be more promising for
diseases with more clearly defined symptom hierarchies[86],[87]. In such cases, adjusting the
weights of symptoms in the model's final layer could improve performance and help pinpoint
which symptoms are truly critical.

Limitations of our study include that that our study population was comprised more females
and older individuals, majorly Caucasian, though this is consistent with the epidemiology of
ME/CFS[21],[88],[89], and was from a single geographic location (Bateman Horne Center). This may
limit our findings to certain populations. In addition, previous RNA sequencing studies have
suggested mitochondrial dysfunction and altered energy metabolism in ME/CFS[90],[91],[92],[93],[94];

thus, incorporating host PBMC RNA or ATAC sequencing in future research could provide deeper
insights into regulatory changes. The typical decades-long disease progression of ME/CFS makes
it challenging for our four-year longitudinal design to capture stable temporal signals - although
separating our short-term (<4 years) and long-term (>10 years) provided valuable insights –
ideally, tracking the same patients over a longer period would likely yield more accurate
trends[95,96]. Long disease history also increases the likelihood of exposure to various diets and
medications[97], which could influence biomarker identification, particularly in metabolomics.
Finally, model-wise, BioMapAI was trained on < 500 samples with fivefold cross-validation,
which is relatively small given the complexity of the outcome matrix; expanding the training
dataset and incorporating more independent validation sets could potentially enhance its
performance and generalizability[98,99].

Main Figure

Note: Figures in Word file are screenshots to reduce file size; Original PDFs attached.

Figure 1: Cohort Summary and Heterogeneity of ME/CFS. A) Cohort Design and 'Omics Profiling. 96 healthy donors and 153 ME/CFS patients were followed over 3-4 years with yearly sampling. Clinical metadata including lifestyle and dietary surveys, blood clinical laboratory measures (N=503), gut microbiome (N=479), plasma metabolome (N=414), and immune profiles (N=489) were collected (Supplemental Table 1 and Supplemental Figure 1A). **B) Heterogeneity and Non-Linear Progression of ME/CFS in Symptom Severity and 'Omics Profiles.** Variability in symptom severity (top) and 'omics profiles (bottom) for 20 representative ME/CFS patients over 3-4 time points. For symptom severity, the 12 major clinical symptoms (x-axis) vs. severity (scaled from 0% to 100%, y-axis) is shown for each patient (each color), with lines showing average severity and shaded areas showing severity range over their timepoints. The widespread highlights the lack of consistent temporal patterns and unique symptomatology of ME/CFS (controls shown in Supplemental Figure 1C). Bottom, PCoA of integrated 'omics data with color dots matching patient timepoints in the symptom plot and grey dots representing the entire cohort. Again, the spread and overlap of the colored space reflect the diversity in 'omics signatures vs. the more consistent pattern typical of controls (Supplemental Figure 1B). **Abbreviations:** ME/CFS, Myalgic Encephalomyelitis/Chronic Fatigue Syndrome; PCoA, Principal Coordinates Analysis. **Supporting Materials:** Supplemental Table 1, Supplemental Figure 1.

472
Figure 2: BioMapAI's Model Structure and Performance. A) Structure of BioMapAI. BioMapAI
is a fully connected deep neural network comprised of an input layer (X), a normalization layer
(not shown), three sequential hidden layers (Z^1, Z^2, Z^3), and one output layer (Y). Hidden layer
1 (Z^1, 64 nodes) and hidden layer 2 (Z^2, 32 nodes), both feature a dropout ratio of 50% to
prevent overfitting (visually represented by dark and light gray nodes). Hidden layer 3 has 12
parallel sub-layers each with 8 nodes ($Z^3 = [z_1^3, z_2^3, ..., z_{12}^3]$) to learn 12 objects in the output
layer ($Y = [y_1, y_2, ..., y_{12}]$) representing key clinical symptoms of ME/CFS. **B) True vs. Predicted**
Clinical Scores highlight BioMapAI's accuracy. Three example density maps (full set,
Supplemental Figure 2A) compare the true score, y (Column 1) against BioMapAI's predictions
generated from different 'omics profiles - $\hat{y}_{immune}, \hat{y}_{species}, \hat{y}_{KEGG}, \hat{y}_{metabolome}, \hat{y}_{omics}$
(Columns 2-6). The color gradient from blue (lower density) to red (higher density) illustrates
the occurrence frequency (e.g., true scores for ~100% of healthy controls' physical health ~ 0 =
red), with dashed lines indicating key statistical percentiles (100%, 75%, 50%, 25%, and 0%).
Note that model's predicted scores a preserve differences between healthy controls and
patients for these three examples, irrespective of 'omics type. **C) 'Omics' Strengths in Symptom**
Prediction. Radar plot shows BioMapAI's performance in predicting the 12 clinical outcomes for
each 'omics datatype. Each of the 12 axes represents a clinical score output ($Y =$
$[y_1, y_2, ..., y_{12}]$), with five colors denoting the 'omics datasets used for model training. The
spread of each color along an axis reflects the normalized mean square error (MSE,
Supplemental Table 2) between the actual, y, and the predicted, $\hat{y}$, outputs, illustrating the
predictive strength or weakness of each 'omics for specific clinical scores. For instance, species
abundance predicted gastrointestinal, emotional, and sleep issues effectively, while the
immune profile was broadly accurate across most scores. **D) BioMapAI's Performance in**
Healthy vs. Disease Classification. ROC curves show BioMapAI's performance in disease
classification using each 'omics dataset separately or combined ('Omics'), with the AUC in
parentheses showing prediction accuracy (full report in Supplemental Table 3). **E) Validation of**
BioMapAI with External Cohorts. External cohorts with microbiome data (Guo et al.[28], Ruud et
al.[29]) and metabolome data (Germain et al.[30], Che et al.[32]) were used to test BioMapAI's model,

underscoring its generalizability (detailed classification matrix, Supplemental Table 4).
Abbreviations: KEGG, Kyoto Encyclopedia of Genes and Genomes; 'Omics' refers to the
combined multi-'omics matrix; MSE, Mean Square Error; ROC curve, Receiver Operating
Characteristic curve; AUC**,** Area Under the Curve; y, True Score; $\hat{y}$, Predicted Score. **Supporting**
Materials: Supplemental Tables 2-4, Supplemental Figures 1-2.

Figure 3: BioMapAI Identifies both Disease- and Symptom-Specific Biomarkers. For Symptom-Specific Biomarkers, A) Circularized Diagram of Species Model with B) Zoomed Segment for Pain. Each circular panel illustrates how the model predicts each of the 12 symptom-specific biomarkers derived from one type of 'omics data (all datatypes shown in Supplemental Figure 4). The x-axis for each panel represents an individual's values for each of the following contributors to the model's performance (from top to bottom): *1. Variance Explained by Biomarker Categories:* Gradients of dark green (100%) to white (0%) show variance explained by the model. For many biomarkers, disease-specific biomarkers account for the greatest proportion of variance, and symptom-specific biomarkers provide additional tailored explanations, with residual accounting for the remaining variance; *2. Aggregated SHAP Values* quantify the contribution of each feature to the model's predictions, with disease-specific biomarkers in grey and symptom-specific in purple. *3. Demography and Cohort Classification:* cohort (controls, white vs. patients, black); age <50 (white) vs. >50 years old (black); sex (male, white vs. female, black); *4. True vs. Predicted Scores* show BioMapAI's predictive performance at the individual sample level, with true in blue and model-predicted scores in orange; *5. Examples of Symptom-Specific Biomarkers:* Line graphs show the contribution of select symptom-specific biomarkers to the model across individuals, e.g., 5 gut species in A). In B), the three features most specific to the pain model include gut microbe *F. prausnitzii,* CD4 memory T, and DC CD1c+ cells. Peaks above 0 (middle line) indicate a positive contribution and below 0 for a negative contribution. For example, the mixed positive and negative contribution peaks of *F. prausnitzii* indicated a biphasic contribution to pain intensity. Disease-Specific Biomarkers are shown in Supplemental Figure 3. **C) Different Correlation Patterns of Biomarkers to**

Symptoms: For pain (other symptoms in Supplemental Figure 5), correlation analysis of raw
abundance (x-axis) of each biomarker with pain score (y-axis) show monotonic (e.g., CD4
memory and DC CD1c+ markers), biphasic (microbial and metabolomic markers), or sparse
(KEGG genes) contribution patterns for those features. Dots represent an individual color-coded
to SHAP value, where the color spectrum indicates negative (blue) to neutral (grey) to positive
(red) contributions to pain prediction. Superimposed trend lines with shaded error bands
represents the predicted correlation trends between biomarkers and pain intensity. Adjacent
bar plots represent the data distribution. **D-E) Examples of Pain-Specific Biomarkers'**
Contributions. SHAP waterfall plots (colors corresponding to gradient in C) illustrate the
contribution of individual features to a model's predictive output. The top 10 features for two
pairs of controls and patients are shown here, illustrating the species and the immune model
(additional examples in Supplemental Figure 4A). The contribution of each feature is shown as a
step (SHAP values provided adjacent), and the cumulative effect of all the steps provides the
final prediction value, $E[f(X)]$. Our example of *F. prausnitzii* exhibits a protective role (negative
SHAP) in controls but exacerbates pain (positive SHAP) in patients – consistent with the biphasic
relationship observed in C). As a second example, all CD4 memory cells in this model have
positive SHAP values, reinforcing the positive monotonic relationship with pain severity
observed in C). Conversely, DC CD1c+ cells contribute negatively and thus may have a
protective role. **Abbreviation:** SHAP, SHapley Additive exPlanations; DNN, Deep Neuron
Network; GBDT, Gradient Boosting Decision Tree; KEGG, Kyoto Encyclopedia of Genes and
Genomes. **Supporting Materials:** Supplemental Table 5-6, Supplemental Figure 3-5.

Figure 4: Microbiome-Immune-Metabolome Crosstalk is Dysbiotic in ME/CFS. A-B)
Microbiome-Immune-Metabolome Network in A) Healthy and B) Patient Subgroups. A
baseline network was established with 200+ healthy control samples (**A**), bifurcating into two
segments: the gut microbiome (species in yellow, genetic modules in orange) and blood

elements (immune modules in green, metabolome modules in purple). Nodes: modules; size: #
of members; colors: 'omics type; edges: interactions between modules, with Spearman
coefficient (adjusted) represented by thickness, transparency, and color - positive (red) and
negative (blue). Here, key microbial pathways (pyruvate, amino acid, and benzoate) interact
with immune and metabolome modules in healthy individuals. Specifically, these correlations
were disrupted in patient subgroups (**B**), as a function of gender, age (young <26 years old vs.
older >50), BMI (normal <26 vs. overweight >26), and health status (individuals with IBS or
infections). Correlations significantly shifted from healthy counterparts (Supplemental Figure
6C) are highlighted with colored nodes and edges indicating increased (red) or decreased (blue)
interactions. **C) Targeted Microbial Pathways and Host Interactions.** Four important microbial
metabolic mechanisms (tryptophan, butyrate, BCAA, benzoate) were further analyzed to
compare control, short and long-term ME/CFS patients, and external cohorts for validation
(Guo[28] and Raijmakers[29]).*1. Microbial Pathway Fold Change:* Key genes were grouped and
annotated in subpathways. Circle size: fold change over control; color: increase (red) or
decrease (blue), p-values (adjusted Wilcoxon) marked. *2. Microbiome-Host Interactions:* Sankey
diagrams visualize interactions between microbial pathways and host immune
cells/metabolites. Line thickness and transparency: Spearman coefficient (adjusted); color: red
(positive), blue (negative). *3. Immune & Metabolites Fold Change:* Pathway-correlated immune
cells and metabolites are grouped by category. *4. Contribution to Disease Symptoms:* Stacked
bar plots show accumulated SHAP values (contributions to symptom severity) for each disease
symptom (1-12, as in Supplemental Table 1). Colors: microbial subpathways and
immune/metabolome categories match module color in fold change maps. X-axis: accumulated
SHAP values (contributions) from negative to positive, with the most contributed symptoms
highlighted. **P-values:** *p < 0.05, **p < 0.01, ***p < 0.001. **Abbreviations:** IBS, Irritable Bowel
Syndrome; BMI, Body Mass Index; BCAA, Branched-Chain Amino Acids; MAIT, Mucosal-
Associated Invariant T cell; SHAP, SHapley Additive exPlanations; GPE,
Glycerophosphoethanolamine; INFγ, Interferon Gamma; CD, Cluster of Differentiation; Th, T
helper cell; TMAO, Trimethylamine N-oxide; KEGG, Kyoto Encyclopedia of Genes and Genomes.
Supporting Materials: Supplemental Table 7-8, Supplemental Figure 6.

584
Figure 5: Overview of Dysbiotic Host-Microbiome Interactions in ME/CFS. This conceptual
diagram visualizes the host-microbiome interactions in healthy conditions (left) and its
disruption and transition into the disease state in ME/CFS (right). The base icons of the figure
remain consistent, while gradients and changes in color and size visually represent the
progression of the disease. Process of production and processing is represented by lines with
arrows, where the color indicates an increase (red) or decrease (blue) in the pathway in
disease; lines without arrows indicate correlations, with red representing positive and blue
representing negative correlations. In healthy conditions, microbial metabolites support
immune regulation, maintaining mucosal integrity and healthy inflammatory responses by
positively regulating Treg and Th22 cell activity, and controlling Th17 activities, including the
secretion of IL17 (purple cells), IL22 (blue), and IFNγ. These microbial metabolites also maintain
many positive interactions with plasma metabolites like lipids, bile acids, vitamins, and phenols.
In ME/CFS, there is a significant decrease in beneficial microbes and a disruption in metabolic
pathways, marked by a decrease in the butyrate (brown-red dots) and BCAA (yellow) pathways
and an increase in tryptophan (green) and benzoate (red) pathways. These changes are linked
to gastrointestinal issues. In ME/CFS, the regulatory capacity of the immune system diminishes,
leading to the loss of health-associated interactions with Th17, Th22, and Treg cells, and an
increase in inflammatory immune activity. Pathogenic immune cells, including CD8 MAIT and
γδT cells, show increased activity, along with the secretion of inflammatory cytokines such as
IFNγ and GzmA, contributing to worsened general health and social functioning. Healthy
interactions between gut microbial metabolites and plasma metabolites weaken or even
reverse in the disease state. A notable strong connection increased in ME/CFS is benzoate
transformation to hippurate, associated with emotional disturbances, sleep issues, and fatigue.
Abbreviations: IFNγ, Interferon gamma; Th17, T helper 17 cells; Th22, T helper 22 cells; Treg,
Regulatory T cells; GzmA, Granzyme A; MAIT, Mucosa-Associated Invariant T cells; γδT, Gamma
delta T cells; BCAA, Branched-Chain Amino Acids; GPE, Glycerophosphoethanolamine.

Methods

Study Design. This was 4-year prospective study. All participants had a physical examination at the baseline visit that included evaluation of vital signs, BMI, orthostatic vital signs, skin, lymphatic system, HEENT, pulmonary, cardiac, abdomen, musculoskeletal, nervous system and fibromyalgia (FM) tender points. We enrolled a total of 153 ME/CFS patients (of which 75 had been diagnosed with ME/CFS <4 years before recruitment and 78 had been diagnosed with ME/CFS >10 years before recruitment) and 96 healthy controls. Among them, 110 patients and 58 healthy controls were followed one year after the recruitment as timepoint 2; 81 patients and 13 healthy controls were followed two years after the recruitment as timepoint 3; and 4 patients were followed four years after the recruitment as timepoint 4. Subject characteristics are shown in Supplemental Table 1 and Supplemental Figure 1A.

Medical history and concomitant medications were documented. Blood samples were obtained prior to orthostatic and cognitive testing. The 10-minute NASA Lean Test and cognitive testing were conducted after the physical examination and blood draw[100]. Cognitive efficiency was tested with the DANA Brain Vital, measuring three reaction time and information processing measurements[101]. The orthostatic challenge was assessed with the 10-minute NASA Lean Test (NLT). Participants rested supine for 10 minutes, and baseline blood pressure (BP) and heart rate (HR) were measured twice during the last 2 minutes of rest[102].

Participants were provided with an at-home stool collection kit at the end of each in-person visit. The following questionnaires were completed at baseline: DePaul Symptom Questionnaire (DSQ), Post-Exertional Fatigue Questionnaire, RAND-36, Fibromyalgia Impact Questionnaire-R, ACR 2010 Fibromyalgia Criteria Symptom Questionnaire, Pittsburgh Sleep Quality Index (PSQI), Stanford Brief Activity Survey, Orthostatic Intolerance Daily Activity Scale, Orthostatic Intolerance Symptom Assessment, Brief Wellness Survey, Hours of Upright Activity (HUA), medical history and family history. All but medical history and family history were administered again when participants came for their annual visit.

Approval was received before enrolling any subjects in the study (The Jackson Laboratory Institutional Review Board, 17-JGM-13). All participants were educated about the study prior to enrollment and signed all appropriate informed consent documents. Research staff followed Good Clinical Practices (GCP) guidelines to ensure subject safety and privacy.

ME/CFS Cohort. Beginning in January 2018, we enrolled ME/CFS patients who had been sick for <4 years or sick for >10 years. No ME/CFS patients with duration ≥4 years and ≤10 years were enrolled in order to have clear distinctions between short and long duration of illness with ME/CFS. All participants were 18 to 65 years old at the time of enrollment. ME/CFS diagnosis according to the Institute of Medicine clinical diagnostic criteria and disease duration of <4 years were confirmed during clinical differential diagnosis and thorough medical work up[103]. Additional inclusion criteria required, 1) a substantial reduction or impairment in the ability to engage in pre-illness levels of occupational, educational, social, or personal activities that persists for more than 6 months and less than 4 years and is accompanied by fatigue, which is often profound, is of new or definite onset (not lifelong), is not the result of ongoing excessive

exertion, and is not substantially alleviated by rest, and 2) post-exertional malaise. Exclusionary
criteria for the <4 year ME/CFS cohort were, 1) morbid obesity BMI>40, 2) other active and
untreated disease processes that explain most of the major symptoms of fatigue, sleep
disturbance, pain, and cognitive dysfunction, 3) untreated primary sleep disorders, 4)
rheumatological disorders, 5) immune disorders, 6) neurological disorders, 7) infectious
diseases, 8) psychiatric disorders that alter perception of reality or ability to communicate
clearly or impair physical health and function, 9) laboratory testing or imaging are available that
support an alternate exclusionary diagnosis, and 10) treatment with short-term (less than 2
weeks) antiviral or antibiotic medication within the past 30 days.
For the >10 year ME/CFS cohort, disease duration of >10 year and clinical criteria was confirmed
to meet the Institute of Medicine criteria for ME/CFS during clinical evaluation and medical
history review[103]. Other than disease duration, inclusion and exclusion criteria were the same as
for <4 year ME/CFS cohort.
668
Healthy Control Cohort. Healthy control participants were also between 18 to 65 years of age
and in general good health. Enrollment began in 2018 and subjects were selected to match the
<4 year ME/CFS cohort by age (within 5 years), race, and sex (~2:1 female to male ratio).
Exclusion criteria for healthy controls included, 1) a diagnosis or history of ME/CFS, 2) morbid
obesity BMI>40, 3) treatment with short-term (less than 2 weeks) antiviral or antibiotic
medication within the past 30 days or 4) treatment long-term (longer than 2 weeks) antiviral
medication or immunomodulatory medications within the past 6 months.
676
Clinical Metadata and Scores. Clinical symptoms and baseline health status were assessed on
the day of physical examination and biological sample collection for both case and control
subjects. For each participant, we collected demographic information (including age, gender,
diet, race, BMI, family, work, and education), medical histories, clinical tests and questionnaires.
From questionnaires and test as described above, we summarized 12 clinical scores to cover
major symptoms of ME/CFS: Scores 1-8 were derived from the RAND36, following standardized
rules [104] and summarized into eight categories: Physical Functioning (also referred to as Daily
Activity in the main contents), Role Limitations due to Physical Health (Physical Limitations),
Role Limitations due to Emotional Problems (Emotional Problems), Energy/Fatigue, Emotional
Wellbeing (Mental Health), Social Functioning (Social Activity), Pain, and General Health (Health
Perception). Cognitive Efficiency was summarized from the DANA Brain Vital test, Orthostatic
Intolerance from the NLT test, Sleeping Problem Score from the Pittsburgh Sleep Quality Index
(PSQI) questionnaire, and Gastrointestinal Problems Score from the Gastrointestinal Symptom
Rating Scale (GSRS) questionnaire. Each score was transformed into a 0–1 scale to facilitate
combination and comparison, where a score of 1 indicates maximum disability or severity and a
score of 0 indicates no disability or disturbance.
693
Plasma Sample collection and Preparation. Healthy and patient blood samples were obtained
from Bateman Horne Center, Salt Lake City, UT and approved by JAX IRB. One 4 mL lavender top
tube (K2EDTA) was collected, and tube slowly inverted 8-10 times immediately after collection.
Blood was centrifuged within 30 minutes of collection at 1000 x g with low brake for 10
minutes. 250 uL of plasma was transferred into three 1 mL cryovial tubes, and tubes were

frozen upright at -80°C. Frozen plasma samples were batch shipped overnight on dry ice to The
Jackson Laboratory, Farmington, CT, and stored at -80°C. One green top tube (Heparin) was
collected, and tube slowly inverted 8-10 times immediately after collection. Heparinized blood
samples were shipped overnight at room temperature. Peripheral blood mononuclear cells
(PBMC) were isolated using Ficoll-paque plus (GE Healthcare) and cryopreserved in liquid
nitrogen.
705
Plasma untargeted metabolome by UPLC-MS/MS. Plasma samples were sent to Metabolon
platform and processed by Ultrahigh Performance Liquid Chromatography-Tandem Mass
Spectroscopy (UPLC-MS/MS) following the CFS cohort pipeline. In brief, samples were prepared
using the automated MicroLab STAR® system from Hamilton Company. The extract was divided
into five fractions: two for analysis by two separate reverse phases (RP)/UPLC-MS/MS methods
with positive ion mode electrospray ionization (ESI), one for analysis by RP/UPLC-MS/MS with
negative ion mode ESI, one for analysis by HILIC/UPLC-MS/MS with negative ion mode ESI, and
one sample was reserved for backup. QA/QC were analyzed with several types of controls were
analyzed including a pooled matrix sample generated by taking a small volume of each
experimental sample (or alternatively, use of a pool of well-characterized human plasma),
extracted water samples, and a cocktail of QC standards that were carefully chosen not to
interfere with the measurement of endogenous compounds were spiked into every analyzed
sample, allowed instrument performance monitoring, and aided chromatographic alignment.
Compounds were identified by comparison to Metabolon library entries of purified standards or
recurrent unknown entities. The output raw data included the annotations and the value of
peaks quantified using area-under-the-curve for metabolites.
722
Immune Profiling: Flow Cytometry Analysis. Frozen PBMC aliquots were thawed, counted and
divided into two parts, one part for day 0 surface staining, and the other part cultured in
complete RPMI 1640 medium (RPMI plus 10% Fetal Bovine Serum (FBS, Atlanta Biologicals) and
1% penicillin/streptomycin (Corning Cellgro) supplemented with IL-2+IL15 (20ng/ml) for Treg
subsets day 1 surface and transcription factors staining after culture with IL-7 (20ng/ml) for day
1 and day 6 intracellular cytokine staining, and a combination of cytokines (20ng/ml IL-12,
20ng/ml IL-15, and 40ng/ml IL-18) for day 1 intracellular cytokine staining (IL-12 from R&D, IL-7
and IL-15 from Biolegend). Surface staining was performed in staining buffer containing PBS +
2% FBS for 30 minutes at 4°C. When staining for chemokine receptors the incubation was done
at room temperature. Antibodies used in the surface staining are 2B4, CD1c, CD14, CD16, CD19,
CD25, CD27, CD31, CD3, CD303, CD38, CD4, CD45RO, CD56, CD8, CD95, CD161, CCR4, CCR6,
CCR7, CX3CR1, CXCR3, CXCR5, γδ TCR bio, HLA-DR, IgG, IgM, LAG3, PD-1, TIM3, Va7.2, Va24Ja18
all were obtained from Biolegend.
736
For intracellular cytokine staining, cells were stimulated with PMA (40ng/ml for overnight
cultured cells and 20ng/ml for 6 days cultured cells) and Ionomycin (500ng/ml) (both from
Sigma-Aldrich) in the presence of GolgiStop (BD Biosciences) for 4 hours at 37°C. For cytokine
secretion after stimulation with IL-12+IL-15+IL-18, GolgiStop was added to the culture on day 1
for 4 hours. For intracellular cytokine and transcription factor staining, PMA+Ionomycin
stimulated cells of unstimulated cells were collected, stained with surface markers including

CD3, CD4, CD8, CD161, PD1, 2B4, Vα7.2, CD45RO, CCR6, and CD27 followed by one wash with
PBS (Phosphate buffer Saline) and staining with fixable viability dye (eBioscience). After surface
staining, cells were fixed and permeabilized using fixation/permeabilization buffers
(eBioscience) according to the manufacturer's instruction. Permeabilized cells were then stained
for intracellular FOXP3, Helios, IL-4, IFNγ, TNFα, IL-17A, IL-22, Granzyme A, GM-CSF, and
Perforin from Biolegend. Flow cytometry analysis was performed on Cytek Aurora (Cytek
Biosciences) and analyzed using FlowJo (Tree Star).
750
Fecal Sample Collection and DNA Extraction. Stool was self-collected at home by volunteers
using a BioCollector fecal collection kit (The BioCollective, Denver, CO) according to
manufacturer instructions for preservation for sequencing prior to sending the sample in a
provided Styrofoam container with a cold pack. Upon receipt, stool and OMNIgene samples
were immediately aliquoted and frozen at –80°C for storage. Prior to aliquoting, OMNIgene
stool samples were homogenized by vortexing (using the metal bead inside the OMNIgene
tube), then divided into 2 microfuge tubes, one with 100μL aliquot and one with 1mL. DNA was
extracted using the Qiagen (Germantown, MD, USA) QIAamp 96 DNA QIAcube HT Kit with the
following modifications: enzymatic digestion with 50μg of lysozyme (Sigma, St. Louis, MO, USA)
and 5U each of lysostaphin and mutanolysin (Sigma) for 30 min at 37 °C followed by bead-
beating with 50 μg 0.1 mm of zirconium beads for 6 min on the Tissuelyzer II (Qiagen) prior to
loading onto the Qiacube HT. DNA concentration was measured using the Qubit high sensitivity
dsDNA kit (Invitrogen, Carlsbad, CA, USA).
764
Metagenomic Shotgun Sequencing. Approximately 50μL of thawed OMNIgene preserved stool
sample was added to a microfuge tube containing 350 μL Tissue and Cell lysis buffer and 100 μg
0.1 mm zirconia beads. Metagenomic DNA was extracted using the QiaAmp 96 DNA QiaCube HT
kit (Qiagen, 5331) with the following modifications: each sample was digested with 5μL of
Lysozyme (10 mg/mL, Sigma-Aldrich, L6876), 1μL Lysostaphin (5000U/mL, Sigma-Aldrich, L9043)
and 1μL oh Mutanolysin (5000U/mL, Sigma-Aldrich, M9901) were added to each sample to
digest at 37°C for 30 minutes prior to the bead-beating in the in the TissueLyser II (Qiagen) for 2
x 3 minutes at 30 Hz. Each sample was centrifuged for 1 minute at 15000 x g prior to loading
200μl into an S-block (Qiagen, 19585) Negative (environmental) controls and positive (in-house
mock community of 26 unique species) controls were extracted and sequenced with each
extraction and library preparation batch to ensure sample integrity. Pooled libraries were
sequenced over 13 sequencing runs using both HiSeq (N=87) and NovaSeq (N=392) platforms.
To address potential biases arising from varying read depths, all samples were down-sampled,
using seqtk[108] (v1.3-r106), to 5 million reads. This threshold corresponds to the 95th percentile
of the read count distribution across the dataset.
780
Sequencing adapters and low-quality bases were removed from the metagenomic reads using
scythe (v0.994) and sickle (v1.33), respectively, with default parameters. Host reads were
removed by mapping all sequencing reads to the hg19 human reference genome using Bowtie2
(v2.3.1), under 'very-sensitive' mode. Unmapped reads (i.e., microbial reads) were used to
estimate the relative abundance profiles of the microbial species in the samples using
MetaPhlAn4.

Taxonomic Profiling (Specie Abundance) and KEGG Gene Profiling. Taxonomic compositions
were profiled using Metaphlan4.0[105] and the species whose average relative abundance > 1e-4
were kept for further analysis, giving 384 species. The gene profiling was computed with
USEARCH[106] (v8.0.15) (with parameters: evalue 1e-9, accel 0.5, top_hits_only) to KEGG
Orthology (KO) database v54, giving a total of 9452 annotated KEGG genes. The reads count
profile was normalized by DeSeq2[107] in R. Genes with a prevalence of over 20% were selected
for downstream analysis.

Confounder Analysis. Confounder analysis was done by R package MaAsLin2[109]. We considered
demographic features (including age, gender, BMI, ethnicity, and race), diet records,
medications (antivirals, antifungals, antibiotics, and probiotics), and self-reported IBS scores as
potential confounders. The analysis followed the model formula:

$$expr \sim \text{ age} + \text{gender} + \text{bmi} + \text{ethnic} + \text{race} + \text{IBS} + \text{diet\_meat} + \text{diet\_sugar} + \text{diet\_veg}$$
$$+ \text{diet\_grains} + \text{diet\_fruit} + \text{antifungals} + \text{antibiotics} + \text{probiotics}$$
$$+ \text{antivirals} + (1|\text{sample\_id\_tp1})$$

where $expr$ refers to the 'omics matrix. For each feature in the 'omics data, we ran this
generalized linear model to identify multivariable associations between each 'omics feature and
each metadata feature. Identified confounders were handled differently based on the type of
data. For species and KEGG genes, any feature with a significant statistical association with any
metadata feature was removed from all subsequent analyses, resulting in the removal of 21
species and 946 microbial genes. For immune profiling and plasma metabolomics, to remove
the effects of identified confounders, each feature was adjusted by retaining the residuals[105],
i.e., the part of the outcome not explained by the confounding factors, from a general linear
model:

$$y' = (y \sim \text{predicted confounders})\$residual$$

Additionally, for network and patient subset analysis (Methods), age, gender, BMI, and IBS were
not included as confounders since we analyzed different age groups, gender groups, weight
groups, and IBS groups separately. However, other identified confounders were still considered
in the residual models.

BioMapAI. The primary goal of BioMapAI is to connect high-dimensional biology data, X to
mixed-type output matrix, Y. Unlike traditional ML or DL classifiers that typically predict a single
outcome, y, BioMapAI is designed to learn multiple objects, $Y = [y_1, y_2, ..., y_n]$, simultaneously
within a single model. This approach allows for the simultaneous prediction of diverse clinical
outcomes - including binary, categorical, continuous variables - with 'omics profiles, thus
address disease heterogeneity by tailoring each patient's specific symptomology.
1. BioMapAI Structure. BioMapAI is a fully connected deep neural network framework
comprising an input layer X, a normalization layer, three sequential hidden layers, Z^1, Z^2, Z^3,and
one output layer Y.
1) Input layer (X) takes high-dimensional 'omics data, such as gene expression, species
abundance, metabolome matrix, or any customized matrix like immune profiling and blood labs.
2) Normalization Layer standardizes the input features to have zero mean and unit variance,
defined as

831
$$X' = \frac{X - \mu}{\sigma}$$

where μ is the mean and σ is the standard deviation of the input features.

3) Feature Learning Module is the core of BioMapAI, responsible for extracting and learning

important patterns from input data. Each fully connected layer (hidden layer 1-3) is designed to

capture complex interactions between features. **Hidden Layer 1 (Z^1)** and **Hidden Layer 2 (Z^2)**

contain 64 and 32 nodes, respectively, both with ReLU activation and a 50% dropout rate,

defined as:

838
$$Z^k = \text{ReLU}(W^k Z^{k-1} + b^k), \qquad k \in \{1,2\}$$

839 **Hidden Layer 3 (Z^3)** has n parallel sub-layers for each object, y_i in Y. Every sub-layer, Z_i^3,

840 contains 8 nodes, represented as:

841
$$Z_i^3 = \text{ReLU}(W_i^3 Z^3 + b_i^3), \qquad i \in \{1,2,\dots,n\}$$

842 All hidden layers used ReLU activation functions, defined as:

843
$$\text{ReLU}(x) = max(0, x)$$

844 **4) Outcome Prediction Module** is responsible for the final prediction of the objects. **The output**

845 **layer (Y)** has n nodes, each representing a different object:

846
$$y_i = \begin{cases} \sigma(W_i^4 Z_i^3 + b_i^4) & \text{for binary object} \\ \text{softmax}(W_i^4 Z_i^3 + b_i^4) & \text{for categorical object} \\ W_i^4 Z_i^3 + b_i^4 & \text{for continuous object} \end{cases}$$

847 The loss functions are dynamically assigned based on the type of each object:

848
$$\mathcal{L} = \begin{cases} \frac{1}{N} \sum_{i=1}^{N} [y_i \log(\hat{y}_i) + (1 - y_i) \log(1 - \hat{y}_i)] & \text{for binary object} \\ -\frac{1}{N} \sum_{i=1}^{N} \sum_{j=1}^{C} y_{ij} \log(\hat{y}_{ij}) & \text{for categorical object} \\ \frac{1}{N} \sum_{i=1}^{N} \begin{cases} 0.5(y_i - \hat{y}_i)^2, & \text{if } |y_i - \hat{y}_i| \leq \delta \\ \delta |y_i - \hat{y}_i| - 0.5\delta^2, & otherwise \end{cases} & \text{for continuous object} \end{cases}$$

During training, the weights are adjusted using the Adam optimizer. The learning rate was set to

0.01, and weights were initialized using the He normal initializer. L2 regularizations were applied

to prevent overfitting.

5) Optional Binary Classification Layer (not used for parameter training)**.** An additional binary

classification layer is attached to the output layer Y to evaluate the model's performance in

binary classification tasks. This layer is not used for training BioMapAI but serves as an auxiliary

component to assess the accuracy of predicting binary outcomes, for example, disease vs.

control. This ScoreLayer takes the predicted scores from the output layer and performs binary

classification:

859
$$y_{binary} = \sigma(W_{binary} Y + b_{binary})$$

860 The initial weights of the 12 scores are derived from the original clinical data, and the weights

861 are adjusted based on the accuracy of BioMapAI's predictions:

862
$$w_{new} = w_{old} - \eta \nabla \mathcal{L}_{MSE}$$

863 where $\nabla \mathcal{L}_{MSE}$ refers to the mean squared error (MSE) between the predicted y' and true y,

864 then adjusts the weights to optimize the accuracy of the binary classification.

2. Training and Evaluation of BioMapAI for ME/CFS – BioMapAI::DeepMECFS. BioMapAI is a framework designed to connect high-dimensional, sparse biological 'omics matrix X to multi-output Y. While BioMapAI is not tailored to a specific disease, it is versatile and applicable to a broad range of biomedical topics. In this study, we trained and validated BioMapAI using our ME/CFS datasets. The trained models are available on GitHub, nicknamed DeepMECFS, for the benefit of the ME/CFS research community.

1) Dataset Pre-Processing Module: Handling Sample Imbalance. To ensure uniform learning for each output y, it is crucial to address sample imbalance before fitting the framework. We recommend using customized sample imbalance handling methods, such as Synthetic Minority Over-sampling Technique (SMOTE)[110], Adaptive Synthetic (ADASYN)[111], or Random Under-Sampling (RUS)[112]. In our ME/CFS dataset, there is a significant imbalance, with the patient data being twice the size of the control data. To effectively manage this class imbalance, we employed RUS as a random sampling method for the majority class. Specifically, we randomly sampled the majority class 100 times. For each iteration i, a different random subset $S_i^{majority}$ was used. This subset $S_i^{majority}$ of the majority class was combined with the entire minority class $S^{minority}$. For each iteration i:

$$S_i^{\text{majority}} \subseteq S^{majortiy}, \qquad S^{minority} = S^{minority}$$
$$S_i = S_i^{majority} \cup S^{minority}$$

where the combined dataset S_i was used for training at each iteration. This approach allows the model to generalize better and avoid biases towards the majority class, improving overall performance and robustness.

2) Cross-Validation and Model Training. DeepMECFS is the name of the trained BioMapAI model with ME/CFS datasets. We trained on five preprocessed 'omics datasets, including species abundances (Feature N=118, Sample N=474) and KEGG gene abundances (Feature N=3959, Sample N=474) from the microbiome, plasma metabolome (Feature N=730, Sample N=407), immune profiling (Feature N=311, Sample N=481), and blood measurements (Feature N=48, Sample N=495). Additionally, an integrated 'omics profile was created by merging the most predictive features from each 'omics model related to each clinical score (SHAP Methods), forming a comprehensive matrix of 154 features, comprising 50 immune features, 32 species, 30 KEGG genes, and 42 plasma metabolites.

To evaluate the performance of BioMapAI, we employed a robust 5-fold cross-validation. Training was conducted over 500 epochs with a batch size of 64 and a learning rate of 0.0005, optimized through grid search. The Adam optimizer was used to adjust the weights during training, chosen for its ability to handle sparse gradients on noisy data. The initial learning rate was set to 0.01, with beta1 set to 0.9, beta2 set to 0.999, and epsilon set to 1e-7 to ensure numerical stability. Dropout layers with a 50% dropout rate were used after each hidden layer to prevent overfitting, and L2 regularization ($\lambda = 0.008$) was applied to the kernel weights, defined as:

$$L_{reg} = \frac{\lambda}{2} \sum_{i=1}^{N} w_i^2$$

3) Model Evaluation. To evaluate the performance of the models, we employed several metrics tailored to both regression and classification tasks. The Mean Squared Error (MSE) was used to

906 evaluate the performance of the reconstruction of each object. For each y_i, MSE was calculated
907 as:

$$MSE_i = \frac{1}{N}\sum_{j=1}^{N}\left(y_i^j - \hat{y}_i^j\right)^2, i = 1,2,\dots,n$$

where y_i^j is the actual values, $\hat{y}_i^j$ is the predicted values, and N is the number of samples, n is
the number of objects. For binary classification tasks (ME/CFS vs control), we utilized multiple
metrics including accuracy, precision, recall, and F1 score to enable a comprehensive evaluation
of the model's performance.
To evaluate the performance of BioMapAI, we compared its binary classification performance
with three traditional machine learning models and one deep neural network (DNN) model. The
traditional machine learning models included: 1) Logistic Regression (**LR**) (C=0.5, saga solver
with Elastic Net regularization); 2) Support Vector Machine (**SVM**) with an RBF kernel (C=2); and
3) Gradient Boosting Decision Trees (**GBDT**) (learning rate = 0.05, maximum depth = 5,
estimators = 1000). **DNN** model employed the same hyperparameters as BioMapAI, except it did
not include the parallel sub-layer, Z_3, thus it only performed binary classification instead of
multi-output predictions. The comparison between BioMapAI and DNN aims to assess the
specific contribution of the spread-out layer, designed for discerning object-specific patterns, in
binary prediction. Evaluation metrics are detailed in Supplemental Table 3.
4) External Validation with Independent Dataset. To validate BioMapAI's robustness in binary
classification, we utilized 4 external cohorts[28],[29],[30],[31] comprising more than 100 samples. For
these external cohorts, only binary classification is available. A detailed summary of data
collection for these cohorts is provided in Supplemental Table 4. For each external cohort, we
processed the raw data (if available) using our in-house pipeline. The features in the external
datasets were aligned to match those used in BioMapAI by reindexing the datasets. The overlap
between the features in the external dataset and BioMapAI's feature set was calculated to
determine feature coverage. Any missing features were imputed with zeros to maintain
consistency across datasets. The input data was then standardized as BioMapAI. We loaded the
pre-trained BioMapAI, GBDT, and DNN for comparison. LR and SVM were excluded because they
did not perform well during the in-cohort training process. The performance of the models was
evaluated using the same binary classification evaluation metrics. Evaluation metrics detailed in
Supplemental Table 4.
3. BioMapAI Decode Module: SHAP. BioMapAI is designed to be explainable, ensuring that it
not only reconstructs and predicts accurately but also is interpretable, which is particularly
crucial in the biological domain. To achieve this, we incorporated SHapley Additive exPlanations
(SHAP) into our framework. SHAP offers a consistent measure of feature importance by
quantifying the contribution of each input feature to the model's output.[113]
We applied SHAP to BioMapAI to interpret the results, following these three steps:
1) Model Reconstruction. BioMapAI's architecture includes two shared hidden layers - Z^1, Z^2-
and one parallel sub-layers - Z_i^3- for each object y_i. To decode the feature contributions for each
object y_i, we reconstructed sub-models from single comprehensive model:

$$Model_i = Z^1 + Z^2 + Z_i^3, i = 1,2,\dots,n$$

946 where n is the number of learned objects.

2) SHAP Kernel Explainer. For each reconstructed model, $Model_i$, we used the SHAP Kernel Explainer to compute the feature contributions. The explainer was initialized with the model's prediction function and the input data X:

$$explainer_i = shap.KernelExplainer(Model_i.predict, X), i = 1,2,...,n$$

Then SHAP values were computed to determine the contribution of each feature to y_i:

$$\phi_i = explainer_i(X), i = 1,2,...,n$$

The kernel explainer is a model-agnostic approach that approximates SHAP by evaluating the model with and without the feature of interest and then assigning weights to these evaluations to ensure fairness. For each $model_i$, with each feature j:

$$\phi_i^j(f,x) = \sum_{S_i \subseteq N_i \setminus \{j\}} \frac{|S_i|!\,(m - |S_i| - 1)!}{m!} \left(Model_i(S_i \cup j) - Model_i(S_i)\right)$$

$$= \frac{1}{m} \sum_{S_i \subseteq N_i \setminus \{j\}} \binom{m-1}{m - |S_i| - 1}^{-1} \left(Model_i(S_i \cup j) - Model_i(S_i)\right), i = 1,2,...,n$$

where n is the number of learned objects, m is the total number of features, ϕ_i^j is the Shapley value for feature j in $model_i$, N_i is the full set of features in $model_i$, S_i is the subset of features not including feature j, $Model_i(S_i)$ is the model prediction for the subset S_i. The SHAP value matrix, ϕ_i, were further reshaped to align with the input data dimensions.

3) Feature Categorization. Analyzing the SHAP value matrices, $[\phi_1, \phi_2, ..., \phi_n]$, features can be roughly assigned to two categories: shared features - important to all outputs; or specific features - specifically important to individual outputs. We set the cutoff at 75%, where features consistently identified as top contributors in 75% of the models were classified as shared important features, termed disease-specific biomarkers. Features that were top contributors in only a few models were classified as specific important features, termed symptom-specific biomarkers.

By reconstructing individual models, $Model_i$, for each object, y_i, and applying SHAP explainer individually, we effectively decoded the contributions of input features to BioMapAI's predictions. This method allowed us to categorize features into shared and specific categories—termed as disease-specific and symptom-specific biomarkers—providing novel interpretations of the 'omics feature contribution to clinical symptoms.

4. Packages and Tools. BioMapAI was constructed by Tensorflow(v2.12.0)[114] and Keras(v2.12.0). ML models were from scikit-learn(v 1.1.2)[115].

WGCNA and Network Analysis. To identify co-expressed patterns of each 'omics, we employed the Weighted Gene Co-expression Network Analysis (WGCNA) using the WGCNA[116] package in R. The analysis was performed on preprocessed omics data (Methods): species abundances (Feature N=373, Sample N=479) and KEGG gene abundances (Feature N=4462, Sample N=479) from the microbiome, plasma metabolome (Feature N=395, Sample N=414), immune profiling (Feature N=311, Sample N=489). Network construction and module detection involved choosing soft-thresholding powers tailored to each dataset: 6 for species, 7 for KEGG, 5 for immune, and 6 for metabolomic. The adjacency matrices were transformed into topological overlap matrices (TOM) to reduce noise and spurious associations. Hierarchical clustering was performed using the TOM, and modules were identified using the dynamic tree cut method with a minimum

module size of 30 genes. Module eigengenes were calculated, and modules with highly similar
eigengenes (correlation > 0.75) were merged. Module-trait relationships were assessed by
correlating module eigengenes with clinical traits, and gene significance (GS) and module
membership (MM) were used to identify hub genes within significant modules.
Network analysis was conducted using igraph[117] in R. Module eigengenes from the WGCNA
analysis were extracted for each dataset. A combined network was constructed by calculating
Spearman correlation coefficients (corrected, Methods) between the module eigengenes of
different datasets, and an adjacency matrix was created based on a threshold of 0.3 (absolute
value) to include only significant associations. Network nodes represented module eigengenes
and edges represented significant correlations. Degree centrality and betweenness centrality
were calculated to identify highly connected and influential nodes. Networks in patient
subgroups were displayed as the correlation differences from their healthy counterparts to
exclude the influence of covariates. For example, correlations in female patients were compared
with female healthy, and correlations in older patients were compared with older healthy.
1001
Statistical Analysis. The dimensionality reduction analysis was conducted by Principal
Correspondence Analysis (PCoA) using sklearn.manifold.MDS function for 'omics. For combined
'omics data, PCoA was applied to combined module eigengenes from WGCNA. Fold change of
species, genes, immune cells, and metabolites were compared between patient and control
groups, short-term and control groups, and long-term and control groups. P values were
computed by Wilcoxon signed-rank test with False Discovery Rate (FDR) correction, adjusted for
multiple group comparisons. Spearman's rank correlation was used to assess correlation
covariant. P-values were adjusted using Holm's method, accounting for multiple group
comparisons. P value annotations: ns: $p > 0.05$, *: $0.01 < p <= 0.05$, **: $0.001 < p <= 0.01$, ***: $p <= 0.001$.
$<= 0.001$.
1012
Longitudinal Analysis. To capture statistically meaningful temporal signals, we employed
various statistical and modeling methods, accounting for both linear and non-linear trends and
intra-individual correlations:
1. Interquartile Range (IQR) and Intraclass Correlation Coefficient (ICC). We initially assessed
statistics at different time points by computing the IQR and ICC. Data were standardized to a
mean of zero and a standard deviation of one to ensure comparability across features with
different scales. The IQR quantified variability, while the ICC assessed the dependence of
repeated measurements[118], indicating the similarity of measurements over time. Data showed
no statistical dependence and no trend of stable variance across time points.
2. Generalized Linear Models (GLMs). GLMs[119] were then used to analyze the effects of time
points, considering age, gender, and their interactions. Time points were included as predictors
to reveal changes in dependent variables over time, with interaction terms exploring variations
based on age and gender. Random effects accounted for intra-individual correlations. Although
12 features out of 5000 showed weak trends over time (slopes < 0.2), they were not deemed
sufficient to be potential longitudinal biomarkers, possibly due to individualized patterns.
3. Repeated Measures Correlation (rmcorr). To better consider individual effects, we employed
rmcorr[120] to assess consistent patterns of association within individuals over time. This method
captured stable within-individual associations across different time points. However, only 30

features out of 5000 showed weak slopes (< 0.3), and these were not considered sufficient to
conclude the presence of longitudinal signals.
4. **Smoothing Spline ANOVA (SS-ANOVA)**. We then considered the longitudinal trends could be
non-linear and more complex. To model complex, non-linear relationships between response
variables and predictors over time, SS-ANOVA[121] was used. SS-ANOVA uncovered non-linear
trends and interactions in the omics data, however, no strong temporal signals were identified.
In conclusion, robust analysis of the longitudinal data, accounting for both linear and non-linear
trends and intra-individual correlations, revealed the difficulty in extracting strong and
statistically meaningful temporal signals. As Myalgic Encephalomyelitis/Chronic Fatigue
Syndrome (ME/CFS) is a disease that usually lasts for decades with non-linear progression, the
four-year tracking period with annual measurements is likely insufficient for capturing
consistent temporal signals, necessitating longer follow-up periods.

Data and Code

Metagenomics data is being deposited under the BioProject submission number SUB14546737 and will be publicly available as of the date of publication. Accession numbers are listed in the key resources table. BioMapAI framework is available at https://github.com/ohlab/BioMapAI/codes/AI. All original code, analyzed data and trained model has been deposited at https://github.com/ohlab/BioMapAI. Other 'omics data and any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Acknowledgements and Funding

We are thankful to the Oh, Unutmaz, and Li laboratories for inspiring discussions and acknowledge the contribution of the Genome Technologies Service at The Jackson Laboratory for expert assistance with sample sequencing for the work described in this publication. We also thank the clinical support team at the Bateman Horne Center and all the individuals who participated in this study. This work was funded by 1U54NS105539. JO is additionally supported by the NIH (1 R01 AR078634-01, DP2 GM126893-01, 1 U19 AI142733, 1 R21 AR075174).

Author Contributions

Conceptualization: DU, JO, SDV, LB, RX; Data Curation: RX, CG, SDV, LB; Formal Analysis: RX; Funding Acquisition: DU, JO, SDV, LB; Clinical sample design and collection: SDV, LB; Investigation: RX, CG, EF, SDV, LB; Project Administration: JO, DU, LB, SDV, CG; Resources: DU, JO, SDV, LB; Supervision: JO; Visualization and Writing: RX, JO; Writing - Review and Editing: RX, CG, SDV, LB, DU, JO.

Competing Interests

Dr. Suzanne D. Vernon is affiliated and has a financial interest with The BioCollective, a company that provided the BioCollector, the collection kit used for at home stool collection discussed in this manuscript. No other authors have competing interests.

Lead Contact

Further information and requests for resources and reagents should be directed to the lead contact, Julia Oh (Julia.Oh@jax.org).

 Supplemental Figure

1077 **Supplemental Figure 1: Data Pairedness Overview and Heterogeneity in Healthy and Patients.**
1078 **A) Cohort Composition and Data Collection.** Over four years, 515 time points were collected:
1079 baseline year from all 249 donors (Healthy N=96, ME/CFS N=153); second year from 168
1080 individuals (Healthy N=58, ME/CFS N=110); third year from 94 individuals (Healthy N=13,
1081 ME/CFS N=81); fourth year from N=4 ME/CFS patients. Nearly 400 collection points included
1082 complete sets of 5 'omics datasets, with others capturing 3-4 'omics profiles. Clinical metadata
1083 and blood measures were collected at all 515 points. Immune profiles from PBMCs were
1084 recorded at 489 points, microbiome data from stool samples at 479 points, and plasma
1085 metabolome data at 414 points. A total of 1,471 biosamples were collected. **B-C) Heterogeneity**
1086 **of B) Healthy Controls and C) All Patients in Symptom Severity and 'Omics Profiles.**
1087 Supplemental information for Figure 1B, which shows examples from 20 patients. Variability in
1088 symptom severity (top) and 'omics profiles (bottom) for all healthy controls and all patients with
1089 3-4 time points. **D) Distribution of 12 Clinical Symptoms in ME/CFS and Control.** Density plots
1090 compare the distributions of 12 clinical scores between control (blue) and ME/CFS patients
1091 (orange) with the y-axis representing severity (scaled from 0% to 100%). Clinical scores include
1092 RAND36 subscales (e.g., Physical Functioning, Emotional Wellbeing), Cognitive Efficiency from

the DANA test, Orthostatic Intolerance from the NLT test, Sleep Problems from the PSQI
questionnaire, and Gastrointestinal Symptoms from the GSRS questionnaire. **E) Principal**
Coordinates Analysis (PCoA) of each 'Omics. PCoA based on Bray-Curtis distance for clinical
scores, immune profiles, plasma metabolome, blood measures, species abundance, and KEGG
gene data. Control samples (blue) and ME/CFS patients (red) show distinct clustering. Here,
except for the clinical scores, controls are indistinguishable from patients, highlighting the
difficulty of building classification models. **Abbreviations:** ME/CFS, Myalgic
Encephalomyelitis/Chronic Fatigue Syndrome; PCoA, Principal Coordinates Analysis; RAND36,
36-Item Short Form Health Survey; DANA, DANA Brain Vital; NLT, NASA Lean Test; PSQI,
Pittsburgh Sleep Quality Index; GSRS, Gastrointestinal Symptom Rating Scale; KEGG, Kyoto
Encyclopedia of Genes and Genomes. **Related to:** Figure 1-2.

Supplemental Figure 2: BioMapAI's Performance at Clinical Score Reconstruction and Disease
Classification. A) Density map of True vs. Predicted Clinical Scores. Supplemental information
for Figure 2B, which shows three examples. Here, the full set of 12 clinical scores compares the
true score, y (Column 1), against BioMapAI's predictions generated from different 'omics
profiles – $\hat{y}_{immune}, \hat{y}_{species}, \hat{y}_{KEGG}, \hat{y}_{metabolome}, \hat{y}_{quest}, \hat{y}_{omics}$ (Columns 2-7). **B) Scatter Plot of**
True vs. Predicted Clinical Scores. Scatter plots display the relationship between true clinical
scores (x-axis) and predicted clinical scores (y-axis) for six different models: Omics, Immune,
Species, KEGG, Metabolome, and Quest Labs. Each plot demonstrates the clinical score
prediction accuracy for each model. **C) ROC Curve for Disease Classification with Original**
Clinical Scores. The Receiver Operating Characteristic (ROC) curve evaluates the performance of
disease classification using the original 12 clinical scores. The mean Area Under the Curve (AUC)
is 0.99, indicating high prediction accuracy, which aligns with the clinical diagnosis of ME/CFS
based on key symptoms. **D) 3D t-SNE Visualization of Hidden Layers.** 3D t-SNE plots show how
BioMapAI progressively distinguishes disease from control across hidden layers for five trained
'omics models: Immune, KEGG, Species, Metabolome, and Quest Labs. Each plot uses the first
three principal components to show the spatial distribution of control samples (blue) and
ME/CFS patients (red). The progression from the input layer (mixed groups) to Hidden Layer 3
(fully separated groups) illustrates how BioMapAI progressively learns to separate ME/CFS from
healthy controls. **Abbreviations:** ROC, Receiver Operating Characteristic; AUC, Area Under the
Curve; t-SNE, t-Distributed Stochastic Neighbor Embedding; PCs, Principal Components; y, True
Score; $\hat{y}$, Predicted Score. **Related to:** Figure 2.

1126
Supplemental Figure 3: Disease-Specific Biomarkers - Top 10 Biomarkers Shared across
Clinical Symptoms and Multiple Models. Through the top 30 high-ranking features for each
score, we discovered that the most critical features for all 12 symptoms were largely shared
and consistently validated across ML and DL models, particularly the foremost 10. Here, this
multi-panel figure presents the top 10 most significant features identified by BioMapAI across
five 'omics profiles, highlighting their importance in predicting clinical symptoms and diagnostic
outcomes across BioMapAI, DNN, and GBDT models, along with their data prevalence. Each
vertical section represents one 'omics profile, with columns of biomarkers ordered by average
feature importance from right to left. From top to bottom: *1. Feature Importance Ranking in*
BioMapAI. Lines depict the rank of feature importance for each clinical score, color-coded by
the 12 clinical scores. Consistency among the top 5 features suggests they are shared disease
biomarkers crucial for all clinical symptoms; *2. Heatmap of SHAP Values from BioMapAI.* This
heatmap shows averaged SHAP values with the 12 scores on the rows and the top 10 features
in the columns. Darker colors indicate a stronger impact on the model's output; *3. Swarm Plot*
of SHAP Values from DNN. This plot represents the distribution of feature contributions from
DNN, which is structurally similar to BioMapAI but omits the third hidden layer (Z^3). SHAP
values are plotted vertically, ranging from negative to positive, showing each feature's influence
on prediction outcomes. Points represent individual samples, with color gradients denoting
actual feature values. For instance, *Dysosmobacteria welbionis*, identified as the most critical
species, shows that greater species relative abundance correlates with a higher likelihood of
disease prediction; *4. Bar Graphs of Feature Importance in GBDT*. GBDT is another machine
learning model used for comparison. Each bar's height indicates a feature's significance within
the GBDT model, providing another perspective on the predictive relevance of each biomarker;
5. Heatmap of Normalized Raw Abundance Data. This heatmap compares biomarker prevalence
between healthy and disease states, with colors representing z-scored abundance values,
highlighting biomarker differences between groups. **Abbreviations:** DNN: Here refer to our
deep Learning model without the hidden 3, 'spread out' layer; GBDT: Gradient Boosting
Decision Tree; SHAP: SHapley Additive exPlanations. **Supporting Materials:** Supplemental Table
5. **Related to:** Figure 3.

Supplemental Figure 4: Symptom-Specific Biomarkers - Immune, KEGG and Metabolome Models. By linking 'omics profiles to clinical symptoms, BioMapAI identified unique symptom-

specific biomarkers in addition to disease-specific biomarkers (Supplemental Figure 3). Each
'omics has a circularized diagram (Figure 3A, Supplemental Figure 4B-D) to display how
BioMapAI use this 'omics profile to predict 12 clinical symptoms and to discuss the contribution
of disease- and symptom-specific biomarkers. Detailed correlation between symptom-specific
biomarkers and their corresponding symptoms is in Supplemental Figure 5. **A) Examples of**
Sleeping Problem-Specific Species' and Gastrointestinal-Specific Species' Contributions.
Supplemental information for Figure 3D, which shows the contribution of pain-specific species.
B-D) Circularized Diagram for Immune, KEGG and Metabolome Models. Supplemental
information for Figure 3A, which shows the species model. **E-F) Zoomed Segment for Pain in**
KEGG and Metabolome Model. Supplemental information for Figure 3B, which shows the
zoomed segment for pain in the species and immune models. **Abbreviations and Supporting**
Materials: Supplemental Figure 5. **Related to:** Figure 3.

1171
Supplemental Figure 5: Symptom-Specific Biomarkers - Different Correlation Patterns of
Biomarkers to Symptom. Supplemental information for Figure 3C, which shows six pain
biomarkers from multiple models. Here for each 'omics, we plotted the correlation of symptom-

specific biomarkers (x-axis) to its related symptom (y-axis), colored by SHAP value (contribution
to the symptom). **Abbreviations:** CD4, Cluster of Differentiation 4; CD8, Cluster of
Differentiation 8; IFNg, Interferon Gamma; DC, Dendritic Cells; MAIT, Mucosal-Associated
Invariant T; Th17, T helper 17 cells; CD4+ TCM, CD4+ Central Memory T cells; DC CD1c+ mBtp+,
Dendritic Cells expressing CD1c+ and myelin basic protein; DC CD1c+ mHsp, Dendritic Cells
expressing CD1c+ and heat shock protein; CD4+ TEM, CD4+ Effector Memory T cells; CD4+ Th17
rfx4+, CD4+ T helper 17 cells expressing RFX4; *F. prausnitzii, Faecalibacterium prausnitzii; A.*
communis, Akkermansia communis; NAD, Nicotinamide Adenine Dinucleotide. **Related to:**
Figure 3.

Supplemental Figure 6: 'Omics WGCNA Modules and Host-Microbiome Network. A) Correlation of WGCNA Modules with Clinical Metadata. Weighted Gene Co-expression Network Analysis (WGCNA) was used to identify co-expression modules for each 'omics layer: species, KEGG, immune, and metabolome. The top dendrograms show hierarchical clustering of 'omics features, with modules identified. The bottom heatmap shows the relationship of module eigengenes (colored as per dendrogram) with clinical metadata – including demographic information and environmental factors - and 12 clinical scores. General linear models were used to determine the primary clinical drivers for each module, with the color gradient representing the coefficients (red = positive, blue = negative). Microbial modules were influenced by disease presence and energy-fatigue levels, while metabolome and immune modules correlated with age and gender. **B-C) Microbiome-Immune-Metabolome Network in B) Patient and C) Healthy Subgroups.** Supplemental information for Figure 4A (Healthy Network) and 4B (Patient Subgroups). Figure 4A is the healthy network; here, Supplemental Figure 6B presented the shifted correlations in all patients. Figure 4B represented the network in patient subgroups; here, Supplemental Figure 6C is the corresponding healthy counterpart, for example, female patients were compared with female controls to exclude gender influences. **D) Differences in Host-Microbiome Correlations between Healthy and Patient Subgroups.** Selected host-microbiome module pairs are grouped on the x-axis (e.g., pyruvate to blood modules, steroids to gut microbiome). Significant positive and negative correlations (top and bottom y-axis) of module members pairs are shown as dots for each subgroup (blue = healthy, orange = patient) (Spearman, adjusted $p < 0.05$), from left to right: Young, Elder, Female, Male, NormalWeight, OverWeight Healthy and Young, Elder, Female, Male, NormalWeight, OverWeight Patient. The middle bars represent the total count of associations. This panel highlights the shifts in host-microbiome networks from health to disease, for example, in patients, the loss of pyruvate to host blood modules correlation and the increase of INFg+ CD4 memory correlation with gut microbiome. **Abbreviations:** WGCNA, Weighted Gene Co-expression Network Analysis; AA, Amino Acids; SCFA, Short-Chain Fatty Acids; IL, Interleukin; GM-CSF, Granulocyte-Macrophage Colony-Stimulating Factor. **Related to**: Figure 4.

Supplemental Table
Supplemental Table 1 Sample Metadata and Clinical Scores
Supplemental Table 2 Model Performance at Reconstructing Twelve Clinical Scores: Averaged
Average Mean Squared Error by Model
Supplemental Table 3 Model Performance in Diagnostic Comparison—Within-Cohort, Cross-
Validated by Various ML and DL Models
Supplemental Table 4 Model Performance in Diagnostic Comparison—Across Independent
Cohorts
Supplemental Table 5 Disease-Specific Biomarker: Averaged Feature Contribution of BioMapAI,
DNN and GDBT
Supplemental Table 6 Symptom-Specific Biomarker: Distinct Sets of Biomarkers for Each
Symptom
Supplemental Table 7 WGCNA Module Eigengene
Supplemental Table 8 Targeted Pathways: Normalized Gene Read Counts and Their Correlation
with Blood Responders

References

References

1.   de Mel, S., Lim, S. H., Tung, M. L. & Chng, W.-J. Implications of Heterogeneity in Multiple
Myeloma. *BioMed Res. Int.* **2014**, 232546 (2014).

2.   Wallstrom, G., Anderson, K. S. & LaBaer, J. Biomarker Discovery for Heterogeneous
Diseases.

3.   Weyand, C. M., McCarthy, T. G. & Goronzy, J. J. Correlation between disease phenotype
and genetic heterogeneity in rheumatoid arthritis. *J. Clin. Invest.* **95**, 2120–2126 (1995).

4.   Xiong, R. *et al.* Multi-'omics of gut microbiome-host interactions in short- and long-term
myalgic encephalomyelitis/chronic fatigue syndrome patients. *Cell Host Microbe* **31**, 273-287.e5
(2023).

5.   Giladi, N., Mirelman, A., Thaler, A. & Orr-Urtreger, A. A Personalized Approach to
Parkinson's Disease Patients Based on Founder Mutation Analysis. *Front. Neurol.* **7**, (2016).

6.   Brown, S. M. *et al.* Consistent Effects of Early Remdesivir on Symptoms and Disease
Progression Across At-Risk Outpatient Subgroups: Treatment Effect Heterogeneity in PINETREE
Study. *Infect. Dis. Ther.* **12**, 1189–1203 (2023).

7.   Iwasaki, T. & Sano, H. Predicting Treatment Responses and Disease Progression in
Myeloma using Serum Vascular Endothelial Growth Factor and Hepatocyte Growth Factor
Levels. *Leuk. Lymphoma* **44**, 1275–1279 (2003).

8.   Hare, P. J., LaGree, T. J., Byrd, B. A., DeMarco, A. M. & Mok, W. W. K. Single-Cell
Technologies to Study Phenotypic Heterogeneity and Bacterial Persisters. *Microorganisms* **9**,
2277 (2021).

9.   Cohen, R. M., Haggerty, S. & Herman, W. H. HbA1c for the Diagnosis of Diabetes and
Prediabetes: Is It Time for a Mid-Course Correction? *J. Clin. Endocrinol. Metab.* **95**, 5203–5206
(2010).

10.   Zhou, W. *et al.* Longitudinal multi-omics of host–microbe dynamics in prediabetes.
Nature **569**, 663–671 (2019).

11.   Hong, S. *et al.* Cancer Statistics in Korea: Incidence, Mortality, Survival, and Prevalence in
2017. *Cancer Res. Treat.* **52**, 335–350 (2020).

12.   Zeeshan, S., Xiong, R., Liang, B. T. & Ahmed, Z. 100 years of evolving gene–disease
complexities and scientific debutants. *Brief. Bioinform.* **21**, 885–905 (2020).

13.   Cortes Rivera, M., Mastronardi, C., Silva-Aldana, C. T., Arcos-Burgos, M. & Lidbury, B. A.
Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Comprehensive Review. *Diagnostics* **9**,
91 (2019).

14.   Sweetman, E. *et al.* Current Research Provides Insight into the Biological Basis and
Diagnostic Potential for Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS).
Diagnostics **9**, 73 (2019).

15.   Noor, N. *et al.* A Comprehensive Update of the Current Understanding of Chronic Fatigue
Syndrome. *Anesthesiol. Pain Med.* **11**, e113629 (2021).

16.   Ruiz-Pablos, M., Paiva, B., Montero-Mateo, R., Garcia, N. & Zabaleta, A. Epstein-Barr
Virus and the Origin of Myalgic Encephalomyelitis or Chronic Fatigue Syndrome. *Front. Immunol.*
12, 656797 (2021).

17.   Su, R. *et al.* The TLR3/IRF1/Type III IFN Axis Facilitates Antiviral Responses against
Enterovirus Infections in the Intestine. *mBio* **11**, 10.1128/mbio.02540-20 (2020).

18.   Anderson, D. E. *et al.* Lack of cross-neutralization by SARS patient sera towards SARS-
CoV-2. *Emerg. Microbes Infect.* **9**, 900–902 (2020).

19.   Poenaru, S., Abdallah, S. J., Corrales-Medina, V. & Cowan, J. COVID-19 and post-
infectious myalgic encephalomyelitis/chronic fatigue syndrome: a narrative review. *Ther. Adv.*
Infect. Dis. **8**, 20499361211009385 (2021).

20.   Reuken, P. A. *et al.* Longterm course of neuropsychological symptoms and ME/CFS after
SARS-CoV-2-infection: a prospective registry study. *Eur. Arch. Psychiatry Clin. Neurosci.* (2023)
doi:10.1007/s00406-023-01661-3.

21.   Bretherick, A. D. *et al.* Typing myalgic encephalomyelitis by infection at onset: A
DecodeME study. *NIHR Open Res.* **3**, 20 (2023).

22.   Bae, J. & Lin, J.-M. S. Healthcare Utilization in Myalgic Encephalomyelitis/Chronic Fatigue
Syndrome (ME/CFS): Analysis of US Ambulatory Healthcare Data, 2000–2009. *Front. Pediatr.* **7**,
(2019).

23.   Zheng, Y. & Zhu, Z. Editorial: Retrieving meaningful patterns from big biomedical data
with machine learning approaches. *Front. Genet.* **14**, (2023).

24.   Leelatian, N. *et al.* Unsupervised machine learning reveals risk stratifying glioblastoma
tumor cells. *eLife* **9**, e56879 (2020).

25.   Su, Q. *et al.* The gut microbiome associates with phenotypic manifestations of post-
acute COVID-19 syndrome. *Cell Host Microbe* **32**, 651-660.e4 (2024).

26.   Bourgonje, A. R., van Goor, H., Faber, K. N. & Dijkstra, G. Clinical Value of Multiomics-
Based Biomarker Signatures in Inflammatory Bowel Diseases: Challenges and Opportunities.
Clin. Transl. Gastroenterol. **14**, e00579 (2023).

27.   Marcos-Zambrano, L. J. *et al.* Applications of Machine Learning in Human Microbiome
Studies: A Review on Feature Selection, Biomarker Identification, Disease Prediction and
Treatment. *Front. Microbiol.* **12**, (2021).

28.   Guo, C. *et al.* Deficient butyrate-producing capacity in the gut microbiome is associated
with bacterial network disturbances and fatigue symptoms in ME/CFS. *Cell Host Microbe* **31**,
288-304.e8 (2023).

29.   Raijmakers, R. P. H. *et al.* Multi-omics examination of Q fever fatigue syndrome identifies
similarities with chronic fatigue syndrome. *J. Transl. Med.* **18**, 448 (2020).

30.   Germain, A. *et al.* Plasma metabolomics reveals disrupted response and recovery
following maximal exercise in myalgic encephalomyelitis/chronic fatigue syndrome. *JCI Insight* **7**,
(2023).

31.   Che, X. *et al.* Metabolomic Evidence for Peroxisomal Dysfunction in Myalgic
Encephalomyelitis/Chronic Fatigue Syndrome. *Int. J. Mol. Sci.* **23**, 7906 (2022).

32.   Liñares-Blanco, J., Fernandez-Lozano, C., Seoane, J. A. & López-Campos, G. Machine
Learning Based Microbiome Signature to Predict Inflammatory Bowel Disease Subtypes. *Front.*
Microbiol. **13**, (2022).

33.   He, F. *et al.* Development and External Validation of Machine Learning Models for
Diabetic Microvascular Complications: Cross-Sectional Study With Metabolites. *J. Med. Internet*
Res. **26**, e41065 (2024).

34.   Hawken, S. *et al.* External validation of machine learning models including newborn
metabolomic markers for postnatal gestational age estimation in East and South-East Asian
infants. Preprint at https://doi.org/10.12688/gatesopenres.13131.2 (2021).

35. Mora-Ortiz, M., Trichard, M., Oregioni, A. & Claus, S. P. Thanatometabolomics: introducing NMR-based metabolomics to identify metabolic biomarkers of the time of death. *Metabolomics* **15**, 37 (2019).

36. Balasubramanian, R. *et al.* Metabolomic profiles associated with all-cause mortality in the Women's Health Initiative. *Int. J. Epidemiol.* **49**, 289–300 (2020).

37. Li, H., Ren, M. & Li, Q. 1H NMR-Based Metabolomics Reveals the Intrinsic Interaction of Age, Plasma Signature Metabolites, and Nutrient Intake in the Longevity Population in Guangxi, China. *Nutrients* **14**, 2539 (2022).

38. Kondoh, H. & Kameda, M. Metabolites in aging and aging-relevant diseases: Frailty, sarcopenia and cognitive decline. *Geriatr. Gerontol. Int.* **24**, 44–48 (2024).

39. Peng, S., Shen, Y., Wang, M. & Zhang, J. Serum and CSF Metabolites in Stroke-Free Patients Are Associated With Vascular Risk Factors and Cognitive Performance. *Front. Aging Neurosci.* **12**, (2020).

40. Duerler, P., Vollenweider, F. X. & Preller, K. H. A neurobiological perspective on social influence: Serotonin and social adaptation. *J. Neurochem.* **162**, 60–79 (2022).

41. Pomrenze, M. B., Paliarin, F. & Maiya, R. Friend of the Devil: Negative Social Influences Driving Substance Use Disorders. *Front. Behav. Neurosci.* **16**, (2022).

42. Laslett, A.-M. Commentary on Bischof et al.: Empirical and conceptual paradigms for studying secondary impacts of a person's substance use. *Addiction* **117**, 3148–3149 (2022).

43. Carco, C. *et al.* Increasing Evidence That Irritable Bowel Syndrome and Functional Gastrointestinal Disorders Have a Microbial Pathogenesis. *Front. Cell. Infect. Microbiol.* **10**, (2020).

44. Saffouri, G. B. *et al.* Small intestinal microbial dysbiosis underlies symptoms associated with functional gastrointestinal disorders. *Nat. Commun.* **10**, 2012 (2019).

45. Liang, S., Wu, X., Hu, X., Wang, T. & Jin, F. Recognizing Depression from the Microbiota–Gut–Brain Axis. *Int. J. Mol. Sci.* **19**, 1592 (2018).

46. Zhu, F., Tu, H. & Chen, T. The Microbiota–Gut–Brain Axis in Depression: The Potential Pathophysiological Mechanisms and Microbiota Combined Antidepression Effect. *Nutrients* **14**, 2081 (2022).

47. Topan, R. & Scott, S. M. Sleep: An Overlooked Lifestyle Factor in Disorders of Gut-Brain Interaction. *Curr. Treat. Options Gastroenterol.* **21**, 435–446 (2023).

48. Moens de Hase, E. *et al.* Impact of metformin and Dysosmobacter welbionis on diet-induced obesity and diabetes: from clinical observation to preclinical intervention. *Diabetologia* **67**, 333–345 (2024).

49. Amabebe, E., Robert, F. O., Agbalalah, T. & Orubu, E. S. F. Microbial dysbiosis-induced obesity: role of gut microbiota in homoeostasis of energy metabolism. *Br. J. Nutr.* **123**, 1127–1137 (2020).

50. Kavanagh, P. *et al.* Tentative identification of the phase I and II metabolites of two synthetic cathinones, MDPHP and α-PBP, in human urine. *Drug Test. Anal.* **12**, 1442–1451 (2020).

51. Wang, J.-H. *et al.* Clinical evidence of the link between gut microbiome and myalgic encephalomyelitis/chronic fatigue syndrome: a retrospective review. *Eur. J. Med. Res.* **29**, 148 (2024).

52. Lenoir, M. *et al.* Butyrate mediates anti-inflammatory effects of Faecalibacterium prausnitzii in intestinal epithelial cells through Dact3. *Gut Microbes* (2020).

53. Sokol, H. *et al.* Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. *Proc. Natl. Acad. Sci.* **105**, 16731–16736 (2008).

54. Quévrain, E. *et al.* Identification of an anti-inflammatory protein from Faecalibacterium prausnitzii, a commensal bacterium deficient in Crohn's disease. *Gut* **65**, 415–425 (2016).

55. Miquel, S. *et al.* Identification of Metabolic Signatures Linked to Anti-Inflammatory Effects of Faecalibacterium prausnitzii. *mBio* **6**, 10.1128/mbio.00300-15 (2015).

56. Vital, M., Howe, A. C. & Tiedje, J. M. Revealing the Bacterial Butyrate Synthesis Pathways by Analyzing (Meta)genomic Data. *mBio* **5**, e00889-14 (2021).

57. Recharla, N., Geesala, R. & Shi, X.-Z. Gut Microbial Metabolite Butyrate and Its Therapeutic Role in Inflammatory Bowel Disease: A Literature Review. *Nutrients* **15**, 2275 (2023).

58. Monteiro, C. R. A. V. *et al.* In Vitro Antimicrobial Activity and Probiotic Potential of Bifidobacterium and Lactobacillus against Species of Clostridium. *Nutrients* **11**, 448 (2019).

59. Zhao, M., Li, G. & Deng, Y. Engineering Escherichia coli for Glutarate Production as the C5 Platform Backbone. *Appl. Environ. Microbiol.* **84**, e00814-18 (2018).

60. Nguyen-Lefebvre, A. T., Selzner, N., Wrana, J. L. & Bhat, M. The hippo pathway: A master regulator of liver metabolism, regeneration, and disease. *FASEB J.* **35**, e21570 (2021).

61. Khan, M. A., Gupta, A., Sastry, J. L. N. & Ahmad, S. Hepatoprotective potential of kumaryasava and its concentrate against CCl4-induced hepatic toxicity in Wistar rats. *J. Pharm. Bioallied Sci.* **7**, 297–299 (2015).

62. Kim, C.-S. Roles of Diet-Associated Gut Microbial Metabolites on Brain Health: Cell-to-Cell Interactions between Gut Bacteria and the Central Nervous System. *Adv. Nutr.* **15**, 100136 (2024).

63. Rebeaud, J., Peter, B. & Pot, C. How Microbiota-Derived Metabolites Link the Gut to the Brain during Neuroinflammation. *Int. J. Mol. Sci.* **23**, 10128 (2022).

64. Ahmad, S. *et al.* Gut microbiome-related metabolites in plasma are associated with general cognition. *Alzheimers Dement.* **17**, e056142 (2021).

65. Ahmed, Z., Zeeshan, S., Xiong, R. & Liang, B. T. Debutant iOS app and gene-disease complexities in clinical genomics and precision medicine. *Clin. Transl. Med.* **8**, e26 (2019).

66. Ahmed, Z., Zeeshan, S., Xiong, R. & Liang, B. T. PAS-Gen: Guide to iOS app with gene-disease.

67. Ahmed, Z., Wan, S., Zhang, F. & Zhong, W. Artificial intelligence for omics data analysis. *BMC Methods* **1**, 4 (2024).

68. Ahmed, Z., Mohamed, K., Zeeshan, S. & Dong, X. Artificial intelligence with multi-functional machine learning platform development for better healthcare and precision medicine. *Database J. Biol. Databases Curation* **2020**, baaa010 (2020).

69. Germain, A., Ruppert, D., Levine, S. M. & Hanson, M. R. Prospective Biomarkers from Plasma Metabolomics of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Implicate Redox Imbalance in Disease Symptomatology. *Metabolites* **8**, 90 (2018).

70. Lim, E.-J. & Son, C.-G. Review of case definitions for myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). *J. Transl. Med.* **18**, 289 (2020).

71.   Germain, A., Barupal, D. K., Levine, S. M. & Hanson, M. R. Comprehensive Circulatory
Metabolomics in ME/CFS Reveals Disrupted Metabolism of Acyl Lipids and Steroids. *Metabolites*
10, 34 (2020).
72.   Martínez-Lavín, M. Holistic Treatment of Fibromyalgia Based on Physiopathology: An
Expert Opinion. *JCR J. Clin. Rheumatol.* **26**, 204 (2020).
73.   López-Hernández, Y. *et al.* The plasma metabolome of long COVID patients two years
after infection. *Sci. Rep.* **13**, 12420 (2023).
74.   Iqbal, M., Elzembely, H. I. & Said, O. M. Letter to the Editor: “Self-Reported
Student Awareness and Prevalence of Computer Vision Syndrome During COVID-19 Pandemic at
Al-Baha University” [Letter]. *Clin. Optom.* **14**, 193–194 (2022).
75.   Sedeh, F. B. *et al.* The correlation between self-reported hand eczema and clinically
based diagnosis in professional cleaners. *Contact Dermatitis* cod.14611 (2024)
doi:10.1111/cod.14611.
76.   Jason, L. A., Yoo, S. & Bhatia, S. Patient perceptions of infectious illnesses preceding
Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Chronic Illn.* **18**, 901–910 (2022).
77.   Hanson, M. R. The viral origin of myalgic encephalomyelitis/chronic fatigue syndrome.
PLOS Pathog. **19**, e1011523 (2023).
78.   Hamine, S., Gerth-Guyette, E., Faulx, D., Green, B. B. & Ginsburg, A. S. Impact of mHealth
Chronic Disease Management on Treatment Adherence and Patient Outcomes: A Systematic
Review. *J. Med. Internet Res.* **17**, e3951 (2015).
79.   Clark, N. M. Management of Chronic Disease by Patients. *Annu. Rev. Public Health* **24**,
289–313 (2003).
80.   Derman, I. D. *et al.* High-throughput bioprinting of the nasal epithelium using patient-
derived nasal epithelial cells. *Biofabrication* **15**, 044103 (2023).
81.   Fleming, E. *et al.* Cultivation of common bacterial species and strains from human skin,
oral, and gut microbiota. *BMC Microbiol.* **21**, 278 (2021).
82.   Ren, J., Cislo, P., Cappelleri, J. C., Hlavacek, P. & DiBonaventura, M. Comparing g-
computation, propensity score-based weighting, and targeted maximum likelihood estimation
for analyzing externally controlled trials with both measured and unmeasured confounders: a
simulation study. *BMC Med. Res. Methodol.* **23**, 18 (2023).
83.   Lynn, J. V., Buchman, L. K., Breuler, C. J. & Buchman, S. R. Surgical Timing and
Neurocognitive Development among Patients with Craniosynostosis: Analysis of Confounders.
Plast. Reconstr. Surg. **151**, 821 (2023).
84.   Karhan, E. *et al.* *Perturbation of Effector and Regulatory T Cell Subsets in Myalgic*
Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS). 2019.12.23.887505
https://www.biorxiv.org/content/10.1101/2019.12.23.887505v1 (2019)
doi:10.1101/2019.12.23.887505.
85.   Krumina, A. *et al.* Clinical Profile and Aspects of Differential Diagnosis in Patients with
ME/CFS from Latvia. *Medicina (Mex.)* **57**, 958 (2021).
86.   Zubcevik, N. *et al.* Symptom Clusters and Functional Impairment in Individuals Treated
for Lyme Borreliosis. *Front. Med.* **7**, (2020).
87.   Costa, G. G., Pereira, A. R. & Carvalho, A. S. Pericardite lúpica: dor torácica e febre em
tempos de COVID-19. *Rev. Port. Med. Geral E Fam.* **38**, 300–4 (2022).

88.   Vyas, J., Muirhead, N., Singh, R., Ephgrave, R. & Finlay, A. Y. Impact of myalgic
encephalomyelitis/chronic fatigue syndrome (ME/CFS) on the quality of life of people with
ME/CFS and their partners and family members: an online cross-sectional survey. *BMJ Open* **12**,
e058128 (2022).
89.   Martinez, A., Okoh, A., Ko, Y.-A. & Wells, B. Racial Differences in FMD.
2023.02.10.23285630 Preprint at https://doi.org/10.1101/2023.02.10.23285630 (2023).
90.   Trivedi, M. S. *et al.* Identification of Myalgic Encephalomyelitis/Chronic Fatigue
Syndrome-associated DNA methylation patterns. *PLOS ONE* **13**, e0201066 (2018).
91.   Bouquet, J. *et al.* Whole blood human transcriptome and virome analysis of ME/CFS
patients experiencing post-exertional malaise following cardiopulmonary exercise testing. *PLOS*
ONE **14**, e0212193 (2019).
92.   Lande, A. *et al.* Human Leukocyte Antigen alleles associated with Myalgic
Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS). *Sci. Rep.* **10**, 5267 (2020).
93.   Almenar-Pérez, E. *et al.* Epigenetic Components of Myalgic Encephalomyelitis/Chronic
Fatigue Syndrome Uncover Potential Transposable Element Activation. *Clin. Ther.* **41**, 675–698
(2019).
94.   Das, S., Taylor, K., Kozubek, J., Sardell, J. & Gardner, S. Genetic risk factors for ME/CFS
identified using combinatorial analysis. *J. Transl. Med.* **20**, 598 (2022).
95.   Caruana, E. J., Roman, M., Hernández-Sánchez, J. & Solli, P. Longitudinal studies. *J.*
Thorac. Dis. **7**, E537–E540 (2015).
96.   White, R. T. & Arzi, H. J. Longitudinal Studies: Designs, Validity, Practicality, and Value.
Res. Sci. Educ. **35**, 137–149 (2005).
97.   Aurora, C., Cecilia, A. & Adina, H. The Role of Diet in the Treatment of Chronic Diseases
Case Study. *ARS Medica Tomitana* **27**, 153–156 (2021).
98.   Therrien, R. & Doyle, S. Role of training data variability on classifier performance and
generalizability. in *Medical Imaging 2018: Digital Pathology* vol. 10581 58–70 (SPIE, 2018).
99.   Zhang, B., Qin, A. K., Pan, H. & Sellis, T. A Novel DNN Training Framework via Data
Sampling and Multi-Task Optimization. in *2020 International Joint Conference on Neural*
Networks (IJCNN) 1–8 (2020). doi:10.1109/IJCNN48605.2020.9207329.
100.   Lathan, C., Spira, J. L., Bleiberg, J., Vice, J. & Tsao, J. W. Defense Automated
Neurobehavioral Assessment (DANA)-psychometric properties of a new field-deployable
neurocognitive assessment tool. *Mil. Med.* **178**, 365–371 (2013).
101.   Resnick, H. E. & Lathan, C. E. From battlefield to home: a mobile platform for assessing
brain health. *mHealth* **2**, 30 (2016).
102.   Lee, J. *et al.* Hemodynamics during the 10-minute NASA Lean Test: evidence of
circulatory decompensation in a subset of ME/CFS patients. *J. Transl. Med.* **18**, 314 (2020).
103.   Committee on the Diagnostic Criteria for Myalgic Encephalomyelitis/ChronicFatigue
Syndrome, Board on the Health of Select Populations, & Institute of Medicine. *Beyond Myalgic*
Encephalomyelitis/Chronic Fatigue Syndrome: Redefining an Illness. (National Academies Press
(US), Washington (DC), 2015).
104.   Monica, 1776 Main Street Santa & California 90401-3208. 36-Item Short Form Survey
(SF-36) Scoring Instructions. https://www.rand.org/health-care/surveys_tools/mos/36-item-
short-form/scoring.html.

105.   Blanco-Míguez, A. *et al.* Extending and improving metagenomic taxonomic profiling with
uncharacterized species using MetaPhlAn 4. *Nat. Biotechnol.* **41**, 1633–1644 (2023).

106.   Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*
26, 2460–2461 (2010).

107.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

108.   Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q
File Manipulation. *PLOS ONE* **11**, e0163962 (2016).

109.   Mallick, H. *et al.* Multivariable association discovery in population-scale meta-omics
studies. *PLoS Comput. Biol.* **17**, e1009442 (2021).

110.   Chawla, N. V., Bowyer, K. W., Hall, L. O. & Kegelmeyer, W. P. SMOTE: Synthetic Minority
Over-sampling Technique. *J. Artif. Intell. Res.* **16**, 321–357 (2002).

111.   Haibo He, Yang Bai, Garcia, E. A., & Shutao Li. ADASYN: Adaptive synthetic sampling
approach for imbalanced learning. in *2008 IEEE International Joint Conference on Neural*
Networks (IEEE World Congress on Computational Intelligence) 1322–1328 (IEEE, Hong Kong,
China, 2008). doi:10.1109/IJCNN.2008.4633969.

112.   Saripuddin, M., Suliman, A., Syarmila Sameon, S. & Jorgensen, B. N. Random
Undersampling on Imbalance Time Series Data for Anomaly Detection. in *Proceedings of the*
2021 4th International Conference on Machine Learning and Machine Intelligence 151–156
(Association for Computing Machinery, New York, NY, USA, 2022).
doi:10.1145/3490725.3490748.

113.   Lundberg, S. & Lee, S.-I. A Unified Approach to Interpreting Model Predictions. Preprint
at https://doi.org/10.48550/arXiv.1705.07874 (2017).

114.   Abadi, M. *et al.* TensorFlow: A system for large-scale machine learning. Preprint at
https://doi.org/10.48550/arXiv.1605.08695 (2016).

115.   Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. Preprint at
https://doi.org/10.48550/arXiv.1201.0490 (2018).

116.   Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**, 559 (2008).

117.   Antonov, M. *et al.* igraph enables fast and robust network analysis across programming
languages. Preprint at https://doi.org/10.48550/arXiv.2311.10260 (2023).

118.   Koo, T. K. & Li, M. Y. A Guideline of Selecting and Reporting Intraclass Correlation
Coefficients for Reliability Research. *J. Chiropr. Med.* **15**, 155–163 (2016).

119.   Nelder, J. A. & Wedderburn, R. W. M. Generalized Linear Models. *J. R. Stat. Soc. Ser. Gen.*
135, 370–384 (1972).

120.   Bakdash, J. Z. & Marusich, L. R. Repeated Measures Correlation. *Front. Psychol.* **8**, (2017).

121.   Gu, C. Smoothing Spline ANOVA Models: R Package gss. *Smoothing Spline ANOVA*
Models.

1526

A

Healthy

Cognitivity
Efficiency

Orthostatic
Intolerance

Disease

Fatigue

Gastro-intestinal
Symptom

Pain

Clinical Metadata

Blood Measure

Microbiome

Plasma Metabolome

Immune Profile

Samples

Features

Temporal events

B

Severity

Symptoms

Omics PCoA 2

Omics PCoA 1