1

Single-nucleus RNA sequencing reveals enduring signatures of acute stress and chronic
exercise in striatal microglia

Meghan G. Connolly[a], Zachary V. Johnson[c]*, Lynna Chu[d]*, Nicholas D. Johnson[e], Trevor J.
Buhr[f,g], Elizabeth M. McNeill[f]*, Peter J. Clark[f,g]*, and Justin S. Rhodes[a,b]*

[a]University of Illinois Urbana-Champaign, Neuroscience Program, Urbana, IL, 61801, USA
[b]University of Illinois Urbana-Champaign, Department of Psychology, Champaign, IL, 61820,
USA
[c]Emory University, Department of Psychiatry and Behavioral Sciences, Atlanta, GA, 30322,
USA
[d]Iowa State University, Department of Statistics, Ames, IA, 50011, USA
[e]Independent Researcher, USA
[f]Iowa State University, Department of Food Science and Human Nutrition, Ames, IA, 50011
USA
[g]Iowa State University, Neuroscience Graduate Program, Ames, IA, 50011, USA

*These authors contributed equally to this work

Emails: meghanc6@illinois.edu, zachary.vernon.johnson@emory.edu, lchu@iastate.edu,
johnsonn573@gmail.com, tjbuhr@iastate.edu, emcneill@iastate.edu, pjclark@iastate.edu,
jrhodes@illinois.edu

Corresponding author:
Meghan G. Connolly
University of Illinois Urbana-Champaign
Neuroscience Program
405 North Mathews Avenue
Urbana, IL 61801
Email: meghanc6@illinois.edu

Abstract

Acute stress has enduring effects on the brain and motivated behavior across species. For

example, acute stress produces persisting decreases in voluntary physical activity as well as

molecular changes in the striatum, a brain region that regulates voluntary physical activity and

other motivated behaviors. Microglia, the primary immune cells of the central nervous system,

are positioned at the interface between neural responses to stress and neural coordination of

voluntary activity in that they respond to stress, sense molecular changes in the striatum, and

modulate neuronal activity. However, the role of striatal microglia in stress-induced long-term

suppression of voluntary activity is unknown. Here we employ single nucleus RNA-sequencing

to investigate how stress and exercise impact the biology of microglia in the striatum. We find

that striatal microglia display altered activation profiles six weeks after an acute stressor.

Furthermore, we show that access to a running wheel is associated with an additional and distinct

microglial activation profile characterized by upregulation of genes related to complement

components and phagocytosis pathways. Lastly, we find that distinct gene sets show expression

changes associated with general access to a running wheel versus variation in running levels.

Taken together, our results deepen our understanding of the diverse molecular states that striatal

microglia assume in response to stress and exercise and suggest that microglia exhibit a broader

range of functional states than previously thought.

Keywords: Single nucleus RNA-sequencing, exercise, acute stress, microglia, inflammation,

motivation, striatum, single cell

1. Introduction

Adverse life events (e.g., psychological traumas or severe stressors) can have lasting consequences on human health[1,2]. Keeping physically active is crucial for maintaining physical and mental health and is considered one of the best ways of coping with stress[3,4]. However, exposure to adverse events can consequently lower physical activity engagement across the lifespan [5,6]. A better understanding of how adverse experiences chronically impede physical activity engagement is critical for conceptualizing new approaches for improving exercise engagement across the lifespan.

Similarly to humans, rodents exposed to stressors of sufficient intensity can develop persistent reductions in physical activity engagement, offering a useful model for investigating the interactions between stress and physical inactivity. Rodents have a natural inclination to run. For example, rats will lever press for access to a running wheel, indicating that they find wheel running to be rewarding[7–10]. However, rats exposed to a single episode of inescapable tail shocks (acute stress) display a four-fold decrease in voluntary wheel running that outlasts the development of all other well-characterized anxiety- and depression-like behaviors by months[11,12]. Moreover, rats exposed to acute stress can perform forced physical activity at high levels, as performance on treadmill running to failure[11], 15 minutes of forced swim[13], and approximately an hour of shuttle box escape testing[14,15] recovers to the levels of non-stressed rats within a week indicating that the ability to exercise is still present. Taken together, these data suggest that long-term reductions in voluntary wheel running following acute stress may represent a persistent deficit in motivation specifically towards physically exertive voluntary activities.

Multiple regions of the brain contribute to motivation for voluntary physical activity[16].

The striatum, both its dorsal and ventral areas, is particularly notable for its dual role in

motivating reward-seeking behaviors and regulating movement[17–21]. A recent study found major

transcriptomic changes in this region in mice that were selectively bred for increased voluntary

wheel running behavior[22]. The striatum is also well known to be sensitive to stress, with several

studies showing stress alters striatal circuitry[23] and striatal secretion of serotonin and dopamine[24].

Additional studies have shown that stress alte rs striatal-dependent processes like motivation,

locomotion, and associative learning[23,25]. Given that the striatum is both sensitive to stress and

regulates reward information processing, decision-making, and motivated behaviors[21], this brain

region is a lead candidate in which the neurobiological impacts of stress may underlie stress-

associated decreases in motivation for voluntary physical activity. However, the specific cell

types and molecular changes in the striatum that remain impacted long after an acute stressor

remain unknown.

Microglia are primely positioned to regulate striatal responses to stress and striatal-

dependent motivated behaviors owing to their roles in neuroinflammation and homeostatic

regulation[26–29]. In response to an acute stressor like inescapable shock, microglia enact a robust

but brief neuroinflammatory response[30–32] that subsides within approximately 48 hours.

However, some microglia remain in a "primed" state for several weeks after exposure to a

stressor [33–37]. This primed state is distinctively marked by increased baseline levels of

inflammation markers and inflammatory mediators, a reduced threshold for activation into a pro-

inflammatory state, and an intensified inflammatory response upon activation [33,35,37,38]. In

addition to their traditional role as mediators of inflammation, microglia are increasingly

recognized for their specialized functions in regulating the extracellular microenvironment in a

way that influences brain-region specific signaling[39]. For instance, striatal microglia detect and

react to neuronal ATP release by increasing production of adenosine, which in turn suppresses

neuronal activity through adenosine receptors[40]. The responses of striatal microglia to stress may

therefore contribute to long-term alterations in striatal function and motivational deficits.

However, the specific molecular profile of microglia long after an acute stressor that occurs in

coincidence with prolonged reduced voluntary physical activity has not been characterized to our

knowledge.

Here we use single nucleus RNA-sequencing to investigate alterations in striatal

microglia six weeks after acute stress in relation to prolonged decreases in motivation for

physical activity. To evaluate microglial signatures of stress and exercise, male rats were given

one session of inescapable shock stress and then allowed access to a running wheel for 42 days.

Compared to controls, we predicted that microglia in the striatum of acutely stressed rats would

show enduring and distinct forms of neuroinflammatory activation, and that physical activity

would promote an anti-inflammatory microglial phenotype. Our results further our understanding

of microglial functional complexity and reveal candidate molecular and cellular mechanisms

underlying long-term stress-associated decreases in motivation for physical activity.

2. Materials and Methods

2.1 Subjects

Adult male Sprague Dawley (SD) rats from Envigo (250-280g) were single-housed upon

arrival to the Iowa State University Vivarium. Male rats were used for this study as the vast

majority of over five decades of research on stress physiology using tail shock model has been

completed in this exact paradigm using male rats, which could provide important behavioral or

physiological context to outcomes in this study[41–46]. Moreover, long-term running deficits

following acute stress have been previously documented in male, but not yet female, rats [11,12].

Chow (ENVIGO, Tekland 2014) and water were provided *ad libitum* throughout the experiment.

Rooms were controlled for temperature ($21 \pm 1°C$) and photoperiod (12:12 light/dark) for the

entire study. All procedures were approved by the Iowa State University Institutional Animal

Care and Use Committee and adhered to NIH guidelines.

2.2 Experimental Design

Upon arrival, twenty-four rats were individually housed in a standard laboratory cage

with (Runner) or without (Sedentary) a locked 13-inch diameter running wheel (STARR Life

Sciences, PA, USA) for one week. Note that rats in the sedentary group were deliberately not

housed in cages with locked running wheels to keep physical activity to a minimum, as rodents

tend to climb on locked wheels[47,48]. Half of the rats in each exercise condition were then

randomly assigned to receive a single episode of 0 (no stress) or 100 (stress) uncontrollable tail

shocks (**Fig. 1A**). The tail shock paradigm used in the present study followed our previous

publications[24,49,50]. Approximately 2-5 h into the light cycle, rats assigned to the stress condition

were restrained in flat bottom Plexiglas tubes with the tail protruding from the back where

electrodes were placed to deliver 100, 1.25mA average current, 5-s tail shocks on a variable 1

min inter-shock interval. This procedure lasted approximately 100 minutes, though the precise

time varied[24,49,50]. Rats that did not receive stress (i.e. 0 shocks) remained undisturbed in home

cages in a different room during the tail shocks.

48 hours later, running wheels of rats in the runner group were unlocked and rats were

allowed to run freely for the six-week duration of the study. The 48-hour delay between tail

shocks and wheel access was done to minimize risk of stressed rats developing an association

between tail shock exposure and free access to running wheels. Wheel running distance was

measured every hour for the entire six-week duration of the experiment. This time period was

chosen because six, but not three-, weeks of running wheel access can prevent the development

of striatum-involved anxiety- and depression-like behavior in rats[51–53]. Moreover, SD rats

continue to display acute stress-induced persistent reductions in voluntary wheel running

behavior beyond six weeks[11,12]. On Day 42, all subjects were euthanized by rapid decapitation

during the peak running activity period (between 1-5 hours into the dark cycle). The final groups

and sample sizes were sedentary no stress (n=6), sedentary stress (n=6), runner no stress (n=6),

and runner stress (n=6).

In addition to measuring wheel revolutions continuously in 1-hour intervals in the rats

that had access to running wheels throughout the entire experiment, we also measured body mass

at the beginning and end of the experiment. We measured food consumption on days 21, 23, 25,

27, and 41 by subtracting the total weight of the food pellets in each cage from the prior day's

weight. These outcomes were measured to investigate whether stress caused long-term

hypophagia.

2.3 Behavioral Analysis

The wheel running data (total revolutions over the entire 42-day experiment) were

analyzed using a repeated-measures ANOVA with day as the within-subjects factor and

treatment group (stress or no stress) as the between-subjects factor. The differences in body

mass between the last day of the experiment and the first day were analyzed using a 2x2

ANOVA, with stress, exercise, and their interaction as factors. The food consumption data were

analyzed using a 2x2 repeated-measures ANOVA with the 5 sampling days as the within-

subjects factor. When interactions were significant, pairwise differences between means were

evaluated using least-significant difference tests.

2.4 Nuclei Isolation, Library Construction, and Sequencing

179   Rat brains were rapidly extracted. The striatum was microdissected on a glass plate over

ice per our previous publications[49,54,55]. The striatum was placed in pre-weighed cryovials, flash-

frozen with liquid nitrogen, and then stored at $-80°C$ until nuclei isolation. For nuclei isolation,

the striata from all six rats within each treatment group (i.e. no stress sedentary, no stress runner,

stress sedentary, or stress runner) were pooled together and placed in a centrifuge tube, lysed,

and dissociated together to yield four samples. The nuclei were extracted and counted following

protocols described previously[56]. Briefly, frozen striatal tissue was placed in lysis buffer (10 mM

Tris-HCL, 10 mM NaCl, 3 mM $MgCl_2$, 0.1% Nonidet P40 Substitute) on ice for 30 min to lyse

tissue followed by addition of Hibernate AB complete media (2% B27 and 0.05mM Glutamax,

BrainBits HAB-500) and homogenization using a hand-held electronic homogenizer. Pooled

homogenates were then centrifuged for 5 min at 600 x g and $4°C$. The supernatant was discarded,

and sample pellets were washed, resuspended, and then strained twice with 40uM strainers to

remove any clumped cells or debris. The nuclei were then stained with 4',6-diamidino-2-

phenylindole (DAPI) and flowed through a fluorescence activated cell sorter (FACS) (BD

FACSCanto, BD Biosciences, Franklin Lakes, NJ, USA) to assess viability. The nuclei were

sorted according to their expression of DAPI and roughly 400,000 viable nuclei were collected

per group. Following successful collection of the nuclei, the samples were given to the Iowa

State DNA Facility who were responsible for making cDNA libraries using the 10x Chromium

kit (3' RNA library prep) (10X Genomics, Pleasanton, CA, USA), and subsequent sequencing on

an S1 lane using the NovaSeq 6000 (Illumina, San Diego, CA, USA).

2.5 snRNA-seq Data Analysis

After sequencing, Raw FASTQ files were aligned to the *Rattus norvegicus* genome using

the CellRanger pipeline (10X Genomics, Pleasanton, CA, USA). In total, 900,107,250 reads

were sequenced from an estimated 44,629 nuclei. Across pools, the median number of Unique

Molecular Identifiers (UMIs) sampled per nucleus ranged from 1,877-2,197, and the median

number of genes sampled per nucleus ranged from 1,200-1,336. Filtered feature matrices

generated by CellRanger were further pre-processed to 1) minimize risk of analyzing dead or

dying nuclei by excluding barcodes associated with extremely few RNA molecules (<750) and

genes (<500), 2) minimize risk of analyzing multiplets by excluding barcodes associated with an

extremely large number of RNA molecules (>20,000) and genes (>5,000) suggesting multiple

nuclei were sampled in the same droplet, 3) excluding barcodes associated with extremely high

mitochondrial gene content (>5% of all genes). Individual rats were recovered from each pooled

sample using the demultiplexing algorithm Souporcell[57]. Souporcell operates without the need

for a reference genome for each individual by identifying putative individual-specific single-

nucleotide polymorphisms (SNPs) associated with droplet-specific barcodes, and then using

these SNPs to separate barcodes predicted to come from distinct individuals. Additionally,

Souporcell uses these SNPs to estimate a probability that each droplet-associated barcode reflects

multiple nuclei derived from genetically distinct individuals. As an additional quality control

measure, we excluded those barcodes that were predicted by Souporcell to come from multiple

individuals.

2.6 Clustering and Identification of Microglial Cluster

A data-driven clustering approach was used to dimensionally reduce the data and assign

nuclei to communities based on transcriptomic similarity in R using Seurat [58]. First, we

normalized the data and identified variable genes using the 'SCTransform' function from the

sctransform package in R. For dimensionality reduction, we performed Principal Components

Analysis using the 'RunPCA' function (with dim=50) and Uniform Manifold Approximation

(UMAP) using the 'RunUMAP' function (dims=1:50, min.dist=0.5, spread=0.2, n.neighbors=50,

n.epochs=1000, metric="euclidean"); both functions can be found in the Seurat package. Lastly,

nearest neighbors were identified using the 'FindNeighbors' function (reduction="pca",

k.param=50, dims=1:50, n.trees=500, prune.SNN=0) and clusters were assigned using the

'FindClusters' function (resolution=0.2, algorithm=2) for Seurat.

This pipeline resulted in 17 clusters. To aid in cluster identification, each parent cluster

was assigned a number (0-16) which can be visualized in UMAP space (**Fig. 2A**). The microglia

cluster was identified using genes specific to microglia (*P2ry12*, *Aif1*, *Itgam*, and *Cx3cr1*; see

Fig. 2B). Re-clustering of microglia followed the same general workflow as above. Since the

number of subclusters within the microglia cluster was not known *a priori*, this number of

clusters was estimated using a non-parametric graph-based approach[59]. The approach involves

constructing a k-minimum spanning tree (K-MST), which embeds nuclei into a similarity graph

structure such that an edge exists between nuclei if they have similar gene expression

information. Since different cluster resolutions lead to different clustering assignments, we

estimated the number of clusters that achieves the greatest separation of cell clusters across

different resolutions. To achieve this, we calculated the standardized within-cluster edge-count

for a grid of resolution values. Cluster resolutions that result in well-separated communities of

nuclei will lead to larger within-cluster edge-counts. It follows that the number of clusters can be

determined using the resolution that maximizes the within-cluster edge-count. This graph-based

approach circumvents the curse-of-dimensionality that is common in distance-based approaches

to estimate the number of clusters, such as the gap statistic[60].

2.7 Differential Gene Expression Analysis

For the microglia cluster, gene expression was compared between treatment groups and

as a function of running distance using a negative binomial mixed effects model to accommodate

for the multiple nuclei measured per individual. This was implemented using the 'glmer'

package in R. The response variable was the number of transcripts of a gene per nucleus. Group

(categorical variable) or running distance (continuous variable) was specified as a fixed effect

and individual as a random effect. Dispersions were estimated using a Gamma-Poisson model

within the 'glmGamPoi' R package[61]. The natural logarithm of the library sizes per nucleus was

entered as an offset term to normalize variation in sequencing depth across the nuclei. To correct

for multiple testing, we defined significance with an FDR cutoff of 0.05 using Storey's q-value

$(q<0.05)$[62]. To determine differentially expressed genes between specific subclusters, the

Wilcoxon Rank-Sum approach, which requires no distributional assumptions, was used within

Seurat's 'FindMarkers' function. In addition to the above, we also analyzed gene expression a

different way by taking the average gene expression using the SCtransform function (which

corrects for library size) across all nuclei within individuals[63]. This metric was used for the

mediation analyses, where we evaluated evidence for potential causal relations between stress,

gene expression, and running levels using the 'mediation' R package[64].

2.7 Gene Ontology

To test for overrepresentation of biological processes in differentially expressed genes

(DEGs), we used the clusterProfiler package in R [65,66]. GO terms were considered significant by

Fisher's Exact Test if FDR-corrected $p<0.05$. In the dot plots generated, GeneRatio is the ratio of

input genes that are annotated in a term, count is the number of genes that belong to a given gene

set, and the activated versus suppressed pathways are based on the normalized enrichment score

(NES). NES is an indicator of the degree of enrichment. A positive NES indicates that members

of the gene set tend to appear disproportionately at the top of a ranked input gene list, supporting

pathway activation, and a negative NES indicates the opposite and is often considered an

indication of pathway suppression [65,66].

2.8 Microglia Activation and Metabolism Scores

To determine changes in states of activation and function within our groups, a score was

generated for the pro-inflammatory microglia state of activation (M1), the anti-inflammatory

state of activation (M2), and microglial metabolism. First, we used sets of genes associated with

M1, M2, and metabolic microglial states outlined in Ruan et al. (2023). The lists of these genes

are in **Supplemental Table 12**. We then generated M1 and M2 scores by summing the

expression (raw number of transcripts) of the predefined M1-associated and M2-associated

genes, respectively, for each nucleus. Similarly, a metabolism score was calculated based on the

sum of expression levels of key metabolic genes. These scores were normalized by the total

transcript counts per nucleus to account for variations in sequencing depth. Statistical analyses,

including linear mixed-effects models, were applied to examine the relationship between these

scores, treatment groups, and microglia subclusters.

2.9 Cellular maturity

To estimate the relative age of microglial subclusters, CytoTRACE (Cellular (Cyto)

Trajectory Reconstruction Analysis using gene Counts and Expression) was used[68]. CytoTRACE

is an algorithm designed to predict the differentiation states of cells based on single-cell RNA

sequencing data. The core principle of CytoTRACE is that less differentiated (immature) cells

exhibit higher transcriptional diversity, whereas more differentiated (mature) cells show more

specific and lower transcriptional activity. The algorithm operates through several key steps: it

begins by normalizing the gene count matrix obtained from single-cell RNA sequencing data to

ensure that variations in sequencing depth across cells do not bias the results. It then identifies

genes with high variability across nuclei, as these are indicative of differentiation status. For each

nucleus, CytoTRACE calculates a measure of transcriptional diversity, which involves

computing the number of unique genes expressed and the expression levels of those genes[68].

Based on this transcriptional diversity, CytoTRACE assigns a score between 0 and 1 to each cell,

with a score of 1 indicating a cell that is predicted to be highly mature with broad gene

expression patterns and a score of 0 indicating a cell that is predicted to be fully mature with

more specialized gene expression patterns[68]. The CytoTRACE scores for the microglia were then

analyzed using a linear mixed model, where the cluster (or subcluster) was treated as the fixed

effect and individual was treated as the random effect. This statistical approach allows for the

evaluation of cluster-specific effects while accounting for variability among individuals,

facilitating estimation of differentiation states across clusters while controlling for potential

confounding factors related to individual differences.

2.10 Nuclei Proportions

Differences in the relative proportions of microglial subclusters among groups were analyzed

using a Chi-square test and Fisher's Exact Test for pairwise comparisons. Significant group

differences were defined using a significance threshold of alpha=0.05.

3   Results

3.1 Acute stress persistently impairs running behavior

The control rats that had access to running wheels (not stressed; **Fig. 1A**) displayed

typical running patterns for Sprague Dawley males, including daily running distances of

approximately 1-7 km that increased over the course of the experiment (42 days)[69–71] (**Fig. 1B**).

This increase over time was supported by analysis of running distances across days, which

revealed a main effect of day on running distance ($F_{(41, 410)}=10.4$, $p<0.001$). Analysis of total

running distances between groups over the course of the experiment revealed a significant main

effect of stress on running distance ($F_{(1, 10)}=24.4$, $p<0.001$), a pattern that was driven by

stressed rats running significantly shorter distances than rats that were not stressed (Fig. 1B).

Notably, the difference in running distance between stressed and non-stressed animals increased

across the days (stress x day interaction, ($F_{(41, 410)}=6.5$, $p<0.0001$). When collapsing running

distances across days, stressed animals ran on average 14.9% as far as non-stressed animals (**Fig.**

1B). Results for body mass and food intake for all rats are reported in Supplementary Results and

shown in **Supplementary Figure 1A-B**.

Figure 1. Persistent decreases in voluntary wheel running behavior are evident six weeks

after an inescapable and acute stressor. (**A**) Male rats were split into four groups (n=6 per

group) that received either one bout of inescapable stress or not and were then given access to a

running wheel or not. Six weeks later, the striatum was dissected for single-nucleus RNA-

sequencing (snRNAseq) (created with BioRender.com). (**B**) Stressed rats ran significantly less

than non-stressed rats over the entire 6-week period (marked by ***; $p < 0.001$).

3.2 Identification of microglia clusters

The clustering algorithm identified 17 clusters based on single nucleus transcriptomes

(**Fig. 2A**). To determine if specific clusters contained nuclei derived from microglia, we

examined expression patterns of several canonical microglial marker genes (*P2ry12*, *Aif1*, *Itgam*,

and *Cx3cr1*) across clusters (**Fig. 2B**). All four genes showed strong and selective expression in

just three clusters (clusters 2, 13, and 14). Analysis of marker genes in each cluster using the

Seurat 'FindAllMarkers' (using a Wilcoxon Rank Sum test) function showed strong expression

of canonical markers of microglia and macrophages within these three clusters as well.

To more closely examine these clusters, we curated lists of microglia-specific (*P2ry12,*

P2ry13, Cd83, C1qa, Trem2, Aif1, Itgam, Plxdc2, Cx3cr1, Tmem119, Slc2a5, Fcer1g, Sirpa,

Selplg, Gpr84, Spi1, Sall1, Hexb, Cst3, Csf1r, Pros1, Slco2b1, Mrc2) and macrophage-specific

marker genes (*Msr1, Clec4e, Slfn4, Itgal, Ptprc, Adgre1, Mrc1, Cd44, Siglec1, Cd274, Fcgr1a,*

Cd68, Ms4a7, Pf4, F13a1) (**Supp. Fig. 2**). Clusters 2 and 14 showed strong and selective

expression of microglia-specific marker genes (**Supp. Fig. 2A**), whereas cluster 13 showed

strong and selective expression of macrophage-specific marker genes (**Supp. Fig. 2B**).

Additionally, we plotted the same set of microglia-specific genes as well as macrophage-specific

genes. This analysis again revealed a clear pattern whereby cluster 13 showed strong and

selective expression of macrophage-specific genes while clusters 2 and 14 showed strong and

selective expression of microglia-specific genes (**Fig. 2C**).

Based on these gene expression patterns and presence of these marker genes within each

cluster, we therefore considered two clusters (clusters 2 and 14) as containing nuclei derived

from microglia. To better understand the biological differences between these two microglial

clusters transcriptomic signatures of cell maturity were investigated using CytoTRACE, a

bioinformatics tool that assigns a differentiation score to each nucleus based on transcriptional

diversity, with a greater score indicating a less differentiated nucleus (see Methods section 2.8).

One cluster (cluster 14) showed a significantly greater CytoTRACE score compared to the other

(cluster 2; β = .47, SE= .013, p<.0001), supporting the idea that it consisted of relatively less

differentiated microglia while the other cluster (cluster 2) consisted of relatively more mature

microglia (**Fig. 2D**) (**Supp. Table 1**). Thus, for the remainder of this paper we focused our

analysis on the relatively mature microglia from cluster 2.

Figure 2. Identification and characterization of striatal microglia subtypes. (**A**) In total, 41,985 striatal nuclei were split into 17 distinct clusters based on their transcriptomes. (**B**) UMAP plots showing strong and selective expression of key microglia-related genes identified clusters 2, 13, and 14. (**C**) A dot plot showing divergent expression patterns microglia-specific and macrophage-specific genes in microglial clusters 2 and 14 versus macrophagic cluster 13. (**D**) CytoTRACE scores supported cluster 14 being composed almost exclusively of relatively immature microglia and cluster 2 containing a more diverse composition of intermediate

maturity scores (marked by ***; p<0.001). The bars represent means of all the nuclei collapsed

across all individuals, and standard errors reflect variation among nuclei not individuals.

3.3 Striatal microglial signatures of stress

To investigate signatures of stress in striatal microglia, gene expression was compared

between the non-stressed sedentary group (no stress control) and the stressed sedentary group six

weeks after acute stress when running deficits are still present (**Fig. 1B**). In microglial cluster 2,

24 stress-related differentially expressed genes (stress-DEGs) were identified, of which 3 were

upregulated in the non-stressed animals while the remaining 21 were upregulated in the stressed

animals (**Fig. 3A**) (**Supp. Table 2**). A chi-square goodness-of-fit test revealed that the relative

proportions of upregulated versus downregulated stress-DEGs in this cluster to be significantly

different from 50:50, i.e., significantly more genes were upregulated versus downregulated in

association with stress ($\chi_1^2 = 13.5$, p < 0.001). Several stress-DEGs that were upregulated in

association with stress have previously been shown to reflect microglia activation, including

Cst3, Apoe, C1qa, and *C1qb* [72–77] (**Fig. 3B**).

To further evaluate differences between the non-stressed and stressed rat, parent cluster 2

(i.e., relatively mature microglia cluster) was re-clustered and showed three distinct sub-clusters

(**Fig. 3C**). CytoTRACE analysis showed that re-clustering differed strongly in predicted cellular

maturity. Based on CytoTRACE scores, subcluster 0 was the least differentiated, followed by

subcluster 2 and then subcluster 1 (**Fig. 3D**) (**Supp. Table 3**). Thus, for the sake of clarity, we

hereafter refer to subclusters 0, 2, and 1 as putatively immature, intermediate, and mature,

respectively.

Significant stress-DEGs were also found when analyzing each of the 3 subclusters of

microglia separately. Despite the reduced sample size of approximately 1/3 the number of nuclei

per subcluster, 1 stress-DEG was detected in the immature cluster (**Supp. Table 4**), 4 in the

mature subcluster (**Supp. Table 5**), and 4 in the intermediate subcluster (**Supp. Table 6**). All

these genes were contained within the 24 stress-DEGs when considering all subclusters together.

Overall, these findings suggest that stress changes gene expression in all subtypes of striatal

microglia, though the majority of the effect is in the mature and intermediate subclusters.

The influence of stress on microglia maturation phenotypes was investigated next by

comparing relative proportions of putatively immature, intermediate, and mature microglia

differed between no stressed and stressed rats. Stressed rats displayed greater relative proportions

of mature microglia and lesser relative proportions of immature microglia compared to the no

stress rats ($\chi^2_2 = 113.88$, p < .0001; pairwise differences indicated stress differed from no stress

in the relative proportions of immature and mature subclusters, p<0.05, but not in the relative

proportions of the intermediate subcluster; **Fig. 3E & 3F**).

To investigate the possible relationship between stress, microglia maturity, and activation

patterns, the expression of genes involved in glial activation and metabolism was scored for each

nucleus. M1 is considered a canonical pro-inflammatory pattern of activation that microglia can

adopt while M2 is a canonical anti-inflammatory pattern adopted by microglia[78]. Furthermore,

studies have shown that microglia polarization to an M1 phenotype is linked to metabolic shifts

from oxidative phosphorylation to aerobic glycolysis, influencing their functional activity

thereby highlighting the importance of generating a metabolism score[78]. However, the M1, M2,

and metabolism scores did not significantly differ among the putatively immature, intermediate,

and mature microglial subclusters. Subsequent analysis further revealed that none of these scores

differed between the stress and no stress rats (**Supp. Fig. 3A-C**). Taken together, these findings

suggest that canonical M1/M2 signatures do not account for the major differences observed

between groups when the data are restricted to non-runners.

Figure 3. Acute stress produces prolonged striatal microglial activation. (**A**) A comparison

of gene expression between the no stress and stress groups revealed 24 stress-DEGs. The

majority stress-DEGs were upregulated in the stress group compared to the no stress group. (**B**)

Mean gene expression (SCTransformed) of stress-DEGs related to microglial activation are

shown. Individual means and standard errors (among nuclei) are shown for all individuals in the

no stress and stress groups. (**C**) Re-clustering of microglia revealed three subclusters: immature

(pink), mature (green), and intermediate (blue). (**D**) These subclusters differed in predicted levels

of cellular maturity estimated by CytoTRACE. (**E**) The relative proportions of each subcluster

are shown for each individual. (**F**) Individuals in the stress group displayed lesser proportions of

relatively immature microglia and greater proportions of relatively mature microglia compared to

individuals in the no stress group (marked with *; p<0.05).

3.4 Striatal microglial signatures of exercise enrichment

We next investigated the impact of exercise enrichment on striatal microglia gene

expression. Exercise enrichment, defined as access to a running wheel (independent of how

much the animal runs), was first studied in the absence of stress (i.e., this analysis was done only

on the no stress sedentary and runner groups). For the purposes of this analysis, we define the no

stress runner group as the 'exercise enrichment' group. We found 201 significant differentially

expressed genes between the no stress sedentary and exercise enrichment groups (exercise

enrichment-DEGs). A disproportionate number of exercise enrichment-DEGs were upregulated

in the sedentary group (n=158) versus the exercise enrichment group (n=43; chi-square

goodness-of-fit test, $\chi_1^2 = 65.796$, p < 0.001) (**Fig. 4A**) (**Supp. Table 18**). Examples of exercise-

enrichment DEGs that were upregulated in the sedentary group and exercise-enrichment DEGs

that were upregulated in the exercise enrichment group are shown in **Figure 4B**. Specific GO

terms that were significantly over-represented among exercise enrichment-DEGs included:

cellular respiration/oxidative phosphorylation (e.g., *Atp7a, Cox5a*), immune response (e.g.,

Cd74, Tnf), synaptic signaling (e.g., *Camk2a, Sorcs3*), and neurogenesis (e.g., *Fgf13, Nrp1*) (**Fig.**

4C) (**Supp. Table 20**). These results are consistent with exercise enrichment shifting striatal

microglia toward functions related to synaptic signaling and remodeling via phagocytosis of

neuronal processes.

To further analyze the effects of exercise enrichment on microglia, parent cluster 2 was

re-clustered containing only the exercise enrichment and sedentary no stress groups. This

analysis revealed the same three subclusters derived in Fig. 3C, along with a fourth distinct

subcluster (**Fig. 4D**). The majority of the cells in this fourth subcluster were from the exercise

enrichment group (**Fig. 4D**) (**Supp. Fig. 4**). Indeed, within the exercise enrichment group, 14.4%

of striatal microglia (in parent cluster 2) were assigned to the exercise enrichment subcluster,

whereas only 1.9% of striatal microglia were assigned to this subcluster in sedentary rats.

CytoTRACE suggested that the remaining three subclusters were distinguished by relative

maturity, mirroring the pattern observed in Fig 3D. Thus, these subclusters are again referred to

as the immature, intermediate, and mature subclusters. The fourth subcluster is referred to as the

'exercise enrichment' subcluster hereafter for the sake of clarity. Notably, the exercise

enrichment subcluster and the intermediate subcluster displayed similar CytoTRACE scores,

supporting that they were both composed of relatively intermediate microglia (**Fig. 4E**) (**Supp.**

Table 8).

The relative proportions of the four microglial subclusters differed between exercise

enriched and sedentary groups ($\chi^2_3 = 157.62$, p < .0001) (**Fig. 4F & 4G**). The greatest difference

was for the exercise enrichment subcluster, as described above, where the proportion of this

subtype of microglia was 7.6-fold higher in runners than the sedentary group. The exercise

enrichment group also had significantly fewer intermediate and immature microglia, and

significantly more mature microglia compared to the sedentary rats (Fig. 4F & 4G) (**Fig. 4F &**

4G). Differential gene expression and gene set enrichment analyses further revealed that the

genes upregulated in the exercise enrichment subcluster relative to the other subclusters were

generally related to microglia activation/immune response (*Cst3, Marcks, Sparc, Cd83, Dab2,*

Tmem176a, Tmem176b, Clec4a1, and Cd74), the complement system (*C1qa, C1qb, C3*), and

microglia phagocytosis (*Ctss, Pltp, RT1-DMa*) (**Fig. 4H**). Lastly, differences among subclusters

were analyzed for M1, M2, and metabolism scores described above (**Supp. Table 12**)[67]. Similar

to our earlier analyses, no significant differences between exercise enrichment and sedentary

groups or among subclusters were observed (**Supp. Fig. 3D-F**). Taken together, these results

suggest that canonical signatures of microglia activation do not sufficiently describe subtypes of

striatal microglia or their responses to exercise enrichment in our paradigm.

Figure 4. Exercise enrichment is associated with a specific striatal microglial activation profile. (**A**) A comparison of gene expression between the sedentary and exercise-enrichment groups found a total of 201 differentially expressed genes, or exercise enrichment-DEGs, the vast majority of which were upregulated in the sedentary group relative to the exercise enrichment group (**B**) Mean gene expression of exemplar genes upregulated in the sedentary and exercise enrichment groups are shown. Individual means and standard errors (among nuclei) are shown for all individuals in the sedentary and exercise enrichment groups. (**C**) Dot plot

highlighting GO terms that were significantly enriched among exercise enrichment-DEGs. (**D**)

Re-clustering identified four distinct subclusters of microglia in the sedentary and exercise

enrichment groups, including a fourth subcluster that was abundant in exercise enriched but not

sedentary rats. (**E**) CytoTRACE analysis revealed a similar maturity structure as the stress vs. no

stress analysis (subcluster 0 = immature, subcluster 1 = mature, and subcluster 2 = intermediate),

with the additional exercise enrichment subcluster showing a relative maturity level similar to

the intermediate subcluster. (**F**) Proportions of nuclei assigned to each subcluster are shown for

each of the individuals. For legend see panel I. (**G**) Individuals in the exercise enrichment group

displayed a significantly lower proportion of relatively immature microglia, a greater proportion

of relatively mature microglia, and an even greater proportion of microglia assigned to the

exercise enrichment subcluster, which was nearly absent in sedentary rats (marked with *,

$p<0.05$). (**H**) Comparison of the exercise enrichment subcluster to the other subclusters revealed

a distinct set of genes related to microglial activation, phagocytosis, and the complement system.

(**I**) Expression patterns of example genes that were upregulated in the exercise enrichment

subcluster.

3.5 Striatal microglial signatures of individual running levels

To further investigate the regulation of gene changes by wheel running, we analyzed gene

expression associated with voluntary wheel running distances in non-stressed rats across multiple

timescales relative to when the striatum was sampled: the previous 40 days, the previous day, the

previous three hours, the period 30-90 minutes before sampling, and the 30-minute period

immediately before sampling (**Supp. Table 9**; **Supp. Tables 13-16, respectively**). Strikingly,

807 genes showed expression associated with total distance run over the entire 6-week period (6-

week distance-DEGs; $q<0.05$), but far fewer distance-DEGs were observed when analyzing total

running distances over the final 24 hours (n=58 distance-DEGs), 3 hours (n=20 distance-DEGs),

30-90 minutes (n=20 distance-DEGs), and 30 minutes (n=9 distance-DEGs). Notably, the most

of the distance-DEGs identified in analyses of these shorter time intervals were also identified in

our analyses of the entire 6-week period, likely due to the positive correlations between 6-week

running distances 6-weekand running distances over the smaller intervals (Pearson's R was 0.47,

0.14, 0.1, and 0.03 for the correlation between running distances over 40 days versus running

distances over 24 hours, 3 hours, 30-90 min, and 30 min respectively). Interestingly, there was

little overlap between running level-DEGs and exercise enrichment-DEGs as indicated by

hypergeometric test (p=0.66), suggesting that different genes are regulated by running levels

versus running wheel access in striatal microglia (**Supp. Fig. 5A, example relationships**

between distance-DEG expression and running distance shown in Supp. Fig. 5B). Taken

together, these results support that striatal microglial gene expression most strongly reflects long-

term as opposed to recent levels of running.

3.6 Striatal microglial signatures of interactions between stress and exercise

To investigate possible interactions of acute stress and exercise on striatal microglial gene

expression, we next analyzed all four experimental groups using a 2x2 design with stress and

exercise as main effects as well as a stress by exercise interaction term. This analysis revealed

207 exercise-DEGs (main effect of exercise), 201 stress-DEGs (main effect of stress), and 17

interaction-DEGs (**Supp. Table 10, example interaction-DEGs are shown in Fig. 5A**). Stress-

DEGs and exercise enrichment-DEGs in this analysis overlapped significantly with the stress-

DEGs and exercise enrichment-DEGs identified earlier when stress and exercise were analyzed

separately (**Supp. Fig. 6A & 6B**; hypergeometric tests, both p<0.05). There was no significant

overlap between stress-DEGs and exercise-DEGs from either analysis (hypergeometric test on

stress-DEGs and exercise-DEGs from analyses using a 2x2 model and analyses of each factor

separately (**Fig. 5B & 5C**). Taken together, these results suggest that stress and exercise have

long-term and robust effects on expression of largely distinct genes in striatal microglia.

534   We additionally ran an analysis of covariance (ANCOVA) to investigate how stress and

long-term running distance may interact to influence striatal microglial gene expression

determine (**Supp. Table 17**). This analysis included only animals housed with a running wheel,

both stressed and not stressed. Results revealed 401 6-week distance-DEGs, 379 stress-DEGs,

and 355 significant stress-by-distance interaction-DEGs. Distance-DEGs identified in the

ANCOVA overlapped significantly with distance-DEGs identified in earlier analyses of the non-

stressed runners only (hypergeometric test, $p < 0.001$; **Supp. Fig. 5D**). Interestingly, stress-

DEGs identified in earlier analyses of stress versus no-stress (all sedentary) (**Supp. Table 2**) did

not significantly overlap with stress-DEGs identified in the ANCOVA ($p > 0.05$) (**Supp. Fig. 6C**),

a pattern that was likely driven by differences in the effects of stress on striatal microgial gene

expression between sedentary and exercising rats. Examples of genes that showed strong effects

of the running covariate, a strong effect of stress, and no interaction are shown in

Supplementary Figure 6E & 6F. Examples of genes that showed a strong interaction between

the running covariate and stress are shown in **Supplementary Figure 6G**. Taken together, these

results support an extraordinary degree of sensitivity to complex interactions between stress and

exercise in striatal microglia.

550   Genes that showed a significant effect of the running covariate, main effect of stress or

interaction in the ANCOVA analyses described above were further analyzed by mediation

analysis. The purpose was to identify candidate genes involved in causing the reduced running

behavior and other genes that are more likely a consequence of the different running levels rather

than a cause. This analysis found 12 genes whose expression levels were significant mediators of

the effects of stress on running levels, i.e., putative genes that cause varying running levels. The

analysis also found 16 genes whose expression levels were mediated by different running levels,

i.e., putative genes that change in response to stress as a consequence of reduced running (**Supp.**

Table 21) (**Supp. Fig. 9**).

Lastly, we extended our analyses of microglial subclusters by re-clustering parent cluster

2 across all four groups. This analysis yielded three distinct subclusters (**Fig. 5D**). Mirroring our

analyses of running rats only, one subcluster (subcluster 2) consisted of nuclei derived

predominantly from runners (**Fig. 5E & Fig. 5F**). Analysis of cell maturity using CytoTRACE

supported that subcluster 1 was relatively immature whereas subclusters 0 and 2 (the exercise

enrichment subcluster) were relatively and comparably mature (**Fig. 5G**) (**Supp. Table 11**). The

proportion of nuclei in each subcluster differed strongly among treatment groups in an overall

Chi-square test (χ_6^2 = 383.59, p < 0.0001). Within each subcluster, all pairwise comparisons

between treatment groups were significant as indicated by Fisher Exact Test (**Fig. 5E**). To

investigate whether these subclusters differed in additional biological dimensions, we analyzed

the same M1, M2, and metabolism scores described above (**Supp. Table 12**). This analysis

revealed differences among subclusters: compared to the immature subcluster 1, the mature

subcluster 0 displayed greater metabolism scores and decreased expression of both M1 and M2

genes, whereas the exercise-enrichment subcluster 2 displayed decreased expression of M1 genes

(**Fig. 5H**). Within subclusters, no significant differences between treatment groups were

observed for the M1, M2, or metabolism scores. Taken together, these results further reinforce

the conclusion that the canonical M1/M2/metabolism metrics are not sufficient to differentiate

the treatment groups, though with the full data set, the subclusters can be differentiated. The fact

that the exercise-enrichment subcluster displays a higher M2 score than the relatively mature

subcluster is indicative of a protective microglial functional state.

Figure 5. Striatal microglial signatures of stress-by-exercise interactions. (**A**) Analysis of all

four groups yielded 207 significant exercise-DEGs (main effect of exercise), 201 stress-DEGs

(main effect of stress), and 17 interaction-DEGs. Mean normalized gene expression for four

exemplar interaction-DEGs are plotted. Individual means and standard errors (among nuclei) are

shown. (**B**) Stress-DEGs and exercise enrichment-DEGs did not significantly overlap (results

shown reflect main effects of stress and exercise in the 2x2 analysis). (**C**) Similarly, stress-DEGs

and exercise enrichment-DEGs from the separate analyses did not significantly overlap. (**D**) Re-

clustering of microglia from all treatment groups resulted in three distinct subclusters, including

a subcluster that is only apparent in runners (subcluster 2). (**E**) Proportions of nuclei in each

subcluster are shown for each treatment group. (**F**) The runner subcluster accounted for a large

percentage of microglia in all individuals that had access to running wheels (both stress and no

stress), regardless of running levels, but was almost entirely absent in individuals without

running wheels. (**G**) Subcluster 1 was relatively more immature than the other clusters (marked

by ***; p<0.001). (**H**) The subclusters were differentiated based on genes related to metabolism,

pro- (M1), or anti-inflammatory (M2) activation. Specifically, the runner subcluster showed a

lower metabolism score and higher M2 score compared to the relatively mature subcluster 1.

4   Discussion

The aim of this study was to define the molecular signatures of stress, exercise, and their

interaction in striatal microglia. Using snRNA-seq, we showed stress-associated differences in

estimated microglial maturity 42 days post-stressor as well as molecular signatures of a

distinctive and non-canonical activation pattern, together underscoring the enduring impacts of

stress on microglial dynamics. Access to a running wheel was also associated with profound

alterations in striatal microglia, including greater relative proportions of a distinct microglial

subpopulation that was nearly absent in sedentary rats. In this subpopulation, exercise

enrichment was associated with increased expression of genes related to complement system-

and phagocytosis-related pathways, indicative of a separate, distinct form of microglial

activation. Even more striking, we observed distinct microglial gene expression signatures

associated with access to a running wheel (i.e., exercise enrichment) compared to individual

running distances. Lastly, we observed evidence for a distinct form of activation in response to

both stress and exercise, potentially reflecting molecular processes that contribute to deficits in

motivation for physical activity. Our findings thus support a broader spectrum of microglial

functional states than previously appreciated and reveal candidate molecular pathways by which

striatal microglia may regulate motivation for exercise following stressors.

We observed that a single session of inescapable foot shock was associated with severely

depressed levels of voluntary wheel running behavior for 42 days after the stressor (**Fig. 1B**).

This extends previous literature where reduced voluntary running was observed up to 28 days

after the stressor[11,12]. We hypothesized that microglia-induced neuroinflammation and/or priming

of microglia in the striatum may contribute to the reduced motivation for running given the role

of microglia in both stress and exercise (as reviewed in [38,79]). Consistent with our hypothesis,

several genes that have been implicated in microglia activation were upregulated in the stress

group: *Cst3, Apoe, Srsf7, Rpl19, C1qa,* and *C1qb* (some of which are plotted in **Fig. 3B**). *Cst3*

and *Apoe* were documented previously as markers of activated microglia[72], disease-associated

microglia[72,73], or injury-associated microglia[74,75]. In contrast, *Srsf7* and *Rpl19* have been found to

be downregulated in microglia in Alzheimer's disease[80] and are associated with homeostatic

microglia[81]. Both *C1qa* and *C1qb* are involved in the complement system which are indicators of

microglia activation as well [76,77].

In addition to displaying differential expression of specific markers of microglia

activation as described above, stress was also associated with greater estimated microglial

maturity (**Fig. 3D & 3E**), a pattern that may reflect a specific form of inflammatory action. Ruan

et al. (2023) used CytoTRACE in their experiment examining the role of leucine-rich alpha-2-

glycoprotein 1 (Lrg1) in cerebral ischemia□reperfusion injury and similarly found that different

clusters of microglia had different states of maturation which they attributed to their

inflammatory action. On the other hand, we did not see evidence of canonical markers of stress-

induced inflammation such as *Il1β*, *Il6*, or *Tnfα* even though transcripts for these genes were

detected in our dataset. Similarly, while we were able to detect expression of many genes that

are considered markers of M1, M2, and metabolism states of microglia, these genes were not

informative for distinguishing subclusters or effects of stress in sedentary rats (**Supp. Fig. 1A-F**).

Together, this supports a unique form of enduring microglial activation following acute stress

that is not captured by classic definitions of microglia activation (i.e., M1 and M2). This is

consistent with previous work showing enduring responses of hippocampal and cortical

microglia to inescapable shock stress[31,32,35,82]. Given the dual role of inescapable shock stress in

producing both microglial activation and reduced motivation for exercise, it is possible that the

reduced running we see in the present study is a result of the sustained striatal microglial

activation following inescapable shock stress. Results of the mediation analysis support this idea

and identify several candidate genes in microglia that may be partially responsible for regulating

running behavior in response to stress (**Supp. Table 21**) (**Supp. Fig. 9**). Future studies are

needed to determine if blocking striatal microglia activation ameliorates deficits in motivation

for physical activity following exposure to a severe, acute stressor.

In addition to exhibiting molecular signatures of stress, microglia also exhibited

molecular signatures of physical activity. Individuals with running wheel access that were not

stressed ran every day throughout the experiment, and total running distance was associated with

expression levels of 807 genes (**Supp. Table 9**). These effects were unlikely driven by acute

effects of running, because more than 10-fold fewer running-level DEGs were observed when

analyzing shorter intervals immediately prior to sampling compared to the total distance run over

previous 40 days. This supports the idea that the cumulative effect of running over multiple days

or weeks results in a robust functional shift in a subset of microglia, perhaps to support striatal

function during running behavior.

In addition to running-level DEGs, we also detected a separate set of non-overlapping

genes associated with running wheel access (**Supp. Fig. 4A**), supporting distinct microglial

profiles associated with running wheel access versus levels of running. Access to a running

wheel is considered a form of enrichment [83–85]. Thus, we believe the DEGs identified in the

comparison of runners to sedentary rats (no stress) reflect the enrichment of the running wheel,

or the deprivation of no running wheel in the sedentary group, rather than differential levels of

physical activity *per se*. Strikingly, we observed an 'exercise enrichment' subcluster that is only

found in rats with access to a running wheel, whether stressed (and thus, running very low levels)

or not (**Fig. 5D-F**). The genes distinguishing this subcluster from other microglial subclusters

were generally related to microglial activation/immune response (*Cst3, Marcks, Sparc, Cd83,*

Dab2, Tmem176a, Tmem176b, Clec4a1, and Cd74) the complement system (*C1qa, C1qb, C3*),

and microglial phagocytosis (*Ctss, Pltp, RT1-DMa*) (**Fig. 4D**). Together, these results are

consistent with microglia being activated and phagocytosing in a complement-dependent manner

more in rats with voluntary wheel running enrichment compared to those deprived of the

enrichment.

Activated microglia can eliminate both damaged cells and dysfunctional synapses, a

phenomenon known as "synaptic stripping" [76,86–89]. C1q and C3, integral components of the

classical complement cascade, attach to the surface of invading pathogens. Subsequently,

phagocytes expressing target complement receptors in the peripheral immune system eliminate

these pathogens [90,91]. Similarly, within the CNS, microglia utilize classical complement cascades

to facilitate phagocytic signaling, leading to the removal of excess synapses [76,77]. Few studies

have evaluated microglial complement and phagocytosis during physical exercise, but persistent

aerobic exercise has the potential to preserve the integrity and functionality of neurovascular

units in aging mice[92]. Exercise achieves this by diminishing C1q-expressing microglia and

enhancing neuroplasticity[92]. Moreover, physical exercise appears to facilitate cellular

communication between microglia and neurons through complement molecules[93]. Thus, the

emergence of a unique C1q+ and C3+ microglial subpopulation with exercise may be involved

in neuron-microglia and/or astrocyte-microglia crosstalk.

In addition to uncovering striatal microglial signatures of traumatic stress and separately,

voluntary wheel running, we were also interested in exploring potential overlap in signatures of

running and stress on microglia. We found that stress-DEGs in rats with running wheel access

did not significantly overlap stress-DEGs in rats without running wheel access (**Supp. Fig. 5C**);

however, we identified a suite of eight genes (*Apoe, C1qa, Cst3, Grm3, Mt.co1, Pde4b, Rpl19,*

and *Tmem176b*) that was differentially expressed in response to both stress and exercise.

Notably, these genes are almost all involved in microglial activation (**Supp. Fig. 6**). Based on

these data, we hypothesize that microglia are generally activated but exhibit different subtypes of

responses to stress depending on exercise enrichment; however, future studies are needed to test

this hypothesis.

We investigated microglial metabolism, pro-inflammatory activation (M1), and anti-

inflammatory activation (M2) by tracking established transcriptional markers of these functional

states. When all four experimental groups were analyzed together in a 2x2 design, microglia

clustered into three subclusters, the exercise enrichment subcluster (more abundant in runners

regardless of stress condition), relatively mature subcluster (more abundant in stressed versus

non-stressed rats), and relatively immature subcluster. The exercise enrichment subcluster

showed an increased M2 score compared to the relatively mature subcluster (**Fig. 5E**), further

supporting distinct microglial signatures of stress and exercise, and supporting the idea that

exercise promotes a shift toward anti-inflammatory activation. However, future research should

validate the role of microglia activation, specifically the anti-inflammatory activation, in stress-

induced decreases in running behavior.

In conclusion, our work demonstrates that a single traumatic stressor decreases voluntary

wheel running and induces signatures of striatal microglial activation six weeks later. Further, we

establish that striatal microglia display distinct molecular signatures associated with acute stress,

exercise enrichment, and long-term running distances. Notably, the observed signature of

microglia activation long after an acute stressor, coupled with the emergence of a unique

exercise-associated subcluster displaying signatures of activation (M2) and phagocytosis,

underscores the complex and multidimensional nature of microglial function. Both the unique

functional profile of the microglia long after stress along with the exercise enrichment subcluster

further support that microglial functional diversity extends beyond traditional M1 and M2

categories. The specific signature of changes in activation six weeks post-stressor also highlights

the temporal complexities of microglial responses and may reflect unique forms of long-term

priming, though future studies are needed to determine if priming has taken place. Future

research should also investigate microglial functional diversity and causally manipulate striatal

microglia to determine their functional role in modulating neural circuits and motivational

deficits associated with stress.

CRediT Author Contributions:

Meghan G. Connolly: Conceptualization, Methodology, Formal Analysis, Software,

Investigation, Writing – Original Draft, Writing – Review & Editing, Visualization, Project

Administration. **Zachary V. Johnson:** Methodology, Formal Analysis, Software, Investigation,

Data Curation, Writing – Review & Editing. **Lynna Chu:** Methodology, Formal Analysis,

Software, Investigation, Data Curation, Writing – Review & Editing. **Nicholas D. Johnson:**

Methodology, Formal Analysis, Software, Writing – Review & Editing. **Trevor J. Buhr:**

Investigation. **Elizabeth M. McNeill:** Resources, Methodology. **Peter J. Clark:**

Conceptualization, Investigation, Methodology, Resources, Writing – Review & Editing,

Funding Acquisition. **Justin S. Rhodes:** Conceptualization, Methodology, Formal Analysis,

Software, Investigation, Writing – Original Draft, Writing – Review & Editing, Supervision.

Conflict of Interest:

The authors declare no conflicts of interest.

Funding:

Iowa State University of Science and Technology Healthy Iowa Interdisciplinary Research Team

Initiative

References:

1.   Meiser-Stedman R, McKinnon A, Dixon C, Boyle A, Smith P, Dalgleish T. Acute stress
disorder and the transition to posttraumatic stress disorder in children and adolescents:
Prevalence, course, prognosis, diagnostic suitability, and risk markers. *Depress Anxiety*.
2017;34(4). doi:10.1002/da.22602

2.   Geoffrion S, Goncalves J, Robichaud I, et al. Systematic Review and Meta-Analysis on
Acute Stress Disorder: Rates Following Different Types of Traumatic Events. *Trauma
Violence Abuse*. 2022;23(1). doi:10.1177/1524838020933844

3.   Fleshner M, Greenwood BN, Yirmiya R. Neuronal-Glial Mechanisms of Exercise-Evoked
Stress Robustness. In: ; 2014:1-12. doi:10.1007/7854_2014_277

4.   Nowacka-Chmielewska M, Grabowska K, Grabowski M, Meybohm P, Burek M, Małecki
A. Running from Stress: Neurobiological Mechanisms of Exercise-Induced Stress
Resilience. *Int J Mol Sci*. 2022;23(21):13348. doi:10.3390/ijms232113348

5.   Zen AL, Whooley MA, Zhao S, Cohen BE. Post-traumatic stress disorder is associated
with poor health behaviors: Findings from the Heart and Soul Study. *Health Psychology*.
2012;31(2):194-201. doi:10.1037/a0025989

6.   de Assis MA, de Mello MF, Scorza FA, et al. Evaluation of Physical Activity Habits in
Patients with Posttraumatic Stress Disorder. *Clinics*. 2008;63(4):473-478.
doi:10.1590/S1807-59322008000400010

7.   Kagan J, Berkun M. The reward value of running activity. *J Comp Physiol Psychol*.
1954;47(2):108-108. doi:10.1037/h0058877

8.   Iversen IH. TECHNIQUES FOR ESTABLISHING SCHEDULES WITH WHEEL
RUNNING AS REINFORCEMENT IN RATS. *J Exp Anal Behav*. 1993;60(1):219-238.
doi:10.1901/jeab.1993.60-219

9.   Collier G, Hirsch E. Reinforcing properties of spontaneous activity in the rat. *J Comp
Physiol Psychol*. 1971;77(1):155-160. doi:10.1037/h0031588

10.   Belke TW. RUNNING AND RESPONDING REINFORCED BY THE OPPORTUNITY
TO RUN: EFFECT OF REINFORCER DURATION. *J Exp Anal Behav*. 1997;67(3):337-
351. doi:10.1901/jeab.1997.67-337

11. Buhr TJ, Reed CH, Wee OM, et al. The persistence of stress-induced physical inactivity in
rats: an investigation of central monoamine neurotransmitters and skeletal muscle
oxidative stress. *Front Behav Neurosci*. 2023;17. doi:10.3389/fnbeh.2023.1169151

12. Moraska A, Fleshner M. Voluntary physical activity prevents stress-induced behavioral
depression and anti-KLH antibody suppression. *American Journal of Physiology-*
Regulatory, Integrative and Comparative Physiology. 2001;281(2):R484-R489.
doi:10.1152/ajpregu.2001.281.2.R484

13. Weiss JM, Goodman PA, Losito BG, Corrigan S, Charry JM, Bailey WH. Behavioral
depression produced by an uncontrollable stressor: Relationship to norepinephrine,
dopamine, and serotonin levels in various regions of rat brain. *Brain Res Rev*.
1981;3(2):167-205. doi:10.1016/0165-0173(81)90005-9

14. Maier SF, Watkins LR. Stressor controllability and learned helplessness: The roles of the
dorsal raphe nucleus, serotonin, and corticotropin-releasing factor. *Neurosci Biobehav*
Rev. 2005;29(4-5):829-841. doi:10.1016/j.neubiorev.2005.03.021

15. Strong PV, Christianson JP, Loughridge AB, et al. 5-hydroxytryptamine 2C receptors in
the dorsal striatum mediate stress-induced interference with negatively reinforced
instrumental escape behavior. *Neuroscience*. 2011;197:132-144.
doi:10.1016/j.neuroscience.2011.09.041

16. Rhodes JS, Garland T, Gammie SC. Patterns of Brain Activity Associated With Variation
in Voluntary Wheel-Running Behavior. *Behavioral Neuroscience*. 2003;117(6):1243-
1256. doi:10.1037/0735-7044.117.6.1243

17. Durieux PF, Schiffmann SN, de Kerchove d'Exaerde A. Differential regulation of motor
control and response to dopaminergic drugs by D1R and D2R neurons in distinct dorsal
striatum subregions. *EMBO J*. 2012;31(3):640-653. doi:10.1038/emboj.2011.400

18. Durieux PF, Bearzatto B, Guiducci S, et al. D2R striatopallidal neurons inhibit both
locomotor and drug reward processes. *Nat Neurosci*. 2009;12(4):393-395.
doi:10.1038/nn.2286

19. Bateup HS, Santini E, Shen W, et al. Distinct subclasses of medium spiny neurons
differentially regulate striatal motor behaviors. *Proceedings of the National Academy of*
Sciences. 2010;107(33):14845-14850. doi:10.1073/pnas.1009874107

20. Cox J, Witten IB. Striatal circuits for reward learning and decision-making. *Nat Rev*
Neurosci. 2019;20(8):482-494. doi:10.1038/s41583-019-0189-2

21.   Balleine BW, Delgado MR, Hikosaka O. The Role of the Dorsal Striatum in Reward and
804   Decision-Making: Figure 1. *The Journal of Neuroscience*. 2007;27(31):8161-8165.
805   doi:10.1523/JNEUROSCI.1554-07.2007

806   22.   Saul MC, Majdak P, Perez S, Reilly M, Garland T, Rhodes JS. High motivation for
807   exercise is associated with altered chromatin regulators of monoamine receptor gene
808   expression in the striatum of selectively bred mice. *Genes Brain Behav*. 2017;16(3):328-
809   341. doi:10.1111/gbb.12347

810   23.   Rodrigues D, Jacinto L, Falcão M, et al. Chronic stress causes striatal disinhibition
811   mediated by SOM-interneurons in male mice. *Nat Commun*. 2022;13(1):7355.
812   doi:10.1038/s41467-022-35028-4

813   24.   Clark PJ, Amat J, McConnell SO, et al. Running Reduces Uncontrollable Stress-Evoked
814   Serotonin and Potentiates Stress-Evoked Dopamine Concentrations in the Rat Dorsal
815   Striatum. *PLoS One*. 2015;10(11):e0141898. doi:10.1371/journal.pone.0141898

816   25.   Robinson OJ, Overstreet C, Charney DR, Vytal K, Grillon C. Stress increases aversive
817   prediction error signal in the ventral striatum. *Proceedings of the National Academy of
818   Sciences*. 2013;110(10):4129-4133. doi:10.1073/pnas.1213923110

819   26.   Woodburn SC, Bollinger JL, Wohleb ES. The semantics of microglia activation:
820   neuroinflammation, homeostasis, and stress. *J Neuroinflammation*. 2021;18(1):258.
821   doi:10.1186/s12974-021-02309-6

822   27.   Felger JC, Treadway MT. Inflammation Effects on Motivation and Motor Activity: Role
823   of Dopamine. *Neuropsychopharmacology*. 2017;42(1):216-241.
824   doi:10.1038/npp.2016.143

825   28.   Vichaya EG, Hunt SC, Dantzer R. Lipopolysaccharide Reduces Incentive Motivation
826   While Boosting Preference for High Reward in Mice. *Neuropsychopharmacology*.
827   2014;39(12):2884-2890. doi:10.1038/npp.2014.141

828   29.   Gellner AK, Sitter A, Rackiewicz M, et al. Stress vulnerability shapes disruption of motor
829   cortical neuroplasticity. *Transl Psychiatry*. 2022;12(1):91. doi:10.1038/s41398-022-
830   01855-8

831   30.   Cheng Y, Pardo M, Armini R de S, et al. Stress-induced neuroinflammation is mediated
832   by GSK3-dependent TLR4 signaling that promotes susceptibility to depression-like
833   behavior. *Brain Behav Immun*. 2016;53:207-222. doi:10.1016/j.bbi.2015.12.012

834   31.   Weber MD, Frank MG, Tracey KJ, Watkins LR, Maier SF. Stress induces the danger-
835   associated molecular pattern HMGB-1 in the hippocampus of male sprague dawley rats: A

priming stimulus of microglia and the NLRP3 inflammasome. *Journal of Neuroscience*.
2015;35(1):316-324. doi:10.1523/JNEUROSCI.3561-14.2015

32. Frank MG, Baratta M V., Sprunger DB, Watkins LR, Maier SF. Microglia serve as a
neuroimmune substrate for stress-induced potentiation of CNS pro-inflammatory cytokine
responses. *Brain Behav Immun*. 2007;21(1):47-59. doi:10.1016/j.bbi.2006.03.005

33. Niraula A, Sheridan JF, Godbout JP. Microglia Priming with Aging and Stress.
Neuropsychopharmacology. 2017;42(1):318-333. doi:10.1038/npp.2016.185

34. Frank MG, Weber MD, Watkins LR, Maier SF. Stress-induced neuroinflammatory
priming: A liability factor in the etiology of psychiatric disorders. *Neurobiol Stress*.
2016;4:62-70. doi:10.1016/j.ynstr.2015.12.004

35. Frank MG, Fonken LK, Watkins LR, Maier SF. Acute stress induces chronic
neuroinflammatory, microglial and behavioral priming: A role for potentiated NLRP3
inflammasome activation. *Brain Behav Immun*. 2020;89. doi:10.1016/j.bbi.2020.05.063

36. Fonken LK, Weber MD, Daut RA, et al. Stress-induced neuroinflammatory priming is
time of day dependent. *Psychoneuroendocrinology*. 2016;66:82-90.
doi:10.1016/j.psyneuen.2016.01.006

37. Fonken LK, Frank MG, Gaudet AD, Maier SF. Stress and aging act through common
mechanisms to elicit neuroinflammatory priming. *Brain Behav Immun*. 2018;73:133-148.
doi:10.1016/j.bbi.2018.07.012

38. Frank MG, Fonken LK, Watkins LR, Maier SF. Microglia: Neuroimmune-sensors of
stress. *Semin Cell Dev Biol*. 2019;94:176-185. doi:10.1016/j.semcdb.2019.01.001

39. Marinelli S, Basilico B, Marrone MC, Ragozzino D. Microglia-neuron crosstalk:
Signaling mechanism and control of synaptic transmission. *Semin Cell Dev Biol*.
2019;94:138-151. doi:10.1016/j.semcdb.2019.05.017

40. Badimon A, Strasburger HJ, Ayata P, et al. Negative feedback control of neuronal activity
by microglia. *Nature*. 2020;586(7829):417-423. doi:10.1038/s41586-020-2777-8

41. Baratta M V., Maier SF. New tools for understanding coping and resilience. *Neurosci Lett*.
2019;693:54-57. doi:10.1016/j.neulet.2017.09.049

42. Fleshner M, Crane CR. Exosomes, DAMPs and miRNA: Features of Stress Physiology
and Immune Homeostasis. *Trends Immunol*. 2017;38(10):768-776.
doi:10.1016/j.it.2017.08.002

43. Maier SF, Seligman MEP. Learned helplessness at fifty: Insights from neuroscience.
Psychol Rev. 2016;123(4):349-367. doi:10.1037/rev0000033

44.   Christianson JP, Greenwood BN. Stress-protective neural circuits: not all roads lead
870   through the prefrontal cortex. *Stress*. 2014;17(1):1-12.
871   doi:10.3109/10253890.2013.794450

872   45.   Frank MG, Watkins LR, Maier SF. Stress-induced glucocorticoids as a neuroendocrine
873   alarm signal of danger. *Brain Behav Immun*. 2013;33:1-6. doi:10.1016/j.bbi.2013.02.004

874   46.   Weiss JM. Effects of coping responses on stress. *J Comp Physiol Psychol*.
875   1968;65(2):251-260. doi:10.1037/h0025562

876   47.   Rhodes JS, Koteja P, Swallow JG, Carter PA, Garland T. Body temperatures of house
877   mice artificially selected for high voluntary wheel-running behavior: repeatability and
878   effect of genetic selection. *J Therm Biol*. 2000;25(5):391-400. doi:10.1016/S0306-
879   4565(99)00112-6

880   48.   Koteja P, Garland T, Sax JK, Swallow JG, Carter PA. Behaviour of house mice artificially
881   selected for high levels of voluntary wheel running. *Anim Behav*. 1999;58(6):1307-1318.
882   doi:10.1006/anbe.1999.1270

883   49.   Reed CH, Bauer EE, Shoeman A, Buhr TJ, Clark PJ. Acute Stress Exposure Alters Food-
884   Related Brain Monoaminergic Profiles in a Rat Model of Anorexia. *J Nutr*.
885   2021;151(12):3617-3627. doi:10.1093/jn/nxab298

886   50.   Clark PJ, Ghasem PR, Mika A, et al. Wheel running alters patterns of uncontrollable
887   stress-induced cfos mRNA expression in rat dorsal striatum direct and indirect pathways:
888   A possible role for plasticity in adenosine receptors. *Behavioural Brain Research*.
889   2014;272:252-263. doi:10.1016/j.bbr.2014.07.006

890   51.   Strong PV, Christianson JP, Loughridge AB, et al. 5-hydroxytryptamine 2C receptors in
891   the dorsal striatum mediate stress-induced interference with negatively reinforced
892   instrumental escape behavior. *Neuroscience*. 2011;197:132-144.
893   doi:10.1016/j.neuroscience.2011.09.041

894   52.   Greenwood BN, Foley TE, Burhans D, Maier SF, Fleshner M. The consequences of
895   uncontrollable stress are sensitive to duration of prior wheel running. *Brain Res*.
896   2005;1033(2):164-178. doi:10.1016/j.brainres.2004.11.037

897   53.   Greenwood BN, Foley TE, Day HEW, et al. Freewheel Running Prevents Learned
898   Helplessness/Behavioral Depression: Role of Dorsal Raphe Serotonergic Neurons. *The
899   Journal of Neuroscience*. 2003;23(7):2889-2898. doi:10.1523/JNEUROSCI.23-07-
900   02889.2003

901   54.   Bauer EE, Shoeman A, Buhr TJ, Daniels KM, Lyte M, Clark PJ. Voluntary
902   binge☐patterned alcohol drinking and sex☐specific influences on monoamine☐related

neurochemical signatures in the mouse gut and brain. *Alcohol Clin Exp Res*.
2021;45(5):996-1012. doi:10.1111/acer.14592

55.   Buhr TJ, Reed CH, Shoeman A, Bauer EE, Valentine RJ, Clark PJ. The Influence of
Moderate Physical Activity on Brain Monoaminergic Responses to Binge-Patterned
Alcohol Ingestion in Female Mice. *Front Behav Neurosci*. 2021;15.
doi:10.3389/fnbeh.2021.639790

56.   Johnson Z V., Hegarty BE, Gruenhagen GW, Lancaster TJ, McGrath PT, Streelman JT.
Cellular profiling of a recently-evolved social behavior in cichlid fishes. *Nat Commun*.
2023;14(1):4891. doi:10.1038/s41467-023-40331-9

57.   Heaton H, Talman AM, Knights A, et al. Souporcell: robust clustering of single-cell
RNA-seq data by genotype without reference genotypes. *Nat Methods*. 2020;17(6):615-
620. doi:10.1038/s41592-020-0820-1

58.   Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell
data. *Cell*. 2021;184(13):3573-3587.e29. doi:10.1016/j.cell.2021.04.048

59.   Bai Y, Chu L. A Graph-based Approach to Estimating the Number of Clusters. *arXiv
preprint*. 2024;arXiv:2402.15600.

60.   Tibshirani R, Walther G, Hastie T. Estimating the Number of Clusters in a Data Set Via
the Gap Statistic. *J R Stat Soc Series B Stat Methodol*. 2001;63(2):411-423.
doi:10.1111/1467-9868.00293

61.   Ahlmann-Eltze C, Huber W. glmGamPoi: fitting Gamma-Poisson generalized linear
models on single cell count data. *Bioinformatics*. 2021;36(24):5701-5702.
doi:10.1093/bioinformatics/btaa1009

62.   Storey JD, Tibshirani R. Statistical significance for genomewide studies. *Proceedings of
the National Academy of Sciences*. 2003;100(16):9440-9445.
doi:10.1073/pnas.1530509100

63.   Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq
data using regularized negative binomial regression. *Genome Biol*. 2019;20(1):296.
doi:10.1186/s13059-019-1874-1

64.   Tingley D, Teppei Yamamoto, Kentaro Hirose, Luke Keele, Kosuke Imai. mediation: R
package for causal mediation analysis. *J Stat Softw*. Published online 2014.

65.   Wu T, Hu E, Xu S, et al. clusterProfiler 4.0: A universal enrichment tool for interpreting
omics data. *Innovation*. 2021;2(3). doi:10.1016/j.xinn.2021.100141

66.   Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R Package for Comparing Biological
936   Themes Among Gene Clusters. *OMICS*. 2012;16(5):284-287. doi:10.1089/omi.2011.0118

937   67.   Ruan Z, Cao G, Qian Y, et al. Single-cell RNA sequencing unveils Lrg1's role in cerebral
938   ischemia☐reperfusion injury by modulating various cells. *J Neuroinflammation*.
939   2023;20(1):285. doi:10.1186/s12974-023-02941-4

940   68.   Gulati GS, Sikandar SS, Wesche DJ, et al. Single-cell transcriptional diversity is a
941   hallmark of developmental potential. *Science (1979)*. 2020;367(6476):405-411.
942   doi:10.1126/science.aax0249

943   69.   Fediuc S, Campbell JE, Riddell MC. Effect of voluntary wheel running on circadian
944   corticosterone release and on HPA axis responsiveness to restraint stress in Sprague-
945   Dawley rats. *J Appl Physiol*. 2006;100(6):1867-1875.
946   doi:10.1152/japplphysiol.01416.2005

947   70.   Droste SK, Chandramohan Y, Hill LE, Linthorst ACE, Reul JMHM. Voluntary Exercise
948   Impacts on the Rat Hypothalamic-Pituitary-Adrenocortical Axis Mainly at the Adrenal
949   Level. *Neuroendocrinology*. 2007;86(1):26-37. doi:10.1159/000104770

950   71.   Lee JR, Parker KE, Tapia M, et al. Voluntary wheel running effects on intra-accumbens
951   opioid high-fat feeding and locomotor behavior in Sprague-Dawley and Wistar rat strains.
952   *Physiol Behav*. 2019;206:67-75. doi:10.1016/j.physbeh.2019.02.015

953   72.   Thrupp N, Sala Frigerio C, Wolfs L, et al. Single-Nucleus RNA-Seq Is Not Suitable for
954   Detection of Microglial Activation Genes in Humans. *Cell Rep*. 2020;32(13):108189.
955   doi:10.1016/j.celrep.2020.108189

956   73.   Mathys H, Davila-Velderrain J, Peng Z, et al. Single-cell transcriptomic analysis of
957   Alzheimer's disease. *Nature*. 2019;570(7761):332-337. doi:10.1038/s41586-019-1195-2

958   74.   Wahane S, Zhou X, Zhou X, et al. Diversified transcriptional responses of myeloid and
959   glial cells in spinal cord injury shaped by HDAC3 activity. *Sci Adv*. 2021;7(9).
960   doi:10.1126/sciadv.abd8811

961   75.   Hammond TR, Dufort C, Dissing-Olesen L, et al. Single-Cell RNA Sequencing of
962   Microglia throughout the Mouse Lifespan and in the Injured Brain Reveals Complex Cell-
963   State Changes. *Immunity*. 2019;50(1):253-271.e6. doi:10.1016/j.immuni.2018.11.004

964   76.   Schafer DP, Lehrman EK, Kautzman AG, et al. Microglia Sculpt Postnatal Neural Circuits
965   in an Activity and Complement-Dependent Manner. *Neuron*. 2012;74(4):691-705.
966   doi:10.1016/j.neuron.2012.03.026

967   77.   Stevens B, Allen NJ, Vazquez LE, et al. The Classical Complement Cascade Mediates
968   CNS Synapse Elimination. *Cell*. 2007;131(6):1164-1178. doi:10.1016/j.cell.2007.10.036

969   78.   Orihuela R, McPherson CA, Harry GJ. Microglial M1/M2 polarization and metabolic
970   states. *Br J Pharmacol*. 2016;173(4):649-665. doi:10.1111/bph.13139

971   79.   Mee-inta, Zhao, Kuo. Physical Exercise Inhibits Inflammation and Microglial Activation.
972   *Cells*. 2019;8(7):691. doi:10.3390/cells8070691

973   80.   Zhou Y, Song WM, Andhey PS, et al. Human and mouse single-nucleus transcriptomics
974   reveal TREM2-dependent and TREM2-independent cellular responses in Alzheimer's
975   disease. *Nat Med*. 2020;26(1):131-142. doi:10.1038/s41591-019-0695-9

976   81.   Li Q, Cheng Z, Zhou L, et al. Developmental Heterogeneity of Microglia and Brain
977   Myeloid Cells Revealed by Deep Single-Cell RNA Sequencing. *Neuron*.
978   2019;101(2):207-223.e10. doi:10.1016/j.neuron.2018.12.006

979   82.   Blandino P, Barnum CJ, Deak T. The involvement of norepinephrine and microglia in
980   hypothalamic and splenic IL-1β responses to stress. *J Neuroimmunol*. 2006;173(1-2):87-
981   95. doi:10.1016/j.jneuroim.2005.11.021

982   83.   Grégoire CA, Bonenfant D, Le Nguyen A, Aumont A, Fernandes KJL. Untangling the
983   Influences of Voluntary Running, Environmental Complexity, Social Housing and Stress
984   on Adult Hippocampal Neurogenesis. *PLoS One*. 2014;9(1):e86237.
985   doi:10.1371/journal.pone.0086237

986   84.   Mustroph ML, Chen S, Desai SC, Cay EB, DeYoung EK, Rhodes JS. Aerobic exercise is
987   the critical variable in an enriched environment that increases hippocampal neurogenesis
988   and water maze learning in male C57BL/6J mice. *Neuroscience*. 2012;219:62-71.
989   doi:10.1016/j.neuroscience.2012.06.007

990   85.   Kobilo T, Liu QR, Gandhi K, Mughal M, Shaham Y, van Praag H. Running is the
991   neurogenic and neurotrophic stimulus in environmental enrichment. *Learning & Memory*.
992   2011;18(9):605-609. doi:10.1101/lm.2283011

993   86.   Lehrman EK, Wilton DK, Litvina EY, et al. CD47 Protects Synapses from Excess
994   Microglia-Mediated Pruning during Development. *Neuron*. 2018;100(1):120-134.e6.
995   doi:10.1016/j.neuron.2018.09.017

996   87.   Scott☐Hewitt N, Perrucci F, Morini R, et al. Local externalization of phosphatidylserine
997   mediates developmental synaptic pruning by microglia. *EMBO J*. 2020;39(16).
998   doi:10.15252/embj.2020105380

999   88.   Paolicelli RC, Bolasco G, Pagani F, et al. Synaptic pruning by microglia is necessary for
1000   normal brain development. *Science (1979)*. 2011;333(6048):1456-1458.
1001   doi:10.1126/science.1202529

1002   89.   Kim HJ, Cho MH, Shim WH, et al. Deficient autophagy in microglia impairs synaptic
1003   pruning and causes social behavioral defects. *Mol Psychiatry*. 2017;22(11):1576-1584.
1004   doi:10.1038/mp.2016.103

1005   90.   Morgan BP, Kavanagh D. Introduction to complement in health and disease: novel aspects
1006   and insights. *Semin Immunopathol*. 2018;40(1):1-2. doi:10.1007/s00281-017-0664-7

1007   91.   Reis ES, Mastellos DC, Hajishengallis G, Lambris JD. New insights into the immune
1008   functions of complement. *Nat Rev Immunol*. 2019;19(8):503-516. doi:10.1038/s41577-
1009   019-0168-x

1010   92.   Soto I, Graham LC, Richter HJ, et al. APOE Stabilization by Exercise Prevents Aging
1011   Neurovascular Dysfunction and Complement Induction. *PLoS Biol*.
1012   2015;13(10):e1002279. doi:10.1371/journal.pbio.1002279

1013   93.   Li F, Geng X, Yun HJ, Haddad Y, Chen Y, Ding Y. Neuroplastic Effect of Exercise
1014   Through Astrocytes Activation and Cellular Crosstalk. *Aging Dis*. 2021;12(7):1644.
1015   doi:10.14336/AD.2021.0325

1016