Interleaved single and bursting spiking resonance in neurons

Cesar C. Ceballos[1], Nourdin Chadly[3,4], Eric Lowet[3], Rodrigo F. O. Pena[1,2, *]

[1] Department of Biological Sciences, Florida Atlantic University, Jupiter, FL 33458, USA
[2] Stiles-Nicholson Brain Institute, Florida Atlantic University, Jupiter, FL 33458, USA
[3] Department of Neuroscience, Erasmus Medical Center, Rotterdam, Netherlands
[4] Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, Netherlands

* Corresponding author
E- mail: penar@fau.edu

Abstract

Under *in vivo* conditions, CA1 pyramidal cells from the hippocampus display transitions from single spikes to bursts. It is believed that subthreshold hyperpolarization and depolarization, also known as down and up-states, play a pivotal role in these transitions. Nevertheless, a central impediment to correlating suprathreshold (spiking) and subthreshold activity has been the technical difficulties of this type of recordings, even with widely used calcium imaging or multielectrode recordings. Recent work using voltage imaging with genetically encoded voltage indicators has been able to correlate spiking patterns with subthreshold activity in a variety of CA1 neurons, and recent computational models have been able to capture these transitions. In this work, we used a computational model of a CA1 pyramidal cell to investigate the role of intrinsic conductances and oscillatory patterns in generating down and up-states and their modulation in the transition from single spiking to bursting. Specifically, the emergence of distinct spiking resonances between these two spiking modes that share the same voltage traces in the presence of theta or gamma oscillatory inputs, a phenomenon we call interleaved single and bursting spiking resonance. We noticed that these resonances do not necessarily overlap in frequency or amplitude, underscoring their relevance for providing flexibility to neural processing. We studied the conductance values of three current types that are thought to be critical for the bursting behavior: persistent sodium current (I_{NaP}) and its conductance G_{NaP}, delayed rectifier potassium (I_{KDR}) and its conductance G_{KDR}, and hyperpolarization-activated current (I_{h}) and its conductance G_{h}. We conclude that the intricate interplay of ionic currents significantly influences the neuronal firing patterns, transitioning from single to burst firing during sustained depolarization. Specifically, the intermediate levels of G_{NaP} and G_{KDR} facilitate spiking resonance at gamma frequency inputs. The resonance characteristics vary between single and burst firing modes, each displaying distinct amplitudes and resonant frequencies. Furthermore, low G_{NaP} and high G_{KDR} values lock bursting to theta frequencies, while high G_{NaP} and low G_{KDR} values lock single spiking to gamma frequencies. Lastly, the duration of quiet intervals plays a crucial role in determining the likelihood of transitioning to either bursting or single spiking modes. We confirmed that the same features were present in previously recorded in vivo voltage-imaging data. Understanding these dynamics provides valuable insights into the fundamental mechanisms underlying neuronal excitability under *in vivo* conditions.

Author summary

Since discovering that neurons in the hippocampus can encode spatial position through phase precession, many experiments have explored how specific theta and gamma oscillations influence location specificity in the brain. However, the individual neuronal properties and dynamics behind these behaviors are still being uncovered. Previously, we found that stereotypical bursting and single-spike firing in pyramidal neurons are linked to these oscillations and further associated with an animal entering or leaving a place field. Advances in voltage-imaging techniques have enabled us to assess these properties more precisely. Our study shows that different frequencies can independently trigger these stereotypical spikes, demonstrating a complex pattern where the same cell can be double-coded: a phenomenon we called interleaved resonance. Additionally, we found that this coding can be modulated by persistent sodium and delayed-rectifier potassium currents. Moreover, these neurons are more likely to burst following long periods of silence. These findings provide new insights into the mechanisms underlying neural coding in the hippocampus and how it relates to behavior.

Introduction

Neurons from the CA1 hippocampus display a diversity of spiking dynamics. In the hippocampus, from the point of view of individual cells, *in vivo* voltage imaging [1, 2, 3], patch clamp recordings [4, 5, 6, 7], and multielectrode recordings [8, 9] reveals that CA1 pyramidal cells exhibit stereotypical single and burst spiking. While investigating their emergence and transitions it was noted that these spike modes appear with different probabilities of occurrence where bursts are more likely to occur after long periods of activity silence [9]. From the point of view of populations, gamma (30-100 Hz) and theta (3-12 Hz) oscillations coexist and have been well characterized in CA1 region of rodent hippocampus [10, 11]. Gamma is modulated by theta during working memory retention, and it is important to notice the cross-frequency coupling of these oscillations for working memory performance [12, 13]. Recent studies have tried to connect those transitions using *in vivo* data, as shown by shifts from single to bursting spikes in the same neurons that are preferentially observed during theta and gamma oscillations, respectively [1]. Behaviorally, these spiking modes have been proposed to encode an animal's position, where burst signals when animals enter a place. In contrast, single spikes signal when the animal leaves a place [1].

Admittedly, understanding spiking patterns transitions is synonymous to understanding ionic current's dynamics. For fixed synaptic input, the composition of ionic currents can produce distinguishable spiking outputs that vary from single to burst of spikes [14, 15, 16]. Combinations of ionic currents can determine these spike modes as in the case of calcium and burst [17, 18, 19], or in combination with the spatial location on the neuron as in the case of persistent sodium current (I_{NaP}) located at or near the soma [14, 20], or only the geometry [21], or combinations of both [22].

Oscillations can cause transitions from low to high firing rate on a neuron. Cells may have amplified response for certain oscillatory frequencies compared to other frequencies. This is known as suprathreshold (spiking) resonance, and it has been observed widely across the brain [23, 24, 25, 26, 27], including CA1 pyramidal cells [28] and the mechanisms underlying this resonance have been intensively investigated using computational models [29, 30]. Similarly, subthreshold resonance can be seen as a transition voltage change of a neuron upon a particular

89   (non-zero) input frequency [31, 32]. Several ionic currents contribute to subthreshold resonance
90   and can be discussed in terms of their chord and derivative conductances [33]. For example,
91   persistent sodium current (I_{NaP}) is an amplifying ionic current and delayed rectifier potassium
92   (I_{KDR}) is a resonant current [31, 34, 35]. Less is known about the role of these ionic currents on
93   suprathreshold (spiking) resonance.
94
It is only natural to assume that CA1 pyramidal neurons embedded in a network can take advantage
of the spiking resonance to preferentially transmit information encoded in the spiking mode (single
or burst spiking) when a particular oscillatory synaptic input (e.g. theta or gamma oscillations) is
received [28, 23, 36, 37]. In addition to that, it is important to understand how transitions at these
different scales connect and modulate them. In this paper, we investigate suprathreshold resonance
of single and burst spiking in a CA1 pyramidal neuron model. We found that spiking resonance
can easily emerge when neurons are stimulated with theta and gamma synaptic inputs and ionic
currents are tuned. Variability of sodium and potassium currents differentially promoted the
presence of resonance according to the firing pattern (single vs. burst), a phenomenon we call
interleaved single and bursting spiking resonance. Our results provide insights that might help to
understand the mechanisms underlying single and burst spiking computation and its role in
connection to *in vivo* processing that is relevant for spike coding in CA1 pyramidal cells for spatial
navigation and memory.
108
Our paper is organized as follows: in the following section, we present studies of how ionic
currents promote transitions from single to burst firing under constant sustained input and later
under oscillatory inputs. For the oscillatory inputs, we show how resonance emerges and depends
on ionic currents. Then we move to a more systematic study where we characterize interleaved
single and bursting resonance in the presence of dendritic I_h current. For the same model, we
characterize phase-locking to gamma and theta and how these main ionic currents shape this
relationship. We conclude by systematically characterizing the probability of bursting spikes in
relation to the periods of silence in the voltage traces. Methods are included in the last section of
the paper.
118

Results

Ionic currents promote a transition from single to burst firing during sustained depolarization

CA1 pyramidal cells are known to exhibit transitions from single (SS) to bursting spikes (BS) *in*
vivo [1, 2, 3, 38]. This transition occurs in the presence of subthreshold oscillations and is
modulated by oscillatory inputs, mostly theta and gamma oscillations. Subthreshold activity is
characterized by transitions from hyperpolarized states to depolarized states (also known as down
and up-states, respectively). Firing mostly occurs on top of the depolarization phases. Input
patterns play a relevant role in the switching, but it is unclear how they can be promoted by the
neuron's intrinsic properties, e.g. ionic currents. We first decided to investigate the firing patterns
occurring during depolarization states and how SS and BS emerge under different ionic
conductances.
132

We first investigated the role of conductances that are mostly expressed in the soma, such as I_{NaP} and I_{KDR}. We used a single compartment computational model of CA1 pyramidal cell (see Methods) and we varied G_{NaP} (persistent sodium conductance) and G_{KDR} (delayed rectifier potassium conductance) for the model receiving a pulse of current that maintains the voltage in a constant depolarized state (Fig 1A). We observed that increasing G_{NaP} makes the firing transits from SS to BS and finally to plateau potential, also known as depolarization block [39] (see voltage traces Fig 1B). The phase plots (dV/dt) in the same figure reveals how the voltage series transits between the different modes. In contrast, the effect of increasing G_{KDR} is a transition from BS to SS (Fig 1C). Single spiking can be explained as a result of having long pauses between spikes due to long afterhyperpolarization, which is facilitated by strong action potential repolarization due to large G_{KDR} or by preventing fast depolarization due to small G_{NaP}.

Fig 1. Ionic currents promote transition from SS to BS during sustained depolarization.
Default values for fixed conductance while varying the other are $G_{\text{NaP}} = 0.3$ and $G_{\text{KDR}} = 6$. **A** Sketch of CA1 model receiving constant current pulse. **B** Example voltage traces and phase plots (dV/dt) when varying G_{NaP} or **C** G_{KDR}. **D** Firing rate and coefficient of variation (CV) heatmaps spanned over G_{NaP} and **E** G_{KDR} vs. Input current.

The firing rate increases with the increase of stimulation amplitude, increase of G_{NaP} and decrease of G_{KDR} (Fig 1D-E). The coefficient of variation (CV) above 2 (reddish to yellowish) determines the area in which the combination of conductance promotes BS behavior (Fig 1D-E). As it is shown in the heatmaps, BS is independent of the input amplitude and mostly occurs in areas around G_{NaP} = 0.35 or G_{KDR} = 6. These values correspond to the edges where the transition from spiking to depolarization block occurs (see dark regions in heatmaps of firing rate). Interestingly, BS occurs in a very narrow range of conductances, meanwhile single spiking is likely to occur in a wide range of values. This set of observations shows the opposite effects of the G_{NaP} and G_{KDR} ionic currents in modulating a switch between SS to BS.

Intermediate G_{NaP} and G_{KDR} generate spiking resonance preferentially at gamma frequency input

We showed previously that G_{NaP} and G_{KDR} have opposite effects on the generated spiking patterns during sustained depolarization. Nevertheless, CA1 pyramidal neurons receive oscillatory inputs in the range from 3 to 100 Hz, where oscillatory gamma is between 30-100 Hz and theta 3-12 Hz frequencies. This induces transitions from hyperpolarized to depolarized states where shorter intervals occur at high input frequencies such as gamma frequencies and longer intervals at theta frequencies. Under this condition, firing during depolarization is not sustained, but rather is alternated by periods of silence. In our paper, we investigate the role of the silent states generated by the interplay between input patterns and ionic currents to create a suprathreshold resonance and its relation to SS and BS.

For stimulation of the cell in this section, we choose varying oscillatory stimulation frequencies covering the range from theta to gamma frequencies (Fig 2A). We investigate the firing rate responses and their preference for a certain input frequency (i.e. suprathreshold resonance). We observe low-pass filtering of the firing rate for low G_{NaP} or high G_{KDR}, suprathreshold resonance for intermediate G_{NaP} or G_{KDR}, and a nearly absent frequency preference for high values of G_{NaP} or low values of G_{KDR} (Fig 2C, E). Interestingly, regardless of the opposite effects of G_{NaP} and

G_{KDR}, we observed that theta input has a stronger effect on the firing rate response than gamma
when the neuron acts as a low-pass filter. In contrast, gamma is stronger than theta for resonance
(Fig 2C, E). These results unveiled an interaction between the input frequency and the filtering
properties that determine the firing output.

Fig 2. Intermediate G_{NaP} and G_{KDR} generates firing rate resonance. Default values for fixed
conductance while varying the other are $G_{NaP} = 0.3$ and $G_{KDR} = 6$. **A** Sketch of simplified CA1
model receiving oscillatory current stimulation. **B** Examples of voltage traces for varying values
of G_{NaP}. **C** Firing rate while varying stimulation frequency of an oscillatory input. Shaded red area:
theta frequencies. Shaded blue area: gamma frequencies. **D** Examples of voltage traces for varying
values of G_{KDR}. **E** Firing rate while varying stimulation frequency of an oscillatory input. **F** Firing
rate and CV heatmap spanned over G_{NaP} and oscillatory input frequency. **G** Firing rate and CV
heatmap spanned over G_{KDR} and oscillatory input frequency.

The CVs above 2 (reddish to yellowish) obtained while changing the oscillatory input frequency
and the conductance indicate the area where BS occurs. The heatmaps show similar patterns for
firing frequency and CV for both G_{NaP} and G_{KDR}, i.e. we observe high firing rates simultaneously
with high values of CV. The high firing rates obtained can be justified by the BS and low firing
rates are by the SS (Fig 2F, G).

We observed BS when G_{NaP} levels were intermediate or high (see the yellowish diagonal in Fig.
2F) and when G_{KDR} was at intermediate levels (Fig. 2G). Additionally, BS was noted at theta
frequencies with low G_{NaP} or high G_{KDR}. Similarly, we observed SS at low G_{KDR} and at gamma
frequencies with low or intermediate G_{NaP} or high G_{KDR}. Interestingly, BS in gamma frequencies
occurs simultaneously with resonance at intermediate values of G_{NaP} or G_{KDR}. These results
indicate that when suprathreshold resonance is present, a linear increase in input frequency and
G_{NaP} shifts the BS presence from theta to gamma frequencies. Similarly, a decrease in G_{KDR}, along
with increased input frequencies, also shifts the BS presence from theta to gamma frequencies (as
shown by the yellowish regions in the heatmaps). A similar pattern is observed for SS, with the
dark regions in the G_{NaP} heatmap and the dark middle-right region in the G_{KDR} heatmap illustrating
this shift.

In summary, for low G_{NaP} and high G_{KDR}, we found high firing rate response to low input
frequencies (i.e. theta) due to weak filtering, and low firing rate response to high input frequencies
(i.e. gamma) due to strong filtering. This suggests that the tuning of ionic currents in interaction
with input frequencies are critical for determining the occurrence of resonance.

Single and burst firing display suprathreshold resonance with
different amplitudes and different resonant frequencies

It is well known that CA1 pyramidal cells have a high expression of the hyperpolarization activated
I_h current [40]. This current is mostly expressed in the dendrites and plays a relevant role in
modulating synaptic inputs [40, 41, 42, 43] and resonance [33]. It prevents excessive
hyperpolarization and causes rebound spiking after hyperpolarizations [44, 45]. Thus, in this
section, we incorporated a dendritic compartment containing I_h current (Fig 3A).

Figure 3. Single and burst firing display suprathreshold resonance with different amplitudes and resonant frequencies. Default values for fixed conductance while varying the other are G_{NaP} = 0.93, G_{KDR} = 5.86, and G_h = 8.46. **A** Sketch of CA1 model with additional dendritic compartment receiving oscillatory current stimulation. **B** Exemplary voltage trace of the simulation with SS and BS indicated. **C1-2** Firing rate vs oscillatory input frequency for SS **C1** and BS **C2** when G_{NaP} values are 0, 0.5 and 1.5. **D1-2** Same as in **C**, but G_{KDR} values are 0, 5 and 10. **E1-2** Same as in **C**, but G_h values are 0, 2 and 5. **F** Representative sketch demonstrating how Δf and ΔA are calculated for the interleaved SS and BS resonances. **G1-2** Δf and ΔA for varying values of G_{NaP}. **H1-2** Same as in **G**, but varying G_{KDR}. **I1-2** Same as in **G**, but varying G_h. Colors mark the magnitude of ionic currents (blue = low, orange = medium, green = high).

We stimulated the cell model with varying oscillatory stimulation frequencies covering the range from theta to gamma frequencies (Fig 3A). Under this condition, the neuron is capable of switching its firing from SS to BS (Fig 3B). We separated our quantitative measures into SS and BS (see Methods). Overall, we observed resonance for all the explored conductances and for both spiking types (SS and BS), except for BS when G_{NaP} is large or when G_{KDR} is low (Fig 3 C2, D2). Finally, SS and BS have resonance for all G_h values (Fig 3 E1, E2).

We further compared the resonance frequencies and their respective firing rate amplitudes between SS and BS (Fig 3F). For low G_{NaP}, SS resonance frequency is larger than BS resonance frequency, but it is the opposite when G_{NaP} is large (Fig 3G1). For low G_{KDR}, SS resonance frequency is smaller than BS resonance frequency, but it is the opposite when G_{KDR} is large (Fig 3H1). Moreover, SS resonance frequency is larger than BS resonance frequency for all G_h values (Fig 3I1). Finally, as expected, firing frequency at the resonant peak was always higher for BS rather than SS (Fig 3G2, H2, I2).

The set of results in this section demonstrates an interesting coding behind SS and BS in our CA1 model. First, it shows that neither the resonance frequencies nor resonance amplitudes between SS and BS necessarily overlap. More interestingly, this phenomenon happens at the same voltage traces (interleaved) and can be modulated by ionic currents. In the next section, we show how these observations are connected to oscillations.

G_{NaP}, G_{KDR} and G_h determine phase locking of SS and BS to theta and gamma input respectively

It has been demonstrated that this neuron model can reproduce coexisting SS and BS phase-locked to gamma and theta oscillations, respectively. The model is derived from [1, 14] and has its conductances fitted to experimental voltage-imaging. In this paper, we modified the model for a similar fitting (see parameters in Methods). Based on the observations above for changing ionic conductances and the theta and gamma relationship to BS and SS, we question the ability of the ionic channels to modulate the phase-locking of the neuron model to these frequencies.

Since most of CA1 neuron inputs come with a mixture of theta and gamma frequencies, it is only natural to study the case where both are present. Thus, we injected both theta and gamma simultaneously into the neuron model (Fig 4A). As observed, the ionic currents embedded in the neuron can generate different phase-locking (Fig 4B). Low values of G_{NaP} are more prompt to BS at theta frequencies. Increasing the values of G_{NaP} removes this phase locking with the generation

of more SS (Fig 4C). In contrast, G_{KDR} had the opposite behavior (Fig 4D). We see only SS for
low values of G_{KDR}, BS at gamma for intermediate values of G_{KDR} and finally BS at theta is
generated for high G_{KDR}. Interestingly, only low and high values of G_h promoted a difference at
the gamma frequencies (Fig 4E).

Fig. 4. G_{NaP}, G_{KDR} and G_h determine phase locking statistic of SS and BS to theta and gamma
input. Default values for fixed conductance while varying the other are $G_{NaP} = 0.93$, $G_{KDR} = 5.86$,
and $G_h = 8.46$. **A** Sketch of CA1 model with somatic and dendritic compartments receiving theta
and gamma oscillatory current stimulation. **B** Examples of voltage traces with SS behavior
transitioning to a BS behavior (compare top to bottom). **C** Phase locking statistic to theta and
gamma input separated into SS and BS with varying G_{NaP}. **D** Phase locking statistic with varying
G_{KDR}. **E** Phase locking statistic with varying G_h. Each bar is the result of 10 different simulations.
t-test with 0.05 level of significance (asterisk).

Taken together, these results show that G_{NaP}, G_{KDR}, and G_h modulate the coexistence of BS and
SS phase locked to theta and gamma differently: low values of G_{NaP} and high values of G_{KDR} lock
BS to theta frequencies. G_h has little effect on theta frequencies and affects gamma frequencies
instead. Furthermore, high values of G_{NaP} and low values of G_{KDR} lock SS to gamma frequencies.

Quiet intervals determine the probability of bursting or single spiking
The *in vivo* activity of CA1 neurons is characterized by subthreshold oscillations that follows the
theta and gamma synaptic inputs. During these oscillations, spiking activity is observed at the
depolarization phase (up-states) and between them it is possible to observe silence intervals at
hyperpolarized phase (down-states). During the up-states, neurons can fire SS or BS. In this
section, we investigate whether the silence intervals modulate the probability of having SS or BS.

According to our observations in Fig 5, high probability of BS occurs for long intervals of silence
(above 100 ms) in spite of variation in the values of the conductances G_{NaP}, G_{KDR}, and G_h (Fig
5D1-F1). Interestingly, we observed that interevent intervals (ISI) are fractured and distributed in
more than one position in the diagram, and that different peaks show up mostly for large G_{NaP} or
low G_{KDR} (Fig 5A and B). We also observed that high probability of SS occurs for short intervals
of silence (below 60 ms) for low G_{NaP}, high G_{KDR} and regardless of the values of G_h (Fig 5D2-F2).
Moreover, we observed high probability of SS occurring for long intervals (above 100 ms) only
when G_{NaP} is high or when G_{KDR} is low (Fig 5D2-E2). In summary, we found that the probability
of BS is robust to variation of the ionic conductances and mostly depends on long silent periods
above 100 ms (i.e. when theta input is present). In contrast, SS is influenced by ionic conductances
and can occur at short or long intervals.

Figure 5. Quiet intervals determine burst and single spikes. Default values for fixed
conductance while varying the other are $G_{NaP} = 0.93$, $G_{KDR} = 5.86$, and $G_h = 8.46$. **A** Histogram of
interevent intervals (ISI) with varying G_{NaP}. **B** Histogram of interevent intervals (ISI) with varying
G_{KDR}. **C** Histogram of interevent intervals (ISI) with varying G_h. **D1-F1** BS and **D2-F2** SS
probabilities for varying G_{NaP}, G_{KDR}, and G_h, respectively. Colors mark the magnitude of ionic
currents (green = low, red = medium, blue = high).

To determine whether the spiking probability is the sum of independent effects of theta and gamma stimulation, or the result of the interaction between both inputs, we calculated the spiking probability when only theta or gamma inputs were used (Fig 6). As expected, when only gamma input is present, we observed that the probability of BS or SS is always larger at short intervals of silence (< 50 ms) (Fig 6A). In contrast, when only theta input is present, we observed that the probability of BS is larger for long intervals of silence (> 100 ms). For SS probability when only theta input is present, we observed the same tendency found above, where high probability of SS occurs for short intervals of silence below 60 ms for low G_{NaP}, high G_{KDR} and regardless the values of G_h (Fig 6B), and high probability of SS occurs for long intervals above 100 ms only when G_{NaP} is high or when G_{KDR} is low.

Figure 6. Burst probability follows the longest quiet intervals according to the slowest oscillatory input. Default values for fixed conductance while varying the other are $G_{NaP} = 0.93$, $G_{KDR} = 5.86$, and $G_h = 8.46$. **A** BS and SS probabilities with only gamma input. **B** BS and SS probabilities with only theta input. Colors mark the magnitude of ionic currents (green = low, red = medium, blue = high).

In summary, we observed that when the input is only at gamma frequencies, the spiking probability is largest for short silent intervals for both SS and BS. Opposite to that, when the input is only oscillating at theta, BS is more likely to occur at long intervals and SS at short intervals. Interestingly, when both theta and gamma are present, the probabilities behave as if only theta was present, i.e., the gamma component is mostly disregarded. These results suggest that BS probability follows the input frequency with the longest intervals of silence, i.e. it follows theta even in the presence of gamma, but it follows gamma in the absence of any other slowest input. Meanwhile, SS would preferentially occur at short intervals.

Voltage imaging burst probability follows the longest quiet intervals

Despite the difficulties in reconciling our computational results with voltage-imaging recordings, where frequency-dependent stimulation is typically absent, it is possible to compare the bursting probability following the longest quiet intervals with spontaneous recordings. Thus, we analyzed voltage imaging traces of 7 neurons (20 s long) obtained from a public repository [3] . Fluorescence images were collected from CA1 pyramidal cells and somatostatin expressing interneurons located in the dorsal CA1 area at depths ~50 to 100 μm (Fig 7A). LED epifluorescence illumination allowed the collection of the voltage indicator fluorescence by a high-speed sCMOS camera at a rate acquisition of 600 Hz. Photobleaching was subtracted by fitting a single exponential decay. We selected the traces from pyramidal cells based on their high signal-to-noise ratio and high frequency of events suggesting good quality of recordings (Fig 7B). These criteria allowed us to properly detect and classify events as SS and BS as done in the computational simulations (Fig 7C). We calculated the probability of finding SS or BS for a certain ISI (Fig 7D). We performed a statistical comparison between BS and SS probabilities for 3 areas marked for low ISI (50 to 150 ms), intermediate ISI (150 to 250 ms), and long ISI (400 to 500 ms).

Figure 7. Voltage imaging burst probability follows the longest quiet intervals. Data from [3]. **A** Sketch of the voltage-imaging experiment. **B** Sample traces that were used for this analysis (right, n = 7). **C** Sample trace showing spike detection algorithm prior to division into BS and SS.

361 **D** Interevent interval (ISI) dependent probabilities of BS and SS (n = 7), and 3 areas marked for
362 low ISI (50 to 150 ms), intermediate ISI (150 to 250 ms), and long ISI (400 to 500 ms). **E** From
363 the marked areas, bar plots with averaged values in each ISI trace (n = 7). t-test with 0.05 level of
364 significance (asterisk).
365
The results showed that SS is more likely at low ISIs, while BS is more likely at long ISIs.
Observations of these probabilities across all traces (Fig. 7D) and statistical significance from the
traces and areas (Fig. 7E) confirm this. The voltage-imaging recordings agree with our simulation
observations, where burst probability follows the longest quiet intervals (Figs 5 and 6). An
independent study observed this phenomenon in behaving rats, focusing on CA1 pyramidal cell
burst activity from electrodes that were surgically implanted within the brain [9].
372

Discussion

In this paper, we investigated the role of ionic conductances in causing suprathreshold resonance
for SS and BS in a CA1 pyramidal neuron model. When studying neuronal resonance, it is
important to assess input-output properties and combinations of intrinsic properties [29, 46, 47,
48, 31, 49]. Our focus lay on specific input and output parameters, with emphasis on oscillatory
theta (3-12 Hz) and gamma (30-100 Hz) frequencies for inputs, and SS and BS for outputs. In
particular, we systematically assessed these quantitates separately and identified interleaved SS
and BS resonance that does not match in frequency nor amplitude, but it is modulated by the ionic
currents. To the best of our knowledge, this is the first paper showing the ability of a single
neuronal voltage trace to carry a double-coded spiking resonance. We have also observed that the
burst probability is associated with longer periods of silence and confirmed this observation
through the analysis of in vivo voltage-imaging data from CA1 pyramidal neurons.
385
More specifically, in terms of ionic currents, we found opposite effects of persistent sodium and
delayed rectifier potassium currents. Regarding the ionic conductances, we investigate how G_{NaP}
and G_{KDR} ionic currents affect the emergence of SS to BS. Accordingly, whereas BS is observed
for large G_{NaP} or low G_{KDR}, SS is observed for low G_{NaP} or large G_{KDR}. We found that under
conditions that allow spiking resonance, BS occurs for theta while SS for gamma. Furthermore, at
low input frequencies (theta), the firing rate response is higher when G_{NaP} is low and G_{KDR} is high
due to weak filtering. Conversely, at high input frequencies (gamma), the firing rate response is
lower for the same conditions (low G_{NaP} and high G_{KDR}) due to strong filtering. This different
strength of filtering is a consequence of the modulation of intrinsic conductances (i.e. G_{NaP} and
G_{KDR}). Both conductances cooperate with each other to set the neuronal filtering at different levels
for the respective input frequency (i.e. theta and gamma). One consequence of these results is that
changes in expression levels in these conductances could allow shifting of the strength of the
output for a certain input frequency. As a result, the neuron can facilitate the transfer of low or
high frequency inputs according to the expression of its ionic conductances [50, 51, 52].
400
We found that the interplay between ionic currents and oscillatory input frequencies determines
the output strength of the firing rate. Specifically, we found that decreasing G_{NaP} increases the
firing at theta frequencies and decreases the firing at gamma. Again, G_{KDR} changes had opposite
results. This suggests that CA1 neurons might be able to tune their output preferences by
downregulating or upregulating the expression of their intrinsic conductances [53]. It also suggests

that pharmacological interactions targeting these ionic channels have the potential to affect
oscillations and BS behavior [54, 55].
408
We previously found that BS and theta frequencies are preferentially linked to the animal entering
a place field, while SS and gamma frequencies are linked to the animal leaving a place field [1].
Thus, BS and SS can be considered coding mechanisms for this type of behavior. Our simulations
demonstrate that different oscillatory frequencies can enhance (resonate with) BS and SS
separately in a complex frequency-dependent manner within this double-coded signal. This
indicates a higher-order interaction where the neuron does not need to code either BS or SS
exclusively but can encode both jointly. It would be interesting to find ways in which one
resonance is kept fixed while the other is varied. This would prove whether they are dependent or
independent.
418
An interesting finding of our work was that ionic currents embedded in the neuron can generate
different phase-locking. These results showed that G_{NaP}, G_{KDR}, and G_h modulate the coexistence
of BS and SS phase locked to theta and gamma differently: while low values of G_{NaP} generate BS
at theta, it requires high values of G_{KDR} to generate BS at theta. G_h has little effect on theta
frequencies and affects gamma frequencies instead. I_h current modulation of up/down states is still
under investigation [56], but clearly a target of a diverse range of neuromodulators [57, 58].
425
Unexpectedly, we found that the likelihood of BS remains consistent despite changes in ionic
conductances and primarily relies on extended silent periods [9], meanwhile SS is influenced by
ionic conductances changes and can occur at any intervals. This BS behavior can be explained by
the degeneracy of ionic conductances. In these circumstances, multiple combinations of different
conductance values, even with significant heterogeneity, can produce the same BS behavior in the
neuron model [53, 59]. Another possible interpretation is that the various conductances within the
neuron model (such as transient sodium, muscarinic-sensitive potassium, A-type potassium, high-
threshold calcium, fast calcium-activated potassium, and slow calcium-activated potassium
currents) might respond differently to prolonged hyperpolarization, primarily inducing a state
conducive to bursting rather than single spiking. Conversely, shorter hyperpolarizations could
result in bistability, where both bursting and single spiking scenarios are plausible [60, 61].
437
The probability of spiking is influenced by the states of the conductances (i.e., the levels of
activation/inactivation) before the onset of depolarization [62]. For the conductances examined in
this paper, it is well known that during extended hyperpolarization (such as under the stimulation
of theta frequency oscillations), G_{KDR} and G_{NaP} remain inactive. However, during the initial phase
of depolarization, both G_{KDR} and G_{NaP} are rapidly activated [34], leading to the onset of bursting.
Similarly, it has been observed that the interplay between potassium and sodium conductances
triggers bursting after extended periods of silence [9, 60]. These findings align with previous
research demonstrating that these currents (e.g., G_{KDR} and G_{NaP}) are crucial for generating bursting
[14, 63]. Finally, it has been observed in cortical neurons that bursts can trigger switches from
intermittent bursting spiking to persistent single spiking activity and vice-versa [64]. These
findings bring new insights into the neural coding mechanisms in the CA1 hippocampus.
449

450 # Methods
451

Computational model

We use a Hodgkin-Huxley model of a CA1 pyramidal neuron described by conductance-based ionic currents [14, 1, 65, 66, 67, 68, 69]. The following differential equation describes how the voltage evolves in time:

$$C\frac{dV}{dt} = -g_L(V - E_L) - \sum I_j,$$

with the currents I_j

$$I_j = g_j m^M h^N (V - E_j).$$

The currents included are I_{NaT} the transient sodium, I_{NaP} the persistent sodium ($M = 3, N = 0$), I_{Kdr} the delayed rectifier K^+ ($M = 0, N = 4$), I_{KM} the muscarinic- sensitive potassium ($M = 1, N = 0$), I_{KA} the A-type K^+ ($M = 3, N = 0$), I_{HCa} the high-threshold Ca^{2+} ($M = 3, N = 0$), I_{fKCa} the fast Ca^{2+}-activated potassium current ($M = 1, N = 0$), and I_{sKCa} the slow Ca^{2+}-activated potassium current ($M = 1, N = 0$). The reversal potentials are $E_L = -70$ mV, $E_K = -90$ mV, $E_{Na} = 55$ mV, and $E_{Ca} = 120$ mV. The remaining parameters are $C = 1$ μF/cm^2, $g_L = 0.04$ mS/cm^2, $g_{NaT} = 35$ mS/cm^2, $g_{NaP} = 0.9$ mS/cm^2, $g_{Kdr} = 5.8$ mS/cm^2, $g_{KA} = 2.8$ mS/cm^2, $g_{KM} = 2.4$ mS/cm^2, $g_{HCa} = 0.17$ mS/cm^2, $g_{sCa} = 2.8$ mS/cm^2. Equations for m and n follow

$$\frac{dx}{dt} = \frac{x_\infty(V) - x}{\tau_x(V)},$$

where $x_\infty(V) = \left\{1 + \exp\left[-\frac{(V - \theta_x)}{\sigma_x}\right]\right\}^{-1}$.

The remaining parameters for these equations are $\theta_m = -30$ mV, $\sigma_m = 9.5$ mV, $\theta_h = -45$ mV, $\sigma_h = -7$ mV, $\theta_n = -35$ mV, $\sigma_n = 10$ mV, $\theta_{nt} = -27$ mV, $\sigma_{nt} = -15$ mV, $\theta_a = -50$ mV, $\sigma_a = 20$ mV, $\theta_b = -80$ mV, $\sigma_b = -6$ mV, $\theta_z = -39$ mV, $\sigma_z = 5$ mV, as in [14].

For the compartmental modeling, we add gap junction currents between i and $i+1$ following $I_{gap} = g_{gap}(V_i - V_{i+1})$, with $g_{gap} = 0.2$ mS/cm^2.

For external currents, we use sinusoidal currents with specific frequencies or combinations thereof, as illustrated in the respective figures. Gamma frequencies are defined as ranging from 30 to 100 Hz, and theta frequencies range from 3 to 12 Hz. In our simulations, we use either 40 Hz for gamma or 10 Hz for theta.

Experimental recordings

To compare the model results with experimentally recorded in vivo voltage-imaging using genetically encoded voltage indicator (acemNeon2) in awake moving mice, we used freely available data from [3] in https://tinyurl.com/kannanEtAlScience2022 (last accessed in June 2024). The recordings are 20 seconds long and the data for this paper contain 7 CA1 pyramidal cells from depths ranging from approximately 50 to 100 μm. Image segmentation was performed using custom-written Matlab scripts as used in previous publications [38, 70, 71, 72]. CA1 pyramidal cells were identified by their expression of the negative polarity relationship of

fluorescence emission to the membrane voltage as expected from the voltage indicator (acemNeon2), whereas somatostatin interneurons were identified by the positive polarity that expressed a different voltage indicator (pACe). The voltage indicator was injected in the entorhinal cortex (EC) and expressed retrogradely in CA1, thus, all CA1 pyramidal neurons reported here are projecting to EC. Data acquisition was performed using LED epifluorescence illumination, and the voltage indicator fluorescence was recorded at a rate of 600 Hz with a high-speed sCMOS camera. To correct for photobleaching, we applied a single exponential decay fit and subtracted it from the data. We specifically selected traces from pyramidal cells that exhibited a high signal-to-noise ratio and a high frequency of events, which indicated good recording quality.

Firing rate and coefficient of variation

We compute the firing rate as the sum of spikes divided by the time window considered. Unless stated otherwise, we adopted 4 s in each simulation. The inter-spike intervals (ISI) are taken as the intervals between spikes in the trace. We also divide intervals between single-spikes and bursts accordingly. ISI probabilities are calculated using the same criteria for separating ISIs from single spikes and bursts. The intervals are measured from the last spike to the next single spike or burst spike, regardless of whether the last spike was part of a single spike or a burst.

Burst detection

Following [4] that showed through intracellular studies that individual spikes with complex spikes can have ISIs of more than 10 ms, we adopted a 14ms ISI time window to separate single-spikes from bursts. As for the detected burst, we separate data from the first single spike only to account for the number of bursts and not the intraburst frequency.

Spike phase-locking computation

We quantify the relative phase of a spike to an oscillation using the phase-locking value (PLV):

$$PLV(f) = \left|\frac{1}{N}\right| \sum_N e^{i\phi(f,n)},$$

with f the frequency and N the number of spikes [73]. For theta oscillations we assume 3-12 Hz and for gamma oscillations we assume 30-100Hz. The phase ϕ was obtained with the help of the Hilbert transform. The phase-locking statistics (PLS) accounts for uneven size in the data between bursting and single-spikes [74, 75]:

$$PLS(f) = \frac{1}{N-1}(PLV(f)^2 \times N - 1).$$

Declaration of competing interests

The authors have declared that no competing interests exist.

Data availability

Data will be made available on request.

530 References

[1] E. Lowet, D. J. Sheehan, U. Chialva, R. D. O. Pena, R. A. Mount, S. Xiao, H. G. Rotstein and X. Han, "Theta and gamma rhythmic coding through two spike output modes in the hippocampus during spatial navigation," *Cell Reports,* vol. 42, 2023.

[2] Y. Adam, J. J. Kim, S. Lou, Y. Zhao, M. E. Xie, D. Brinks, ... and A. E. Cohen, "Voltage imaging and optogenetics reveal behaviour-dependent changes in hippocampal dynamics," *Nature,* vol. 569, pp. 413-417, 2019.

[3] M. Kannan, G. Vasan, S. Haziza, C. Huang, R. Chrapkiewicz, J. Luo, J. A. Cardin, M. J. Schnitzer and V. A. Pieribone, "Dual-polarity voltage imaging of the concurrent dynamics of multiple neuron types," *Science,* vol. 378, p. eabm8797, 2022.

[4] J. Epsztein, M. Brecht and A. K. Lee, "Intracellular Determinants of Hippocampal CA1 Place and Silent Cell Activity in a Novel Environment," *Neuron,* vol. 70 , p. 109–120, 2011.

[5] J. D. Cohen, M. Bolstad and A. K. Lee, "Experience-dependent shaping of hippocampal CA1 intracellular activity in novel and familiar environments," *Elife,* vol. 6, p. e23040, 2017.

[6] C. D. Harvey, F. Collman, D. A. Dombeck and D. W. Tank, "Intracellular dynamics of hippocampal place cells during virtual navigation," *Nature,* vol. 461, pp. 941-946, 2009.

[7] K. C. Bittner, A. D. Milstein, C. Grienberger, S. Romani and J. C. Magee, "Behavioral time scale synaptic plasticity underlies CA1 place fields," *Science,* vol. 357, pp. 1033-1036, 2017.

[8] J. Csicsvari, H. Hirase, A. Czurko and G. Buzsáki, "Reliability and state dependence of pyramidal cell–interneuron synapses in the hippocampus: an ensemble approach in the behaving rat," *Neuron,* vol. 21, pp. 179-189, 1998.

[9] K. D. Harris, H. Hirase, X. Leinekugel, D. A. Henze and G. Buzsáki, "Temporal interaction between single spikes and complex spike bursts in hippocampal pyramidal cells," *Neuron, ,* vol. 32, pp. 141-149, 2001.

[10] G. Buzsáki, "Theta oscillations in the hippocampus," *Neuron,* vol. 33, pp. 325-340, 2002.

[11] G. Buzsáki and X. J. Wang, "Mechanisms of gamma oscillations," *Annual review of neuroscience,* vol. 35, pp. 203-225, 2012.

[12] N. Axmacher, M. M. Henseler, O. Jensen, I. Weinreich, C. E. Elger and J. Fell, "Cross-frequency coupling supports multi-item working memory in the human hippocampus," *Proceedings of the National Academy of Sciences,* vol. 107 , pp. 3228-3233, 2010.

[13] J. E. Lisman and O. Jensen, "The theta-gamma neural code," *Neuron,* vol. 77, pp. 1002-1016, 2013.

[14] D. Golomb, C. Yue and Y. Yaari, "Contribution of persistent Na+ current and M-type K+ current to somatic bursting in CA1 pyramidal cells: combined experimental and modeling study," *Journal of Neurophysiology,* vol. 96, pp. 1912-1926, 2006.

[15] J. H. Cocatre-Zilgien and F. Delcomyn, "Identification of bursts in spike trains," *Journal of Neuroscience Methods,* vol. 41, pp. 19-30, 1992.

[16] D. Terman, "The transition from bursting to continuous spiking in excitable membrane models," *Journal of Nonlinear Science,* vol. 2 , pp. 135-182, 1992 .

[17] S. M. Cain and T. P. Snutch, "T-type calcium channels in burst-firing, network synchrony, and epilepsy," *Biochimica et Biophysica Acta (BBA)-Biomembranes,* vol. 1828, pp. 1572-1578, 2013.

[18] D. Jeanmonod, M. Magnin and A. Morel, "Low–threshold calcium spike bursts in the human thalamus: Common physiopathology for sensory, motor and limbic positive symptoms," *Brain,* vol. 119, pp. 363-375, 1996.

[19] M. L. Molineux, J. E. McRory, B. E. McKay, J. Hamid, W. H. Mehaffey, R. Rehak, ... and R. W. Turner, "Specific T-type calcium channel isoforms are associated with distinct burst phenotypes in deep cerebellar nuclear neurons," *Proceedings of the National Academy of Sciences,* vol. 103, pp. 5555-5560, 2006 .

[20] C. A. Del Negro, N. Koshiya, R. J. Butera Jr and J. C. Smith, "Persistent sodium current, membrane properties and bursting behavior of pre-botzinger complex inspiratory neurons in vitro," *Journal of Neurophysiology,* vol. 88, pp. 2242-2250, 2002.

[21] Z. F. Mainen and T. J. Sejnowski, "Influence of dendritic structure on firing pattern in model neocortical neurons," *Nature,* vol. 382, pp. 363-366, 1996.

[22] H. Su, G. Alroy, E. D. Kirson and Y. Yaari, "Extracellular calcium modulates persistent sodium current-dependent burst-firing in hippocampal pyramidal neurons," *Journal of Neuroscience,* vol. 21, pp. 4173-4182, 2001.

[23] E. Stark, R. Eichler, L. Roux, S. Fujisawa, H. G. Rotstein and G. Buzsáki, "Inhibition-induced theta resonance in cortical circuits," *Neuron,* vol. 80, pp. 1263-1276, 2013 .

[24] M. H. Higgs and W. J. Spain, "Conditional bursting enhances resonant firing in neocortical layer 2–3 pyramidal neurons," *Journal of Neuroscience,* vol. 29, pp. 1285-1299, 2009.

[25] J. A. Beatty, S. C. Song and C. J. Wilson, "Cell-type-specific resonances shape the responses of striatal neurons to synaptic input," *Journal of Neurophysiology,* vol. 113, pp. 688-700, 2015.

[26] S. Ostojic, G. Szapiro, E. Schwartz, B. Barbour, N. Brunel and V. Hakim, "Neuronal morphology generates high-frequency firing resonance," *Journal of Neuroscience,* vol. 35, pp. 7056-7068, 2015.

[27] T. Deemyad, J. Kroeger and M. J. Chacron, "Sub-and suprathreshold adaptation currents have opposite effects on frequency tuning," *The Journal of Physiology,* vol. 590, pp. 4839-4858, 2012.

[28] T. Broicher, P. Malerba, A. D. Dorval, A. Borisyuk, F. R. Fernandez and J. A. White, "Spike phase locking in CA1 pyramidal neurons depends on background conductance and firing rate," *Journal of Neuroscience,* vol. 32, pp. 14374-14388, 2012.

[29] M. J. Richardson, N. Brunel and V. Hakim, "From subthreshold to firing-rate resonance," *Journal of Neurophysiology,* vol. 89, pp. 2538-2554, 2003.

[30] E. Stark, A. Levi and H. G. Rotstein, "Network resonance can be generated independently at distinct levels of neuronal organization," *PLoS computational biology,* vol. 18, p. e1010364, 2022 .

[31] B. Hutcheon and Y. Yarom, "Resonance, oscillation and the intrinsic frequency preferences of neurons," *Trends in Neurosciences,* vol. 23, pp. 216-222, 2000 .

[32] R. F. Pena and H. G. Rotstein, "The voltage and spiking responses of subthreshold resonant neurons to structured and fluctuating inputs: persistence and loss of resonance and variability," *Biological Cybernetics,* vol. 116, pp. 163-190, 2022.

[33] R. F. Pena, C. C. Ceballos, V. Lima and A. C. Roque, "Interplay of activation kinetics and the derivative conductance determines resonance properties of neurons," *Physical Review E,* vol. 97, p. 042408, 2018.

[34] C. C. Ceballos, A. C. Roque and R. M. Leao, "A negative slope conductance of the persistent sodium current prolongs subthreshold depolarizations," *Biophysical Journal,* vol. 113, pp. 2207-2217, 2017.

[35] C. C. Ceballos, A. C. Roque and R. M. Leão, "The role of negative conductances in neuronal subthreshold properties and synaptic integration," *Biophysical Reviews,* vol. 9, pp. 827-834, 2017.

[36] A. Kepecs and J. Lisman, "Information encoding and computation with spikes and bursts," *Network: Computation in Neural Systems,* vol. 14, pp. 103-118, 2003.

[37] J. E. Lisman, "Bursts as a unit of neural information: making unreliable synapses reliable," *Trends in Neurosciences,* vol. 20, pp. 38-43, 1997.

[38] K. D. Piatkevich, S. Bensussen, H. A. Tseng, S. N. Shroff, V. G. Lopez-Huerta, D. Park, ... and X. Han, "Population imaging of neural activity in awake behaving mice," *Nature,* vol. 574, pp. 413-417, 2019.

[39] J. I. Kass and I. M. Mintz, "Silent plateau potentials, rhythmic bursts, and pacemaker firing: three patterns of activity that coexist in quadristable subthalamic neurons," *Proceedings of the National Academy of Sciences,* vol. 103, pp. 183-188, 2006.

[40] J. C. Magee, "Dendritic hyperpolarization-activated currents modify the integrative properties of hippocampal CA1 pyramidal neurons," *Journal of Neuroscience,* vol. 18, pp. 7613-7624, 1998.

[41] A. Lörincz, T. Notomi, G. Tamás, R. Shigemoto and Z. Nusser, "Polarized and compartment-dependent distribution of HCN1 in pyramidal cell dendrites," *Nature Neuroscience,* vol. 5, p. 1185–1193, 2002.

[42] E. Campanac, G. Daoudal, N. Ankri and D. Debanne, "Downregulation of dendritic Ih in CA1 pyramidal neurons after LTP," *Journal of Neuroscience ,* vol. 28, pp. 8635-8643, 2008.

[43] C. C. Ceballos, R. F. Pena and A. C. Roque, "Impact of the activation rate of the hyperpolarization-activated current Ih on the neuronal membrane time constant and synaptic potential duration," *The European Physical Journal Special Topics,* vol. 230, pp. 2951-2961, 2021.

[44] G. A. Ascoli, S. Gasparini, V. Medinilla and M. Migliore, " Local control of postinhibitory rebound spiking in CA1 pyramidal neuron dendrites," *Journal of Neuroscience,* vol. 30, pp. 6434-6442, 2010.

[45] C. C. Ceballos, S. Li, A. C. Roque, T. Tzounopoulos and R. M. Leão, "Ih equalizes membrane input resistance in a heterogeneous population of fusiform neurons in the dorsal cochlear nucleus," *Frontiers in Cellular Neuroscience,* vol. 10, p. 249, 2016.

[46] G. Hahn, A. F. Bujan, Y. Frégnac, A. Aertsen and A. Kumar, "Communication through resonance in spiking neuronal networks," *PLoS Computational Biology,* vol. 10, p. e1003811, 2014.

[47] J. S. Sherfey, S. Ardid, J. Hass, M. E. Hasselmo and N. J. Kopell, "Flexible resonance in prefrontal networks with strong feedback inhibition," *PLoS Computational Biology,* vol. 14, p. e1006357, 2018.

[48] J. Sherfey, S. Ardid, E. K. Miller, M. E. Hasselmo and N. J. Kopell, "Prefrontal oscillations modulate the propagation of neuronal activity required for working memory," *Neurobiology of Learning and Memory,* vol. 173, p. 107228, 2020.

[49] R. F. Pena, V. Lima, R. O. Shimoura, C. C. Ceballos, H. G. Rotstein and A. C. Roque, "Asymmetrical voltage response in resonant neurons shaped by nonlinearities," *Chaos: An Interdisciplinary Journal of Nonlinear Science,* vol. 29, p. 103135, 2019.

[50] G. S. Newkirk, D. Guan, N. Dembrow, W. E. Armstrong, R. C. Foehring and W. J. Spain, "Kv2. 1 potassium channels regulate repetitive burst firing in extratelencephalic neocortical pyramidal neurons," *Cerebral Cortex,* vol. 32, pp. 1055-1076, 2022.

[51] S. Chen, H. Su, C. Yue, S. Remy, M. Royeck, D. Sochivko, ... and Y. Yaari, "An increase in persistent sodium current contributes to intrinsic neuronal bursting after status epilepticus," *Journal of Neurophysiology,* vol. 105, pp. 117-129, 2011 .

[52] C. L. Hsu, X. Zhao, A. D. Milstein and N. Spruston, "Persistent sodium current mediates the steep voltage dependence of spatial coding in hippocampal pyramidal neurons," *Neuron,* vol. 99, pp. 147-162, 2018.

[53] R. Roy and R. Narayanan, "Ion-channel degeneracy and heterogeneities in the emergence of complex spike bursts in CA3 pyramidal neurons," *The Journal of Physiology,* vol. 601, pp. 3297-3328, 2023.

[54] K. A. Alkadhi and L. M. Tian, "Veratridine-enhanced persistent sodium current induces bursting in CA1 pyramidal neurons," *Neuroscience,* vol. 71 , pp. 625-632, 1996.

[55] P. Müller, A. Draguhn and A. V. Egorov, "Persistent sodium current modulates axonal excitability in CA 1 pyramidal neurons," *Journal of Neurochemistry,* vol. 146, pp. 446-458, 2018.

[56] L. Dalla Porta, A. Barbero-Castillo, J. M. Sanchez-Sanchez, N. Cancino and M. V. Sanchez-Vives, "H-current modulation of cortical Up and Down states," *bioRxiv,* pp. 2024-04, 2024.

[57] H. C. Pape, "Queer current and pacemaker: the hyperpolarization-activated cation current in neurons," *Annual Review of Physiology,* vol. 58, pp. 299-327, 1996.

[58] C. He, F. Chen, B. Li and Z. Hu, "Neurophysiology of HCN channels: from cellular functions to multiple regulations," *Progress in Neurobiology,* vol. 112, pp. 1-23, 2014.

[59] G. Drion, T. O'Leary and E. Marder, "Ion channel degeneracy enables robust and tunable neuronal firing rates," *Proceedings of the National Academy of Sciences,* vol. 112, pp. E5361-E5370, 2015.

[60] F. S. Borges, P. R. Protachevicz, D. L. Souza, C. F. Bittencourt, E. C. Gabrick, L. E. Bentivoglio, J. D. Szezech, A. Batista, I. Caldas, S. Dura-Bernal and R. F. Pena, "The Roles of Potassium and Calcium Currents in the Bistable Firing Transition," *Brain Sciences,* vol. 13, p. 1347, 2023.

[61] P. R. Protachevicz, F. S. Borges, E. L. Lameu, P. Ji, K. C. Iarosz, A. L. Kihara, I. L. Caldas, J. D. J. Szezech, M. S. Baptista, E. E. N. Macau, C. G. Antonopoulos, A. M. Batista and J. Kurths, "Bistable firing patterns in a neural network model," *Frontiers in Computational Neuroscience,* vol. 13, p. 19, 2019.

[62] J. Yamada-Hanff and B. P. Bean, "Activation of Ih and TTX-sensitive sodium current at subthreshold voltages during CA1 pyramidal neuron firing," *Journal of Neurophysiology,* vol. 114, pp. 2376-2389, 2015.

[63] A. Compte, M. V. Sanchez-Vives, D. A. McCormick and X. J. Wang, "Cellular and network mechanisms of slow oscillatory activity (< 1 Hz) and wave propagations in a cortical network model," *Journal of Neurophysiology,* vol. 89, pp. 2707-2725, 2003.

[64] C. Li, Y. T., M. M. Poo and Y. Dan, "Burst spiking of a single cortical neuron modifies global brain state," *Science,* vol. 324, pp. 643-646, 2009 .

[65] R. Azouz, M. S. Jensen and Y. Yaari, "Ionic basis of spike after-depolarization and burst generation in adult rat hippocampal CA1 pyramidal cells," *The Journal of Physiology,* vol. 492, pp. 211-223, 1996.

[66] M. Bazhenov, I. Timofeev, M. Steriade and T. J. Sejnowski, "Potassium model for slow (2-3 Hz) in vivo neocortical paroxysmal oscillations," *Journal of Neurophysiology,* vol. 92, pp. 1116-1132, 2004.

[67] R. Bertram, M. J. Butte, T. Kiemel and A. Sherman, "Topological and phenomenological classification of bursting oscillations," *Bulletin of Mathematical Biology,* vol. 57, pp. 413-439, 1995.

[68] I. A. Fleidervish, A. Friedman and M. J. Gutnick, "Slow inactivation of Na+ current and slow cumulative spike adaptation in mouse and guinea-pig neocortical neurones in slices," *The Journal of Physiology,* vol. 493, pp. 83-97, 1996.

[69] A. Kepecs and X. J. Wang, "Analysis of complex bursting in cortical pyramidal neuron models," *Neurocomputing,* vol. 32, pp. 181-187, 2000.

[70] S. Xiao, E. Lowet, H. J. Gritton, P. Fabris, Y. Wang, J. Sherman, ... and X. Han, "Large-scale voltage imaging in behaving mice using targeted illumination," *Iscience,* vol. 24, 2021.

[71] E. Lowet, K. Kondabolu, S. Zhou, R. A. Mount, Y. Wang, C. R. Ravasio and X. Han, "Deep brain stimulation creates informational lesion through membrane depolarization in mouse hippocampus," *Nature Communications,* vol. 13, p. 7709, 2022.

[72] S. Xiao, W. J. Cunningham, K. Kondabolu, E. Lowet, M. V. Moya, R. A. Mount, ... and J. Mertz, "Large-scale deep tissue voltage imaging with targeted-illumination confocal microscopy," *Nature Methods ,* pp. 1-9, 2024.

[73] J. P. Lachaux, A. Lutz, D. Rudrauf, M. Cosmelli, J. Le Van Quen, F. Martinerie and F. Varela, "Estimating the time-course of coherence between single-trial brain signals: an introduction to wavelet coherence," *Clinical Neurophysiology,* vol. 32, p. 157–174, 2002.

[74] M. Vinck, M. van Wingerden, T. Womelsdorf, P. Fries and C. M. A. Pennartz, "The pairwise phase consistency: A bias-free measure of rhythmic neuronal synchronization," *Neuroimage,* vol. 51, p. 112–122 , 2010.

[75] S. Aydore, D. Pantazis and R. M. Leahy, "A note on the phase locking value and its properties," *Neuroimage,* vol. 74, p. 231–244, 2013.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

Only gamma input

B

Only theta input

Figure 6

Figure 7