Interleaved single and bursting spiking resonance in neurons

Cesar C. Ceballos[1], Nourdin Chadly[3,4], Eric Lowet[3], Rodrigo F. O. Pena[1,2, *]

[1] Department of Biological Sciences, Florida Atlantic University, Jupiter, FL 33458, USA
[2] Stiles-Nicholson Brain Institute, Florida Atlantic University, Jupiter, FL 33458, USA
[3] Department of Neuroscience, Erasmus Medical Center, Rotterdam, Netherlands
[4] Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, Netherlands

* Corresponding author
E- mail: penar@fau.edu

Abstract

Under *in vivo* conditions, CA1 pyramidal cells from the hippocampus display transitions from single spikes to bursts. It is believed that subthreshold hyperpolarization and depolarization, also known as down and up-states, play a pivotal role in these transitions. Nevertheless, a central impediment to correlating suprathreshold (spiking) and subthreshold activity has been the technical difficulties of this type of recordings, even with widely used calcium imaging or multielectrode recordings. Recent work using voltage imaging with genetically encoded voltage indicators has been able to correlate spiking patterns with subthreshold activity in a variety of CA1 neurons, and recent computational models have been able to capture these transitions. In this work, we used a computational model of a CA1 pyramidal cell to investigate the role of intrinsic conductances and oscillatory patterns in generating down and up-states and their modulation in the transition from single spiking to bursting. Specifically, the emergence of distinct spiking resonances between these two spiking modes that share the same voltage traces in the presence of theta or gamma oscillatory inputs, a phenomenon we call interleaved single and bursting spiking resonance. We noticed that these resonances do not necessarily overlap in frequency or amplitude, underscoring their relevance for providing flexibility to neural processing. We studied the conductance values of three current types that are thought to be critical for the bursting behavior: persistent sodium current (I_{NaP}) and its conductance G_{NaP}, delayed rectifier potassium (I_{KDR}) and its conductance G_{KDR}, and hyperpolarization-activated current (I_h) and its conductance G_h. We conclude that the intricate interplay of ionic currents significantly influences the neuronal firing patterns, transitioning from single to burst firing during sustained depolarization. Specifically, the intermediate levels of G_{NaP} and G_{KDR} facilitate spiking resonance at gamma frequency inputs. The resonance characteristics vary between single and burst firing modes, each displaying distinct amplitudes and resonant frequencies. Furthermore, low G_{NaP} and high G_{KDR} values lock bursting to theta frequencies, while high G_{NaP} and low G_{KDR} values lock single spiking to gamma frequencies. Lastly, the duration of quiet intervals plays a crucial role in determining the likelihood of transitioning to either bursting or single spiking modes. We confirmed that the same features were present in previously recorded in vivo voltage-imaging data. Understanding these dynamics provides valuable insights into the fundamental mechanisms underlying neuronal excitability under *in vivo* conditions.

Author summary

Since discovering that neurons in the hippocampus can encode spatial position through phase precession, many experiments have explored how specific theta and gamma oscillations influence location specificity in the brain. However, the individual neuronal properties and dynamics behind these behaviors are still being uncovered. Previously, we found that stereotypical bursting and single-spike firing in pyramidal neurons are linked to these oscillations and further associated with an animal entering or leaving a place field. Advances in voltage-imaging techniques have enabled us to assess these properties more precisely. Our study shows that different frequencies can independently trigger these stereotypical spikes, demonstrating a complex pattern where the same cell can be double-coded: a phenomenon we called interleaved resonance. Additionally, we found that this coding can be modulated by persistent sodium and delayed-rectifier potassium currents. Moreover, these neurons are more likely to burst following long periods of silence. These findings provide new insights into the mechanisms underlying neural coding in the hippocampus and how it relates to behavior.

Introduction

Neurons from the CA1 hippocampus display a diversity of spiking dynamics. In the hippocampus, from the point of view of individual cells, *in vivo* voltage imaging [1, 2, 3], patch clamp recordings [4, 5, 6, 7], and multielectrode recordings [8, 9] reveals that CA1 pyramidal cells exhibit stereotypical single and burst spiking. While investigating their emergence and transitions it was noted that these spike modes appear with different probabilities of occurrence where bursts are more likely to occur after long periods of activity silence [9]. From the point of view of populations, gamma (30-100 Hz) and theta (3-12 Hz) oscillations coexist and have been well characterized in CA1 region of rodent hippocampus [10, 11]. Gamma is modulated by theta during working memory retention, and it is important to notice the cross-frequency coupling of these oscillations for spatial navigation and memory performance [12, 13]. Recent studies have tried to connect those transitions using *in vivo* data, as shown by shifts from single to bursting spikes in the same neurons that are preferentially observed during theta and gamma oscillations, respectively [1]. Here we use the term burst to define cluster of spikes with intervals lower than 14 ms, according to studies where this variation is found ~6-14 ms [9, 4, 14]. Behaviorally, these spiking modes have been proposed to encode an animal's position, where burst signals when animals enter a place. In contrast, single spikes signal when the animal leaves a place [1].

Admittedly, understanding spiking patterns transitions is synonymous to understanding ionic current's dynamics. For fixed synaptic input, the composition of ionic currents can produce distinguishable spiking outputs that vary from single to burst of spikes [15, 16, 17]. Combinations of ionic currents can determine these spike modes as in the case where expression of calcium is linked to emergence of burst [18, 19, 20], or in combination with the spatial location on the neuron as in the case of persistent sodium current (I_{NaP}) located at or near the soma [15, 21], or only the geometry [22], or combinations of both [23]. Moreover, potassium currents such as I_h have been investigated in terms of their effects on subthreshold oscillations and resonance [24, 25, 26, 27], although their influence on bursting has still not been clarified (but see [28] for modeling efforts on its influence on this direction).

Oscillations can cause transitions from low to high firing rate on a neuron. Cells may have
amplified response for certain oscillatory frequencies compared to other frequencies. This is
known as suprathreshold (spiking) resonance, and it has been observed widely across the brain
[29, 30, 31, 32, 33], including CA1 pyramidal cells [34] and the mechanisms underlying this
resonance have been intensively investigated using computational models [35, 36]. Similarly,
subthreshold resonance can be seen as a transition voltage change of a neuron upon a particular
(non-zero) input frequency [37, 38]. Several ionic currents contribute to subthreshold resonance
and can be discussed in terms of their chord and derivative conductances [39]. For example,
persistent sodium current (I_NaP) is an amplifying ionic current associated to bursting in CA1 [40]
and delayed rectifier potassium (I_KDR) is a resonant current [37, 41, 42]. Less is known about the
role of these ionic currents on suprathreshold (spiking) resonance.
99
It is only natural to assume that CA1 pyramidal neurons embedded in a network can take advantage
of the spiking resonance to preferentially transmit information encoded in the spiking mode (single
or burst spiking) when a particular oscillatory synaptic input (e.g. theta or gamma oscillations) is
received [34, 29, 43, 44]. In addition to that, it is important to understand how transitions at these
different scales connect and modulate them. In this paper, we investigate suprathreshold resonance
of single and burst spiking in a CA1 pyramidal neuron model. We found that spiking resonance
can easily emerge when neurons are stimulated with theta and gamma synaptic inputs and ionic
currents are tuned. Variability of sodium and potassium currents differentially promoted the
presence of resonance according to the firing pattern (single vs. burst), a phenomenon we call
interleaved single and bursting spiking resonance. Our results provide insights that might help to
understand the mechanisms underlying single and burst spiking computation and its role in
connection to *in vivo* processing that is relevant for spike coding in CA1 pyramidal cells for spatial
navigation and memory.
113
Our paper is organized as follows: in the following section, we present studies of how ionic
currents promote transitions from single to burst firing under constant sustained input and later
under oscillatory inputs. For the oscillatory inputs, we show how resonance emerges and depends
on ionic currents. Then we move to a more systematic study where we characterize interleaved
single and bursting resonance in the presence of dendritic I_h current. For the same model, we
characterize phase-locking to gamma and theta and how these main ionic currents shape this
relationship. We conclude by systematically characterizing the probability of bursting spikes in
relation to the periods of silence in the voltage traces. Methods are included in the last section of
the paper.
123

Results

Ionic currents promote a transition from single to burst firing during sustained depolarization

CA1 pyramidal cells are known to exhibit transitions from single (SS) to bursting spikes (BS) *in*
vivo [1, 2, 3, 45]. This transition occurs in the presence of subthreshold oscillations and is
modulated by oscillatory inputs, mostly theta and gamma oscillations. Subthreshold activity is
characterized by transitions from hyperpolarized states to depolarized states (also known as down
and up-states, respectively). Firing mostly occurs on top of the depolarization phases. Input

patterns play a relevant role in the switching, but it is unclear how they can be promoted by the
neuron's intrinsic properties, e.g. ionic currents. We first decided to investigate the firing patterns
occurring during depolarization states and how SS and BS emerge under different ionic
conductances.
137
We first investigated the role of conductances that are mostly expressed in the soma, such as I_{NaP}
and I_{KDR}. In this section, we used a single compartment computational model of CA1 pyramidal
cell (see Methods) and we varied G_{NaP} (persistent sodium conductance) and G_{KDR} (delayed rectifier
potassium conductance) for the model receiving a step function of current that maintains the
voltage in a constant depolarized state (Fig 1A). We observed that increasing G_{NaP} makes the firing
transits from SS to BS and finally to plateau potential, also known as depolarization block [46]
(see voltage traces Fig 1B1). The phase plots (dV/dt) in the same figure reveals how the voltage
series transits between the different modes (Fig 1B2). In contrast, the effect of increasing G_{KDR} is
a transition from BS to SS (Fig 1C1) and this can be seen in the phase plots (Fig 1C2). Single
spiking can be explained as a result of having long pauses between spikes due to long
afterhyperpolarization, which is facilitated by strong action potential repolarization due to large
G_{KDR} or by preventing fast depolarization due to small G_{NaP}.
150

151
Fig 1. Ionic currents promote transition from SS to BS during sustained depolarization.
Default values for fixed conductance while varying the other are $G_{NaP} = 0.3$ and $G_{KDR} = 6$. **A** Sketch
of CA1 model receiving a step function of current. **B1** Example voltage traces and **B2** phase plots
(dV/dt) when varying G_{NaP}. **C1** Example voltage traces and **C2** phase plots (dV/dt) when varying
G_{KDR}. **D1** Firing rate and **D2** coefficient of variation (CV), a measure the activity of the neuron
and the presence of bursts, displayed as heatmaps spanned over G_{NaP}. **E1** Firing rate and **E2**
coefficient of variation (CV) displayed as heatmaps spanned over G_{KDR} vs. Input current. For phase
plots we remove the first 100 ms, but for all other panels we show the traces from the beginning
with stimulation application starting at 0 ms.
161

The firing rate increases with the increase of stimulation amplitude, increase of G_{NaP} and decrease
of G_{KDR} (Fig 1D1 and 1E1, respectively). As a measure of the burstiness we adopted the coefficient
of variation (CV), as described in the methods. The CV above 2 (reddish to yellowish) determines
the area in which the combination of conductance promotes BS behavior (Fig 1D2 and 1E2). As
it is shown in the heatmaps, BS is independent of the input amplitude and mostly occurs in areas
around $G_{NaP} = 0.35$ or $G_{KDR} = 6$ (Fig 1D2 and 1E2, respectively). These values correspond to the
edges where the transition from spiking to depolarization block occurs (see dark regions in
heatmaps of firing rate). Interestingly, BS occurs in a very narrow range of conductances,
meanwhile single spiking is likely to occur in a wide range of values. This set of observations
shows the opposite effects of the G_{NaP} and G_{KDR} ionic currents in modulating a switch between SS
to BS. We have also investigated the effect of the additional ionic currents I_{KA} the A-type K^+, I_{KM}
the muscarinic- sensitive potassium, I_{sKCa} the slow Ca^{2+}-activated potassium current, and I_{fKCa} the
fast Ca^{2+} -activated potassium current (Fig. S1). Only G_{KM} caused a significant effect on the
modulation of bursting behavior.
176

Intermediate G_{NaP} and G_{KDR} generate spiking resonance
preferentially at gamma frequency input
We showed previously that G_{NaP} and G_{KDR} have opposite effects on the generated spiking patterns
during sustained depolarization. Nevertheless, CA1 pyramidal neurons receive oscillatory inputs
in the range from 3 to 100 Hz, where oscillatory gamma is between 30-100 Hz and theta 3-12 Hz
frequencies. This induces transitions from hyperpolarized to depolarized states where shorter
intervals occur at high input frequencies such as gamma frequencies and longer intervals at theta
frequencies. Under this condition, firing during depolarization is not sustained, but rather is
alternated by periods of silence. In our paper, we investigate the role of the silent states generated
by the interplay between input patterns and ionic currents to create a suprathreshold resonance and
its relation to SS and BS.
188

For stimulation of the cell in this section, we choose varying oscillatory stimulation frequencies
covering the range from theta to gamma frequencies (Fig 2A). In Figs 2B1, B2, D1, and D2 we
present voltage-traces for variations of G_{NaP} and G_{KDR} under theta or gamma (see captions). We
investigate the firing rate responses and their preference for a certain input frequency (i.e.
suprathreshold resonance). We observe low-pass filtering of the firing rate for low G_{NaP} or high
G_{KDR}, suprathreshold resonance for intermediate G_{NaP} or G_{KDR}, and a nearly absent frequency
preference for high values of G_{NaP} or low values of G_{KDR} (Fig 2C, E). Interestingly, regardless of
the opposite effects of G_{NaP} and G_{KDR}, we observed that theta input has a stronger effect on the
firing rate response than gamma when the neuron acts as a low-pass filter. In contrast, gamma is
stronger than theta for resonance (Fig 2C, E). These results unveiled an interaction between the
input frequency and the filtering properties that determine the firing output.
200

201
Fig 2. Intermediate G_{NaP} and G_{KDR} generates firing rate resonance. Default values for fixed
conductance while varying the other are $G_{NaP} = 0.3$ and $G_{KDR} = 6$. **A** Sketch of simplified CA1
model receiving oscillatory current stimulation. **B1** Examples of voltage traces for varying values
of G_{NaP} under gamma stimulation. **B2** Examples of voltage traces for varying values of G_{NaP} under
theta stimulation. **C** Firing rate while varying stimulation frequency of an oscillatory input. Shaded
red area: theta frequencies. Shaded blue area: gamma frequencies. **D1** Examples of voltage traces
for varying values of G_{KDR} under gamma stimulation. **D2** Examples of voltage traces for varying
values of G_{KDR} under theta stimulation. **E** Firing rate while varying stimulation frequency of an
oscillatory input. **F1** Firing rate and **F2** CV heatmap spanned over G_{NaP} and oscillatory input
frequency. **G1** Firing rate and **G2** CV heatmap spanned over G_{KDR} and oscillatory input frequency.
In all panels, stimulation starts at 0 ms. To allow slow but non-zero frequencies to be studied we
cut-off the simulations at 2 Hz.

214
The CVs above 2 (reddish to yellowish) obtained while changing the oscillatory input frequency
and the conductance indicate the area where BS occurs (Fig 2F2 and 2G2). The heatmaps show
similar patterns for firing frequency and CV for both G_{NaP} and G_{KDR}, i.e. we observe high firing
rates simultaneously with high values of CV (Fig 2F1, 2F2, 2G1 and 2G2). The high firing rates
obtained can be justified by the BS and low firing rates are by the SS (Fig 2F1, 2G1). Note that
for these panels we used a cut-off of 2 Hz to allow slow frequencies to be tested but different from
zero. Also note that certain conductance value choices may lead the neuron into non-physiological
states, which are most likely to occur at low G_{KDR} values and high G_{NaP} values which is acceptable
for exploratory purposes.

224
We observed BS at gamma frequencies when G_{NaP} levels were intermediate or high (see the
yellowish diagonal in Fig. 2F2) and when G_{KDR} was at intermediate levels (Fig. 2G2).
Additionally, BS was noted at theta frequencies with low G_{NaP} or high G_{KDR}. Similarly, we
observed SS at low G_{KDR} and at gamma frequencies with low or intermediate G_{NaP} or high G_{KDR}.
Interestingly, BS in gamma frequencies occurs simultaneously with resonance at intermediate
values of G_{NaP} or G_{KDR}. These results in Figs 2F1 and 2F2 indicate that when suprathreshold
resonance is present, a linear increase in input frequency and G_{NaP} shifts the BS presence from
theta to gamma frequencies. Similarly, a decrease in G_{KDR}, along with increased input frequencies,
also shifts the BS presence from theta to gamma frequencies (as shown by the yellowish regions
in the heatmaps, Fig 2G1 and 2G2). A similar pattern is observed for SS, with the dark regions in
the G_{NaP} heatmap and the dark middle-right region in the G_{KDR} heatmap illustrating this shift (Fig
2F2 and 2G2, respectively). We have also investigated the effect of the additional ionic currents
I_{KA} the A-type K^+, I_{KM} the muscarinic- sensitive potassium, I_{sKCa} the slow Ca^{2+}-activated
potassium current, and I_{fKCa} the fast Ca^{2+}-activated potassium current (Fig. S2). Only G_{KM} caused
a significant effect on the modulation of bursting behavior. However, the effect of G_{KM} was
restricted to gamma band, meanwhile for G_{NaP} and G_{KDR} we observed effects in both gamma and
theta. Combined with the results in Fig. S1, we therefore decided to focus on the effect of G_{NaP}
and G_{KDR} for the remainder of the paper.
243
244 In summary, for low G_{NaP} and high G_{KDR}, we found high firing rate response to low input
245 frequencies (i.e. theta) due to weak filtering, and low firing rate response to high input frequencies
246 (i.e. gamma) due to strong filtering. This suggests that the tuning of ionic currents in interaction
247 with input frequencies are critical for determining the occurrence of resonance.
248

Single and burst firing display suprathreshold resonance with different amplitudes and different resonant frequencies

It is well known that CA1 pyramidal cells have a high expression of the hyperpolarization activated
I_h current [47]. This current is mostly expressed in the dendrites and plays a relevant role in
modulating synaptic inputs [47, 48, 49, 50] and resonance [39]. It prevents excessive
hyperpolarization and causes rebound spiking after hyperpolarizations [51, 52]. Thus, in this
section, we incorporated a dendritic compartment containing I_h current (Fig 3A). Following
previous work [1], I_h is also necessary for the transition single-spike/bursting behavior and
therefore it is added in the next sections.
258

259
Figure 3. Single and burst firing display suprathreshold resonance with different amplitudes and resonant frequencies. Default values for fixed conductance while varying the other are G_{NaP} = 0.93, G_{KDR} = 5.86, and G_h = 8.46. **A** Sketch of CA1 model with additional dendritic compartment receiving oscillatory current stimulation. **B** Exemplary voltage trace of the simulation with SS and BS indicated. **C1-2** Firing rate vs oscillatory input frequency for SS **C1** and BS **C2** when G_{NaP} values are 0, 0.5 and 1.5. **D1-2** Firing rate vs oscillatory input frequency for SS **D1** and BS **D2** when G_{KDR} values are 0, 5 and 10. **E1-2** Firing rate vs oscillatory input frequency for SS **E1** and BS **E2** when G_h values are 0, 2 and 5. **F** Representative sketch demonstrating how Δf and ΔA are calculated for the interleaved SS and BS resonances. **G1-2** Δf and ΔA for varying values of G_{NaP}. **H1-2** Δf and ΔA for varying values of G_{KDR}. **I1-2** Δf and ΔA for varying values of G_h. Colors mark the magnitude of ionic currents (blue = low, orange = medium, green = high).

We stimulated the cell model with varying oscillatory stimulation frequencies covering the range from theta to gamma frequencies (Fig 3A). Under this condition, the neuron is capable of switching its firing from SS to BS (Fig 3B). We separated our quantitative measures into SS and BS (see

275 Methods). Overall, we observed resonance for all the explored conductances and for both spiking
276 types (SS and BS), except for BS when G_{NaP} is large or when G_{KDR} is low (Fig 3 C2, D2). Finally,
277 SS and BS have resonance for all G_h values (Fig 3 E1, E2).
278
We further compared the resonance frequencies and their respective firing rate amplitudes between
SS and BS (Fig 3F). For low G_{NaP}, SS resonance frequency is larger than BS resonance frequency,
but it is the opposite when G_{NaP} is large (Fig 3G1). For low G_{KDR}, SS resonance frequency is
smaller than BS resonance frequency, but it is the opposite when G_{KDR} is large (Fig 3H1).
Moreover, SS resonance frequency is larger than BS resonance frequency for all G_h values (Fig
3I1). Finally, as expected, firing frequency at the resonant peak was always higher for BS rather
than SS (Fig 3G2, H2, I2).
286
287 The set of results in this section demonstrates an interesting coding behind SS and BS in our CA1
288 model. First, it shows that neither the resonance frequencies nor resonance amplitudes between SS
289 and BS necessarily overlap. More interestingly, this phenomenon happens at the same voltage
290 traces (interleaved) and can be modulated by ionic currents.
291
To gain more insight about the mechanism underlying the generation of interleaved single spikes
and burst, in particular for currents that were not studied in depth in our analysis, we compared
conditions where the I_{KM} is present and when it is zeroed, the latter a simulation of the experimental
effect of channel blocking. Voltage traces show that decreasing I_{KM} affect mostly single spiking,
specifically converting them into bursts (Fig S2E; notice first, third, and seventh cycle where single
spike become burst). Furthermore, we characterized the time evolution of activation variables of
I_{KDR}, I_{KM} and I_{NaP} (Fig S3). We observed different time scales, with I_{NaP} being the fastest current,
I_{KDR} being slower and I_{KM} the slowest current. This was expected from the way their kinetics was
set up (see Methods).
301
We also investigated the effect of the background noise on the interleaved resonance from single
spiking to burst. We found the interleaved resonance is sensitive to the noise level (Fig S4).
Altogether, these results suggest that the interleaving from single spiking to burst is a consequence
of the interplay of three ionic currents (I_{KDR}, I_{KM} and I_{NaP}) working at different timescales and
modulated by the presence of noise. In the next section, we show how these observations are
connected to oscillations.
308

G_{NaP}, G_{KDR} and G_h determine phase locking of SS and BS to theta and gamma input respectively

It has been demonstrated that this neuron model can reproduce coexisting SS and BS phase-locked
to gamma and theta oscillations, respectively, and that these frequency preference (resonance) of
SS and BS is interleaved. The model is derived from [1, 15] and has its conductance fitted by
obtaining values that generate voltages that match experimental voltage imaging. In this paper, we
modified the model for a similar fitting (see parameters in Methods). Based on the observations
above for changing ionic conductances and the theta and gamma relationship to BS and SS, we
question the ability of the ionic channels to modulate the phase-locking of the neuron model to
these frequencies.
319

Since most of CA1 neuron inputs come with a mixture of theta and gamma frequencies, it is only
natural to study the case where both are present. Thus, we injected both theta and gamma
simultaneously into the neuron model (Fig 4A). As observed, the ionic currents embedded in the
neuron can generate different phase-locking (Fig 4B). Low values of G_{NaP} are more prone to BS
at theta frequencies (Fig 4C1, Cohen's d = 1.66; Fig 4C2, Cohen's d = 2.77). Increasing the values
of G_{NaP} removes this phase locking with the generation of more SS (Fig 4C3). In contrast, G_{KDR}
had the opposite behavior. We see only SS for low values of G_{KDR} (Fig 4D1), BS at gamma for
intermediate values of G_{KDR} and finally BS at theta is generated for high G_{KDR} (Fig 4D2, Cohen's
d =0.54; Fig 4D3, Cohen's d = 2.41). Interestingly, only low and high values of G_h promoted a
difference at the gamma frequencies (Fig 4E1, Cohen's d = 0.68; Fig 4E3, Cohen's d = -0.26).
330

331
Fig. 4. G_{NaP}, G_{KDR} and G_h determine phase locking statistics (PLS) of SS and BS to theta and
gamma input. Default values for fixed conductance while varying the other are $G_{NaP} = 0.93$, G_{KDR}

= 5.86, and G_h = 8.46. **A** Sketch of CA1 model with somatic and dendritic compartments receiving
theta and gamma oscillatory current stimulation. **B** Examples of voltage traces with SS behavior
transitioning to a BS behavior (compare top to bottom). **C1** Phase locking statistic (PLS) to theta
and gamma input separated into SS and BS with varying G_{NaP} = 0 (asterisk has Cohen's d 1.66),
C2 when G_{NaP} = 0.5 (asterisk has Cohen's d 2.77) and **C3** when G_{NaP} = 1.5. **D1** Phase locking
statistic (PLS) with varying G_{KDR} = 0, **D2** when G_{KDR} = 5 (asterisk has Cohen's d 0.54) and **D3**
when G_{KDR} = 10 (asterisk has Cohen's d 2.41). **E1** Phase locking statistic (PLS) with varying G_h
= 0 (asterisk has Cohen's d 0.68), **E2** when G_h = 2 and **E3** G_h = 5 (asterisk has Cohen's d -0.26).
Each bar is the result of 10 different simulations. t-test with 0.05 level of significance (asterisk).
343
344 Taken together, these results show that G_{NaP}, G_{KDR}, and G_h modulate the coexistence of BS and
345 SS phase locked to theta and gamma differently: low values of G_{NaP} (Fig 4C1) and high values of
346 G_{KDR} (Fig 4D3) lock BS to theta frequencies. G_h has little effect on theta frequencies and affects
347 gamma frequencies instead. Furthermore, low values of G_{KDR} lock SS to gamma frequencies (Fig
348 4D1).
349

350
351 **Fig 5. G_{KA}, G_{KDR}, G_{KM}, and G_{NaP} determine firing rate of SS and BS.** Heatmaps of firing rate
352 for all the conductances for different input frequencies for each pattern (SS and BS). **A1** BS for
353 G_{KA} **A2** SS for G_{KA}. **B1** BS for G_{fKCa} **B2** SS for G_{fKCa}. **C1** BS for G_{KDR} **C2** SS for G_{KDR}. **D1** BS
354 for G_{KM} **D2** SS for G_{KM}. **E1** BS for G_{NaP}. **E2** SS for G_{NaP}. **F1** BS for G_{sKCa}. **F2** SS for G_{sKCa}.
355
As previously observed for the remaining ionic currents, we also investigated the separate role of
each ionic current in the model and how it would shape the emergence of SS or BS. This would
mean either performing a sensitivity analysis or fixing all other parameters and further examining
the contribution of all conductances (G_{KA}, G_{fKCa}, G_{KDR}, G_{KM}, G_{NaP}, and G_{sKCa}) to firing frequency
and its dependence on input frequency for each pattern (SS and BS). Those results are presented
in Fig 5 where heatmaps of firing rates (SS and BS) are displayed as conductance values vary. Our
results indicate that calcium-dependent potassium currents (G_{fKCa} and G_{sKCa}) do not influence the
firing properties of SS and BS. In contrast, G_{KA}, G_{KDR}, G_{KM}, and G_{NaP} exhibit significant effects.
Notably, all potassium conductances follow a consistent pattern: increasing their values reduces

the input frequency at which SS and BS reach maximum firing. Conversely, G_{NaP} shows the
opposite effect.
367
Quiet intervals determine the probability of bursting or single spiking
369 The *in vivo* activity of CA1 neurons is characterized by subthreshold oscillations that follows the
370 theta and gamma synaptic inputs. During these oscillations, spiking activity is observed at the
371 depolarization phase and between them it is possible to observe silence intervals at hyperpolarized
372 phase. During the depolarization, neurons can fire SS or BS. In this section, we investigate whether
373 the silence intervals modulate the probability of having SS or BS.
374
According to our observations in Fig 6, high probability of BS occurs for long intervals of silence
(above 100 ms) in spite of variation in the values of the conductances G_{NaP}, G_{KDR}, and G_h (Fig
6D1, 6E1 and 6F1). Interestingly, we observed that interevent intervals (ISI) are fractured and
distributed in more than one position in the diagram, and that different peaks show up mostly for
large G_{NaP} or low G_{KDR} (Fig 6A3 and 6B1). We also observed that high probability of SS occurs
for short intervals of silence (below 60 ms) for low G_{NaP}, high G_{KDR} and regardless of the values
of G_h (Fig 6D2, 6E2 and 6F2). Moreover, we observed high probability of SS occurring for long
intervals (above 100 ms) only when G_{NaP} is high or when G_{KDR} is low (Fig 6D2 and 6E2,
respectively). In summary, we found that the probability of BS is robust to variation of the ionic
conductances and mostly depends on long silent periods above 100 ms (i.e. when theta input is
present). In contrast, SS is influenced by ionic conductances and can occur at short or long
intervals.
387

388
Figure 6. Quiet intervals determine burst and single spikes. Default values for fixed
conductance while varying the other are $G_{NaP} = 0.93$, $G_{KDR} = 5.86$, and $G_h = 8.46$. **A1-A3**
Histogram of interevent intervals (ISI) with varying G_{NaP}. **B1-B3** Histogram of interevent intervals

(ISI) with varying G_{KDR}. **C1-C3** Histogram of interevent intervals (ISI) with varying G_h. **D1, E1**
and F1 BS and **D2, E2 and F2** SS probabilities for varying G_{NaP}, G_{KDR}, and G_h, respectively.
Colors mark the magnitude of ionic currents (green = low, red = medium, blue = high).
395
To determine whether the spiking probability is the sum of independent effects of theta and gamma
stimulation, or the result of the interaction between both inputs, we calculated the spiking
probability when only theta or gamma inputs were used (Fig 7). As expected, when only gamma
input is present, we observed that the probability of BS or SS is always larger at short intervals of
silence (< 50 ms) (Fig 7A1-7A6). In contrast, when only theta input is present, we observed that
the probability of BS is larger for long intervals of silence (> 100 ms) (Fig 7B1-7B3). For SS
probability when only theta input is present, we observed the same tendency found above, where
high probability of SS occurs for short intervals of silence below 60 ms for low G_{NaP}, high G_{KDR}
and regardless the values of G_h, and high probability of SS occurs for long intervals above 100 ms
only when G_{NaP} is high or when G_{KDR} is low (Fig 7B4-7B6).
406

407
408 **Figure 7. Burst probability follows the longest quiet intervals according to the slowest**
409 **oscillatory input**. Default values for fixed conductance while varying the other are $G_{NaP} = 0.93$,
410 $G_{KDR} = 5.86$, and $G_h = 8.46$. **A1-A3** BS and **A4-A6** SS probabilities with only gamma input. **B1-**
411 **B3** BS and **B4-B6** SS probabilities with only theta input. Colors mark the magnitude of ionic
412 currents (green = low, red = medium, blue = high).
413
In summary, we observed that when the input is only at gamma frequencies, the spiking probability
is largest for short silent intervals for both SS and BS. Opposite to that, when the input is only
oscillating at theta, BS is more likely to occur at long intervals and SS at short intervals.
Interestingly, when both theta and gamma are present, the probabilities behave as if only theta was
present, i.e., the gamma component is mostly disregarded even though the stimulation contains
both theta and gamma (Fig 7). These results suggest that BS probability follows the input
frequency with the longest intervals of silence, i.e. it follows theta even in the presence of gamma,

but it follows gamma in the absence of any other slowest input. Meanwhile, SS would
preferentially occur at short intervals.
423

Voltage imaging burst probability follows the longest quiet intervals

Despite the difficulties in reconciling our computational results with voltage-imaging recordings,
where frequency-dependent stimulation is typically absent, it is possible to compare the bursting
probability following the longest quiet intervals with spontaneous recordings. Thus, we analyzed
voltage imaging traces of 7 neurons (20 s long) obtained from a public repository [3]. We used a
difference source than [1] who were using a general synapsin promoter, hence that was not specific
for pyramidal neurons. For our purposes, we chose the dataset available at [3] because the neurons
are identified as pyramidal neurons using the CamKII promoter. Fluorescence images were
collected from CA1 pyramidal cells and somatostatin expressing interneurons located in the dorsal
CA1 area at depths ~50 to 100 μm (Fig 8A). LED epifluorescence illumination allowed the
collection of the voltage indicator fluorescence by a high-speed sCMOS camera at a rate
acquisition of 600 Hz. We observed waveform distortions in the voltage imaging traces due to low
pass filtering from the slower voltage indicator kinetics relative to the action potential. While this
limits the technique, it does not affect our results, as our focus is on peak detection rather than
spike kinetics. Photobleaching was subtracted by fitting a single exponential decay. We selected
the traces from pyramidal cells based on their high signal-to-noise ratio and high frequency of
events suggesting good quality of recordings (Fig 8B). These criteria allowed us to properly detect
and classify events as SS and BS as done in the computational simulations (Fig 8C). We calculated
the probability of finding SS or BS for a certain ISI (Fig 8D). We performed a BS and SS
comparison statistical comparison between the set of ISIs falling in the 3 areas marked for low ISI
(100 to 150 ms), intermediate ISI (150 to 250 ms), and long ISI (400 to 500 ms).
445

Figure 8. Voltage imaging burst probability follows the longest quiet intervals. Data from [3].
A Sketch of the voltage-imaging experiment. **B** Sample traces that were used for this analysis
(right, n = 7). **C** Sample trace showing spike detection algorithm prior to division into BS and SS.
D Interevent interval (ISI) probability density function of BS and SS (n = 7), and 3 areas marked
for low ISI (100 to 150 ms), intermediate ISI (150 to 250 ms), and long ISI (400 to 500 ms). **E**
From the marked areas, bar plots with averaged values in each ISI trace (n = 7). t-test perfomed
with the data in the particular area that delivers 0.05 level of significance (asterisk). Cohen's d = -
1.02 and p=0.03 for Area 1, Cohen's d = -0.24 and p=0.68 for Area 2, and Cohen's d = 1.54 and
p=0.02 for Area 3.
456
The results showed that SS is more likely at low ISIs, while BS is more likely at long ISIs.
Observations of these probabilities across all traces (Fig. 8D) and statistical significance from the
traces and areas (Fig. 8E) confirm this. The voltage-imaging recordings agree with our simulation
observations, where burst probability follows the longest quiet intervals (Figs 6 and 7). An
independent study observed this phenomenon in behaving rats, focusing on CA1 pyramidal cell
burst activity from electrodes that were surgically implanted within the brain [9].
463

Discussion

In this paper, we investigated the role of ionic conductances in causing suprathreshold resonance
for SS and BS in a CA1 pyramidal neuron model. When studying neuronal resonance, it is
important to assess input-output properties and combinations of intrinsic properties [35, 53, 54,
55, 37, 56]. Our focus lay on specific input and output parameters, with emphasis on oscillatory
theta (3-12 Hz) and gamma (30-100 Hz) frequencies for inputs, and SS and BS for outputs. In
particular, we systematically assessed these quantitates separately and identified interleaved SS
and BS resonance that does not match in frequency nor amplitude, but it is modulated by the ionic
currents. To the best of our knowledge, this is the first paper showing the ability of a single
neuronal voltage trace to carry a double-coded spiking resonance, where single spikes and burst
were separated and shown to have different frequency preferences: interleaved resonance. We have
also observed that the burst probability is associated with longer periods of silence and confirmed
this observation through the analysis of in vivo voltage-imaging data from CA1 pyramidal
neurons.
478
More specifically, in terms of ionic currents, we found opposite effects of persistent sodium and
delayed rectifier potassium currents. Regarding the ionic conductances, we investigate how G_{NaP}
and G_{KDR} ionic currents affect the emergence of SS to BS. Accordingly, whereas BS is observed
for large G_{NaP} or low G_{KDR}, SS is observed for low G_{NaP} or large G_{KDR}. We found that under
conditions that allow spiking resonance, BS occurs for theta while SS for gamma. Furthermore, at
low input frequencies (theta), the firing rate response is higher when G_{NaP} is low and G_{KDR} is high
due to weak low-pass filtering. Conversely, at high input frequencies (gamma), the firing rate
response is lower for the same conditions (low G_{NaP} and high G_{KDR}) due to strong filtering. This
different strength of filtering is a consequence of the modulation of intrinsic conductances (i.e.
G_{NaP} and G_{KDR}). Both conductances cooperate with each other to set the neuronal filtering at
different levels for the respective input frequency (i.e. theta and gamma). One consequence of
these results is that changes in expression levels in these conductances could allow shifting of the
strength of the output for a certain input frequency. As a result, the neuron can facilitate the transfer
of low or high frequency inputs according to the expression of its ionic conductances [57, 58, 59].

Future studies should investigate this phenomenon from the point of view of other ionic currents,
in particular G_{KM} which seems to promote modulation of the bursting behavior (Fig S1) but have
not been studied in this work. Other currents such as the calcium-dependent potassium seem to
have a minor effect. In addition, we acknowledge that the amplitude current values used in this
work are larger than what one would expect experimentally, which is an artifact of the fact that the
potassium reversal potential was established at -90 mV. Therefore, values can go as low as -90
mV and around it. However, we have tested smaller amplitude currents (reduced by 2-fold), which
cause peak-to-peak membrane potential oscillations of less than 20 mV. This produced similar
qualitative results with different quantitative values, such as shifting the resonance frequency (see
Fig. S6).
503
We found that the interplay between ionic currents and oscillatory input frequencies determines
the output strength of the firing rate. Specifically, we found that decreasing G_{NaP} increases the
firing at theta frequencies and decreases the firing at gamma. Again, G_{KDR} changes had opposite
results. This suggests that CA1 neurons might be able to tune their output preferences by
downregulating or upregulating the expression of their intrinsic conductances [60]. It also suggests
that pharmacological interactions targeting these ionic channels have the potential to affect
oscillations and BS behavior [61, 62].
511
We previously found that BS and theta frequencies are preferentially linked to the animal entering
a place field, while SS and gamma frequencies are linked to the animal leaving a place field [1].
Thus, BS and SS can be considered coding mechanisms for this type of behavior. Our simulations
demonstrate that different oscillatory frequencies can enhance (resonate with) BS and SS
separately in a complex frequency-dependent manner within this double-coded signal. This
indicates a higher-order interaction where the neuron does not need to code either BS or SS
exclusively but can encode both jointly. It would be interesting to find ways in which one
resonance is kept fixed while the other is varied. This would prove whether they are dependent or
independent.
521
An interesting finding of our work was that ionic currents embedded in the neuron can generate
different phase-locking. These results showed that G_{NaP}, G_{KDR}, and G_h modulate the coexistence
of BS and SS phase locked to theta and gamma differently: while low values of G_{NaP} generate BS
at theta, it requires high values of G_{KDR} to generate BS at theta. G_h has little effect on theta
frequencies and affects gamma frequencies instead (Fig 4 E1-E3). I_h current modulation of
up/down states is still under investigation [63], but clearly a target of a diverse range of
neuromodulators [64, 65].
529
Unexpectedly, we found that the likelihood of BS remains consistent despite changes in ionic
conductances and primarily relies on extended silent periods [9], meanwhile SS is influenced by
ionic conductances changes and can occur at any intervals. This BS behavior can be explained by
the degeneracy of ionic conductances. In these circumstances, multiple combinations of different
conductance values, even with significant heterogeneity, can produce the same BS behavior in the
neuron model [60, 66]. Another possible interpretation is that the various conductances within the
neuron model (such as transient sodium, muscarinic-sensitive potassium, A-type potassium, high-
threshold calcium, fast calcium-activated potassium, and slow calcium-activated potassium
currents) might respond differently to prolonged hyperpolarization, primarily inducing a state

conducive to bursting rather than single spiking. Conversely, shorter hyperpolarizations could
result in bistability, where both bursting and single spiking scenarios are plausible [67, 68].
541
The probability of spiking is influenced by the states of the conductances (i.e., the levels of
activation/inactivation) before the onset of depolarization [69]. For the conductances examined in
this paper, it is well known that during extended hyperpolarization (such as under the stimulation
of theta frequency oscillations), G_{KDR} and G_{NaP} remain inactive. However, during the initial phase
of depolarization, both G_{KDR} and G_{NaP} are rapidly activated [41], leading to the onset of bursting.
Similarly, it has been observed that the interplay between potassium and sodium conductances
triggers bursting after extended periods of silence [9, 67]. These findings align with previous
research demonstrating that these currents (e.g., G_{KDR} and G_{NaP}) are crucial for generating bursting
[15, 70]. Finally, it has been observed in cortical neurons that bursts can trigger switches from
intermittent bursting spiking to persistent single spiking activity and vice-versa [71]. These
findings bring new insights into the neural coding mechanisms in the CA1 hippocampus.
553

Methods

Computational model

557 We use a Hodgkin-Huxley model of a CA1 pyramidal neuron described by conductance-based
558 ionic currents [15, 1, 72, 73, 74, 75, 76]. The following differential equation describes how the
559 voltage evolves in time:

560
$$C\frac{dV}{dt} = -g_L(V - E_L) - \Sigma I_j + I_{app},$$

561 with the currents I_j
562
563
$$I_j = g_j m^M h^N (V - E_j).$$

The currents included are I_{NaT} the transient sodium (M = 3, N = 1), I_{NaP} the persistent sodium (M
= 1, N = 0), I_{Kdr} the delayed rectifier K^+ (M = 0, N = 4), I_{KM} the muscarinic- sensitive potassium
(M = 1, N = 0), I_{KA} the A-type K^+ (M = 3, N = 0), I_{HCa} the high-threshold Ca^{2+} (M = 2, N = 0),
I_{fKCa} the fast Ca^{2+} -activated potassium current (M = 1, N = 0), and I_{sKCa} the slow Ca^{2+}-activated
potassium current (M = 1, N = 0). In addition, when the dendrite is present it contains an I_h current
(M=1, N=0). The reversal potentials are $E_L = -70$ mV, $E_K = -90$ mV, $E_{Na} = 55$ mV, and $E_{Ca} = 120$
mV. The remaining parameters are $C = 1$ μF/cm^2, $g_L = 0.04$ mS/cm^2, $g_{NaT} = 35$ mS/cm^2, $g_{NaP} = 0.9$
mS/cm^2, $g_{Kdr} = 5.8$ mS/cm^2, $g_{KA} = 2.8$ mS/cm^2, $g_h = 8.46$ mS/cm^2, $g_{KM} = 2.4$ mS/cm^2, $g_{HCa} = 0.17$
mS/cm^2, $g_{sKCa} = 2.8$ mS/cm^2, $g_{fKCa} = 9.05$ mS/cm^2. Throughout the paper, all conductance values
will be displayed in mS/cm^2, but we will omit the units unless necessary for clarity. All currents
in the model have units μA/cm^2, but we will also omit units. Equations for m and n follow
576
577
$$\frac{dx}{dt} = \frac{x_\infty(V) - x}{\tau_x(V)},$$

578 where $x_\infty(V) = \left\{1 + \exp\left[-\frac{(V-\theta_x)}{\sigma_x}\right]\right\}^{-1}$. This is the same for all ionic currents except to I_{fKCa}

579 which has d variable governed by calcium following $d_\infty([Ca^{2+}]_i) = \{1 + a_c/[Ca^{2+}]_i\}^{-1}$ and

580 I_{sKCa} which has a q variable governed by calcium following $q_\infty([Ca^{2+}]_i) = \{1 + a_q^4/[Ca^{2+}]_i^4\}^{-1}$.

582 For the dynamics of the calcium concentration inside the cell $[Ca^{2+}]_i$ we assume

584
$$\frac{d[Ca^{2+}]_i}{dt} = -\nu I_{HCa} - \frac{[Ca^{2+}]_i}{\tau_{HCa}},$$

where $\nu = 0.13$ cm²/(ms× μA), and $\tau_{HCa} = 13$ ms.

The remaining parameters for these equations are $\theta_m = -30$ mV, $\sigma_m = 9.5$ mV, $\theta_h = -45$ mV,

$\sigma_h = -7$ mV, $\theta_p = -35$ mV, $\sigma_p = 3$ mV, $\theta_n = -35$ mV, $\sigma_n = 10$ mV, $\theta_{nt} = -27$ mV, $\sigma_{nt} =$

-15 mV, $\theta_a = -50$ mV, $\sigma_a = 20$ mV, $\theta_b = -80$ mV, $\sigma_b = -6$ mV, $\theta_z = -39$ mV, $\sigma_z = 5$ mV,

$\theta_r = -20$ mV, $\sigma_r = 19$ mV, $\theta_c = 30$ mV, $\sigma_c = -7$ mV as in [15].

For the compartmental modeling, we add axial conductance between i and $i+1$ following the ohmic

current $I_{ac} = g_{ac}(V_i - V_{i+1})$, with $g_{ac} = 0.2$ mS/cm²

For external currents, we use sinusoidal currents with specific frequencies or combinations thereof,

as illustrated in the respective figures. Gamma frequencies are defined as ranging from 30 to 100

Hz, and theta frequencies range from 3 to 12 Hz. In our simulations, we use either 40 Hz for gamma

or 10 Hz for theta. We chose sinusoidal current injections over conductance-based synapses

because our previous work [38] showed that cells exhibit impedance peaks (Z-resonance)

and voltage envelope resonance in response to sinusoidal inputs but not to square-wave or

synaptic-like inputs. This applies to both conductance- and current-based inputs. Thus, we

used sinusoidal stimulation, consistent with prior experimental approaches [1].

A complete description of the model can be found in the table below.

Ionic currents	Time constant, ms	Parameters
$I_{NaT}(V,h) = g_{NaT}m_\infty^3(V)h(V - V_{Na})$	$\tau_h = 0.1 + 0.75 \times \left\{1 + \exp\left[-\frac{V - \theta_{ht}}{\sigma_{ht}}\right]\right\}^{-1}$, $m = m_\infty$	$\theta_m = -30$ mV, $\sigma_m = 9.5$ mV, $\theta_h = -45$ mV, $\sigma_h = -7$ mV
$I_{NaP}(V) = g_{NaP}p_\infty(V)(V - V_{Na})$	$p = p_\infty$	$\theta_p = -35$ mV, $\sigma_p = 3$ mV
$I_{Kdr}(V,n) = g_{Kdr}n^4(V - V_K)$	$\tau_n = 0.1 + 0.5 \times \left\{1 + \exp\left[-\frac{V - \theta_{nt}}{\sigma_{nt}}\right]\right\}^{-1}$	$\theta_n = -35$ mV, $\sigma_n = 10$ mV
$I_{KM}(V,z) = g_{KM}z(V - V_K)$	$\tau_z = 75$	$\theta_z = -39$ mV, $\sigma_z = 5$ mV
$I_{KA}(V,b) = g_{KA}a^3(V)b(V - V_K)$	$a = a_\infty$ $\tau_b = 15$	$\theta_a = -50$ mV, $\sigma_a = 20$ mV, $\theta_b = -80$ mV, $\sigma_h = 15$ mV
$I_h(V,) = g_h t(V)(V - V_K)$	$\tau_t = \{\exp[-17.9 + 0.116V] + \exp[-1.84 + 0.09V]\}^{-1} + 0.1$	$\theta_a = -84$ mV, $\sigma_a = 10.2$ mV
$I_{HCa}(V,r) = g_{HCa}r^2(V - V_{Ca})$	$\tau_r = 1$	$\theta_r = -20$ mV, $\sigma_r = 19$ mV
$I_{fKCa}(V,c) = g_{fKCa}d_\infty([Ca^{2+}]_i)c(V - V_K)$	$\tau_c = 2$	$\theta_c = 30$ mV, $\sigma_c = -7$ mV, $a_c = 6$ mV

$I_{sKCa}(V, q) = g_{sKCa}q(V - V_K)$	$\tau_q = 450$	$a_q = 2$ mV
Current between compartments		
$I_{ac} = g_{ac}(V_i - V_{i+1})$		$g_{ac} = 0.2$
External current		
$I_{sin} = I_\theta + I_\gamma$	I_θ is a sinusoidal 10Hz current and I_γ is 40Hz, with amplitudes 1 μA/cm^2 and 2 μA/cm^2, respectively.	
I_{app}	Amplitude 1 μA/cm^2	
External Noise		
$I_\eta = \sqrt{2D}\xi(t)$	The function $\xi(t)$ is a zero-mean white Gaussian noise with $\langle\xi(t)\xi(t')\rangle = \delta(t - t')$.	$D = 9.6 \times 10^{-6}$

606   Table 1. Summary of parameters and variables used to build the computational model.
607
608   The code for this model is available at https://github.com/rodrigo-pena-lab/interleaved_resonance.
609

Experimental recordings

To compare the model results with experimentally recorded in vivo voltage-imaging using
genetically encoded voltage indicator (acemNeon2) in awake moving mice, we used freely
available data from [3] in https://tinyurl.com/kannanEtAlScience2022 (last accessed in June
2024). The recordings are 20 seconds long and the data for this paper contain 7 CA1 pyramidal
cells from depths ranging from approximately 50 to 100 μm. Image segmentation was performed
using custom-written Matlab scripts as used in previous publications [45, 77, 78, 79]. CA1
pyramidal cells were identified by their expression of the negative polarity relationship of
fluorescence emission to the membrane voltage as expected from the voltage indicator
(acemNeon2), whereas somatostatin interneurons were identified by the positive polarity that
expressed a different voltage indicator (pACe). The voltage indicator was injected in the entorhinal
cortex (EC) and expressed retrogradely in CA1, thus, all CA1 pyramidal neurons reported here are
projecting to EC. Data acquisition was performed using LED epifluorescence illumination, and
the voltage indicator fluorescence was recorded at a rate of 600 Hz with a high-speed sCMOS
camera. To correct for photobleaching, we applied a single exponential decay fit and subtracted it
from the data. We specifically selected traces from pyramidal cells that exhibited a high signal-to-
noise ratio and a high frequency of events, which indicated good recording quality.
627

Firing rate and coefficient of variation

629   We compute the firing rate as the sum of spikes divided by the time window considered. Unless
630   stated otherwise, we adopted 4 s in each simulation. The inter-spike intervals (ISI) are taken as
631   the intervals between spikes in the trace. From the ISI we define the coefficient of variation (CV)
632   which points to burstiness in the traces for CV>2, and is defined as the standard deviation over the
633   mean of the ISI:
634

$$CV = std(ISI)/mean(ISI).$$

636
637   We also divide intervals between single-spikes and bursts accordingly. ISI probabilities are
638   calculated using the same criteria for separating ISIs from single spikes and bursts. The intervals
639   are measured from the last spike to the next single spike or burst spike, regardless of whether the
640   last spike was part of a single spike or a burst.

Burst detection

Following [4] that showed through intracellular studies that individual spikes with complex spikes can have ISIs of more than 10 ms, we adopted a 14ms ISI time window to separate single-spikes from bursts. At least two spikes are necessary for this analysis, otherwise the spike-train is disregarded. As for the detected burst, we separate data from the first single spike only to account for the number of bursts and not the intraburst frequency.

Spike phase-locking computation

We quantify the relative phase of a spike to an oscillation using the phase-locking value (PLV):

$$PLV(f) = \left| \frac{1}{N} \right| \sum_N e^{i\phi(f,n)},$$

with f the frequency and N the number of spikes [80]. For theta oscillations we assume 3-12 Hz and for gamma oscillations we assume 30-100Hz. The phase ϕ was obtained with the help of the Hilbert transform. The phase-locking statistics (PLS) accounts for uneven size in the data between bursting and single-spikes [81, 82]:

$$PLS(f) = \frac{1}{N-1}(PLV(f)^2 \times N - 1).$$

Resonance frequencies and amplitudes

We compare resonance frequencies and their amplitudes using the following equations: $\Delta f = f_{SS} - f_{BS}$ and $\Delta A = A_{SS} - A_{BS}$, where f_{SS} is the resonance frequency of single spikes and f_{BS} is the resonance frequency of bursts (in Hz). Also, A_{SS} is the firing frequency at the resonance peak (also known as firing rate amplitude) of single spikes and A_{BS} is the firing frequency at the resonance peak of bursts (in Hz).

Declaration of competing interests

The authors have declared that no competing interests exist.

Data availability

Data will be made available on request.

References

[1] E. Lowet, D. J. Sheehan, U. Chialva, R. D. O. Pena, R. A. Mount, S. Xiao, S. L. Zhou, H. Tseng, H. Gritton, S. Shroff, K. Kondabolu, C. Cheung, Y. Wang, K. D. Piatkevich, E. D. Boyden, J. Mertz, M. E. Hasselmo, H. G. Rotstein and X. Han, "Theta and gamma rhythmic coding through two spike output modes in the hippocampus during spatial navigation," *Cell Reports,* vol. 42, 2023.

[2] Y. Adam, J. J. Kim, S. Lou, Y. Zhao, M. E. Xie, D. Brinks, H. Wu, M. Mostajo-Radji, S. Kheifets, V. Parot, S. Chettih, K. J. Williams, B. Gmeiner, S. L. Farhi, L. Madisen, E. K. Buchanan, I. Kinsella, D. Zhou, L. Paninski, C. D. Harvey, . and A. E. Cohen, "Voltage imaging and optogenetics reveal behaviour-dependent changes in hippocampal dynamics," *Nature,* vol. 569, pp. 413-417, 2019.

[3] M. Kannan, G. Vasan, S. Haziza, C. Huang, R. Chrapkiewicz, J. Luo, J. A. Cardin, M. J. Schnitzer and V. A. Pieribone, "Dual-polarity voltage imaging of the concurrent dynamics of multiple neuron types," *Science,* vol. 378, p. eabm8797, 2022.

[4] J. Epsztein, M. Brecht and A. K. Lee, "Intracellular Determinants of Hippocampal CA1 Place and Silent Cell Activity in a Novel Environment," *Neuron ,* vol. 70 , p. 109–120, 2011.

[5] J. D. Cohen, M. Bolstad and A. K. Lee, "Experience-dependent shaping of hippocampal CA1 intracellular activity in novel and familiar environments," *Elife,* vol. 6, p. e23040, 2017.

[6] . D. Harvey, . Collman, . A. Dombeck and D. W. Tank, "Intracellular dynamics of hippocampal place cells during virtual navigation," *Nature,* vol. 461, pp. 941-946, 2009.

[7] . C. Bittner, . D. Milstein, . Grienberger, . Romani and J. C. Magee, "Behavioral time scale synaptic plasticity underlies CA1 place fields," *Science,* vol. 357, pp. 1033-1036, 2017.

[8] J. Csicsvari, H. Hirase, A. Czurko and G. Buzsáki, "Reliability and state dependence of pyramidal cell–interneuron synapses in the hippocampus: an ensemble approach in the behaving rat," *Neuron,* vol. 21, pp. 179-189, 1998.

[9] K. D. Harris, H. Hirase, X. Leinekugel, D. A. Henze and G. Buzsáki, "Temporal interaction between single spikes and complex spike bursts in hippocampal pyramidal cells.," *Neuron, ,* vol. 32, pp. 141-149, 2001.

[10] G. Buzsáki, "Theta oscillations in the hippocampus," *Neuron,* vol. 33, pp. 325-340, 2002.

[11] G. Buzsáki and X. J. Wang, "Mechanisms of gamma oscillations.," *Annual review of neuroscience,* vol. 35, pp. 203-225, 2012.

[12] N. Axmacher, M. M. Henseler, O. Jensen, I. Weinreich, C. E. Elger and J. Fell, "Cross-frequency coupling supports multi-item working memory in the human hippocampus," *Proceedings of the National Academy of Sciences,* vol. 107 , pp. 3228-3233, 2010.

[13] J. E. Lisman and O. Jensen, "The theta-gamma neural code," *Neuron,* vol. 77, pp. 1002-1016, 2013.

[14] K. Mizuseki, S. Royer, K. Diba and G. Buzsáki, "Activity Dynamics and Behavioral Correlates of CA3 and CA1 Hippocampal Pyramidal Neurons," *Hippocampus ,* vol. 22, p. 1659–1680, 2012.

[15] D. Golomb, C. Yue and Y. Yaari, "Contribution of persistent Na+ current and M-type K+ current to somatic bursting in CA1 pyramidal cells: combined experimental and modeling study," *Journal of Neurophysiology,* vol. 96, pp. 1912-1926, 2006.

[16] J. H. Cocatre-Zilgien and F. Delcomyn, "Identification of bursts in spike trains," *Journal of Neuroscience Methods,* vol. 41, pp. 19-30, 1992.

[17] D. Terman, "The transition from bursting to continuous spiking in excitable membrane models," *Journal of Nonlinear Science,* vol. 2 , pp. 135-182, 1992 .

[18] S. M. Cain and T. P. Snutch, "T-type calcium channels in burst-firing, network synchrony, and epilepsy," *Biochimica et Biophysica Acta (BBA)-Biomembranes,* vol. 1828, pp. 1572-1578, 2013.

[19] D. Jeanmonod, M. Magnin and A. Morel, "Low–threshold calcium spike bursts in the human thalamus: Common physiopathology for sensory, motor and limbic positive symptoms," *Brain,* vol. 119, pp. 363-375, 1996.

[20] M. L. Molineux, J. E. McRory, B. E. McKay, J. Hamid, W. H. Mehaffey, R. Rehak, ... and R. W. Turner, "Specific T-type calcium channel isoforms are associated with distinct burst phenotypes in deep cerebellar nuclear neurons," *Proceedings of the National Academy of Sciences,* vol. 103, pp. 5555-5560, 2006 .

[21] C. A. Del Negro, N. Koshiya, R. J. Butera Jr and J. C. Smith, "Persistent sodium current, membrane properties and bursting behavior of pre-botzinger complex inspiratory neurons in vitro," *Journal of Neurophysiology,* vol. 88, pp. 2242-2250, 2002.

[22] Z. F. Mainen and T. J. Sejnowski, "Influence of dendritic structure on firing pattern in model neocortical neurons," *Nature,* vol. 382, pp. 363-366, 1996.

[23] H. Su, G. Alroy, E. D. Kirson and Y. Yaari, "Extracellular calcium modulates persistent sodium current-dependent burst-firing in hippocampal pyramidal neurons," *Journal of Neuroscience,* vol. 21, pp. 4173-4182, 2001.

[24] R. Klink and A. Alonso, "Ionic mechanisms for the subthreshold oscillations and differential electroresponsiveness of medial entorhinal cortex layer II neurons," *Journal of Neurophysiology,* vol. 70, pp. 144-157, 1993.

[25] C. T. Dickson, J. Magistretti, M. H. Shalinsky, E. Fransén, M. E. Hasselmo and A. Alonso, "Properties and role of Ih in the pacing of subthreshold oscillations in entorhinal cortex layer II neurons," *Journal of Meurophysiology ,* vol. 83, pp. 2562-2579, 2000.

[26] E. Fransén, A. A. Alonso, C. T. Dickson, J. Magistretti and M. E. Hasselmo, "Ionic mechanisms in the generation of subthreshold oscillations and action potential clustering in entorhinal layer II stellate neurons," *Hippocampus,* vol. 14, pp. 368-384, 2004.

[27] R. Narayanan and D. Johnston, "The h channel mediates location dependence and plasticity of intrinsic phase response in rat hippocampal neurons," *Journal of Neuroscience,* vol. 28, pp. 5846-5860, 2008.

[28] S. A. Neymotin, M. M. Hilscher, T. C. Moulin, Y. Skolnick, M. T. Lazarewicz and W. W. Lytton, "Ih tunes theta/gamma oscillations and cross-frequency coupling in an in silico CA3 model," *PLoS One,* vol. 8, p. e76285, 2013.

[29] E. Stark, R. Eichler, L. Roux, S. Fujisawa, H. G. Rotstein and G. Buzsáki, "Inhibition-induced theta resonance in cortical circuits," *Neuron,* vol. 80, pp. 1263-1276, 2013 .

[30] M. H. Higgs and W. J. Spain, "Conditional bursting enhances resonant firing in neocortical layer 2–3 pyramidal neurons," *Journal of Neuroscience,* vol. 29, pp. 1285-1299, 2009.

[31] J. A. Beatty, S. C. Song and C. J. Wilson, "Cell-type-specific resonances shape the responses of striatal neurons to synaptic input," *Journal of Neurophysiology,* vol. 113, pp. 688-700, 2015.

[32] S. Ostojic, G. Szapiro, E. Schwartz, B. Barbour, N. Brunel and V. Hakim, "Neuronal morphology generates high-frequency firing resonance," *Journal of Neuroscience,* vol. 35, pp. 7056-7068, 2015.

[33] T. Deemyad, J. Kroeger and M. J. Chacron, "Sub-and suprathreshold adaptation currents have opposite effects on frequency tuning," *The Journal of Physiology,* vol. 590, pp. 4839-4858, 2012.

[34] T. Broicher, P. Malerba, A. D. Dorval, A. Borisyuk, F. R. Fernandez and J. A. White, "Spike phase locking in CA1 pyramidal neurons depends on background conductance and firing rate," *Journal of Neuroscience,* vol. 32, pp. 14374-14388, 2012.

[35] M. J. Richardson, N. Brunel and V. Hakim, "From subthreshold to firing-rate resonance," *Journal of Neurophysiology,* vol. 89, pp. 2538-2554, 2003.

[36] E. Stark, A. Levi and H. G. Rotstein, "Network resonance can be generated independently at distinct levels of neuronal organization," *PLoS computational biology, ,* vol. 18, p. e1010364, 2022 .

[37] B. Hutcheon and Y. Yarom, "Resonance, oscillation and the intrinsic frequency preferences of neurons," *Trends in Neurosciences,* vol. 23 , pp. 216-222, 2000 .

[38] R. F. Pena and H. G. Rotstein, "The voltage and spiking responses of subthreshold resonant neurons to structured and fluctuating inputs: persistence and loss of resonance and variability," *Biological Cybernetics,* vol. 116, pp. 163-190, 2022.

[39] R. F. Pena, C. C. Ceballos, V. Lima and A. C. Roque, "Interplay of activation kinetics and the derivative conductance determines resonance properties of neurons," *Physical Review E,* vol. 97, p. 042408, 2018.

[40] C. Yue, S. Remy, H. Su, H. Beck and Y. Yaari, "Proximal persistent Na+ channels drive spike afterdepolarizations and associated bursting in adult CA1 pyramidal cells," *Journal of Neuroscience,* vol. 25(42), pp. 9704-9720, 2005 .

[41] C. C. Ceballos, A. C. Roque and R. M. Leao, "A negative slope conductance of the persistent sodium current prolongs subthreshold depolarizations," *Biophysical Journal,* vol. 113, pp. 2207-2217, 2017.

[42] C. C. Ceballos, A. C. Roque and R. M. Leão, "The role of negative conductances in neuronal subthreshold properties and synaptic integration," *Biophysical Reviews,* vol. 9, pp. 827-834, 2017.

[43] . Kepecs and J. Lisman, "Information encoding and computation with spikes and bursts," *Network: Computation in Neural Systems,* vol. 14, pp. 103-118, 2003.

[44] J. E. Lisman, "Bursts as a unit of neural information: making unreliable synapses reliable," *Trends in Neurosciences,* vol. 20, pp. 38-43, 1997.

[45] K. D. Piatkevich, S. Bensussen, H. A. Tseng, S. N. Shroff, V. G. Lopez-Huerta, D. Park, ... and X. Han, "Population imaging of neural activity in awake behaving mice," *Nature,* vol. 574, pp. 413-417, 2019.

[46] J. I. Kass and I. M. Mintz, "Silent plateau potentials, rhythmic bursts, and pacemaker firing: three patterns of activity that coexist in quadristable subthalamic neurons," *Proceedings of the National Academy of Sciences,* vol. 103, pp. 183-188, 2006.

[47] J. C. Magee, "Dendritic hyperpolarization-activated currents modify the integrative properties of hippocampal CA1 pyramidal neurons," *Journal of Neuroscience,* vol. 18, pp. 7613-7624, 1998.

[48] A. Lörincz, T. Notomi, G. Tamás, R. Shigemoto and Z. Nusser, "Polarized and compartment-dependent distribution of HCN1 in pyramidal cell dendrites," *Nature Neurosciences,* vol. 5, p. 1185–1193, 2002.

[49] E. Campanac, G. Daoudal, N. Ankri and D. Debanne, "Downregulation of dendritic Ih in CA1 pyramidal neurons after LTP," *Journal of Neuroscience ,* vol. 28, pp. 8635-8643, 2008.

[50] C. C. Ceballos, R. F. Pena and A. C. Roque, "Impact of the activation rate of the hyperpolarization-activated current I h on the neuronal membrane time constant and synaptic potential duration," *The European Physical Journal Special Topics,* vol. 230, pp. 2951-2961, 2021.

[51] G. A. Ascoli, S. Gasparini, V. Medinilla and M. Migliore, " Local control of postinhibitory rebound spiking in CA1 pyramidal neuron dendrites," *Journal of Neuroscience,* vol. 30, pp. 6434-6442, 2010.

[52] C. C. Ceballos, S. Li, A. C. Roque, T. Tzounopoulos and R. M. Leão, "Ih equalizes membrane input resistance in a heterogeneous population of fusiform neurons in the dorsal cochlear nucleus," *Frontiers in Cellular Neuroscience,* vol. 10, p. 249, 2016.

[53] G. Hahn, A. F. Bujan, Y. Frégnac, A. Aertsen and A. Kumar, "Communication through resonance in spiking neuronal networks," *PLoS Computational Biology,* vol. 10, p. e1003811, 2014.

[54] J. S. Sherfey, S. Ardid, J. Hass, M. E. Hasselmo and N. J. Kopell, "Flexible resonance in prefrontal networks with strong feedback inhibition," *PLoS Computational Biology,* vol. 14, p. e1006357, 2018.

[55] J. Sherfey, S. Ardid, E. K. Miller, M. E. Hasselmo and N. J. Kopell, "Prefrontal oscillations modulate the propagation of neuronal activity required for working memory," *Neurobiology of Learning and Memory,* vol. 173, p. 107228, 2020.

[56] R. F. Pena, V. Lima, R. O. Shimoura, C. C. Ceballos, H. G. Rotstein and A. C. Roque, "Asymmetrical voltage response in resonant neurons shaped by nonlinearities," *Chaos: An Interdisciplinary Journal of Nonlinear Science, ,* vol. 29, p. 103135, 2019.

[57] G. S. Newkirk, D. Guan, N. Dembrow, W. E. Armstrong, R. C. Foehring and W. J. Spain, "Kv2. 1 potassium channels regulate repetitive burst firing in extratelencephalic neocortical pyramidal neurons," *Cerebral Cortex,* vol. 32, pp. 1055-1076, 2022.

[58] S. Chen, H. Su, C. Yue, S. Remy, M. Royeck, D. Sochivko, ... and Y. Yaari, "An increase in persistent sodium current contributes to intrinsic neuronal bursting after status epilepticus," *Journal of Neurophysiology,* vol. 105, pp. 117-129, 2011 .

[59] C. L. Hsu, X. Zhao, A. D. Milstein and N. Spruston, "Persistent sodium current mediates the steep voltage dependence of spatial coding in hippocampal pyramidal neurons," *Neuron,* vol. 99, pp. 147-162, 2018.

[60] R. Roy and R. Narayanan, "Ion-channel degeneracy and heterogeneities in the emergence of complex spike bursts in CA3 pyramidal neurons," *The Journal of Physiology,* vol. 601, pp. 3297-3328, 2023.

[61] K. A. Alkadhi and L. M. Tian, "Veratridine-enhanced persistent sodium current induces bursting in CA1 pyramidal neurons," *Neuroscience,* vol. 71 , pp. 625-632, 1996.

[62] P. Müller, A. Draguhn and A. V. Egorov, "Persistent sodium current modulates axonal excitability in CA 1 pyramidal neurons," *Journal of Neurochemistry,* vol. 146, pp. 446-458, 2018.

[63] L. Dalla Porta, A. Barbero-Castillo, J. M. Sanchez-Sanchez, N. Cancino and M. V. Sanchez-Vives, "H-current modulation of cortical Up and Down states," *bioRxiv,* pp. 2024-04, 2024.

[64] H. C. Pape, "Queer current and pacemaker: the hyperpolarization-activated cation current in neurons," *Annual Review of Physiology,* vol. 58, pp. 299-327, 1996.

[65] C. He, F. Chen, B. Li and Z. Hu, "Neurophysiology of HCN channels: from cellular functions to multiple regulations," *Progress in Neurobiology, ,* vol. 112, pp. 1-23, 2014.

[66] G. Drion, T. O'Leary and E. Marder, "Ion channel degeneracy enables robust and tunable neuronal firing rates.," *Proceedings of the National Academy of Sciences,* vol. 112, pp. E5361-E5370, 2015.

[67] F. S. Borges, P. R. Protachevicz, D. L. Souza, C. F. Bittencourt, E. C. Gabrick, L. E. Bentivoglio, J. D. Szezech, A. Batista, I. Caldas, S. Dura-Bernal and R. F. Pena, "The Roles of Potassium and Calcium Currents in the Bistable Firing Transition," *Brain Sciences,* vol. 13, p. 1347, 2023.

[68] P. R. Protachevicz, F. S. Borges, E. L. Lameu, P. Ji, K. C. Iarosz, A. L. Kihara, I. L. Caldas, J. D. J. Szezech, M. S. Baptista, E. E. N. Macau, C. G. Antonopoulos, A. M. Batista and J. Kurths, "Bistable firing patterns in a neural network model," *Frontiers in Computational Neuroscience,* vol. 13, p. 19, 2019.

[69] J. Yamada-Hanff and B. P. Bean, "Activation of Ih and TTX-sensitive sodium current at subthreshold voltages during CA1 pyramidal neuron firing," *Journal of Neurophysiology,* vol. 114, pp. 2376-2389, 2015.

[70] A. Compte, M. V. Sanchez-Vives, D. A. McCormick and X. J. Wang, "Cellular and network mechanisms of slow oscillatory activity (< 1 Hz) and wave propagations in a cortical network model," *Journal of Neurophysiology,* vol. 89, pp. 2707-2725, 2003.

[71] C. Li, Y. T., M. M. Poo and Y. Dan, "Burst spiking of a single cortical neuron modifies global brain state," *Science,* vol. 324, pp. 643-646, 2009 .

[72] R. Azouz, M. S. Jensen and Y. Yaari, "Ionic basis of spike after-depolarization and burst generation in adult rat hippocampal CA1 pyramidal cells," *The Journal of Physiology,* vol. 492 , pp. 211-223, 1996.

[73] M. Bazhenov, I. Timofeev, M. Steriade and T. J. Sejnowski, "Potassium model for slow (2-3 Hz) in vivo neocortical paroxysmal oscillations," *Journal of Neurophysiology,* vol. 92, pp. 1116-1132, 2004.

[74] R. Bertram, M. J. Butte, T. Kiemel and A. Sherman, "Topological and phenomenological classification of bursting oscillations," *Bulletin of Mathematical Biology,* vol. 57, pp. 413-439, 1995.

[75] I. A. Fleidervish, A. Friedman and M. J. Gutnick, "Slow inactivation of Na+ current and slow cumulative spike adaptation in mouse and guinea-pig neocortical neurones in slices," *The Journal of Physiology,,* vol. 493, pp. 83-97, 1996.

[76] A. Kepecs and X. J. Wang, "Analysis of complex bursting in cortical pyramidal neuron models," *Neurocomputing,* vol. 32, pp. 181-187, 2000.

[77] S. Xiao, E. Lowet, H. J. Gritton, P. Fabris, Y. Wang, J. Sherman, ... and X. Han, "Large-scale voltage imaging in behaving mice using targeted illumination," *Iscience,* vol. 24, 2021.

[78] E. Lowet, K. Kondabolu, S. Zhou, R. A. Mount, Y. Wang, C. R. Ravasio and X. Han, "Deep brain stimulation creates informational lesion through membrane depolarization in mouse hippocampus," *Nature Communications,* vol. 13, p. 7709, 2022.

[79] S. Xiao, W. J. Cunningham, K. Kondabolu, E. Lowet, M. V. Moya, R. A. Mount, ... and J. Mertz, "Large-scale deep tissue voltage imaging with targeted-illumination confocal microscopy," *Nature Methods ,* pp. 1-9, 2024.

[80] J. P. Lachaux, A. Lutz, D. Rudrauf, M. Cosmelli, J. Le Van Quen, F. Martinerie and F. Varela, "Estimating the time-course of coherence between single-trial brain signals: an introduction to wavelet coherence," *Clinical Neurophysiology,* vol. 32, p. 157–174, 2002.

[81] M. Vinck, M. van Wingerden, T. Womelsdorf, P. Fries and C. M. A. Pennartz, "The pairwise phase consistency: A bias-free measure of rhythmic neuronal synchronization," *Neuroimage,* vol. 51, p. 112–122 , 2010.

[82] S. Aydore, D. Pantazis and R. M. Leahy, "A note on the phase locking value and its properties," *Neuroimage ,* vol. 74, p. 231–244, 2013.

691 **Supporting information for**

693 **Interleaved single and bursting spiking resonance in neurons**

695 Cesar C. Ceballos[1], Nourdin Chadly[3,4], Eric Lowet[3], Rodrigo F. O. Pena[1,2,*]

696
697 [1] Department of Biological Sciences, Florida Atlantic University, Jupiter, FL 33458, USA
698 [2] Stiles-Nicholson Brain Institute, Florida Atlantic University, Jupiter, FL 33458, USA
699 [3] Department of Neuroscience, Erasmus Medical Center, Rotterdam, Netherlands
700 [4] Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, Netherlands

701
* Corresponding author
E- mail: penar@fau.edu

Fig S1. Ionic currents effect on neuronal response during sustained depolarization. Firing
rate and coefficient of variation (CV), a measure the activity of the neuron and the presence of
bursts, displayed as heatmaps spanned over different ionic current variation. **A** G_{KA} vs. Input
current. **B** G_{KM} vs. Input current. **C** G_{fKCa} vs. Input current. **D** G_{sKCa} vs. Input current.

712
713 **Fig S2. Ionic currents effect on neuronal response during oscillatory input.** Firing rate and
714 coefficient of variation (CV), a measure the activity of the neuron and the presence of bursts,
715 displayed as heatmaps spanned over different frequency input variation. **A** G_{KA} vs. Input current.
716 **B** G_{KM} vs. Input current. **C** G_{fKCa} vs. Input current. **D** G_{sKCa} vs. Input current. **E** Voltage traces of
717 control vs after blocking M-type potassium current.

Fig S3. Time evolution of activation variables. A Voltage trace showing interleaved single spiking and bursting during oscillatory stimulation. Asterisks represent spike peaks. **B.** Time evolution of activation variables of I_{KDR}, I_{KM} and I_{NaP}. Note the different temporal scales.

730
731
732

Fig S4. Effect of noise on interleaved resonance. Default values for fixed conductance while varying the other are $G_{NaP} = 0.93$, $G_{KDR} = 5.86$, and $G_h = 8.46$. **A1-2** Δf and ΔA for varying values of G_{NaP}. **B1-2** Δf and ΔA for varying values of G_{KDR}. **C1-2** Δf and ΔA for varying values of G_h. Colors mark the magnitude of noise (blue = low, orange = medium, green = high).

Fig S5. Somatic and dendritic traces under gamma and theta oscillations input. Somatic
traces (top) and dendritic traces (bottom).

752
753

Fig S6. Intermediate G_{NaP} and G_{KDR} generates firing rate resonance even at low input amplitudes. Same simulation as in Fig 2, but for lower input amplitudes (2x). We have also raised the holding potential by to be closer to threshold for this experiment (offset amplitude: 2 μA/cm^2) **A** Firing rate while varying stimulation frequency of an oscillatory input for $G_{KDR} = 5$. **B** Firing rate while varying stimulation frequency of an oscillatory input for $G_{NaP} = 0.35$.