Linalool's Multifaceted Antimicrobial Potential: Unveiling its Antimicrobial Efficacy and Immunomodulatory Role Against *Saprolegnia parasitica*

Tao Tang[1], Weiming Zhong[1], Puyu Tang[3], Rongsi Dai[1], Jiajing Guo[2]* and Zhipeng Gao[1]**

[1] Hunan Engineering Technology Research Center of Featured Aquatic Resources Utilization, Fisheries College, Hunan Agricultural University

Changsha, Hunan, 410128, China

E-mail: gaozhipeng627@163.com

[2] Hunan Agriculture Product Processing Institute, Hunan Academy of Agricultural Sciences

Changsha, Hunan, 410125, China

E-mail: guojiajing1986@163.com

[3] State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of Hydrobiology, Chinese Academy of Sciences

Wuhan, Hubei, 430072, China

Abstract

Saprolegnia parasitica (*S. parasitica*) is one of the most virulent oomycete species in freshwater aquatic environments, causing severe saprolegniasis and leading to significant economic losses in the aquaculture industry. Thus far, the prevention and control of saprolegniasis face a shortage of medications. Linalool, a natural antibiotic alternative found in various essential oils, exhibits promising antimicrobial activity against a wide range of pathogens. In this study, the specific role of linalool in protecting *S. parasitica* infection at both *in vitro* and *in vivo* levels was investigated. Linalool showed multifaceted anti-oomycetes potential by both of antimicrobial efficacy and immunomodulatory efficacy. For *in vitro* test, linalool exhibited strong anti-oomycetes activity and mode of action included: (1) Linalool disrupted the cell membrane of the mycelium, causing the intracellular components leak out; (2) Linalool prohibited ribosome function, thereby inhibiting protein synthesis and ultimately affecting mycelium growth. Surprisingly, meanwhile we found the potential immune protective mechanism of linalool in the *in vivo* test: (1) Linalool enhanced the complement and coagulation system which in turn activated host immune defense and lysate *S. parasitica* cells; (2) Linalool promoted wound healing, tissue repair, and phagocytosis to cope with *S. parasitica* infection; (3) Linalool positively modulated the immune response by increasing the abundance of beneficial Actinobacteriota; (4) Linalool stimulated the production of inflammatory cytokines and chemokines to lyse *S. parasitica* cells. In all, our findings showed that linalool possessed multifaceted anti-oomycetes potential which would be a promising natural antibiotic alternative to cope with *S. parasitica* infection in the aquaculture industry.

Keywords: *Saprolegnia parasitica*, linalool, anti-oomycetes mechanism, immune protective mechanism, antibiotic alternative

1. Introduction

Saprolegniasis is one of the most serious oomycete diseases in the aquaculture industry all over the world, infecting a wide host range (fish, crustaceans, amphibians, and their eggs et al.) with a conspicuous symptom (grayish-white cotton wool-like covering on the surface of host)[1, 2]. Oomycete infections lead to significant economic losses and serious damage to natural ecosystems[1]. As the infection intensifies, saprolegniasis can cause muscle ulceration, compromised immune system, respiratory problems, secondary infection, and finally result in host death[3]. *Saprolegnia*, *Achlya*, and *Leptolegnia* are key pathogens responsible for Saprolegniasis outbreaks[4], among them, *Saprolegnia* is proven to be the most virulent[5, 6]. *S. parasitica*, *S. ferax*, *S. delica*, and *S. diclina* are the main pathogens causing saprolegniasis[7], as a highly virulent group belonging to the phylum Oomycota, *S. parasitica* serves as the primary species in fish infections[8].

In the past, malachite green was highly effective in controlling *saprolegnia* infections, however, it has been banned worldwide for its carcinogenic and toxicological effects, which led to a sharp re-emergence of *Saprolegnia* infections in aquaculture industry[9]. Recently, various methods have been established to deal with *S. parasitica* infections, mainly including physical (water quality management, temperature control, and so on), chemical (hydrogen peroxide, natural antimicrobial compounds, and so on), and biological (probiotics, antagonistic microbe, and so on) methods[10]. Among these, antibiotics have served as the primary therapeutic and prophylactic tools for Saprolegniasis management in aquaculture. However, their overuse and abuse have resulted in antibiotic resistance among various aquatic pathogens[11]. Therefore, the increasing need for sustainable and eco-friendly strategies to combat *saprolegnia* infections has driven research into natural antimicrobial agents, due to their advantages in terms of avoiding antibiotic resistance issues, health safety, and ecological sustainability[12].

In our earlier studies, the anti-oomycetes activities of 12 plant essential oils (EOs) and 5 of their major compounds (linalool, pinene, limonene, myrcene, and terpinene)

were evaluated, and we found that linalool was one of the strongest anti-oomycetes

compounds and selected for further study. Linalool, a monoterpene alcohol found in

various EOs, has shown promising antimicrobial activity against a wide range of

pathogens[13]. Our groups and other research teams have reported that linalool exhibits

antimicrobial activity against bacteria such as *Escherichia coli*[12], *Listeria*

monocytogenes[14], *Elizabethkingia miricola*, *Streptococcus pyogenes*[15],

Pseudomonas fragi[16], *staphylococcus aureus*[17], *Brochothrix thermosphacta*[18] and

Aeromonas hydrophila[19], as well as against oomycetes like *Colletotrichum*

lagenarium[20] and *Phytophthora capsici*[21], and fungi including *Aspergillus flavus*[22],

Penicillium citrinum, *Chaetomium globosum*[23], *Colletotrichum gloesporioides*,

Fusarium oxysporum[24], *Botrytis cinerea*[25] and *Candida albicans*[26]. Although the

antimicrobial activity of linalool has been extensively studied *in vitro*, the potential

immune mechanism *in vivo* remains poorly understood.

Therefore, in this study, for the first time, we aimed to investigate the specific role

of linalool in protecting *S. parasitica* infection at both of *in vitro* and *in vivo* level, and

hope to unravel its potential molecular mechanisms through comprehensive analyses,

including *in situ* fluorescence observation, microstructure observation, transcriptome

profiling, histological examinations, gut microbiota analysis, and deep bioinformatics

analysis. We believe this study will shed light on the comprehensive of anti-oomycete

of linalool and contribute to the development of novel antibiotic alternative strategies

to cope with *S. parasitica* infection in aquaculture industry.

2. Results

2.1 The *in vitro* anti-oomycete activity of linalool against *S. parasitica* CQT2

Figure 1A illustrated the anti-oomycetes activities of EOs and their major

components against *S. parasitica* CQT2. Notably, linalool and the EO extracted from

daidai flowers were particularly effective in restraining mycelium growth.

Linalool exhibited the MIC and MFC of 0.025% and 0.025% (v/v, 0.216 mg/mL)

against spores, respectively (Figure 1B). Besides, both the MIC and MFC against

mycelium growth were 0.05% (v/v, 0.432 mg/mL) and 0.1% (v/v, 0.864 mg/mL),

respectively. Various linalool doses were examined against *S. parasitica* CQT2 on linalool-containing plates to evaluate their impact on inhibiting mycelium growth. As shown in Figure 1C-D, the findings indicated the anti-oomycetes activity of linalool against *S. parasitica* CQT2 was dose-dependent. Figure 1E displayed mycelium growth on PDA plates under varying linalool concentrations. The viability of mycelium following linalool treatment was assessed using a PI staining assay, where non-viable (dead) mycelium appeared red after PI uptake. Untreated mycelium showed no detectable red fluorescent, indicating 100% viability. In contrast, distinct red fluorescent mycelium was observed after linalool treatment at both MIC and 2×MIC levels (Figure 1F).

2.2 Effects of linalool on cell membrane damage of *S. parasitica* CQT2

The *in situ* fluorescence observations were displayed in Figure 2A, with linalool treatment for 0 minutes, the mycelium structure remained undamaged, and the red and green fluorescence showed an even distribution. After 2 min, red fluorescence started appearing within the mycelium, accompanied by the entry of the styryl dye FM4-64, indicating damage to the cell membrane. As the duration of linalool treatment increased, the intensity of red fluorescence within the mycelium also increased, indicating a rise in the number of mycelial cell deaths along with severe damage to the cell membrane. When examined by SEM, the mycelium exhibited a smooth and intact surface in the control group (Figure 2B), while in the treated groups (Figure 2C), it appeared wrinkled, deformed, and shrunken (Red dashed quadrilateral). When examined by TEM, the untreated mycelium cell exhibited clear and intact ultrastructure in the cell membrane, cell wall, mitochondria, and rough endoplasmic reticulum (Figure 2D). However, after treatment with linalool, the cell membrane was wrinkled and seriously damaged, and mitochondria began to dissolve (Figure 2E). As shown in Figure 2F-G, the cell membrane structure disappeared, the cell wall became thinner, and some exhibited severe deformation and bending. Vacuoles appeared in the cytoplasm, and mitochondria and rough endoplasmic reticulum ruptured and dissolved (red circles). No intact organelles were present within the cytoplasm.

As shown in the GO classification analysis (Figure 2H), the cellular components

were distributed across 8 groups, with "cell part" and "membrane part" being the most

prevalent. The DEPs related to intrinsic component of membrane were screened out,

categorizing them into 4 classes: related to ribosome function, related to DNA/RNA

function, related to signal transduction and regulation, and related to cell metabolism

(Figure 2I). Therefore, the cell membrane may be a potential target for linalool.

2.3 The *in vitro* anti-oomycete mechanisms revealed by transcriptome analysis

As shown in Figure S1A-B, a total of 4142 differentially expressed genes (DEGs)

were discovered between the control and treatment groups, with 2016 showing

upregulation and 2108 downregulation. Figure 2H presented the Gene Ontology (GO)

analysis, providing a comprehensive evaluation of the functional aspects of genes and

their products. These DEGs were categorized into 20 GO categories, with 6 associated

with biological activities, where cellular and metabolic functions had the most DEGs.

There were also 6 molecular function categories, with catalytic activity and binding

containing the majority of DEGs. Additionally, Figure S1C demonstrated the results

of Kyoto Encyclopedia of Genes and Genomes (KEGG) annotation, revealing 10

metabolic (M) pathways, 2 environmental information processing (EIP) pathways, 2

cellular processes (CP), and 5 genetic information processing (GIP).

2.3.1 The ribosome biogenesis and protein synthesis of *S. parasitica* were

prohibited by linalool

Figure 3A-B displayed the results of GO enrichment analysis, revealing the top 20

biological processes that differed significantly between the control and treated groups.

Notably, in Figure 3B, the downregulated biological processes are highlighted, and

remarkably, 19 of them were associated with "ribosome", such as ribosome

biogenesis, ribonucleoprotein complex biogenesis, pre ribosome, and small-subunit

process some. In addition, KEGG enrichment analysis in Figure 3C revealed that

"Ribosome biogenesis in eukaryotes" was the most enriched pathway, aligning with

the GO enrichment analysis results mentioned earlier. The pathway of ribosome

biogenesis in eukaryotes (Figure 3D) showed that linalool influenced DNA

transcription, tRNA transport, rRNA processing and maturation (5.8S, 18S, and 25S),

and the biogenesis and assembly of ribosome subunits (40S and 60S) in the cell,

which might lead to the reduction of *S. parasitica* growth. We found that many DEGs

related to ribosome biogenesis and RNA polymerase (Figure 3E) are down expressed.

Figure 3A highlighted the upregulated biological processes, and notably, 11 of them

were linked to "amino acids", such as the sulfur amino acid, L-phenylalanine, alpha-

amino acid, tyrosine, and homocysteine. Amino acids serve essential roles within the

cell, acting as both the foundational components for synthesizing new proteins and as

precursors for various metabolic processes[27].

Molecular docking was carried out to explore the specific binding protein of

linalool with ribosome. The binding energy of linalool was -6.2 kcal/mol with NOP, -

4.8 kcal/mol with SNU13, and -5.0 kcal/mol with DKC1 (Table S1). Consequently,

NOP1 was selected as the receptor protein for subsequent molecular docking. Figure

3F illustrated the tertiary structure of ribosomal protein NOP1. Linalool formed

hydrogen bonds with glutamic acid (Glu155) and phenylalanine (Phe156) of NOP1

while exhibiting hydrophobic interactions with alanine (Ala181), arginine (Arg182),

valine (Val201), and isoleucine (Ile209) (Figure 3G). In summary, this interference

may hinder the ribosome's ability to execute normal biological processes, such as

protein synthesis, thereby impacting the growth and reproduction of oomycetes.

2.3.2 The global metabolisms of *S. parasitica* were regulated by linalool

In KEGG enrichment analysis, we found that 6 processes were associated with

metabolic pathways, such as cysteine and methionine metabolism, starch and sucrose

metabolism, and sulfur metabolism. Moreover, metabolic pathways were the most

annotated pathway which was in line with the results of KEGG annotation analysis.

As many changes were linked to metabolism, the Interactive Pathways Explorer

(iPath) was employed to comprehend the global differential metabolic response in *S.*

parasitica CQT2 (Figure 3I) which identified that linalool affects the global metabolic

regulation of *S. parasitica* CQT2.

2.3.3 ABC transporters of *S. parasitica* were affected by linalool

Some other important pathways were significantly enriched in our study, such as ABC transporter and sulfur relay system. The DEGs involved in ABC transporters (ABCA3, ABCB1, ABCC1, ABCG2, and PDR5) were shown in Figure 3H. In brief, these results demonstrated that cell membrane, protein synthesis, functions related to metabolism, and ribosome were significantly impacted by linalool and were potential drug targets of linalool.

2.4 The *in vivo* protective effect of linalool against *Saprolegniosis*

To study the protective effects of linalool on *S. parasitica* infection in fish, linalool was added to grass carp after *S. parasitica* infection (linalool therapeutic group, LT) or 48 h before infection (linalool prophylactic group, LP), and a positive control group (PC) without linalool and a negative control group (NC) without infection and linalool were set up (Figure 4A). The results showed that the survival rate of *S. parasitica* infected grass carp was significantly improved after treatment with linalool (Figure 4C), which proved greatly anti-*Saprolegnia* activity of linalool. In Figure 4B, there was no observable mycelium in both LT and LP groups, while the PC group displayed significant mycelium growth along with inflammation on the dorsum. The LZM activity was significantly increased in LP and LT group on the seventh day (Figure 4E), yet the AKP activity (Figure 4D) was reduced. However, there was no significant difference in the SOD activity (Figure 4F) between LP and LT groups during the whole feeding trial.

To explore the histomorphology alterations caused by *S. parasitica* infection and linalool treatment, we performed histomorphology evaluations through HE staining (Figure 4G). The findings revealed significant pathological changes in the kidney, liver, spleen, and skin of grass carp due to *S. parasitica* infection. In PC group, severe kidney damage was evident, characterized by enlarged glomeruli, glomerular cysts, interstitial bleeding, and infiltration of inflammatory cells. In addition, extensive vacuolation and expansion of intercellular space was observed in hepatocytes, with nuclei displaced toward one side. Cytoplasmic pyknosis and structural disintegration were observed in the pancreatic acinar cells. Meanwhile, the red and white pulp of the

spleen was poorly demarcated, the splenic corpuscle was irregularly shaped and arranged, and the white pulp was structurally disrupted. Furthermore, there was a notable increase in the number of macrophages and an expansion in the size of melanoma-macrophage centers. *S. parasitica* not only caused damage to both the dermal and epidermal layers but also induced muscle cell degeneration. Continuous treatment of linalool for 7 days significantly improved kidney, liver, spleen, and skin morphological integrity.

In the LP and LT groups, linalool alleviated tissue damage caused by *S. parasitica* infection on the dorsal surface of grass carp, enhancing the healing capacity (Figure 4B). Histomorphology demonstrated that linalool could improve the integrity of organ morphology in grass carp (Figure 4G). Transcriptomics of the spleen revealed that linalool promoted wound healing, tissue repair, and phagocytosis to cope with *S. parasitica* infection (Figure 5E). In all, our results indicated that linalool promoted wound healing in grass carp.

2.5 The *in vivo* Saprolegniasis protective mechanism of linalool revealed by comparative transcriptome analysis

To analyze the contrasting gene-level responses in grass carp spleens, a vital immune organ, following *S. parasitica* infection and linalool treatment, spleen samples were collected for transcriptome analysis. Compared to PC group, 2029 genes in LT group exhibited significant difference, with 809 DEGs up regulated and 1220 down regulated. Meanwhile, 671 genes were identified as significantly different, with 609 DEGs up-regulated and 618 downregulated in LP group relative to PC group (Figure S2A-B).

Based on the GO annotation analysis, a total of 2219 unigenes in the LT group and 520 unigenes in the LP group were classified into 3 primary functional categories (Figure S2C). Within the biological process category, "cellular process," "metabolic process," and "biological regulation" were the most prominent subcategories. In terms of cellular components, "membrane part" and "cell part" were highly represented. "Binding" and "catalytic activity" were the most prevalent subcategories within

molecular functions. Regarding KEGG annotations, the largest cluster within

organismal systems was the immune system, encompassing 209 genes (42.47%) in

the LT group (Figure S2D) and 38 genes (50.00%) in the LP group (Figure S2E). In

the environmental information processing category, most unigenes were associated

with signal transduction, with 227 (65.60%) in the LT group and 38 genes (52.00%) in

the LP group. These results suggested that the immune response and signal

transduction pathways were crucial for grass carp in dealing with *S. parasitica*

infections.

The GO enrichment analysis of the spleens illustrated in Figure 5A-B to reveal

distinct biological process. Those down regulated biological process were shown in

Figure 5A, compared to PC group, the "positive regulation of immune system

process" term (GO:0002684) and "regulation of body fluid levels" term

(GO:0050878) were enriched with the greatest number of DEGs in LT group. It is

worth noting that there are 9 terms related to "body fluids (especially blood)", such as

regulation of coagulation (GO:0050818), blood coagulation (GO:0007596), and

regulation of hemostasis (GO:1900046). In addition, those up regulated biological

process, the "ion binding" term (GO:0043167) was enriched with the greatest number

of DEGs in LT group. Figure 5B illustrated the GO terms with the largest number of

down regulated DEGs in LP group was "small molecule binding" (GO:0036094).

KEGG enrichment analysis was used to explore the functional pathways involving

DEGs. The 20 significantly enriched up regulated and down regulated pathways of

each group are shown in Figure 5C-D. Multiple pathways associated with immune

response were enriched in down regulated genes after linalool treatment in LT group,

including complement and coagulation cascades, Toll-like receptor signaling pathway,

IL-17 signaling pathway, C-type lectin receptor signaling pathway, T cell receptor

signaling pathway, B cell receptor signaling pathway, and so on (Figure 5C). Toll-like

receptor signaling pathway, RIG-I-like receptor signaling pathway, Complement and

coagulation cascades, and Chemokine signaling pathway were enriched with mostly

up-regulated genes. Furthermore, metabolic pathways were enriched with mostly

down-regulated genes in LP group. Some pathways associated with immune response were significantly enriched, such as Antigen processing and presentation, Fc epsilon RI signaling pathway, Intestinal immune network for IgA production, and so on (Figure 5D). GO and KEGG enrichment analyses revealed the presence of numerous immune-related genes within corresponding functional processes and pathways. The expression of *fb*, *c1q*, *c2*, *c3*, *c4*, *c6*, *c7*, *c8A*, *c8b*, and *c9* genes were up expressed. Additionally, there was a significant up expressed in the gene expression levels of coagulation factors II (thrombin), VII, VIIi, VIII, X, coagulation factor IXb and IXa, serpin peptidase inhibitor clade C/D/G/F, *fga*, *fgb*, *fgg*, protein C, plasminogen, and plasminogen activator (Figure 5E). The differential expression of immune-related DEGs implied their role in host defense against *S. parasitica* infection.

To ensure the regulation of immune-related DEGs by linalool, the expression of several vital immune related genes (*cfh*, *masp2*, *c1r*, *c3*, *cfb*, *ap-1*, *il-1β*, and *il-6*) in different groups were tested by RT-qPCR assay. The results showed that the expression of these genes in the LP and LT groups significantly exceeded those in the PC group (P < 0.05) (Figure 5H). Further exploration of immune-related genes indicated that linalool treatment promoted the expression of genes linked to the complement system and inflammatory factors. Additionally, analysis of 8 genes revealed that the real-time fluorescence quantitative analysis results were consistent with the expression trend of transcriptome analysis genes.

2.6 Linalool regulated gut microbiota composition

In our study, 862 OTUs were obtained from 12 samples through 16S sequencing analysis, with 48 common OTUs shared across all samples (Figure 6E). Alpha diversity analysis (Figure 6A-D) showed that the Simpson, Shannon, Chao1, and Ace index of gut microbiota decreased after linalool treatment which indicated that linalool reduced the abundance and diversity of gut microbes. PCoA analysis showed a significant difference in microbial composition among the PC, LP, and LT groups (Figure 6F).

The composition of gut microbiota in Figure 6G showed that the main dominant

bacteria in the intestinal tract are Proteobacteria, Actinobacteriota, Verrucomicrobiota,

Firmicutes, and Bacteroidota at the phylum level. Among them, compared with PC

group, an obvious tendency of greater relative abundances of phyla Proteobacteria and

Actinobacteriota was evidenced in LP and LT groups, the relative abundance of

Verrucomicrobiota was decreased. From the composition of the genus level, the

Neochlamydia decreased, while *Achromobacter* increased. It is worth noting that the

genus *Aurantimicrobium* (classified as Actinobacteriota) was increased in linalool

prophylactic and linalool therapeutic groups. Notably, among biomarkers with an

LDA SCORE exceeding 3, the genus *Actinomycetales* bacterium held the highest

number of positions, which contributed to the major differences between the positive

control and linalool treatment groups. Moreover, the potential metabolic pathways of

gut microbiota primarily showed enrichment in functions related to metabolism,

including carbohydrate, amino acid, and cofactors and vitamins metabolism (Figure

6I).

2.7 Correlation analysis of potential related immune genes, specific microbial

species, and physical characteristics

Correlation analysis showed that complement system were significantly positively

correlated to Actinobacteriota, AKP, and SR (survival rate), which, in turn, a

significantly negatively correlated with Verrucomicrobiota. Additionally, there was a

significant positive correlation between PPRs and AKP. Notably, IgA related immune

genes were correlated significantly with Proteobacteria (Figure 6J). In summary, our

findings suggested that linalool effectively modulated gut microbiota during *S.*

parasitica infection which was associated with immune response.

3. Discussion

3.1 *In vitro* mode of action of linalool against *S. parasitica*

3.1.1 Membrane disruption

The cell membrane maintains cell integrity, regulates substance entry and exit,

while the cell wall provides structural support, protection, and promotes growth and

cell-to-cell connections[28]. The mycelium forms a dense structure to protect *S.*

parasitica spores from external factors. CLSM, SEM, and TEM observations

suggested that the cell membrane of the mycelium treated with linalool was damaged,

causing the intracellular components to leak out, and could not develop completely

(Figure 2A-G). Meanwhile, 678 genes related to the "intrinsic component of

membrane" showed significant expressed in GO enrichment analysis, suggesting that

linalool may enhance the expression of intracellular membrane components, reflecting

its impact on cell membrane structure or function. Gao et al. illustrated that linalool

caused significant damage to the cell wall/membrane of *E. coli*, resulting in the lysis

and death of bacterial cells[12]. Gao et al. demonstrated that linalool[14], *Changshan-*

huyou Y.B. Chang EO[29], and Fingered Citron EO[30] can disrupt the cell membrane of

L. monocytogenes. Lemon essential oil nanoemulsions altered the permeability of *E.*

coli cell membrane, affecting membrane potential, integrity, and efflux function[31].

Cruz et al suggested that alkamides in *Echinacea* disrupted the cell wall/membrane

complex, making it an ideal target for specifically inhibiting pathogens[32]. Hinokitiol

caused mycelium malformation by inducing cell component leakage and influencing

chitinase activity[33]. Therefore, linalool may act anti-oomycete effects against *S.*

parasitica by targeting the cell membrane.

ABC transporters utilize ATP hydrolysis energy to facilitate the export of harmful

compounds or essential lipids across bacterial and eukaryotic membranes[34]. The

DEGs involved in ABC transporters (ABCA3, ABCB1, ABCC1, ABCG2, and PDR5)

in our study, almost all of which have been linked to illness states usually involving

loss of function[35]. PDR5-related proteins in fungi played a role in excreting

xenobiotics, including antifungal agents[36]. Multidrug resistance-associated protein

(MRP) transporters were typically associated with multidrug resistance, cellular

detoxification, and drug metabolism[37]. To sum up, linalool may greatly inhibit

mycelium growth by interfering with protein synthesis from the "inside" and

damaging the integrity and permeability of the cell membrane and wall from the

"outside."

3.1.2 Reduction of ribosome biogenesis and RNA polymerase

To further investigate the mechanism of linalool against *S. parasitica*, the gene expression profiles of *S. parasitica* were compared (with or without exposed to linalool). Many DEGs related to ribosome biogenesis and RNA polymerase were down expressed, indicating that the linalool may influence translation, replication, and repair processes, leading to increased protein diversity and potential disruption of corresponding functions[38]. Linalool influenced DNA transcription, tRNA transport, rRNA processing and maturation (5.8S, 18S, and 25S), and the biogenesis and assembly of ribosome subunits (40S and 60S) in the cell, which might lead to the reduction of *S. parasitica* growth. Ribosomes are assembled logistically by more than 200 ribosomal proteins (such as NOP, RIOK, and UTP) and RNA factors in a highly coordinated manner[39]. Depletion of Nop-7-associated 2 (NSA2) led to decreased nascent rRNA synthesis, reduced protein synthesis, and impacted cell proliferation due to its critical role in ribosome biogenesis regulation[40]. The inhibitory effect of myriocin on *Fusarium oxysporum* includes both membrane damage and interactions with intracellular targets, such as proteins like RIOK2, within the ribosome biogenesis pathway[41]. Innate immunity in *Caenorhabditis elegans* has been discovered to be negatively regulated by NOL-6 (also named UTP22)[42]. The growth of *Aspergillus niger* may be inhibited by cyclosporin A because it may prevent the transcription and translation of the RNA polymerase III complex[43]. Rifampicin can prevent RNA transcription in bacteria by interacting with the β-subunit of DNA-dependent RNA polymerase, which can then inhibit protein biosynthesis[44]. These results indicated that the down-regulation genes may limit the proteins necessary for normal cell biogenesis and growth. Therefore, ribosome is a promising potential target of linalool against *S. parasitica*.

3.1.3 The influence on the metabolism function of *S. parasitica*

Metabolism function is crucial for cells to maintain homeostasis, survive, and proliferate[45]. Linalool caused the activation of energy metabolism pathways, compensating for the insufficient energy needed for cell survival[46]. In our study,

linalool was found to enhance amino acid metabolism, carbohydrate metabolism,

energy metabolism, and so on in mycelial cells, which suggested that cells may boost

functions related to energy metabolism to acquire additional energy for repairing

damage caused by linalool. Hence, linalool treatment caused intracellular energy

metabolism disruption, resulting in impediments to protein synthesis, cellular signal

transduction, DNA synthesis, and ultimately affecting cell viability and proliferation.

3.2 *In vivo* mode of action of linalool against *S. parasitica* infection

Recent studies demonstrated that the spleen serves antibacterial functions in various

fish species, with a high quantity of melano-macrophages that assist in phagocytosis

during the immune response[47]. The rise of macrophages and the change in the

pathological and physiological circumstances in fish are correlated[48]. Similarly,

histopathological analysis demonstrated that linalool effectively reduced *S. parasitica*

infection-induced damage in the spleen, indicating its potential immunomodulatory

effects on immune organs. Here we used comparative transcriptome analysis to

investigate potential genes and signaling pathways regulated by linalool in the spleen

of grass carp. Several genes associated with both innate and adaptive immunity, such

as complement molecules, pattern recognition receptors, inflammatory cytokines,

immunoglobulins, and chemokines, were differentially expressed as shown in Figure

5E-G.

3.2.1 Enhancement of complement and coagulation system

It is widely acknowledged that the complement and coagulation system in fish is an

important immune defense mechanism for fighting pathogens and maintaining

immune homeostasis[49]. The complement system is made up of over 30 different

proteins (complement receptors, internal components, and regulatory proteins) and 3

activation pathways (the classic pathway, the lectin pathway, and the alternative

pathway)[50]. In our study, the expression of *fb*, *c1q*, *c2*, *c3*, *c4*, *c6*, *c7*, *c8a*, *c8b*, and

c9 genes were all up regulated after linalool therapeutic. Meanwhile, the c3a

anaphylatoxin chemotactic receptor (*c3aar1*) and mannan-binding lectin serine

protease 2 (*masp2*) genes were up expressed. During complement activation, *c3* and

c5 split into molecules such as *c3a*, *c3b*, *c5a*, and *c5b*, of which *c3a* and *c5a* are

considered strong inflammatory mediators because of their ability to attract

leukocytes and promote inflammation and pathogen clearance[51]. *masp2* activates

complement components such as *c4* and *c2*, triggering a complement cascade reaction

that ultimately leads to pathogen rupture[52]. *c8b* is a subunit of the complement

component 8 (*c8*) protein, which helps form the membrane attack complex (MAC) for

cell lysis and immune defense against microbes[53]. In addition, high expression of

complement factor H like 3 (*chfl3*) prevents overactivation of the complement system

to avoid damage to their own tissues[54]. The effective antimicrobial role of the *c5a-*

c5ar1 signaling pathway in the complement system has been identified. It functions

by enhancing neutrophil phagocytosis of fungi and macrophage-mediated fungicidal

activity to combat fungal infections[55]. In all, our findings suggested that linalool may

enhance the complement system which in turn activated host immune defense and

lysate *S. parasitica* cells.

The coagulation cascades are connected with the complement system. Linalool

significantly increased the expression of fibrinogen alpha (*fga*), beta (*fgb*), and

gamma (*fgg*) chain, which together encoded fibrinogen[56], fibrinogen is a crucial

component of blood clots to prevent blood loss, and also important in the initial stages

of wound healing[57]. Secondly, activated prothrombin *f2* is central in clot formation,

platelet aggregation, and activating other clotting factors. It also supports tissue repair,

cell growth, angiogenesis, and vascular integrity[53]. Thirdly, an increase in

plasminogen (PLG) levels can cause an immune response[58], and serve as a regulator

of the natural immune system by promoting the phagocytosis of phagocytes[59]. Taken

together, the upregulation of fibrinogen, prothrombin, and plasminogen in the

coagulation system suggested that linalool may promote wound healing, tissue repair,

and phagocytosis to cope with *S. parasitica* infection.

3.2.2 Regulation of gut microbiota and increase the abundance of

Actinobacteriota

Gut microbiota controls host growth and health status through various pathways[60].

Firstly, in our study, the function of gut microbiota is mainly enriched in metabolic-

related pathways linalool increased the abundance of the beneficial phylum

Actinobacteriota and decreased the abundance of the harmful phylum

Verrucomicrobiota. Actinobacteriota plays a crucial role because Actinobacteria can

produce enzymes, antibiotics, Short-chain fatty acids (SCFAs), signaling molecules,

and immunomodulators[61]. Meanwhile, Actinobacteriota correlated positive with the

complement and coagulation system according to the correlation analysis. Secondly,

intestinal IgA production was correlated significantly with Proteobacteria which are

the primary inducers of IgA production by B cells[62]. In summary, our findings

suggested that linalool may positively modulate the immune response by increasing

the abundance of beneficial phylum Actinobacteriota to *S. parasitica* infection.

3.2.3 Regulation of Inflammatory Factors

Pattern Recognition Receptors (PPRs) detect and initiate multiple signaling

pathways upon recognizing Pathogen-Associated Molecular Patterns (PAMPs) on the

surface of pathogens (bacteria, viruses, fungi, etc.), inducing subsequent host immune

responses[63]. Myeloid differentiation primary-response protein 88 (MyD88) serves as

the crucial signaling adaptor protein for Toll-like and interleukin-1 receptors, which is

responsible for the conduction of multiple downstream signaling pathways[64]. The

expression of *ap-1*, *p-38*, and *pI3k* were up regulate in downstream signaling

pathways, causing higher expression of inflammatory factors and enhanced cellular

oxidation and antioxidant levels[65]. Meanwhile, in our study, the gene expression

levels of *il-1β*, *il-6*, *ccl19*, and *ccl5* which are closely related to the inflammasome,

were up expression. Inflammasomes are protein complexes capable of detecting

diverse inflammatory stimuli and are vital for eliminating pathogens or injured

cells[66]. Therefore, linalool might have influenced the immune system response in

grass carp, regulating the inflammatory processes.

4. Conclusions

In summary, our study proved that linalool is an effective natural antimicrobial agent

against *S. parasitica*. Linalool disrupted the cell membrane, resulting in the leakage of

cellular components and restricting ribosomal function, thereby inhibiting protein

synthesis. Comprehensive analyses revealed the specific role of linalool in protecting

S. parasitica infection skin inflammation in grass carp. Meanwhile, the complement

and coagulation systems, inflammatory factors, along with Actinobacteriota,

significantly involved in this protective effect (Figure 7). Further investigation would

be carried out to explore and refine how linalool affects the inflammatory effects of *S.*

parasitica on fish through signaling pathways and mechanisms. We believe that this

study will contribute to developing an innovative alternative strategy, distinct from

antibiotics, to cope with *S. parasitica* infection in the aquaculture industry.

5. Materials and methods

5.1 Oomycete strains and materials

S. parasitica used in this study was from our lab, which was isolated from grass

carp (*Ctenopharyngodon idella*). Linalool (0.862g/mL), pinene (0.874g/mL),

limonene (0.842g/mL), myrcene (≥95%), terpinene (≥95%), resazurin dye, and

Prodium iodide (PI) were purchased from Sigma-Aldrich (USA). SYTO 9 and FM4-

64 were purchased from ThermoFisher (USA). Potato dextrose agar (PDA) and

glucose yeast extract (GY) were purchased from Guangdong Huankai Microbial Sci.

and Tech. Co., Ltd, China.

5.2 Agar diffusion assay

The anti-oomycetes effects of EOs and 5 components (linalool, terpinene,

limonene, myrcene, and pinene) were assessed by using the agar diffusion method.

Initially, a 6 mm mycelium plug was inoculated at the center of the plates.

Subsequently, 6 mm sterile filter paper discs were soaked in EOs and components for

more than 12 h and placed symmetrically at a distance of 2.5 cm from the target

mycelium plug. Sterile distilled water was used as negative control. Finally, the width

of the colonies was measured once the mycelium in the control group was completely

covered the plate.

Inhibition rate (%) = [(P1 − P2)/P1] × 100%, where, P1 = colony growth diameter

(mm) of control group, P2 = colony growth diameter (mm) of treated group.

5.3 Minimum inhibitory concentration (MIC) and minimum fungicidal concentration (MFC) assays.

To determine the MIC and MFC of linalool on mycelium, 6 mm mycelium plugs

were inoculated into a 24 well cell culture plate. The wells contained linalool at

concentrations of 0.8%, 0.4%, 0.2%, 0.1%, 0.05%, 0.025%, 0.0125%, and 0.00625%,

dissolved in 1% Tween 20 and potato dextrose broth (PDB). Wells without mycelium

plugs were served as negative controls. After incubating at 25°C for 48 h, the lowest

concentration of linalool with no visible mycelium growth was recorded as MIC.

Mycelium plugs showing no growth at this MIC concentration were then inoculated

onto PDA plates and incubated for 7 days at 25°C to determine MFC.

The method for spores induction is similar as previously described by our previous

research[67]. In a 96-well cell culture plate, 100 μL of linalool at concentrations of

0.4%, 0.2%, 0.1%, 0.05%, 0.025%, 0.0125%, 0.00625%, and 0.003125% were

dissolved in 1% Tween 20 and GY were added initially to each well, followed by the

addition of 100 μL of spore suspension at a concentration of $1×10^6$ spores/mL to each

well. Malachite green and Tween 20 served as positive and negative controls,

respectively. Following a 24-hour incubation at 20°C, 5 μL of resazurin dye was

added. When the color of the wells turned to pink, the linalool concentration in the

first non-pink well corresponded to the MIC. The culture liquid from wells with no

visible mycelium growth was streaked onto new plates. After incubating at 25°C for 7

days, the well with the lowest linalool concentration with no visible mycelium growth

was determined as MFC.

5.4 Effect of linalool on the mycelium radial growth rate of *S. parasitica*

544   Different concentrations of linalool were added to molten PDA plates at 50°C for

545   quantitative assessment. A similarly sized mycelium plug was placed at the center of

546   each plate. Once the mycelium in the control group without linalool reached the edge

547   of the plate (approximately 60 h), the radial diameter of the mycelium was measured.

The calculation of growth inhibition was determined by assessing the percentage inhibition of radial growth, as below:

PIRG % = [(R1− R2)/R1] × 100%, where, R1 = radial growth of control plate (mm), R2 = radial growth of treatment plate (mm)

5.5 Confocal laser scanning microscopy (CLSM) assay

2.5.1 Analysis of viability of *S. parasitica*

The mycelium was treated with linalool at MIC (0.05%) and 2×MIC (0.1%) for 2 h, and untreated control group was prepared simultaneously. The mycelium resuspended in PBS was further treated with PI at a final concentration of 2.5 g/mL for 15 minutes. PI is excited at a wavelength of 488 nm and emits light at 620 nm. Fluorescence images were observed using CLSM (LSM880, Zeiss, Germany) at 40× magnification.

2.5.2 Analysis of membrane damage of *S. parasitica*

Mycelium cultured for 18 h was collected and sequentially treated with SYTO 9 at a concentration of 20 μmol/mL, FM4-64 at a concentration of 10 μmol/mL, and linalool at a concentration of 3%. Record observations at 0, 2, 8, and 20 min. Both SYTO 9 and FM4-64 have an excitation wavelength of 488 nm and emit light at 620 nm and 520 nm, respectively. Fluorescence images were observed using CLSM (LSM880, Zeiss, Germany) at 40× magnification.

5.6 Scanning Electron Microscope (SEM) assay and transmission electron microscopy (TEM) assay

Mycelium exposed to MIC (0.05%) linalool for 2 h underwent fixation with 2.5% glutaraldehyde at 4°C for 4 h to examine the impact of linalool on membrane damage. Following a rinse with phosphate-buffered saline (PBS; pH 7.2), the fixed samples experienced dehydration for 15 min using a graded series of ethanol concentrations (10%, 30%, 50%, 70%, 90%, and 100%), followed by an additional 20 min in 100% ethanol. Post-dehydration, ethanol was exchanged twice for 20 min each with tertiary butanol. Subsequently, the samples were subjected to vacuum freeze-drying, gold-spraying, and examination with SEM (Hitachi, SU8010). After fixation, the samples

were ultrathin-sectioned, subjected to negative staining using 1% phosphotungstic

acid for 5 min and observed by TEM (Hitachi, HT-7700).

5.7 RNA-Seq and bioinformatics analysis

Total RNA was isolated from linalool treated (exposed to 0.5×MIC linalool for 10

minutes) and untreated control groups using the TRIzol method. The messenger RNA

(mRNA) was enriched and fragmented into 200 bp fragments. These mRNA

fragments were reverse-transcribed into cDNA using random primers and reverse

transcriptase. The cDNA ends were modified and ligated to an adaptor. After PCR

amplification and purification, Illumina sequencing was performed. Quality control

removed low-quality reads. The mapped reads were assembled using StringTie. Gene

expression was quantified with the transcripts-per-million-reads method to identify

differentially expressed genes (DEGs) using a threshold of |log twofold change| $\geq$ 2

and FDR < 0.05. Afterward, functional enrichment analysis was performed using the

Kyoto Encyclopedia of Genes and Genomes (KEGG) database. KEGG pathway

analysis was conducted using Goatools. The Majorbio cloud platform was used for

data analysis[68].

5.8 Molecular docking

Linalool may interact with specific domains of the ribosome, potentially

influencing its structure and function. Therefore, our investigation aimed to determine

whether linalool could bind to the ribosome-associated upstream proteins of *S.*

parasitica. As no crystal structure of *S. parasitica* ribosome proteins existed in the

Protein Data Bank (PDB), we performed homology modeling. The protein sequences

NOP1 (XP_012204452.1), SNU13 (XP_012198008.1), and DKC1

(XP_012193765.1) were obtained from GenBank, and homology modeling was

conducted using AlphaFold2. The most optimal configuration identified from the

homologous modeling results was chosen as the receptor protein for subsequent

molecular docking. The model was then evaluated using ProSA-web and

PROCHECK server. The secondary structure of linalool was then loaded on

PubChem. Both the ligand (linalool) and target proteins underwent processing with

AutoDockTools 1.5.6, including energy minimization, dehydration, and
hydrogenation. The conformation exhibiting the lowest energy was selected as the
most likely binding state. The ultimate docking outcome was visualized utilizing the
open-source software PyMOLv2 and Discovery Studio Visualizer V2021.

5.9 Animal Experiments

5.9.1 Ethics statement

All experiments involving animals were conducted according to the ethical policies
and procedures approved by the Institutional Animal Care and Use Committee of
Hunan Agricultural University, China (Approval No. 430516).

5.9.2 Linalool therapeutic and prophylactic assays

In this study, 4 groups were established: (1) Positive control (10 fish infected with
S. parasitica). (2) Linalool therapeutic group (10 fish infected with *S. parasitica*,
soaked in 0.00039% linalool in a 20L tank for 7 days). (3) Linalool prophylactic
group (10 uninfected fish soaked in 0.00039% linalool in a 20L tank for 2 days,
followed by the addition of 1×10^6 spores/mL secondary zoospores). (4) Negative
control (10 uninfected fish without linalool treatment). Each group had 3 replicate
tanks. The number of fish without visible mycelium on day 7 and the cumulative
survival rate were recorded. The fish samples from the negative control, positive
control, linalool therapeutic, and linalool prophylactic groups were collected for
histopathological examinations. Tissue samples underwent fixation, sectioning,
hematoxylin and eosin (H&E) staining, and microscopic examination using a
microscope (BX53, Olympus).

5.9.3 16S rRNA and transcriptome sequencing and analysis

Genomic DNA was extracted from grass carp intestinal content samples and stored
at -20°C following the instructions of the 16S rRNA gene pyrosequencing DNA kit.
We conducted paired-end sequencing on the Illumina MiSeq platform, amplifying the
V3-V4 regions with specific barcoded primers using a thermocycler PCR system. To
unveil the intricate microbial diversity within the samples, we employed Quantitative
Insights Into Microbial Ecology (QIIME) which can compute and visualize α-

diversity, β-diversity, and principal coordinate analysis (PCoA). Each sample was

measured 3 times to ensure robust results.

Spleen tissue was obtained from 3 fish in each group on day 7. These collected

spleen tissues were promptly flash-frozen in liquid nitrogen and then stored at -80°C

until they were ready for RNA isolation. RNA-Seq and bioinformatics analysis was

carried out as shown in Section 2.6.

In each group, 8 fish were utilized for immunological assays, and on day 7, blood

samples were collected from the tail veins using heparinized syringes and left to

coagulate overnight at 4°C. After centrifugation at 3500×g for 10 minutes at 4°C,

serum was obtained from each fish and stored at -80°C for subsequent analysis. Kits

from Nanjing Jiancheng Institute (Nanjing, China) were used to measure lysozyme

(LZY) activity, superoxide dismutase (SOD) activity, and alkaline phosphatase (AKP)

activity.

5.10 Detection of immune-related gene expression

Spleen tissues RNA extraction utilized the EZNA Total RNA Kit II (OMEGA

R6934-02). RNA concentrations were quantified using a spectrophotometer

(Eppendorf BioSpectrometer basic, Germany), and quality assessment was conducted

through 1% gel electrophoresis. Subsequent cDNA synthesis employed reverse

transcription with the RevertAid First Strand cDNA Synthesis Kit (Thermo Fisher

Scientific, USA). RNA concentrations were quantified via spectrophotometry

(Eppendorf BioSpectrometer basic, Germany). Subsequently, cDNA was synthesized

via reverse transcription using the RevertAid First Strand cDNA Synthesis Kit

(Thermo Fisher Scientific, USA). Expression levels of complement factor H (*cfh*),

mannan-binding lectin serine protease 2 (*masp2*), complement component 1, r

subcomponent (*c1r*), complement component 3 (*c3*), complement factor B (*cfb*),

transcription factor AP-1 (*ap-1*), interleukin 1 beta (*il-1β*) and interleukin 6 (*il-6*) were

detected. The PCR procedure was described as follows: 95 °C for 30 s, followed by

40 cycles of 95 °C for 5 s and 60 °C for 40 s. Employing the CFX96 Touch Real-time

PCR Detection System (Bio-rad, USA), gene expression levels were assessed using

the $2^{-\Delta\Delta Ct}$ method, with β-actin serving as the reference gene. The primers in qPCR analysis are listed in Table 1.

5.11 Statistical analysis

The data was expressed as mean ± standard error, and all experiments were conducted in triplicate. Pearson correlation analysis was employed to investigate the correlation between potential immune-related genes in grass carp and their characteristics, the method was implemented in R Programming language. Statistical analysis was performed using GraphPad Prism 9 software, and the student t-test was conducted.

Acknowledgments

This research was funded by National Natural Science Foundation of China (32073020, 32201960), Science and Technology Innovation Program of Hunan Province (2022RC1150), Hunan Provincial Natural Science Foundation of China (2023JJ40364), and Agricultural Science and Technology Innovation Fund of Hunan (2023CX49).

Author contributions

Tao Tang was in charge of conceptualization, methodology, writing-original draft, writing-review & editing.

Weiming Zhong was in charge of data curation.

Puyu Tang was in charge of methodology.

Rongsi Dai was in charge of data curation.

Jiajing Guo was in charge of writing - review & editing.

Zhipeng Gao was in charge of funding acquisition, supervision, writing-original draft, writing-review & editing.

Declaration of interests

The authors declare no competing interests.

Data and code availability

The data that support the findings of this study are available in the supplementary material of this article and are available from the corresponding author upon reasonable request.

References

Derevnina, L., Petre, B., Kellner, R., Dagdas, Y. F., Sarowar, M. N., Giannakopoulou, A., De la Concepcion, J. C., Chaparro-Garcia, A., Pennington, H. G., van West, P., et al. (2016) Emerging Oomycete Threats to Plants and Animals. Philosophical Transactions of the Royal Society B-Biological Sciences 371, 9 20150459. https://doi.org/10.1098/rstb.2015.0459.

Tandel, R. S., Chadha, N. K., Dash, P., Sawant, P. B., Pandey, N. N., Chandra, S., Bhat, R. A. H., and Thakuaria, D. (2021) An *in Vitro* Study of Himalayan Plant Extracts against Oomycetes Disease Saprolegniasis in Rainbow Trout (*Oncorhynchus Mykiss*). Journal of Environmental Biology 42, 1008-18. https://doi.org/10.22438/jeb/42/4/MRN-1623.

Wang, H. Z., Li, D. L., Fan, M. H., Zhu, J. K., and Han, X. (2021) Identification and Expression Analysis of Ricin B-Like Lectin Genes in the Fish Pathogen *Saprolegnia Parasitica*. Aquaculture International 29, 1853-68. https://doi.org/10.1007/s10499-021-00722-5.

Costa, S., and Lopes, I. (2022) Saprolegniosis in Amphibians: An Integrated Overview of a Fluffy Killer Disease. Journal of Fungi 8, 30 537. https://doi.org/10.3390/jof8050537.

Sandoval-Sierra, J. V., Latif-Eugenin, F., Martín, M. P., Zaror, L., and Diéguez-Uribeondo, J. (2014) *Saprolegnia* Species Affecting the Salmonid Aquaculture in Chile and Their Associations with Fish Developmental Stage. Aquaculture 434, 462-69. https://doi.org/10.1016/j.aquaculture.2014.09.005.

Khulbe, R. D., Joshi, C., and Bisht, G. S. (1995) Fungal Diseases of Fish in Nanak Sagar, Naini Tal, India. Mycopathologia 130, 71-4. https://doi.org/10.1007/bf01103452.

Engblom, C., Landor, L., Sjöqvist, C., Korkea-aho, T., Viljamaa-Dirks, S., Paulin, L., and Wiklund, T. (2023) Identification and Genetic Characterization of *Saprolegnia Parasitica*, Isolated from Farmed and Wild Fish in Finland. Journal of Fish Diseases 46, 849-60. https://doi.org/10.1111/jfd.13795.

Money, N. P., Davis, C. M., and Ravishankar, J. P. (2004) Biomechanical Evidence for Convergent Evolution of the Invasive Growth Process among Fungi and Oomycete Water Molds. Fungal Genetics and Biology 41, 872-76. https://doi.org/10.1016/j.fgb.2004.06.001.

Van West, Pieter. (2006) *Saprolegnia Parasitica*, an Oomycete Pathogen with a Fishy Appetite:: New Challenges for an Old Problem. Mycologist 20, 99-104. https://doi.org/10.1016/j.mycol.2006.06.004.

van den Berg, A. H., McLaggan, D., Diéguez-Uribeond, J., and van West, P. (2013) The Impact of the Water Moulds *Saprolegnia Diclina* and *Saprolegnia Parasitica* on Natural Ecosystems and the Aquaculture Industry. Fungal Biology Reviews 27, 33-42. https://doi.org/10.1016/j.fbr.2013.05.001.

Yu, J. J., Hu, Y. L., Liu, C. Z., Wu, S. B., Zheng, Z. J., Cui, Z. H., Chen, L., Wei, T., Sun, S. K., Ning, J., et al. (2023) Arscp: An Antimicrobial Residue Surveillance Cloud Platform for Animal-Derived Foods. Science of the Total Environment 858, 7 159807. https://doi.org/10.1016/j.scitotenv.2022.159807.

Gao, Zhipeng, Jiang, Sifan, Zhong, Weiming, Liu, Ting, and Guo, Jiajing. (2023) Linalool Controls the Viability of *Escherichia Coli* by Regulating the Synthesis and Modification of Lipopolysaccharide, the Assembly of Ribosome, and the Expression of Substrate Transporting Proteins. Food Research International 164, 112337.

Peana, A. T., D'Aquila, P. S., Panin, F., Serra, G., Pippia, P., and Moretti, M. D. L. (2002) Anti-Inflammatory Activity of Linalool and Linalyl Acetate Constituents of Essential Oils. Phytomedicine 9, 721-26. https://doi.org/10.1078/094471102321621322.

Zhipeng, Gao, Weiming, Zhong, Ting, Liu, Tianyu, Zhao, and Jiajing, Guo. (2021) Global Proteomic Analysis of *Listeria Monocytogenes'* Response to Linalool. Foods 10, 2449. https://doi.org/10.3390/foods10102449.

Praseetha, S., Sukumaran, S. T., Dan, M. T., Augustus, A. R., Pandian, S. K., and Sugathan, S. (2023) The Anti-Biofilm Potential of Linalool, a Major Compound from *Hedychium Larsenii*, against *Streptococcus Pyogenes* and Its Toxicity Assessment in *Danio Rerio*. Antibiotics-Basel 12, 18 545. https://doi.org/10.3390/antibiotics12030545.

Li, Y. S., He, R. R., Chen, H. M., Chen, D., and Chen, W. X. (2022) Respiratory Depression as Antibacterial Mechanism of Linalool against *Pseudomonas Fragi* Based on Metabolomics. International Journal of Molecular Sciences 23, 13 11586. https://doi.org/10.3390/ijms231911586.

Aytac, Z., Yildiz, Z. I., Kayaci-Senirmak, F., Tekinay, T., and Uyar, T. (2017) Electrospinning of Cyclodextrin/Linalool-Inclusion Complex Nanofibers: Fast-Dissolving Nanofibrous Web with

Prolonged Release and Antibacterial Activity. Food Chemistry 231, 192-201. https://doi.org/10.1016/j.foodchem.2017.03.113.

Rongrong, He, Qiuping, Zhong, Weijun, Chen, Ming, Zhang, Jianfei, Pei, Haiming, Chen, and Wenxue, Chen. (2022) Antimicrobial Mechanism of Linalool against *Brochothrix Thermosphacta* and Its Application on Chilled Beef. Food Research International 157, 111407. https://doi.org/10.1016/j.foodres.2022.111407.

Zhong, W. M., Tang, P. Y., Liu, T., Zhao, T. Y., Guo, J. J., and Gao, Z. P. (2021) Linalool Nanoemulsion Preparation, Characterization and Antimicrobial Activity against *Aeromonas Hydrophila*. International Journal of Molecular Sciences 22, 10 11003. https://doi.org/10.3390/ijms222011003.

Jiang, Y., Pan, X. Y., Li, Y. M., Yang, Y. Z., Jia, Y. N., Lei, B., Feng, J. T., Ma, Z. Q., Liu, X. L., and Yan, H. (2023) Linalool Induces Resistance against Tobacco Mosaic Virus in Tobacco Plants. Plant Disease 107, 2144-52. https://doi.org/10.1094/pdis-09-22-2246-re.

Yang, J. J., Wang, Q. Z., Li, L. W., Li, P. R., Yin, M., Xu, S., Chen, Y., Feng, X., and Wang, B. (2022) Chemical Composition and Antifungal Activity of *Zanthoxylum Armatum* Fruit Essential Oil against *Phytophthora Capsici*. Molecules 27, 17 8636. https://doi.org/10.3390/molecules27238636.

Li, Y. N., Zhang, S. B., Lv, Y. Y., Zhai, H. C., Cai, J. P., and Hu, Y. S. (2022) Linalool, the Main Volatile Constituent from *Zanthoxylum Schinifolium* Pericarp, Prevents Growth of *Aspergillus Flavus* in Post-Harvest Grains. Food Control 137, 13 108967. https://doi.org/10.1016/j.foodcont.2022.108967.

Satoh, M., Kusumoto, N., Matsui, N., Makino, R., Hashida, K., Arai, D., Iiduka, Y., and Ashitani, T. (2022) Antitermitic and Antifungal Properties of Enantiopure Linalool and Furanoid Linalool Oxide Confirmed in *Lindera Umbellata* Var. *Membranacea*. Journal of Wood Chemistry and Technology 42, 37-45. https://doi.org/10.1080/02773813.2021.2004166.

Gomes, P. R. B., Cunha, M. J. O., de Sousa, D. A., Reis, J. B., Melo, A. V., de Freitas, A. C., Hunaldo, V. K. L., Fontenele, M. A., de Paula, M. D., Louzeiro, H. C., et al. (2019) Chemical Study and Antifungal Activity of the Essential Oil of the Branches of *Aniba Duckei* Kostermans. Journal of Essential Oil Bearing Plants 22, 1554-61. https://doi.org/10.1080/0972060x.2019.1700829.

Shen, Q., Li, H. S., Wang, Q. F., Wang, J. Q., Ge, J. R., Yang, X. Y., Wang, X. Y., Li, X. M., Zhang, Y., Zhang, R. M., et al. (2022) Alleviating Effects of Linalool Fumigation on *Botrytis Cinerea* Infections in Postharvest Tomato Fruits. Horticulturae 8, 15 1074. https://doi.org/10.3390/horticulturae8111074.

Marcos-Arias, C., Eraso, E., Madariaga, L., and Quindós, G. (2011) In Vitro Activities of Natural Products against Oral *Candida* Isolates from Denture Wearers. Bmc Complementary and Alternative Medicine 11, 7 119. https://doi.org/10.1186/1472-6882-11-119.

Yan, L. J., and Lamb, R. F. (2012) Amino Acid Sensing and Regulation of Mtorc1. Seminars in Cell & Developmental Biology 23, 621-25. https://doi.org/10.1016/j.semcdb.2012.02.001.

Athanasopoulos, A., André, B., Sophianopoulou, V., and Gournas, C. (2019) Fungal Plasma Membrane Domains. Fems Microbiology Reviews 43, 642-73. https://doi.org/10.1093/femsre/fuz022.

Jiajing, Guo, Zhipeng, Gao, Gaoyang, Li, Fuhua, Fu, Zengenni, Liang, Hui, Zhu, and Yang, Shan. (2019) Antimicrobial and Antibiofilm Efficacy and Mechanism of Essential Oil from Citrus Changshan-Huyou Y. B. Chang against *Listeria Monocytogenes*. Food Control 105, 256-64. https://doi.org/10.1016/j.foodcont.2019.06.014.

Zhipeng, Gao, Weiming, Zhong, Kangyong, Chen, Puyu, Tang, and Jiajing, Guo. (2020) Chemical Composition and Anti-Biofilm Activity of Essential Oil from *Citrus Medica* L. Var. *Sarcodactylis Swingle* against *Listeria Monocytogenes*. Industrial Crops and Products 144, 112036. https://doi.org/10.1016/j.indcrop.2019.112036.

Wenbin, Xiao, Gao, Zhipeng, Liu, Ting, Zhong, Weiming, Jiang, Sifan, He, Mingwang, Fu, Fuhua, Li, Gaoyang, Su, Donglin, and Guo, Jiajing. (2023) Lemon Essential Oil Nanoemulsions: Potential Natural Inhibitors against *Escherichia Coli*. Food Microbiology, 104459.

Cruz, I., Cheetham, J. J., Arnason, J. T., Yack, J. E., and Smith, M. L. (2014) Alkamides from *Echinacea* Disrupt the Fungal Cell Wall-Membrane Complex. Phytomedicine 21, 435-42. https://doi.org/10.1016/j.phymed.2013.10.025.

Meng, F. Y., Liu, X., Li, C., Peng, X. D., Wang, Q., Xu, Q., Sui, J. L., Zhao, G. Q., and Lin, J. (2023) Hinokitiol Inhibits *Aspergillus Fumigatus* by Interfering with the Cell Membrane and Cell Wall. Frontiers in Microbiology 14, 11 1132042. https://doi.org/10.3389/fmicb.2023.1132042.

Ford, R. C., and Beis, K. (2019) Learning the Abcs One at a Time: Structure and Mechanism of Abc Transporters. Biochemical Society Transactions 47, 23-36. https://doi.org/10.1042/bst20180147.

Dean, M., Hamon, Y., and Chimini, G. (2001) The Human Atp-Binding Cassette (Abc) Transporter Superfamily. Journal of Lipid Research 42, 1007-17.

van den Brule, Sybille, Muller, Axel, Fleming, Andrew J., and Smart, Cheryl C. (2002) The Abc Transporter Sptur2 Confers Resistance to the Antifungal Diterpene Sclareol. The Plant journal : for cell and molecular biology 30, 649-62. https://doi.org/10.1046/j.1365-313X.2002.01321.x.

Rees, D. C., Johnson, E., and Lewinson, O. (2009) Abc Transporters: The Power to Change. Nature Reviews Molecular Cell Biology 10, 218-27. https://doi.org/10.1038/nrm2646.

Chaillou, Thomas, Kirby, Tyler J., and McCarthy, John J. (2014) Ribosome Biogenesis: Emerging Evidence for a Central Role in the Regulation of Skeletal Muscle Mass. Journal of Cellular Physiology 229, 1584-94. https://doi.org/https://doi.org/10.1002/jcp.24604.

Qian, W. D., Wang, W. J., Zhang, J. N., Fu, Y. T., Liu, Q. M., Li, X. C., Wang, T., and Zhang, Q. (2022) Exploitation of the Antifungal and Antibiofilm Activities of Plumbagin against *Cryptococcus Neoformans*. Biofouling 38, 558-74. https://doi.org/10.1080/08927014.2022.2094260.

Xing, J. H., Nan, X., Cui, Q., Ma, W. P., and Zhao, H. S. (2018) Nop-7-Associated 2 (Nsa2) Is Required for Ribosome Biogenesis and Protein Synthesis. Biochemical and Biophysical Research Communications 505, 249-54. https://doi.org/10.1016/j.bbrc.2018.09.041.

Wang, H. X., Wang, Z. G., Xu, W. H., and Wang, K. X. (2021) Comprehensive Transcriptomic and Proteomic Analyses Identify Intracellular Targets for Myriocin to Induce *Fusarium Oxysporum* F. Sp. Niveum Cell Death. Microbial Cell Factories 20, 15 69. https://doi.org/10.1186/s12934-021-01560-z.

Lieschke, G. J., Oates, A. C., Crowhurst, M. O., Ward, A. C., and Layton, J. E. (2001) Morphologic and Functional Characterization of Granulocytes and Macrophages in Embryonic and Adult Zebrafish. Blood 98, 3087-96. https://doi.org/10.1182/blood.V98.10.3087.

Li, F. M., Lv, Z. C., Zhong, Z. J., Mao, L. T., Chua, L. S., Xu, L. X., and Huang, R. M. (2023) The Effect of Cyclosporin a on *Aspergillus Niger* and the Possible Mechanisms Involved. Foods 12, 11 567. https://doi.org/10.3390/foods12030567.

Abdelwahab, H., Del Campo, J. S. M., Dai, Y. M., Adly, C., El-Sohaimy, S., and Sobrado, P. (2016) Mechanism of Rifampicin Inactivation in *Nocardia Farcinica*. Plos One 11, 23 e0162578. https://doi.org/10.1371/journal.pone.0162578.

Gasmi, A., Bjorklund, G., Peana, M., Mujawdiya, P. K., Pivina, L., Ongenae, A., Piscopo, S., and Severin, B. (2022) Phosphocalcic Metabolism and the Role of Vitamin D, Vitamin K2, and Nattokinase Supplementation. Critical Reviews in Food Science and Nutrition 62, 7062-71. https://doi.org/10.1080/10408398.2021.1910481.

He, R. R., Zhong, Q. P., Chen, W. J., Zhang, M., Pei, J. F., Chen, H. M., and Chen, W. X. (2023) Transcriptomic and Proteomic Investigation of Metabolic Disruption in *Listeria Monocytogenes* Triggered by Linalool and Its Application in Chicken Breast Preservation. Lwt-Food Science and Technology 176, 12 114492. https://doi.org/10.1016/j.lwt.2023.114492.

Gao, C. B., Cai, X., Lymbery, A. J., Ma, L., Cao, M., and Li, C. (2023) Integrative Transcriptomic Profiling Reveals the Key Pathways in the Regulation Mechanism of Fish Intestine-Spleen Immunity During the Bacterial Challenges. Aquaculture 568, 15 739320. https://doi.org/10.1016/j.aquaculture.2023.739320.

Xu, C., Li, E. C., Suo, Y. T., Su, Y. J., Lu, M. H., Zhao, Q., Qin, J. G., and Chen, L. Q. (2018) Histological and Transcriptomic Responses of Two Immune Organs, the Spleen and Head Kidney, in Nile Tilapia (Oreochromis Niloticus) to Long-Term Hypersaline Stress. Fish & Shellfish Immunology 76, 48-57. https://doi.org/10.1016/j.fsi.2018.02.041.

Oikonomopoulou, K., Ricklin, D., Ward, P. A., and Lambris, J. D. (2012) Interactions between Coagulation and Complement-Their Role in Inflammation. Seminars in Immunopathology 34, 151-65. https://doi.org/10.1007/s00281-011-0280-x.

Pan, L. Y., Peng, C., Wang, L., Li, L. L., Huang, S., Fei, C. Y., Wang, N., Chu, F. R., Peng, D. Y., and Duan, X. C. (2022) Network Pharmacology and Experimental Validation-Based Approach to Understand the Effect and Mechanism of Taohong Siwu Decoction against Ischemic Stroke. Journal of Ethnopharmacology 294, 12 115339. https://doi.org/10.1016/j.jep.2022.115339.

Kobayashi, H., Albarracin, L., Sato, N., Kanmani, P., Kober, Akmh, Ikeda-Ohtsubo, W., Suda, Y., Nochi, T., Aso, H., Makino, S., et al. (2016) Modulation of Porcine Intestinal Epitheliocytes Immunetranscriptome Response by Lactobacillus Jensenii Tl2937. Beneficial Microbes 7, 769-82. https://doi.org/10.3920/bm2016.0095.

Krarup, A., Wallis, R., Presanis, J. S., Gal, P., and Sim, R. B. (2007) Simultaneous Activation of Complement and Coagulation by Mbl-Associated Serine Protease 2. Plos One 2, 8 e623. https://doi.org/10.1371/journal.pone.0000623.

Xie, Y., Zhang, H. Z., and Huang, T. Q. (2022) Quantitative Proteomics Reveal Three Potential Biomarkers for Risk Assessment of Acute Myocardial Infarction. Bioengineered 13, 4939-50. https://doi.org/10.1080/21655979.2022.2037365.

Ferluga, J., Kouser, L., Murugaiah, V., Sim, R. B., and Kishore, U. (2017) Potential Influences of Complement Factor H in Autoimmune Inflammatory and Thrombotic Disorders. Molecular Immunology 84, 84-106. https://doi.org/10.1016/j.molimm.2017.01.015.

Desai, J. V., Kumar, D., Freiwald, T., Chauss, D., Johnson, M. D., Abers, M. S., Steinbrink, J. M., Perfect, J. R., Alexander, B., Matzaraki, V., et al. (2023) C5a-Licensed Phagocytes Drive Sterilizing Immunity During Systemic Fungal Infection. Cell 186, 2802-+. https://doi.org/10.1016/j.cell.2023.04.031.

de Willige, S. U., de Visser, M. C. H., Houwing-Duistermaat, J. J., Rosendaal, F. R., Vos, H. L., and
Bertina, R. M. (2005) Genetic Variation in the Fibrinogen Gamma Gene Increases the Risk for
Deep Venous Thrombosis by Reducing Plasma Fibrinogen Gamma ' Levels. Blood 106, 4176-
83. https://doi.org/10.1182/blood-2005-05-2180.

Wang, Y., Guo, W. J., Xie, S. X., Liu, Y., Xu, D. D., Chen, G., and Xu, Y. J. (2021) Multi-Omics
Analysis of Brain Tissue Metabolome and Proteome Reveals the Protective Effect of Gross
Saponins of *Tribulus Terrestris* L. Fruit against Ischemic Stroke in Rat. Journal of
Ethnopharmacology 278, 11 114280. https://doi.org/10.1016/j.jep.2021.114280.

Bryer, Scott C., Fantuzzi, Giamila, Van Rooijen, Nico, and Koh, Timothy J. (2008) Urokinase-Type
Plasminogen Activator Plays Essential Roles in Macrophage Chemotaxis and Skeletal Muscle
Regeneration1. The Journal of Immunology 180, 1179-88.
https://doi.org/10.4049/jimmunol.180.2.1179.

Das, Riku, Ganapathy, Swetha, Settle, Megan, and Plow, Edward F. (2014) Plasminogen Promotes
Macrophage Phagocytosis in Mice. Blood 124, 679-88. https://doi.org/10.1182/blood-2014-
01-549659.

Zhu, B. P., Zhou, J. C., Wang, Z. Q., Hu, Y. J., Cai, M. L., Yang, L. L., Dai, J. H., and Hu, Y. (2022)
Interactions between Intestinal Morphology, Digestion, Inflammatory Responses, and Gut
Microbiota of Juvenile Channel Catfish Elicited by Dietary Enzymatic Rice Protein. Fish &
Shellfish Immunology 127, 155-65. https://doi.org/10.1016/j.fsi.2022.06.018.

Xin, W. G., Li, X. D., Lin, Y. C., Jiang, Y. H., Xu, M. Y., Zhang, Q. L., Wang, F., and Lin, L. B. (2022)
Whole Genome Analysis of Host-Associated *Lactobacillus Salivarius* and the Effects on
Hepatic Antioxidant Enzymes and Gut Microorganisms of *Sinocyclocheilus Grahami*.
Frontiers in Microbiology 13, 15 1014970. https://doi.org/10.3389/fmicb.2022.1014970.

Mirpuri, J., Raetz, M., Sturge, C. R., Wilhelm, C. L., Benson, A., Savani, R. C., Hooper, L. V., and
Yarovinsky, F. (2014) Proteobacteria-Specific Iga Regulates Maturation of the Intestinal
Microbiota. Gut Microbes 5, 28-39. https://doi.org/10.4161/gmic.26489.

Khan, S., Godfrey, V., and Zaki, M. H. "Cytosolic Nucleic Acid Sensors in Inflammatory and
Autoimmune Disorders." In *Nucleic Acid Sensing and Immunity, Pt A*, edited by C.
VanpouilleBox and L. Galluzzi. International Review of Cell and Molecular Biology, 215-53.
London: Academic Press Ltd-Elsevier Science Ltd, 2019.

Minderman, M., Lantermans, H., van der Zwaan, C., Hoogendijk, A. J., van den Biggelaar, M.,
Kersten, M. J., Spaargaren, M., and Pals, S. T. (2023) The Oncogenic Human B-Cell
Lymphoma Myd88 L265p Mutation Genocopies Activation by Phosphorylation at the
Toll/Interleukin-1 Receptor (Tir) Domain. Blood Cancer Journal 13, 10 125.
https://doi.org/10.1038/s41408-023-00896-6.

Wang, S. T., Zhang, K. L., Song, X. T., Huang, Q. Y., Lin, S., Deng, S. L., Qi, M. Y., Yang, Y. C., Lu,
Q., Zhao, D. W., et al. (2023) Tlr4 Overexpression Aggravates Bacterial Lipopolysaccharide-
Induced Apoptosis Via Excessive Autophagy and Nf-Kb/Mapk Signaling in Transgenic
Mammal Models. Cells 12, 19 1769. https://doi.org/10.3390/cells12131769.

Li, M., Xu, C. J., Li, D., Wu, G. F., Wu, G. Q., Yang, C. Y., Pan, Y. F., Pan, Z. Q., Tan, G. L., and Liu,
Y. Y. (2021) Transcriptome Analysis of the Spleen Provides Insight into the Immunoregulation
of *Cyprinus Carpio* Koi under *Aeromonas Veronii* Infection. Aquaculture 540, 10 736650.
https://doi.org/10.1016/j.aquaculture.2021.736650.

Tang, Tao, Zhong, Weiming, Yang, Linlin, He, Mingwang, Jiang, Sifan, Yin, Dan, Guo, Jiajing, and
943   Gao, Zhipeng. (2023) *In Vitro* and *in Vivo* Anti-Oomycetes Activities and Mechanisms of
944   Linalool against *Saprolegnia Ferax*. Aquaculture, 740031.
945   Guo, J. J., Hu, X., Gao, Z. P., Li, G. Y., Fu, F. H., Shang, X. B., Liang, Z. E. N., and Shan, Y. (2021)
946   Global Transcriptomic Response of *Listeria Monocytogenes* Exposed to Fingered Citron
947   (*Citrus Medica* L. Var. *Sarcodactylis Swingle*) Essential Oil. Food Research International 143,
948   11 110274. https://doi.org/10.1016/j.foodres.2021.110274.

Figure 1. The *in vitro* anti-oomycetes activity of linalool against *S. parasitica* CQT2.
A) Inhibition rate (IR) of EOs and their major components against *S. parasitica*
CQT2. B) Determination of MIC of linalool against *S. parasitica* spores, employing
malachite green (2.5 mg/mL) as a positive control (PC) and Tween 20 as a negative
control (NC). C) Effects of linalool on mycelium growth inhibition over a 6 h period
at varying concentrations. D) Examination of PIRG % in mycelium treated with
linalool after 60 hours. E) Effects of different concentrations of linalool on mycelium
growth on PDA plates. F) Effects of linalool on the viability of *S. parasitica* CQT2

through a PI staining assay was conducted under 3 conditions: Control (no linalool),

1×MIC (0.05%) linalool treatment, and 2×MIC (0.1%) linalool treatment.

Figure 2. Effects of linalool on the cell membrane integrity of *S. parasitica* CQT2. A)

Observation of the effect of linalool on the cell membrane of *S. parasitica* CQT2

using the FM4-64 and SYTO 9 staining assay. B-C) SEM images of *S. parasitica*

CQT2 mycelium without (B) and with (C) linalool treatment. D-G) TEM images of *S.*

parasitica CQT2 mycelium without (D) and with (E-G) linalool treatment. CM: cell

membrane; CW: cell wall; ER: rough endoplasmic reticulum; MI: mitochondria; VA:

vacuoles. H) The GO classification of DEGs. I) DEPs involved in intrinsic component

of membrane.

Figure 3. The *in vitro* anti-oomycete mechanisms revealed by transcriptome analysis.

A-B) The GO enrichment of up and down regulated DEGs. C) The KEGG enrichment

of DEGs. Significant enrichment was labeled as "*". P < 0.05, P < 0.01, and P < 0.001

were labeled as "*", "**", and "***", respectively. D) Comparison of ribosome

biogenesis in eukaryotes pathway between the linalool treated mycelium and the

control group. E) Comparison of RNA polymerase pathway between the linalool

treated mycelium and the control group. F) The tertiary structure of NOP1. G)

Molecular docking of linalool with NOP1 involved enlarging the regions binding to

the NOP1 activation pocket to showcase the detailed amino acid structures. H)

Comparison of ABC transporters pathway between the linalool treated mycelium and

the control group. The red squares represented up-regulated genes, and the blue

squares represented down-regulated genes. I) Visual analysis of metabolic pathways

with iPath3.0. The figure represented gene set annotated pathways, red and green

represented pathways annotated by genes in different gene sets, respectively, and blue

represented pathways co-annotated by genes in two gene sets.

Figure 4. Protective effect of linalool on grass carp infected with *S. parasitica*. A) The experimental design. Grass carp were raised for 2 weeks without feeding, fish without infection and linalool (Group NC), fish infected with *S. parasitica* (Group PC), and 10 fish uninfected soaked water containing linalool for 2 days and then 1×10^6 spores /mL secondary zoospores were added (Group LP), and fish infected with *S. parasitica* soaked for 7 days in water containing linalool (Group LT). B) The symptoms of *S. parasitica* infection in grass carps of different groups. C) The survival rates of grass carp infected with *S. parasitica* of different groups. D-F) Alkaline phosphatase (AKP), lysozyme (LYZ), and superoxide dismutase (SOD) activities in serum of grass carp of different groups. G) Histopathological analysis grass carp tissues in different groups. The arrows of different colors indicated: inflammatory cell infiltration in the kidney (blue arrow), cytoplasmic pyknosis (red arrow), nuclei displaced toward one side (yellow arrow), the red and white pulp was poorly demarcated, and a larger volume of melano-macrophage centers (black area), critical damage to the epithelium (green arrow) and myofiber (white arrow).

Figure 5. Global transcriptomic analysis after *S. parasitica* infection and linalool

treatment in grass carp and in-depth analysis of crucial KEGG pathway and DEGs. A-

B) The GO enrichment of up and down regulated DEGs in LT and LP groups. C-D)

The KEGG enrichment of up and down regulated DEGs in LT and LP groups. E)

Complement and coagulation cascades pathway between *S. parasitica* infection and

linalool treatment in the spleens of grass carp. F) 31 DEGs related to Toll-like

receptor signaling pathway. G) 29 DEGs related to chemokine signaling pathway. The

red squares represented up-regulated genes, and the blue squares represented down-

regulated genes. H) Expression levels of *cfh, masp2*, *c1r*, *c3*, *cfb*, *ap-1*, *il-1β*, and *il-6*

in the spleen revealed variations among the PC, LP, and LT groups. For RT-qPCR, the

results were presented as the means ± SD and were analyzed using independent t-tests

(**$p < 0.05$, ** $p < 0.001$, ****$p < 0.0001$).

Figure 6. The effect of linalool on regulating gut microbiota of grass carp infected

with *S. parasitica* and correlation analysis. A-D) The α diversity index comparison

among the different groups. E) The OTUs petal map of PC, LP, and LT groups. F) PCoA using Bray–Curtis distance revealed variations among the PC, LP, and LT groups (ANOSIM R = 0.3086, P = 0.061). G) Relative abundance of the top 10 species in the gut from the different groups (phylum and genus levels). H) Column chart of LDA value distribution. Discriminative biomarkers identified by linear discriminant analysis effect size (LEfSe) with logarithmic LDA score greater than 3.0. I) Heat map of the differences in predicted functional metabolisms within gut bacterial KEGG pathways. J) The correlation layout of potential related immune genes, specific microbial species, and physical characteristics. Pairwise comparisons of characterizations are presented, with a color gradient and block size denoting Pearson's correlation coefficients.

Figure 7. Model diagram of the mode of action of linalool on *S. parasitica* and grass carp. *In vitro*, (1) linalool influenced DNA transcription, tRNA transport, rRNA processing and maturation (5.8S, 18S, and 25S), and the biogenesis and assembly of ribosome subunits (40S and 60S) in the cell, which might lead to the reduction of S. parasitica growth; (2) Linalool disrupted the cell membrane, and the upregulation of

glycerophospholipid metabolism likely represents the cell's response to cope with this

damage; (3) ABC transporters contributed to metabolic resistance by pumping linalool

out of the cell. *In vivo*, (1) Linalool enhanced the complement and coagulation system

which in turn activated host immune defense and lysate *S. parasitica* cells; (2)

Linalool promoted wound healing, tissue repair, and phagocytosis to cope with *S.*

parasitica infection; (3) Linalool positively modulated the immune response by

increasing the abundance of beneficial Actinobacteriota; (4) Linalool stimulated the

production of inflammatory cytokines (*il-1β* and *il-6*) and chemokines (*ccl19* and

ccl5) to lyse *S. parasitica* cells.

Table 1. Primers for qPCR were used in this study.

Primer name	Primer sequence 5′-3′
c3-F	ACAAGCCCATCTACACGCCT
c3-R	GAAACCTCAGGAATGGCGTA
cfb-F	AGAATGGAGAGGTGACCCCA
cfb-R	CCGCTATCCTGACACACTCG
c1r-F	GGGTTTCGGAGGGTTTGA
c1r-R	CCTGTATTGCCACATCTTTCA
cfh-F	ATAGGAGGGAAATGTGGGC
cfh-R	TCCCGTCCACGCCATC
masp2-F	TGGGCACCAGGTTCATCG
masp2-R	GCTGTTTGACTACGCCCATCTT
il-1b-F	ATCTCCTCGTCTGCTGGGTG
il-1b-R	TTGAGAGTTTCAGTGACCTCCTTC
il-6-F	AAAATCTGACCTCAACCCTGG
il-6-R	TGTCCACCCTTCCTCTTGCT
ap-1-F	TCTCACTCAACAGATGCCAAAC
ap-1-R	CTTAGCGGCTGCGAGTTTAT
β-ActinF	GCTATGTGGCTCTTGACTTCG
β-ActinR	GGGCACCTGAACCTCTCATT

1052