Spatial fingerprint of consumer body size and habitat preference on resource

distribution

Chelsea J. Little[1,2*], Pierre Etienne Banville[1,3], Adam T. Ford[4], Rachel M. Germain[2]

[1] School of Environmental Science, Simon Fraser University, Burnaby, B.C., Canada V5A 1S6

[2] Department of Zoology, Biodiversity Research Center, University of British Columbia,

Vancouver, B.C., Canada V6T 1Z4

[3] Department of Geography, Simon Fraser University, Burnaby, B.C., Canada V5A 1S6

[4] Department of Biology, University of British Columbia, Kelowna, B.C., Canada V1V 1V7

* Corresponding author: Chelsea_little@sfu.ca

Abstract

Consumers shape spatial patterns on landscapes by amplifying or dampening environmental

heterogeneity through feeding, excretion, and movement of resources. The degree to which the

environment is modified by consumers depends on species' traits, including body mass,

movement behavior, sociality, and habitat specialization. Global change is altering the size and

traits of consumer populations, but our understanding of how this may impact resource

heterogeneity is limited. Here, we developed an individual-based model of habitat specialists'

and generalists' movement and activity in a patchy landscape and investigated the impact of

changes in population and mean body sizes on landscape-scale resource heterogeneity. We found

that consumers specializing on low-resource habitats (a common risk avoidance strategy)

increased spatial resource heterogeneity regardless of their population and body size. By

contrast, generalists eroded differences among habitats, and we further found that resource

heterogeneity decreased with the average body size of generalist consumers, even while

controlling for total consumer biomass. These nuanced spatial outcomes of consumer-resource

interactions emerge from the unique metabolic demands of specialists vs. generalists, which

scale nonlinearly with body size. Since global change disproportionately impacts larger species

and specialists, indirect consequences on ecosystems may arise via biotic processes, affecting

spatial heterogeneity of future landscapes.

Introduction

Resource landscapes are often thought of as an environmental template that organisms respond
to, but this neglects the role that organisms play in shaping spatial distributions of resources.
Animal consumers serve as important resource vectors in landscapes [1–4]. The deposition of
resources via feces, urine, seeds, placenta, and carcasses contribute to resource dispersal in the
landscape [5–10]. Organismal activity will therefore impact spatial resource heterogeneity,
which in turn facilitates the coexistence of species [11] and thus ecosystem functioning [12]. The
distribution of resources in space is therefore an important landscape characteristic that shapes
patterns of biodiversity and ecosystem functioning [13–16]. With global change altering abiotic
conditions and consumer populations, it is imperative to gain a better understanding of the
processes shaping spatial resource heterogeneity.

The traits of species making up consumer communities may provide a key to
understanding organismal effects on resource landscapes. A handful of studies have established
that species' traits, such as behavior and body size, determine the amount, stoichiometry, and
scale of consumer-driven resource cycling [10,17,18]. For example, social aggregation creates
spatial clustering of consumer-driven resource subsidies [19]. Other traits such as diet
preferences and risk avoidance strategies may concentrate consumer activity into specific habitat
types, leading to divergent rates of resource cycling across the landscape [20]. Meanwhile,
consumer body size influences resource dispersal through at least two pathways. First, energetic
requirements scale nonlinearly with body size [21], and consequently so should resource
consumption and excretion. Second, body size is associated with differences in movement ability
and perceptual range [2,22], and thus the spatial dispersion of foraging and resource deposition.

Predicted shifts in consumer populations and communities due to global change may

rewire resource distributions, yet understanding how they do so is made more challenging

because species' responses to global change vary, in part due to their traits. Larger animals are

more impacted by climate change and land use change [23], and specialist species more than

generalists [24]. Moreover, as population sizes shift with global change [25], we can anticipate

shifts in patterns of resource consumption and deposition across landscapes; however, our

understanding of the net effect of these shifts on spatial resource distribution is limited. Because

body size and population size often trade off [26], changes in body size might offset the effects

of changes in population size and lead to unique spatial patterns of resource distributions that

could not be predicted from body size alone. In one of the few examples of research addressing

this topic, Ferraro *et al.* [27] used an individual-based model of foraging caribou and showed that

higher densities of caribou increased nutrient spatial heterogeneity through carcass depositions

(resource hotspots) and over-grazed dead zones (resource coldspots). At even larger spatial

scales, higher densities of bison increase the amount and change the timing of nutrients available

in different regions of grassland landscapes [28].

In this study, we present an individual-based model of consumer movement and activity

where we factorially manipulated i) population sizes, ii) mean body size, iii) risk-based habitat

preferences, and iv) sociality, and investigated effects on spatial resource heterogeneity in a

patchy landscape. Our model was parameterized to capture key biological features of two

consumer species—a generalist, and a specialist that avoids high-resource patches (a common

predator avoidance strategy)—with scaled movement distances, metabolic requirements, and

perceptual ranges associated with body size.

We hypothesized that larger populations (compared to smaller populations) of generalists

would preferentially graze high-quality areas until they resemble lower-quality areas, reducing

spatial heterogeneity. By contrast, we predicted that specialists would remove resources from

lower-quality areas and accentuate the differences between nutrient-rich and nutrient-poor parts

of the landscape, and thus larger populations (compared to smaller populations) of specialists

would increase spatial heterogeneity. We also expected that, while keeping the total consumer

biomass constant, body size would have contrasting effects on resource redistribution by

generalists and specialists: decreasing the mean body size of generalists would reduce resource

heterogeneity, while reducing the mean body size of specialists would amplify resource

heterogeneity. This is because the metabolic load of smaller individuals is higher, and thus more

resources should be consumed by a population of many small individuals than an equivalent

biomass of fewer large consumers. Therefore, we predicted that the largest differences in the

impacts of consumers on resource heterogeneity would occur between small-bodied specialists

(which would amplify heterogeneity) and small-bodied generalists (which would dampen

heterogeneity). Finally, we predicted that spatial aggregation of consumers would increase

clustering of consumers' impacts on the landscape, thus increasing spatial resource

heterogeneity.

Methods

We used NetLogo v6.0.4 [29] to model generalists and specialists moving in a landscape, where

consumers withdraw resources by foraging and deposit resources through feces and carcasses. Key

parameters are indicated in Table 1, and a detailed explanation is found in Appendix I. This

landscape is divided into a 141 x 141 grid of equal-sized pixels, each pixel representing 10 m^2.

The landscape has two discrete habitat types (**Fig. 1**), approximating a Kenyan savanna (see [30]):

open, high-nutrient glades as clusters of pixels (each cluster forming a "Patch"), in a "Matrix" of

bushy, lower-quality grassland [31]. Each pixel is classified as Patch or Matrix and is assigned a

forage amount drawn from a normal distribution centered on 50 g/m^2 for Patches and 33 g/m^2 for

Matrix, with forage containing 6% nitrogen (N) in Patches and 4% in Matrix. Landscapes

contained an average of four Patches, making up ~5% of the landscape area (**Fig. 1**).

Individuals in the model are either generalists or specialists. Specialist consumers,

resembling Guenther's dik-diks, stay in the Matrix to avoid predators [32,33], while generalists,

akin to impalas, can use both habitat types, but favor Patches [30,33]. Animals forage based on

their metabolic needs and move within their perceptual range, both of which vary as a function of

their body size [34,35]. Foraging behavior adheres to the marginal value theorem, with animals

selecting pixels with better-than-average forage available [36]. Patch pixels are not considered by

specialists, while generalists can choose any pixel (Patch or Matrix) and preferentially graze

Patches at the start of each simulation because of the higher forage and N concentration. Model

timesteps represent 3 hours of movement/activity (see Appendix I for details on movement,

foraging, excretion, and death), and simulations were run for three months, or the length of one

growing season.

Figure 1. N distributions within Patches and Matrix at the beginning (top left) and end (right) of the growing season in simulated landscapes, which are based on a Kenyan savanna (bottom left) with habitat specialists that forage only in Matrix, and generalists that forage across the whole landscape but prefer Patches. In our model, consumers realistically increase perceptual range, movement ability, and metabolic needs with body size. N ranges from low (yellow) to intermediate (orange) and high (green). On right, N distributions after grazing, movement, and nutrient deposition for three example scenarios out of the study's total 64 (see Table S1): here, by equal numbers of large (40kg) specialists, large generalists, or small (10kg) generalists in example simulations. Percent changes in Moran's I, a measure of spatial autocorrelation [37], and in the nitrogen content of each pixel in Patches and Matrix are indicated for each simulation. In the upper right panel, note the waves of N distribution in the Matrix created by foraging specialists compared to the initial landscape. Small dark green spots at the end of the growing season represent nutrient deposition by carcasses of consumers that died during the simulation, which is most evident for larger consumers. Real landscape image © 2024 Maxar Technologies.

We designed scenarios to look at the impact of changes in population size (500 - 6000 kg total biomass) and mean body size (10, 25, and 40 kg) on nitrogen (N) heterogeneity when

individual consumers choose foraging locations independently, as well as for consumers that

forage in social groups (see Appendices II and III; Table S1). While consumers were inspired by

dik-diks (small and solitary) and impala (larger and herd-living), we varied all traits factorially,

and thus implemented sociality independently of specialists/generalist status or body size. For each

scenario, we performed 25 replicate simulations using the R package 'nlrx' [38], which interfaces

with Netlogo.

For each simulation, we then analyzed non-spatial changes in N before and after the

growing season by calculating the four statistical moments (i.e. mean, variance, skewness, and

kurtosis across all pixels). We also analyzed variables providing insight into animal behavior, such

as the net change in N (N deposited minus N eaten) in Patches and in Matrix. To analyze spatial

changes in N distribution, we calculated Global Moran's I ("Moran's I"), a statistical test

measuring spatial autocorrelation (Moran 1950), using the package 'spdep' [39]. Here, Moran's I

quantifies the clustering of nutrients, with higher values indicating stronger spatial autocorrelation.

For each scenario, we present the mean and standard deviation of 25 replicates for all statistics.

We performed all statistical analysis using R version 4.2.2. Full analysis details can be found in

Appendix II.

Table 1: Key parameters influencing the initial distribution of N on the landscape and the withdrawals and

deposits of N by animals.

Parameter (units)	Value	Reference
Mean forage per Patch pixel (g)	5000	[31]
Mean forage per Matrix pixel (g)	3300	[31]
Mean N per Patch pixel (g)	300 (6% of forage)	[31]

Mean N per matrix patch (g)	132 (4% of forage)	[31]
N in consumers' body (g)	2% * bodymass	[40]
Daily metabolic needs of consumers (g)	(7.94 * (bodymass) ^ 0.646 / 11.5)	[35]
Yearly % chance of death for animals	10%	[41,42]
N deposited on landscape per defecation event (g)	50% of N absorbed since last defecation event	[43]
N deposited on landscape per death event (g)	All body N + 100% N absorbed since last defecation event	

Results

Spatial distribution of N varied across simulated landscapes at the end of the growing season. As

predicted, solitary habitat specialists foraging only in the Matrix accentuated differences between

habitat types (note increase in Moran's I in **Fig. 2a,** top panel); for these specialists, larger

population sizes and larger body sizes were associated with greater increases in spatial

heterogeneity. Also as predicted, solitary generalists, who preferentially forage in Patches,

homogenized the landscape, reducing differences between Patches and Matrix (note decrease in

Moran's I in **Fig. 2a,** bottom panel). Although larger body sizes and larger population sizes lead

to greater N depletion in the total landscape for both specialists and generalists (**Fig. S1**),

changes in the spatial heterogeneity of N resulted from the interplay between N depletion in

Patches and Matrix. Here, we have presented results only from scenarios where consumers are

solitary. Results from herd simulations were qualitatively similar, but produced slightly higher

levels of spatial resource heterogeneity (**Fig. S2**), in line with our predictions.

Figure 2: Changes in the spatial distribution of nitrogen (N) from beginning to end of simulations with
population mean body mass of 10 kg, 25 kg, and 40 kg. In (a), bars show mean change in Moran's I and
standard deviation of 25 simulations for each scenario. Note different y-axis scales for generalists and
specialists. Panels (b) through (d) include only generalists, and colors differentiate simulations with 50,
100, and 150 individuals; each point is one replicate simulation. (b) Relationship between total forage
consumed in Patches and total forage consumed in Matrix. (c) and (d) Relationship between mean Net N
in Patches (c) and Matrix (d), respectively, with the change in Moran's I.

For small generalists, larger population sizes led to greater homogenization of N

distribution over the course of the simulations than that caused by smaller population sizes. Since

small generalists, even at larger population sizes, did not remove enough forage from the Patches

to make them of comparable quality as the Matrix, N removal from the Matrix was negligible

(i.e., the first panel of **Fig. 2d**) and larger population sizes were nearly linearly associated with

reductions in Moran's I (i.e., the first panel in **Fig. 2c**). However, for large generalists, the

reduction in Moran's I was consistent across population sizes. Larger generalists depleted

Patches faster than smaller generalists, subsequently removing N from the Matrix (**Fig. 2b**).

However, the increased N depletion of Patches by the largest population sizes of large generalists

(x-axis in rightmost panel of **Fig. 2c**) was of a similar magnitude to their increased removal of N

from the Matrix (x-axis in rightmost panel of **Fig. 2d**), and as a result the difference between

Patch and Matrix quality (and thus Moran's I) was fairly stable across population sizes of large

generalists (**Fig. 2a**, top panel).

Figure 3: Changes in the spatial distribution of nitrogen (N) from beginning to end of simulations with total
population biomass of 1000kg, 2000kg, and 4000kg of. Bars and y-axis in (a) are as for Fig. 2. (b-d) show
only generalists. (b) is as for Fig. 2. (c) and (d) Relationship between mean time individuals spent in
Patches (c) and Matrix (d) with the total forage consumed in the respective habitat type.

Comparing scenarios with equal total population biomass but with many small versus

fewer large consumers revealed the effects of nonlinear scaling of metabolic requirements on

spatial distribution of N (**Fig. 3**). Total forage consumption was higher with many small

consumers compared to few large ones (**Fig. S3**). At lower total biomass, generalists of all body

sizes spent most of their time in Patches (**Fig. 3c**), but small generalists consumed more forage

there (**Fig. 3c**), leading to a stronger homogenizing effect than their larger counterparts (**Fig. 3a**)

as hypothesized. However, at larger total biomass, smaller generalists depleted and left Patches

and spent upwards of 60% of their time in the Matrix (**Fig. 3d**), consuming markedly more

forage in the Matrix than populations of large generalists did despite similar forage consumption

in Patches (**Fig. 3b**), and thus changes in Moran's I were less differentiated.

Discussion

The role of animals in structuring resource patterns is poorly understood. Here, using a spatially-

explicit, individual-based model, we show that organismal traits can make the difference in

whether consumers amplify or erode spatial resource heterogeneity. We found that a key trait

determining whether consumers amplify or erode resource patchiness is species' activity in the

different habitats, modeled here to represent different risk avoidance strategies. Patchy

landscapes have driven the development of ecological theory [44–46], yet to date, models

assessing the effects of animal movement, foraging, and resource deposition have primarily

focused on landscapes without spatial structure in resource availability [19,27,47,48]. When

spatial structure is included, over seasonal time scales, we showed that specialists on lower-

quality (Matrix) habitat amplified existing resource heterogeneity between the resource hotspots

(which they avoided) and the low-quality habitat. Conversely, generalists who could take

advantage of high-resource Patches eroded the landscape's spatial structure. This is a particularly

important insight in an era of global change because generalists often thrive where specialist

species are more vulnerable to habitat loss and fragmentation [24,49], and because of the

proliferation of non-native species and the association between generalism and invasion success

[50,51].

The specialists in our simulations provided an example of how risk avoidance strategies

can reinforce spatial structure and patchiness of landscapes by strengthening existing resource

gradients (for an empirical example, see [52]). In addition to accentuating the difference between

Patch and Matrix habitat, these same specialists also created finer-scale spatial resource

heterogeneity where it was previously lacking: our landscapes featured no initial spatial

clustering of resources *within* a habitat type, but over the course of our simulated growing

season, spatial heterogeneity in the Matrix increased. This was because the removal of nutrients

by foraging followed individuals' movement patterns, which were spatially autocorrelated at the

scale of the individual's perceptual range. This increase in fine-scale heterogeneity was not

captured by Moran's I, which better reflected the larger differences in nutrient concentrations

between Matrix and Patches, but is visible as waves of resource concentrations in landscape

depictions (**Fig. 1**, "150 Large Specialists"). Thus, the effect of organismal activity on nutrient

spatial heterogeneity depends on both the existing structure of the landscape and the scale of

observation.

Globally, many species of large-bodied vertebrates are experiencing range contractions,

whereas smaller-bodied vertebrates are more likely to be experiencing range expansions [53],

and the largest consumer species are also at higher risk of extinction than are their smaller

counterparts [54]. Our results imply that in patchy landscapes, the impact of species turnover

from large to small generalists depends on both initial community biomass and whether shifts in

body size are accompanied by altered total biomass (i.e., the summed biomass of all individuals

in the population). For the same total biomass, smaller consumers consume more forage due to

higher mass-specific metabolic needs [21]. In terms of spatial ramifications, at small body sizes

and low total biomass, larger populations of generalists homogenized the landscape more than

smaller populations by depleting resource hotspots. However, at larger body sizes and greater

total biomass, foraging behavior whereby generalists left depleted Patches to forage in the Matrix

meant that the further depletion of resource hotspots by a larger population was offset by the

increased consumption of resources in the lower quality habitat—that is, spatial homogenization

plateaued. Thus, we suggest that where generalists are already dominant, a shift from large to

small body size may have minimal effects on spatial structure if total biomass remains the same,

however, such a shift would increase overall rates of consumption and recycling of resources.

This is because smaller individuals have higher metabolic demands per unit mass compared to

larger individuals (a universal feature of metabolic scaling).

Our model did not include population dynamics of either consumers (beyond mortality)

or forage resources, and only considered the spatial dynamics of net resource availability. This

increased computational feasibility and enabled us to isolate the impact of vertebrates on nutrient

distributions. Despite simplifications, our finding that generalists (which we based on impalas)

homogenize N distributions by reducing N in Patches aligns with empirical results in an

intriguing way: impalas increase the relative density of a well-defended plant species [30] which

has less foliar N than a poorly-defended congener [55] that is abundant where impala are rare,

thus altering spatial heterogeneity of N in plants.

Another caveat is that in our model, not all N consumed in forage is returned to the

landscape. Some N is incorporated into consumer bodies, and consumer death was not high

enough in our model to offset this. This dynamic is not unrealistic over the course of a single wet

season, based on mortality estimates from the literature [41,42]. We also did not explicitly model

processes such as predator and scavenger export of nutrients, leaching, or nutrient volatilization,

instead incorporating assumptions about how much N was returned to the landscape by

excretion, egestion, and carcass deposition (see Appendix I, and **Fig. S4** for a sensitivity test).

Here, we showed through simulations that animals have the ability to shape resource

heterogeneity on the landscape over short periods of time such as a single season, thus shifting

the environmental template for subsequent consumer-resource interactions. Landscape

heterogeneity is important because it creates different niches that can serve to sustain

biodiversity [57]. Our results suggest that species' traits and abundance are key mediators of

their effect on resource distributions, which could lead to varying impacts of future biodiversity

change. We propose two future directions for spatially-explicit, agent-based modeling to explore

the impacts of these projected changes.The first is to explicitly model different chemical forms

of nutrients, which are more or less bioavailable to organisms [56]. This approach would account

for the fact that nutrients deposited through dung and urine are not immediately available to

consumers, but must first be transformed into plant shoots. We expect that coupling these

considerations with longer simulations that account for consumer population dynamics would

lead to new spatial patterns of foraging movement by consumers, and thus new emergent

dynamics of nutrient spatial distributions. Second, further work should also explore how

competing consumers and trophic interactions modify the patterns found in this study. For

example, we modeled species based on known risk-avoidance strategies rather than including

predators in the models, but doing so may lead to behavioral shifts and emergent dynamics,

additionally because trophic role influences resource deposition [58]. Incorporating additional

ecological complexity into spatially-explicit models could enable a better understanding of how

organisms will shape their own fates via resource dynamics in future landscapes.

Acknowledgments

289   This research was enabled in part by support from the BC DRI Group, Simon Fraser University's

290   Supercomputer Cedar (www.sfu.ca/research/supercomputer-cedar), and the Digital Research

291   Alliance of Canada (www.alliancecan.ca). We thank Matteo Rizzuto and members of the Little

292   Ecology Group for valuable feedback on the manuscript.

Funding

295   This work was supported by a Killam Postdoctoral Research Fellowship from the Killam Trusts

296   and a Discovery Grant from the Natural Science and Engineering Research Council of Canada.

References

1. Earl JE, Zollner PA. 2017 Advancing research on animal-transported subsidies by

integrating animal movement and ecosystem modelling. *Journal of Animal Ecology* **86**, 987–

997. (doi:10.1111/1365-2656.12711)

2. Little CJ *et al.* 2022 Movement with meaning: integrating information into meta-ecology.

Oikos **2022**. (doi:10.1111/oik.08892)

3.   McInturf AG, Pollack L, Yang LH, Spiegel O. 2019 Vectors with autonomy: what

distinguishes animal-mediated nutrient transport from abiotic vectors? *Biological Reviews*

94, 1761–1773. (doi:10.1111/brv.12525)

4.   Pringle RM *et al.* 2023 Impacts of large herbivores on terrestrial ecosystems. *Current*

Biology **33**, R584–R610. (doi:10.1016/j.cub.2023.04.024)

5.   Bump JK, Webster CR, Vucetich JA, Peterson RO, Shields JM, Powers MD. 2009 Ungulate

Carcasses Perforate Ecological Filters and Create Biogeochemical Hotspots in Forest

Herbaceous Layers Allowing Trees a Competitive Advantage. *Ecosystems* **12**, 996–1007.

(doi:10.1007/s10021-009-9274-0)

6.   Ferraro KM, Welker L, Ward EB, Schmitz OJ, Bradford MA. 2023 Plant mycorrhizal

associations mediate the zoogeochemical effects of calving subsidies by a forest ungulate.

Journal of Animal Ecology **92**, 2280–2296. (doi:10.1111/1365-2656.14002)

7.   McIntyre PB, Flecker AS, Vanni MJ, Hood JM, Taylor BW, Thomas SA. 2008 Fish

distributions and nutrient cycling in streams: Can fish create biogeochemical hotspots?

Ecology **89**, 2335–2346. (doi:10.1890/07-1552.1)

8.   le Roux E, van Veenhuisen LS, Kerley GIH, Cromsigt JPGM. 2020 Animal body size

distribution influences the ratios of nutrients supplied to plants. *Proceedings of the National*

Academy of Sciences of the United States of America **117**, 22256–22263.

(doi:10.1073/pnas.2003269117)

9.   Schmitz OJ, Wilmers CC, Leroux SJ, Doughty CE, Atwood TB, Galetti M, Davies AB,

Goetz SJ. 2018 Animals and the zoogeochemistry of the carbon cycle. *Science* **362**.

(doi:10.1126/science.aar3213)

10. Sitters J, Bakker ES, Veldhuis MP, Veen GF, Venterink HO, Vanni MJ. 2017 The

327 stoichiometry of nutrient release by terrestrial herbivores and its ecosystem consequences.

328 *Frontiers in Earth Science* **5**, 1–8. (doi:10.3389/feart.2017.00032)

329 11. Amarasekare P. 2003 Competitive coexistence in spatially structured environments: a

330 synthesis. *Ecology Letters* **6**, 1109–1122. (doi:10.1046/j.1461-0248.2003.00530.x)

331 12. Loreau M, Mouquet N, Gonzalez A. 2003 Biodiversity as spatial insurance in heterogeneous

332 landscapes. *Proc. Natl. Acad. Sci. U.S.A.* **100**, 12765–12770.

333 (doi:10.1073/pnas.2235465100)

334 13. Gonzalez A *et al.* 2020 Scaling-up biodiversity-ecosystem functioning research. *Ecology*

335 *Letters* **23**, 757–776. (doi:10.1111/ele.13456)

336 14. Rizzuto M, Leroux SJ, Schmitz OJ, Vander Wal E, Wiersma YF, Heckford TR. 2024

337 Animal-vectored nutrient flows across resource gradients influence the nature of local and

338 meta-ecosystem functioning. *Ecological Modelling* **488**, 110570.

339 (doi:10.1016/j.ecolmodel.2023.110570)

340 15. Seiferling I, Proulx R, Wirth C. 2014 Disentangling the environmental-heterogeneity–

341 species-diversity relationship along a gradient of human footprint. *Ecology* **95**, 2084–2095.

342 (doi:10.1890/13-1344.1)

343 16. Tylianakis JM, Rand TA, Kahmen A, Klein A-M, Buchmann N, Perner J, Tscharntke T.

344 2008 Resource Heterogeneity Moderates the Biodiversity-Function Relationship in Real

345 World Ecosystems. *PLoS Biol* **6**, e122. (doi:10.1371/journal.pbio.0060122)

346 17. Veldhuis MP, Gommers MI, Olff H, Berg MP. 2018 Spatial redistribution of nutrients by

347 large herbivores and dung beetles in a savanna ecosystem. *Journal of Ecology* **106**, 422–433.

348 (doi:10.1111/1365-2745.12874)

349 18. Subalusky AL, Post DM. 2019 Context dependency of animal resource subsidies. *Biological*

Reviews **94**, 517–538. (doi:10.1111/brv.12465)

19. Bampoh DK, Earl JE, Zollner PA. 2021 Simulating the relative effects of movement and

sociality on the distribution of animal-transported subsidies. *Theor Ecol* **14**, 57–70.

(doi:10.1007/s12080-020-00480-9)

20. Peller T, Marleau JN, Guichard F. 2022 Traits affecting nutrient recycling by mobile

consumers can explain coexistence and spatially heterogeneous trophic regulation across a

meta-ecosystem. *Ecology Letters* **25**, 440–452. (doi:10.1111/ele.13941)

21. Brown JH, Gillooly JF, Allen AP, Savage VM, West GB. 2004 Toward a metabolic theory

of ecology. *Ecology* **85**, 1771–1789.

22. Straus S *et al.* 2024 Macroecological constraints on species' 'movement profiles': Body

mass does not explain it all. *Global Ecol Biogeogr* **33**, 227–243. (doi:10.1111/geb.13786)

23. Albaladejo-Robles G, Böhm M, Newbold T. 2023 Species life-history strategies affect

population responses to temperature and land-cover changes. *Global Change Biology* **29**,

97–109. (doi:10.1111/gcb.16454)

24. Clavel J, Julliard R, Devictor V. 2011 Worldwide decline of specialist species: toward a

global functional homogenization? *Frontiers in Ecol & Environ* **9**, 222–228.

(doi:10.1890/080216)

25. Spooner FEB, Pearson RG, Freeman R. 2018 Rapid warming is associated with population

decline among terrestrial birds and mammals globally. *Global Change Biology* **24**, 4521–

4531. (doi:10.1111/gcb.14361)

26. Gjoni V, Glazier DS. 2020 A Perspective on Body Size and Abundance Relationships across

Ecological Communities. *Biology* **9**, 42. (doi:10.3390/biology9030042)

27. Ferraro KM, Schmitz OJ, McCary MA. 2022 Effects of ungulate density and sociality on

landscape heterogeneity: a mechanistic modeling approach. *Ecography* **2022**, ecog.06039.

(doi:10.1111/ecog.06039)

28. Geremia C, Merkle JA, Eacker DR, Wallen RL, White PJ, Hebblewhite M, Kauffman MJ.

2019 Migrating bison engineer the green wave. *Proceedings of the National Academy of*

Sciences **116**, 25707–25713. (doi:10.1073/pnas.1913783116)

29. Wilensky U. 1999 NetLogo. https://ccl.northwestern.edu/netlogo/

30. Ford AT, Goheen JR, Otieno TO, Bidner L, Isbell LA, Palmer TM, Ward D, Woodroffe R,

Pringle RM. 2014 Large carnivores make savanna tree communities less thorny. *Science*

346, 346–349.

31. Augustine DJ. 2004 Influence of Cattle Management on Habitat Selection By Impala on

Central Kenyan Rangeland. *Journal of Wildlife Management* **68**, 916–923.

(doi:10.2193/0022-541x(2004)068[0916:iocmoh]2.0.co;2)

32. Kingswood SC, Kumamoto AT. 1996 *Madoqua guentheri*. *Mammalian Species* **539**, 1–10.

33. Otieno TO, Goheen JR, Webala PW, Mwangi A, Osuga IM, Ford AT. 2019 Human- and

risk-mediated browsing pressure by sympatric antelope in an African savanna. *Biological*

Conservation **232**, 59–65. (doi:10.1016/j.biocon.2019.01.028)

34. Mech SG, Zollner PA. 2002 Using body size to predict perceptual range. *Oikos* **98**, 47–52.

(doi:10.1034/j.1600-0706.2002.980105.x)

35. Nagy KA. 2001 Food requirements of wild animals: predictive equations for free-living

mammals, reptiles, and birds. *Nutrition Abstracts and Reviews*

36. Charnov EL. 1976 Optimal foraging, the marginal value theorem. *Theoretical Population*

Biology **9**, 129–136. (doi:10.1016/0040-5809(76)90040-X)

37. Moran PAP. 1950 Notes on Continuous Stochastic Phenomena. *Biometrika* **37**, 17.

(doi:10.2307/2332142)

38. Salecker J, Sciaini M, Meyer KM, Wiegand K. 2019 The nlrx r package: A next-generation

framework for reproducible NetLogo model analyses. *Methods Ecol Evol* **10**, 1854–1863.

(doi:10.1111/2041-210X.13286)

39. Bivand RS, Wong DWS. 2018 Comparing implementations of global and local indicators of

spatial association. *TEST* **27**, 716–748. (doi:10.1007/s11749-018-0599-x)

40. Pace N, Rathbun EN. 1945 Studies on Body Composition. *Journal of Biological Chemistry*

158, 685–691. (doi:10.1016/S0021-9258(19)51345-X)

41. Ogutu JO, Piepho HP, Kanga E. 2012 Dynamics of an Insularized and Compressed Impala

Population: Rainfall, Temperature and Density Influences. *TOECOLJ* **5**, 1–17.

(doi:10.2174/1874213001205010001)

42. Owen-Smith N. 2008 Changing vulnerability to predation related to season and sex in an

African ungulate assemblage. *Oikos* **117**, 602–610. (doi:10.1111/j.0030-1299.2008.16309.x)

43. Singer FJ, Schoenecker KA. 2003 Do ungulates accelerate or decelerate nitrogen cycling?

Forest Ecology and Management **181**, 189–204. (doi:10.1016/S0378-1127(03)00133-6)

44. Hanski I. 1999 *Metapopulation Ecology*. Oxford University Press.

45. Loreau M, Mouquet N, Holt RD. 2003 Meta-ecosystems: a theoretical framework for a

spatial ecosystem ecology. *Ecology Letters* **6**, 673–679. (doi:10.1046/j.1461-

0248.2003.00483.x)

46. Macarthur RH, Wilson EO. 1967 *The Theory of Island Biogeography*. Princeton University

Press.

47. Bampoh D, Earl JE, Zollner PA. 2019 Examining the relative influence of animal movement

patterns and mortality models on the distribution of animal transported subsidies. *Ecological*

Modelling **412**, 108824. (doi:10.1016/j.ecolmodel.2019.108824)

48. Earl JE, Zollner PA. 2014 Effects of animal movement strategies and costs on the

distribution of active subsidies across simple landscapes. *Ecological Modelling* **283**, 45–52.

(doi:10.1016/j.ecolmodel.2014.03.020)

49. Keinath DA, Doak DF, Hodges KE, Prugh LR, Fagan W, Sekercioglu CH, Buchart SHM,

Kauffman M. 2017 A global analysis of traits predicting species sensitivity to habitat

fragmentation. *Global Ecology and Biogeography* **26**, 115–127. (doi:10.1111/geb.12509)

50. Ord TJ, Hundt PJ. 2020 Crossing extreme habitat boundaries: Jack-of-all-trades facilitates

invasion but is eroded by adaptation to a master-of-one. *Functional Ecology* **34**, 1404–1415.

(doi:10.1111/1365-2435.13600)

51. Romanuk TN, Zhou Y, Brose U, Berlow EL, Williams RJ, Martinez ND. 2009 Predicting

invasion success in complex ecological networks. *Philosophical Transactions of the Royal*

Society B: Biological Sciences **364**, 1743–1754. (doi:10.1098/rstb.2008.0286)

52. Germain RM, Johnson L, Schneider S, Cottenie K, Gillis EA, MacDougall AS. 2013 Spatial

Variability in Plant Predation Determines the Strength of Stochastic Community Assembly.

The American Naturalist **182**, 169–179. (doi:10.1086/670928)

53. Pacifici M, Rondinini C, Rhodes JR, Burbidge AA, Cristiano A, Watson JEM, Woinarski

JCZ, Di Marco M. 2020 Global correlates of range contractions and expansions in terrestrial

mammals. *Nat Commun* **11**, 2840. (doi:10.1038/s41467-020-16684-w)

54. Enquist BJ, Abraham AJ, Harfoot MBJ, Malhi Y, Doughty CE. 2020 The megabiota are

disproportionately important for biosphere functioning. *Nat Commun* **11**, 699.

(doi:10.1038/s41467-020-14369-y)

55. Coetsee C, Wigley BJ, Sankaran M, Ratnam J, Augustine DJ. 2023 Contrasting Effects of

Grazing vs Browsing Herbivores Determine Changes in Soil Fertility in an East African

Savanna. *Ecosystems* **26**, 161–173. (doi:10.1007/s10021-022-00748-7)

56. Hempson GP *et al.* 2015 Ecology of grazing lawns in Africa. *Biological Reviews* **90**, 979–

994. (doi:10.1111/brv.12145)

57. Ryser R, Hirt MR, Häussler J, Gravel D, Brose U. 2021 Landscape heterogeneity buffers

biodiversity of simulated meta-food-webs under global change through rescue and drainage

effects. *Nat Commun* **12**, 4716. (doi:10.1038/s41467-021-24877-0)

58. Mackay J, Nikiforuk K, Szojka M, Little CJ, Fleri JR, Germain RM. 2021 Animals connect

plant species and resources in a meta-ecosystem. *Landscape Ecol* **36**, 1621–1629.

(doi:10.1007/s10980-021-01244-4)