Spatial fingerprint of consumer body size and habitat preference on resource distribution

Chelsea J. Little[1,2] *, Pierre Etienne Banville[1,3], Adam T. Ford[4], Rachel M. Germain[2]

[1] School of Environmental Science, Simon Fraser University, Burnaby, B.C., Canada V5A 1S6

[2] Department of Zoology, Biodiversity Research Center, University of British Columbia,

Vancouver, B.C., Canada V6T 1Z4

[3] Department of Geography, Simon Fraser University, Burnaby, B.C., Canada V5A 1S6

[4] Department of Biology, University of British Columbia, Kelowna, B.C., Canada V1V 1V7

* corresponding author: chelsea_little@sfu.ca

Abstract

Consumers shape spatial patterns on landscapes by amplifying or dampening environmental

heterogeneity through feeding, excretion, and movement of resources. The degree to which the

environment is modified by consumers depends on species' traits, including body mass,

movement behavior, sociality, and habitat specialization. Global change is altering the size and

traits of consumer populations, but our understanding of how this may impact resource

heterogeneity is limited. Here, we developed an individual-based model of habitat specialists'

and generalists' movement and activity in a patchy landscape and investigated the impact of

changes in population and mean body sizes on landscape-scale resource heterogeneity. We found

that consumers specializing on low-resource habitats (a common risk avoidance strategy)

increased spatial resource heterogeneity regardless of their population and body size. By

contrast, generalists eroded differences among habitats, and we further found that resource

heterogeneity decreased with the average body size of generalist consumers, even while

controlling for total consumer biomass. These nuanced spatial outcomes of consumer-resource

interactions emerge from the unique metabolic demands of specialists vs. generalists, which

scale nonlinearly with body size. Since global change disproportionately impacts larger species

and specialists, indirect consequences on ecosystems may arise via biotic processes, affecting

spatial heterogeneity of future landscapes.

Introduction

Resource landscapes are often thought of as an environmental template that organisms respond

to, but this neglects the role that organisms play in shaping spatial distributions of resources.

Animal consumers serve as important resource vectors in landscapes (Earl and Zollner 2017,

McInturf et al. 2019, Little et al. 2022, Pringle et al. 2023). The deposition of resources via feces,

urine, seeds, placenta, and carcasses contribute to resource dispersal in the landscape (McIntyre

et al. 2008, Bump et al. 2009, Sitters et al. 2017, Schmitz et al. 2018, le Roux et al. 2020, Ferraro

et al. 2023). Organismal activity will therefore impact spatial resource heterogeneity, which in

turn facilitates the coexistence of species (Amarasekare 2003) and thus ecosystem functioning

(Loreau et al. 2003b). The distribution of resources in space is therefore an important landscape

characteristic that shapes patterns of biodiversity and ecosystem functioning (Tylianakis et al.

2008, Seiferling et al. 2014, Gonzalez et al. 2020, Rizzuto et al. 2024). With global change

altering abiotic conditions and consumer populations, it is imperative to gain a better

understanding of the processes shaping spatial resource heterogeneity.

The traits of species making up consumer communities may provide a key to

understanding organismal effects on resource landscapes. A handful of studies have established

that species' traits, such as behavior and body size, determine the amount, stoichiometry, and

scale of consumer-driven resource cycling (Sitters et al. 2017, Veldhuis et al. 2018, Subalusky

and Post 2019). For example, social aggregation creates spatial clustering of consumer-driven

resource subsidies (Bampoh et al. 2021). Other traits such as diet preferences and risk avoidance

strategies may concentrate consumer activity into specific habitat types, leading to divergent

rates of resource cycling across the landscape (Peller et al. 2022). Meanwhile, consumer body

size influences resource dispersal through at least two pathways. First, energetic requirements

scale nonlinearly with body size (Brown et al. 2004), and consequently so should resource

consumption and excretion. Second, body size is associated with differences in movement ability

and perceptual range (Little et al. 2022, Albaladejo-Robles et al. 2023, Straus et al. 2024), and

thus the spatial dispersion of foraging and resource deposition.

Predicted shifts in consumer populations and communities due to global change may

rewire resource distributions, yet understanding how they do so is made more challenging

because species' responses to global change vary, in part due to their traits. Larger animals are

more impacted by climate change and land use change (Albaladejo-Robles et al. 2023), and

specialist species more than generalists (Clavel et al. 2011). Moreover, as population sizes shift

with global change (Spooner et al. 2018), we can anticipate shifts in patterns of resource

consumption and deposition across landscapes; however, our understanding of the net effect of

these shifts on spatial resource distribution is limited. Because body size and population size

often trade off (Gjoni and Glazier 2020), changes in body size might offset the effects of changes

in population size and lead to unique spatial patterns of resource distributions that could not be

predicted from body size alone. In one of the few examples of research addressing this topic,

Ferraro *et al.* (2022) used an individual-based model of foraging caribou and showed that higher

densities of caribou increased nutrient spatial heterogeneity through carcass depositions

(resource hotspots) and over-grazed dead zones (resource coldspots). At even larger spatial

scales, higher densities of bison increase the amount and change the timing of nutrients available

in different regions of grassland landscapes (Geremia et al. 2019).

In this study, we present an empirically-calibrated, individual-based model of consumer

movement and activity where we factorially manipulated i) population sizes, ii) mean body size,

iii) risk-based habitat preferences, and iv) sociality, and investigated effects on spatial resource

heterogeneity in a patchy landscape. Our model was parameterized to capture key biological

features of two consumer species—a generalist, and a specialist that avoids high-nutrient

resource patches (a common predator avoidance strategy)—with scaled movement distances,

metabolic requirements, and perceptual ranges associated with body size.

We hypothesized that larger populations (compared to smaller populations) of generalists

would preferentially graze high-quality areas until they resemble lower-quality areas, reducing

spatial heterogeneity. By contrast, we predicted that by removing resources from lower-quality

areas, specialists would accentuate the differences between nutrient-rich and nutrient-poor parts

of the landscape, and thus larger populations (compared to smaller populations) of specialists

would increase spatial heterogeneity. We also expected that, while keeping the total consumer

biomass constant, body size would have contrasting effects on resource redistribution by

generalists and specialists: decreasing the mean body size of generalists would reduce resource

heterogeneity, while reducing the mean body size of specialists would amplify resource

heterogeneity. This is because the metabolic load of smaller individuals is higher, and thus more

resources should be consumed by a population of many small individuals than an equivalent

biomass of fewer large consumers. Therefore, we predicted that for a given population biomass,

the largest differences in the impacts of consumers on resource heterogeneity would occur

between small-bodied specialists (which would amplify heterogeneity) and small-bodied

generalists (which would dampen heterogeneity). Finally, we predicted that spatial aggregation

of consumers would increase clustering of consumers' impacts on the landscape, thus increasing

spatial resource heterogeneity.

Methods

99 *Modeling approach*

100 We used NetLogo v6.0.4 (Wilensky 1999) to model generalists and specialists moving in a

101 landscape, where consumers withdraw resources by foraging and deposit resources through feces

102 and carcasses. Key parameters are indicated in Table 1, and a detailed explanation is found in

103 Appendix I. This landscape is divided into a 141 x 141 grid of equal-sized pixels, each pixel

104 representing 10 m^2. The landscape has two discrete habitat types (**Fig. 1**), approximating a

105 Kenyan savanna (see Ford et al. [2014]): open, high-nutrient glades as clusters of pixels (each

106 cluster forming a "Patch"), in a "Matrix" of bushy, lower-quality grassland (Augustine 2004).

107 Each pixel is classified as Patch or Matrix and is assigned a forage amount drawn from a normal

108 distribution centered on 50 g/m^2 for Patches and 33 g/m^2 for Matrix, with forage containing 6%

109 nitrogen (N) in Patches and 4% in Matrix. Landscapes contained an average of four Patches,

110 making up ~5% of the landscape area (**Fig. 1**).

111  Individuals in the model are either generalists or specialists. Specialist consumers,

112 resembling Guenther's dik-diks, stay in the Matrix to avoid predators (Kingswood and Kumamoto

113 1996, Otieno et al. 2019), while generalists, akin to impalas, can use both habitat types, but favor

114 Patches (Ford et al. 2014, Otieno et al. 2019). Animals forage based on their metabolic needs and

115 move within their perceptual range, both of which vary as a function of their body size (Nagy

116 2001, Mech and Zollner 2002). Foraging behavior adheres to the marginal value theorem, with

117 animals selecting pixels with better-than-average forage available (Charnov 1976). Patch pixels

118 are not considered by specialists, while generalists can choose any pixel (Patch or Matrix) and

119 preferentially graze Patches at the start of each simulation because of the higher forage and N

120 concentration. Model timesteps represent 3 hours of movement/activity (see Appendix I for details

121   on movement, foraging, excretion, and death), and simulations were run for three months, or the

122   length of one growing season.

Figure 1. N distributions within Patches and Matrix at the beginning (top left) and end (right) of
the growing season in simulated landscapes, which are based on a Kenyan savanna (bottom left)
with habitat specialists that forage only in Matrix, and generalists that forage across the whole
landscape but prefer Patches. In our model, consumers realistically increase perceptual range,
movement ability, and metabolic needs with body size. N ranges from low (yellow) to intermediate
(orange) and high (green). On right, N distributions after grazing, movement, and nutrient
deposition for three example scenarios out of the study's total 64 (see Table S1): here, by equal
numbers of large (40kg) specialists, large generalists, or small (10kg) generalists in example
simulations. Percent changes in Moran's I, a measure of spatial autocorrelation (Moran 1950), and
in the nitrogen content of each pixel in Patches and Matrix are indicated for each simulation. In
the upper right panel, note the waves of N distribution in the Matrix created by foraging specialists
compared to the initial landscape. Small dark green spots at the end of the growing season represent
nutrient deposition by carcasses of consumers that died during the simulation, which is most
evident for larger consumers. Real landscape image © 2024 Maxar Technologies.

Simulation Scenarios

We designed our scenarios (Table S1) to look at the impact of 4 different factors on landscape

heterogeneity: species role (habitat generalists or specialists), aggregation type (solitary or herd),

body size, and total biomass on the landscape where the total biomass on the landscape is

determined by the mean body size multiplied by the number of individuals. We tested

combinations of aggregation type (solitary or herd), body size (10kg, 25kg or 40kg), and total

biomass on the landscape (varying between 500kg and 6000kg) for each of the species. We

varied the total biomass to allow for comparisons where the total number of individuals is the

same (50, 100, or 150 individuals) and where the total biomass is the same (1000kg, 2000kg, or

4000kg). We varied those factors to test all combinations possible. This resulted in 64 scenarios

as described in Table S1. All scenarios were replicated 25 times. The Netlogo simulations were

processed through R version 4.2.2 using the package 'nlrx' (Salecker *et al.* 2019). They were

performed on the high-performance computer Cedar from Digital Research Alliance of Canada.

The outputs from the simulation experiments focus on changes that happen in the

distribution of nutrients on the landscape and are recorded at the pixel level. They include the

initial and final forage amount, the initial and final N amount, the N eaten through foraging, the

N deposited through defecation or animal death, the coordinates and whether the pixel is Patch or

Matrix. We also recorded the number of times a pixel was grazed, defecated on, and the number

of deaths that occurred on a given pixel to help with results interpretation.

Table 1: Key parameters influencing the initial distribution of N on the landscape and the
withdrawals and deposits of N by animals.

Parameter (units)	Value	Reference
Mean forage per Patch pixel (g)	5000	(Augustine 2004)
Mean forage per Matrix pixel (g)	3300	(Augustine 2004)
Mean N per Patch pixel (g)	300 (6% of forage)	(Augustine 2004)
Mean N per matrix patch (g)	132 (4% of forage)	(Augustine 2004)
N in consumers' body (g)	2% * bodymass	(Pace and Rathbun 1945)
Daily metabolic needs of consumers (g)	$(7.94 * (bodymass)^{0.646} / 11.5)$	(Nagy 2001)
Yearly % chance of death for animals	10%	(Ogutu et al. 2012, Owen-Smith and Goodall 2014)
N deposited on landscape per defecation event (g)	50% of N absorbed since last defecation event	(Singer and Schoenecker 2003)
N deposited on landscape per death event (g)	All body N + 100% N absorbed since last defecation event	

Model Analysis

We analyzed changes in the spatial and non-spatial nutrient distribution across the landscape to

determine the effect of species role, aggregation type, body size, and total biomass on landscape

heterogeneity. The key variables of interest were the initial and final N amount of each pixel. For

each individual simulation, we calculated the four statistical moments (i.e. mean, variance,

skewness and kurtosis) for the initial and final N amount across all pixels. Comparing the value

of those statistics for the initial N and final N amount indicates the change in the non-spatial N

distribution across the landscape resulting from the simulation. For each scenario, we then

calculated the mean and standard deviation of the 25 replicates for all statistics calculated at the

individual simulation level. For a given statistic, we then plotted the mean with a standard

deviation error bar of the specific scenarios that we wanted to compare, such as different body

sizes or different number of individuals on the landscape. All statistical analyses were performed

using R version 4.2.2 and the skewness and kurtosis were calculated using the package 'geodiv'

(Smith et al. 2021).

We also explored the spatial changes in N distribution by calculating the Global Moran's

I ("Moran's I") on initial and final N for each individual simulation. The Moran's I is a statistical

test that measures spatial autocorrelation, indicating how similar are the pixels from their

neighbors (Moran 1950). The Moran's I sums covariances between pixels and, in its standard

form using Queen connectivity (pixels are neighbors if they share a border or a point), gives

equal weight to the 8 closest neighbors (first order neighbors) of a given pixel in the calculations.

An increase in Moran's I means that the pixels are more autocorrelated, indicating spatial

clustering of resources and increased spatial heterogeneity. A decrease in Moran's I means that

the pixels are less autocorrelated and the distribution of N is less clustered on the landscape.

To determine if the relative impact of body size on spatial autocorrelation varies

depending on the size of the neighborhood considered, we also calculated Moran's I using 4th

order neighbors (80 neighbors), 10th order neighbors (440 neighbors) and 17th order neighbors

(1224 neighbors) with an inverse distance spatial weight (i.e. the correlation between the closest

neighbors weighs more than the correlation between neighbors further away). The 4th, 10th, and

17th order neighbors were selected to approximate the perceptual range of individuals with a

body size of 10kg, 25kg, and 40kg respectively. Individuals with larger body sizes can access

pixels that are located further away than individuals with smaller body sizes, but are more likely

to select nearby pixels to minimize energy costs. We found no qualitative differences in the

results between the "queen" configuration Moran's I and Moran's I with 4th, 10th, and 17th order

neighbors, and hence present results with the standard Moran's I only. The Moran's I analysis

was conducted using the package 'spdep' (Bivand and Wong 2018) in R version 4.2.2.

To better understand the drivers of the changes in the spatial and non-spatial N

distribution at the landscape level, we analyzed the variables making up the final amount of N

amount (for a given pixel, Final N = Initial N + N Deposited – N Eaten) at the landscape level

and separately for Patches and Matrix. We calculated the mean and variance for N deposited, N

eaten, and net N (difference between N Deposited and N Eaten) for the landscape as a whole and

for Patches and Matrix separately for each individual simulation. As potential explanatory

variables for N dynamics, we calculated the mean and variance in the number of times a pixel

was grazed and defecated on, and the total number of death events at the landscape level for each

individual simulation.

Results

Spatial distribution of N varied across simulated landscapes at the end of the growing season.

Although larger body sizes and larger population sizes led to more forage consumed (**Fig. 2**) and

greater N depletion (**Fig. S1**) in the total landscape for both specialists and generalists, changes

in the spatial heterogeneity of N resulted from the interplay between N depletion in Patches and

Matrix.

Figure 2. Total forage consumed during simulations with different combinations of parameters.
(a) and (b) show the effect of increasing population size (on the x-axis) at small, medium, or
large mean individual body size (left, center, and right panels), when the landscape is populated
either with specialists that forage only in Matrix (a) or generalists that forage in both habitats (b).
(c) and (d) show the effect of increasing mean body size (on the x-axis) when the total
population biomass on the landscape is 1000 kg (left column), 2000 kg (middle column), or 4000
kg (right column) of specialists (c) or generalists (d). Bars show mean ± one standard deviation
of 25 simulations for each scenario.

As predicted, solitary habitat specialists foraging only in the Matrix accentuated

differences between habitat types (note increase in Moran's I in **Fig. 3,** top panels); for these

specialists, larger population sizes and larger body sizes were associated with greater increases in

spatial heterogeneity. Also as predicted, solitary generalists, who preferentially forage in

Patches, homogenized the landscape, reducing differences between Patches and Matrix (note

decrease in Moran's I in **Fig. 3,** bottom panels). Here, we have presented results only from

scenarios where consumers are solitary. Results from herd simulations were qualitatively similar,

but produced slightly higher levels of spatial resource heterogeneity (**Fig. S2**), in line with our

predictions.

Figure 3. Change in Moran's I for the distribution of nitrogen between beginning and end of
simulations. (a) and (b) show the effect of increasing population size (on the x-axis) at small,
medium, or large mean individual body size (left, center, and right panels), when the landscape is
populated either with specialists that forage only in Matrix (a) or generalists that forage in both
habitats (b). (c) and (d) show the effect of increasing mean body size (on the x-axis) when the
total population biomass on the landscape is 1000 kg (left column), 2000 kg (middle column), or
4000 kg (right column) of specialists (c) or generalists (d). Bars show mean ± one standard
deviation of 25 simulations for each scenario. Note different y-axes in top and bottom panels.

241   For small generalists, larger population sizes led to greater homogenization of N

242 distribution over the course of the simulations than that caused by smaller population sizes. Since

243 small generalists, even at larger population sizes, did not remove enough forage from the Patches

244 to make them of comparable quality as the Matrix, N removal from the Matrix was negligible

245 (i.e., the first panel of **Fig. 4c**) and larger population sizes were nearly linearly associated with

246 reductions in Moran's I (i.e., the first panel in **Fig. 4b**). However, for large generalists, the

247 reduction in Moran's I was consistent across population sizes. Larger generalists depleted

248 Patches faster than smaller generalists, subsequently removing N from the Matrix (**Fig. 4a**).

249 However, the increased N depletion of Patches by the largest population sizes of large generalists

250 (x-axis in rightmost panel of **Fig. 4b**) was of a similar magnitude to their increased removal of N

251 from the Matrix (x-axis in rightmost panel of **Fig. 4c**), and as a result the difference between

252 Patch and Matrix quality (and thus Moran's I) was fairly stable across population sizes of large

253 generalists (**Fig. 3a**).

Figure 4. Changes in forage consumed in Patches and Matrix and in the spatial distribution of nitrogen in simulations with varying generalist body size and population size. Each dot represents one simulation, with the color of the dots indicating the number of individuals in the simulation. Within a panel, the leftmost graph shows results from simulations with 10 kg as the generalists' mean body size, 25 kg for middle graph, and 40 kg for right graph. (a) Relationship between the total forage consumed in Patches and the total forage consumed in the Matrix. (b) Change in Moran's I varies nearly linearly with net nitrogen loss from Patches when generalists are small, but less so when generalists are large. (c) When generalists are small, net nitrogen loss varies only minimally with population size, yet the spatial distribution of that nitrogen substantially changes at larger population sizes; when generalists are large, there is much greater variation in net nitrogen loss in the Matrix depending on population size, but less impact of population size on the spatial clustering of nitrogen distribution.

Comparing scenarios with equal total population biomass but with many small versus

fewer large consumers revealed the effects of nonlinear scaling of metabolic requirements on

spatial distribution of N (**Fig. 5**). Total forage consumption was higher with many small

consumers compared to few large ones (**Fig. 2**). At lower total biomass, generalists of all body

sizes spent most of their time in Patches (**Fig. 5b)**, but small generalists consumed more forage

there (**Fig. 5b**), leading to a stronger homogenizing effect than their larger counterparts (**Fig. 3c**)

as hypothesized. However, at larger total biomass, smaller generalists depleted and left Patches

and spent upwards of 60% of their time in the Matrix (**Fig. 5c**), consuming markedly more

forage in the Matrix than populations of large generalists did despite similar forage consumption

in Patches (**Fig. 5a**), and thus changes in Moran's I were less differentiated.

Figure 5. Changes in forage consumed in Patches and Matrix with different amounts of total generalist biomass on the landscape (shown in left, center, and right panels), and when that biomass is made up of different average body sizes (indicated by color of points). (a) Relationship between the total forage consumed in Patches and total forage consumed in the Matrix. (b) Relationship between the mean percentage of time individuals spent in Patches and the total forage consumed in Patches. (c) Relationship between the mean percentage of time individuals spent in the Matrix and total forage consumed in the Matrix.

Discussion

The role of animals in structuring resource patterns is poorly understood. Here, using a spatially-

explicit, individual-based model, we show that organismal traits can make the difference in

whether consumers amplify or erode spatial resource heterogeneity. We found that a key trait

determining whether consumers amplify or erode resource patchiness is species' activity in the

different habitats, modeled here to represent different risk avoidance strategies. Patchy

landscapes have driven the development of ecological theory (Macarthur and Wilson 1967,

Hanski 1999, Loreau et al. 2003a), yet to date, models assessing the effects of animal movement,

foraging, and resource deposition have primarily focused on landscapes without spatial structure

in resource availability (Earl and Zollner 2014, Bampoh et al. 2019, 2021, Ferraro et al. 2022).

When spatial structure is included, over seasonal time scales, we showed that specialists on

lower-quality (Matrix) habitat amplified existing resource heterogeneity between the resource

hotspots (which they avoided) and the low-quality habitat. Conversely, generalists who could

take advantage of high-resource Patches eroded the landscape's spatial structure. This is a

particularly important insight in an era of global change because generalists often thrive where

specialist species are more vulnerable to habitat loss and fragmentation (Clavel et al. 2011,

Keinath et al. 2017), and because of the proliferation of non-native species and the association

between generalism and invasion success (Romanuk et al. 2009, Ord and Hundt 2020).

The specialists in our simulations provided an example of how risk avoidance strategies

can reinforce spatial structure and patchiness of landscapes by strengthening existing resource

gradients (for an empirical example, see [Germain et al. 2013]). In addition to accentuating the

difference between Patch and Matrix habitat, these same specialists also created finer-scale

spatial resource heterogeneity where it was previously lacking: our landscapes featured no initial

spatial clustering of resources *within* a habitat type, but over the course of our simulated growing

season, spatial heterogeneity in the Matrix increased. This was because the removal of nutrients

by foraging followed individuals' movement patterns, which were spatially autocorrelated at the

scale of the individual's perceptual range. This increase in fine-scale heterogeneity was not

captured by Moran's I, which better reflected the larger differences in nutrient concentrations

between Matrix and Patches, but is visible as waves of resource concentrations in landscape

depictions (**Fig. 1**, "150 Large Specialists"). Thus, the effect of organismal activity on nutrient

spatial heterogeneity depends on both the existing structure of the landscape and the scale of

observation.

Globally, many species of large-bodied vertebrates are experiencing range contractions,

whereas smaller-bodied vertebrates are more likely to be experiencing range expansions (Pacifici

et al. 2020), and the largest consumer species are also at higher risk of extinction than are their

smaller counterparts (Enquist et al. 2020). Our results imply that in patchy landscapes, the

impact of species turnover from large to small generalists depends on both initial community

biomass and whether shifts in body size are accompanied by altered total biomass (i.e., the

summed biomass of all individuals in the population). For the same total biomass, smaller

consumers consume more forage due to higher mass-specific metabolic needs (Brown et al.

2004). In terms of spatial ramifications, at small body sizes and low total biomass, larger

populations of generalists homogenized the landscape more than smaller populations by

depleting resource hotspots. However, at larger body sizes and greater total biomass, foraging

behavior whereby generalists left depleted Patches to forage in the Matrix meant that the further

depletion of resource hotspots by a larger population was offset by the increased consumption of

resources in the lower quality habitat—that is, spatial homogenization plateaued. Thus, we

suggest that where generalists are already dominant, a shift from large to small body size may

have minimal effects on spatial structure if total biomass remains the same, however, such a shift

would increase overall rates of consumption and recycling of resources. This is because smaller

individuals have higher metabolic demands per unit mass compared to larger individuals (a

universal feature of metabolic scaling).

Our model did not include population dynamics of either consumers (beyond mortality)

or forage resources, and only considered the spatial dynamics of net resource availability. This

increased computational feasibility and enabled us to isolate the impact of vertebrates on nutrient

distributions. Despite simplifications, our finding that generalists (which we based on impalas)

homogenize N distributions by reducing N in Patches aligns with empirical results in an

intriguing way: impalas increase the relative density of a well-defended plant species (Ford et al.

2014) which has less foliar N than a poorly-defended congener (Coetsee et al. 2023) that is

abundant where impala are rare, thus altering spatial heterogeneity of N in plants.

Another caveat is that in our model, not all N consumed in forage is returned to the

landscape. Some N is incorporated into consumer bodies, and consumer death was not high

enough in our model to offset this. This dynamic is not unrealistic over the course of a single wet

season, based on mortality estimates from the literature (Ogutu et al. 2012, Owen-Smith and

Goodall 2014). We also did not explicitly model processes such as predator and scavenger export

of nutrients, leaching, or nutrient volatilization, instead incorporating assumptions about how

much N was returned to the landscape by excretion, egestion, and carcass deposition (see

Appendix I, and **Fig. S3** for a sensitivity test).

Here, we showed through simulations that animals have the ability to shape resource

heterogeneity on the landscape over short periods of time such as a single season, thus shifting

the environmental template for subsequent consumer-resource interactions. Landscape

heterogeneity is important because it creates different niches that can serve to sustain

biodiversity (Ryser et al. 2021). Our results suggest that species' traits and abundance are key

mediators of their effect on resource distributions, which could lead to varying impacts of future

biodiversity change. We propose two future directions for spatially-explicit, agent-based

modeling to explore the impacts of these projected changes. The first is to explicitly model

different chemical forms of nutrients, which are more or less bioavailable to organisms

(Hempson et al. 2015). This approach would account for the fact that nutrients deposited through

dung and urine are not immediately available to consumers, but must first be transformed into

plant shoots. We expect that coupling these considerations with longer simulations that account

for consumer population dynamics would lead to new spatial patterns of foraging movement by

consumers, and thus new emergent dynamics of nutrient spatial distributions. Second, further

work should also explore how competing consumers and trophic interactions modify the patterns

found in this study. For example, we modeled species based on known risk-avoidance strategies

rather than including predators in the models, but doing so may lead to behavioral shifts and

emergent dynamics, additionally because trophic role influences resource deposition (Mackay et

al. 2021). Furthermore, consumers exist in multi-species communities and trait-based differences

among co-occurring species could lead to complementarity of resource use, altering the patterns

that we found here. Incorporating additional ecological complexity into spatially-explicit models

could enable a better understanding of how organisms will shape their own fates via resource

dynamics in future landscapes.

Data / code availability

Data and code are available to peer reviewers at

https://vault.sfu.ca/index.php/s/YT0XQKrzX0gZqlV. Data and code will be deposited in Dryad

before publication of the manuscript.

Acknowledgments

This research was enabled in part by support from the BC DRI Group, Simon Fraser University's

Supercomputer Cedar (www.sfu.ca/research/supercomputer-cedar), and the Digital Research

Alliance of Canada (www.alliancecan.ca). We thank Matteo Rizzuto and members of the Little

Ecology Group for valuable feedback on the manuscript.

Funding

This work was supported by a Killam Postdoctoral Research Fellowship from the Killam Trusts

and a Discovery Grant from the Natural Science and Engineering Research Council of Canada.

References

Albaladejo-Robles, G., Böhm, M. and Newbold, T. 2023. Species life-history strategies affect population responses to temperature and land-cover changes. - Global Change Biology 29: 97–109.

Amarasekare, P. 2003. Competitive coexistence in spatially structured environments: a synthesis. - Ecology Letters 6: 1109–1122.

Augustine, D. J. 2004. Influence of Cattle Management on Habitat Selection By Impala on
Central Kenyan Rangeland. - Journal of Wildlife Management 68: 916–923.

Bampoh, D., Earl, J. E. and Zollner, P. A. 2019. Examining the relative influence of animal
movement patterns and mortality models on the distribution of animal transported
subsidies. - Ecological Modelling 412: 108824.

Bampoh, D. K., Earl, J. E. and Zollner, P. A. 2021. Simulating the relative effects of movement
and sociality on the distribution of animal-transported subsidies. - Theor Ecol 14: 57–70.

Bivand, R. S. and Wong, D. W. S. 2018. Comparing implementations of global and local
indicators of spatial association. - TEST 27: 716–748.

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. and West, G. B. 2004. Toward a
metabolic theory of ecology. - Ecology 85: 1771–1789.

Bump, J. K., Webster, C. R., Vucetich, J. A., Peterson, R. O., Shields, J. M. and Powers, M. D.
2009. Ungulate Carcasses Perforate Ecological Filters and Create Biogeochemical
Hotspots in Forest Herbaceous Layers Allowing Trees a Competitive Advantage. -
Ecosystems 12: 996–1007.

Charnov, E. L. 1976. Optimal foraging, the marginal value theorem. - Theoretical Population
Biology 9: 129–136.

Clavel, J., Julliard, R. and Devictor, V. 2011. Worldwide decline of specialist species: toward a
global functional homogenization? - Frontiers in Ecol & Environ 9: 222–228.

Coetsee, C., Wigley, B. J., Sankaran, M., Ratnam, J. and Augustine, D. J. 2023. Contrasting
Effects of Grazing vs Browsing Herbivores Determine Changes in Soil Fertility in an
East African Savanna. - Ecosystems 26: 161–173.

Earl, J. E. and Zollner, P. A. 2014. Effects of animal movement strategies and costs on the
distribution of active subsidies across simple landscapes. - Ecological Modelling 283:
45–52.

Earl, J. E. and Zollner, P. A. 2017. Advancing research on animal-transported subsidies by
integrating animal movement and ecosystem modelling. - Journal of Animal Ecology 86:
987–997.

Enquist, B. J., Abraham, A. J., Harfoot, M. B. J., Malhi, Y. and Doughty, C. E. 2020. The
megabiota are disproportionately important for biosphere functioning. - Nat Commun 11:
699.

Ferraro, K. M., Schmitz, O. J. and McCary, M. A. 2022. Effects of ungulate density and sociality
on landscape heterogeneity: a mechanistic modeling approach. - Ecography 2022:
ecog.06039.

Ferraro, K. M., Welker, L., Ward, E. B., Schmitz, O. J. and Bradford, M. A. 2023. Plant
mycorrhizal associations mediate the zoogeochemical effects of calving subsidies by a
forest ungulate. - Journal of Animal Ecology 92: 2280–2296.

Ford, A. T., Goheen, J. R., Otieno, T. O., Bidner, L., Isbell, L. A., Palmer, T. M., Ward, D.,
Woodroffe, R. and Pringle, R. M. 2014. Large carnivores make savanna tree communities
less thorny. - Science 346: 346–349.

Geremia, C., Merkle, J. A., Eacker, D. R., Wallen, R. L., White, P. J., Hebblewhite, M. and
Kauffman, M. J. 2019. Migrating bison engineer the green wave. - Proceedings of the
National Academy of Sciences 116: 25707–25713.

Germain, R. M., Johnson, L., Schneider, S., Cottenie, K., Gillis, E. A. and MacDougall, A. S.
2013. Spatial Variability in Plant Predation Determines the Strength of Stochastic
Community Assembly. - The American Naturalist 182: 169–179.

Gjoni, V. and Glazier, D. S. 2020. A Perspective on Body Size and Abundance Relationships
across Ecological Communities. - Biology 9: 42.

Gonzalez, A., Germain, R. M., Srivastava, D. S., Filotas, E., Dee, L. E., Gravel, D., Thompson,
P. L., Isbell, F., Wang, S., Kéfi, S., Montoya, J., Zelnik, Y. R. and Loreau, M. 2020.
Scaling-up biodiversity-ecosystem functioning research. - Ecology Letters 23: 757–776.

Hanski, I. 1999. Metapopulation Ecology. - Oxford University Press.

Hempson, G. P., Archibald, S., Bond, W. J., Ellis, R. P., Grant, C. C., Kruger, F. J., Kruger, L.
M., Moxley, C., Owen-Smith, N., Peel, M. J. S., Smit, I. P. J. and Vickers, K. J. 2015.
Ecology of grazing lawns in Africa. - Biological Reviews 90: 979–994.

Keinath, D. A., Doak, D. F., Hodges, K. E., Prugh, L. R., Fagan, W., Sekercioglu, C. H.,
Buchart, S. H. M. and Kauffman, M. 2017. A global analysis of traits predicting species
sensitivity to habitat fragmentation. - Global Ecology and Biogeography 26: 115–127.

Kingswood, S. C. and Kumamoto, A. T. 1996. Madoqua guentheri. - Mammalian Species 539:
1–10.

le Roux, E., van Veenhuisen, L. S., Kerley, G. I. H. and Cromsigt, J. P. G. M. 2020. Animal
body size distribution influences the ratios of nutrients supplied to plants. - Proceedings
of the National Academy of Sciences of the United States of America 117: 22256–22263.

Little, C. J., Rizzuto, M., Luhring, T. M., Monk, J. D., Nowicki, R. J., Paseka, R. E., Stegen, J.
C., Symons, C. C., Taub, F. B. and Yen, J. D. L. 2022. Movement with meaning:
integrating information into meta-ecology. - Oikos in press.

Loreau, M., Mouquet, N. and Holt, R. D. 2003a. Meta-ecosystems: a theoretical framework for a
spatial ecosystem ecology. - Ecology Letters 6: 673–679.

Loreau, M., Mouquet, N. and Gonzalez, A. 2003b. Biodiversity as spatial insurance in
heterogeneous landscapes. - Proc. Natl. Acad. Sci. U.S.A. 100: 12765–12770.

Macarthur, R. H. and Wilson, E. O. 1967. The Theory of Island Biogeography. - Princeton
University Press.

Mackay, J., Nikiforuk, K., Szojka, M., Little, C. J., Fleri, J. R. and Germain, R. M. 2021.
Animals connect plant species and resources in a meta-ecosystem. - Landscape Ecol 36:
1621–1629.

McInturf, A. G., Pollack, L., Yang, L. H. and Spiegel, O. 2019. Vectors with autonomy: what
distinguishes animal-mediated nutrient transport from abiotic vectors? - Biological
Reviews 94: 1761–1773.

McIntyre, P. B., Flecker, A. S., Vanni, M. J., Hood, J. M., Taylor, B. W. and Thomas, S. A.
2008. Fish distributions and nutrient cycling in streams: Can fish create biogeochemical
hotspots? - Ecology 89: 2335–2346.

Mech, S. G. and Zollner, P. A. 2002. Using body size to predict perceptual range. - Oikos 98:
47–52.

Moran, P. A. P. 1950. Notes on Continuous Stochastic Phenomena. - Biometrika 37: 17.

Nagy, K. A. 2001. Food requirements of wild animals: predictive equations for free-living
mammals, reptiles, and birds. - Nutrition Abstracts and Reviews in press.

Ogutu, J. O., Piepho, H. P. and Kanga, E. 2012. Dynamics of an Insularized and Compressed
Impala Population: Rainfall, Temperature and Density Influences. - TOECOLJ 5: 1–17.

Ord, T. J. and Hundt, P. J. 2020. Crossing extreme habitat boundaries: Jack-of-all-trades
facilitates invasion but is eroded by adaptation to a master-of-one. - Functional Ecology
34: 1404–1415.

Otieno, T. O., Goheen, J. R., Webala, P. W., Mwangi, A., Osuga, I. M. and Ford, A. T. 2019.
Human- and risk-mediated browsing pressure by sympatric antelope in an African
savanna. - Biological Conservation 232: 59–65.

Owen-Smith, N. and Goodall, V. 2014. Coping with savanna seasonality: comparative daily
activity patterns of African ungulates as revealed by GPS telemetry. - Journal of Zoology
293: 181–191.

Pace, N. and Rathbun, E. N. 1945. Studies on Body Composition. - Journal of Biological
Chemistry 158: 685–691.

Pacifici, M., Rondinini, C., Rhodes, J. R., Burbidge, A. A., Cristiano, A., Watson, J. E. M.,
Woinarski, J. C. Z. and Di Marco, M. 2020. Global correlates of range contractions and
expansions in terrestrial mammals. - Nat Commun 11: 2840.

Peller, T., Marleau, J. N. and Guichard, F. 2022. Traits affecting nutrient recycling by mobile
consumers can explain coexistence and spatially heterogeneous trophic regulation across
a meta-ecosystem. - Ecology Letters 25: 440–452.

Pringle, R. M., Abraham, J. O., Anderson, T. M., Coverdale, T. C., Davies, A. B., Dutton, C. L.,
Gaylard, A., Goheen, J. R., Holdo, R. M., Hutchinson, M. C., Kimuyu, D. M., Long, R.
A., Subalusky, A. L. and Veldhuis, M. P. 2023. Impacts of large herbivores on terrestrial
ecosystems. - Current Biology 33: R584–R610.

Rizzuto, M., Leroux, S. J., Schmitz, O. J., Vander Wal, E., Wiersma, Y. F. and Heckford, T. R.
2024. Animal-vectored nutrient flows across resource gradients influence the nature of
local and meta-ecosystem functioning. - Ecological Modelling 488: 110570.

Romanuk, T. N., Zhou, Y., Brose, U., Berlow, E. L., Williams, R. J. and Martinez, N. D. 2009.
Predicting invasion success in complex ecological networks. - Philosophical Transactions
of the Royal Society B: Biological Sciences 364: 1743–1754.

Ryser, R., Hirt, M. R., Häussler, J., Gravel, D. and Brose, U. 2021. Landscape heterogeneity
buffers biodiversity of simulated meta-food-webs under global change through rescue
and drainage effects. - Nat Commun 12: 4716.

Schmitz, O. J., Wilmers, C. C., Leroux, S. J., Doughty, C. E., Atwood, T. B., Galetti, M., Davies,
A. B. and Goetz, S. J. 2018. Animals and the zoogeochemistry of the carbon cycle. -
Science in press.

Seiferling, I., Proulx, R. and Wirth, C. 2014. Disentangling the environmental-heterogeneity–
species-diversity relationship along a gradient of human footprint. - Ecology 95: 2084–
2095.

Singer, F. J. and Schoenecker, K. A. 2003. Do ungulates accelerate or decelerate nitrogen
cycling? - Forest Ecology and Management 181: 189–204.

Sitters, J., Bakker, E. S., Veldhuis, M. P., Veen, G. F., Venterink, H. O. and Vanni, M. J. 2017.
The stoichiometry of nutrient release by terrestrial herbivores and its ecosystem
consequences. - Frontiers in Earth Science 5: 1–8.

Smith, A. C., Dahlin, K. M., Record, S., Costanza, J. K., Wilson, A. M. and Zarnetske, P. L.
2021. The GEODIV R package: Tools for calculating gradient surface metrics. - Methods
Ecol Evol 12: 2094–2100.

Spooner, F. E. B., Pearson, R. G. and Freeman, R. 2018. Rapid warming is associated with
population decline among terrestrial birds and mammals globally. - Global Change
Biology 24: 4521–4531.

Straus, S., Forbes, C., Little, C. J., Germain, R. M., Main, D. A., O'Connor, M. I., Thompson, P.
L., Ford, A. T., Gravel, D. and Guzman, L. M. 2024. Macroecological constraints on
species' 'movement profiles': Body mass does not explain it all. - Global Ecol Biogeogr
33: 227–243.

Subalusky, A. L. and Post, D. M. 2019. Context dependency of animal resource subsidies. -
Biological Reviews 94: 517–538.

Tylianakis, J. M., Rand, T. A., Kahmen, A., Klein, A.-M., Buchmann, N., Perner, J. and
Tscharntke, T. 2008. Resource Heterogeneity Moderates the Biodiversity-Function
Relationship in Real World Ecosystems (M Loreau, Ed.). - PLoS Biol 6: e122.

Veldhuis, M. P., Gommers, M. I., Olff, H. and Berg, M. P. 2018. Spatial redistribution of
nutrients by large herbivores and dung beetles in a savanna ecosystem. - Journal of
Ecology 106: 422–433.

Wilensky, U. 1999. NetLogo.

548